

Research Article

Synthesis, Characterization, and Magnetic and Thermal Studies on Some Metal(II) Thiophenyl Schiff Base Complexes

Aderoju Amoke Osowole

Inorganic Chemistry Unit, Department of Chemistry, University of Ibadan, Ibadan, Nigeria

Correspondence should be addressed to Aderoju Amoke Osowole, aderoju30@yahoo.com

Received 7 November 2010; Revised 22 December 2010; Accepted 31 December 2010

Academic Editor: Maurizio Peruzzini

Copyright © 2011 Aderoju Amoke Osowole. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

4-(Thiophen-3-yl)-aniline undergoes condensation with *o*-vanillin to form an ONS donor Schiff base, 2-methoxy-6-[(4-thiophene-3-yl-phenylimino)-methyl]-phenol, which forms complexes of the type $[ML_2]xH_2O$ (where M = Mn, Co, Ni, Cu, Zn, Pd). These complexes are characterized by elemental analysis, 1H nmr, electronic, mass, and IR spectroscopies and conductance measurements. The electronic, IR and CHN data are supportive of a 4-coordinate tetrahedral geometry for Mn(II), Co(II), Ni(II), and Zn(II) complexes and square-planar geometry for Cu(II) and Pd(II) complexes, with the chromophores N_2O_2 . The magnetic data reveals that the complexes are magnetically dilute and mononuclear with exception of the Cu(II) complex, which exhibits some anti-ferromagnetisms. The complexes are air-stable solids, and none is an electrolyte in nitro methane.

1. Introduction

My group has in the last five years been actively involved in the synthesis, characterization, and thermal and biological properties of various Schiff bases and their M(II) chelates (M = VO, Mn, Co, Ni, Cu, Zn, Pd), with the objectives of deriving Schiff base chelates which can be used as precursors in metal-organic chemical vapor depositions (MOCVD) and those with good *in vitro* antimicrobial activities as surface cleaning agents [1–5]. Thiophenyl Schiff bases are particularly interesting because of their wide range of activities such as anticancer activity as shown by benzyl-N-[1-(thiophenyl-3-yl)ethylidene] hydrazine carbodithioate [6], antibacterial and antifungal activities typified by thiophenyl-azetidinones, -cephalexins and -vinyl anilines [7–16] as well as optical exhibited by thiophene-2-aldazine [17]. Other activities include structural activity exemplified by the hexacarbonyliron complex of N-(2-thienylmethylidene)aniline, which is composed of two thienyl moieties derived from the original thienyl imine and coupled together by a C–C bond [18]. Furthermore, Palladium imine complexes of 2-thiophenecarboxaldehyde are used as catalysts in the Suzuki cross-coupling of aryl bromides with phenyl boric acid [19]. Extensive literature reviews show that no studies are

reported on the Schiff base derived from *o*-vanillin and 4-(thiophene-3-yl)-aniline and its metal(II) chelates [6–19]. Thus, the objective of this work is to synthesize, characterize, and investigate the magnetic and thermal properties of the Schiff base, 2-methoxy-6-[(4-thiophene-3-yl-phenylimino)-methyl]-phenol and its Mn(II), Co(II), Ni(II), Cu(II), Zn(II) and Pd(II) complexes. These metal complexes and its ligands are new, being reported here for the first time.

2. Experimental Details

2.1. Materials and Physical Measurements. Reagent grade *o*-vanillin, 4-(thiophene-3-yl)-aniline, manganese(II) nitrate hydrate, cobalt(II) nitrate hexahydrate, nickel(II) nitrate hexahydrate, copper(II) nitrate hexahydrate, zinc(II) nitrate hexahydrate, and palladium(II) chloride are purchased from BDH and Aldrich chemicals and are used as received. Solvents are dried and distilled before use according to standard procedures. Melting points (uncorrected) are determined using the Stuart scientific melting point SMP1 machine, and conductivities of 10^{-3} M solutions of the complexes are measured in nitromethane at 25°C using a MC-1, Mark V conductivity meter with a cell constant of 1.0. The solid

reflectance spectra are recorded on a Perkin-Elmer λ 20 spectrophotometer while infrared spectra are measured as KBr discs on a Perkin-Elmer FTIR paragon 1000 spectrometer in the range 4000–400 cm^{-1} . The elemental analyses C, H, and N are recorded on GmbH VarioEl analyser, and manganese, cobalt, nickel, copper, zinc, and palladium are determined titrimetrically and by atomic absorption spectroscopy [20]. The ^1H nmr spectra are recorded on a 300 MHz Oxford Varian NMR instrument in CDCl_3 at 295 K. ^1H chemical shifts are referenced to the residual signals of the protons of CDCl_3 and are quoted in ppm. Magnetic susceptibilities are measured on Johnson Matthey magnetic susceptibility balance, and diamagnetic corrections are calculated using Pascal's constants [21]. Thermogravimetric analyses are done in static air, using a T6 Linseis thermal analyser with a heating rate of 10°C/min in the range 30–700°C. The MALDI-TOF mass and atomic absorption spectra are obtained using a Bruker Daltonic Reflex TOF spectrometer with graphite as matrix and Perkin Elmer Analyst 200 coupled to Winlab 32 software assembly, respectively.

2.2. Preparation of the Schiff Base (2-methoxy-6-[(4-thiophene-3-yl-phenylimino)-methyl]-phenol). A 20 mL solution of 8.71 mmol (1.33 g) *o*-vanillin in absolute ethanol is added dropwise to a stirring solution of 8.71 mmol (1.53 g) of 4-(thiophene-3-yl)-aniline in 30 mL of absolute ethanol. The resulting orange-colored solution is refluxed for 4 h after addition of 4 drops of acetic acid. The orange product formed on cooling to room temperature is filtered and recrystallized from ethanol. The yield of the title compound is 1.88 g (70%). ^1H nmr (ppm) δ 11.13 (s, 1H, OH), 9.93 (s, 1H, HCN), 8.70 (1H, s, C_2HS), 7.35–7.43 (m, 3H, C_6H_3); 7.03–7.24 (m, 4H, C_6H_4); 6.91–7.01 (m, 2H, $\text{C}_2\text{H}_2\text{S}$); 3.96 (s, 3H, OCH_3).

2.3. Preparation of the Metal(II) Complexes. The various complexes are prepared by gradual addition of 0.35 mmol (0.06–0.11 g) $\text{M}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (M = Mn, Co, Ni, Cu, Zn) neat to a stirring 0.7 mmol (0.22 g) of the ligand in 30 mL of absolute ethanol. The resulting solutions are then buffered with 0.7 mmol (0.10 mL) of triethylamine and refluxed for 6 h during which the products formed. The precipitated solids are filtered, washed with ethanol, and dried over anhydrous calcium chloride. The yields are 0.12 g (50%), 0.17 g (70%), 0.17 g (70%), 0.17 g (70%), and 0.14 g (60%), respectively.

The Pd(II) complex is prepared using a similar method. 0.36 mmol (0.064 g) of Pd(II) chloride, in 10 mL of absolute ethanol, is added dropwise to a stirring solution of 0.72 mmol (0.23 g) of the ligand in 30 mL of absolute ethanol. The resulting solution is then buffered with 0.72 mmol (0.11 mL) of triethylamine and refluxed for 6 h during which the product is formed. The product is filtered, washed with ethanol, and dried over anhydrous calcium chloride. The yield is 0.18 g (70%).

Proton nmr measurements are done only for the diamagnetic Zn(II) and Pd(II) complexes. Zn(II) complex: ^1H nmr (ppm) 8.70 (s, 1H, HCN), 8.34 (1H, s, C_2HS), 7.28–7.43

(m, 3H, C_6H_3); 7.01–7.10 (m, 4H, C_6H_4); 6.63–6.94 (m, 2H, $\text{C}_2\text{H}_2\text{S}$); 3.94 (s, 3H, OCH_3).

Pd(II) complex: ^1H nmr (ppm) 8.67 (s, 1H, HCN), 8.30 (1H, s, C_2HS), 7.48–7.66 (m, 3H, C_6H_3); 7.00–7.47 (m, 4H, C_6H_4); 6.88–6.98 (m, 2H, $\text{C}_2\text{H}_2\text{S}$); 3.94 (s, 3H, OCH_3).

3. Results and Discussion

The equations for the formation of the complexes are



(where M = Zn(II), Cu(II), Mn(II), Co(II), Ni(II))



All complexes adopt $[\text{ML}_2]$ stoichiometry, except Mn(II), Co(II) and Ni(II) complexes that form as $[\text{ML}_2]_x\text{H}_2\text{O}$, where $x = 1$ and 0.5, respectively. Proposed structures for the ligand and the Cu(II) complex are shown in Figure 1. The formation of this ligand is confirmed by microanalysis and ^1H nmr. The colors, melting points, and room temperature magnetic moments (μ_{eff}) of the compounds are presented in Table 1. Attempts to isolate suitable crystals for single X-ray structural determination have not been successful so far.

3.1. Infrared Spectra. The relevant infrared bands of the compounds are presented in Table 2. The broadband at 3360 cm^{-1} in the ligand, which is conspicuously absent in the spectra of the metal(II) Schiff base complexes, is assigned as νOH stretching frequency, and it confirms involvement of the phenol O in chelation. It is broad due to intramolecular hydrogen bonding, usually very strong in Schiff bases [1]. The new broadband at 3500 cm^{-1} in the spectra of Co(II), Ni(II), and Mn(II) complexes is assigned to the νOH frequency of crystallization H_2O . The uncoordinated C=N stretching vibrations are observed as three bands between 1614–1521 cm^{-1} in the ligand [2–5] and 1639–1503 cm^{-1} in the metal complexes with exceptions of the Cu(II) and Pd(II) complex which have two bands. The bathochromic/hypsochromic shifts of these bands in the complexes are attributed to the involvement of N atom of C=N in coordination to the metal ions. Moreover, it has been documented that square planar Pd(II) and Cu(II) Schiff base complexes do exhibit geometric isomerism [22], with the trans isomer showing two $\nu\text{C}=\text{N}$ bands and the cis isomer a lone $\nu\text{C}=\text{N}$ band. The spectra of the Cu(II) and Pd(II) complexes in this work show two $\nu\text{C}=\text{N}$ bands and are consequently in the trans-isomeric form. The $\nu\text{Ph}/\text{C}-\text{O}$ and $\delta\text{C}-\text{H}$ vibrations of the ligand are observed at 1461–1364 and 972 cm^{-1} , respectively. These suffer bathochromic shifts to 1298–1177 and 896–720 cm^{-1} in the Schiff base complexes due to the coordination of the phenol oxygen atom and pseudoaromatic nature of the chelates [5, 6]. The observation of new bands at 480–405 and 581–542 cm^{-1} due to $\nu(\text{M}-\text{O})$ and $\nu(\text{M}-\text{N})$ [11, 14, 22] is further evidence of coordination.

3.2. Electronic Spectra and Magnetic Moments. The electronic spectral data for the complexes are presented in Table 2.




FIGURE 1: Proposed structures for the ligand (a) and its Cu(II) complex (b).

TABLE 1: Analytical data for the ligand and its complexes.

Compound (empirical formula)	Formula mass	Color	m/z (100%)	μ_{eff}	% Yield	Λ_m^*	M.p (°C)	Analysis (calculated)			
								% C	% H	% N	% M
HL (C ₁₈ H ₁₅ NO ₂ S)	309.38	Orange	309	—	80	—	150–151	69.82 (69.88)	4.53 (4.89)	4.35 (4.53)	—
[MnL ₂]H ₂ O (MnC ₃₆ H ₃₀ N ₂ S ₂ O ₅)	689.73	green	671	5.70	50	10.0	338–340	62.54 (62.69)	4.32 (4.38)	4.02 (4.06)	8.02 (7.97)
[CoL ₂]H ₂ O (CoC ₃₆ H ₃₀ N ₂ S ₂ O ₅)	693.71	Brick Red	675	4.33	70	20.0	290–292	62.36 (62.33)	3.64 (4.36)	4.06 (4.04)	8.48 (8.50)
[NiL ₂]1/2H ₂ O (NiC ₃₆ H ₃₀ N ₂ S ₂ O _{4.5})	684.48	Yellow	675	3.10	70	15.0	193–195	63.07 (63.17)	3.86 (4.27)	4.17 (4.09)	8.55 (8.58)
[CuL ₂] ₂ (Cu ₂ C ₇₂ H ₅₆ N ₄ S ₄ O ₈)	680.31	Brown	680	1.56	70	9.0	205–207	64.14 (63.56)	4.06 (4.15)	4.01 (4.12)	9.35 (9.34)
[ZnL ₂] (ZnC ₃₆ H ₂₈ N ₂ S ₂ O ₄)	681.76	Yellow	681	D	60	17.0	316–318	63.13 (63.42)	3.84 (4.14)	3.81 (4.11)	9.52 (9.53)
[PdL ₂] (PdC ₃₆ H ₂₈ N ₂ S ₂ O ₄)	723.18	green	720	D	70	12.0	148–150	60.37 (59.79)	4.32 (3.90)	3.86 (3.87)	14.70 (14.72)

* $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$, D: diamagnetic.

The Mn(II) Schiff base complex shows two absorption bands at 15.00 and 23.11 kK, respectively, consistent with a four-coordinate, tetrahedral geometry and are assigned to ${}^6A_1 \rightarrow {}^4E_1$ (ν_1) and ${}^6A_1 \rightarrow {}^4A_1$ (ν_2) transitions [23]. A room temperature moment of 5.92 B.M is usually observed for the Mn(II) compounds, regardless of stereochemistry because the ground term is an 6A_1 , and thus, orbital contribution

is nil. This Mn(II) complex has a moment of 5.70 B.M. complementary of tetrahedral geometry [1].

The cobalt(II) Schiff base complex gives two absorption bands at 10.82 and 18.20 kK, respectively, typical of a four-coordinate tetrahedral geometry and is assigned to ${}^4A_2 \rightarrow {}^4T_1(P)$ (ν_2) and ${}^4A_2 \rightarrow {}^4T_1(P)$ (ν_3). The transition ${}^4A_2 \rightarrow {}^4T_2$ (ν_1) in the range 5–7 kK is not observed as usual

TABLE 2: Relevant infrared and electronic spectral data of the ligand and its complexes.

Compound	ν_{OH}	$\nu(\text{C}=\text{N}) + \nu(\text{C}=\text{C})$	$\nu_{\text{Ph/C-O}}$	$\delta_{\text{C-H}}$	$\nu(\text{M-N})$	$\nu(\text{M-O})$	Electronic transitions (kK)
HL	3360b	1614s 1593s 1521s	1461s 1364s	972s	—	—	33.50, 36.25, 40.20
[MnL ₂]H ₂ O	3500b	1612s 1589s 1551s	1298s 1194s	848s 720s	580m 561s	450m 420s	15.00, 23.11, 30.26, 33.45, 40.20
[CoL ₂]H ₂ O	3500b	1609s 1557s 1503s	1231m 1180m	840s 777m	575s 537m	475m 415m	10.82, 18.20, 32.0, 37.40, 41.0
[NiL ₂]1/2H ₂ O	3500b	1599s 1585s 1537s	1233s 1192s	853s 780m	581s 564s	460s 453s	14.29, 20.21, 30.72, 36.25, 40.40
[CuL ₂]	—	1587s 1538s	1237m 1193m	856s 795s	573s 557s	465s 410s	14.09, 21.60, 30.92, 34.35, 41.20
[ZnL ₂]	—	1609s 1582s 1504s	1231m 1197m	896s 731m	579s 562m	480m 405m	28.20, 35.00, 42.00
[PdL ₂]	—	1592s 1540s	1240s 1177s	843m 783m	562m 542m	450m 420m	18.20, 26.32, 30.25, 39.20, 43.00

b: broad, m: medium, s: strong., 1 kK: 1000 cm⁻¹.

since it lies in the infrared region [10]. This geometry is corroborated by a moment of 4.33 B.M [11].

Nickel(II) complexes are known to exhibit complicated equilibria between coordination numbers six (octahedral) to four (square planar/tetrahedral) [24]. The Ni(II) Schiff base complex exhibits two absorption bands at 14.29 and 20.21 kK typical of a 4-coordinate tetrahedral geometry, assigned to $^3T_1(\text{F}) \rightarrow ^3T_2, (\nu_2)$ and $^3T_1(\text{F}) \rightarrow ^3A_2, (\nu_3)$ transitions. Its moment of 3.10 B.M is complimentary of tetrahedral geometry, since moments of 3.1–3.5 B.M. are reported for distorted tetrahedral complexes [12].

The copper(II) complex displays two bands at 14.09 and 21.60 kK, assigned to $^2B_{1g} \rightarrow ^2A_{1g}$ and $^2B_{1g} \rightarrow ^2E_{1g}$ transitions of 4-coordinate, square planar geometry [13]. A moment of 1.9–2.2 B.M. is usually observed for mononuclear copper(II) complexes, regardless of stereochemistry [14]. A magnetic moment of 1.56 B.M. is observed for this complex, indicative of the presence of some anti-ferromagnetic interactions, operating through Cu–Cu interactions [25]. However, this could not be probed further due to lack of facilities for variable temperature magnetic measurements and nonsuitable crystal for single X-ray diffraction measurement (Figure 1).

The Zn(II) complex expectedly shows only charge transfer transition from M → L and π – π^* transitions, as no d-d transition is expected at 28.20, 35.00, and 42.0 kK, respectively. This complex is diamagnetic, confirming its tetrahedral geometry [12–14].

The Pd(II) complex shows absorption bands at 18.20 and 26.32 kK, typical of square planar geometry and is assigned to $^1A_{1g} \rightarrow ^1B_{1g}$ and $^1A_{1g} \rightarrow ^1E_{2g}$ transitions. This complex is expectedly diamagnetic [26].

3.3. ¹H nmr Spectra. The ligand shows the phenolic proton as a singlet at δ 11.13 ppm (s, 1H, OH), the imine (s, 1H, HCN), and 2-thiophenyl (s, H, C₂HS) protons resonate as singlets at 9.93 and 8.70 ppm, respectively. The *o*-vanillin (m, 3H, C₆H₃) and phenyl (aniline) protons (m, 4H, C₆H₄) are both observed as multiplets at δ 7.35–7.43 and 7.03–7.24 ppm, respectively. The two protons of thiophenyl ring at 4 and 5 positions come up as a multiplet at 6.91–7.01 ppm (m, 2H, –C₂H₂S), and the methoxy protons are observed as a singlet at 3.96 ppm (s, 3H, OCH₃).

In the Zn(II) complex spectrum, the phenolic proton disappears, an indication of coordination of the phenolic oxygen to the Zn(II) ion, and other protons are all upfield shifted in comparison to the ligand. The imine hydrogen resonates as a singlet at 8.70 ppm (s, 1H, HCN) while the 2-thiophenyl proton is seen at 8.34 ppm as a singlet (s, H, C₂HS).

The *o*-vanillin protons resonate as a multiplet at δ 7.28–7.43 ppm (m, 3H, C₆H₃), and the phenyl(aniline) protons are seen as multiplets at 7.01–7.10 ppm (m, 4H, C₆H₄). The two protons of thiophenyl ring at 4 and 5 positions resonate as a multiplet at 6.63–6.94 ppm (m, 2H, –C₂H₂S), and the methoxy protons are seen as a singlet at 3.94 ppm (s, 3H, OCH₃). This shift shows deshielding, a consequence of coordination of the imine nitrogen atom [27].

The Pd(II) complex spectrum reveals the absence of the phenolic proton. This is indicative of involvement of the phenol oxygen in coordination to the Pd(II) ion. The other protons are all upfield shifted in comparison to the ligand with exceptions of the *o*-vanillin and phenyl(aniline) protons which are downfield shifted. The imine hydrogen (s, 1H, HCN) and 2-thiophenyl (s, H, C₂HS) protons resonate as singlets at 8.67 and 8.30 ppm, respectively. The *o*-vanillin (m, 3H, C₆H₃), phenyl(aniline) (m, 4H, C₆H₄), and two protons of thiophenyl ring (m, 2H, –C₂H₂S) at 4 and 5 positions are all seen as multiplets at δ 7.48–7.66, 7.00–7.47, and 6.88–6.98 ppm, respectively. The methoxy protons resonate as a singlet at 3.94 ppm (s, 3H, OCH₃). These shifts are indicative of coordination of the imine nitrogen atom to the Pd(II) ion [26].

3.4. Mass Spectroscopy and Thermal Studies. The mass spectra of ligand and the complexes showed peaks attributed to the molecular ions m/z at 309 [L]⁺; 671 [MnL₂-2H]⁺; 675 [CoL₂-2H]⁺; 675 [NiL₂-2H]⁺; 680 [CuL₂-2H]⁺; 681 [ZnL₂-2H]⁺ and 720 [PdL₂], respectively, and are presented in Table 1.

The thermal degradation of the ligand and complexes is presented in Table 3. The ligand, HL, decomposes in three steps. First, the loss of the fragment C₂H₂ and 0.5 mol N₂ at 30–220°C, with mass losses of (obs. = 12.96%, calc. = 12.93%). The next step involves the loss of the organic fraction, C₁₁H₈O₂S, with mass losses of (obs. = 66.32%,

TABLE 3: Thermal data for the ligand and its complexes.

Compound	Temperature range (°C)	TG weight loss (%)		Assignments
		Calc.	Found	
HL (C ₁₈ H ₁₅ NSO ₂)	30–220	12.93	12.96	C ₂ H ₂ + 0.5N ₂
	220–420	65.94	66.32	C ₁₁ H ₈ O ₂ S
	420–700	22.78	21.01	C ₅ H ₅
[MnL ₂]H ₂ O (MnC ₃₆ H ₃₀ N ₂ S ₂ O ₅)	30–200	9.57	9.40	H ₂ O + 1.5O ₂
	210–450	44.07	44.45	C ₂₀ H ₁₈ NS
	450–700	22.04	22.53	C ₈ H ₁₀ OS Mn (residue)
[CoL ₂]H ₂ O (CoC ₃₆ H ₃₀ N ₂ S ₂ O ₅)	30–200	4.90	4.77	CH ₄ + H ₂ O
	200–400	20.76	21.45	C ₆ H ₈ S ₂
	400–700	57.08	57.50	C ₂₄ H ₁₆ O ₄ N ₂ Co (residue)
[NiL ₂]1/2H ₂ O (NiC ₃₆ H ₃₀ N ₂ S ₂ O _{4.5})	30–240	10.67	10.56	0.5H ₂ O + SO ₂
	240–420	23.08	22.73	C ₁₀ H ₁₀ N ₂
	420–700	55.81	55.51	C ₂₅ H ₁₈ SO ₂ Ni (residue)
[CuL ₂] (CuC ₃₆ H ₂₈ N ₂ S ₂ O ₄)	30–200	17.93	17.86	C ₈ H ₁₂ N
	200–400	33.81	33.63	C ₁₃ H ₁₀ O ₂ S
	400–700	38.81	42.00	C ₁₅ H ₆ NO ₂ S Cu (residue)
[ZnL ₂] (ZnC ₃₆ H ₂₈ N ₂ S ₂ O ₄)	30–260	5.00	5.17	H ₂ S
	260–440	27.87	27.80	C ₁₁ H ₁₀ OS
	440–700	52.22	52.21	C ₂₂ H ₁₆ N ₂ O ₃ Zn (residue)
[PdL ₂] (PdC ₃₆ H ₂₈ N ₂ S ₂ O ₄)	30–210	3.87	3.50	C ₂ H ₄
	210–400	40.79	41.33	C ₁₈ H ₁₅ SO ₂
	400–700	33.32	33.42	C ₁₃ H ₉ N ₂ SO PdO (residue)

calc. = 65.94%) at 220–420°C. The final step is the loss of the fragment C₅H₅, with mass losses of (obs. = 22.78%, calc. = 21.01%) at 420–700°C.

The Mn(II) complex decomposes in three phases. The first phase corresponds to the loss of 1.5 moles of O₂ and H₂O between 30–200°C with mass losses of (obs. = 9.40%, calc. = 9.57%). The second phase is from 210 to 450°C and is attributed to the loss of the organic moiety C₂₀H₁₈NS with mass losses of (obs. = 44.37%, calc. = 44.07%). The final phase shows the loss of the organic moiety, C₈H₁₀OS, at 450–700°C with mass losses of (obs. = 22.53%, calc. = 22.33%) leaving Mn as the final product, and the fragment C₈N is lost as 8CO₂ and 0.5N₂.

The decomposition of the Co(II) complex also occurred in three steps. The first step is due to the loss of a mole of water and CH₄ at 30–200°C, with mass losses of (obs. = 4.77%, calc. = 4.90%). The successive decomposition occurs within a temperature range of 200–400°C and is attributed to the loss of the organic moiety C₆H₈S₂ with mass losses of (obs. = 21.45%, calc. = 20.76%). The last step involves the loss of the organic moiety, C₂₄H₁₆N₂O₄, at 400–700°C with mass losses of (obs. = 57.50%, calc. = 57.08%). The final product is Co, and the C₅ fragment is lost as 5CO₂.

The TGA curve of the Ni(II) complex reveals a three-step decomposition. The first is the loss of 0.5 mole of water and SO₂ at 30–240°C, with mass losses of (obs. = 10.56%, calc. = 10.67%). The second step ranges from 240 to 420°C and is assigned to the loss of the organic moiety, C₁₀H₁₀N₂ with mass losses (obs. = 22.73%, calc. = 23.08%). The final step is within a temperature range of 420–700°C and is attributed to the loss of the organic moiety C₂₅H₁₈SO₂ (obs. = 55.51%, calc. = 55.81%). The remaining fraction is Ni residue, and the fragment CH is lost as CO₂ + 0.5H₂.

Cu(II) complex decomposes in three steps. The first step is attributed to the loss of the fragment C₈H₁₂N, with mass losses of (obs. = 17.86%, calc. = 17.89%) at 30–200°C. The second step ranges from 200 to 400°C and is attributed to the loss of the fragment C₁₃H₁₀O₂S, with mass losses of (obs. = 33.63%, calc. = 33.74%). The final step is from 400 to 700°C corresponding to the loss of the organic moiety C₁₅H₄O₂NS, with mass losses of (obs. = 42.0%, calc. = 38.72%). The remaining residue is Cu.

The Zn(II) complex decomposes in three steps. Step one is between 30–260°C, which indicates the loss of H₂S, with mass losses of (obs. = 5.17%, calc. = 5.00%). The second step involves the loss of the organic moiety C₁₁H₁₀OS,

from 260 to 440°C, with mass losses of (obs. = 27.80%, calc. = 27.87%). The final step is attributed to the loss of the organic moiety, C₂₂H₁₆N₂O₃, at 440–700°C, with mass losses of (obs. = 52.21%, calc. = 52.22%), leaving behind the Zn residue, and the C₃ fragment is lost as 3CO₂.

The Pd(II) complex also decomposes in three phases. The first phase is between 30–210°C and is attributed to loss of C₂H₄, with mass losses of (obs. = 3.50%, calc. = 3.87%). The second phase involves the loss of the organic moiety, C₁₈H₁₅O₂S at 210–400°C, with mass losses of (obs. = 41.33%, calc. = 40.79%) while the final stage involves the loss of the organic moiety, C₁₃H₉N₂SO, from 400 to 700°C, with mass losses of (obs. = 33.42%, calc. = 33.32%), leaving behind PdO residue, and the fragment C₃ is lost as 3CO₂.

In all cases with the exceptions of the ligand and the Cu(II) complex, the decomposition pattern showed the loss of carbon fragments which got oxidized to CO₂, and hydrogen or nitrogen which were lost as gases. Thus, the decomposition pattern corroborates the proposed formulation of the complex.

3.5. Conductance. The molar conductances of the complexes in nitromethane are below 20.0 ohm⁻¹ cm² mol⁻¹ confirming their covalent nature. A value in the range 75–90 ohm⁻¹ cm² mol⁻¹ is expected for a 1 : 1 electrolyte [12].

4. Conclusion

The Schiff-base ligand coordinates to the Mn(II), Ni(II), Co(II), Cu(II), Pd(II), and Zn(II) ions in a tetradentate manner using the N₂O₂ chromophores. The assignment of a 4-coordinate square-planar geometry to Cu(II) and Pd(II) complexes and tetrahedral geometry to Mn(II), Ni(II), Co(II), and Zn(II) complexes is corroborated by elemental analysis, thermal, magnetic, and electronic spectral measurements. The Cu(II) and Pd(II) complexes exhibit geometric isomerism and are in the trans-isomeric form as confirmed by their infrared spectra. Furthermore, the Cu(II) complex exhibits some anti-ferromagnetic interactions, operating through a dimeric structure while the other complexes are mononuclear.




Acknowledgment

The author thanks The Alexander von Humboldt (AvH) Foundation for a Georg Forster Fellowship.

References

- [1] A. A. Osowole, G. A. Kolawole, R. Kempe, and O. E. Fagade, "Spectroscopic, magnetic and biological studies on some metal(II) complexes of 3-(4,6-Dimethyl-2-Pyrimidinylamino)-1-Phenyl-2-Butenone and their adducts with 2,2-Bipyridine and 1,10-Phenanthroline," *Synthesis and Reactivity in Inorganic, Metal-Organic and Nano-Metal Chemistry*, vol. 39, no. 3, pp. 165–174, 2009.
- [2] A. A. Osowole, G. A. Kolawole, and O. E. Fagade, "Synthesis, characterization and biological studies on unsymmetrical Schiff-base complexes of nickel(II), copper(II) and zinc(II) and adducts with 2,2'-dipyridine and 1,10-phenanthroline," *Journal of Coordination Chemistry*, vol. 61, no. 7, pp. 1046–1055, 2008.
- [3] A. A. Osowole, B. C. Ejelolu, and S. A. Balogun, "Spectroscopic, magnetic and antibacterial properties of some metal(II) unsymmetric Schiff-base complexes and their mixed-ligand analogs," *Journal of Ultra Chemistry*, vol. 20, no. 3, pp. 549–558, 2008.
- [4] A. A. Osowole and O. E. Fagade, "Synthesis, characterization and biopotency of some metal(II) β -ketoimines and their mixed-ligand complexes," *Polish Journal of Chemistry*, vol. 81, no. 12, pp. 2039–2048, 2007.
- [5] A. A. Osowole, G. A. Kolawole, and O. E. Fagade, "Synthesis, physicochemical, and biological properties of nickel(II), copper(II), and zinc(II) complexes of an unsymmetrical tetradentate Schiff base and their adducts," *Synthesis and Reactivity in Inorganic, Metal-Organic and Nano-Metal Chemistry*, vol. 35, no. 10, pp. 829–836, 2005.
- [6] M. H. E. Chan, K. A. Crouse, M. I. M. Tahir, R. Rosli, N. Umar-Tsafe, and A. R. Cowley, "Synthesis and characterization of cobalt(II), nickel(II), copper(II), zinc(II) and cadmium(II) complexes of benzyl N-[1-(thiophen-2-yl)ethylidene] hydrazine carbodithioate and benzyl N-[1-(thiophen-3-yl)ethylidene] hydrazine carbodithioate and the X-ray crystal structure of bisbenzyl N-[1-(thiophen-2-yl)ethylidene] hydrazine carbodithioatenickel(II)," *Polyhedron*, vol. 27, no. 4, pp. 1141–1149, 2008.
- [7] P. S. Kendrekar, S. V. More, P. S. Patil, S. R. S. Bhusare, and R. P. Pawar, "Synthesis of 2-(2-hydroxy-3-iodo-5-bromo phenyl)-3-(substituted phenyl)-4-thiazolidinones as antibacterial agents," *Oriental Journal of Chemistry*, vol. 18, no. 3, pp. 595–597, 2002.
- [8] N. Idrees, M. Siddique, A. G. Doshi, and A. W. Raut, "Synthesis of N-substituted phenyl-4-thiophenyl-2-azetidinones and its antimicrobial activity," *Oriental Journal of Chemistry*, vol. 17, no. 1, pp. 143–146, 2001.
- [9] N. Idrees, M. Siddique, S. D. Patil, A. G. Doshi, and A. W. Raut, "Synthesis of Schiff bases of thiophene-2-carboxaldehyde and its antimicrobial activity," *Oriental Journal of Chemistry*, vol. 17, no. 1, pp. 131–133, 2001.
- [10] B. N. Reddy, P. G. Avaji, P. S. Badami, and S. A. Patil, "Synthesis, spectral and biological studies of cobalt(II), nickel(II) and copper(II) complexes with 1,5-bis(thiophenylidene) thiocarbonylhydrazones," *Journal of Saudi Chemical Society*, vol. 11, no. 2, pp. 253–268, 2007.
- [11] Z. H. Chohan, H. Pervez, K. M. Khan, A. Rauf, and C. T. Supuran, "Binding of transition metal ions [cobalt, copper, nickel and zinc] with furanyl-, thiophenyl-, pyrrolyl-, salicylyl- and pyridyl-derived cephalosporins as potent antibacterial agents," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 19, no. 1, pp. 51–56, 2004.
- [12] Z. H. Chohan, H. Pervez, K. M. Khan, A. Rauf, G. M. Maharvi, and C. T. Supuran, "Antifungal cobalt(II), copper(II), nickel(II) and zinc(II) complexes of furanyl-thiophenyl-, pyrrolyl-, salicylyl- and pyridyl-derived cephalosporins," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 19, no. 1, pp. 85–90, 2004.
- [13] Z. H. Chohan, H. Pervez, A. Rauf, A. Scozzafava, and C. T. Supuran, "Antibacterial Co(II), Cu(II), Ni(II) and Zn(II) complexes of thiazazole derived furanyl, thiophenyl and pyrrolyl schiff bases," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 17, no. 2, pp. 117–122, 2002.

- [14] Z. H. Chohan and S. Kausar, "Synthesis, characterization and biological properties of tridentate NNO, NNS and NNN donor thiazole-derived furanyl, thiophenyl and pyrrolyl Schiff bases and their Co(II), Cu(II), Ni(II) and Zn(II) metal chelates," *Metal-Based Drugs*, vol. 7, no. 1, pp. 17–22, 2000.
- [15] A. I. P. Sinha and M. Bala, "Coordination behavior of herbicidal Schiff bases derived from 2-amino-6-ethoxybenzothiazole towards copper(II)," *Asian Journal of Chemistry*, vol. 3, no. 1, pp. 45–51, 1991.
- [16] P. Mittal, S. Joshi, V. Panwar, U. Vatsala, and V. Uma, "Biologically active Co (II), Ni (II), Cu (II) and Mn(II) Complexes of Schiff bases derived from vinyl aniline and heterocyclic aldehydes," *International Journal of ChemTech Research*, vol. 1, no. 2, pp. 225–232, 2009.
- [17] M. Ghazzali, V. Langer, C. Lopes, A. Eriksson, and L. Öhrström, "Syntheses, crystal structures, optical limiting properties, and DFT calculations of three thiophene-2-aldazine Schiff base derivatives," *New Journal of Chemistry*, vol. 31, no. 10, pp. 1777–1784, 2007.
- [18] D. L. Wang, W. S. Hwang, L. C. Liang, L. I. Wang, L. Lee, and M. Y. Chiang, "Reaction of a thienyl schiff base with diiron nonacarbonyl: characterization and structures of [μ -N-(((2,3- η : η)-5-methyl-2-thienyl)methyl)- η : η (N)-anilino]hexacarbonyldiiron and [μ -N-((anilino(2-thienyl)methyl) ((2,3- η : η)-2-thienyl)methyl)- η : η (N)," *Organometallics*, vol. 16, no. 14, pp. 3109–3113, 1997.
- [19] J. Wiedermann, K. Mereiter, and K. Kirchner, "Palladium imine and amine complexes derived from 2-thiophenecarboxaldehyde as catalysts for the Suzuki cross-coupling of aryl bromides," *Journal of Molecular Catalysis A*, vol. 257, no. 1-2, pp. 67–72, 2006.
- [20] J. Bassett, R. C. Denney, G. H. Jeffery, and J. Mendham, *Vogel's Textbook of Quantitative Inorganic Analysis*, ELBS, London, UK, 1978.
- [21] A. Earnshaw, *The Introduction to Magnetochemistry*, Academic Press, London, UK, 1980.
- [22] A. A. Nejo, G. A. Kolawole, A. R. Opoku, C. Muller, and J. Wolowska, "Synthesis, characterization, and insulin-enhancing studies of unsymmetrical tetradentate Schiff-base complexes of oxovanadium(IV)," *Journal of Coordination Chemistry*, vol. 62, no. 21, pp. 3411–3424, 2009.
- [23] A. B. Lever, *Inorganic Electronic Spectroscopy*, Elsevier, London, UK, 1980.
- [24] E. Kwiatkowski and M. Kwiatkowski, "Unsymmetrical schiff base complexes of nickel(II) and palladium(II)," *Inorganica Chimica Acta*, vol. 42, no. C, pp. 197–202, 1980.
- [25] M. G. B. Drew and P. C. Yates, "Synthetic, structural and molecular mechanics investigation of some mono- and dinuclear copper(I) and copper(II) complexes of macrocyclic and related acyclic ligands," *Inorganica Chimica Acta*, vol. 118, no. 1, pp. 37–47, 1986.
- [26] S. Cherayath, J. Alice, and C. P. Prabhakaran, "Palladium(II) complexes of Schiff bases derived from 5-amino-2,4-(1H, 3H)pyrimidinedione (5-aminouracil) and 1,2-dihydro-1,5-dimethyl-2-phenyl-4-amino-3H-pyrazol-3-one," *Transition Metal Chemistry*, vol. 15, no. 6, pp. 449–453, 1990.
- [27] X. Han, Z. L. You, Y. T. Xu, and X. M. Wang, "Synthesis, characterization and crystal structure of a mononuclear zinc(II) complex derived from 2-methoxy-6-[(3-cyclohexylamino-propylimino)methyl]phenol," *Journal of Chemical Crystallography*, vol. 36, no. 11, pp. 743–746, 2006.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

