

Synthesis of conduritols and related compounds


Metin BALCI

Atatürk University, Department of Chemistry, Faculty of Science, Erzurum, TURKEY

Abstract: New and stereospecific syntheses for all conduritol isomers have been developed starting from appropriate 1,3-cyclohexadiene derivatives. Oxygen functionalities were introduced by photooxygenation. Application of the ene-reaction of singlet oxygen to 1,4-cyclohexadiene and its derivatives afforded diene system which can be easily trapped by second mol of singlet oxygen to give 49 and 50. Thiourea reduction of the peroxide linkages followed by oxidation gave the corresponding pentol derivatives. Furthermore, reaction of unsaturated endoperoxides with 1,2,4,5-tetraazine derivatives 59-63 afforded new bicyclic endoperoxides with unusual structures.

Introduction


Conduritols¹ are 1,2,3,4-cyclohexenetetrol isomers and interesting inhibitors for glycosidases². A number of conduritol derivatives have been found to possess antifeedant³, antibiotic, antileukemic, and growth-regulating¹ activity. There are also many plant metabolites like crotopoxide, senepoxide, pipoxide, (Scheme 1) which have four oxygen bonded substituents on a cyclohexene-ring, exhibiting interesting biological properties including tumour-inhibitory, antileukaemic, and antibiotic activities. Therefore, the synthesis of these compounds gain interest. In this paper, we describe a new synthetic methodology leading to the synthesis of various conduritols and related compounds. Before synthesis of conduritol derivatives we have developed a synthetic methodology for tetrahydroxy-cyclohexan derivatives. Therefore, we want to discuss at first the synthesis of some cyclitols derived from cyclohexane.




Cyclitols: Cyclohexanetetrols

Theoretically six cyclohexanetetrols diastereoisomers are possible (Scheme 2). Some of these occur in the nature⁴. These compounds have been synthesized either by reduction of the corresponding conduritols⁵ or by direct hydroxylation of 1,3-cyclohexadiene⁶ (Scheme 3). Direct hydroxylation of the appropriate cyclic dienes is not a suitable method for the synthesis of the corresponding tetrols.

Our synthetic sequence is based on introduction of the two oxygen functionality's by photooxygenation of 1,3-cyclohexadiene. The other oxygen functionality's are introduced through the classical peracid epoxidation reaction⁷. Oxidation of the endoperoxide **2** obtained by photooxygenation⁸ of cyclohexadiene, with *m*-chloroperbenzoic acid gives two isomeric epoxy endoperoxides in a 55:45 ratio. These isomers were separated by column chromatography on silica gel. The ¹H-NMR spectra of these epoxy endoperoxides are consistent with the symmetry of their structures. However we were not able to distinguish between the two isomers **3** and **4** on the basis of the spectral data. The configurational assignment of syn and anti to the isomeric epoxy endoperoxides is easily made by chemical means. We synthesized the anti isomer **4** by an independent route from oxepine-benzene oxide **6** as shown below.



Foster and Berchtold⁹ reported that reaction of oxepine-benzene oxide with singlet oxygen afforded the unsaturated anti epoxy endoperoxide **5**. Reduction of the double bond in **5** with diimide occurs without reduction of the peroxide bond to give anti epoxy endoperoxide **4** quantitatively, which was identical with the major product obtained by photooxygenation of 1,3-cyclohexadiene followed by epoxidation of the formed bicyclic endoperoxide **2** (Scheme 1).




Catalytic hydrogenation of **3** and **4** gave epoxy diols **7** and **9**. Since only the oxygen-oxygen bond breaks in this reaction⁸, the configuration at all four carbon atoms must be the same as in the starting material. For further characterization, **7** and **9** were converted to the corresponding diacetates **8** and **10** with acetic anhydride in pyridine. In the trans epoxide ring opening by addition of water in a classic S_N2 mechanism, both epoxides **7** and **9** gave the cyclitol **11**¹⁰. (Scheme 2).

After completing the synthesis of **11** we were interested in the synthesis of the anti-symmetrical diol **14** which opens entry to tetrols **15** and **16**. Therefore we studied the NEt₃-catalyzed rearrangement of **3** and **4**. Base-catalyzed reaction of bicyclic endoperoxides results generally in the formation of hydroxyketones. Treatment of **3** and **4** with triethylamine led within a few hours to the expected epoxy ketols **12** and **13** in high yields. While catalytic hydrogenation of **3** and **4** affords only symmetrical epoxy diol **7** and **9**, reduction of **12** and **13** with NaBH₄ in methanol provides a mixture of diols in both cases

(7/14 and 9/14). Newly formed epoxy diol **14** has been separated by column chromatography and identified by spectroscopic data.


Ring opening of **14** from which a pair of diastereoisomers is expected, with acidified water produced a mixture of **15** and **16** (Scheme 3). Spectral data of these cyclohexanetetrols were completely in agreement with those reported in the literature⁷.




Scheme 3

CONDURITOLS

Conduritols are 1,2,3,4-cyclohexenetetrol isomers. Theoretically, six conduritol diastereoisomers are possible. To avoid ambiguity, the diastereoisomers have been called with letters A,B,C,D,E,F¹. In the nature the occurrence of only two conduritols, namely Conduritol-A and Conduritol-B has been established.




Conduritol-A: In 1908 Kübler¹¹ isolated from the bark of the vine *Marsdenia Condurango* the first known cyclohexenetetrol which was named as Conduritol-A. The first successful and non-stereospecific synthesis of conduritol-A was carried out by Nakajima et al.¹²



Our synthetic strategy is based on the introduction of two oxygen functionalities at the C₂ and C₃ positions by KMnO₄-oxidation and the other two oxygen functionalities at the C₁ and C₄ positions by photooxygenation¹³. The key compound **17** in the synthesis of conduritol-A was synthesized as described on Scheme 4. Photooxygenation of **17** at room temperature afforded **18** in a yield of 95%. The exact configuration of **18** was determined at the final step.

Selective reduction of the peroxide linkage was performed with thiourea under very mild condition to give **19** in 80%. Since only the oxygen-oxygen bond breaks in this reaction, it preserves the configuration. Deketalisation of **19** was carried out in acidified methanol solution in quantitative yield. The spectroscopic properties of **20** compared well with those of the previously reported conduritol-A¹⁴ **20**.


Conduritol-F and Conduritol-B: In 1962, Plouvier¹⁵ discovered a new optically active cyclitol-isomer from *Crysanthemum Leucanthemitol* which was isomer with conduritol-A and he named this new conduritol isomer as L-Leucanthemitol (Conduritol-F). Conduritol-F can be detected at least in traces in almost green plants. We have developed a new and stereo specific synthesis for conduritol-F by two different approaches¹⁶



For the first approach we used the known trans-benzene diacetate **22**¹⁷ (Scheme 5). Photooxygenation of trans-benzene diacetate afforded the endoperoxide **23** in 56% yield. At this step we introduced all possible oxygen functionalities in correct configuration as desired in conduritol-F. Selective reduction of the peroxide linkage with thiourea followed by deacetylation of **23** was carried out with ammonia in methanol to give conduritol-F **24** which was identical with those reported in the literature.


In the second approach, we started from the endoperoxide **5** which has been synthesized by photooxygenation of the system benzene oxide-oxepine. Reduction of the peroxide-linkage in **5** with thiourea and acid-catalyzed ring opening reaction of **25** afforded conduritol-F **24** which was identical with the compound obtained from the first sequence.

In another reaction we converted epoxy diol **25** into the corresponding epoxy diacetate **26**. Epoxy diacetate was submitted to acid-catalyzed ring-opening reaction in acetic anhydride. Analysis of the reaction



mixture has revealed that the product was consisting from two isomers, conduritol-F **24** and conduritol B **28** in a ratio of 2:1. The formation of conduritol-B has a likely explanation on the basis of the involving of the neighboring acetoxy group on the course of the epoxide ring-opening as shown on scheme 6.


Conduritol-C and Conduritol-E: The first synthesis of conduritol-C was carried out by McCasland and Reeves¹⁸ from epiinositol. Yurev and Zefirov^{19a} have described a short synthetic way to conduritol-C starting from cycloaddition product of furan and ethylene carbonate. In this paper we describe two different synthetic ways leading to the synthesis of conduritol-C and conduritol-E^{19b}.



cis-Hydroxylation of the known compound **30**²⁰ led to **31** as the sole isomer. The tetraacetate **32** gave upon reaction with zinc-DMSO the unsaturated tetraacetate from which the free tetrol conduritol-C **33** was obtained by ammonolysis (Scheme 7).

The first synthesis of conduritol-E was described by Nakajima et al.¹² in 1957 which was not stereospecific and not useful. A stereospecific synthesis for conduritol-E was described by Angyal and Gilham²¹ and later again by Nakajima²². We used by our synthesis **34**, whose oxidation with *m*-chloroperbenzoic acid (*m*-CPBA) gave the epoxy compound **35**. In the trans epoxide ring-opening by addition of water, in a classic substitution mechanism, **35** should give two different opening products. However, we obtained only **36** which is the precursor of the conduritol-E. For further reaction we converted the tetrol into the readily obtained di-O-isopropylidene derivative **37**. This cyclic ketal reacted smoothly with zinc-DMSO and gave conduritol-E **38**.

Aminoconduritols; Conduramine F₄ : Aminoconduritols show also interesting inhibitor activity for some glycosidases. Synthesis of some amino conduritols has been reviewed²³. The synthetic methodology developed by us opened up entry to the synthesis of various aminocyclitols. We describe the synthesis of conduramin F₄ **41** which has the same configuration as conduritol-F.




The key compound was the endoperoxide **18**. Triethylphosphite reduction of the endoperoxide **18** was resulted in the formation of epoxy ketal **39** with definitive configuration. Acidic hydrolysis of this

compound gave conduritol-F as the sole product. Nucleophilic substitution with NH_3 followed by acidic hydrolysis afforded conduramin F₄ **41** as the sole product (Scheme 8).


Conduritol Analogues

In this section we want to discuss the synthesis of compounds derived from bicyclooctatriene which have similar structures to conduritols. Because of biological activity of the conduritol and aminoconduritol derivatives, we undertook the synthesis of the isomeric tetrols starting from bicyclooctatriene in order to see whether this kind of molecules show biological activity or not. Bicyclic endoperoxide **43**²⁴ synthesized by photooxygenation of the corresponding diene (Scheme 9) was submitted to thiourea reduction. The formed diol was converted to the corresponding diacetate for further characterization. Epoxidation of **44** resulted in the formation of only one isomer **45** whose exact configuration could not be determined. Zn-elimination followed by acidic hydrolysis gave the unsaturated cyclitol **48** in high yield. The orientation of the four hydroxyl groups in **48** is completely in agreement with conduritol-F. ¹H- and ¹³C-NMR spectral studies indicates also the asymmetric structure.




After successful synthesis of **48** we turned our attention to the synthesis of other conduritol analogues. Dibromo diacetate **44** served as the starting material. *cis*-Hydroxylation was carried out with KMnO_4 . The formed compound was characterized as the tetraacetate. The tetraacetate gave upon reaction with zinc-acetic acid unsaturated tetra acetate from which the free tetrol **47** was obtained by ammonolysis (Scheme 9). The exact configuration and the symmetry in the molecule was determined by ¹H- and ¹³C-NMR spectral measurements. Finally, the X-ray analysis of the corresponding tetraacetate indicated to our surprise that the formed compound has conduritol-D structure not the expected conduritol-A.

Cyclohexanepentols: Quercitols




The first known cyclohexanepentol was a dextrorotatory cyclitol obtained from the acorns of *Quercus* species (oaks) hence the name (+) quercitol. The structure and configuration were elucidated²⁶. The synthesis of quercitol is reported in the literature. We describe a short, efficient and stereospecific synthesis for racemic quercitol (Scheme 10). Tetraphenylporphine-sensitized photooxygenation of 1,4-cyclohexadiene gave the endoperoxide **50**. We assume that singlet oxygen undergoes an ene-reaction to give **49**. The conjugated diene unit in **49** can be easily trapped by singlet oxygen²⁷. Reduction of the peroxide linkage in **50** with thiourea followed by KMnO_4 oxidation provided racemic *proto*-quercitol **52** in high yield. A second route²⁸ for the synthesis of **52** based on photooxygenation of **53** to give **54**. Reduction of the peroxide with sodium borohydride and followed by trans opening of the epoxide ring resulted in the formation of the key compound **51**.




Application of the ene reaction of singlet oxygen to ketal **55** formed the hydroperoxide **56** whose configuration was established by NOE-measurements²⁹. KMnO_4 oxidation of the double bond in **56** formed the correct configuration of *talo*-quercitol **58**. Suitable hydrolysis of the acetates and ketal ring gave **58**.

Miscellaneous



More recently, we succeeded in the synthesis of some interesting bicyclic endoperoxides **59-63**³⁰ which contain heteroaromatic ring. The addition of 1,2,4,5-tetraazin derivatives to the appropriate unsaturated bicyclic endoperoxides followed by nitrogen extrusion gave dihydro derivative **65**. Aromatization proceeded smoothly where the peroxide linkage was remained. Chemistry and application of these compounds to the synthesis of some natural products are under investigation.



Acknowledgement: The author are indebted to the Atatürk University (Grant No. AFP 1989/31, 1991/6) and Department of Chemistry and Scientific and Technical Research Council of Turkey (DPT-TBAG-6, TBAG-1063,) for providing the necessary funds to enable our recent exploration into the chemistry of cyclitols. Furthermore, I am grateful to my students whose names are given in the references.

References

- 1) For a recent review of conduritols see: M. Balcı, Y. Sütbeyaz, H. Seçen, *Tetrahedron*, **46**, 3715 (1990) and references cited therein.
- 2) a) S. Atsumi, K. Umezawa, H. Iinuma, H. Naganawa, H. Nakamura, Y. Iitaka, T. Takeuchi, *J. Antibiot.* **43**, 49 (1989). b) G. Legler, E. Bause, *Carbohydr. Res.* **28**, 45 (1973). c) G. Legler, W. Lotz.,; *Z. Physiol. Chem.* **354**, 243 (1973).
- 3) N. Chida, K. Yamada, S. Ogawa, *S. J. Chem Soc. Chem. Commun.* 588 (1991).
- 4) Z. Zeying, Z. Mingzhe, *Jiegou Huaxue, J. Struct. Chem.* **6**, 128 (1987).
- 5) G. E. McCasland, E.C. Horswill., *J. Am. Chem. Soc.* **75**, 4020 (1953).
- 6) a) H. Z. Sable, K.A. Powel, H. Katchian, C. B. Niewoehner, S. B. Kadlec, *Tetrahedron*, **26**, 1509 (1970). b) J. A. Franks, B. Tolbert, R. Steyn, H. Z. Sable, *J. Org. Chem.* **30**, 1440 (1995). c) R. Steyn, H. Z. Sable, H.Z. *Tetrahedron*, **25**, 3579 (1969).
- 7) N. Akbulut, M. Balcı, *J. Org. Chem.* **53**, 3338 (1988).
- 8) M. Balcı, *Chem. Rev.* **81**, 91 (1981).
- 9) C. Foster, G. A. Berchtold, *J. Org. Chem.* **40**, 3743 (1975).
- 10) T. Posternak, H. Friedli, *Helv. Chim. Acta.* **36**, 251 (1953)
- 11) a) K. Kübler, *Arch. Pharm. Ber. Stsch. Pharm.* **246**, 620 (1908). b) see also: P. E. Manni, J. E. Sinsheimer, *J. Pharm. Sci.* **54**, 1541 (1965).
- 12) a) M. Nakajima, I. Tomida, S. Takei, *Chem. Ber.* **90**, 246 (1957). b) S. Knapp, R. M. Ornaf, K. E. Rodrigues, *J. Am. Chem. Soc.* **105**, 5494 (1983).
- 13) Y. Sütbeyaz, H. Seçen, M. Balcı, *M. J. Chem. Soc. Chem. Commun.* 1330 (1988).
- 14) R. J. Abraham, H. Gottschalck, H. Paulsen, W. A. Thomas, *J. Chem. Soc.* **1** 6268 (1965)
- 15) V. Plouvier, *C.r. hebdomad. Sc' Acad. Sci. Paris*, 360 (1962).
- 16) H. Seçen, Y. Sütbeyaz, M. Balcı, *Tetrahedron Lett.* **31**, 1323 (1990).
- 17) M. R. Demuth, P. E. Garret, J. D. White, *J. Am. Chem. Soc.* **98**, 634 (1976).
- 18) G. E. McCasland, J. M. Reeves, *J. Am. Chem. Soc.* **77**, 1812 (1955).
- 19) a) Y. K. Yurev, K. Zefirov, *Zh. Obshchei Khim.* **31**, 685 (1961). b) H. Seçen, A. Maraş, Y. Sütbeyaz, M. Balcı, *Synth. Commun.* **22**, 2613 (1992).
- 20) H. J. Altenbach, H. Stegelmeier, E. Vogel, *E. Tetrahedron Lett.* 3333 (1978).
- 21) S. J. Angyal, P. T. Gilham, *J. Chem. Soc.* 375 (1955).
- 22) M. Nakajima, I. Tomida, S. Takei, *Chem. Ber.* **92**, 163 (1959).
- 23) T. Posternak, *The Cyclitols* Hermann, Paris, (1965).
- 24) a) M. Oda, Y. Kayama, Y. Kitahara, *Y. Tetrahedron Lett.* 2019 (1974).
- 25) a) A. C. Cope, N. A. Nelson, D. S. Smith, *J. Am. Chem. Soc.* **76**, 1100 (1954). b) R. M. Carlson, R. K. Hill, *Org. Synth.* **50**, 24 (1970).
- 26) G. E. McCasland, M. O. Naumann, L. J. Durham, *J. Org. Chem.* **33**, 4220 (1968).
- 27) a) H. Seçen, E. Salamcı, Y. Sütbeyaz, *Synlett*, 609 (1993).
b) For similar reaction see: I. Saito, K. Tamoto, A. Katsumura, *Chem. Lett.* 127 (1978).
- 28) Gültekin, S.; Salamcı, E.; Sütbeyaz, Y.; Balcı, M. unpublished results.
- 29) A. Maraş, M. Balcı, unpublished results.
- 30) N. Saraçoğlu, A. Menzek, M. Balcı, unpublished results.