

11-27-2006

Synthesis of Indolizines and Benzoindolizines by Annulation of Donor-Acceptor Cyclopropanes with Electron-Deficient Pyridines and Quinolines

Nicholas A. Morra

Christian L. Morales

Barbora Bajtos

Xin Wang

Hyosook Jang

See next page for additional authors

Follow this and additional works at: <https://ir.lib.uwo.ca/chempub>

 Part of the [Chemistry Commons](#)

Citation of this paper:

Morra, Nicholas A.; Morales, Christian L.; Bajtos, Barbora; Wang, Xin; Jang, Hyosook; Wang, Jian; and Pagenkopf, Brian, "Synthesis of Indolizines and Benzoindolizines by Annulation of Donor-Acceptor Cyclopropanes with Electron-Deficient Pyridines and Quinolines" (2006). *Chemistry Publications*. 60.
<https://ir.lib.uwo.ca/chempub/60>

Authors

Nicholas A. Morra, Christian L. Morales, Barbora Bajtos, Xin Wang, Hyosook Jang, Jian Wang, and Brian Pagenkopf


Synthesis of Indolizines and Benzoindolizines by Annulation of Donor-Acceptor Cyclopropanes with Electron Deficient Pyridines and Quinolines

Nicholas A. Morra, Christian L. Morales, Barbora Bajtos, Xin Wang, Hyosook Jang, Jian Wang, Ming Yu and Brian L. Pagenkopf*

The University of Western Ontario, Department of Chemistry, 1151 Richmond Street
London, Ontario, N6A 5B7 Canada
Fax: (+1) 519 661-3022
bpagenko@uwo.ca

Supporting Information for this article is available at <http://asc.wiley-vch.de/home/>


Abstract



The formal [3+2] dipolar cycloaddition (or annulation) of donor-acceptor cyclopropanoate esters with pyridines and 5-nitroquinoline is reported. Electron deficient pyridine dipolarophiles (R = CN, CO₂Et, COMe) participate in the annulation whereas electron rich species do not. The product 2,3-dihydroindolizines undergo rapid autooxidation, and the X-ray structures for two of the aromatic products are reported.

Donor-acceptor alkoxy substituted cyclopropanoate esters are important synthetic intermediates that when suitably activated by Lewis acids can undergo fruitful annulation or [3+2] dipolar cycloaddition processes with electron rich aldehydes, imines and related pi systems.^[1] We recently found success with nitrile dipolarophiles in such reactions,^[2] and subsequently developed a general pyrrole synthesis.^[3,4] While investigating the scope of the pyrrole synthesis with heteroaromatic nitriles^[4] we examined the reaction between 4-cyano pyridine **1** and cyclopropane **2a** (Scheme 1) under the standard reaction conditions (Me₃SiOTf, MeNO₂ solvent, 0 °C → rt).^[3] However, the anticipated pyrrole **3** was not observed, and instead the intensely purple 2,3-dihydroindolizine **5a** and the pale yellow indolizine **6a** were obtained by flash chromatography on silica gel in 52% and 9% isolated yield, respectively. The product **5a** is apparently tautomerized from the initial alkene **4**, and the fully aromatic indolizine **6a** is formed by autooxidation during workup and isolation.

Scheme 1



This report describes our results from an exploration of this new synthetic pathway for preparing indolizines, the skeleton of which appears in biologically active natural products,^[5] medicinal chemistry,^[6] dyes^[7] and agrochemicals.^[8] Many creative methods have been devised for their synthesis.^[9,10,11,12] Annulations between pyridines and cyclopropanones are known,^[11,12] as are formal [3+2] dipolar cycloadditions of pyridines with certain 1,3-dipoles, such as nitrile ylides^[13] and nitrile imines.^[14]

The reaction of more electron rich pyridines, including pyridine itself and 4-methoxypyridine, with cyclopropane **2a** under the above conditions gave complex reaction mixtures. A trace amount of product was found from reaction with pyridine (3%), and nothing identifiable from 4-methoxypyridine. The original Lewis acid and solvent system for the reaction appear somewhat incompatible with pyridines. For example, reaction of Me₃SiOTf and pyridine with the nitromethane solvent might lead to formation of the silylnitronate, or the interaction of the nucleophilic pyridine and electrophilic triflate might sequester them as their pyridinium salt. Therefore, efforts to optimize the reaction were undertaken that included adjusting stoichiometry and surveying various solvents (PhNO₂, CH₂Cl₂, CCl₄, EtOAc, PhMe) and Lewis acids (Yb(OTf)₃,^[15] TBSOTf, Me₃SiOMs, ⁱPr₃SiOTf, BF₃•OEt₂, ZnCl₂). Additionally, different workup procedures that were tested ranged from strongly acid to basic and both aqueous and anhydrous. Unfortunately, none of these modifications proved generally more useful than the original combination of Me₃SiOTf in MeNO₂.

The reaction scope was then evaluated by combining 4-cyanopyridine with additional cyclopropanes offering different substitution patterns (Figure 1), including those with: the R¹ and R² substituents tied back as a fused cyclohexane ring (**2b**), a phenyl ring at the alkoxy carbon (**2c**), a methyl at the alkoxy carbon (**2d**), no substituents (**2e**) and a fused pyranose structure (**2f**).¹⁶ The fused cyclohexane proved compatible and gave the annulation highly air sensitive 2,3-dihydroindolizine along with some aromatic product (Table 1, entry b). Therefore to facilitate isolation and characterization the annulation products were immediately subjected to oxidizing conditions (MnO₂ or DDQ),¹⁷ and the yields reported in Table 1 are for the two step sequence. The substrate bearing a phenyl group at the alkoxy carbon provided an additional class of compatible cyclopropanes (entry c), and the oxidized annulation product was obtained in a modest 35% yield (over two steps). With a methyl in the same position the yield dropped to 29% (entry d). No identifiable products could be isolated from complex decomposition residues after reaction with cyclopropanes **2e** and **2f**.

Figure 1. Representative Cyclopropanes





Table 1. Annulation and oxidation of DA cyclopropanes and pyridines




Entry	Cyclopropane	Pyridine	Product	Yield ^a
a		4-CN		61%
b		4-CN		43%
c		4-CN		35%
d		4-CN		29%
e		4-CO2Et		57%
f		4-CO2Et		46%
g		4-CO2Et		42%
h		4-CO2Et		18%
i		4-COMe		8%
j		4-COMe		15%
k		4-COMe		33%
l		3-CO2Me		10%
m		3-CO2Me		12% ^b

^a Isolated yield over two steps. ^b PhNO₂ used as solvent.

Other electron deficient pyridines were examined, and ethyl isonicotinate performed comparably well with members of the representative set of cyclopropanes bearing untethered aliphatic substituents (entry e, 57%), a fused cyclohexyl ring (entry f, 46%), an aromatic substituent (entry g, 42%) and a methyl group (entry h, 18%). A 4-acylpyridine participates, but with test cyclopropanes **2a** and **2b** the yield was markedly lower at 8% and 15% (entries i and j). The 33% yield from reaction with aryl substituted cyclopropane **2c** (entry k) is, however, comparable to results with the other pyridines.

Reactions with non-symmetric methyl nicotinate took about five hours longer for complete consumption of cyclopropane (TLC), and yields were disappointing (entries l and m). Only the regioisomer shown with the pyridine substituent distal to the newly formed carbon-carbon bond was detected in these reactions. In the case of entry m a 6% yield was improved to 12% by replacing the MeNO₂ solvent with PhNO₂, and the oxidized product gave crystals of quality suitable for X-ray analysis (Figure 2).^[18]

Figure 2. ORTEP of **6m**.^[18]



Among all of the pyridines examined the best yields were obtained with cyclopropane **2a**, whereas only intractable mixtures were encountered with cyclopropanes **2e** and **2f**. One structural feature that appears necessary from this set of examples is the requirement that the alkoxy substituent be bonded to a quaternary carbon. It is noteworthy that the cyclopropanes that fail here excel in annulation reactions with nitriles.^[3]

Quinolines were evaluated as annulation partners and quinoline, like the electron rich pyridines, gave complex reaction mixtures under the standard conditions.^[12,19] Therefore commercially available 5-nitroquinoline was chosen as an electron deficient substrate, and with cyclopropane **2b** a more satisfying 85% isolated yield of the oxidized product **8** was obtained (Scheme 2). The structure was confirmed by single crystal X-ray analysis (Figure 3).

Scheme 2







Figure 3. ORTEP of **8**.^[18]


The improved reaction efficiency with 5-nitroquinoline fortunately extends to those cyclopropanes that proved incompatible in the pyridine series. For example, cyclopropanes **2e** and **2f** both failed to give any identifiable products with the pyridines in Table 1, whereas they were successful here. Specifically, reaction with **2e** furnished the aminal **9** and the non-oxidized elimination product **10** in a 1.1 to 1 ratio and 79% combined isolated yield (Scheme 3). The aminal **9** was cleanly converted to the dihydrobenzoindolizine **10** by treatment with dilute aqueous HCl. Reaction with cyclopropane **2f** was equally successful, and afforded the aminal **11** and the elimination product **12** in a 1 to 1.7 ratio and 83% yield (Scheme 4). No aminal products were detected during work with the pyridines.

Scheme 3**Scheme 4**

It is clear from these results that reaction efficiency increases with electron deficient pyridines, a trend that is opposite to that observed with nitriles.^[2-4] A mechanistic hypothesis that is consistent with these observations is provided in Scheme 5. Activation of the cyclopropane by Me_3SiOTf generates an electrophilic oxocarbenium ion (or equivalent) that, in a stepwise mechanism, is intercepted by a nucleophilic pyridine (step 1) to give **13**. Electron donating groups (X) will improve nucleophilicity and facilitate step 1. However, the same group's stabilization of **13'** will critically render the pyridinium ion less susceptible to attack by

the ester enolate (step 2), and the reaction will fail. This mechanistic speculation implies that pyridines of a particular electronic disposition (i.e., an electron deficient X) are required to balance the opposing demands of the two bond forming steps. Quizzically, the observed electronic trends may be an unfortunate artifact resulting from complications due to the incompatibility of the Me_3SiOTf and other reagents as discussed in an introductory paragraph. Furthermore, a dipolar cycloaddition mechanism is plausible.

Scheme 5



In summary, a new operationally simple annulation between DA-cyclopropanes with pyridines and quinolines for the synthesis of indolizines and benzoindolizines has been described. Electron poor pyridines and quinolines are superior substrates for this reaction at its current state of development. Despite some limitations in substrate scope, this methodology provides convenient new pathways that access a variety of indolizine scaffolds. It is hoped that future catalyst systems will result in greater substrate diversity and improved yields.

Experimental Section

General annulation procedure. A degassed solution of cyclopropane (1.0 mmol) and pyridine (2.0 mmol) in MeNO_2 (0.6 mL) at 0 °C was treated with Me_3SiOTf (1.1 mmol) and the resulting mixture was allowed to warm to room temperature. After two hours the solution was diluted with EtOAc (8 mL) and then poured into a vigorously stirred solution of saturated aqueous NaHCO_3 (40 mL). The heterogeneous mixture was extracted with EtOAc (3 x 10 mL) and the combined organic layers were washed with brine and dried (Na_2SO_4). After filtration through a thin pad of celite the solution was divided into two equal portions. One was used for the (attempted) isolation and characterization of **5**. The other portion was concentrated under reduced pressure, dissolved in CH_2Cl_2 (10 mL) and treated with a mixture of powdered molecular sieves (4 Å, 1.2 g) and MnO_2 (5 mmol, 0.43 g, 10 equivalents). The reaction was heated to reflux and monitored periodically by TLC (ca. 3 h). The cooled reaction mixture was filtered through a pad of silica gel and concentrated under reduced pressure. Purification by flash chromatography on silica gel using EtOAc-hexanes for elution provided the indolizine.

Key words: annulation, autooxidation, donor-acceptor systems, cyclopropanes, cycloadditions, zwitterions.

Acknowledgements. We thank the donors of the American Chemical Society Petroleum Research fund, Johnson and Johnson, NSERC and the University of Western Ontario for financial assistance. We thank Vincent Lynch for determination of the X-ray structures.

References

- [1] Donor-acceptor cyclopropane reviews: (a) M. Yu; B. L. Pagenkopf, *Tetrahedron* **2005**, 61, 321-347. (b) F. Gnad; O. Reiser, *Chem. Rev.* **2003**, 103, 1603-1624. (c) H.-U. Reissig; R. Zimmer, *Chem. Rev.* **2003**, 103, 1151-1196. (d) A. de Meijere; S. I. Kozhushkov; L. P. Hadjiarapoglou, *Top. Curr. Chem.* **2000**, 207, 149-227. (e) A. de Meijere; S. I. Kozhushkov; A. F. Khlebnikow, *Top. Curr. Chem.* **2000**, 207, 89-147. (f) J. Salaün, *Top. Curr. Chem.* **2000**, 207, 1-67. (g) G. S. Cousins; J. O. Hoberg *Chem. Soc. Rev.* **2000**, 29, 165-174. (h) Carbocyclic Three- and Four-Membered Ring Compounds. In *Houben-Weyl, Methods of Organic Chemistry*, 4th ed.; A. de Meijere, Ed.; Thieme Verlag: Stuttgart, 1997; Vol. E17a, E17b, E17c, E17d. (j) H. N. C. Wong; M.-Y. Hon; C.-W. Tse; Y.-C. Yip; J. Tanko; T. Hudlicky, *Chem. Rev.* **1989**, 89, 165-198. (k) H.-U. Reissig, *Top. Curr. Chem.* **1988**, 144, 73-135. (l) H.-U. Reissig In *The Chemistry of the Cyclopropyl Group*; Z. Rappoport, Ed.; John Wiley & Sons: Chichester, 1987; Chapter 8, pp 375-443. (m) L. A. Paquette, *Chem. Rev.* **1986**, 86, 733-750. (n) E. Wenkert, *Acc. Chem. Res.* **1980**, 13, 27-31. (o) S. Danishefsky, *Acc. Chem. Res.* **1979**, 12, 66-72.
- [2] M. Yu; B. L. Pagenkopf, *J. Am. Chem. Soc.* **2003**, 125, 8122-8123.
- [3] M. Yu; B. L. Pagenkopf, *Org. Lett.* **2003**, 5, 5099-5101.
- [4] M. Yu; G. D. Pantos; J. L. Sessler and B. L. Pagenkopf, *Org. Lett.* **2004**, 6, 1057-1059.
- [5] J. P. Michael, *Nat. Prod. Rep.* **2005**, 603-626.
- [6] (a) T. Weide; L. Arve; H. Prinz; H. Waldmann; H. Kessler, *Bioorg. Med. Chem. Lett.* **2006**, 16, 59-63. (b) P. Sonnet, *et al. Bioorg. Med. Chem. Lett.* **2000**, 8, 945-955. (c) S. Hagishita *et al. J. Med. Chem.* **1996**, 39, 3636-3658. (d) J. Gubin, *et al. Boorg. Med. Chem.* **1993**, 36, 1425-1433.
- [7] C. H. Weidner; D. H. Wadsworth; S. L. Bender; D. J. Beltman, *J. Org. Chem.* **1989**, 54, 3660-3664.
- [8] S. C. Smith; E. D. Clarke; S. M. Ridley; D. Bartlett; D. T. Greenhow; H. Glithro; A. Y. Klong; G. Mitchell; G. W. Mullier *Pest. Man. Sci.* **2005**, 61, 16-24.
- [9] Recent synthetic strategies: (a) J. Kaloko, Jr.; A. Hayford *Org. Lett.* **2005**, 7, 4305-4308. (b) C.-H. Park; V. Ryabova; I. V. Seregin; A. W. Sromek; V. Gevorgyan *Org. Lett.* **2004**, 6, 1159-1162. (c) A. R. Katritzky; G. Qiu; B. Yang; H.-Y. He, *J. Org. Chem.* **1999**, 64, 7618-7621.
- [10] Review of indolizine synthetic methods, see: (a) T. Uchida; K. Matsumoto *Synthesis* **1976**, 209-236. (b) W. Flitsch In *Comprehensive Heterocyclic Chemistry*; A. R. Katritzky, C. W. Rees, Eds.; Pergamon: Oxford, 1984; Vol.4, p.443.
- [11] C. H. Weidner; F. M. Michaels; D. J. Beltman; C. J. Montgomery; D. H. Wadsworth; B. T. Briggs; M. L. Picone, *J. Org. Chem.* **1991**, 56, 5594-5598.
- [12] (a) J. W. Lown; K. Matsumoto, *Can. J. Chem.* **1971**, 49, 1165-1175. (b) C. Weber; H. Durr; R. Bonneau, *Synthesis* **1999**, 1005-1012.
- [13] P. Caramella; A. G. Invernizzi; E. Pastormerlo; P. Quadrelli; A. Corsaro, *Heterocycles* **1995**, 40, 515-520.

- [14] (a) L. Grubert; W. Jugelt; H. J. Bress; H. Koppel; U. Strietzel; A. Dombrowski, *Liebigs Ann. Chem.* **1992**, 885-894; (b) L. Grubert; M. Patzel; W. Jugelt; B. Riemer; J. Liebscher, *Liebigs Ann. Chem.* **1994**, 1005-1011;
- [15] C. A. Carson; M. A. Kerr, *J. Org. Chem.* **2005**, *70*, 8242-8244.
- [16] The cyclopropanes were used as a mixture of diastereomers. No difference in reactivity was observed from deployment of single diastereomers
- [17] T. Chandra; S. Zou.; K. L. Brown, *Tetrahedron Lett.* **2004**, *45*, 7783-7786. DDQ was used to oxidize reaction products from cyclopropane **2e**.
- [18] CCDC 611519 (**6m**) and CCDC 611520 (**8**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.
- [19] E. Bacu.; D. Samson-Belei; G. Nowogrocki; A. Couture; P. Grandclaoudon, *Org. Biomol. Chem.* **2003**, *1*, 2377-2382.