

Synthesis of quinoline/naphthalene-containing azaphenothiazines and their potent in vitro antioxidant properties

Małgorzata Jeleń · Eugenia I. Bavavea · Maria Pappa ·
Angeliki P. Kourounakis · Beata Morak-Młodawska ·
Krystian Pluta

Received: 11 June 2014 / Accepted: 27 August 2014 / Published online: 12 September 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract New tetracyclic and pentacyclic azaphenothiazines containing one or two quinoline rings instead of benzene rings were obtained in the original reactions of isomeric diquinodithiins, dichlorodiquinoliny sulfides, and disulfide with aromatic amines. The type of ring fusion in the azaphenothiazine system was concluded from the ^1H NMR spectra. The obtained azaphenothiazines were evaluated in vitro for their antioxidant activity on rat hepatic microsomal membranes for protection of non-enzymatic lipid peroxidation promoted by the Fe^{2+} /ascorbic acid redox system. Most compounds exhibited a very significant antioxidant activity with IC_{50} values between 1 and 23 μM . The degree of antioxidant activity depends on the lipophilicity and molecular size as well as the (non)substitution of the thiazine nitrogen atom and type of ring system fusion. It is the first time to our knowledge that azaphenothiazines are shown to exhibit such potent antioxidant activity.

Keywords Phenothiazine derivatives ·
NH-azaphenothiazines · Quinonaphthothiazines ·
Diquinothiazines · Lipid peroxidation · Lipophilicity

Introduction

Phenothiazines are an important class of drugs exhibiting antipsychotic, antihistaminic, antitussive, and anti-emetic activities (Gupta and Kumar, 1988). The most significant modifications of the phenothiazine structure are the introduction of new pharmacophoric substituents at the thiazine nitrogen atom and the substitution of the benzene rings with other homoaromatic or heteroaromatic ones. Recently studied phenothiazines exhibit promising antibacterial, antifungal, anticancer, antiviral, anti-inflammatory, anti-malarial, antifilarial, trypanocidal, anticonvulsant, analgesic, immunosuppressive, and multidrug resistance reversal properties (Aaron *et al.*, 2009; Dasgupta *et al.*, 2008; Motohashi *et al.*, 2006; Pluta *et al.*, 2011). In our study of new azaphenothiazines, we elaborated the synthesis of new types of phenothiazines containing the heterocyclic rings of pyridine or quinoline. Some of those azaphenothiazines exhibited promising immunosuppressive and anticancer activities against cell lines of ten types of human cancer in vitro: leukemia, non-small cell lung cancer, melanoma, as well as colon, CNS, ovarian, renal, prostate, breast, and skin cancer (Jeleń *et al.*, 2013; Pluta *et al.*, 2010; Zimecki *et al.*, 2009).

Free radicals, generated in many redox processes, may induce oxidative damage of proteins, lipids, and DNA. They affect living cells and mediate the pathogenesis of many chronic diseases, such as atherosclerosis, Parkinson's and Alzheimer's diseases, stroke, and arthritis, acting by various mechanisms. A recent trend in the field of antioxidant development focuses on multipotent antioxidant agents that not only can prevent biological substrates from radical induced oxidative damage but also possess additional pharmacological properties (Zhang *et al.*, 2006). The study of antioxidant activity among N-heterocycles has

M. Jeleń · B. Morak-Młodawska · K. Pluta (✉)
Department of Organic Chemistry, School of Pharmacy with the
Division of Laboratory Medicine, The Medical University of
Silesia, Jagiellońska 4, 41-200 Sosnowiec, Poland
e-mail: pluta@sum.edu.pl

E. I. Bavavea · M. Pappa · A. P. Kourounakis
Department of Medicinal Chemistry, Faculty of Pharmacy,
University of Athens, 15771 Athens, Greece

attracted attention. One such heterocyclic structural scaffold is the 1,4-thiazine ring present in the multi-target phenothiazines. Therefore, recent reports on promising antioxidant compounds deal with classical and new phenothiazines (Asghar *et al.*, 2012; Borges *et al.*, 2010; Liu *et al.*, 2009; Naik *et al.*, 2012;) and their derivatives, benzothiazines (Matralis *et al.*, 2011), and azaphenothiazines (Kumar *et al.*, 2010; Morak-Młodawska *et al.*, 2010).

Our previous work (Morak-Młodawska *et al.*, 2010) revealed that tricyclic azaphenothiazines being dipyrrothiazines have a variable degree of antioxidant activity depending on substitution at the thiazine nitrogen atom, with the unsubstituted compound being the most active. In this study, we obtained eleven tetracyclic and pentacyclic (linearly and angularly fused) azaphenothiazines containing one or two quinoline rings instead of the benzene rings and determined their antioxidant properties to find an influence of the number of rings, their type of fusion, and their substituents.

Materials and methods

General techniques

Melting points were determined in open capillary tubes on a Boetius melting point apparatus and were uncorrected. The ^1H NMR spectra were recorded on a Bruker Fourier 300 and a Bruker DRX spectrometer at 500 MHz in CDCl_3 and $\text{DMSO}-d_6$ with tetramethylsilane as the internal standard. The ^{13}C NMR spectra were recorded at 75 MHz. Electron impact (EI MS) mass spectra were run on a Finnigan MAT 95 spectrometer at 70 eV. The thin-layer chromatography was performed on aluminum oxide 60 F₂₅₄ neutral (type E, Merck 1.05581) with CH_2Cl_2 and on silica gel 60 F₂₅₄ (Merck 1.05735) with CHCl_3 -EtOH (10:1 v/v) as eluents.

Synthesis of substrates **1**, **2**, **7**, **8**, **10**, and **11**

The substrates for the title compounds, i.e., diquinodithiins **1**, **7**, **10**, sulfides **8**, **11**, and disulfide **2**, were obtained as described previously (Nowak *et al.*, 2002, 2003, 2007; Pluta, 1994).

Quino[3,2-b]benzo[1,4]thiazines (**3a–c**)

From diquino-1,4-dithiin **1**

A mixture of diquino-1,4-dithiin **1** (0.16 g, 0.5 mmol) and hydrochloride of aniline, or *p*-chloroaniline or *p*-methoxyaniline (2.5 mmol) was finely powdered together and then heated on an oil bath at 200–205 °C for 4 h and

after cooling water was added (10 ml) and the insoluble solid was filtered off. The filtrate was alkalized with 5 % aqueous sodium hydroxide to pH 10, and the resulting solid was filtered off and washed with water. The combined solids were purified by column chromatography (silica gel, CHCl_3) to give quinobenzothiazines **3a–c**.

6H-Quinobenzothiazine (3a) 0.06 g (24 %), yellow, mp 169–170 °C (mp 169–170 °C, Jeleń and Pluta, 2009). ^1H NMR (CDCl_3) δ : 6.62 (m, 1H, H-7), 6.87 (m, 1H, H-9), 7.03 (m, 2H, H-8, H-10), 7.26 (t, 1H, H-2), 7.47 (m, 2H, H-1, H-3), 7.53 (s, 1H, H-12), 7.56 (d, 1H, H-4). ^{13}C NMR (CDCl_3) δ : 115.57 (C-7), 116.49 and 116.69 (C-10a, C-11a), 122.95 (C-9), 124.19 (C-2), 125.86 (C-10), 126.04 and 126.45 (C-1, C-8), 126.56 (C-12a), 127.57 (C-4), 129.52 (C-3), 131.69 (C-12), 138.45 (C-6a), 145.40 (C-4a), 150.98 (C-5a).

6H-9-Chloroquinobenzothiazine (3b) 0.08 g (28 %), yellow, mp 224–225 °C (mp 224–225 °C, Jeleń and Pluta, 2009). ^1H NMR (CDCl_3) δ : 6.63 (d, 1H, H-7), 6.99 (s, 1H, H-10), 7.01 (d, 1H, H-8), 7.33 (t, 1H, H-2), 7.51 (d, 1H, H-1), 7.52 (t, 1H, H-3), 7.59 (d, 1H, H-4), 7.60 (s, 1H, H-12). ^{13}C NMR (CDCl_3) δ : 115.80 (C-11a), 116.71 (C-7), 118.19 (C-10a), 124.84 and 124.91 (C-8, C-10), 125.65 (C-2), 126.13 (C-12a), 126.61 (C-1), 127.59 (C-4), 128.56 (C-9), 130.31 (C-3), 132.35 (C-12), 136.29 (C-6a), 143.81 (C-4a), 150.04 (C-5a),

6H-9-Methoxyquinobenzothiazine (3c) 0.09 g (32 %), orange, mp 159–160 °C.

^1H NMR (CDCl_3) δ 3.76 (s, 3H, CH_3), 6.54 (d, 1H, H-7), 6.63 (d, 1H, H-10), 6.76 (d, 1H, H-8), 7.29 (t, 1H, H-2), 7.46 (d, 1H, H-1), 7.52 (t, 1H, H-3), 7.55 (s, 1H, H-12), 7.57 (d, 1H, H-4). ^{13}C NMR (CDCl_3) δ : 111.59 (C-10), 113.22 (C-8), 116.41 (C-11a), 116.82 (C-7), 117.39 (C-10a), 124.36 and 124.49 (C-1, C-2), 125.80 (C-12a), 126.55 (C-4), 130.10 (C-3), 130.60 (C-6a), 132.07 (C-12), 143.40 (C-4a), 150.36 (C-5a), 156.12 (C-9). EIMS m/z : 280 (M^+ , 100), 265 ($\text{M}-\text{CH}_3$, 90). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{N}_2\text{OS}$: C, 68.55; H, 4.31; N, 9.99. Found: C, 68.45; H, 4.36; N, 9.82.

From 2,2'-dichloro-3,3'-diquinoliny disulfide (**2**)

A solution of disulfide **2** (0.20 g, 0.5 mmol) and *p*-methoxyaniline (0.25 g, 2 mmol) in monomethyl ether of diethylene glycol (MEDG) (5 ml) was refluxed for 3 h. After cooling, the solution was poured into water (20 ml) and alkalized with 5 % aqueous sodium hydroxide to pH 10. The resulting solid was filtered off, washed with water, and purified by column chromatography (silica gel, CHCl_3) to give 0.18 g (64 %) of 6H-9-methoxyquinobenzothiazine (**3c**).

Quino[3,2-b]naphtho[1',2'-e][1,4]thiazine (4)

Diquinodithiin **1** (0.16 g, 0.5 mmol) was finely powdered together with 1-naphthylamine hydrochloride (0.45 g, 2.5 mmol) on an oil bath at 200–205 °C for 4 h. After cooling, the solution was poured into water (10 ml) and alkalized with 5 % aqueous sodium hydroxide to pH 10. The resulting solid was filtered off, washed with water, and purified by column chromatography (Al₂O₃, CHCl₃) to give 0.08 g (27 %) of 14*H*-quinonaphthothiazine (**4**), orange, mp 147–148 °C.

¹H NMR (CDCl₃) δ: 7.01 (d, 1H, H-6), 7.30 (t, 1H, H-10), 7.47 (m, 4H, H-3, H-4, H-5, H-9), 7.52 (t, 1H, H-2), 7.56 (s, 1H, H-8), 7.60 (t, 1H, H-11), 7.64 (d, 1H, H-12), 7.75 (d, 1H, H-1). ¹³C NMR (CDCl₃) δ: 110.98 (C-6a), 116.91 (C-7a), 118.43 (C-1), 121.89 (C-14b), 122.87 (C-6), 123.70 (C-5), 124.49 (C-10), 125.93, 126.45 and 126.83 (C-2, C-3, C-9), 126.90 (C-8a), 128.92 and 129.65 (C-4, C-12), 131.54 (C-11), 132.55 (C-4a), 133.04 (C-8), 135.07 (C-14a), 145.23 (C-12a), 150.98 (C-13a). EIMS *m/z*: 300 (M⁺, 100), 268 (M-S, 45). Anal. Calcd. for C₁₉H₁₂N₂S: C, 75.97; H, 4.03; N, 9.33. Found: C, 75.82; H, 4.07; N, 9.21.

Quino[3,2-b]naphtho[2',1'-e][1,4]thiazine (5)

Diquinodithiin **1** (0.16 g, 0.5 mmol) was finely powdered together with 2-naphthylamine hydrochloride (0.45 g, 2.5 mmol) on an oil bath at 200–205 °C for 4 h. After cooling, the solution was poured into water (10 ml) and alkalized with 5 % aqueous sodium hydroxide to pH 10. The resulting solid was filtered off, washed with water, and purified by column chromatography (Al₂O₃, CHCl₃) to give 0.12 g (40 %) of 7*H*-quinonaphthothiazine (**5**), greenish, mp 244–245 °C.

¹H NMR (CDCl₃) δ: 7.06 (d, 1H, H-6), 7.37 (t, 1H, H-11), 7.47 (t, 1H, H-3), 7.57 (m, 3H, H-2, H-10, H-12), 7.65 (d, 1H, H-5), 7.66 (d, 1H, H-4), 7.72 (s, 1H, H-13), 7.80 (m, 2H, H-9, H-1). ¹³C NMR (CDCl₃) δ: 107.94 (C-14a), 115.77 (C-13a), 116.04 (C-6), 121.32 (C-1), 123.33, 123.66 and 123.89 (C-3, C-9, C-11), 125.23 (C-12a), 125.62 (C-2), 126.36, 126.99 and 127.56 (C-4, C-5, C-12), 128.73 (C-4a), 129.22 (C-10), 129.62 (C-14b), 131.51 (C-13), 133.54 (C-6a), 142.13 (C-8a), 149.64 (C-7a). EIMS *m/z*: 300 (M⁺, 100), 268 (M-S, 50). Anal. Calcd. for C₁₉H₁₂N₂S: C, 75.97; H, 4.03; N, 9.33. Found: C, 75.88; H, 4.05; N, 9.19.

Diquino[3,2-b;6',5'-e][1,4]thiazine (6)

Diquinodithiin **1** (0.16 g, 0.5 mmol) was finely powdered together with 6-aminoquinoline hydrochloride (0.46 g, 2.5 mmol) on an oil bath at 200–205 °C for 4 h. After cooling, the solution was poured into water (10 ml) and

alkalized with 5 % aqueous sodium hydroxide to pH 10. The resulting solid was filtered off, washed with water, and purified by column chromatography (Al₂O₃, CHCl₃) to give 0.10 g (33 %) of 7*H*-diquinothiazine (**6**), brown, mp 260–261 °C.

¹H NMR (CDCl₃) δ: 7.44 (t, 1H, H-11), 7.49 (d, 1H, H-6), 7.57 (m, 2H, H-2, H-12), 7.64 (t, 1H, H-10), 7.70 (d, 1H, H-9), 7.75 (s, 1H, H-13), 8.10 (d, 1H, H-5), 8.19 (d, 1H, H-1), 8.90 (d, 1H, H-3). ¹³C NMR (CDCl₃) δ: 107.62 (C-14a), 114.59 (C-13a), 119.33 (C-6), 120.76 (C-2), 124.05 (C-11), 124.37 and 125.45 (C-12a, C-14b), 125.65 (C-12), 128.27, 129.24, 129.62 and 129.64 (C-1, C-5, C-9, C-10), 131.80 (C-13), 134.54 (C-6a), 144.53 (C-7a), 147.55 (C-3), 149.49 and 149.55 (C-4a, C-8a). EIMS *m/z*: 301 (M⁺, 100), 269 (M-S, 45). Anal. Calcd. for C₁₈H₁₁N₃S: C, 71.74; H, 3.68; N, 13.94. Found: C, 71.59; H, 3.71; N, 13.72.

Diquino[3,2-b;2',3'-e][1,4]thiazines (9)

6*H*-Diquinothiazine **9a**

This compound was obtained in the reaction of diquinodithiin **7** with acetamide (Nowak *et al.*, 2007), orange, mp > 300 °C (mp > 300 °C, Nowak *et al.*, 2007). ¹H NMR (CDCl₃) δ: 7.42 (t, 2H, H-2, H-10), 7.55 (d, 2H, H-1, H-11), 7.62 (t, 2H, H-3, H-9), 7.72 (s, 2H, H-12, H-14), 7.86 (d, 2H, H-4, H-8). ¹³C NMR (DMSO-*d*₆) δ: 124.83 (C-12a, C-13a), 127.29 (C-2, C-10), 128.00 (C-11a, C-14a), 128.16 and 128.28 (C-1, C-11 and C-4, C-8), 131.29 (C-3, C-9), 135.26 (C-12, C-14), 146.58 (C-4a, C-7a), 156.22 (C-5a, C-6a).

6-(*p*-Fluorophenyl)diquinothiazine (**9b**)

From diquinodithiin **7** Diquinodithiin **7** (0.16 g, 0.5 mmol) was finely powdered together with *p*-fluoroaniline hydrochloride (0.37 g, 2.5 mmol), and the mixture was heated on an oil bath at 200–205 °C for 3 h. After cooling, water (10 ml) was added to the reaction mixture and the resulting solid was filtered off, washed with water, air-dried, and purified by column chromatography (Al₂O₃, CH₂Cl₂) to give 0.14 g (35 %) of 6-(*p*-fluorophenyl)diquinothiazine (**9b**), yellow, mp 248–249 °C.

From 2,2'-dichloro-3,3'-diquinoliny sulfide **8** A solution of sulfide **8** (0.18 g, 0.5 mmol) and *p*-fluoroaniline (0.17 g, 1.5 mmol) in MEDG (5 ml) was refluxed for 3 h. After cooling, the solution was poured into water (20 ml) and alkalized with 5 % aqueous sodium hydroxide to pH = 10. The resulting solid was filtered off, washed with water, and purified by column chromatography (Al₂O₃, CH₂Cl₂) to give 0.16 g (81 %) 6-(*p*-fluorophenyl)diquinothiazine (**9b**), yellow, mp 248–249 °C.

^1H NMR (CDCl_3) δ : 7.31 (m, 4H, H-2, H-10, C_6H_2), 7.47 (m, 4H, H-3, H-9, C_6H_2), 7.56 (d, 2H, H-1, H-11), 7.67 (d, 2H, H-4, H-8), 7.83 (s, 2H, H-12, H-14). ^{13}C NMR (CDCl_3) δ : 115.85 ($J = 22.6$ Hz, *m*-C of $\text{C}_6\text{H}_4\text{F}$), 115.98 (C-12a, C-13a), 125.16 (C-2, C-10), 125.78 (C-11a, C-14a), 125.96 (C-1, C-11), 128.07 (C-4, C-8), 129.37 (C-3, C-9), 132.07 (C-12, C-14), 132.40 ($J = 7.5$ Hz, *o*-C of $\text{C}_6\text{H}_4\text{F}$), 135.59 ($J = 2.5$ Hz, *ipso*-C of $\text{C}_6\text{H}_4\text{F}$), 145.13 (C-4a, C-7a), 150.98 (C-5a, C-6a), 161.83 ($J = 244.6$ Hz, *p*-C of $\text{C}_6\text{H}_4\text{F}$).

EIMS m/z : 395 (M^+ , 75), 394 (M-1, 100), 363 (M-S, 5). Anal. Calcd. for $\text{C}_{24}\text{H}_{14}\text{FN}_3\text{S}$: C, 72.89; H, 3.57; N, 10.63. Found: C, 72.80; H, 3.55; N, 10.41.

Diquino[3,4-b;4',3'-e][1,4]thiazines (**12a–c**)

6*H*-Diquinothiazine (**12a**) and 6-methyldiquinothiazine (**12b**) were obtained from the reaction of sulfide **11** with ammonia and methylamine in hot phenol (Pluta, 1997).

6*H*-Diquinothiazine (**12a**)

Beige, mp 200–201 °C (mp 200–201 °C, Pluta, 1997). ^1H NMR (CDCl_3) δ : 7.64 (t, 2H, H-2, H-12), 7.71 (t, 2H, H-3, H-11), 7.81 (d, 2H, H-4, H-10), 8.04 (d, 2H, H-1, H-13), 8.40 (s, 2H, H-6, H-8). ^{13}C NMR (CDCl_3) δ : 109.10 (C-6a, C-7a), 117.18 (C-13a, C-14b), 117.41 (C-1, C-13), 127.25 (C-2, C-12), 129.49 (C-3, C-11), 130.78 (C-4, C-10), 142.21 (C-4a, C-9a), 147.94 (C-6, C-8), 148.07 (C-13b, C-14a).

6-Methyldiquinothiazine (**12b**)

Yellow, mp 156–157 °C (mp 156–157 °C, Pluta, 1997). ^1H NMR (CDCl_3) δ : 3.54 (s, 3H, CH_3), 7.66 (t, 2H, H-2, H-12), 7.72 (t, 2H, H-3, H-11), 8.11 (d, 2H, H-4, H-10), 8.34 (d, 2H, H-1, H-13), 8.66 (s, 2H, H-6, H-8). ^{13}C NMR (CDCl_3) δ : 43.63 (CH_3), 122.09 (C-1, C-13), 124.17 and 124.42 (C-6a, C-7a and C-13a, C-14b), 127.46 (C-2, C-12), 129.44 (C-3, C-11), 130.11 (C-4, C-10), 148.33 (C-6, C-8), 148.76 and 148.85 (C-4a, C-9a and C-13b, C-14a).

14-(*p*-Fluorophenyl)diquinothiazine (**12c**)

From diquinodithiin **10**

Diquinodithiin **10** (0.16 g, 0.5 mmol) was finely powdered together with *p*-fluoroaniline hydrochloride (0.37 g, 2.5 mmol), and the mixture was heated on an oil bath at 200–205 °C for 3 h. After cooling, water (10 ml) was added to the reaction mixture and the resulting solid was filtered off, washed with water, air-dried, and purified by column chromatography (Al_2O_3 , CH_2Cl_2) to give 0.12 g

(30 %) of 14-(*p*-fluorophenyl)diquinothiazine (**12c**), beige, mp 315–316 °C.

From 4,4'-dichloro-3,3'-diquinoliny sulfide (**11**)

A solution of sulfide **11** (0.18 g, 0.5 mmol) and *p*-fluoroaniline (0.17 g, 1.5 mmol) in MEDG (5 mL) was refluxed for 3 h. After cooling, the solution was poured into water (20 ml) and alkalized with 5 % aqueous sodium hydroxide to pH 10. The resulting solid was filtered off, washed with water and purified by column chromatography (Al_2O_3 , CHCl_3) to give 0.17 g (86 %) of 14-(*p*-fluorophenyl)diquinothiazine (**12c**), beige, mp 315–316 °C.


^1H NMR (CDCl_3) δ : 6.43 (dd, 2H, C_6H_2), 6.77 (m, 2H, C_6H_2), 7.75 (t, 2H, H-2, H-12), 7.85 (t, 2H, H-3, H-11), 8.34 (d, 2H, H-4, H-10), 8.39 (d, 2H, H-1, H-13), 9.06 (s, 2H, H-6, H-8). ^{13}C NMR (CDCl_3) δ : 115.75 ($J = 22.5$ Hz, *m*-C of $\text{C}_6\text{H}_4\text{F}$), 116.30 ($J = 7.5$ Hz, *o*-C of $\text{C}_6\text{H}_4\text{F}$), 122.87 (C-1, C-13), 126.82 (C-13a, C-14b), 128.51 (C-2, C-12), 129.89 (C-6a, C-7a), 130.13 (C-3, C-11), 130.25 (C-4, C-10), 140.57 ($J = 2.5$ Hz, *ipso*-C of $\text{C}_6\text{H}_4\text{F}$), 145.54 (C-13b, C-14a), 147.98 (C-4a, C-9a), 149.49 (C-6, C-8), 158.07 ($J = 238.5$ Hz, *p*-C of $\text{C}_6\text{H}_4\text{F}$). EIMS m/z : 395 (M^+ , 100), 363 (M-S, 20), 300 (M-C $_6\text{H}_4\text{F}$, 17). Anal. Calcd. for $\text{C}_{24}\text{H}_{14}\text{FN}_3\text{S}$: C, 72.89; H, 3.57; N, 10.63. Found: C, 72.77; H, 3.59; N, 10.46.

In vitro lipid peroxidation

Heat-inactivated hepatic microsomes from untreated rats were prepared as described (Rekka *et al.*, 1989). The incubation mixture contained microsomal fraction (corresponding to 2.5 mg of hepatic protein per ml or 4 mM fatty acid residues), ascorbic acid (0.2 mM) in Tris-HCl/KCl buffer (50 mM/150 mM, pH 7.4), and the studied compounds (50–1 μM) dissolved in DMSO. The reaction was initiated by addition of a freshly prepared FeSO_4 solution (10 μM), and the mixture was incubated at 37 °C for 45 min. Lipid peroxidation of aliquots was assessed spectrophotometrically (535 against 600 nm) as TBAR. Both compounds and solvents were found not to interfere with the assay. Each assay was performed in duplicate, and IC_{50} values represent the mean concentration of compounds that inhibit the peroxidation of control microsomes by 50 % after 45 min of incubation. All standard errors are within 10 % of the respective reported values.

Calculation of lipophilicity, molecular mass, surface area, and molecular volume

Lipophilicity (as cLogP), molecular mass (M), surface area (S), and molecular volume (V_M) were calculated using CS



Scheme 1 Reactants: **a** C₆H₅NH₂·HCl (*p*-ClC₆H₄NH₂·HCl, *p*-CH₃OC₆H₄NH₂·HCl), 200–205 °C, 4 h; **b** *p*-CH₃OC₆H₄NH₂, MEDG, reflux, 3 h; **c** 1-naphthylamine·HCl, 200–205 °C, 4 h; **d** 2-naphthylamine·HCl, 200–205 °C, 4 h; **e** 6-aminoquinoline·HCl, 200–205 °C, 4 h

Chem 3D Ultra 7.0 (CambridgeSoft) and Spartan'04 (Wavefunction, Inc. Irvine, CA).

Results and discussion

Synthesis


The synthesis of the title azaphenothiazines was based on the reactions of isomeric diquinodithiins, dichlorodiquinolyl sulfides, and disulfide with amines, ammonia, and acetamide. The fusion reactions of linearly condensed diquinodithiin **1** with hydrochlorides of aniline and its *p*-substituted derivatives such as *p*-chloroaniline and *p*-methoxyaniline led to tetracyclic 9-substituted 6*H*-quinobenzothiazines **3a–c** (Scheme 1). 9-Methoxy-6*H*-quinobenzothiazine **3c** was obtained in better yield in the reaction of 2,2'-dichloro-3,3'-diquinolyl disulfide **2** with *p*-methoxyaniline in monomethyl ether of diethylene glycol. The similar reaction of diquinodithiin **1** with hydrochlorides of 1-naphthylamine, 2-naphthylamine, and 6-aminoquinoline gave pentacyclic 7*H*-quinonaphthothiazine **4**, 14*H*-quinonaphthothiazine **5**, and 7*H*-diquinobenzothiazine **6**. The reaction of isomeric diquinodithiin **7** with acetamide and *p*-fluoroaniline hydrochloride gave linearly condensed pentacyclic 6*H*-diquinobenzothiazines **9a** and 6-(*p*-

fluorophenyl)diquinobenzothiazine **9b** (Scheme 2). Analogous reaction of another isomeric diquinodithiin **10** with *p*-fluoroaniline hydrochloride led to angularly condensed diquinobenzothiazine **12c**. Better yields of the fluoroaniline products **9b** and **12c** were achieved when *x,x'*-dichloro-3,3'-diquinolyl sulfides **8** and **11** (*x* = 2 and 4) were used. Sulfide **11** reacted also with ammonia or methylamine in hot phenol to give diquinobenzothiazines **12a, b**.

The described syntheses were monitored by TLC analysis. All chromatograms of new compounds showed characteristic for azaphenothiazines (Jeleń *et al.*, 2011) color changing during irradiation with UV light from blue to yellow (**4, 9b**), from yellow to green (**5, 6**), from orange to yellow (**12c**), and from yellow to orange (**7c**).

Structure

It is well known that the synthesis of phenothiazines can proceed via the Smiles rearrangement of the S–N type of the appropriate sulfide (Pluta *et al.*, 2009). The identification of the product structures was based on the spectroscopic ¹H NMR and MS analysis. In the case of the reactions of sulfides **7** and **11**, the products **9** and **12** possessed the C_{2v} symmetry (the left part was a mirror image of the right one) what excluded the stage of rearrangement. The reactions of diquinodithiin **1** and disulfide **2** with



Scheme 2 Reactants: **a** CH₃CONH₂, K₂CO₃, 180 °C, 0.5 h; **b** *p*-FC₆H₄NH₂·HCl, 200–205 °C, 3 h; **c** *p*-FC₆H₄NH₂, MEDG, reflux, 3 h; **d** NH₃ (CH₃NH₂), phenol, 180 °C, 1 h

anilines proceeded similarly without the stage of rearrangement to give tetracyclic quinobenzothiazines **3a–c** (Jeleń and Pluta, 2009). The reaction with 1-naphthylamine gave pentacyclic quinonaphthothiazine **4**. On the contrary, the reactions with 2-naphthylamine and 6-aminoquinoline were more complex as there were two possibilities of the thiazine ring formation. The ¹H NMR analysis of the reaction products pointed at compounds **5** and **6** excluding compounds **13** and **14**, as evidenced from coupling constants; the H-5 and H-6 protons in compounds **5** and **6** showed a coupling constant J_{ortho} , whereas analogous protons in compounds **13** and **14** (H-7/H-12 and H-5/H-14, respectively) would have shown a coupling constant J_{para} , which is very small (i.e., $J_{1,4} = 0.6\text{--}0.8$ Hz in naphthalene (Hamm and von Philipsborn, 1971; Lucchini and Wells, 1976) and $J_{5,8} = 0.5\text{--}0.8$ Hz in quinoline (Hamm and von Philipsborn, 1971; Jones, 1977)). We did not observe such small values of coupling constants in the reaction products **5** and **6**.

Antioxidant activity

The effect of the new derivatives on non-enzymatic lipid peroxidation of rat hepatic microsomal membrane lipids was investigated in vitro. Most of the studied derivatives demonstrated significant antioxidant activity, with IC₅₀ values between 1 and 23 μM (Table 1). It is worthwhile to mention that under the same experimental conditions known potent antioxidants, trolox ((S)-(-)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid) and probucol (4,4'-[(1-methylethylidene)bis(thio)]bis[2,6-bis(1,1-dimethylethyl)phenol]), exhibited IC₅₀ values of 25 μM and >1 mM, respectively (Kourounakis *et al.*, 2008).

Table 1 IC₅₀ values for in vitro lipid peroxidation (LP), LogP, molecular volume (V_M), and molecular mass (M) as well as surface area (S) of the tested compounds

Compound	LP IC ₅₀ (μM)	LogP	M	S (Å ²)	V _M (Å ³)
3a	23	3.37	250.06	253.13	246.02
3b	3	3.93	284.02	268.84	259.50
3c	2	3.25	280.07	283	273.38
4	2	4.37	300.07	297.74	296.96
5	6	4.37	300.07	297.68	296.87
6	16	3.46	301.07	293.28	291.10
9a	>1000	4.20	301.07	295.91	291.54
9b	>1000	6.00	395.09	374.91	379.66
12a	1	2.71	301.07	291.11	290.87
12b	500	4.77	315.08	317.08	321.82
12c	>1000	4.51	395.09	359.77	375.69

Further, all of the active new derivatives were significantly much more potent than previously studied tricyclic dipyr-dithiazines (IC₅₀ of most active compounds was between 64 and 470 μM) (Morak-Młodawska *et al.*, 2010). The time course of lipid peroxidation, as affected by various concentrations of representative compounds, is depicted in Fig. 1.

Tetracyclic NH-azaphenothiazines **3a–c** exhibited significant activity dependent on the substitution (H, Cl, and OCH₃) on the benzene ring (Table 1). From the pentacyclic compounds, the angularly fused with unsubstituted, the thiazine nitrogen atom (**4–6** and **12a**) exhibited very significant activity with most active compound **12a**, which showed an IC₅₀ of 1 μM. The change of the quinoline moiety into naphthalene (compare compounds **4** and **5** with




Fig. 1 Representative *graphs* of the time course of lipid peroxidation as affected by various concentrations of compounds **3a–c**, **5**, **6**, and **12a**. IC_{50} values are calculated according to these results as the

concentration showing 50 % inhibition of the lipid peroxidation reaction at 45 min incubation time

6) marginally increased activity. However, compounds with a linearly fused ring system (**9a** and **9b**) and/or a large aryl substituent at the thiazine nitrogen atom (**9b** and **12c**) did not show any antioxidant activity, while compound **12b**, with a small substituent, exhibited very weak activity.

Considering three isomers (**6**, **9a**, and **12a**), one can find that their antioxidant activity increased with decreasing lipophilic character represented by the logP values. On the other hand, the least active compounds (**9b**, **12b**, and **12c**) exhibited high values of molecular descriptors such as molecular mass ($M > 315$), molecular volume ($V_M > 321$), and surface area ($S > 317$, Table 1). However, attempts to correlate the activity with those properties turned out to be unsatisfactory.

In conclusion, eleven tetracyclic and pentacyclic (linearly or angularly condensed) azaphenothiazines were synthesized, and structure–(antioxidant)activity relationships were investigated. The type of the ring fusion was concluded from the 1H NMR spectra. The degree of antioxidant activity of these derivatives seems to depend on their lipophilicity and molecular mass. The non-substitution of the thiazine nitrogen atom, the type of ring system fusion, and the nature of substituents promote activity. Finally, it is the first time to our knowledge that azaphenothiazines are shown to exhibit such potent antioxidant activity.

Acknowledgments The synthesis and the structure elucidation are supported by the Medical University of Silesia (Grant KNW-1-032/K/3/0).

Conflict of interest Authors have no financial/commercial conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Aaron JJ, Gaye Seye MD, Trajkovska S, Motohashi N (2009) Bioactive phenothiazines and benzo[a]phenothiazines: spectroscopic studies and biological and biomedical properties and applications. *Top Heterocycl Chem* 16:153–231
- Asghar MN, Alam Q, Augusten S (2012) Fluphenazine hydrochloride radical cation assay: a new, rapid and precise method to determine in vitro total antioxidant capacity of fruit extracts. *Chin Chem Lett* 23:1271–1274
- Borges MBD, Dos Santos CG, Yokomizo CH, Sood RR, Vitovic PP, Kinnunen KJ, Rodrigues T, Nantes IL (2010) Characterization of hydrophobic interaction and antioxidant properties of the phenothiazine nucleus in mitochondrial and model membranes. *Free Radical Res* 44:1054–1063
- Dasgupta A, Dastridara SG, Shirataki Y, Motohashi Y (2008) Antibacterial activity of artificial phenothiazines and isoflavones from plants. *Top Heterocycl Chem* 15:67–132
- Gupta RR, Kumar M (1988) Synthesis, properties and reactions of phenothiazines. In: Gupta RR (ed) *Phenothiazines and 1,4-benzothiazines—chemical and biological aspects*. Elsevier, Amsterdam, pp 1–161
- Hamm P, von Philipsborn W (1971) Protonenresonanzspektren von aromatischen *N*-Oxiden Berechnung der chemischen Verschiebungen, verursacht durch die Feldeffekte der *N*-O-gruppe. *Helv Chim Acta* 54:2363–2401
- Jeleń M, Pluta K (2009) Synthesis of quinobenzo-1,4-thiazines from diquino-1,4-dithiin and 2,2'-dichloro-3,3'-diquinoliny disulfide. *Heterocycles* 78:2325–2336
- Jeleń M, Morak-Młodawska B, Pluta K (2011) Thin-layer chromatographic detection of new azaphenothiazines. *J Pharm Biomed Anal* 55:466–471

- Jeleń M, Pluta K, Zimecki M, Morak-Młodawska B, Artym J, Kocięba M (2013) Synthesis and selected immunological properties of substituted quino[3,2-b]benzo[1,4]thiazines. *Eur J Med Chem* 63:444–456
- Jones G (1977) In: Jones G (ed), *The chemistry of heterocyclic compounds*, vol 32, Quinolines, Part 1. Wiley, London, p 11–12
- Kourounakis AP, Charitos C, Rekka EA, Kourounakis PN (2008) Lipid-lowering (hetero)aromatic tetrahydro-1,4-oxazine derivatives with antioxidant and squalene synthase inhibitory activity. *J Med Chem* 51:5861–5865
- Kumar M, Sharma K, Samarth RM, Kumar A (2010) Synthesis and antioxidant activity of quinobenzothiazinones. *Eur J Med Chem* 45:4467–4472
- Liu Z-Q, Tang Y-Z, Wu D (2009) Antioxidant effects of phenothiazine, phenoxazine, and iminostilbene on free-radical-induced oxidation of linoleic acid and DNA. *J Phys Org Chem* 22:1009–1014
- Lucchini V, Wells PR (1976) Proton magnetic resonance spectra of monosubstituted naphthalenes. *Org Magn Reson* 8:137–140
- Matralis AN, Katselou MG, Nikitakis A, Kourounakis AP (2011) Novel benzoxazine and benzothiazine derivatives as multifunctional antihyperlipidemic agents. *J Med Chem* 54:5583–5591
- Morak-Młodawska B, Pluta K, Matralis AN, Kourounakis AP (2010) Antioxidant activity of newly synthesized 2,7-diazaphenothiazines. *Arch Pharm Chem Life Sci* 343:268–273
- Motohashi N, Kawase M, Satoh K, Sakagami H (2006) Cytotoxic potential of phenothiazines. *Curr Drug Targets* 7:1055–1066
- Naik N, Kumar HV, Veena V (2012) Novel phenothiazine analogues: synthesis and a new perceptivity into their antioxidant potential. *Pharmacia Lett* 4:786–794
- Nowak M, Pluta K, Suwińska K (2002) Synthesis of novel heteropentacenes containing nitrogen, sulfur and oxygen or selenium. *New J Chem* 26:1216–1220
- Nowak M, Pluta K, Kloc K, Siegriest T (2003) Synthesis and X-ray analysis of isomeric diazadithiapentacenes. *Heterocycles* 60:2045–2056
- Nowak M, Pluta K, Suwińska K, Straver L (2007) Synthesis of new pentacyclic diquinothiazines. *J Heterocycl Chem* 44:543–550
- Pluta K (1994) Synthesis and NMR assignment of 1,4-oxathiino[3,2-c:5,6-c']diquinoline. *J Heterocycl Chem* 31:557–560
- Pluta K (1997) Synthesis and properties of 14-substituted 1,4-thiazinodiquinolines. *Phosphorus Sulfur Silicon* 126:145–156
- Pluta K, Morak-Młodawska B, Jeleń M (2009) Synthesis and properties of diaza-, triaza- and tetraazaphenothiazines. *J Heterocycl Chem* 46:355–391
- Pluta K, Jeleń M, Morak-Młodawska B, Zimecki M, Artym J, Kocięba M (2010) Anticancer activity of newly synthesized azaphenothiazines in NCI's anticancer screening. *Pharmacol Rep* 62:319–332
- Pluta K, Morak-Młodawska B, Jeleń M (2011) Recent progress in biological activities of synthesized phenothiazines. *Eur J Med Chem* 46:3179–3189
- Rekka E, Kolstee J, Timmerman H, Bast A (1989) The effect of some H₂-receptor antagonists on rat hepatic microsomal cytochrome P-450 and lipid peroxidation in vitro. *Eur J Med Chem* 24:43–54
- Zhang H-Y, Yang D-P, Tang G-Y (2006) Multipotent antioxidants: from screening to design. *Drug Discov Today* 11:749–754
- Zimecki M, Artym J, Kocięba M, Pluta K, Morak-Młodawska B, Jeleń M (2009) Immunosuppressive activities of newly synthesized azaphenothiazines in human and mouse models. *Cell Mol Biol Lett* 14:622–635