

# Synthetic transporters for sulfate: a new method for the direct detection of lipid bilayer sulfate transport†

Cite this: *Chem. Sci.*, 2014, 5, 1118Nathalie Busschaert,<sup>a</sup> Louise E. Karagiannidis,<sup>a</sup> Marco Wenzel,<sup>‡,a</sup> Cally J. E. Haynes,<sup>a</sup> Neil J. Wells,<sup>a</sup> Philip G. Young,<sup>b</sup> Damjan Makuc,<sup>cd</sup> Janez Plavec,<sup>\*cde</sup> Katrina A. Jolliffe<sup>\*b</sup> and Philip A. Gale<sup>\*a</sup>

The transmembrane transport of anions by small synthetic molecules is a growing field in supramolecular chemistry and has focussed mainly on the transmembrane transport of chloride. On the other hand, the transport of the highly hydrophilic sulfate anion across lipid bilayers is much less developed, even though the inability to transport sulfate across cellular membranes has been linked to a variety of genetic diseases. Tris-thioureas possess high sulfate affinities and have been shown to be excellent chloride and bicarbonate transporters. Herein we report the sulfate transport abilities of a series of tris-ureas and tris-thioureas based on a tris(2-aminoethyl)amine or cyclopeptide scaffold. We have developed a new technique based on <sup>33</sup>S NMR that can be used to monitor sulfate transport, using <sup>33</sup>S-labelled sulfate and paramagnetic agents such as Mn<sup>2+</sup> and Fe<sup>3+</sup> to discriminate between intra- and extravesicular sulfate. Reasonable sulfate transport abilities were found for the reported tris-ureas and tris-thioureas, providing a starting point for the development of more powerful synthetic sulfate transporters that can be used in the treatment of certain channelopathies or as a model for biological sulfate transporters.

Received 17th July 2013  
Accepted 9th August 2013

DOI: 10.1039/c3sc52006d

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

## Introduction

The lipid bilayer transport of anionic species such as chloride and bicarbonate are key processes in biology mediated by transmembrane ion channels that regulate membrane potential, cell volume and intracellular pH.<sup>1</sup> Synthetic molecules and assemblies that can replicate the transport activity of these natural transport systems either *via* carrier, relay or channel mechanisms have been reported recently.<sup>2</sup> This area is attracting interest as this generation of compounds can replicate the activity of faulty anion channels, and may have future therapeutic potential in the treatment of diseases caused by anion dysregulation (such as cystic fibrosis)<sup>3</sup> or in disruption of pH

gradients in cells and triggering apoptosis, and hence potential anti-cancer activity.<sup>4,5</sup>

Sulfate transport through lipid bilayers is challenging due to the high hydrophilicity of this anion ( $\Delta G_h = -1080 \text{ kJ mol}^{-1}$  as compared to  $-340 \text{ kJ mol}^{-1}$  for chloride).<sup>6</sup> In fact sulfate is often used as a tool in transmembrane transport experiments to determine the transport mechanism as it can usually be assumed that SO<sub>4</sub><sup>2-</sup> cannot be transported and hence the use of sulfate as extravesicular anion allows antiport and co-transport processes to be discriminated.<sup>7</sup> Although there is some evidence for sulfate transport through synthetic membrane-spanning channels,<sup>8</sup> transport of sulfate by mobile carriers that require full or partial dehydration of the anion prior to transport has remained largely unexplored. On the other hand, much effort has been devoted to the problem of selective extraction of sulfate<sup>9</sup> from mixtures due to the relevance of this process to nuclear waste remediation.<sup>10</sup> A number of groups have also synthesised selective receptors for sulfate in order to meet the challenge of mimicking the selectivity of biological sulfate binding sites.<sup>11</sup>

From a biological perspective, sulfate is one of the most abundant anions in human plasma and is the major sulfur source in many organisms.<sup>12</sup> Sulfate transporters are therefore important biological proteins and misregulation of sulfate transport is present in a number of disease states. For example, defects in the diastrophic dysplasia sulfate transporter (DTDST) can lead to various chondrodysplasia due to insufficient

<sup>a</sup>Chemistry, University of Southampton, Southampton, SO17 1BJ, UK. Tel: +44 (0)23 8059 3332. E-mail: philip.gale@soton.ac.uk

<sup>b</sup>School of Chemistry (F11), The University of Sydney, 2006 NSW, Australia. E-mail: kate.jolliffe@sydney.edu.au; Fax: +61 2 9351 3329; Tel: +61 2 9351 2297

<sup>c</sup>Slovenian NMR Centre, National Institute of Chemistry, Hajdrihova 19, SI-1000 Ljubljana, Slovenia. E-mail: janez.plavec@ki.si

<sup>d</sup>EN-FIST Centre of Excellence, Dunajska 156, SI-1000 Ljubljana, Slovenia


<sup>e</sup>Faculty of Chemistry and Chemical Technology, University of Ljubljana, SI-1000 Ljubljana, Slovenia

† Electronic supplementary information (ESI) available: Experimental procedures, Hill plots, cholesterol test, <sup>13</sup>C NMR tests, symport tests, HPTS pH change tests, calcein leakage assay, lucigenin tests, <sup>33</sup>S NMR tests on all compounds with Mn<sup>2+</sup> and Fe<sup>3+</sup>, SEM-EDX experiments. See DOI: 10.1039/c3sc52006d

‡ Present address: Fraunhofer IKTS, 01217, Dresden, Germany.




sulfation of cartilage proteoglycans, resulting in symptoms such as dwarfism, spinal deformations and other abnormalities of the joints.<sup>13</sup> We therefore decided to examine whether small molecules with high sulfate affinities – that could also encapsulate sulfate within a three dimensional framework shielding it from the hydrophobic interior of the lipid bilayer – could function as transmembrane sulfate transporters. Gale and co-workers have previously reported a series of fluorinated ureas and thioureas based on tris(2-aminoethyl)amine (*tren*), including compounds 1–6, that showed strong affinity and selectivity for sulfate in DMSO-*d*<sub>6</sub>/0.5% water.<sup>14,15</sup> It was also observed that compound 4 (and to a lesser extent 3 and 5) were able to transport chloride out of unilamellar phospholipid vesicles in the presence of external sulfate. It was suggested that this was due to the Cl<sup>-</sup>/SO<sub>4</sub><sup>2-</sup> (or HCl/HSO<sub>4</sub><sup>-</sup>) antiport ability of 3–5, but at the time of publication only indirect evidence for this claim could be obtained.



- |   |     |                                                                            |   |     |                                                                            |
|---|-----|----------------------------------------------------------------------------|---|-----|----------------------------------------------------------------------------|
| 1 | X=O | R=C <sub>6</sub> H <sub>5</sub>                                            | 2 | X=S | R=C <sub>6</sub> H <sub>5</sub>                                            |
| 3 | X=O | R=C <sub>6</sub> F <sub>5</sub>                                            | 4 | X=S | R=C <sub>6</sub> F <sub>5</sub>                                            |
| 5 | X=O | R= <i>m</i> -C <sub>6</sub> H <sub>3</sub> (CF <sub>3</sub> ) <sub>2</sub> | 6 | X=S | R= <i>m</i> -C <sub>6</sub> H <sub>3</sub> (CF <sub>3</sub> ) <sub>2</sub> |

Receptors 1–6 still showed higher transport activity for chloride, nitrate and bicarbonate than for sulfate and we therefore decided to investigate the transmembrane anion transport ability of related compounds with higher sulfate binding affinities. Cage-like anion receptors often have high selectivity for a specific guest and tris-amide cages have previously been used as chloride/nitrate transporters.<sup>16</sup> We therefore decided to select potential sulfate transporters from known cage-like sulfate receptors. The recently reported tripod<sup>17</sup> and cryptand-like<sup>18,19</sup> anion receptors 7–9, based on a *Lissoclinum*-type cyclic peptide scaffold, have been found to bind sulfate ions with high affinity ( $K_a > 10^4 \text{ M}^{-1}$  in DMSO-*d*<sub>6</sub>/0.5% water for all three receptors) and selectivity. This is attributed to the ability of these receptors to form nine hydrogen bonds to sulfate ions – six from the thiourea groups and three from the cyclic peptide backbone amides. In contrast, spherical anions such as chloride bind through the thiourea groups only.<sup>17,18</sup> Notably, receptor 8 binds sulfate strongly even in highly competitive solvents ( $K_a = 1.3 \times 10^3 \text{ M}^{-1}$  in DMSO-*d*<sub>6</sub>/20% water at 330 K) indicating that the anion is well shielded from the external environment (compounds 7 and 9 were not soluble enough to allow determination of binding constants under these conditions). In this article we report the anion transport abilities of cyclopeptides 7–9, including the ability of compound 9 to transport the highly hydrophilic sulfate anion across a phospholipid bilayer. We also provide a new method that can directly prove the occurrence of sulfate transport by both the *tren*-based compounds 1–6 and cyclopeptides 7–9 using <sup>33</sup>S NMR.



## Results and discussion

### Chloride, nitrate and bicarbonate transport by 7–9

The chloride, nitrate and bicarbonate transport abilities of *tren*-based compounds 1–6 have been previously reported by Gale and co-workers.<sup>14,15</sup> It was found that in all cases tris-thioureas (2, 4 and 6) are more potent anion transporters than their analogous tris-ureas (1, 3 and 5) and that the fluorinated transporters (3–6) are more active than their unfluorinated counterparts and have the ability to transport chloride out of phospholipid vesicles at concentrations of 1 : 250 000 carrier-to-lipid (on average one transporter per vesicle).<sup>15,20</sup> Gale and co-workers also showed that 1–6 display high affinity for sulfate ( $K_a > 10^4 \text{ M}^{-1}$ ), moderate affinity for chloride ( $K_a \sim 500 \text{ M}^{-1}$ ) and no affinity for nitrate in DMSO-*d*<sub>6</sub>/0.5% water. Likewise, Jolliffe and co-workers have found that cyclopeptide based tris-thioureas 7–9 also bind sulfate with high affinity and chloride with moderate affinity (no binding was observed for nitrate).<sup>17–19</sup> This similarity in anion binding behaviour between tris-(thio)ureas 1–6 and the structurally related tris-thioureas 7–9, has prompted us to investigate the trans-membrane anion transport ability of cyclopeptides 7–9.

Initially, chloride transport properties were studied using a series of unilamellar 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) vesicles loaded with NaCl (489 mM) and suspended in an external NaNO<sub>3</sub> (489 mM) solution using the previously reported procedures utilised for 1–6.<sup>15,21</sup> A sample of receptor 7–9 (4 mol% carrier to lipid) was added as a DMSO solution and the resultant chloride efflux was monitored using a chloride selective electrode. After 300 s, the vesicles were lysed by addition of detergent and the final reading of the electrode was used to calibrate 100% release of chloride. The results are shown in Fig. 1 and reveal significant chloride transport by the






Fig. 1 Chloride efflux promoted by 7–9 (4 mol% carrier to lipid) from unilamellar POPC vesicles loaded with 489 mM NaCl buffered to pH 7.2 with 5 mM sodium phosphate salts. The vesicles were dispersed in 489 mM NaNO<sub>3</sub> buffered to pH 7.2 with 5 mM sodium phosphate salts. At the end of the experiment, detergent was added to lyse the vesicles and calibrate the electrode to 100% chloride efflux. Each point represents the average of three trials. DMSO was used as a control.

cryptand-like compounds **8** and **9**. The benzyl bridged receptor **9** shows a higher transport activity compared to receptor **8**, whilst tripodal receptor **7** was found to be not active. The chloride transport ability of **8** and **9** were further quantified by means of Hill plot,<sup>22</sup> where the chloride efflux out of the vesicles is monitored at various concentrations of transporter to yield EC<sub>50,270s</sub> values (*i.e.* the concentration of transporter (in mol%) required to obtain 50% chloride efflux in 270 s). The EC<sub>50,270s</sub> value of compound **9** (1.13 mol%) is three times lower than of compound **8** (3.11 mol%), which points at a substantial influence of the variation in the bridgehead group between the two cryptand-like receptors. Presumably the enhanced lipophilicity of **9** and the larger cavity size, allowing quicker movement of the anions in and out of the cavity (necessary for uptake and release of the anion during transport), result in higher transport activity for receptor **9**.

In order to investigate the potential mechanisms of chloride release we repeated the transport experiments with 70 : 30 POPC : cholesterol vesicles. The addition of cholesterol decreases the fluidity of the membrane and can therefore have a profound effect on the transport activity of a mobile carrier which relies on diffusion through the bilayer.<sup>23</sup> Under these conditions a reduction in chloride transport rate was observed for compounds **8** and **9** (see ESI†), evidence in support of the cyclopeptides functioning as discrete molecular carriers rather than as membrane spanning channels. Furthermore, Hill analyses of transport data reveal *n* values of 1.1 and 1.0 for **8** and **9** respectively, and also point at a mobile carrier mechanism (see ESI† for Hill plots).<sup>24</sup>

In order to determine whether these results were due to a Cl<sup>-</sup>/NO<sub>3</sub><sup>-</sup> antiport mechanism, we carried out comparable experiments with sulfate and bicarbonate as external anions. The anions differ in their hydration energy ( $\Delta G_{\text{hydr}}(\text{SO}_4^{2-}) - 1080 \text{ kJ mol}^{-1}$ ;  $\Delta G_{\text{hydr}}(\text{HCO}_3^-) - 335 \text{ kJ mol}^{-1}$ ;  $\Delta G_{\text{hydr}}(\text{NO}_3^-) - 300 \text{ kJ mol}^{-1}$ )<sup>6</sup> and their topology (NO<sub>3</sub><sup>-</sup> and HCO<sub>3</sub><sup>-</sup> planar;

SO<sub>4</sub><sup>2-</sup> tetrahedral). Consequently an altered transport activity is expected for an antiport mechanism. In this test a series of unilamellar POPC vesicles were prepared loaded with 450 mM NaCl and suspended in an isotonic external Na<sub>2</sub>SO<sub>4</sub> solution (162 mM). Transport is initiated by the addition of a DMSO solution of the receptors (4 mol% with respect to lipid) and two minutes after the addition of the transporters NaHCO<sub>3</sub> was added to the external Na<sub>2</sub>SO<sub>4</sub> solution. The results of this combined sulfate and bicarbonate assay are shown in Fig. 2. It can be seen that the addition of NaHCO<sub>3</sub> led to a marked acceleration of chloride efflux for receptor **9** while only minor transport is observed for receptor **8**, confirming the substantial influence of the bridgehead group on chloride transport (Fig. 2). In addition, a significant chloride efflux was observed in the presence of sulfate as external anion for **9**. We have observed such behaviour previously for the open tripodal receptors **1–6** (ref. 15) and this prompted us to investigate the sulfate transport behaviour of these receptors in more detail (*vide infra*).

Although the marked difference in transport rate between external nitrate, bicarbonate and sulfate is evidence in support of an antiport mechanism, there is no direct evidence that any anion other than chloride is transported across the phospholipid bilayer by 7–9. In order to provide direct evidence for Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> antiport, we performed the <sup>13</sup>C NMR experiment that was developed by Davis, Gale and Quesada<sup>25</sup> and that was also used to prove bicarbonate transport by receptors **1** and **2**.<sup>14</sup> In these experiments large POPC vesicles (5 μm) were prepared, filled with <sup>13</sup>C-labeled NaHCO<sub>3</sub> and suspended in an isotonic Na<sub>2</sub>SO<sub>4</sub> solution. One <sup>13</sup>C-NMR signal was observed (*ca.* 165.5 ppm) and this was assigned as the signal for intravesicular H<sup>13</sup>CO<sub>3</sub><sup>-</sup> (Fig. 3). In order to promote Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchange 100 mM NaCl was added, followed by a DMSO solution of the transporters (4 mol% with respect to lipid) and this mixture was allowed to incubate for 1 hour at room temperature with stirring.




Fig. 2 Chloride efflux promoted by 7–9 (4 mol% carrier to lipid) from unilamellar POPC vesicles loaded with 450 mM NaCl buffered to pH 7.2 with 20 mM sodium phosphate salts. The vesicles were dispersed in 162 mM Na<sub>2</sub>SO<sub>4</sub> buffered to pH 7.2 with 20 mM sodium phosphate salts. At *t* = 120 s, a solution of NaHCO<sub>3</sub> was added to give a 40 mM external concentration. At the end of the experiment, detergent was added to lyse the vesicles and calibrate the electrode to 100% chloride efflux. Each point represents the average of three trials. DMSO was used as a control.






Fig. 3  $^{13}\text{C}$  NMR experiments to detect bicarbonate transport by 7–9 with DMSO used as a control. The left-hand column shows the experimental set-up, the 4 subsequent columns show results for  $\text{HCO}_3^-/\text{Cl}^-$  exchange and the last column shows the result for  $\text{HCO}_3^-/\text{SO}_4^{2-}$  exchange (no addition of NaCl). All spectra were recorded in 9 : 1 water :  $\text{D}_2\text{O}$  solutions and are shown at absolute intensity.  $^{13}\text{C}$  NMR spectra of (i) POPC vesicles loaded with 155 mM  $\text{NaH}^{13}\text{CO}_3$ , 20 mM phosphate buffer, pH 7.4, and dispersed in 50 mM  $\text{Na}_2\text{SO}_4$ , 20 mM phosphate buffer, pH 7.4; (ii) after the addition of the 100 mM NaCl pulse; (iii) after 1 hour incubation with transporter or DMSO (4 mol% transporter to lipid); (iv) after the addition of  $\text{MnCl}_2$  (1.5 mol%  $\text{Mn}^{2+} : \text{Cl}^-$  ratio); (v) after the addition of detergent.

The result of this NMR test is shown in Fig. 3 for compounds 7–9 (the results for 1–6 can be found in ESI $^\dagger$ ). It can be seen that the addition of NaCl did not lead to any leakage of the vesicles as the  $^{13}\text{C}$  NMR spectrum remained unchanged. However, when active transporter 9 was added, a new peak appeared 2.5 ppm downfield of the original  $^{13}\text{C}$  NMR signal (ca. 163.0 ppm) and this was assigned as the extravesicular  $\text{H}^{13}\text{CO}_3^-$  signal. In order to prove that this new signal is indeed extravesicular  $\text{H}^{13}\text{CO}_3^-$  a small amount of paramagnetic  $\text{Mn}^{2+}$  was added (1.5 mol% with respect to NaCl). As  $\text{Mn}^{2+}$  cannot cross the lipid bilayer, only extravesicular signals are broadened, while intravesicular signals remain sharp. Fig. 3 clearly shows that only cyclopeptide 9 is able to transport  $\text{HCO}_3^-$  out of the vesicle, while cyclopeptides 7 and 8 showed no evidence for bicarbonate transport. The combination of this experiment (Fig. 3), where 9 is able to transport  $\text{HCO}_3^-$  out of the vesicle in the presence of external chloride, and the previous experiments (Fig. 2), where 9 is able to transport  $\text{Cl}^-$  out of the vesicles in the presence of external bicarbonate, suggests that the receptor 9 functions as a  $\text{Cl}^-/\text{HCO}_3^-$  antiporter. Furthermore, compound 9 is also able to transport  $\text{HCO}_3^-$  out of the vesicles when sulfate is the only external anion (no addition of NaCl, Fig. 3 – right hand column), albeit to a smaller extent than in the presence of chloride (again confirming antiport). This suggests that the cyclopeptides can perform the biologically relevant  $\text{HCO}_3^-/\text{SO}_4^{2-}$  exchange $^{12}$  and it gave us another reason to investigate the sulfate transport abilities of these tris-(thio)ureas in more detail.

#### Indirect evidence for sulfate transport by 1–9

In order to investigate the sulfate transport behaviour of 1–9 in more detail, we started by looking at the chloride efflux

mediated by 1–9 (4 mol% carrier to lipid) in the presence of external sulfate. In this test a series of unilamellar POPC vesicles are prepared, loaded with 450 mM NaCl and suspended in an isotonic external  $\text{Na}_2\text{SO}_4$  solution (162 mM). Transport is initiated by the addition of a DMSO solution of the receptors (4 mol% with respect to lipid) and chloride efflux is monitored for 5 minutes using a chloride selective electrode. The result of this test is given in Fig. 4 and shows that compounds 3, 4, 5 and 9 show significant chloride transport in the presence of sulfate




Fig. 4 Chloride efflux promoted by 1–9 (4 mol% carrier to lipid) from unilamellar POPC vesicles loaded with 450 mM NaCl buffered to pH 7.2 with 20 mM sodium phosphate salts. The vesicles were dispersed in 162 mM  $\text{Na}_2\text{SO}_4$  buffered to pH 7.2 with 20 mM sodium phosphate salts. At the end of the experiment, detergent was added to lyse the vesicles and calibrate the electrode to 100% chloride efflux. Each point represents the average of three trials. DMSO was used as a control.





(>50% chloride efflux in 5 minutes) and are therefore the best candidates for  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport.

There are several possible explanations for the chloride transport observed in Fig. 4, including (1)  $\text{M}^+/\text{Cl}^-$  symport; (2) non-specific leaking or destruction of the vesicles; (3)  $\text{H}^+/\text{Cl}^-$  symport,  $\text{OH}^-/\text{Cl}^-$  antiport; (4)  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport. The first option,  $\text{M}^+/\text{Cl}^-$  symport, was excluded when repeats with encapsulated KCl or CsCl (instead of NaCl) showed no difference in chloride transport and proved that the counter cation does not play a significant role in the chloride transport by 1–9 (see ESI†). The second option, leaking of the vesicles, was discarded using a calcein leakage assay.<sup>26</sup> Calcein is a large and highly charged fluorescent dye that self-quenches at high concentration. Thus, when calcein is encapsulated inside vesicles the fluorescence intensity is initially low and increases when it leaks out of the vesicles or when the vesicles are destroyed. However, none of the receptors 1–9 showed any sign of leakage, even after 12 hours (see ESI†). The third alternative explanation,  $\text{H}^+/\text{Cl}^-$  symport or the equivalent  $\text{OH}^-/\text{Cl}^-$  antiport, can be investigated by monitoring the internal pH of the vesicle using the pH-sensitive dye 8-hydroxy-1,3,6-pyrenetrisulfonate (HPTS).<sup>27</sup> This test has been previously reported for 1–6 and showed either a minor pH increase (1, 2, 3, 6) or a large pH decrease (4, 5). This decrease in pH cannot be explained *via* an  $\text{H}^+/\text{Cl}^-$  symport or  $\text{OH}^-/\text{Cl}^-$  antiport mechanism and it was postulated that the observed pH change could be due to  $\text{HSO}_4^-/\text{HCl}$  exchange.<sup>15</sup> The results for the cyclopeptide-based compounds 7–9 also showed no significant change in the internal pH (see ESI†) and therefore an  $\text{H}^+/\text{Cl}^-$  symport or  $\text{OH}^-/\text{Cl}^-$  antiport mechanism can be excluded for all tris-(thio)ureas 1–9 reported in this article.

The only plausible remaining explanation for the transport observed in Fig. 4 is therefore  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport. We have previously developed an indirect method based on the halide sensitive dye lucigenin to provide evidence for sulfate transport.<sup>15</sup> This method provided evidence that led us to suggest that compounds 4 and 5 have considerable sulfate transport ability and we now used this method to investigate sulfate transport by compounds 8 and 9. In brief, we prepared a series of vesicles loaded with NaCl and lucigenin (to monitor the intravesicular chloride concentration)<sup>28</sup> that were suspended in a NaCl solution (100 mM). The lack of both a chloride and pH gradient rules out HCl co-transport. The experiment was initiated by the addition of various anions ( $\text{Na}_2\text{SO}_4$ ,  $\text{NaNO}_3$  or NaCl) followed by the addition of a methanol solution of 9. The result depicted in Fig. 5 shows a significant increase in fluorescence upon the addition of 9 when sulfate was added, indicating that chloride is transported out of the vesicle. When  $\text{NaNO}_3$  was added instead of  $\text{Na}_2\text{SO}_4$ , a much faster increase in fluorescence was observed, as would be expected for an antiport process of a more lipophilic anion (Fig. 5). As a control, the experiment was repeated with the addition of NaCl instead of  $\text{Na}_2\text{SO}_4$ . In this case no change in fluorescence was observed, which is evidence in support of an antiport mechanism. No change in fluorescence was observed when compound 8 or methanol was added (see ESI†), indicating that 9 is the only cyclopeptide-based tris-thiourea able to transport sulfate.




Fig. 5 Unilamellar POPC vesicles were loaded with 100 mM NaCl and 2 mM lucigenin buffered to pH 7.2 with 20 mM sodium phosphate salts and dispersed in a 100 mM NaCl solution (buffered to pH 7.2). At  $t = 30$  s, a solution of the appropriate anion was added (final concentration of 40 mM  $\text{NaNO}_3$ , 40 mM  $\text{Na}_2\text{SO}_4$  or 40 mM NaCl). At  $t = 60$  s, a methanol solution of the transporter 9 (4 mol% carrier to lipid) was added. At the end of the experiment (300 s), detergent was added to lyse the vesicles. The blank measurement refers to the addition of  $\text{Na}_2\text{SO}_4$ , followed by the addition of methanol. Each point represents the average of three trials.

#### Direct evidence of sulfate transport by 1–9 using $^{33}\text{S}$ NMR

Even though all other possibilities apart from  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport have been excluded, there is still no direct evidence that the highly hydrophilic sulfate anion is transported across the membrane because all tests were conducted by following the movement of chloride. The majority of the biological characterisation of sulfate transporting proteins use  $^{35}\text{S}$  radio-labelled sulfate.<sup>12,29</sup> However, this method requires the separation of the internal and external solutions prior to measuring radioactive intensities, which can be quite challenging for small dilute vesicles. NMR techniques that can discriminate between the internal and external vesicular environment, without the need for separation, are therefore more suitable and practical for real-time experiments. Thus, we decided to design a method for the direct detection of sulfate transport using  $^{33}\text{S}$  NMR techniques similar to the  $^{13}\text{C}$  NMR techniques used for the detection of bicarbonate transport in Fig. 3. Due to low natural abundance of  $^{33}\text{S}$ , relatively large quadrupole moment and low sensitivity, the use of  $^{33}\text{S}$  isotopically labelled sulfate is required for this technique.  $^{33}\text{S}$  with its low gyromagnetic ratio exhibits a resonance frequency of 46.0 MHz on a 600 MHz NMR spectrometer. Initially vesicles were prepared containing  $^{33}\text{S}$  labelled sodium sulfate and they were dispersed in a sodium chloride solution. Unfortunately, it became apparent that the  $^{33}\text{S}$  signal of internal sodium sulfate was at the same  $^{33}\text{S}$  chemical shift as that observed in a reference solution of  $^{33}\text{Na}_2\text{SO}_4$  in the absence of vesicles and therefore the position of the peak cannot be used to identify the location of the sulfate anion. Shift reagents are not readily available for sulfate and therefore a different approach was adopted to circumvent this problem, once again using the relaxation properties of paramagnetic  $\text{Mn}^{2+}$ .



In a first set of tests, vesicles were prepared containing  $^{33}\text{S}$  labelled  $\text{Na}_2\text{SO}_4$  and they were dispersed in a  $\text{NaCl}$  solution containing 0.5 mol%  $\text{MnCl}_2$  (with respect to  $^{33}\text{SO}_4^{2-}$ ). A small amount of transporter was added as a  $\text{DMSO}$  solution and the mixture was allowed to incubate for 2 hours at room temperature to allow transport. The results for a selection of compounds are represented in Fig. 6. Before the addition of transporter the  $^{33}\text{S}$  NMR spectra consists of a single peak due to internal sulfate. When sulfate is transported out of the vesicle the signal is quenched due to the presence of external  $\text{Mn}^{2+}$ . As can be seen in Fig. 6, the addition of compounds 4 and 9 leads to a significant reduction in the intensity of the  $^{33}\text{S}$  NMR peak, but the peak never disappears into the baseline. The addition of detergent to lyse the vesicles and release all the encapsulated sulfate, however, does lead to a complete disappearance of the  $^{33}\text{S}$  NMR signal, indicating that the reduction in intensity was due to incomplete transport rather than insufficient  $\text{Mn}^{2+}$  to quench all the sulfate signals. These results seem to suggest that the proposed compounds are indeed able to transport sulfate. However, integration of a single peak without internal reference is not accurate in NMR spectroscopy and we therefore optimised the procedure to obtain more reliable direct evidence of sulfate transport.

In a second set of experiments, vesicles were prepared containing  $\text{Na}_2^{33}\text{SO}_4$  and  $\text{MnSO}_4$  (0.5 mol%  $\text{Mn}^{2+}$  with respect to  $^{33}\text{SO}_4^{2-}$ ) and were dispersed in a  $\text{NaCl}$  solution. This way there is no  $^{33}\text{S}$  signal prior to the addition of transporter. Addition of transporter could lead to a number of processes: (1)  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport – which would separate the sulfate anions from the  $\text{Mn}^{2+}$  ions and would lead to the appearance of a  $^{33}\text{S}$  peak; (2)  $\text{Na}^+/\text{Cl}^-$  symport – which would have no effect on the NMR spectra; (3)  $\text{Na}^+$  or  $\text{Mn}^{2+}/\text{SO}_4^{2-}$  symport – which would probably mean that the sulfate signal remains quenched due to the co-transport of manganese; (4) non-specific leaking – where both  $\text{SO}_4^{2-}$  and  $\text{Mn}^{2+}$  leak out of the vesicle and the signal remains quenched. In other words the appearance of a new  $^{33}\text{S}$  NMR peak after incubation with transporter can almost certainly only be due to  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport (or  $\text{Na}^+/\text{SO}_4^{2-}$  co-transport, which is still sulfate transport). The results for a selection of

compounds are depicted in Fig. 7 (results of the remaining compounds are in ESI†). It can be seen that the addition of compounds 3 or 4 leads to the appearance of a new  $^{33}\text{S}$  NMR signal and for these compounds all of the tests performed indicate sulfate transport. Similarly, the addition of compounds 5 or 6 also led to the appearance of a new  $^{33}\text{S}$  signal, while compounds 1, 2 and 7–9 proved to be inactive (see ESI†). Whereas most of these results are consistent with the results observed in Fig. 4, in the case of compound 9 no  $^{33}\text{S}$  NMR signal was observed, even after 12 hours incubation with the receptor, whilst all previous tests suggested sulfate transport was mediated by this receptor. Possible reasons for this observation could be the simultaneous transport of sulfate and  $\text{Mn}^{2+}$  (or leaking of the vesicles, but this possibility was already ruled out using calcein leakage assays, *vide supra*). This co-transport may occur due to  $\text{Mn}^{2+}$  and  $\text{SO}_4^{2-}$  both being doubly charged and hence in the presence of compound 9 forming a neutral molecular ensemble that is lipid soluble.

This hypothesis can be tested by repeating the  $^{33}\text{S}$  NMR experiments whilst using  $\text{Fe}^{3+}$  as the paramagnetic agent. In this case there is a mismatch between the charges of  $\text{SO}_4^{2-}$  and  $\text{Fe}^{3+}$  and co-transport is therefore less likely. To this end, lipid vesicles were prepared loaded with  $\text{Na}_2^{33}\text{SO}_4$  and  $\text{Fe}_2(\text{SO}_4)_3$  and buffered to pH 7.2 using sodium phosphate salts and dispersed in an isotonic  $\text{NaCl}$  solution. Preliminary tests showed that the addition of 5 mol%  $\text{Fe}^{3+}$  (with respect to  $^{33}\text{SO}_4^{2-}$ ) was sufficient to broaden the  $^{33}\text{S}$  NMR signal, meaning that the initial  $^{33}\text{S}$  NMR would show no signal and only the addition of a transporter capable of transporting sulfate would result in a  $^{33}\text{S}$  signal. Once detergent is added to lyse the vesicles, it is expected that the  $^{33}\text{S}$  NMR would disappear again, proving that the vesicles did not leak during the experiment. Fig. 8 shows the results of these tests where the vesicles were incubated for 2 hours with a selection of compounds (the results of the other compounds can be found in the ESI†). In accordance with the previous tests compounds 3–6 proved to be able to transport sulfate out of the vesicles, while compounds 1, 2, 7 and 8 appeared to be inactive. Receptor 9 on the other hand, which appeared to be inactive when  $\text{Mn}^{2+}$  was used as paramagnetic




Fig. 6  $^{33}\text{S}$  NMR experiments indicating that some of the receptors are able to facilitate  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport (external  $\text{Mn}^{2+}$ ). All spectra were recorded in 9 : 1 water :  $\text{D}_2\text{O}$  solutions and are shown at absolute intensity.  $^{33}\text{S}$  NMR spectra of (i) the POPC vesicles loaded with 162 mM  $\text{Na}_2^{33}\text{SO}_4$ , 20 mM phosphate buffer, pH 7.2, and dispersed in 450 mM  $\text{NaCl}$  with  $\text{MnCl}_2$  (0.5 mol%  $\text{Mn}^{2+}$  :  $^{33}\text{SO}_4^{2-}$  ratio), 20 mM phosphate buffer, pH 7.2; (ii) after 2 hour incubation with transporter 2, 3, 4, 7 or 9 (4 mol% transporter to lipid); (iii) after the addition of detergent to lyse all vesicles.






Fig. 7  $^{33}\text{S}$  NMR experiments indicating that some of the receptors are able to facilitate  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport (internal  $\text{Mn}^{2+}$ ). All spectra were recorded in 9 : 1 water :  $\text{D}_2\text{O}$  solutions and are shown at absolute intensity.  $^{33}\text{S}$  NMR spectra of (i) the POPC vesicles loaded with 162 mM  $\text{Na}_2^{33}\text{SO}_4$  with  $\text{MnSO}_4$  (0.5 mol%  $\text{Mn}^{2+}$  :  $^{33}\text{SO}_4^{2-}$  ratio), 20 mM phosphate buffer, pH 7.2, and dispersed in 450 mM NaCl, 20 mM phosphate buffer, pH 7.2; (ii) after 2 hour incubation with transporter 2, 3, 4, 7 or 9 (4 mol% transporter to lipid); (iii) after the addition of detergent to lyse all vesicles.




Fig. 8  $^{33}\text{S}$  NMR experiments indicating that some of the receptors are able to facilitate  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport (internal  $\text{Fe}^{3+}$ ). All spectra were recorded in 9 : 1 water :  $\text{D}_2\text{O}$  solutions and are shown at absolute intensity.  $^{33}\text{S}$  NMR spectra of (i) the POPC vesicles loaded with 162 mM  $\text{Na}_2^{33}\text{SO}_4$  with  $\text{Fe}_2(\text{SO}_4)_3$  (5 mol%  $\text{Fe}^{3+}$  :  $^{33}\text{SO}_4^{2-}$  ratio), 20 mM phosphate buffer, pH 7.2, and dispersed in 450 mM NaCl, 20 mM phosphate buffer, pH 7.2; (ii) after 2 hours incubation with transporter 2, 3, 4, 7 or 9 (4 mol% transporter to lipid); (iii) after the addition of detergent to lyse all vesicles.

agent, now showed evidence of sulfate transport when  $\text{Fe}^{3+}$  was used as paramagnetic agent (Fig. 8). These results suggest that compound 9 is able to transport sulfate *via* either a  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport process or a  $\text{Mn}^{2+}/\text{SO}_4^{2-}$  symport process, while compounds 3–6 mainly function *via* a  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport mechanism and compounds 1, 2, 7 and 8 are unable to transport sulfate. However, Fig. 8 also shows a potential problem when using  $\text{Fe}^{3+}$  as paramagnetic agent, namely that the  $^{33}\text{S}$  signal increases upon the addition of detergent whereas it was expected to disappear. It was also observed that the addition of detergent led to the formation of a large amount of yellow precipitate, which was identified as iron(III)phosphate using SEM-EDX (see ESI†). It is proposed that disintegration of the vesicles triggers the precipitation of  $\text{FePO}_4$  (presumably due to excess phosphate buffer upon lysis), thereby removing the majority of paramagnetic  $\text{Fe}^{3+}$  out of solution and leading to an increased  $^{33}\text{S}$  NMR signal. This is however an advantage, as it clearly shows that the  $^{33}\text{S}$  signal appearing upon addition of transporter is due to  $\text{SO}_4^{2-}$  being transported and not  $\text{Fe}^{3+}$ , because the latter would have led to the formation of a  $\text{FePO}_4$

precipitate prior to the addition of detergent. Therefore, the combination of  $^{33}\text{S}$  NMR experiments using  $\text{Mn}^{2+}$  or  $\text{Fe}^{3+}$  as paramagnetic agents is an excellent way of directly proving sulfate transport, able to discriminate between  $\text{Cl}^-/\text{SO}_4^{2-}$  antiport,  $\text{M}^{n+}/\text{SO}_4^{2-}$  symport and indiscriminate leaking of the vesicles.

Table 1 gives an overview of all the tests that were performed on compounds 1–9 and the most likely anion transport mechanisms that can be derived from this information. It can be seen that for the tren-based compounds 1–6, the transporters that are the most potent  $\text{Cl}^-/\text{NO}_3^-$  antiporters (lowest  $\text{EC}_{50,270\text{s}}$  values) are also the most potent bicarbonate and sulfate transporters. This means that these receptors are better at transporting any anion, rather than having an increased selectivity for sulfate. This suggests that the reason why these transporters are better at transporting sulfate is mainly due to the fact that they are more able to screen any hydrophilic anion from the lipid environment (increased lipophilicity, see log  $P$  values) and not because they possess increased sulfate affinities. The data in Table 1 shows that it is mainly the lipophilic receptors 3–5 that are able to



**Table 1** Overview of the various anion transport tests conducted on 1–9 and the subsequent anion transport mechanisms that can be derived from them

	C log <i>P</i>	EC <sub>50,270s</sub> <sup>a</sup> (Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> )	<sup>13</sup> C NMR <sup>b</sup> (Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> )	<sup>13</sup> C NMR <sup>c</sup> (SO <sub>4</sub> <sup>2-</sup> /HCO <sub>3</sub> <sup>-</sup> )	HPTS <sup>d</sup> (ΔpH)	Calcein <sup>e</sup> (leakage)	% Cl <sup>-</sup> efflux <sup>f</sup> (Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> )	<sup>33</sup> S NMR <sup>g</sup> (Mn <sup>2+</sup> )	<sup>33</sup> S NMR <sup>h</sup> (Fe <sup>3+</sup> )	Proposed transport mechanisms
1	2.06 <sup>i</sup>	5.60 <sup>j</sup>	Partial <sup>k</sup>	n.m. <sup>l</sup>	+0.15	No	8.8	No	No	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> antiport and minimal Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> antiport
2	5.50 <sup>i</sup>	0.31 <sup>j</sup>	Yes <sup>k</sup>	n.m. <sup>l</sup>	+0.15	No	33.4	No	No	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> and Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> antiport
3	4.43 <sup>i</sup>	0.24 <sup>j</sup>	Yes	Partial	+0.00	No	84.3	Signal	Signal	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> and HCO <sub>3</sub> <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> antiport
4	7.87 <sup>i</sup>	0.032 <sup>j</sup>	Yes	Partial	-1.00	No	96.4	Signal	Signal	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> and HCO <sub>3</sub> <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> antiport (sulfate probably transported as HSO <sub>4</sub> <sup>-</sup> )
5	7.59 <sup>i</sup>	0.0044 <sup>j</sup>	Yes	Partial	-0.80	No	63.3	Signal	Signal	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> and HCO <sub>3</sub> <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> antiport (sulfate probably transported as HSO <sub>4</sub> <sup>-</sup> )
6	11.03 <sup>i</sup>	0.042 <sup>j</sup>	Partial	Partial	+0.15	No	16.0	Signal	Signal	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> antiport and small amounts of Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> and HCO <sub>3</sub> <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> antiport
7	6.97 <sup>m</sup>	n.m. <sup>l</sup>	No	No	n.m. <sup>l</sup>	No	4.7	No	No	No transport
8	-3.62 <sup>m</sup>	3.11	No	No	+0.10	No	16.4	No	No	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> antiport
9	0.71 <sup>m</sup>	1.13	Yes	Partial	+0.00	No	80.2	No	Signal	Cl <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /HCO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> and HCO <sub>3</sub> <sup>-</sup> /SO <sub>4</sub> <sup>2-</sup> antiport and Mn <sup>2+</sup> /SO <sub>4</sub> <sup>2-</sup> symport

<sup>a</sup> EC<sub>50,270s</sub> = concentration (in mol% carrier to lipid) needed to obtain 50% chloride efflux in 270 s, obtained through Hill plot using vesicles containing NaCl dispersed in a NaNO<sub>3</sub> solution (see Fig. 1 and ESI†). <sup>b</sup> Evidence for HCO<sub>3</sub><sup>-</sup> transport in the presence of external NaCl (yes/no or partial transport) using <sup>13</sup>C NMR techniques (see Fig. 3 and ESI†). <sup>c</sup> Evidence for HCO<sub>3</sub><sup>-</sup> transport in the presence of external Na<sub>2</sub>SO<sub>4</sub> (yes/no or partial transport) using <sup>13</sup>C NMR techniques (see Fig. 3 and ESI†). <sup>d</sup> The magnitude of the intravesicular pH change upon the addition of transporter, determined using vesicles containing NaCl and HPTS and dispersed in a Na<sub>2</sub>SO<sub>4</sub> solution (see ref. 15 and ESI†). <sup>e</sup> Evidence for leakage of the vesicles (yes or no) using a calcein leakage assay (see ESI†). <sup>f</sup> Amount of chloride efflux (in %) observed 300 seconds after the addition of 4 mol% of transporter (with respect to lipid) from vesicles loaded with NaCl and dispersed in Na<sub>2</sub>SO<sub>4</sub> (see Fig. 4 and ESI†). <sup>g</sup> Evidence for sulfate transport (no or signal) using <sup>33</sup>S NMR techniques and Mn<sup>2+</sup> as paramagnetic agent (see Fig. 7 and ESI†). <sup>h</sup> Evidence for sulfate transport (no or signal) using <sup>33</sup>S NMR techniques and Fe<sup>3+</sup> as paramagnetic agent (see Fig. 8 and ESI†). <sup>i</sup> C log *P* values calculated using Spartan '08 for Macintosh (Ghose-Crippen model) (values taken from ref. 15). <sup>j</sup> EC<sub>50,270s</sub> values taken from ref. 15. <sup>k</sup> Result from <sup>13</sup>C NMR experiment taken from ref. 14. <sup>l</sup> n.m. = not measured. <sup>m</sup> C log *P* values calculated using Spartan '10 for Macintosh (Ghose-Crippen model).

transport sulfate across lipid bilayers, usually *via* an antiport mechanism (Cl<sup>-</sup>/SO<sub>4</sub><sup>2-</sup> or HCO<sub>3</sub><sup>-</sup>/SO<sub>4</sub><sup>2-</sup>), which is unsurprising as there are many reports about the importance of lipophilicity in anion transport.<sup>30</sup> It can also be seen that the data obtained using a chloride selective electrode corresponds well to the <sup>33</sup>S NMR data, *i.e.* sulfate transport can only be observed using <sup>33</sup>S NMR when the chloride efflux in the presence of external sulfate is higher than 50% efflux in 300 seconds. The only exception is compound 6, where the sulfate transport ability is lower than initially expected from log *P*. It has previously been suggested that this receptor is too lipophilic (log *P* > 11) and therefore it can get “stuck” in the inner layer of the membrane and is not able to pick up any anion from the aqueous phase or the interphase.<sup>15</sup> This is probably why the bicarbonate and sulfate transporting abilities of this receptor are impaired.

In contradistinction to these results, even though the EC<sub>50,270s</sub> values for 8–9 are much higher than for tren

compounds 1–6, compound 9 is still one of the best sulfate transporters in the series. This implies that compound 9 is more selective for transporting sulfate than tren-based compounds 3–5 that still preferentially transport the more lipophilic chloride and nitrate anions. The reason for this increased selectivity for sulfate transport is probably due to the increased selectivity for sulfate binding, due to the cage-like structure of 9 that is ideal for tetrahedral anions such as sulfate that can be bound inside the cavity *via* nine hydrogen bonds.<sup>17–19</sup> The difference in transport ability between the various cyclopeptides 7, 8 and 9 is most likely a combination between lipophilicity, anion affinity and size of the cage. Compound 7 is the most lipophilic receptor, but the large open cage results in reduced anion affinity and a reduced ability to screen the anion from the lipid environment and therefore it is unable to transport anions. The anion transport ability of cyclopeptide 8 on the other hand might be hindered by the





highly polar character of this receptor (and hence reduced membrane partitioning) and by the small cage size in which anions can be too tightly bound and cannot be released quickly once the transporter crosses the membrane. Finally, compound **9** shows the correct balance between lipophilicity, high anion affinity and a cage that is small enough to screen the anion from the membrane, but large enough to allow fast in-and-out movement of the anion during transport. Furthermore, compound **9** is the only receptor that can also function *via* a  $\text{Mn}^{2+}/\text{SO}_4^{2-}$  symport mechanism, a process that can be aided by the overall neutral complex of this ion-pair and the many metal-coordinating N and O atoms in the cyclopeptide ring. It is still possible that the activity of **9** is due to the formation of a channel or pore that is small enough to keep calcein inside but large enough to cause leakage of small ions, but the Hill coefficient of 1.1 and the difference in bicarbonate transport ability in the presence or absence of chloride (Fig. 3) and the difference in sulfate transport in the presence of  $\text{Mn}^{2+}$  or  $\text{Fe}^{3+}$  are evidence that lead us to suggest otherwise. Even though all tris-ureas and tris-thioureas **1–9** still transport chloride, nitrate and bicarbonate better than sulfate (with varying degrees of selectivity), it is a first step in developing sulfate selective anion carriers that could be used in the treatment of sulfate transport related diseases and it shows that both lipophilicity and sulfate affinity/selectivity are of the utmost importance in the transport of the highly hydrophilic sulfate anion.

## Conclusions

In conclusion, we have developed a novel method using  $^{33}\text{S}$  NMR that can be employed to directly monitor sulfate transport in vesicles. By taking advantage of the paramagnetic properties of  $\text{Mn}^{2+}$  or  $\text{Fe}^{3+}$  it is possible to determine whether [ $^{33}\text{S}$ ]sulfate is located inside or outside the vesicle and hence to establish sulfate transport. We have used this technique to show for the first time that trans-membrane sulfate transport may be mediated by small molecules based on tren and cyclic peptide scaffolds. We believe this work provides a good starting point for the development of new sulfate transporters and provides an unambiguous method to characterise their transport properties in vesicles.

## Acknowledgements

The authors thank the European Cooperation in Science and Technology (COST) action CM1005 "Supramolecular Chemistry in Water" and EU FP7 project East-NMR (grant no. 228461) for funding. PAG thanks the University of Southampton and A\*STAR for a post-graduate scholarship (NB) and the EPSRC for post-doctoral fellowships (MW & CJEH) (EP/J009687/1 and EP/G002576/1) and a studentship (LEK). PAG also thanks the Royal Society and the Wolfson Foundation for a Royal Society Wolfson Research Merit Award. DM and JP thank the Slovenian Research Agency (ARRS, P1-0242 and J1-4020). KAJ thanks the Australian Research Council (DP0877726) and the University of Sydney International Program Development Fund for financial

support. The authors would also like to thank Alistair J. Clark from the University of Southampton for his help with SEM-EDX.

## Notes and references

- 1 *Advances in Molecular and Cell Biology*, ed. E. E. Bittar and P. Michael, Elsevier, 2006, vol. 38; T. J. Jentsch, V. Stein, F. Weinreich and A. A. Zdebik, *Physiol. Rev.*, 2002, **82**, 503; F. M. Ashcroft, *Ion Channels and Disease*, Academic Press, San Diego and London, 2000.
- 2 For recent reviews: J. T. Davis, O. Okunola and R. Quesada, *Chem. Soc. Rev.*, 2010, **39**, 3843; P. R. Brotherhood and A. P. Davis, *Chem. Soc. Rev.*, 2010, **39**, 3633; P. A. Gale, *Acc. Chem. Res.*, 2011, **44**, 216; S. Matile, A. Vargas Jentzsch, J. Montenegro and A. Fin, *Chem. Soc. Rev.*, 2011, **40**, 2453.
- 3 C. Higgins, *Nature*, 1992, **358**, 536; J. Y. Choi, D. Muallem, K. Kiselyov, M. G. Lee, P. J. Thomas and S. Muallem, *Nature*, 2001, **410**, 94.
- 4 For recent reviews: N. Busschaert and P. A. Gale, *Angew. Chem., Int. Ed.*, 2013, **52**, 1374; P. A. Gale, R. Pérez-Tomás and R. Quesada, *Acc. Chem. Res.*, 2013, **46**, 2801–2813.
- 5 I. Alfonso and R. Quesada, *Chem. Sci.*, 2013, **4**, 3009; S. J. Moore, C. J. E. Haynes, J. Gonzalez, J. L. Sutton, S. J. Brooks, M. E. Light, J. Herniman, G. J. Langley, V. Soto-Cerrato, R. Pérez-Tomás, I. Marques, P. J. Costa, V. Félix and P. A. Gale, *Chem. Sci.*, 2013, **4**, 103; S. J. Moore, M. Wenzel, M. E. Light, R. Morley, S. J. Bradberry, P. Gómez-Iglesias, V. Soto-Cerrato, R. Pérez-Tomás and P. A. Gale, *Chem. Sci.*, 2012, **3**, 2501; S. Matsuyama, J. Llopis, Q. L. Deveraux, R. Y. Tsien and J. C. Reed, *Nat. Cell Biol.*, 2000, **2**, 318; J. L. Sessler, L. R. Eller, W.-S. Cho, S. Nicolaou, A. Aguilar, J. T. Lee, V. M. Lynch and D. J. Magda, *Angew. Chem., Int. Ed.*, 2005, **44**, 5989; T. Sato, H. Konno, Y. Tanaka, T. Kataoka, K. Nagai, H. H. Wasserman and S. Ohkuma, *J. Biol. Chem.*, 1998, **273**, 21455; S. Ohkuma, T. Sato, M. Okamoto, H. Matsuya, K. Arai, T. Kataoka, K. Nagai and H. H. Wasserman, *Biochem. J.*, 1998, **334**, 731.
- 6 Y. Marcus, *J. Chem. Soc., Faraday Trans.*, 1991, **87**, 2995.
- 7 S. K. Berezin and J. T. Davis, *J. Am. Chem. Soc.*, 2009, **131**, 2458; B. A. McNally, A. V. Koulov, T. N. Lambert, B. D. Smith, J.-B. Joos, A. L. Sisson, J. P. Clare, V. Sgarlata, L. W. Judd, G. Magro and A. P. Davis, *Chem.-Eur. J.*, 2008, **14**, 9599.
- 8 P. Talukdar, G. Bollot, J. Mareda, N. Sakai and S. Matile, *J. Am. Chem. Soc.*, 2005, **127**, 6528.
- 9 I. Ravikumar and P. Ghosh, *Chem. Soc. Rev.*, 2012, **41**, 3077; L. R. Eller, M. Stępień, C. J. Fowler, J. T. Lee, J. L. Sessler and B. A. Moyer, *J. Am. Chem. Soc.*, 2007, **129**, 11020; R. Custelcean, P. Remy, P. V. Bonnesen, D.-e. Jiang and B. A. Moyer, *Angew. Chem., Int. Ed.*, 2008, **47**, 1866; R. Custelcean, A. Bock and B. A. Moyer, *J. Am. Chem. Soc.*, 2010, **132**, 7177; B. Akhuli, I. Ravikumar and P. Ghosh, *Chem. Sci.*, 2012, **3**, 1522.
- 10 G. Lumetta, in *Fundamentals and Applications of Anion Separations*, ed. B. Moyer and R. Singh, Springer, USA, 2004, pp. 107–114; B. A. Moyer, R. Custelcean, B. P. Hay,



- J. L. Sessler, K. Bowman-James, V. W. Day and S.-O. Kang, *Inorg. Chem.*, 2013, **52**, 3473.
- 11 For examples: S. O. Kang, V. W. Day and K. Bowman-James, *Org. Lett.*, 2009, **11**, 3654; P. A. Gale, J. R. Hiscock, C. Z. Jie, M. B. Hursthouse and M. E. Light, *Chem. Sci.*, 2010, **1**, 215; J.-i. Kim, H. Juwarker, X. Liu, M. S. Lah and K.-S. Jeong, *Chem. Commun.*, 2010, **46**, 764.
- 12 D. Markovich, *Physiol. Rev.*, 2001, **81**, 1499.
- 13 J. Hästbacka, A. de la Chapelle, M. M. Mahtani, G. Clines, M. P. Reeve-Daly, M. Daly, B. A. Hamilton, K. Kusumi, B. Trivedi, A. Weaver, A. Coloma, M. Lovett, A. Buckler, I. Kaitila and E. S. Lander, *Cell*, 1994, **78**, 1073; A. Superti-Furga, A. Rossi, B. Steinmann and R. Gitzelmann, *Am. J. Med. Genet.*, 1996, **63**, 144.
- 14 N. Busschaert, P. A. Gale, C. J. E. Haynes, M. E. Light, S. J. Moore, C. C. Tong, J. T. Davis and W. A. Harrell, Jr, *Chem. Commun.*, 2010, **46**, 6252.
- 15 N. Busschaert, M. Wenzel, M. E. Light, P. Iglesias-Hernández, R. Pérez-Tomás and P. A. Gale, *J. Am. Chem. Soc.*, 2011, **133**, 14136.
- 16 I. Martí, J. Rubio, M. Bolte, M. I. Burguete, C. Vicent, R. Quesada, I. Alfonso and S. V. Luis, *Chem.-Eur. J.*, 2012, **18**, 16728.
- 17 P. G. Young and K. A. Jolliffe, *Org. Biomol. Chem.*, 2012, **10**, 2664.
- 18 P. G. Young, J. K. Clegg, M. Bhadbhade and K. A. Jolliffe, *Chem. Commun.*, 2011, **47**, 463.
- 19 P. G. Young, J. K. Clegg and K. A. Jolliffe, *Supramol. Chem.*, 2011, **24**, 77.
- 20 B. D. Smith and T. N. Lambert, *Chem. Commun.*, 2003, 2261.
- 21 A. V. Koulov, T. N. Lambert, R. Shukla, M. Jain, J. M. Boon, B. D. Smith, H. Li, D. N. Sheppard, J.-B. Joos, J. P. Clare and A. P. Davis, *Angew. Chem., Int. Ed.*, 2003, **42**, 4931.
- 22 A. V. Hill, *Biochem. J.*, 1913, **7**, 471.
- 23 T. P. W. McMullen, R. N. A. H. Lewis and R. N. McElhaney, *Curr. Opin. Colloid InterfaceSci.*, 2004, **8**, 459;
- J. C. M. Holthuis, G. van Meer and K. Huitema, *Mol. Membr. Biol.*, 2003, **20**, 231; W. F. D. Bennett, J. L. MacCallum and D. P. Tieleman, *J. Am. Chem. Soc.*, 2009, **131**, 1972.
- 24 S. Bhosale and S. Matile, *Chirality*, 2006, **18**, 849.
- 25 J. T. Davis, P. A. Gale, O. A. Okunola, P. Prados, J. C. Iglesias-Sánchez, T. Torroba and R. Quesada, *Nat. Chem.*, 2009, **1**, 138.
- 26 R. R. C. New, in *Liposomes: a Practical Approach*; IRL Press, Oxford, 1990, pp. 131–134; V. Sidorov, F. W. Kotch, G. Abdrakhmanova, R. Mizani, J. C. Fettinger and J. T. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 2267; T. N. Lambert, J. Middleton Boon, B. D. Smith, M. Nieves Perez-Payan and A. P. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 5276; S. Licen, C. Coppola, J. D'Onofrio, D. Montesarchio and P. Tecilla, *Org. Biomol. Chem.*, 2009, **7**, 1060.
- 27 K. Kano and J. H. Fendler, *Biochim. Biophys. Acta*, 1978, **509**, 289; N. R. Clement and J. M. Gould, *Biochemistry*, 1981, **20**, 1534.
- 28 J. Biwersi, B. Tulk and A. S. Verkman, *Anal. Biochem.*, 1994, **219**, 139.
- 29 E. G. Schneider, J. C. Durham and B. Sacktor, *J. Biol. Chem.*, 1984, **259**, 14591; J. B. Pritchard and J. L. Renfro, *Proc. Natl. Acad. Sci. U. S. A.*, 1983, **80**, 2603; H. Lohi, M. Kujala, S. Mäkelä, E. Lehtonen, M. Kestilä, U. Saarialho-Kere, D. Markovich and J. Kere, *J. Biol. Chem.*, 2002, **277**, 14246.
- 30 C. J. E. Haynes, S. J. Moore, J. R. Hiscock, I. Marques, P. J. Costa, V. Félix and P. A. Gale, *Chem. Sci.*, 2012, **3**, 1436; V. Saggiomo, S. Otto, I. Marques, V. Felix, T. Torroba and R. Quesada, *Chem. Commun.*, 2012, **48**, 5274; N. Busschaert, S. J. Bradberry, M. Wenzel, C. J. E. Haynes, J. R. Hiscock, I. L. Kirby, L. E. Karagiannidis, S. J. Moore, N. J. Wells, J. Herniman, G. J. Langley, P. N. Horton, M. E. Light, I. Marques, P. J. Costa, V. Felix, J. G. Frey and P. A. Gale, *Chem. Sci.*, 2013, **4**, 3036.

