


NATL INST OF STAND & TECH



A11107 223934

11-20-57 ✓

REFERENCE

NSRDS-NBS 9



Tables of Bimolecular Gas Reactions

U.S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS



R7503299

R7503299

Tables of Bimolecular Gas Reactions

A. F. Trotman-Dickenson* and G. S. Milne*

*Prepared under contract at the
Edward Davies Chemical Laboratory
Aberystwyth, Wales



NSRDS-NBS 9
National Standard Reference Data Series-
National Bureau of Standards 9

(Category 6—Chemical Kinetics)

Issued October 27, 1967

Library of Congress Catalog Card Number: 67-60026

Foreword

The National Standard Reference Data System is a government-wide effort to give to the technical community of the United States optimum access to the quantitative data of physical science, critically evaluated and compiled for convenience. This program was established in 1963 by the President's Office of Science and Technology, acting upon the recommendation of the Federal Council for Science and Technology. The National Bureau of Standards has been assigned responsibility for administering the effort. The general objective of the System is to coordinate and integrate existing data evaluation and compilation activities into a systematic, comprehensive program, supplementing and expanding technical coverage when necessary, establishing and maintaining standards for the output of the participating groups, and providing mechanisms for the dissemination of the output as required.

The NSRDS is conducted as a decentralized operation of nation-wide scope with central coordination by NBS. It comprises a complex of data centers and other activities, carried on in government agencies, academic institutions, and nongovernmental laboratories. The independent operational status of existing critical data projects is maintained and encouraged. Data centers that are components of the NSRDS produce compilations of critically evaluated data, critical reviews of the state of quantitative knowledge in specialized areas, and computations of useful functions derived from standard reference data.

For operational purposes, NSRDS compilation activities are organized into seven categories as listed below. The data publications of the NSRDS, which may consist of monographs, loose-leaf sheets, computer tapes, or any other useful product, will be classified as belonging to one or another of these categories. An additional "General" category of NSRDS publications will include reports on detailed classification schemes, lists of compilations considered to be Standard Reference Data, status reports, and similar material. Thus, NSRDS publications will appear in the following eight categories:

| <i>Category</i> | <i>Title</i> |
|-----------------|--|
| 1 | General |
| 2 | Nuclear Properties |
| 3 | Atomic and Molecular Properties |
| 4 | Solid State Properties |
| 5 | Thermodynamic and Transport Properties |
| 6 | Chemical Kinetics |
| 7 | Colloid and Surface Properties |
| 8 | Mechanical Properties of Materials |

Within the National Bureau of Standards publication program a new series has been established, called the National Standard Reference Data Series. The present report, which is in Category 6 of the above list, is Number 9 of the new series and is designated NSRDS-NBS 9.

A. V. ASTIN, *Director.*

Contents

| | Page |
|---|------|
| Foreword..... | iii |
| Preface..... | 1 |
| Notes on the tables..... | 1 |
| Tables | |
| Atom reactions | |
| Hydrogen atoms (H)..... | 4 |
| Fluorine atoms (F)..... | 15 |
| Chlorine atoms (Cl)..... | 16 |
| Bromine atoms (Br)..... | 22 |
| Iodine atoms (I)..... | 25 |
| Sulphur atoms (S)..... | 27 |
| Sodium atoms (Na)..... | 28 |
| Potassium atoms (K)..... | 37 |
| Radicals containing one carbon atom | |
| Methyne (CH)..... | 38 |
| Methylene (CH ₂)..... | 39 |
| Methyl (CH ₃)..... | 40 |
| Difluoromethylene (CF ₂)..... | 56 |
| Trifluoromethyl (CF ₃)..... | 57 |
| Misc. halogenated methyls (CFH ₂ , CF ₂ Cl, CCl ₃)..... | 62 |
| Formyl (HCO)..... | 64 |
| Methoxy (CH ₃ O)..... | 65 |
| Methylthio (CH ₃ S)..... | 67 |
| Chloroformyl (ClCO)..... | 68 |
| Cyano (CN)..... | 69 |
| Radicals containing two carbon atoms | |
| Ethynyl (C ₂ H)..... | 70 |
| Ethyl (C ₂ H ₅)..... | 71 |
| Perfluoroethyl (C ₂ F ₅)..... | 75 |
| Misc. halogenated ethyls (C ₂ H ₄ Cl, C ₂ H ₃ Cl ₂ , C ₂ H ₂ Cl ₃ , C ₂ HCl ₄ , C ₂ Cl ₅ , C ₂ F ₄ Cl, C ₂ F ₃ Cl ₂ , C ₂ F ₂ Cl ₃ , C ₂ F ₄ Br, C ₂ H ₄ Br)..... | 76 |
| Carbonylcarbene (CCO)..... | 78 |
| Acetyl and trifluoroacetyl (CH ₃ CO, CF ₃ CO)..... | 79 |
| Ethoxy (C ₂ H ₅ O)..... | 80 |
| Radicals containing three carbon atoms | |
| Allyl (CH ₂ CHCH ₂)..... | 81 |
| <i>n</i> -Propyl (CH ₃ CH ₂ CH ₂)..... | 82 |
| <i>iso</i> -Propyl (CH ₃ CHCH ₃)..... | 83 |
| Perfluoropropyl (CF ₃ CF ₂ CF ₂)..... | 84 |
| Misc. halogenated propyls (CCl ₃ CH ₂ CH ₂ , ICH ₂ CH ₂ CH ₂)..... | 85 |
| Isopropoxy ((CH ₃) ₂ CHO)..... | 86 |
| Radicals containing four carbon atoms | |
| Butyls (CH ₃ CH ₂ CH ₂ CH ₂ , CH ₃ CHCH ₂ CH ₃ , (CH ₃) ₂ CHCH ₂ , (CH ₃) ₃ C)..... | 87 |
| <i>t</i> -Butoxy ((CH ₃) ₃ CO)..... | 88 |

| | Page |
|---|------|
| Miscellaneous carbon-containing radicals | |
| Peracid radicals (CH_3CO_3 , $\text{C}_2\text{H}_5\text{CO}_3$)..... | 89 |
| Peroxy radicals ($(\text{CH}_3)_3\text{CO}_2$, cyclo- $\text{C}_6\text{H}_9\text{O}_2$, $\text{C}_6\text{H}_5\text{C}(\text{CH}_3)_2\text{O}_2$)..... | 90 |
| Phenyl radicals (C_6H_5)..... | 91 |
| Tolyl ($\text{CH}_3\text{C}_6\text{H}_4$)..... | 92 |
| Inorganic radicals | |
| Hydroxyl (OH)..... | 93 |
| Hydroperoxyl (HO_2)..... | 96 |
| Imino (NH)..... | 97 |
| Amino (NH_2)..... | 98 |
| Difluoroamino (NF_2)..... | 99 |
| Nitrate (NO_3)..... | 100 |
| Thiyl (HS)..... | 101 |
| Chloromonoxy (ClO)..... | 102 |
| Borine (BH_3)..... | 103 |
| Fluosulfates (F_3SO , FSO_3)..... | 104 |
| Radical–radical reactions | |
| Radical disproportionation..... | 105 |
| Disproportionation/combination ratios..... | 108 |
| Molecule–molecule reactions | |
| Reactions involving hydrogen and hydrogen halides..... | 118 |
| Reactions involving organic radicals..... | 120 |
| Reactions involving nitric oxide..... | 122 |
| Reactions involving nitrogen dioxide..... | 124 |
| Misc. bimolecular molecule–molecule reactions..... | 126 |
| Odds and ends | |
| Termolecular reactions..... | 127 |
| Inversion (group transfer) reactions..... | 128 |

Tables of Bimolecular Gas Reactions

A. F. Trotman-Dickenson and G. S. Milne

This survey covers the kinetics of bimolecular and termolecular gas reactions that do not involve atoms or molecules in electronically excited states. Bimolecular reactions are here defined as reactions in which two molecules are involved as reactants, that yield two or more molecules as products. Those reactions in which two molecules combine to form one molecule are most usefully considered as the reverse of unimolecular reactions which will be dealt with in another survey. Reactions of oxygen and nitrogen atoms have been omitted as they will also form the subject of another survey.

The literature from 1954 to December 31, 1965 has been exhaustively searched and it is hoped that for this period nothing has been omitted that should have been included.

The survey of earlier work has been based on one of the writers' books on "Gas Kinetics" which covered the literature to 1954. Use of the book for over ten years has revealed few omissions and these have been included in these tables. Data for the period January to August 1966 has been included where possible.

Key Words: Chemical kinetics, gas, bimolecular, reactions, tables, rate constants, activation energies, Arrhenius equation, data.

Preface

This survey covers the kinetics of bimolecular and termolecular gas reactions that do not involve atoms or molecules in electronically excited states. Bimolecular reactions are here defined as reactions in which two molecules are involved as reactants, that yield two or more molecules as products. Those reactions in which two molecules combine to form one molecule are most usefully considered as the reverse of unimolecular reactions which will be dealt with in another survey. Reactions of oxygen and nitrogen atoms have been omitted as they will also form the subject of another survey.

The literature from 1954 to December 31, 1965 has been exhaustively searched and it is hoped that for this period nothing has been omitted that should have been included.

The survey of earlier work has been based on one of the writers' books on "Gas Kinetics" which covered the literature to 1954. Use of the book for over ten years has revealed few omissions and these have been included in these tables. Data for the period January to August 1966 has been included where possible.

Originally it was hoped to make this survey a critical study from which reliable data could be distilled. As the work proceeded it became evident that this was not a realistic goal. Only a small fraction of the reactions has been the subject of investigation by several workers. When several studies have been made the results fall into two classes. In the first class are those reactions that have been repeatedly investigated. It is then usually found that the discrepancies are greater than the random errors and that some systematic differences exist. It is not possible to decide which value is most

nearly correct. In the second class are those reactions that have been investigated under widely different conditions. In these cases the best result for one purpose may not be the best for another, so that no useful choice can be made. We therefore decided to make the tables as comprehensive as possible so that users can readily discover what information is available.

It is likely that some reactions have been omitted and that other errors have crept in. We shall be grateful if readers will write to tell us of suggested improvements. It is hoped to publish supplements to these tables every two or three years.

Notes on the Tables

All the rate data is in the simple Arrhenius form $k = Ae^{-E/RT}$. "A" factors and rate constants are in cm. mole. sec. units throughout. Activation energies are in kcal/mole and temperatures in degrees centigrade.

Where data has been represented by equations of the type

$$k = BT^n e^{-H/RT} \quad \left(n = 1, \frac{1}{2}, \frac{1}{4} \right),$$


B and H have been converted to the simple form by means of the equations

$$A = e^n BT^n \text{ and } E = H + nRT,$$

where T is the absolute temperature at the mid-point of the experimental range.

Where the rate equation is reported as a more complex function of temperature, it has been quoted as such in the tables.

Most of the data is for simple radical/molecule metathetical reactions of the type



where R_1 is an atom or radical attacking a molecule R_2X . X is the atom transferred in the reaction.

For each radical R_1 there is a separate table (see contents). The first column of each table consists of a list of molecules R_2X with which R_1 reacts. Columns three to five contain the Arrhenius values and the experimental temperature ranges. Column six gives an indication of the nature of the experimental method used to obtain the data.

The following abbreviations have been used in describing the radical source:

- P photolysis.
- F.P. flash photolysis
- T thermal reaction
- S.T. shock tube
- D.F. diffusion flame
- M.D.F. modified diffusion flame
- M.L. mean life method
- Dil. F. dilute flame
- M.B. molecular beam
- + H Hg photosensitized addition of H atoms to olefines.
- H Hg photosensitized removal of H atoms from saturated compounds.
- + CH₃ CH₃ radical attack on the parent compound.


The other radical source abbreviations used, such as H₂/discharge and H₂/O₂ flame, are self explanatory.

The table for each attacking species R_1 is split into six sections defined by the atom X being transferred. In the order used in the tables these

- are: (i) hydrogen atom transfer
- (ii) fluorine atom transfer
- (iii) chlorine atom transfer
- (iv) bromine atom transfer
- (v) iodine atom transfer
- (vi) oxygen atom transfer

Only in the case of methyl radicals are all six categories used.

In the few cases not covered by this classification e.g., transfer of an atom from the radical to the molecule as in



either the reaction is written in full, or the products of the reaction are indicated in a footnote.

A seventh category of reactions not already covered (e.g., group transfer and inversion reactions) is included at the end of each table.

Simple ratios of rate constants (of the form k_1/k_2) are presented in separate tables, these proved much more difficult to classify, and wherever necessary, explanations of the construction of these tables are included at the head of the table.

Rate data is available in the literature as complex functions of several rate constants: this data has not been included, but references as to where it is available are given at the appropriate sections in the tables.

Where data is considered to be doubtful this is indicated in a footnote. As far as possible, an attempt has been made to include criticism of data by cross reference. Thus, where several references are given for one set of Arrhenius values, the first reference contains the values themselves; the other references indicate the sources of supplementary data and/or comment on the values shown.

Where a sufficient quantity of data has been collected for any one reaction, recommended Arrhenius values are given. They are compatible in all cases with the values for the reverse reaction (where these are known).

List of Tables

| 1. Atom Reactions | Page |
|---|------|
| 1. Hydrogen atoms..... H..... | 4 |
| 2. Fluorine atoms..... F..... | 15 |
| 3. Chlorine atoms..... Cl..... | 16 |
| 4. Bromine atoms..... Br..... | 22 |
| 5. Iodine atoms..... I..... | 25 |
| 6. Sulphur atoms..... S..... | 27 |
| 7. Sodium atoms..... Na..... | 28 |
| 8. Potassium atoms..... K..... | 37 |
| | |
| 2. Radicals Containing One Carbon Atom | |
| 1. Methyne..... CH..... | 38 |
| 2. Methylene..... CH ₂ | 39 |
| 3. Methyl..... CH ₃ | 40 |
| 4. Difluoromethylene..... CF ₂ | 56 |

List of Tables – Continued

| | Page |
|---|------|
| 2. Radicals Containing One Carbon Atom – Continued | |
| 5. Trifluoromethyl..... | 57 |
| 6. Misc. halogenated methyls..... | 62 |
| 7. Formyl..... | 64 |
| 8. Methoxy..... | 65 |
| 9. Methylthio..... | 67 |
| 10. Chloroformyl..... | 68 |
| 11. Cyano..... | 69 |
| 3. Radicals Containing Two Carbon Atoms | |
| 1. Ethynyl..... | 70 |
| 2. Ethyl..... | 71 |
| 3. Perfluoroethyl..... | 75 |
| 4. Misc. halogenated ethyls..... | 76 |
| 5. Carbonylcarbene..... | 78 |
| 6. Acetyl and trifluoroacetyl..... | 79 |
| 7. Ethoxy..... | 80 |
| 4. Radicals Containing Three Carbon Atoms | |
| 1. Allyl..... | 81 |
| 2. <i>n</i> -propyl..... | 82 |
| 3. <i>iso</i> -propyl..... | 83 |
| 4. Perfluoropropyl..... | 84 |
| 5. Misc. halogenated propyls..... | 85 |
| 6. Isopropoxy..... | 86 |
| 5. Radicals Containing Four Carbon Atoms | |
| 1. Butyls..... | 87 |
| 2. <i>t</i> -Butoxy..... | 88 |
| 6. Miscellaneous Carbon-Containing Radicals | |
| 1. Peracid radicals..... | 89 |
| 2. Peroxy radicals..... | 90 |
| 3. Phenyl radicals..... | 91 |
| 4. Tolyl..... | 92 |
| 7. Inorganic Radicals | |
| 1. Hydroxyl..... | 93 |
| 2. Hydroperoxyl..... | 96 |
| 3. Imino..... | 97 |
| 4. Amino..... | 98 |
| 5. Difluoroamino..... | 99 |
| 6. Nitrate..... | 100 |
| 7. Thiyl..... | 101 |
| 8. Chloromonoxo..... | 102 |
| 9. Borine..... | 103 |
| 10. Fluorosulfates..... | 104 |
| 8. Radical-Radical Reactions | |
| 1. Radical disproportionation..... | 105 |
| 2. Disproportionation/combination ratios..... | 108 |
| 9. Molecule-Molecule Reactions | |
| 1. Reactions involving hydrogen and hydrogen halides..... | 118 |
| 2. Reactions involving organic radicals..... | 120 |
| 3. Reactions involving nitric oxide..... | 122 |
| 4. Reactions involving nitrogen dioxide..... | 124 |
| 5. Miscellaneous bimolecular molecule-molecule reactions..... | 126 |
| 10. Odds and Ends | |
| 1. Termolecular reactions..... | 127 |
| 2. Inversion (group transfer) reactions..... | 128 |

Reaction of Hydrogen Atoms

Index

| | Page |
|---|------|
| (a) Hydrogen atom transfer | |
| Reaction with | |
| 1. Hydrogen..... | 5 |
| 2. Alkanes..... | 5 |
| 3. Cyclo-alkanes..... | 6 |
| 4. Aromatic hydrocarbons..... | 6 |
| 5. Aldehydes and ketones..... | 6 |
| 6. Metal alkyl..... | 7 |
| 7. Halogenated alkanes..... | 7 |
| 8. Hydrogen halides..... | 8 |
| 9. Hydrides of oxygen and nitrogen..... | 8 |
| (b) Fluorine atom transfer..... | 8 |
| (c) Chlorine atom transfer..... | 8 |
| (d) Bromine atom transfer..... | 9 |
| (e) Iodine atom transfer..... | 9 |
| (f) Oxygen atom transfer..... | 9 |
| (g) Ratios of rate constants | |
| 1. Reactions of the type (1) $H + RH = H_2 + R$ (2) $H + Ol = OlH$ where RH is any hydrogen containing compound and Ol is an olefine..... | 10 |
| 2. Reactions with hydrocarbons and hydrocarbon derivatives..... | 11 |
| 3. Reactions of the type (1) $H + HX = H_2 + X$ (2) $H + X_2 = HX + X$ where X is a halogen atom..... | 12 |
| 4. Miscellaneous ratios..... | 12 |

Metathetical Reactions of Hydrogen Atoms With Hydrogen

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference | |
|--------------------|---------|--|--|-------------------|---------------------------------|---------------------------------|----|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | | |
| H + H ₂ | (a) | 6.68 | 13.78 | | | 1 | |
| | (b) | 7.5 ± 1 | 13.7 | 10-730 | | 2 | |
| | (d) (c) | 8.0 ± 0.5 | 14.05 | 10-730 | | 8 | |
| | | $\log_{10} k = 15.45 - 3.49 \times 10^3/T + 3.84 \times 10^5/T^2$ | | | 27-171 | H ₂ /tungsten spiral | 11 |
| | (e) (f) | $k = 1.18 \times 10^{15} T^{-1/2} \Gamma \exp(-9.21 \times 10^3/RT)$ | | | 27-171 | H ₂ /tungsten spiral | 12 |
| H + HD | (a) | 6.96 | 13.12 | | | 1 | |
| H + HD | (a) | 7.04 | 13.06 | | | 1 | |
| H + D ₂ | (a) | 7.48 | 13.32 | | | 1 | |
| | | 7.30 ± 0.1 | 12.64 ± 0.05 | 95-195 | H ₂ /tungsten spiral | 13 | |
| D + H ₂ | (a) | 6.14 | 13.37 | | | 1 | |
| | (f) | $k = 3.63 \times 10^{15} T^{-1/2} \Gamma \exp(-9.4J \times 10^3/RT)$ | | 1-195 | D ₂ /tungsten spiral | 12 | |
| D + HD | (a) | 6.40 | 12.99 | | | 1 | |
| D + HD | (a) | 6.48 | 12.93 | | | 1 | |
| D + D ₂ | (a) | 6.09 | 13.49 | | | 1 | |

Notes

- (a) A self-consistent set of values derived from absolute rate calculations.
 (b) Review of literature data (refs. 3 to 6).
 (c) Review of literature data (refs. 3 to 7).
 (d) For a critical summary of the (then) available experimental data, see refs. 9 and 10.
 (e) Recalculation of the data from ref. 11.
 (f) Γ is the tunneling factor.

References

1. Polanyi, J. Chem. Phys. **23**, 1505 (1955).

2. Trotman-Dickenson, "Gas Kinetics," p. 169 et seq. (Butterworths, London, 1955).
 3. Farkas and Farkas, Proc. Roy. Soc. **152A**, 124 (1935).
 4. Geib and Harteck, Z. Phys. Chem. Bodenstein Festband **849** (1931).
 5. Melville and Robb, Proc. Roy. Soc. **196A**, 445 (1949).
 6. Van Meersche, Bull. Soc. Chim. Belg. **60**, 99 (1951).
 7. Boata, Careri, Cimino, Molinari, and Volpi, J. Chem. Phys. **24**, 783 (1956).
 8. Weston, J. Chem. Phys. **31**, 892 (1959).
 9. Shavitt, J. Chem. Phys. **31**, 1359 (1959).
 10. Rapp and Weston, J. Chem. Phys. **36**, 2807 (1962).
 11. Schulz and Le Roy, J. Chem. Phys. **42**, 3869 (1965).
 12. Ridley, Schulz, and Le Roy, J. Chem. Phys. **44**, 3344 (1966).
 13. Schulz and Le Roy, Can. J. Chem. **42**, 2480 (1964).

Metathetical Reactions of Hydrogen Atoms

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|-----------|-------|---------------------------------|--|-------------------|----------------|-----------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |

HYDROGEN ATOM TRANSFER

| | | | | | | |
|-----------------|-----|-----------|-------------|---------|-----------------------------------|---|
| Alkanes | | | | | | |
| CH ₄ | (a) | 7.0 ± 1.5 | | 130-290 | | 1 |
| | | 6.6 | 10.5 | 99-163 | H ₂ /tungsten filament | 2 |
| | | 4.5 ± 1.2 | 10.01 ± 0.2 | 99-163 | H ₂ /tungsten filament | 3 |

Metathetical Reactions of Hydrogen Atoms—Continued

| Reactants | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|---------|-----------------------------------|--|-------------------|---|-----------|
| | | (<i>kcal mole⁻¹</i>) | (<i>cm³ mole⁻¹ sec⁻¹</i>) | °C | | |
| | (a) | 9.0 | 12.5 | 130–420 | | 4 |
| | (a) | 8.0 | 12.6 | 477–547 | | 5 |
| | | 11.5 | 14.3 | 950–1530 | CH ₄ /O ₂ flame | 6 |
| | | 15.1 | 14.53 | 400–480 | H ₂ /O ₂ ignition | 7 |
| | (b) | 11.05 | 14.19 | | | 8 |
| | | | $k = 2.9 \times 10^{10}$ | 610 | H ₂ /O ₂ ignition | 9 |
| | | 7.4 ± 1.1 | 11.8 | 212–530 | H ₂ /discharge | 10 |
| | | 8.0 | 13.9 | 697–863 | H ₂ O ₂ ignition | 11 |
| | (w) (q) | 13.5 | | 450 | D ₂ /discharge | 12 |
| | (w) (c) | 7.8 | 13.0 | 347–465 | D ₂ CO photolysis | 13 |
| | (r) | 9.6 | 12.6 | | | |
| C ₂ H ₂ | (d) | 1.5 | | | | 14 |
| C ₂ H ₄ | (e) | 6.6 ± 1 | 13.26 | 500–625 | H ₂ /O ₂ ignition | 15 |
| C ₂ H ₆ | | 6.8 | 12.53 | 80–163 | H ₂ /tungsten filament | 16 |
| | | 12.0 | 13.17 | 410–550 | H ₂ O ₂ ignition | 17 |
| | (e) | 14 ± 1 | 14.76 | 500–625 | H ₂ /O ₂ ignition | 15 |
| | (q) | 9.0 ± 0.2 | | 24 ± 2 | H ₂ /discharge | 18 |
| | | 12.2 | 14.52 | 410–550 | H ₂ /O ₂ ignition | 7 |
| | (f) | 9.9 ± 1.0 | 14.20 ± 0.4 | 25–1230 | | 19 |
| | | 9.7 ± 2 | 14.1 | 720–1160 | C ₂ H ₆ /H ₂ /O ₂ flame | 105 |
| | (g) | | $k = 3.0 \times 10^9$ | 630 | | 20 |
| | (h) | 9.0 | 13.5 | 50–250 | C ₂ H ₆ /C ₃ H ₆ radiolysis | 21 |
| | (f) | 9.71 ± 0.58 | 14.12 ± 0.25 | 31–1227 | | 22 |
| | | 6.20 | 13.6 | 752–917 | H ₂ /O ₂ ignition | 11 |
| | (w) (i) | 9.0 | 14.4 | 30–306 | D ₂ S/C ₂ H ₆ photolysis | 23 |
| C ₃ H ₆ | (e) | 1.1 ± 1 | 12.45 | 500–625 | H ₂ /O ₂ ignition | 15 |
| | (w) (i) | 5.0 | 13.7 | 25–205 | D ₂ S + C ₃ H ₆ photolysis | 23 |
| C ₃ H ₈ | | 8.5 | 13.43 | 410–550 | H ₂ /O ₂ ignition | 17 |
| | (e) | 8.3 ± 1 | 13.56 | 500–625 | H ₂ /O ₂ ignition | 15 |
| | | 8.8 | 13.80 | 410–550 | H ₂ /O ₂ ignition | 7 |
| | (h) | 7.4 | 13.7 | 57–217 | C ₃ H ₈ radiolysis | 24 |
| | | 8.2 | 14.12 | 95–170 | H ₂ /tungsten filament | 25 |
| | (j) | | $k = 4.53 \times 10^{11}$ | 520 | H ₂ /O ₂ ignition | 26 |
| | (f) | 7.83 ± 0.79 | 13.81 ± 0.37 | 28–520 | | 26 |
| | (w) (q) | 8.0 | | Room temp. | H ₂ /discharge | 27 |
| | (w) (i) | 7.2 | 14.5 | 30–250 | D ₂ S + C ₃ H ₈ photolysis | 23 |
| CH ₃ CH:CHCH ₃ | (w) (i) | 5.0 | 14.1 | 28–230 | D ₂ S + C ₄ H ₈ photolysis | 23 |
| <i>n</i> -C ₄ H ₁₀ | (q) | 8.9 | | Room temp. | H ₂ /discharge | 27 |
| | (j) | | $k = 5.1 \times 10^{11}$ | 520 | H ₂ /O ₂ ignition | 28 |
| | (f) | 7.5 ± 0.5 | 13.9 ± 0.2 | 27–520 | | 28 |
| | (h) | 6.7 | 13.3 | 50–250 | <i>n</i> -C ₄ H ₁₀ radiolysis | 21 |
| | (k) | 7.1 | 13.82 | 70–170 | H ₂ /tungsten filament | 29 |
| | (w) (q) | 7.9 | | Room temp. | H ₂ /discharge | 27 |
| | (w) (i) | 7.1 | 14.5 | 30–250 | D ₂ S + C ₄ H ₁₀ photolysis | 23 |
| <i>i</i> -C ₄ H ₁₀ | (q) | 9.3 | | | H ₂ /discharge | 30 |
| | | 6.0 | 13.47 | 410–550 | H ₂ /O ₂ ignition | 17 |
| | (e) | 5.6 ± 1 | 12.89 | 500–625 | H ₂ /O ₂ ignition | 15 |
| | | 5.1 | 12.76 | 410–510 | H ₂ /O ₂ ignition | 7 |
| | (h) | 5.1 | 13.1 | 50–250 | <i>i</i> -C ₄ H ₁₀ radiolysis | 21 |
| | (j) | | $k = 9.2 \times 10^{11}$ | 520 | H ₂ /O ₂ ignition | 28 |

Metathetical Reactions of Hydrogen Atoms – Continued

| Reactants | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|------------------------------|---------|---------------------|------------------------------------|-------------------|----------------------------------|-----------|
| | | $(kcal\ mole^{-1})$ | $(cm^3\ mole^{-1}\ sec^{-1})$ | $^{\circ}C$ | | |
| | (f) | 6.8 ± 0.5 | 14.0 ± 0.2 | 27–520 | | 28 |
| | (w) (i) | 6.3 | 14.5 | 30–350 | $D_2S + C_4H_{10}$ photolysis | 23 |
| $n-C_3H_{12}$ | (q) | 8.5 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | 7.8 | | Room temp. | D_2 /discharge | 27 |
| neo- C_5H_{12} | (q) | 9.2 | | Room temp. | H_2 /discharge | 31 |
| $n-C_6H_{14}$ | (q) | 9.1 | | Room temp. | H_2 /discharge | 27 |
| | (f) | 5.9 | | 25–305 | | 32 |
| | (w) (q) | 8.0 | | Room temp. | D_2 /discharge | 27 |
| Cyclo Alkanes | | | | | | |
| cyclo- C_3H_6 | (q) | 9.5 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | 9.3 | | Room temp. | D_2 /discharge | 27 |
| cyclo- C_4H_8 | (q) | 8.2 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | 7.7 | | Room temp. | D_2 /discharge | 27 |
| cyclo- C_6H_{10} | (q) | 7.5 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | 6.6 | | Room temp. | D_2 /discharge | 27 |
| cyclo- C_6H_{12} | (q) | 8.0 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | 7.2 | | Room temp. | D_2 /discharge | 27 |
| Aromatic hydrocarbons | | | | | | |
| C_6H_6 | (q) | < 7 | | Room temp. | H_2 /discharge | 27 |
| | (w) (q) | < 6 | | Room temp. | D_2 /discharge | 27 |
| Aldehydes and ketones | | | | | | |
| HCHO | (l) | 2.6 | 13.21 | 252–397 | H_2CO/D_2 photolysis | 33 |
| | (f) | 3.49 | 13.46 | 327–727 | | 34 |
| | (f) | 3.26 | 13.29 | 250–400 | | 34 |
| | | | $k = (2.7 \pm 0.7) \times 10^{10}$ | 27 | H_2 /discharge | 35 |
| | | | $k = 3.5 \times 10^{12}$ | 547 | HCHO pyrolysis | 36 |
| | (w) (l) | 2.7 | 13.21 | 252–397 | H_2CO/D_2 photolysis | 33 |
| DCDO | (l) | 3.6 | 13.21 | 252–397 | H_2CO/D_2CO photolysis | 33 |
| | (w) (m) | 3.0 | 13.3 | 308–341 | D_2CO photolysis | 13 |
| | (w) (l) | 3.6 | 13.21 | 252–397 | H_2CO/D_2CO photolysis | 33 |
| CH_3CHO | (q) | ≤ 6 | | 25 | H_2 /discharge | 37 |
| CH_3COCH_3 | (q) | 8.8 | | Room temp. | H_2 /discharge | 38 |
| | (w) (q) | 8.0 | | Room temp. | D_2 /discharge | 38 |
| Metal alkyl | | | | | | |
| CH_3HgCH_3 | (q) | ≤ 6 | | 25 | H_2 /discharge | 106 |
| Halogenated methane | | | | | | |
| CF_3H | (a) | 5.0 | 12.7 | 832–1011 | H_2/O_2 ignition | 11 |

Metathetical Reactions of Hydrogen Atoms – Continued

| Reactants | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|---------|-------------------------------|---|-------------------|--|-----------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| Hydrogen halides | | | | | | |
| H F | (n) | 35.0 | 13.00 | 3527–5027 | HF shock wave decomposition | 39 |
| H Cl | | 5.2 | 13.4 | 628–798 | H ₂ /HCl thermal reaction | 40 |
| | (a) | 4.5 | | | | 41 |
| H Br | | 2.2 | 13.79 | 832–1011 | H ₂ /O ₂ ignition | 11 |
| | | 0.9 | 13.1 | 548–711 | H ₂ /HBr thermal reaction | 42 |
| | (o) | 3.7 | 14.04 | 700–1400 | | 43 |
| | (r) | 2.9 | 13.8 | | | |
| HI | | 0.7 ± 0.25 | 13.70 ± 0.07 | 394–527 | H ₂ /I ₂ thermal reaction | 44 |
| | | 1.17 ± 0.35 | 13.8 ± 0.2 | 394–465 | H ₂ /I ₂ thermal reaction | 107 |
| Hydrides of oxygen and nitrogen | | | | | | |
| H ₂ O | | $k = 6.6 \pm 0.9 \times 10^9$ | | 799 | H ₂ /O ₂ /N ₂ O flame | 45 |
| | (a) | 20.4 | 13.96 | 37–642 | | 108 |
| | (a) | 21.6 | 14.84 | 687–807 | H ₂ /O ₂ ignition | 46 |
| | (f) | 21.1 ± 1.5 | 14.48 ± 1.0 | 27–1727 | | 47 |
| D ₂ O | (a) | 25.5 | 15 | 1012–1227 | | 109 |
| | | $k = 5.0 \times 10^9$ | | 799 | H ₂ /O ₂ flame | 48 |
| | | $k = 3.6 \times 10^9$ | | 799 | H ₂ /O ₂ flame | 49 |
| | (p) (f) | 21.8 | 13.93 | 799 | H ₂ /O ₂ flame | 45 |
| H ₂ O ₂ | (a) | 18.8 | 16.39 | 687–807 | H ₂ O ₂ ignition | 46 |
| N ₂ H ₄ | | 2.0 | 11.54 | 25–150 | H ₂ / discharge | 50 |

FLUORINE ATOM TRANSFER

| | | | | | | |
|-----------------|--|--------|------|-----------|---|----|
| SF ₆ | | 30 ± 5 | 15.3 | 1030–1670 | H ₂ /O ₂ and C ₂ H ₂ /O ₂ flames | 51 |
|-----------------|--|--------|------|-----------|---|----|

CHLORINE ATOM TRANSFER

| | | | | | | |
|------------------|-----|-----------|--|--------|--|----|
| HCl | | 0.9 ± 1.3 | | 22–152 | HCl/D ₂ and DCl/H ₂ photolysis | 52 |
| Cl ₂ | | > 2.1 | | 25–75 | H ₂ /Cl ₂ /O ₂ photolysis | 53 |
| | | > 2.9 | | | | 54 |
| | | > 1.8 | | 25–100 | H ₂ /Cl ₂ /O ₂ photolysis | 55 |
| | | > 1.6 | | 0–85 | H ₂ /Cl ₂ /O ₂ photolysis | 56 |
| | | < 2.5 | | | | 57 |
| CCl ₄ | (s) | ~ 5 | | 2034 | Molecular beam | 58 |

Metathetical Reactions of Hydrogen Atoms—Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-----------|----------------------------|---|--------------------------------------|---|-----------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| BROMINE ATOM TRANSFER | | | | | | |
| Br ₂ | | 3.7 | 14.97 | 700–1400 | | 43 |
| CF ₃ Br | | 17.45 | 15.64 | 832–1011 | H ₂ /O ₂ ignition | 11 |
| C ₂ F ₄ Br ₂ | | 14.5 | 16.0 | 855–1013 | H ₂ /O ₂ ignition | 11 |
| IODINE ATOM TRANSFER | | | | | | |
| I ₂ | | 0.0 ± 0.5 | 14.6 ± 0.2 | 394–465 | H ₂ /I ₂ thermal reaction | 107 |
| OXYGEN ATOM TRANSFER | | | | | | |
| O ₂ | | | $k = 5.0 \times 10^9$ | 485 | H ₂ /O ₂ ignition | 60 |
| | | | $k = 6.7 \times 10^9$ | 520 | H ₂ /O ₂ ignition | 60 |
| | | | $k = 2.7 \times 10^9$ | 520 | H ₂ /O ₂ ignition | 61 |
| | | | $k = 6.6 \times 10^8$ | 520 | H ₂ /O ₂ ignition | 62 |
| | | 15.1 | 13.75 | | H ₂ /O ₂ ignition | 63 |
| | | 20.0 ± 2.0 | 14.9 | 1380 | H ₂ /O ₂ ignition | 64 |
| | | | $k = 1.43 \times 10^{12}$ | 1012–1227 | H ₂ /O ₂ flame | 109 |
| | | | $k = 1.5 \times 10^{11}$ | 827 | H ₂ /O ₂ flame | 110 |
| | | (t) 14.9 | 13.72 | | H ₂ /O ₂ ignition | 65 |
| | | 15.9 ± 0.8 | 13.89 | 570–660 | H ₂ /CO/O ₂ ignition | 66 |
| | | | $k = 6.0 \times 10^9$ | 540 | H ₂ /O ₂ ignition | 34 |
| | | 15.0 ± 0.6 | 13.59 | 590–660 | C ₂ H ₆ /CO/O ₂ ignition | 111 |
| | | 16.3 | | 552–1089 | H ₂ /O ₂ ignition | 67 |
| | | 17.4 ± 0.7 | 14.82 ± 0.2 | 1853–2772 | H ₂ /O ₂ flame | 68 |
| | | (a) | $k = 1.83 \times 10^3$ | 20 | | 69 |
| | | (a) | $k = 3.28 \times 10^3$ | 37 | | 70 |
| | | (f) 17.8 ± 1.0 | 15.08 ± 0.8 | 27–327 | | 70 |
| | | (f) 16.9 ± 1.0 | 14.86 ± 0.8 | 1227–1427 | | 70 |
| | | 18.9 ± 0.9 | | 620–1080 | H ₂ /O ₂ ignition | 71 |
| | | 15.6 ± 0.6 | 13.66 | 570–660 | H ₂ /CO/O ₂ ignition | 72 |
| | | (f) 16.49 ± 0.70 | 14.31 ± 0.28 | 20–1377 | | 73 |
| | | 17.0 | 13.69 | 999–1923 | C ₂ H ₂ /O ₂ oxidation | 74 |
| | | 17.75 | 15.0 | 687–807 | H ₂ /O ₂ ignition | 46 |
| | | $k = 1.6 \times 10^{10}$ | 642 | H ₂ /O ₂ flame | 108 | |
| | (f) 16.50 | 14.34 | 20–1400 | | 112 | |
| O ₃ | | | $k = (2.4 \pm 0.3) \times 10^{12}$ | Room temp. | Diffusion flame | 75 |
| | | | $k = (1.6 \pm 0.3) \times 10^{12}$ | Room temp. | H ₂ discharge | 76 |
| H ₂ O ₂ | (u) | | $k = (1.9 - 3.0) \times 10^{11}$ | 447 | H ₂ O ₂ ignition | 77 |
| | (u) | | $k = 1.1 \times 10^{12}$ | 500 | H ₂ /O ₂ slow reaction | 78 |
| N ₂ O | | 16.3 ± 2.0 | 14.62 | 990–1510 | H ₂ /N ₂ O/O ₂ flame | 110 |
| | | | $k = 4.3 \times 10^{10}$ | 627 | H ₂ /N ₂ O/O ₂ flame | 108 |
| | | | $k = 2.9 \times 10^{11}$ | 1084 | H ₂ /N ₂ O/O ₂ flame | 108 |
| | | | $k = (4.3 \pm 1.5) \times 10^{10}$ | 627 | H ₂ /N ₂ O/O ₂ flame | 45 |
| | (f) | 13 ± 1.5 | 13.7 ± 0.4 | 150–1487 | | 45 |

Metathetical Reactions of Hydrogen Atoms – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|-----------------|-------|------------------------------------|--|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| NO ₂ | | $k = 3.2 \times 10^{13}$ | | 227–267 | H ₂ /Cl ₂ NO ₂ thermal reaction | 79 |
| | | $k = 2.9 \times 10^{13}$ | | 25 | H ₂ / discharge | 80 |
| | (v) | $k = (2.9 \pm 0.3) \times 10^{13}$ | | Room temp. | H ₂ / discharge | 76 |
| | | $k = 5.5 \times 10^{13}$ | | 360 | H ₂ /O ₂ ignition | 81 |
| CO ₂ | (b) | 33.3 | 15.48 | 944–1072 | H ₂ /O ₂ flame | 113 |
| | | 29.7 | 14.89 | 1000–1500 | | 8 |
| | | | $k = 8.4 \times 10^8$ | 799 | H ₂ /O ₂ flame | 49 |

Ratios of Rate Constants (Hydrogen Atoms)

- (i) Reactions of the type (1) $H + RH = H_2 + R$ where RH is any hydrogen containing compound.
 (2) $H + OL = OLH$ where OL is an olefine.

| RH | OL | Notes | k_1/k_2 | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | Temperature range | Radical source | Reference |
|--|---|-------|--|---------------------------------|---------------------|-------------------|---|-----------|
| | | | | <i>(kcal mole⁻¹)</i> | | °C | | |
| CH ₄ | C ₂ H ₄ | | 2.6×10^{-6} | | | 120 | CH ₄ -H | 82 |
| C ₂ H ₄ | C ₂ H ₄ | | -1.0×10^{-3} (i.e. = 0) | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| C ₂ H ₆ | C ₂ H ₄ | | 2.5×10^{-5} | | | 25 | C ₂ H ₆ radiolysis | 84 |
| C ₃ H ₆ | C ₃ H ₆ | | 4.5×10^{-2} $(4.10 \pm 0.37) \times 10^{-2}$ | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| | | | | | | 25 | <i>n</i> -C ₄ H ₁₀ -H | 85 |
| C ₃ D ₆ | C ₃ D ₆ | | $(6.3 \pm 2.3) \times 10^{-2}$ | | | 25 | <i>n</i> -C ₄ H ₁₀ -H | 85 |
| C ₃ H ₈ | C ₂ H ₄ | | 1.9×10^{-3} 1.5×10^{-3} | | | 25 | C ₃ H ₈ -H | 86 |
| | | | | | | 24 | C ₃ H ₈ -H | 87 |
| C ₃ H ₈ | C ₃ H ₆ | | 5.2×10^{-4} | | | 25 | C ₃ H ₈ radiolysis | 88 |
| 1,3-C ₄ H ₆ | 1,3-C ₄ H ₆ | | 3.2×10^{-2} | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| 1-C ₄ H ₈ | 1-C ₄ H ₈ | | 8.1×10^{-2} | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| <i>i</i> -C ₄ H ₈ | <i>i</i> -C ₄ H ₈ | | 2.0×10^{-2} | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| cis-C ₄ H ₈ -2 | cis-C ₄ H ₈ -2 | | 6.6×10^{-2} | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| trans-C ₄ H ₈ -2 | trans-C ₄ H ₈ -2 | | 11.7×10^{-2} | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| <i>n</i> -C ₄ H ₁₀ | C ₃ H ₆ | | 1.4×10^{-3} | | | 25 | <i>n</i> -C ₄ H ₁₀ -H | 85 |
| <i>n</i> -C ₄ H ₁₀ | C ₃ D ₆ | | 1.3×10^{-3} | | | 25 | <i>n</i> -C ₄ H ₁₀ -H | 85 |
| <i>i</i> -C ₄ H ₁₀ | C ₂ H ₄ | | 8.6×10^{-3} 8.1×10^{-3} | | | 25 | <i>i</i> -C ₄ H ₁₀ -H | 86 |
| | | | | | | 24 | <i>i</i> -C ₄ H ₁₀ -H | 87 |

Ratios of Rate Constants (Hydrogen Atoms) – Continued

| RH | OL | Notes | k_1/k_2 | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | Temperature range | Radical source | Reference |
|--|--|-------|--|---------------------------------|---------------------|-------------------|--|-----------|
| | | | | <i>(kcal mole⁻¹)</i> | | °C | | |
| cyclo-C ₅ H ₁₀ | cyclo-C ₅ H ₈ | | 5.7×10^{-3} | | | 24 | cyclo-C ₅ H ₁₀ -H | 89 |
| <i>n</i> -C ₅ H ₁₂ | C ₂ H ₄ | | 3.7×10^{-3} 2.2×10^{-3} | | | 25 24 | <i>n</i> -C ₅ H ₁₂ -H <i>n</i> -C ₅ H ₁₂ -H | 86 87 |
| <i>n</i> -C ₅ H ₁₂ | C ₅ H ₁₀ | (y) | $\geq 4.5 \times 10^{-3}$ | | | 21 | <i>n</i> -C ₅ H ₁₂ -H | 114 |
| <i>i</i> -C ₅ H ₁₂ | C ₂ H ₄ | | 10.6×10^{-3} | | | 25 | <i>i</i> -C ₅ H ₁₂ -H | 86 |
| (CH ₃) ₂ C: C(CH ₃) ₂ | (CH ₃) ₂ C: C(CH ₃) ₂ | | -2.5×10^{-2} (i.e. = 0) | | | 24 | <i>n</i> -C ₄ H ₁₀ -H | 83 |
| <i>n</i> -C ₆ H ₁₄ | C ₂ H ₄ | | 3.7×10^{-3} | | | 25 | <i>n</i> -C ₆ H ₁₄ -H | 86 |
| neo-C ₆ H ₁₄ | C ₂ H ₄ | | 1.8×10^{-3} | | | 25 | neo-C ₆ H ₁₄ -H | 86 |
| (CH ₃) ₂ CHCH ₂ CH ₂ CH ₃ | C ₂ H ₄ | | 11.6×10^{-3} | | | 25 | (CH ₃) ₂ CHCH ₂ CH ₂ CH ₃ -H | -H 86 |
| (CH ₃) ₂ CHCH (CH ₃) ₂ | C ₂ H ₄ | | 21.0×10^{-3} | | | 25 | (CH ₃) ₂ CHCH (CH ₃) ₂ -H | 86 |
| H ₂ S | C ₂ H ₄ | | 0.5 | | | 30 | H ₂ S photolysis | 90 |
| CH ₃ SH | C ₂ H ₄ | | | 0.54 ± 0.12 | 0.59 | 50–220 | CH ₃ SH photolysis | 91 |

(ii) Miscellaneous reactions of hydrogen atoms with hydrocarbons and hydrocarbon derivatives.

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|---------------------|-----------|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) H + CCl ₃ D = CCl ₃ + HD (2) H + CCl ₃ D = CCl ₂ D + HCl | | | | 1.1 | Room temp. | H ₂ / discharge | 92 |
| (1) H + CCl ₃ Br = CCl ₃ + HBr (2) H + CCl ₃ Br = CCl ₂ Br + HCl | | | | 0.49 | Room temp. | H ₂ / discharge | 115 |
| (1) H + CCl ₃ F = CCl ₃ + HF (2) H + CCl ₃ F = CCl ₂ F + HCl | | | | 0.071 | Room temp. | H ₂ / discharge | 115 |
| (1) H + CCl ₂ BrH = CCl ₂ Br + H ₂ (2) H + CCl ₂ BrH = CClBrH + HCl | | | | 1.7 | Room temp. | H ₂ / discharge | 92 |
| (1) H + CCl ₂ BrH = CCl ₂ H + HBr (2) H + CCl ₂ BrH = CClBrH + HCl | | | | 0.8 | Room temp. | H ₂ / discharge | 92 |
| (1) H + C ₃ H ₆ = C ₃ H ₅ + H ₂ (2) H + C ₃ H ₈ = C ₃ H ₇ + H ₂ | | | | 32 | 25 | C ₃ H ₈ radiolysis | 88 |
| (1) H + (CH ₃) ₃ CD = C ₄ H ₉ + HD (2) H + (CH ₃) ₃ CD = C ₄ H ₈ D + H ₂ | | | | 1.2 | 25 | <i>i</i> -C ₄ H ₉ D photolysis | 93 |

Ratios of Rate Constants (Hydrogen Atoms)–Continued

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|---------------------|--|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) $D + (CH_3)_3CD = C_4H_9 + D_2$ (2) $D + (CH_3)_3CD = C_4H_8D + HD$ | z | | | 1.2 | 25 | <i>i</i> -C ₄ H ₉ D photolysis | 93 |
| (1) $H + i-C_5H_{12} = C_5H_{11} + H_2$ (2) $H + C_5H_{11} = C_5H_{10} + H_2$ | (z) | | | 5.2×10^{-2} | 25 | <i>i</i> -C ₅ H ₁₂ -H | 94 |
| (1) $H + 1-C_5H_{10} = C_5H_9 + H_2$ (2) $H + n-C_5H_{12} = C_5H_{11} + H_2$ | | | | 8.8 | 21 | <i>n</i> -C ₅ H ₁₂ -H | 88 |
| (1) $H + i-C_5H_{12} = C_5H_{11} + H_2$ (2) $H + C_5H_{11} = C_5H_{12}$ | (z) | | | 5.9×10^{-3} | 25 | <i>i</i> -C ₅ H ₁₂ -H | 94 |
| (1) $H + n-C_6H_{14} = C_6H_{13} + H_2$ (2) $H + C_6H_{13} = C_6H_{14}$ | | | | 5.3×10^{-3} 1.3×10^{-3} | 25 25 | <i>n</i> -C ₆ H ₁₄ -H <i>n</i> -C ₆ H ₁₄ /1-C ₆ H ₁₂ -H | 94 94 |
| (1) $H + CH_3CH:CHCHO = CH_3CH:CHCO + H_2$ (2) $H + CH_3CH:CHCHO = CH_3CH:CH_2 + HCO$ | | ~ 0.5 | | | 210–350 | (CH ₃) ₂ CO/ CH ₃ CH:CHCHO photolysis | 95 |

(iii) Reactions of the type (1) $H + HX = H_2 + X$

(2) $H + X_2 = HX + X$ where X is a halogen

| | | | | | | | |
|---|------|-----------------|-----------------|--|--------------------------------|---|------------------------|
| (1) $H + HCl = H_2 + Cl$ (2) $H + Cl_2 = HCl + Cl$ | | 1.54 ± 0.13 | -0.84 ± 0.1 | ≤ 0.05 | 25 0–62 | H ₂ /Cl ₂ photolysis H ₂ /Cl ₂ photolysis | 96 97 |
| (1) $H + HBr = H_2 + Br$ (2) $H + Br_2 = HBr + Br$ | (aa) | | | 0.12 0.1 0.12 ± 0.009 | 327–1197 162–310 25–300 | H ₂ /Br ₂ thermal reaction H ₂ /Br ₂ photolysis H ₂ /Br ₂ thermal reaction | 98 99 100 |
| (1) $H + HI = H_2 + I$ (2) $H + I_2 = HI + I$ | | | | 0.099 ± 0.007 0.070 \pm 0.02 0.082 ± 0.012 0.06 | 1027–1427 394 527 700 | H ₂ /Br ₂ thermal reaction H ₂ /I ₂ thermal reaction H ₂ /I ₂ thermal reaction | 101 44 44 102 |
| (1) $D + DI = D_2 + I$ (2) $D + I_2 = DI + I$ | | | | 0.073 | 527 | D ₂ /I ₂ thermal reaction | 116 |

(iv) Miscellaneous Ratios

| | | | | | | | |
|--|--|--|--|-----------------------------|-----------|--------------------------------|----------|
| (1) $H + D_2 = HD + D$ (2) $H + HCl = H_2 + Cl$ | | | | 14 ± 4 | 22 | HCl/D ₂ photolysis | 52 |
| (1) $D + HD = D_2 + H$ (2) $D + DCl = D_2 + Cl$ | | | | 65 | 22 | H ₂ /DCl photolysis | 52 |
| (1) $D + HCl = HD + Cl$ (2) $D + HCl = DCl + H$ | | | | 0.5 ± 0.4 $5. \pm 4$ | 22 152 | D ₂ /HCl photolysis | 52 52 |

Ratios of Rate Constants (Hydrogen Atoms)–Continued

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|---------------------|--|-------------------|----------------------------------|------------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) $H + D_2 = HD + D$ (2) $H + HBr = H_2 + Br$ | | | | 0.55 | 22 | D_2/HBr photolysis | 52 |
| (1) $H + NO_2 = NO + OH$ (2) $H + Cl_2 = HCl + Cl$ | | -3.13 | 0.42 | | 227-267 | $H_2/Cl_2/NO_2$ thermal reaction | 79 |
| (1) $D + C_2H_6 = C_2H_5 + HD$ (2) $D + DI = D_2 + I$ | | | | 1.2×10^{-3} 8.7×10^{-3} | 25 100 | DI photolysis DI photolysis | 103 103 |
| (1) $H + H_2S = HS + H_2$ (2) $H + O_2 = HO_2$ | | | | 0.60 0.137 0.171 | 50 93 100 | H_2S photo-oxidation | 104 |

Notes

- (a) Estimated from the reverse reaction.
 (b) This value was used by the authors in the region 1000–1500 °C, but its origin was not stated.
 (c) Measured relative to $D + D_2CO = D_2 + DCO$ for which $k = 10^{13.30} \exp(-3000/RT)$.
 (d) Private communication from Asaba.
 (e) Taken from Tikhomirova and Voevodsky, "Chain oxidation of Hydrocarbons in the Gas Phase" (Russ), Moscow, 1955.
 (f) Critical survey of literature data.
 (g) "A rough estimate."
 (h) Measured relative to $H + C_3H_6 = C_3H_7$ for which $k = 10^{13.5} \exp(-2600/RT)$.
 (i) Measured relative to $D + H_2 = HD + H$ for which $k = 10^{13.4} \exp(-5000/RT)$.
 (j) Measured relative to $H + O_2 = OH + O$ for which $k = 5.1 \times 10^9$ at 520 °C.
 (k) Measured relative to $H + C_3H_8 = C_3H_7 + H_2$ for which $k = 10^{14.11} \exp(-8200/RT)$.
 (l) Measured relative to $D + H_2 = HD + H$ for which $k = 10^{13.92} \exp(-6000/RT)$.
 (m) Measured relative to $D + H_2 = HD + H$ for which $k = 10^{13.46} \exp(-5400/RT)$.
 (n) Rate constant estimated in computer calculations. E value assumed and A factor varied.
 (o) Estimated from kinetic data.
 (p) The value of E was assumed, and A calculated from the rate constant.
 (q) A steric factor of 0.1 was assumed.
 (r) Calculated from the recommended value for the reverse reaction.
 (s) Steric factor = 0.13.
 (t) Used at flame temperatures.
 (u) Measured relative to $H + O_2 = OH + O$ for which $k = 10^{14.31} \exp(-16500/RT)$.
 (v) Measured relative to $H + O_2 + H_2 = HO_2 + H_2$ for which $k = 5.5 \times 10^{13} \text{ cm}^6 \text{ mole}^2 \text{ sec}^{-1}$.
 (w) Reactions with D atoms.
 (x) Calculated from the recommended value for the reverse reaction.
 (y) C_5H_{10} refers to the mixture of pentenes formed in the primary reaction.

(z) C_3H_{11} is a radical formed in the primary process, probably either $CH_3C(CH_3)CH_2CH_3$ or $CH_3CH(CH_3)CHCH_3$.

(aa) The author shows that this value is compatible with his results in the temperature range shown.

References

1. Majury and Steacie, *Disc. Faraday Soc.* **14**, 45 (1953).
2. Le Roy, *Disc. Faraday Soc.* **14**, 120 (1953).
3. Berlie and Le Roy, *Can. J. Chem.* **32**, 650 (1954).
4. Pritchard, Pyke, and Trotman-Dickenson, *J. Am. Chem. Soc.* **77**, 2629 (1955).
5. Benson and Jain, *J. Chem. Phys.* **31**, 1008 (1959).
6. Fenimore and Jones, *J. Phys. Chem.* **65**, 2200 (1961).
7. Gorban and Nalbandyan, *Russ. J. Phys. Chem.* **36**, 946 (1962).
8. Intezarova, Kondratiev, and Mukhoyan, *Kinetika i Kataliz* **5**, 585 (1964).
9. Azatyan, Nalbandyan, and Tsui, *Kinetika i Kataliz* **5**, 201 (1964).
10. Jamieson and Brown, *Can. J. Chem.* **42**, 1638 (1964).
11. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
12. Steacie, *Can. J. Res.* **15B**, 264 (1937).
13. Klein, McNesby, Scheer, and Schoen, *J. Chem. Phys.* **30**, 58 (1959).
14. Chase and Weinberg, *Proc. Roy. Soc.* **275A**, 411 (1963).
15. Voevodsky and Kondratiev, "Progress in Reaction Kinetics," p. 41 (Pergamon Press, 1961).
16. Berlie and Le Roy, *Disc. Faraday Soc.* **14**, 50 (1953).
17. Gorban and Nalbandyan, *Dokl. Akad. Nauk. S.S.S.R.* **132**, 1335 (1960).
18. Trost and Steacie, *J. Chem. Phys.* **16**, 361 (1948).
19. Baldwin, 9th Int. Comb. Symp., p. 604 (Academic Press, N.Y., 1963).
20. Blackmore and Hinshelwood, *Proc. Roy. Soc.* **271A**, 34 (1963).
21. Yang, *J. Phys. Chem.* **67**, 562 (1963).
22. Baldwin and Melvin, *J. Chem. Soc.* 1785 (1964).
23. Darwent and Roberts, *Disc. Faraday Soc.* **14**, 55 (1953).
24. Yang, *J. Am. Chem. Soc.* **84**, 3795 (1962).
25. Kazmi, Diefendorf, and Le Roy, *Can. J. Chem.* **41**, 690 (1963).
26. Baldwin, *Trans. Faraday Soc.* **60**, 527 (1964).
27. Schiff and Steacie, *Can. J. Chem.* **29**, 1 (1951).
28. Baldwin and Walker, *Trans. Faraday Soc.* **60**, 1236 (1964).
29. Kazmi and Le Roy, *Can. J. Chem.* **42**, 1145 (1964).

30. White, Winkler, and Kenalty, *Can. J. Res.* **20B**, 255 (1942).
31. Trost and Steacie, *J. Chem. Phys.* **16**, 361 (1948).
32. Norrish and Purnell, *Proc. Roy. Soc.* **243A**, 449 (1958).
33. McNesby, Scheer, and Klein, *J. Chem. Phys.* **32**, 1814 (1960).
34. Baldwin and Cowe, *Trans. Faraday Soc.* **58**, 1768 (1962).
35. Brennen, Gay, Glass, and Niki, *J. Chem. Phys.* **43**, 2569 (1965).
36. Klein, Scheer, and Schoen, *J. Am. Chem. Soc.* **78**, 50 (1956).
37. Trost, Darwent, and Steacie, *J. Chem. Phys.* **16**, 353 (1948).
38. Harris and Steacie, *J. Chem. Phys.* **13**, 554 (1945).
39. Jacobs, Giedt, and Cohen, *J. Chem. Phys.* **43**, 3688 (1965).
40. Steiner and Rideal, *Proc. Roy. Soc.* **173A**, 503 (1939).
41. Bigeleisen, Klein, Weston, and Wolfsberg, *J. Chem. Phys.* **30**, 1340 (1959).
42. Steiner *Proc. Roy. Soc.* **173A**, 531 (1939).
43. Cooley and Anderson, *Ind. and Eng. Chem.* **44**, 1402 (1952).
44. Sullivan, *J. Chem. Phys.* **36**, 1925 (1962).
45. Dixon-Lewis, Sutton, and Williams, *J. Chem. Soc.* 5724 (1965).
46. Skinner and Ringrose, *J. Chem. Phys.* **42**, 2190 (1965).
47. Kaufman and Del Greco, 9th Int. Comb. Symp., p. 659 (Academic Press, N.Y., 1963).
48. Dixon-Lewis, Sutton, and Williams, *Disc. Faraday Soc.* **33**, 205 (1962).
49. Dixon-Lewis, Sutton, and Williams, *Trans. Faraday Soc.* **61**, 255 (1965).
50. Schiavello and Volpi, *J. Chem. Phys.* **37**, 1510 (1962).
51. Fenimore and Jones, *Comb. and Flame* **8**, 231 (1964).
52. DeVries and Klein, *J. Chem. Phys.* **41**, 3428 (1964).
53. Hertel, *Z. Phys. Chem.* **15B**, 325 (1931).
54. Padoa and Butirone, *Accad. Lincei* **25**, 215 (1916); and *Gazz. Chim. Ital.* **47**, 6 (1917).
55. Porter, Bardwell, and Lind, *J. Am. Chem. Soc.* **48**, 2603 (1926).
56. Krauskopf and Rollefson, *J. Am. Chem. Soc.* **56**, 327 (1934).
57. Trotman-Dickenson, "Gas Kinetics," p. 184 (Butterworths, London, 1955).
58. Seidel, Martin, and Mietzner, *Z. Phys. Chem.* **47B**, 348 (1965).
59. Campbell and Hirschfelder, Univ. of Wisconsin report CF-2108 NORD 9938.
60. Semenoff, *Acta. Physicochim.* **20**, 291 (1945).
61. Baldwin, *Trans. Faraday Soc.* **52**, 1344 (1956).
62. Baldwin and Simmons, *Trans. Faraday Soc.* **53**, 964 (1957).
63. Semenoff, Nalbandyan, and Karmilova, *Zhur. Phys. Chem.* **32**, 1193 (1957).
64. Schott and Kinsey, *J. Chem. Phys.* **29**, 1177 (1958).
65. Kondratiev, 7th Int. Comb. Symp., p. 41 (Butterworths, London, 1959).
66. Azatyan, Voevodskii, and Nalbandyan, *Kinetika i Kataliz* **2**, 340 (1961).
67. Fujimoto, *Bull. Chem. Soc. Japan* **36**, 1233 (1963).
68. Lovachev, *Comb. and Flame* **7**, 388 (1963).
69. Clyne, 9th Int. Comb. Symp., p. 211 (Academic Press, N.Y., 1963).
70. Kaufman and Del Greco, 9th Int. Comb. Symp., p. 659 (Academic Press, N.Y., 1963).
71. Miyama and Takeyama, *J. Chem. Phys.* **41**, 2287 (1964).
72. Azatyan, Nalbandyan, and Tsui, *Kinetika i Kataliz* **5**, 201 (1964).
73. Baldwin and Melvin, *J. Chem. Soc.* 1785 (1964).
74. Glass, Kistiakowsky, Michael, and Niki, *J. Chem. Phys.* **42**, 608 (1965).
75. Garvin and McKinley, *J. Chem. Phys.* **24**, 1256 (1956).
76. Phillips and Schiff, *J. Chem. Phys.* **37**, 1233 (1962).
77. Forst and Giguere, *J. Phys. Chem.* **62**, 340 (1958).
78. Baldwin and Mayor, *Trans. Faraday Soc.* **56**, 103 (1960).
79. Rosser and Wise, *J. Phys. Chem.* **65**, 532 (1961).
80. Schiff, *Disc. Faraday Soc.* **33**, 285 (1962).
81. Ashmore and Tyler, *Trans. Faraday Soc.* **58**, 1108 (1962).
82. Back and Van der Auwera, *Can. J. Chem.* **40**, 2339 (1962).
83. Jennings and Cvetanovic, *J. Chem. Phys.* **35**, 1233 (1961).
84. Yang and Gant, *J. Phys. Chem.* **65**, 1861 (1961).
85. Takahasi and Cvetanovic, *Can. J. Chem.* **40**, 1037 (1962).
86. Holroyd and Klein, *J. Phys. Chem.* **67**, 2273 (1963).
87. Holroyd and Pierce, *J. Phys. Chem.* **68**, 1392 (1964).
88. Back, *J. Phys. Chem.* **64**, 124 (1960).
89. Stock and Gunning, *Can. J. Chem.* **38**, 2295 (1960).
90. Arthur and Bell, *J. Chem. Soc.* 4866 (1962).
91. Inaba and Darwent, *J. Phys. Chem.* **64**, 1431 (1960).
92. Clark and Tedder, *Trans. Faraday Soc.* **62**, 405 (1966).
93. Okabe and Becker, *J. Am. Chem. Soc.* **84**, 4004 (1962).
94. Kuntz, *J. Phys. Chem.* **69**, 2291 (1965).
95. Pitts, Thompson, and Woolfolk, *J. Am. Chem. Soc.* **80**, 66 (1958).
96. Bigeleisen, Klein, Weston, and Wolfsberg, *J. Chem. Phys.* **30**, 1340 (1959).
97. Klein and Wolfsberg, *J. Chem. Phys.* **34**, 1494 (1961).
98. Levy, *J. Phys. Chem.* **62**, 570 (1958).
99. Bodenstein and Lutkemeyer, *Z. Phys. Chem.* **114**, 208 (1924).
100. Bodenstein and Jung, *Z. Phys. Chem.* **121**, 127 (1926).
101. Britton and Cole, *J. Phys. Chem.* **65**, 1302 (1961).
102. Horie, Ishii, and Amano, *J. Phys. Chem.* **68**, 1264 (1964).
103. Carter, Hamill, and Williams, *J. Am. Chem. Soc.* **77**, 6457 (1955).
104. Darwent and Krasnansky, 7th Int. Comb. Symp., p. 3 (Butterworths, London, 1959).
105. Fenimore and Jones, 9th Int. Comb. Symp., p. 597 (Academic Press, N.Y., 1963).
106. Harris and Steacie, *J. Chem. Phys.* **13**, 559 (1945).
107. Sullivan, *J. Chem. Phys.* **30**, 1292 (1959).
108. Dixon-Lewis, Sutton, and Williams, 10th Int. Comb. Symp., p. 495 (Comb. Inst. Pittsburgh, 1965).
109. Fenimore and Jones, *J. Phys. Chem.* **62**, 693 (1958).
110. Fenimore and Jones, *J. Phys. Chem.* **63**, 1154 (1959).
111. Azatyan, Nalbandyan, and Tsui, *Dokl. Akad. Nauk. S.S.S.R.* **147**, 361 (1962).
112. Westenberg and Fristrom, 10th Int. Comb. Symp., p. 473 (Comb. Inst. Pittsburgh, 1965).
113. Fenimore and Jones, *J. Phys. Chem.* **62**, 1578 (1958).
114. Back, *Trans. Faraday Soc.* **54**, 512 (1958).
115. Clark and Tedder, *Trans. Faraday Soc.* **62**, 399 (1966).
116. Sullivan, *J. Chem. Phys.* **39**, 3001 (1963).

Review

1. Thrush, "Progress in Reaction Kinetics," Vol. 3, p. 65 (Pergamon Press, London, 1965).

Metathetical Reactions of Fluorine Atoms

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H ₂ | (a) | 1.71 ± 0.22 | 14.07 | 25-150 | CH ₄ /H ₂ /F ₂ P | 1, 4 |
| Alkanes | | | | | | |
| CH ₄ | (a) | 1.21 ± 0.80 | 14.09 | -75-78 | CH ₄ /C ₂ H ₆ /F ₂ T | 2, 4 |
| C ₂ H ₆ | | 0.28 | 13.78 | -60-20 | standard T | 2, 4 |
| CH ₃ CH ₂ CH ₃ | (a) | 0.00 ± 0.25 | 13.52 | -60-20 | C ₂ H ₆ /C ₃ H ₈ /F ₂ T | 2, 4 |
| CH ₃ CH ₂ CH ₃ | (a) | 0.00 ± 0.25 | 13.10 | -60-20 | C ₂ H ₆ /C ₃ H ₈ /F ₂ T | 2, 4 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (a) | 0.00 ± 0.70 | 13.46 | -60-20 | <i>n</i> -C ₄ H ₁₀ /C ₃ H ₈ /F ₂ T | 2, 3, 4 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (a) | 0.00 ± 0.65 | 13.37 | -60-20 | <i>n</i> -C ₄ H ₁₀ /C ₃ H ₈ /F ₂ T | 2, 3, 4 |
| (CH ₃) ₃ CH | (a) | 0.00 ± 0.70 | 13.65 | -60-20 | <i>n</i> -C ₄ H ₁₀ / <i>i</i> -C ₄ H ₁₀ /F ₂ T | 2, 3, 4 |
| (CH ₃) ₃ CH | (a) | 0.00 ± 0.80 | 12.84 | -60-20 | <i>n</i> -C ₄ H ₁₀ / <i>i</i> -C ₄ H ₁₀ /F ₂ T | 2, 3, 4 |
| (CH ₃) ₄ C | (a) | 0.00 ± 0.80 | 13.75 | -60-20 | <i>i</i> -C ₄ H ₁₀ /(CH ₃) ₄ C/F ₂ T | 2, 4 |
| Cyclo-Alkane | | | | | | |
| cyclo-C ₃ H ₆ | (a) | 0.00 ± 0.10 | 13.46 | | cyclo-C ₃ H ₆ /C ₃ H ₈ /F ₂ T | 2, 4 |
| CHLORINE ATOM TRANSFER | | | | | | |
| CCl ₄ | | $k = 2 \times 10^{13}$ | | 20 | CCl ₄ /F ₂ T | 5 |
| OXYGEN ATOM TRANSFER | | | | | | |
| O ₃ | | 2.8 | | 0-20 | F ₂ /O ₃ P | 6 |

Notes

(a) These values are based on the assumed value $k = 10^{13.78} \exp(-280/RT)$ for the reaction $F + C_2H_6 = C_2H_5 + HF$.

References

1. Mercer and Pritchard, *J. Phys. Chem.* **63**, 1468 (1959).
2. Fettis, Knox, and Trotman-Dickenson, *J. Chem. Soc.* 1064 (1960).
3. Anson, Fredricks, and Tedder, *J. Chem. Soc.* 918 (1959).
4. Fettis and Knox, "Progress in Reaction Kinetics," Vol. 2, p. 3 (Pergamon Press, N.Y., 1964).
5. Clark and Tedder, *J. Phys. Chem.* **68**, 2018 (1964).

6. Starico, Sicre, and Schumacher, *Anales Assoc. Quim. Argentina* **50**, 120 (1962).

Reviews

1. Fettis and Knox, "Progress in Reaction Kinetics," Vol. 2, p. 3 (Pergamon Press, N.Y., 1964).
2. Fettis, Knox, and Trotman-Dickenson, *Can. J. Chem.* **38**, 1643 (1960).

Reaction of Chlorine Atoms

Index

| | |
|---|-----------|
| (a) Hydrogen atom transfer | Page |
| Reaction with | |
| 1. Hydrogen..... | 17 |
| 2. Alkanes..... | 17 |
| 3. Cyclo-alkanes..... | 17 |
| 4. Halogenated alkanes..... | 18 |
| 5. Hydrogen halide..... | 19 |
| (b) Chlorine atom transfer | |
| Reaction with | |
| 1. Halogenated alkanes..... | 19 |
| 2. Halogens and pseudo-halogens..... | 19 |
| 3. Carbonyls, etc..... | 19 |
| (c) Bromine atom transfer..... | 20 |
| (d) Oxygen atom transfer..... | 20 |
| (e) Miscellaneous reactions..... | 20 |

Metathetical Reactions of Chlorine Atoms

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|-------|----------------------------|---|-------------------|---|-------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| H ₂ | | ≥ 5.9 | | 25-75 | H ₂ /Cl ₂ | P 1 |
| | | ≥ 5.8 | | -73-23 | H ₂ /Cl ₂ | P 2 |
| | | 6.1 ± 1.0 | 14.3 | 0-25 | Cl ₂ discharge | 3, 38 |
| | | | $k = (4.8 \pm 0.4) \times 10^{11}$ | 250 | H ₂ /Cl ₂ /NO | T 5, 6 |
| | | 5.9 | | 628-798 | H ₂ /HCl | T 4 |
| | (a) | 5.48 ± 0.14 | 13.92 ± 0.03 | 0-798 | | 7, 5 |
| | | 5.48 | 13.92 | | | |
| HD | (b) | 5.97 | 13.83 | -33-77 | H ₂ /HD/Cl ₂ | P 8, 41 |
| HT | (b) | 6.03 | 13.79 | -38-72 | H ₂ /HT/Cl ₂ | P 9 |
| | (b) | 6.28 | 13.82 | -30-70 | H ₂ /HD/HT/Cl ₂ | P 41 |
| D ₂ | (b) | 6.60 | 13.78 | 0-32 | H ₂ /CO/Cl ₂ | P 10, 11, 7 |
| | (b) | 6.70 | 13.84 | 30-178 | H ₂ /D ₂ /Cl ₂ | P 12 |
| | (b) | 6.61 | 13.76 | -30-70 | HD/D ₂ /Cl ₂ | P 41 |
| DT | (b) | 6.90 | 13.73 | -30-70 | DT/D ₂ /Cl ₂ | P 41 |
| T ₂ | (b) | 7.17 | 13.73 | -30-70 | T ₂ /D ₂ /Cl ₂ | P 41 |
| Alkanes | | | | | | |
| CH ₄ | (b) | 3.85 ± 0.18 | 13.42 | 20-211 | CH ₄ /H ₂ /Cl ₂ | P 13, 16 |
| CD ₄ | (i) | 5.76 | 13.73 | 31-188 | CH ₄ /CD ₄ /Cl ₂ | P 12 |
| C ₂ H ₆ | (b) | 1.04 ± 0.08 | 13.96 | -41-385 | CH ₄ /C ₂ H ₆ /Cl ₂ | P 14, 15 |
| | (b) | 1.00 | 14.08 | 76-290 | CH ₄ /C ₂ H ₆ /Cl ₂ | P 16 |
| C ₂ D ₆ | (i) | 1.32 | 13.75 | 30-160 | C ₂ H ₆ /C ₂ D ₆ /Cl ₂ | P 12 |
| C ₃ H ₈ | (b) | 0.67 | 14.28 | 25-211 | C ₃ H ₈ /C ₂ H ₆ /Cl ₂ | P 16 |
| CH ₃ CH ₂ CH ₃ | (b) | 0.98 | 14.03 | -70-230 | C ₂ H ₆ /C ₃ H ₈ /Cl ₂ | P 14, 15 |
| CH ₃ CH ₂ CH ₃ | (b) | 0.66 | 13.87 | -70-230 | C ₂ H ₆ /C ₃ H ₈ /Cl ₂ | P 14, 15 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (b) | 0.77 | 13.93 | -70-200 | C ₂ H ₆ / <i>n</i> -C ₄ H ₁₀ /Cl ₂ | P 14, 17 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (b) | 0.25 | 13.95 | -70-200 | C ₂ H ₆ / <i>n</i> -C ₄ H ₁₀ /Cl ₂ | P 14, 17 |
| (CH ₃) ₃ CH | (b) | 0.86 | 14.29 | 25-211 | C ₂ H ₆ / <i>i</i> -C ₄ H ₁₀ /Cl ₂ | P 16 |
| (CH ₃) ₃ CH | (b) | 0.80 | 14.09 | -80-100 | C ₂ H ₆ / <i>i</i> -C ₄ H ₁₀ /Cl ₂ | P 14, 17 |
| (CH ₃) ₃ CH | (b) | 0.02 | 13.24 | -80-100 | C ₂ H ₆ / <i>i</i> -C ₄ H ₁₀ /Cl ₂ | P 14, 17 |
| (CH ₃) ₄ C | (b) | 0.90 | 14.25 | -70-230 | C ₃ H ₈ /(CH ₃) ₄ C/Cl ₂ | P 14 |
| | (b) | 0.70 | 14.09 | 25-211 | C ₂ H ₆ /(CH ₃) ₄ C/Cl ₂ | P 16 |
| Cyclo-alkanes | | | | | | |
| cyclo-C ₃ H ₆ | (b) | 4.12 | 13.73 | 20-260 | C ₂ H ₆ /cyclo-C ₃ H ₆ /Cl ₂ | P 14 |

Metathetical Reactions of Chlorine Atoms – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|-------|---------------------------------|--|-------------------|--|--------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| cyclo-C ₄ H ₈ | (b) | 0.80 | 14.43 | -30-105 | C ₃ H ₈ /cyclo-C ₄ H ₈ /Cl ₂ | P 14 |
| cyclo-C ₅ H ₁₀ | (b) | 0.58 | 14.47 | 25-211 | C ₂ H ₆ /cyclo-C ₅ H ₁₀ /Cl ₂ | P 16 |
| Halogenated alkanes | | | | | | |
| CH ₃ Cl | (b) | 3.36 | 13.76 | 25-211 | CH ₄ /CH ₃ Cl/Cl ₂ | P 16 |
| | (b) | 3.3 ± 0.1 | 13.5 ± 0.7 | 0-300 | CH ₃ Cl/vari- ous RH/Cl ₂ | P 18 |
| | (c) | 3.08 | 13.5 | 85-180 | C ₂ Cl ₄ /CH ₃ Cl/Cl ₂ | P 19 |
| CH ₂ Cl ₂ | | 5.5 | 14.6 | | | 28 |
| | | 3.0 | 13.43 | 0-300 | CH ₂ Cl ₂ /vari- ous RH/Cl ₂ | P 18 |
| | (c) | 3.12 | 13.4 | 110-210 | C ₂ Cl ₄ /CH ₂ Cl ₂ /Cl ₂ | P 19 |
| CHCl ₃ | | 6.5 | 14.6 | | | 28 |
| | | 3.35 | 12.84 | 0-300 | CH ₄ /CHCl ₃ /Cl ₂ | P 18, 27 |
| | (c) | 3.33 | 13.2 | 140-210 | C ₂ Cl ₄ /CHCl ₃ /Cl ₂ | P 19, 20, 21 |
| CDCl ₃ | (d) | 4.06 ± 0.2 | 12.69 | -20-180 | CHCl ₃ /CDCl ₃ /Cl ₂ | P 22 |
| | (d) | 4.79 | 13.30 | 109-203 | CHCl ₃ /CDCl ₃ /Cl ₂ | P 12 |
| C ₂ H ₅ Cl | (b) | 1.50 ± 0.2 | 13.66 | 25-211 | C ₂ H ₆ /C ₂ H ₅ Cl/Cl ₂ | P 16 |
| CH ₂ ClCH ₂ Cl | | 3.0 | 13.4 | | | 23 |
| C ₂ H ₃ Cl ₃ | | 3.5 | 13.5 | | | 23 |
| CHCl ₂ CHCl ₂ | | 3.3 | 13.8 | | | 23 |
| C ₂ HCl ₅ | | 5.4 | 13.9 | | | 28 |
| | | 3.3 | 12.8 | 110-210 | C ₂ Cl ₄ /C ₂ HCl ₅ /Cl ₂ | P 19, 24 |
| | | 3.4 | 12.7 | | | 18 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (e) | 0.77 ± 0.10 | 13.0 | 0-230 | C ₄ H ₉ F/Cl ₂ | P 25, 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (e) | 0.62 ± 0.15 | 13.1 | 0-230 | C ₄ H ₉ F/Cl ₂ | P 25, 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (e) | (0.37 ± 0.10) | (13.3) | 0-230 | C ₄ H ₉ F/Cl ₂ | P 25, 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (e) | 0.77 | 13.1 | 0-230 | C ₄ H ₉ F/Cl ₂ | P 25, 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ Cl | (e) | 0.77 ± 0.20 | 13.4 | 35-146 | C ₄ H ₉ Cl/Cl ₂ | P 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ Cl | (e) | 0.30 ± 0.40 | 13.3 | 35-146 | C ₄ H ₉ Cl/Cl ₂ | P 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ Cl | (e) | 0.22 ± 0.10 | 13.7 | 35-146 | C ₄ H ₉ Cl/Cl ₂ | P 26, 29 |
| CH ₃ CH ₂ CH ₂ CH ₂ Cl | (e) | (0.77) | (13.4) | 35-146 | C ₄ H ₉ Cl/Cl ₂ | P 26, 29 |
| CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ | (e) | 2.0 | 12.2 | 0-230 | CF ₃ C ₄ H ₉ /Cl ₂ | P 25 |
| CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ | (e) | 0.6 | 13.0 | 0-230 | CF ₃ C ₄ H ₉ /Cl ₂ | P 25 |

Metathetical Reactions of Chlorine Atoms – Continued

| Reactants | Notes | E | $\log_{10} A$ | Temperature range | Radical source | References |
|------------------------|-------|-----------------------|---------------------------------|-------------------|---------------------|------------|
| | | ($kcal\ mole^{-1}$) | ($cm^3\ mole^{-1}\ sec^{-1}$) | $^{\circ}C$ | | |
| $CF_3CH_2CH_2CH_2CH_3$ | (e) | (0.3) | (13.3) | 0–230 | $CF_3C_4H_9/Cl_2$ P | 25 |
| $CF_3CH_2CH_2CH_2CH_3$ | (e) | 0.5 | 12.8 | 0–230 | $CF_3C_4H_9/Cl_2$ P | 25 |
| Hydrogen Halide | | | | | | |
| HCl | | 6.57 ± 0.49 | | 39–150 | $HCl/D_2/Cl_2$ P | 39 |

CHLORINE ATOM TRANSFER

| | | | | | | |
|-------------------------------------|---------|----------------|----------------|-----------|-------------|--------|
| Halogenated alkanes | | | | | | |
| CH_3Cl | (f) (g) | 25.0 | 14.0 | | | 23, 30 |
| CH_2Cl_2 | (f) (g) | 21.4 | 14.0 | | | 23, 30 |
| $CHCl_3$ | (f) (g) | 21.0 | 14.0 | | | 23, 30 |
| CCl_4 | (f) | 18.9 | 14.3 | | | 23 |
| | (h) | 20.0 | 14.0 | | | 28 |
| C_2H_5Cl | (f) | 21.5 | 14.3 | | | 23, 30 |
| CH_2ClCH_2Cl | (f) | 21.3 | 14.3 | | | 23, 30 |
| $CHCl_2CH_2Cl$ | (f) | 20.6 | 14.3 | | | 23, 30 |
| $CHCl_2CHCl_2$ | (f) | 20.4 | 14.3 | | | 23, 30 |
| $CHCl_2CCl_3$ | (h) | 19.0 | 14.5 | | | 28 |
| | (f) | 18.3 | 14.3 | | | 23, 30 |
| | (f) | 17.9 ± 1.0 | 13.8 ± 0.5 | | | 31 |
| C_2Cl_6 | (f) | 19.5 | 14.4 | | | 23 |
| | (h) | 18.0 | 13.5 | | | 28 |
| Halogens and pseudo-halogens | | | | | | |
| BrCl | (h) | 1.1 ± 0.4 | 12.3 | 20–60 | | 32 |
| ICl | | 4.5 | 11.7 | 30–60 | ICl P | 33, 35 |
| CNCl | | 34 | 14 | 1727–2527 | CICN S.T. | 34 |
| Carbonyls, etc. | | | | | | |
| COCl | | 0.83 | 14.6 | 25–55 | CO/Cl_2 P | 36 |
| $COCl_2$ | (h) | 19.9 | 14.4 | 25–55 | | 28 |
| | (a) | 23.5 | 14.5 | 15–450 | | 43 |

Metathetical Reactions of Bromine Atoms

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | References |
|---|---------|---------------------------------|--|-------------------|--|------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H ₂ | | 19.4 ± 0.2 | 14.36 | 227–302 | H ₂ /Br ₂ T | 1, 2 |
| | | 18.6 ± 0.6 | 13.93 | 277–327 | H ₂ /Br ₂ T | 3, 2 |
| | | 17.6 | 13.86 | 200–300 | H ₂ /Br ₂ P | 4 |
| | (a) | 19.8 | 13.36 | 700–1400 | | 6 |
| | | | $k = (0.89 \text{ and } 1.03) \times 10^{11}$ | 1123 | H ₂ /Br ₂ T | 7 |
| | | | $k = (0.91 \text{ and } 0.79) \times 10^{11}$ | 1168 | H ₂ /Br ₂ T | 7 |
| | (b) | 17.8 | 13.7 | | | 18 |
| | | 18.3 | 14.03 | 327–1197 | H ₂ /Br ₂ T | 8 |
| | | 19.2 ± 1.8 | 14.25 | 1027–1427 | H ₂ /Br ₂ T | 2 |
| | (b) | 19.7 ± 0.4 | 14.43 ± 0.14 | 227–1427 | | 9 |
| | | 19.17 | 14.24 | 832–1011 | H ₂ /O ₂ /CF ₃ Br ignition | 32 |
| | | 19.7 | 14.43 | | | |
| HD | (i) | 20.3 | 14.37 | 168–350 | HD/H ₂ /Br ₂ T, P | 34 |
| HT | (i) | 20.4 | 14.34 | 168–350 | HT/H ₂ /Br ₂ T, P | 34 |
| D ₂ | | 20.2 | 14.89 | 277–377 | H ₂ /Br ₂ T | 3, 2 |
| | | 20.4 ± 1.0 | 14.08 | 1027–1427 | H ₂ /Br ₂ T | 2 |
| | (i) | 21.0 | 14.31 | 168–350 | H ₂ /D ₂ /Br T, P | 34 |
| | (b) | 21.4 ± 0.4 | 14.29 | 277–1427 | | 9 |
| | | 21.4 | 14.29 | | | |
| Alkanes | | | | | | |
| CH ₄ | (c) (d) | 18.2 ± 0.5 | 13.8 | 150–210 | CH ₄ /Br ₂ P | 10 |
| | | 18.3 | 14.0 | 204–341 | CH ₄ /CH ₃ Cl/Br ₂ T, P | 11 |
| | | 17.3 | 13.15 | 150–300 | | 22 |
| | | 18.2 | 13.8 | | | |
| C ₂ H ₆ | (d) | 13.6 ± 0.5 | | 35–90 | C ₂ H ₆ /Br ₂ P | 13 |
| | (e) | 13.4 ± 0.1 | 13.90 ± 0.04 | 59–199 | C ₂ H ₆ /CH ₃ Br/Br ₂ T | 11, 14 |
| | | 12.3 | 13.29 | 25–121 | CH ₃ CHF ₂ /C ₂ H ₆ /Br ₂ T | 15 |
| C ₃ H ₈ | (d) | 10.15 ± 0.14 | 13.71 ± 0.07 | 13–145 | C ₃ H ₈ /C ₂ H ₆ /Br ₂ T | 11, 14 |
| <i>n</i> -C ₄ H ₁₀ | (d) | 10.23 ± 0.23 | 13.22 ± 0.14 | –6–98 | <i>n</i> -C ₄ H ₁₀ / <i>i</i> -C ₄ H ₁₀ /Br ₂ T | 11, 14, 20 |
| <i>i</i> -C ₄ H ₁₀ | (d) | 7.51 ± 0.20 | 13.30 ± 0.11 | 34–148 | <i>i</i> -C ₄ H ₁₀ /C ₃ H ₈ /Br ₂ T | 11, 14, 20 |
| | (h) | 11.7 | 17.6 | 40–95 | <i>i</i> -C ₄ H ₁₀ /Br ₂ P | 16, 17 |
| neo-C ₅ H ₁₂ | (d) | 14.29 ± 0.13 | 14.24 ± 0.06 | 57–200 | neo-C ₅ H ₁₂ /C ₂ H ₆ /Br ₂ T | 11 |
| | (h) | 18.2 | 17.0 | 98–152 | neo-C ₅ H ₁₂ /Br ₂ P | 19, 17 |
| Aromatic hydrocarbon | | | | | | |
| C ₆ H ₅ CH ₃ | (h) | 7.6 | 13.5 | 82–132 | C ₆ H ₅ CH ₃ /Br ₂ P | 21, 25, 17 |
| Halogenated alkanes | | | | | | |
| CH ₃ F | (e) | 14.8 | 12.86 | 150–300 | CH ₃ /CH ₃ F/Br ₂ P | 12 |
| CH ₂ F ₂ | (e) | 15.3 | 12.52 | 150–300 | CH ₃ /CH ₂ F ₂ /Br ₂ P | 12 |

Metathetical Reactions of Bromine Atoms – Continued

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | References |
|---|---------|----------------------------|---|--------------------|--|--------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CHF ₃ | (e) | 21.1 23.0 | 12.28 13.46 | 150–300 361–431 | CHF ₃ /C ₂ F ₅ H/Br ₂ P CF ₃ H/Br ₂ T | 12 22 |
| CH ₃ Cl | | 14.45 ± 0.16 | 13.62 ± 0.06 | 59–200 | CH ₃ Cl/C ₂ H ₆ /Br ₂ T, P | 11 |
| CHCl ₃ | | 9.3 10. | 12.36 | 147–182 | CHCl ₃ /Br ₂ T CHCl ₃ /Br ₂ P | 23, 17 24 |
| CH ₃ Br | | 16.05 | 13.7 | 150–230 | CH ₃ Br/Br ₂ P | 10, 11, 14 |
| CH ₃ CH ₂ F | (e) | 10.3 | 12.38 | 39–121 | C ₂ H ₆ /C ₂ H ₅ F/Br ₂ P | 15 |
| CH ₃ CHF ₂ | (e) | 13.3 | 12.52 | 96–230 | CH ₃ /C ₂ H ₄ F ₂ /Br ₂ P | 15 |
| CH ₃ CF ₃ | (e) | 22.2 | 13.18 | 243–379 | CF ₃ H/CH ₃ CF ₃ /Br ₂ P | 15 |
| CF ₃ CH ₂ F | (e) | 18.2 | 12.73 | 130–322 | CH ₂ F ₂ /CF ₃ CH ₂ F/ Br ₂ P | 15 |
| CF ₂ HCF ₂ H | (e) | 18.1 | 12.75 | 118–283 | CH ₂ F ₂ /CF ₃ CH ₂ F/ Br ₂ P | 15 |
| CF ₃ CF ₂ H | (e) | 18.0 | 12.20 | 150–300 | CH ₄ /C ₂ F ₅ H/Br ₂ | 12 |
| CF ₃ CF ₂ CF ₂ H | | 17.8 | 12.08 | 146–286 | CH ₄ /C ₃ F ₇ H/Br ₂ T | 15 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (f) | 11.6 | 13.2 | 62–185 | CH ₃ CH ₂ CH ₂ CH ₂ F/ Br ₂ P | 26 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (f) | 11.5 | 13.1 | 62–185 | CH ₃ CH ₂ CH ₂ CH ₂ F/ Br ₂ P | 26 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (f) | (10.2) | (13.6) | 62–185 | CH ₃ CH ₂ CH ₂ CH ₂ F/ Br ₂ P | 26 |
| CH ₃ CH ₂ CH ₂ CH ₂ F | (f) | 13.0 | 13.0 | 62–185 | CH ₃ CH ₂ CH ₂ CH ₂ F/ Br ₂ P | 26 |
| CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ | (g) (f) | 11.0 | 13.2 | 0–230 | CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ / Br ₂ P | 26 |
| CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ | (f) | (10.2) | (13.6) | 0–230 | CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ / Br ₂ P | 26 |
| CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ | (f) | 13.0 | 13.1 | 0–230 | CF ₃ CH ₂ CH ₂ CH ₂ CH ₃ / Br ₂ P | 26 |
| Alcohols | | | | | | |
| CH ₃ OH | | 6.3 | 11.65 | 76–135 | CH ₃ OH/Br ₂ P | 27 |
| CH ₃ CH ₂ OH | | 2.6 ± 2.0 | | 70–150 | C ₂ H ₅ OH/Br ₂ P | 28 |

CHLORINE ATOM TRANSFER

| | | | | | | |
|-----------------|--|-----------|-------------|-------|------------------------------------|----|
| Cl ₂ | | 6.9 ± 0.4 | 12.65 ± 0.2 | 20–60 | Br ₂ /Cl ₂ P | 29 |
|-----------------|--|-----------|-------------|-------|------------------------------------|----|

BROMINE ATOM TRANSFER

| | | | | | | |
|---------------------------------|--|------|------|---------|---------------------------------------|--------|
| Alkyl halides | | | | | | |
| CH ₃ Br | | 22.9 | 13.7 | 150–297 | CH ₃ /Br ₂ P, T | 10, 17 |
| CH ₂ Br ₂ | | | 14.0 | | | 10, 17 |

Metathetical Reactions of Bromine Atoms—Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|------------------------|-------|---------------------------------|--|--------------------|---------------------------------------|-------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| CCl ₃ Br | | 10.3 | 13.91 | 146–183 | CCl ₃ Br/Br ₂ T | 30 |
| Hydrogen halide | | | | | | |
| HBr | (a) | 43.8 41.7 | 14.4 13.9 | 700–1400 25–302 | H ₂ /Br ₂ T | 6 33, 17 |
| Halogen | | | | | | |
| BrCl | | $k = (3.1 \pm 0.8) \times 10^3$ | | 20 | Br ₂ /Cl ₂ P | 29 |

OXYGEN ATOM TRANSFER

| | | | | | | |
|------------------|--|----|------|---------|------------------------------------|----|
| N ₂ O | | 37 | 14.3 | 603–700 | N ₂ O/Br ₂ T | 31 |
|------------------|--|----|------|---------|------------------------------------|----|

MISCELLANEOUS REACTIONS

| | | | | | | |
|-------------------------------------|--|--------------------------|--|-------|--------------------------|----|
| Br + Na ₂ = NaBr + Na | | $k = 1.5 \times 10^{14}$ | | ~ 300 | Br ₂ /Na D.F. | 35 |
|-------------------------------------|--|--------------------------|--|-------|--------------------------|----|

Notes

- (a) Estimated from a general review of kinetic data.
 (b) Critical survey of literature data.
 (c) These values are less reliable than those given for other compounds, by the same authors.
 (d) All values are related to Br + CH₃Br = CH₂Br + HBr as standard, for which $k = 10^{13.73} \exp(-16050/RT)$.
 (e) Measured relative to CH₄ + Br = CH₃ + HBr for which $k = 10^{13.15} \exp(-17300/RT)$. This is a recalculation of data from ref. 10. To compare these values with the others, *E* should be increased by 1 kcal/mole and log *A* by 0.65 units.
 (f) The values at the δ or γ positions (bracketed) are assumed the same as for the 2 position in *n*-butane (ref. 11). The other values are calculated relative to this. These values must be considered very approximate.
 (g) Very small quantities of CF₃CHBrCH₂CH₂CH₃ were formed, but not in sufficient quantities for the Arrhenius factors to be estimated.
 (h) These values are believed to be in error (ref. 17)
 (i) Measured relative to Br + H₂ = H + HBr, for which $k = 10^{14.43} \exp(-19,700/RT)$.

References (Bromine Atoms)

1. Bodenstein and Lind, *Z. Phys. Chem.* **57**, 168 (1906).
2. Britton and Cole, *J. Phys. Chem.* **65**, 1302 (1961).
3. Bach, Bonhoeffer, and Moelwyn-Hughes, *Z. Phys. Chem.* **27B**, 71 (1935).
4. Bodenstein and Lutkemeyer, *Z. Phys. Chem.* **114**, 208 (1924).
5. Steiner, *Proc. Roy. Soc.* **173A**, 531 (1939).
6. Cooley and Anderson, *Ind. and Eng. Chem.* **44**, 1402 (1952).
7. Britton and Davidson, *J. Chem. Phys.* **23**, 2461 (1955).
8. Levy, *J. Phys. Chem.* **62**, 570 (1958).
9. Fettis and Knox, *Progress in Reaction Kinetics*, Vol. 2, p. 3 (Pergamon Press, 1964).
10. Kistiakowsky and Van Artsdalen, *J. Chem. Phys.* **12**, 469 (1944).
11. Fettis, Knox, and Trotman-Dickenson, *J. Chem. Soc.* 4177 (1960).

12. Tarr, Coomber, and Whittle, *Trans. Faraday Soc.* **61**, 1182 (1965).
13. Andersen and Van Artsdalen, *J. Chem. Phys.* **12**, 479 (1944).
14. Fettis and Trotman-Dickenson, *J. Am. Chem. Soc.* **81**, 5260 (1959).
15. Coomber and Whittle, *Trans. Faraday Soc.* **62**, 1553 (1966).
16. Eckstein, Scheraga, and Van Artsdalen, *J. Chem. Phys.* **22**, 28 (1954).
17. Benson and Buss, *J. Chem. Phys.* **28**, 301 (1958).
18. Campbell and Fristom, *Chem. Rev.* **58**, 173 (1958).
19. Hormats and Van Artsdalen, *J. Chem. Phys.* **19**, 778 (1951).
20. Anson, Fredricks, and Tedder, *J. Chem. Soc.* 918 (1959).
21. Anderson, Scheraga, and Van Artsdalen, *J. Chem. Phys.* **21**, 1258 (1953).
22. Corbett, Tarr, and Whittle, *Trans. Faraday Soc.* **59**, 1609 (1963).
23. Sullivan and Davidson, *J. Chem. Phys.* **19**, 143 (1951).
24. Braunworth and Schumacher, *Kolloidzshr.* **89**, 184 (1939).
25. Swegler, Scheraga, and Van Artsdalen, *J. Chem. Phys.* **19**, 135 (1951).
26. Galiba, Tedder, and Watson, *J. Chem. Soc.* 1321 (1964).
27. Buckley and Whittle, *Trans. Faraday Soc.* **58**, 536 (1962).
28. Tarr and Whittle, *Trans. Faraday Soc.* **60**, 2039 (1964).
29. Christie and Thrush, *Trans. Faraday Soc.* **55**, 1139 (1959).
30. Davidson and Sullivan, *J. Chem. Phys.* **17**, 176 (1949).
31. Kaufman, Gerri, and Pascale, *J. Chem. Phys.* **24**, 32 (1956).
32. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
33. Bodenstein and Jung, *Z. Phys. Chem.* **121**, 127 (1926).
34. Timmons and Weston, *J. Chem. Phys.* **41**, 1654 (1964).
35. Polanyi, "Atomic Reactions." (Williams and Norgate, London, 1932).

Review

1. Fettis and Knox, "Progress in Reaction Kinetics," Vol. 2, p. 3 (Pergamon Press, London, 1964).

Metathetical Reactions of Iodine Atoms

| Reactants | Notes | E | $\log_{10}A$ | Temperature range | Radical source | References |
|--|-------|---------------------|-------------------------------|-------------------|---|------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | $(kcal\ mole^{-1})$ | $(cm^3\ mole^{-1}\ sec^{-1})$ | $^{\circ}C$ | | |
| H ₂ | | 33.9 ± 0.3 | 14.32 ± 0.11 | 105-360 | H ₂ /I ₂ T | 1 |
| | | 33.5 ± 0.2 | 14.20 ± 0.07 | 394-527 | H ₂ /I ₂ T | 2 |
| | | | $k = 4.6 \times 10^6$ | 700 | HI and H ₂ /I ₂ T | 3, 4 |
| D ₂ | | 34.5 ± 0.3 | 14.06 ± 0.10 | 394-527 | D ₂ /I ₂ T | 5 |
| Alkanes | | | | | | |
| CH ₄ | (a) | 33.5 | 14.70 | 260-316 | | 6, 7 |
| | | 34.1 | 14.70 | 260-316 | CH ₄ /I ₂ T | 7 |
| | | 35.0 ± 1.1 | 14.95 | 275-345 | CH ₄ /I ₂ T | 8 |
| C ₂ H ₆ | | 27.9 | 14.22 | 263-303 | C ₂ H ₅ I/HI T | 9 |
| C ₃ H ₈ | | 25.5 ± 1 | 14.53 ± 0.36 | 307-340 | C ₃ H ₇ /I ₂ T | 10 |
| CH ₃ CH ₂ CH ₃ | | 25.0 | 14.22 | 307-340 | C ₃ H ₇ /I ₂ T | 10 |
| <i>i</i> -C ₄ H ₁₀ | | 21.4 ± 0.5 | 13.88 ± 0.15 | 252-310 | <i>i</i> -C ₄ H ₉ /I ₂ T | 11 |
| Alkenes | | | | | | |
| CH ₂ :CHCH ₃ | | 18.04 ± 0.32 | 13.25 ± 0.14 | 208-300 | CH ₂ :CHCH ₃ /I ₂ T | 22 |
| 1-CH ₂ :CHCH ₂ CH ₃ | | 13.2 ± 0.4 | 12.83 ± 0.15 | 204-253 | CH ₂ :CHCH ₂ CH ₃ /I ₂ T | 12 |
| Aromatic hydrocarbon | | | | | | |
| C ₆ H ₅ CH ₃ | | | $k = 4.8 \times 10^7$ | 501 | C ₂ H ₅ I/C ₆ H ₅ CH ₃ T | 13 |
| Aldehyde | | | | | | |
| CH ₃ CHO | | 15.7 | 13.3 | 222-268 | CH ₃ COI/HI T | 14 |
| Alcohol | | | | | | |
| (CH ₃) ₂ CHOH | | 20.5 | 14.07 | 207-300 | (CH ₃) ₂ CHOH/I ₂ T | 23 |

IODINE ATOM TRANSFER

| | | | | | | |
|--|--|------------|--------------------------|---------|---|--------|
| Alkyl halides | | | | | | |
| CH ₃ I | | 19.8 | 14.3 | 270-320 | CH ₃ I/HI T | 15, 16 |
| | | ≥ 20.5 | $(k = 1.60 \times 10^3)$ | 280 | CH ₃ I/HI T | 17, 16 |
| | | 20.5 ± 0.5 | 14.4 ± 0.2 | 260-316 | CH ₃ I/HI T | 6 |
| | | 19.2 | 13.71 | | CH ₃ I/HI T | 18 |
| CF ₃ I | | 17.60 | 13.8 | 167-485 | CF ₃ I/HI T | 18 |
| C ₂ H ₅ I | | 16.7 | 13.62 | 250-300 | C ₂ H ₅ I/HI T | 15, 16 |
| | | ≥ 18.2 | $(k = 6.8 \times 10^3)$ | 260 | C ₂ H ₅ I/HI T | 17, 16 |
| | | 17.1 ± 0.7 | 14.01 ± 0.28 | 263-303 | C ₂ H ₅ I/HI T | 9 |
| <i>n</i> -C ₃ H ₇ I | | ≥ 18.5 | $(k = 12.9 \times 10^3)$ | 290 | <i>n</i> -C ₃ H ₇ I/HI T | 17, 16 |
| ICH ₂ CH ₂ CH ₂ I | | 19.5 | 14.0 | | | 19 |
| (CH ₃) ₃ CI | | 13.0 ± 0.7 | 13.7 ± 0.4 | 252-310 | (CH ₃) ₃ CH/I ₂ T | 11 |

Metathetical Reactions of Iodine Atoms – Continued

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | References |
|--------------------------------|-------|---------------------------------|--|--------------------|--|------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| Acetyl halide | | | | | | |
| CH ₃ COI | | 14.6 | 13.9 | 222–268 | CH ₃ COI/HI T | 14 |
| Hydrogen halide | | | | | | |
| HI | | 36.4 36.5 | 14.32 ± 0.20 | 105–360 394–527 | H ₂ /I ₂ T H ₂ /I ₂ T | 1 2 |
| OXYGEN ATOM TRANSFER | | | | | | |
| N ₂ O | | 38 | 14.45 | 603–700 | N ₂ O/I ₂ T | 20 |
| MISCELLANEOUS REACTIONS | | | | | | |
| I + Na ₂ = NaI + Na | | | $k = 2.6 \times 10^{13}$ | ~ 300 | I ₂ /Na D.F. | 21 |

Notes

- (a) Calculated from the reverse reaction.
 (b) Estimated from thermodynamic data.

References (Iodine Atoms)

1. Sullivan, J. Chem. Phys. **30**, 1292 (1959).
2. Sullivan, J. Chem. Phys. **36**, 1925 (1962).
3. Horie, Ishii, and Amano, J. Phys. Chem. **68**, 1264 (1964).
4. Graven, J. Am. Chem. Soc. **78**, 3297 (1956).
5. Sullivan, J. Chem. Phys. **39**, 3001 (1963).
6. Flowers and Benson, J. Chem. Phys. **38**, 882 (1963).
7. Golden, Walsh, and Benson, J. Am. Chem. Soc. **87**, 4053 (1965).
8. Goy and Pritchard, J. Phys. Chem. **69**, 3040 (1965).
9. Hartley and Benson, J. Chem. Phys. **39**, 132 (1963).
10. Nangia and Benson, J. Am. Chem. Soc. **86**, 2773 (1964).
11. Teranishi and Benson, J. Am. Chem. Soc. **85**, 2887 (1963).
12. Benson, Bose, and Nangia, J. Am. Chem. Soc. **85**, 1388 (1963).

13. Yang and Conway, J. Chem. Phys. **43**, 1296 (1965).
14. O'Neal and Benson, J. Chem. Phys. **37**, 540 (1962).
15. Sullivan, J. Phys. Chem. **65**, 722 (1961).
16. Ogg, J. Am. Chem. Soc. **56**, 526 (1934).
17. Benson and O'Neal, J. Chem. Phys. **34**, 514 (1961).
18. Body, Downs, Gow, and Horrex, J. Phys. Chem. **67**, 719 (1963).
19. Benson, J. Chem. Phys. **34**, 521 (1961).
20. Kaufman, Gerri, and Pascale, J. Chem. Phys. **24**, 32 (1956).
21. Polanyi, "Atomic Reactions" (Williams and Norgate, London, 1932).
22. Golden, Rodgers, and Benson, J. Am. Chem. Soc. **88**, 3196 (1966).
23. Walsh and Benson, J. Am. Chem. Soc. **88**, 3480 (1966).

Review

1. Fettis and Knox, "Progress in Reaction Kinetics," Vol. 2, p. 3 (Pergamon Press, London, 1964).

Ratios of Rate Constants (Sulphur Atoms)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature Range | Radical Source | References |
|--|-------|---------------------------------|---------------------|-------------|-------------------|----------------------|------------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) S(¹ D) + COS = CO + S ₂ (2) S(¹ D) + CH ₄ = CH ₃ SH | (a) | | | 17.8 | 25 | COS P | 1 |
| (1) S(¹ D) + COS = CO + S ₂ (2) S(¹ D) + C ₂ H ₆ = C ₂ H ₅ SH | | | | 2.2 2.04 | 25 25 | COS P COS P | 2 1 |
| (1) S(¹ D) + COS = CO + S ₂ (2) S(¹ D) + C ₃ H ₈ = C ₃ H ₇ SH | | | | 1.9 | 25 | COS P | 2 |
| (1) S(¹ D) + COS = CO + S ₂ (2) S(¹ D) + <i>i</i> -C ₄ H ₁₀ = C ₄ H ₉ SH | | | | 2.04 | 25 | COS P | 1 |

Notes

- (a) Ratio decreases with pressure.
 (b) Data also given (ref. 2) on S atom insertion into cyclo-C₃H₆, cyclo-C₄H₈, and cyclo-C₅H₁₀.

References

1. Knight, Strauss, and Gunning, *J. Am. Chem. Soc.* **85**, 2349 (1963).
2. Knight, Strauss, Malm, and Gunning, *J. Am. Chem. Soc.* **86**, 4243 (1964).

Reactions of Sodium Atoms

Index

| | Page |
|--|-----------|
| (a) Fluorine Atom Transfer | |
| Reactions with | |
| 1. Halogenated alkanes..... | 29 |
| 2. Halogenated alkenes..... | 29 |
| 3. Halogenated acids and acid halides..... | 29 |
| 4. Halogenated aromatic derivatives..... | 29 |
| 5. Polyfluorides..... | 29 |
| (b) Chlorine Atom Transfer | |
| Reactions with | |
| 1. Halogenated alkanes..... | 29 |
| 2. Halogenated alkenes..... | 31 |
| 3. Halogenated aromatic derivatives..... | 31 |
| 4. Cyanides..... | 32 |
| 5. Halogenated acids, acid chlorides, esters, etc..... | 32 |
| 6. Polyhalides..... | 32 |
| 7. Oxychlorides..... | 33 |
| 8. Halogens and pseudo-halogens..... | 33 |
| 9. Hydrogen halide..... | 33 |
| (c) Bromine Atom Transfer | |
| Reactions with | |
| 1. Halogenated alkanes..... | 33 |
| 2. Halogenated alkenes..... | 34 |
| 3. Halogenated aromatic derivatives..... | 34 |
| 4. Acid bromides..... | 35 |
| 5. Halogens and pseudo-halogens..... | 35 |
| 6. Hydrogen halides..... | 35 |
| (d) Iodine Atom Transfer | |
| Reactions with | |
| 1. Halogenated alkanes..... | 35 |
| 2. Halogenated alkenes..... | 36 |
| 3. Halogenated aromatic derivatives..... | 36 |
| 4. Halogens..... | 36 |
| 5. Hydrogen halides..... | 36 |
| (e) Cyanide Group Transfer..... | 36 |

Metathetical Reactions of Sodium Atoms

| Reactants | Notes | (a) E | $\log_{10} k$ | Temperature range | Radical source | References |
|--|----------------|----------------------------|--|-------------------|----------------|--------------|
| FLUORINE ATOM TRANSFER | | | | | | |
| Halogenated alkanes | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CF ₂ | (b) (e) (e) | | 6.5 log (k_f) ^{1/2} 5.0 log (k_f) ^{1/2} | 313 310 | D.F. D.F. | 4 5 |
| CFH ₂ | (b) (c) | | 6.1 log (k_c) ^{1/2} | 313 | D.F. | 4 |
| CF ₂ H | (b) (d) | | 6.2 log (k_c) ^{1/2} | 313 | D.F. | 4 |
| CF ₃ | (b) | | 13.83 13.97 | 313 310 | D.F. D.F. | 4 5 |
| CFH ₃ | | > 25 ~ 18.5 | < 8.7 7.3 | 240 247 | D.F. | 1, 6 2, 3 |
| CH ₂ F ₂ | | 14.0 | 8.8 | 247 | | 2, 3 |
| CHF ₃ | | 14.0 | 8.8 | 247 | | 2, 3 |
| CF ₄ | | 12.6 | 9.4 | 247 | | 2, 3 |
| cyclo-C ₆ F ₁₂ | | 7.1 | 11.7 | 247 | | 2, 3 |
| cyclo-C ₆ F ₁₁ CF ₃ | | 6.7 | 12.0 | 247 | | 2, 3 |
| Alkenes | | | | | | |
| C ₂ F ₄ | | 8.8 | 11.0 | 247 | | 2, 3 |
| Acids and acid halides | | | | | | |
| CF ₃ COOH | | 4.0 | 13.0 | 247 | | 2, 3 |
| CF ₃ COF | | 7.6 | 11.5 | 247 | | 2, 3 |
| Halogenated aromatics | | | | | | |
| C ₆ H ₅ F | | | < 8.7 | 247 | | 19 |
| Miscellaneous | | | | | | |
| SF ₆ | | 3.3 | 13.3 | 247 | | 2, 3 |

CHLORINE ATOM TRANSFER

| | | | | | | |
|----------------------------|-----|------------|----------------------|-------------------|----------------------|----------------|
| Halogenated alkanes | | | | | | |
| CH ₃ Cl | | 7.5 | 10.7 10.9 11.8 | 240 270 325 | D.F. M.D.F. | 1, 6 7 8 |
| | (f) | 9.8 8.2 | 11.1 11.8 | 313 220 | D.F. D.F. M.L. | 9 10 11 |

Metathetical Reactions of Sodium Atoms—Continued

| Reactants | Notes | ^(a) <i>E</i> (<i>kcal mole</i> ⁻¹) | $\log_{10} k$ (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) | Temperature range °C | Radical source | References |
|--|---------|---|--|-----------------------------|----------------|------------|
| CH ₂ Cl ₂ | | | 11.75 | 275 | D.F. | 1, 6 |
| | | | 12.4 | 270 | | 7 |
| | | | 11.8 | 250 | D.F. | 12 |
| CHCl ₃ | | | 12.7 | 275 | D.F. | 1, 6 |
| | | | 13.5 | 270 | | 7 |
| | | | 13.0 | 250 | D.F. | 12 |
| CCl ₄ | | | 13.3 | 275 | D.F. | 1, 6 |
| | | | 14.5 | 270 | | 7 |
| | | | 14.0 | 250 | D.F. | 12 |
| | (g) (f) | 8.4 | 12.17 | 310 | M.D.F. | 13 |
| | | | 14.35 | 247 | D.F. | 14 |
| CH ₂ FCI | (g) (f) | 10.1 | 10.97 | 313 | D.F. | 9 |
| CHF ₂ Cl | | | 10.0 | 247 | | 3 |
| | (g) (f) | 10.0 | 11.0 | 313 | D.F. | 9 |
| CF ₃ Cl | | | 10.7 | 247 | | 3 |
| | | 9.2 | 11.3 | 313 | D.F. | 9 |
| | (f) | 7.4 | | 275–315 | D.F. | 15 |
| | (g) (f) | 10.2 | 10.9 | 310 | M.D.F. | 13 |
| CF ₂ Cl ₂ | (g) (f) | 9.0 to 9.5 | 11.5 to 11.7 | 310 | M.D.F. | 13 |
| CFCl ₃ | (g) (f) | 8.7 to 9.2 | 11.8 to 11.9 | 310 | M.D.F. | 13 |
| C ₂ H ₅ Cl | | | 10.9 | 275 | D.F. | 6, 1 |
| | | 10.2 | | 260–380 | D.F. | 16, 17 |
| | | 7.8 | | 292–391 | D.F. | 10 |
| CH ₂ ClCH ₂ Cl | | | 11.8 | 275 | D.F. | 6 |
| | (a) | 7.0 | 12.0 | 285 | D.F. | 18 |
| CH ₃ CHCl ₂ | | | 11.9 | 275 | D.F. | 6 |
| CH ₃ CCl ₃ | | | 13.6 | 247 | D.F. | 14 |
| CH ₃ CH ₂ CH ₂ Cl | (a) | 9.1 | 11.1 | 275 | D.F. | 6 |
| | (a) | 9.0 | 11.0 | 261 | D.F. | 18 |
| | | | 10.7 | 267 | D.F. | 19 |
| CH ₃ CHClCH ₃ | | | 11.2 | 275 | D.F. | 6 |
| CH ₂ ClCH ₂ CH ₂ Cl | | | 11.7 | 275 | D.F. | 6 |
| CH ₃ CHClCH ₂ Cl | | | 12.0 | 275 | D.F. | 6 |
| CH ₃ CH ₂ CHCl ₂ | | | 12.2 | 275 | D.F. | 6 |
| (CH ₃) ₂ CCl ₂ | | | 12.4 | 275 | D.F. | 6 |
| | | | 12.6 | 247 | D.F. | 14 |
| CH ₃ CH ₂ CH ₂ CH ₂ Cl | | | 11.2 | 275 | D.F. | 6 |
| CH ₃ CH ₂ CHClCH ₃ | | | 11.4 | 275 | D.F. | 6 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) <i>E</i> | $\log_{10} k$ | Temperature range | Radical source | References |
|---|-------|---------------------------------|--|-------------------|----------------------|---------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| (CH ₃) ₂ CHCH ₂ Cl | | | 11.2 | 275 | D.F. | 6 |
| (CH ₃) ₃ CCl | | | 11.5 11.4 | 275 247 | D.F. D.F. | 6 14 |
| CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ Cl | | | 11.4 | 275 | D.F. | 6 |
| (CH ₃) ₂ CHCH ₂ CH ₂ Cl | | | 11.4 | 275 | D.F. | 6 |
| (CH ₃) ₂ CClCH ₂ CH ₃ | | | 11.9 | 275 | D.F. | 6 |
| Alkenes | | | | | | |
| CH ₂ :CHCl | | | 10.7 | 275 | D.F. | 6 |
| trans-CHCl:CHCl | | | 11.3 | 275 | D.F. | 6 |
| cis-CHCl:CHCl | | | 11.4 | 275 | D.F. | 6 |
| CH ₂ :CHCH ₂ Cl | | | 12.3 12.5 | 275 260 | D.F. D.F. | 6 18 |
| CH ₂ :CClCH ₃ | | | 11.0 | 275 | D.F. | 6 |
| C ₆ H ₅ CH:CHCl | | | 13.0 | 275 | D.F. | 18 |
| C ₆ H ₅ CH:CHCH ₂ Cl | | | 13.9 | 275 | D.F. | 18 |
| Aromatic chlorides and their derivatives | | | | | | |
| C ₆ H ₅ Cl | | | 9.4 11.2 11.8 | 247 244 270 | D.F. M.L. D.F. | 19 25 1 |
| C ₆ H ₅ CH ₂ Cl | | | 14.7 13.7 | 275 284 | D.F. D.F. | 6 18 |
| C ₆ H ₅ CH ₂ CH ₂ Cl | | | 13.4 | 285 | D.F. | 18 |
| <i>o</i> -FC ₆ H ₄ Cl | | | 10.3 | 247 | D.F. | 19 |
| <i>m</i> -FC ₆ H ₄ Cl | | | 9.8 | 247 | D.F. | 19 |
| <i>p</i> -FC ₆ H ₄ Cl | | | 9.1 | 247 | D.F. | 19 |
| <i>o</i> -ClC ₆ H ₄ Cl | | | 11.2 | 247 | D.F. | 19 |
| <i>o</i> -CH ₃ OOC ₆ H ₄ Cl | | | 13.4 | 247 | D.F. | 19 |
| 2:Cl-pyridine | | | 12.7 | 247 | D.F. | 19 |
| 3:Cl-pyridine | | | 10.0 | 247 | D.F. | 19 |
| Cyanides | | | | | | |
| CNCH ₂ Cl | | | 13.9 | 285 | D.F. | 18 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) <i>E</i> | $\log_{10} k$ | Temperature range | Radical source | References |
|--|-------|------------------------------------|---|-------------------|----------------|------------|
| | | (<i>kcal mole</i> ⁻¹) | (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) | °C | | |
| CNCH ₂ CH ₂ Cl | | | 11.8 | 285 | D.F. | 18 |
| Acids, acid chlorides, esters, etc. | | | | | | |
| CH ₂ ClCOOH | | | 13.3 | 247 | | 3 |
| CH ₃ COCl | | | 12.7 | 275 | D.F. | 6, 1 |
| | 0 | | 13.4 | 247 | D.F. | 3, 2 |
| | | | | | D.F. | 10 |
| C ₆ H ₅ COCl | | | 14.7 | 240 | D.F. | 1 |
| ClCOOC ₂ H ₅ | | | 11.6 | 247 | D.F. | 3, 2 |
| CH ₂ ClCOOC ₂ H ₅ | | | 13.2 | 247 | D.F. | 3, 2 |
| ClCH ₂ CH ₂ OH | | | 11.7 | 285 | D.F. | 18 |
| CH ₃ COCH ₂ Cl | | | 13.7 | 275 | D.F. | 6 |
| | | | 14.0 | 247 | D.F. | 2 |
| CH ₃ OCH ₂ Cl | | | 11.3 | 247 | | 3, 2 |
| Polyhalides | | | | | | |
| BCl ₃ | | | 11.7 | 270 | D.F. | 7 |
| SiCl ₄ | | | 11.6 | 270 | D.F. | 7 |
| | | | 11.8 | 247 | D.F. | 14 |
| CH ₃ SiCl ₃ | | | 9.8 | 247 | D.F. | 14 |
| (CH ₃) ₂ SiCl ₂ | | | 9.3 | 247 | D.F. | 14 |
| (CH ₃) ₃ SiCl | | | 9.4 | 247 | D.F. | 14 |
| PCl ₃ | | | 14.2 | 270 | D.F. | 7 |
| TiCl ₄ | | | 14.0 | 270 | D.F. | 7 |
| GeCl ₄ | | | 14.7 | 270 | D.F. | 7 |
| AsCl ₃ | | | 14.2 | 270 | D.F. | 7 |
| SnCl ₄ | | | 14.5 | 270 | D.F. | 7 |
| SCl ₂ | | | 14.8 | 270 | D.F. | 7 |
| S ₂ Cl ₂ | | | 14.2 | 270 | D.F. | 7 |
| Oxychlorides | | | | | | |
| COCl ₂ | | | 13.9 | 270 | D.F. | 7 |
| CrO ₂ Cl ₂ | | | 14.1 | 270 | D.F. | 7 |
| POCl ₃ | | | 14.4 | 270 | D.F. | 7 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) E | $\log_{10} k$ | Temperature range | Radical source | References |
|-------------------------------------|-------|----------------------------|---|-------------------|----------------|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| Halogens and pseudo-halogens | | | | | | |
| Cl ₂ | | | 14.6 | | Dil. F. | 20 |
| CNCl | | | 14.0 | 285 | D.F. | 18 |
| | | | 13.9 | 250 | D.F. | 1 |
| Hydrogen halide | | | | | | |
| HCl | | | 12.4 | 327 | Dil. F. | 21, 20 |

BROMINE ATOM TRANSFER

| | | | | | | |
|----------------------------------|-----|-----|------|-----|------|------|
| Halogenated alkanes | | | | | | |
| CH ₃ Br | (h) | 3.4 | 13.3 | 240 | D.F. | 1, 6 |
| | | | 12.8 | 270 | | 7 |
| | | | 12.6 | 255 | D.F. | 12 |
| | | | 12.2 | 200 | M.L. | 11 |
| | | | 12.8 | 242 | M.L. | 11 |
| | | | | | | 10 |
| CH ₂ Br ₂ | | | 13.6 | 255 | D.F. | 12 |
| | | | 13.5 | 255 | M.L. | 12 |
| CHBr ₃ | | | 14.6 | 255 | D.F. | 12 |
| | | | 13.9 | 255 | M.L. | 12 |
| CFBr ₃ | | | 14.3 | 255 | D.F. | 12 |
| CF ₃ Br | (a) | 2.3 | | 285 | D.F. | 15 |
| CHFBr ₂ | | | 13.7 | 255 | M.L. | 12 |
| CH ₂ ClBr | | | 13.3 | 255 | D.F. | 12 |
| | | | 13.3 | 255 | M.L. | 12 |
| CHClBr ₂ | | | 14.3 | 255 | D.F. | 12 |
| | | | 13.7 | 255 | M.L. | 12 |
| CHCl ₂ Br | | | 14.0 | 255 | D.F. | 12 |
| | | | 13.6 | 255 | M.L. | 12 |
| CCl ₃ Br | | | 14.5 | 255 | D.F. | 12 |
| | | | 14.0 | 255 | M.L. | 12 |
| C ₂ H ₅ Br | | | 12.8 | 240 | D.F. | 1 |
| | | | 12.4 | 247 | D.F. | 19 |
| | | | 12.7 | 263 | D.F. | 18 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) <i>E</i> | $\log_{10} k$ | Temperature range | Radical source | References |
|---|-------|----------------------------|---|-------------------|----------------|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CH ₂ BrCH ₂ Br | | | 13.3 | 247 | | 3 |
| cyclo-C ₃ H ₅ Br | | | 12.0 | 247 | D.F. | 22 |
| cyclo-C ₄ H ₇ Br | | | 12.6 | 247 | D.F. | 22 |
| cyclo-C ₅ H ₉ Br | | | 13.1 | 247 | D.F. | 22 |
| CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ Br | | | 12.8 | 247 | D.F. | 22 |
| cyclo-C ₆ H ₁₁ Br | | | 12.7 | | | 23 |
| | | | 12.7 | 247 | D.F. | 22 |
| Alkenes | | | | | | |
| CH ₂ :CHBr | | | 12.4 | 269 | D.F. | 18 |
| | | | 11.2 | 247 | D.F. | 22 |
| C ₆ H ₅ CH:CHBr | | | 13.1 | 270 | D.F. | 18 |
| C ₆ H ₅ CH:CHCH ₂ Br | | | 14.6 | 285 | D.F. | 18 |
| Aromatic bromides and their derivatives | | | | | | |
| C ₆ H ₅ Br | | | 12.3 | 255 | M.L. | 24 |
| | | | 11.4 | 247 | D.F. | 19 |
| | | | 13.2 | 244 | M.L. | 25 |
| | | | 13.4 | 260 | D.F. | 1 |
| | | | 12.1 | 247 | D.F. | 14 |
| <i>o</i> -ClC ₆ H ₄ Br | | | 12.6 | 247 | D.F. | 19 |
| <i>m</i> -ClC ₆ H ₄ Br | | | 11.9 | 247 | D.F. | 19 |
| <i>p</i> -ClC ₆ H ₄ Br | | | 11.6 | 247 | D.F. | 19 |
| <i>o</i> -BrC ₆ H ₄ Br | | | 13.4 | 247 | D.F. | 19 |
| <i>o</i> -CNC ₆ H ₄ Br | | | 13.3 | 247 | D.F. | 19 |
| <i>m</i> -CNC ₆ H ₄ Br | | | 12.6 | 247 | D.F. | 19 |
| <i>p</i> -CNC ₆ H ₄ Br | | | 12.8 | 247 | D.F. | 19 |
| <i>o</i> -HOC ₆ H ₄ Br | | | 12.2 | 247 | D.F. | 19 |
| <i>m</i> -HOC ₆ H ₄ Br | | | 11.8 | 247 | D.F. | 19 |
| <i>o</i> -CH ₃ C ₆ H ₄ Br | | | 11.5 | 247 | D.F. | 19 |
| <i>m</i> -CH ₃ C ₆ H ₄ Br | | | 11.2 | 247 | D.F. | 19 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) E | $\log_{10} k$ | Temperature range | Radical source | References |
|--|-------|---------------------|-------------------------------|-------------------|----------------|------------|
| | | $(kcal\ mole^{-1})$ | $(cm^3\ mole^{-1}\ sec^{-1})$ | $^{\circ}C$ | | |
| <i>p</i> -CH ₃ C ₆ H ₄ Br | | | 11.3 | 247 | D.F. | 19 |
| <i>o</i> -CH ₃ OC ₆ H ₄ Br | | | 11.9 | 247 | D.F. | 19 |
| <i>p</i> -CH ₃ OC ₆ H ₄ Br | | | 11.6 | 247 | D.F. | 19 |
| <i>o</i> -CH ₃ OOC ₆ H ₄ Br | | | 13.8 | 247 | D.F. | 19 |
| <i>m</i> -CH ₃ OOC ₆ H ₄ Br | | | 12.6 | 247 | D.F. | 19 |
| <i>p</i> -CH ₃ OOC ₆ H ₄ Br | | | 12.6 | 247 | D.F. | 19 |
| α -Bromonaphthalene | | | 12.7 | 247 | | 3 |
| β -Bromonaphthalene | | | 12.1 | 247 | | 3 |
| 2:Br-pyridine | 4 | | 13.3 | 247 | D.F. | 19 |
| 3:Br-pyridine | | | 11.9 | 247 | D.F. | 19 |
| C ₆ H ₅ CH ₂ CH ₂ Br | | | 13.6 | 280 | D.F. | 18 |
| Acid bromide | | | | | | |
| CH ₃ COBr | | | 13.9 | 300 | D.F. | 1 |
| Halogens and pseudo-halogens | | | | | | |
| CNBr | | | 14.1 | 273 | D.F. | 18 |
| Hydrogen halide | | | | | | |
| HBr | | | 13.8 | 327 | Dil. F. | 21, 20 |

IODINE ATOM TRANSFER

| Halogenated alkanes | Notes | (a) E | $\log_{10} k$ | Temperature range | Radical source | References |
|---|-------|---------------------|-------------------------------|-------------------|----------------|------------|
| | | $(kcal\ mole^{-1})$ | $(cm^3\ mole^{-1}\ sec^{-1})$ | $^{\circ}C$ | | |
| CH ₃ I | | | 14.7 | 240 | D.F. | 6 |
| | | | 14.5 | 240 | M.L. | 11 |
| | | | 13.7 | 250 | M.L. | 12 |
| | | | 13.6 | 255 | D.F. | 12 |
| | | | 13.6 | 277 | D.F. | 23 |
| | 0 | | | | D.F. | 10 |
| CF ₃ I | | | 14.0 | 247 | | 3 |
| | | | | 285–295 | D.F. | 15 |
| C ₂ H ₅ I | | | 14.0 | 240–270 | D.F. | 1 |
| | | | 14.3 | 276 | D.F. | 18 |
| CH ₃ CH ₂ CH ₂ I | | | 13.4 | 240 | D.F. | 1 |

Metathetical Reactions of Sodium Atoms – Continued

| Reactants | Notes | (a) E | $\log_{10} k$ | Temperature range | Radical source | References |
|---------------------------------|-------|----------------------------|---|-------------------|----------------|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| Alkenes | | | | | | |
| CH ₂ :CHI | | | 13.4 | 263 | D.F. | 18 |
| Aromatic iodides | | | | | | |
| C ₆ H ₅ I | | | 14.7 | 240 | D.F. | 1 |
| | | | 14.3 | 244 | M.L. | 25 |
| Halogens | | | | | | |
| I ₂ | | | 14.8 | | Dil. F. | 20 |
| Hydrogen halide | | | | | | |
| HI | | | 14.7 | 327 | Dil. F. | 20, 21 |

CYANIDE GROUP TRANSFER

| | | | | | | |
|--|--|--|-------|-----|--|---|
| CH ₃ CN | | | < 6.6 | 247 | | 3 |
| C ₆ H ₅ CH ₂ CN | | | 9.8 | 247 | | 3 |
| CH ₂ (CN)COOC ₂ H ₅ | | | 11.6 | 247 | | 3 |

Notes

(a) The activation energies E are calculated from the formula $k = 10^{14.7} \exp(-E/RT)$.

(b) These values are lower limits.

(c) k_e is the rate constant for the combination reaction $2CFH_2 = C_2F_2H_4$.

(d) k_c is the rate constant for the combination reaction $2CF_2H = C_2F_4H_2$.

(e) k_f is the rate constant for the combination reaction $2CF_2 = C_2F_4$.

(f) Calculated assuming a steric factor of unity.

(g) These values refer to the rate constants for the primary process.

(h) Estimated from a critical survey of literature data.

References

- Von Hartel and Polanyi, *Z. Phys. Chem.* **11B**, 97 (1930).
- Warhurst, *Quart. Rev. Chem. Soc.* **5**, 44 (1951).
- Trotman-Dickenson, "Gas Kinetics," p. 219 (Butterworths, London, 1955).
- Reed and Rabinovitch, *J. Phys. Chem.* **61**, 598 (1957).
- Kaufman and Reed, *J. Phys. Chem.* **67**, 896 (1963).
- Hartel, Meer, and Polanyi, *Z. Phys. Chem.* **19B**, 139 (1933).
- Heller and Polanyi, *Trans. Faraday Soc.* **32**, 633 (1936).
- Reed and Rabinovitch, *J. Chem. Phys.* **27**, 988 (1957).

- Reed and Rabinovitch, *J. Phys. Chem.* **61**, 598 (1957).
- Kerr, Lissi, and Trotman-Dickenson, *J. Chem. Soc.* 1673 (1964).

- Frommer and Polanyi, *Trans. Faraday Soc.* **30**, 519 (1934).

- Haresnape, Stevels, and Warhurst, *Trans. Faraday Soc.* **36**, 465 (1940).

- Kaufman and Reed, *J. Phys. Chem.* **67**, 896 (1963).

- Gowenlock and Thomas, *J. Chem. Soc.* 5068 (1965).

- Hodgins and Haines, *Can. J. Chem.* **30**, 473 (1952).

- Cvetanovic and Le Roy, *J. Chem. Phys.* **20**, 1016 (1952).

- Cvetanovic and Le Roy, *Can. J. Chem.* **29**, 597 (1951).

- Evans and Walker, *Trans. Faraday Soc.* **40**, 384 (1944).

- Riding, Scanlon, and Warhurst, *Trans. Faraday Soc.* **52**, 1354 (1956).

- Polanyi, "Atomic Reactions" (Williams and Norgate, London, 1932).

- Schay, *Z. Phys. Chem.* **11B**, 291 (1930).

- Friswell, Gowenlock, and Thomas, *J. Chem. Soc.* 6323 (1965).

- Whittle (Ph. D. Thesis, Manchester, 1951).

- Warhurst, *Trans. Faraday Soc.* **35**, 674 (1939).

- Fairbrother and Warhurst, *Trans. Faraday Soc.* **31**, 987 (1935).

Reviews

- Polanyi, "Atomic Reactions" (Williams and Norgate, London, 1932).
- Bawn, *Ann. Rep. Chem. Soc.* **39**, 36 (1942).
- Warhurst, *Quart. Rev. Chem. Soc.* **5**, 44 (1951).

Metathetical Reactions of Potassium Atoms

| Reactants | Notes | <i>E</i> | $\log_{10} k$ | Temperature range | Radical source | Reference |
|---|------------|----------|---------------------------------------|-------------------|----------------|-----------|
| CHLORINE ATOM TRANSFER | | | | | | |
| $(kcal\ mole^{-1})$ HCl Cl ₂ | (b) | | $(cm^3\ mole^{-1}\ sec^{-1})$ 14.0 | °C 327 | Dil F. | 1, 2 |
| BROMINE ATOM TRANSFER | | | | | | |
| HBr Br ₂ | (a) (b) | 3.4 | 14.6 | 327 | Dil F. M.B. | 1, 2 3 |
| IODINE ATOM TRANSFER | | | | | | |
| HI CH ₃ I | (a) | < 0.3 | 15.3 | 327 | Dil F. M.B. | 1, 2 5 |

Notes

- (a) These quantities cannot be identified with normal Arrhenius factors.
 (b) See reference 4.

Ratios of Rate Constants (Potassium Atoms)

The table refers to ratios of rate constants for reactions of the following types:



where X is any halogen atom.

| <i>R₁X</i> | <i>R₂X</i> | Notes | <i>E₁ - E₂</i> | <i>A₂/A₁</i> | <i>k₂/k₁</i> | Temperature range | Radical source | Reference |
|----------------------------------|--------------------------------------|-------|--------------------------------------|------------------------------------|------------------------------------|-------------------|----------------|-----------|
| CH ₃ Cl | C ₂ H ₅ Cl | | $(kcal\ mole^{-1})$ 0.4 ± 0.09 | 0.83 ± 0.10 | | °C 217-333 | D.F. | 6 |
| CH ₃ Cl | CH ₃ COCl | | | | 360 | 285 | D.F. | 6 |
| C ₂ H ₅ Cl | (CH ₃) ₂ CHCl | | -0.13 ± 0.14 | 2.18 ± 0.25 | | 238-325 | D.F. | 7 |
| C ₂ H ₅ Cl | (CH ₃) ₃ CCl | | 2.27 ± 0.55 | 0.78 ± 0.4 | | 229-275 | D.F. | 7 |
| CH ₃ Br | C ₂ H ₅ Cl | | 4.4 | 0.52 | | 230-316 | D.F. | 6 |
| CH ₃ Br | CH ₃ COCl | | 1.4 | 3.5 | | 229-318 | D.F. | 6 |
| CH ₃ I | C ₂ H ₅ Cl | | | | 38 | 285 | D.F. | 6 |
| CH ₃ I | CH ₃ COCl | | 0 | 7.0 | | 238-322 | D.F. | 6 |

References

1. Schay, Z. Phys. Chem. **11B**, 291 (1930).
2. Polanyi, "Atomic Reactions," p. 46 (Williams and Norgate, London, 1932).
3. Taylor and Datz, J. Chem. Phys. **23**, 1711 (1955).
4. Krocsak and Schay, Z. Phys. Chem. **19B**, 344 (1932).
5. Herschbach, Kwei, and Norris, J. Chem. Phys. **34**, 1842 (1961).
6. Kerr, Lissi, and Trotman-Dickenson, J. Chem. Soc. 1673 (1964).
7. Emovon and Lissi, J. Chem. Soc. 3509 (1964).

Metathetical Reactions of Methyne Radicals

| Reaction | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|-------|---|--|-------------------------------------|---|-----------|
| $\text{CH} + \text{NH}_3 = \text{HCN} + \text{H}_2 + \text{H}$ | | (kcal mole^{-1}) $k > 6 \times 10^{10}$ | $(\text{cm}^3 \text{ mole}^{-1} \text{ sec}^{-1})$ | $^{\circ}\text{C}$ no indication | $\text{C}_2\text{H}_2/\text{NH}_3$ flames | 1 |

Reference

- (1) Safrany, Reeves, and Harteck, J. Am. Chem. Soc. **86**, 3160 (1964).

Ratios of Rate Constants (Methylene Radicals)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|------------|---------------------------|---------------------|---------------|-----------------------------------|--|-------------------------------------|
| (1) $\text{CH}_2 + \text{CH}_2\text{CO} = \text{C}_2\text{H}_4 + \text{CO}$ (2) $\text{CH}_2 + \text{CO} = \text{CH}_2\text{CO}$ | (a) (b) | (kcal mole^{-1}) | | 2.3 1.25 | $^{\circ}\text{C}$ 6-9 4-16 | $\text{CH}_2\text{CO}/^{13}\text{CO}$ P | 1 1 |
| (1) $\text{CH}_2 + \text{CH}_2\text{CO} = \text{C}_2\text{H}_4 + \text{CO}$ (2) $\text{CH}_2 + \text{C}_2\text{H}_4 = \text{C}_3\text{H}_6$ | | | 4.8 | | 28-81 | | $\text{CH}_2\text{CO}/\text{O}_2$ P |
| (1) $\text{CH}_2 + \text{CH}_2\text{CO} = \text{C}_2\text{H}_4 + \text{CO}$ (2) $\text{CH}_2 + \text{C}_2\text{H}_2 = \text{C}_3\text{H}_4$ | (c) | | | 0.9 ± 0.1 | no indication | $\text{CH}_2\text{CO}/\text{C}_2\text{H}_2$ F.P. | 3 |
| (1) $\text{CH}_2 + \text{CH}_2\text{CO} = \text{C}_2\text{H}_4 + \text{CO}$ (2) $\text{CH}_2 + \text{H}_2 = \text{CH}_3 + \text{H}$ | | -0.8 | | | -40-99 | $\text{CH}_2\text{CO}/\text{H}_2$ P | 4, 5, 7 |
| (1) $\text{CH}_2 + \text{CH}_2\text{CO} = \text{C}_2\text{H}_4 + \text{CO}$ (2) $\text{CH}_2 + \text{CD}_4 = \text{CH}_2\text{D}_2 + \text{CD}_2$ | | -0.9 | 0.15 | | 27-298 | $\text{CH}_2\text{CO}/\text{CD}_4$ P | 6, 7 |

Notes

(a) and (b) represent two methods of estimating k_1/k_2 : the discrepancy lies outside the experimental error.

(c) k_2 represents the sum of the rate constants for the reactions $\text{CH}_2 + \text{C}_2\text{H}_2 = \text{HC}:\text{CCH}_3$ and $\text{CH}_2 + \text{C}_2\text{H}_2 = \text{H}_2\text{C}:\text{C}:\text{CH}_2$.

References

1. Wilson and Kistiakowsky, J. Am. Chem. Soc. **80**, 2934 (1958).

2. Holroyd and Noyes, J. Am. Chem. Soc. **78**, 4831 (1956).
3. Terao, Sakai, and Shida, J. Am. Chem. Soc. **85**, 3919 (1963).
4. Gesser and Steacie, Can. J. Chem. **34**, 113 (1956).
5. Chanmugam and Burton, Can. J. Chem. **34**, 1021 (1956).
6. Chanmugam and Burton, J. Am. Chem. Soc. **78**, 509 (1956).
7. Bell and Kistiakowsky, J. Am. Chem. Soc. **84**, 3417 (1962).

Review

1. Frey, "Progress in Reaction Kinetics," Vol. 2, p. 133 (Pergamon Press, London, 1964).

Methyl Radical Reactions

Index

| | <i>Page</i> |
|--|-------------|
| (a) Hydrogen atom transfer | |
| Reaction with | |
| 1. Hydrogen..... | 41 |
| 2. Alkanes..... | 41 |
| 3. Cyclo-alkanes..... | 43 |
| 4. Alkenes..... | 43 |
| 5. Alkynes..... | 43 |
| 6. Aromatic hydrocarbons..... | 44 |
| 7. Halogenated aromatics..... | 44 |
| 8. Alcohols..... | 45 |
| 9. Thiols..... | 45 |
| 10. Amines..... | 45 |
| 11. Imines..... | 46 |
| 12. Amides, azines, etc..... | 46 |
| 13. Aldehydes..... | 47 |
| 14. Ketones..... | 48 |
| 15. Esters..... | 48 |
| 16. Acids and acid anhydrides..... | 49 |
| 17. Ethers and epoxides..... | 49 |
| 18. Peroxides..... | 49 |
| 19. Nitriles..... | 49 |
| 20. Metal alkyls..... | 49 |
| 21. Halogenated alkanes..... | 50 |
| 22. Hydrides of nitrogen, oxygen, and sulphur..... | 50 |
| 23. Hydrogen halides..... | 50 |
| (b) Fluorine atom transfer..... | 50 |
| (c) Chlorine atom transfer..... | 51 |
| (d) Bromine atom transfer..... | 51 |
| (e) Iodine atom transfer..... | 51 |
| (f) Oxygen atom transfer..... | 51 |
| (g) Group transfer reactions..... | 51 |
| (h) Ratios of rate constants | |
| 1. Reactions with halogens and hydrogen halides..... | 52 |
| 2. Miscellaneous reactions..... | 53 |

Metathetical Reactions of Methyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---------------------------------|---------|----------------------------|---|-------------------|---|-------------------------------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| H ₂ | (d) | 10.0 ± 0.5 | 11.5 | 130-290 | CH ₃ COCH ₃ P | 1, 2, 3 |
| | | 9.9 ± 0.5 | 11.7 | 25-250 | Hg(CH ₃) ₂ T | 4, 3 |
| | | 13 ± 2 | 13.4 | 50-250 | Cd(CH ₃) ₂ P | 5 |
| | (a) | 13.2 ± 1.0 | 12.5 | 136-318 | CH ₃ COCH ₃ P | 6 |
| | | 10.2 ± 0.5 | 11.52 | 99-207 | CH ₂ CO P | 7 |
| | (b) | 12.85 | 12.75 | | | 8 |
| | | | $k = 4.1 \times 10^8$ | | 507 | CH ₃ OCH ₃ T |
| | | 10.2 | 12.50 | 697-863 | H ₂ /O ₂ ignition | 10 |
| | (y) | 10.2 ± 0.2 | 11.7 | 130-290 | CD ₃ COCD ₃ P | 1, 2 |
| | | 11.1 | 12.0 | 150-300 | CD ₃ COCD ₃ P | 3 |
| | | 10.0 | 11.5 | | | |
| | HD | | 10.0 | 11.1 | 140-296 | CH ₃ COCH ₃ P |
| (y) | | 10.7 | 11.4 | 137-298 | CD ₃ COCD ₃ P | 3 |
| HD | | 11.5 | 11.5 | 135-296 | CH ₃ COCH ₃ P | 3 |
| | (y) | 10.7 | 11.2 | 137-299 | CD ₃ COCD ₃ P | 3 |
| D ₂ | (d) | 11.8 ± 0.1 | 11.8 | 130-290 | CH ₃ COCH ₃ P | 1, 2, 3 |
| | | 15.3 ± 1.0 | 14.8 | 258-451 | CH ₃ CHO P | 6 |
| | (a) | 14.3 ± 0.6 | 12.5 | 150-458 | CH ₃ COCH ₃ P | 6 |
| | | 12.7 ± 0.3 | 12.3 | 27-253 | Hg(CH ₃) ₂ P | 12 |
| | | | $k = 2.0 \times 10^8$ | 507 | CH ₃ OCH ₃ T | 9 |
| | | 12.1 ± 0.6 | 11.79 | 151-299 | CH ₃ COCH ₃ P | 13 |
| | | 11.9 | 11.4 | 140-425 | CH ₃ COCH ₃ P | 14 |
| | (y) | 10.9 ± 0.3 | 11.4 | 130-290 | CD ₃ COCD ₃ P | 1, 2 |
| | | | | | | |
| Alkanes | | | | | | |
| CH ₄ | | 14.9 | 12.00 | 200-350 | CH ₃ COCH ₃ P | 16, 17 |
| | (y) | | $k = 1.7 \times 10^5$ | 182 | CD ₃ COCD ₃ P | 18, 19 |
| | (y) | 14.3 | 11.5 | 350-525 | CD ₃ COCD ₃ P, T | 20 |
| | (y) | | $k = 3.6 \times 10^6$ | 320 | CD ₃ COCD ₃ P | 21 |
| | (y) | 14.1 | 11.8 | 200-350 | | 17 |
| CH ₃ D | (y) (e) | 12.8 | 10.42 | 27-327 | | 22 |
| CD ₃ H | (e) | 13.8 | 11.21 | 27-327 | | 22 |
| CD ₄ | | 12.93 ± 0.65 | 11.26 | 153-428 | CH ₃ COCH ₃ P | 23, 17 |
| | (y) | 18.4 | 12.95 | 200-350 | CD ₃ COCD ₃ P | 17 |
| C ₂ H ₆ | | 12.0 | 10.6 | 770-890 | C ₂ H ₆ T | 24 |
| | | | $k = 1.4 \times 10^8$ | 630 | C ₂ H ₆ T | 25 |
| | (y) (f) | 9.9 ± 1.1 | | 500-560 | CD ₃ CDO T | 29, 51 |
| | | 10.4 ± 0.4 | 11.3 | 116-294 | CD ₃ COCD ₃ P | 19, 26 |
| | (y) | 11.8 | 12.21 | 162-341 | CD ₃ COCD ₃ P | 21 |
| | (y) | 12.1 | 12.3 | 260-490 | CD ₃ COCD ₃ P | 27 |
| | (y) | 11.5 ± 0.2 | 11.9 | 246-524 | CD ₃ COCD ₃ P, T | 28 |
| | | | | | | |
| CH ₃ CD ₃ | (y) | 12.2 | 12.0 | 260-490 | CD ₃ COCD ₃ P | 27, 39 |
| CH ₃ CD ₃ | (y) | 14.1 | 12.1 | 260-490 | CD ₃ COCD ₃ P | 27, 39 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|---------|----------------------------|---|-----------------------|--|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| C ₂ D ₆ | (y) | 14.8 ± 0.3 | 12.3 | 328–507 | CH ₃ COCH ₃ P, T | 28 |
| | | 13.6 | 12.3 | 260–490 | CD ₃ COCD ₃ P | 27 |
| C ₃ H ₈ | (g) | 10.3 | $k = 3.3 \times 10^9$ | 580 | C ₃ H ₈ T | 29 |
| | | | | 300–460 | CD ₃ COCD ₃ P | 30 |
| CH ₃ CD ₂ CH ₃ | (y) | 11.6 | 12.08 | 300–450 | CD ₃ COCD ₃ P | 31, 39 |
| CH ₃ CD ₂ CH ₃ | (y) | 11.6 | 11.85 | 300–450 | CD ₃ COCD ₃ P | 31, 39 |
| <i>n</i> -C ₄ H ₁₀ | (h) | 8.3 ± 0.2 | 11.0 | 122–198 | CH ₃ COCH ₃ P | 19, 26 |
| | | 8.2 ± 0.5 | 11.3 | 130–220 | Hg(CH ₃) ₂ P | 32 |
| | | 9.5 ± 0.5 | 11.5 | | Hg(CH ₃) ₂ P | 32 |
| | (h) | 8.6 ± 0.3 | 11.2 | 92–223 | Hg(CH ₃) ₂ P | 33 |
| | | 9.1 ± 0.3 | 11.4 | 182 | Hg(CH ₃) ₂ P | 34 |
| | (y) (f) | 9.6 | 11.92 | 79–162 | (CH ₃) ₂ N ₂ P | 35 |
| | | 10.3 ± 0.2 | | 250–449 | CD ₃ COCD ₃ P | 36 |
| | | | 500–560 | CD ₃ CDO T | 29 | |
| CH ₃ CD ₂ CD ₂ CH ₃ | (y) | 11.7 | 12.09 | 356–450 | CD ₃ COCD ₃ P | 36 |
| CH ₃ CD ₂ CD ₂ CH ₃ | (y) | 11.7 | 12.12 | 356–450 | CD ₃ COCD ₃ P | 36 |
| <i>iso</i> -C ₄ H ₁₀ | (y) | 7.6 ± 0.2 | 11.0 | 76–194 | CH ₃ COCH ₃ P | 19, 26 |
| | | 7.4 ± 0.3 | 10.8 | 93–220 | Hg(CH ₃) ₂ P | 33 |
| | | 6.6 ± 0.3 | 10.3 | 25–169 | (CH ₃) ₂ N ₂ P | 35 |
| | | 7.7 | 10.3 | 111–146 | D.T.B.P. T | 37 |
| | (y) (f) | 9.0 ± 0.6 | | 198 | CD ₃ COCD ₃ P | 38 |
| | | | | 500–560 | CD ₃ CDO T | 29 |
| (CH ₃) ₃ CH | (y) | 8.2 | 11.47 | 300–460 | CD ₃ COCD ₃ P | 30 |
| (CH ₃) ₃ CD | (y) | 11.7 | 12.26 | 300–460 | CD ₃ COCD ₃ P | 30, 39 |
| (CH ₃) ₃ CD | (y) | 9.8 | 11.57 | 300–460 | CD ₃ COCD ₃ P | 30, 39 |
| <i>n</i> -C ₅ H ₁₂ | (y) | 8.1 ± 0.2 | 11.0 | 89–178 | CH ₃ COCH ₃ P | 19, 26 |
| | | | $k = 1.0 \times 10^9$ | 540 | <i>n</i> -C ₅ H ₁₂ T | 40 |
| | | 9.9 ± 2.7 | | 540–580 | <i>n</i> -C ₅ H ₁₂ T | 29 |
| <i>iso</i> -C ₅ H ₁₂ | (y) | | $k = 0.9 \times 10^9$ | 540 | <i>iso</i> -C ₅ H ₁₂ T | 40 |
| <i>neo</i> -C ₅ H ₁₂ | (y) (f) | 10.0 ± 0.3 | 11.3 | 138–292 | CH ₃ COCH ₃ P | 19, 26 |
| | | 10.4 ± 0.3 | 11.3 | 131–251 | Hg(CH ₃) ₂ P | 33 |
| | (y) (f) | 10.8 ± 0.2 | | 540 | <i>neo</i> -C ₅ H ₁₂ T | 40 |
| | | | $k = 0.2 \times 10^9$ | 500–560 | CD ₃ CDO T | 29 |
| <i>n</i> -C ₆ H ₁₄ | (y) | 8.1 ± 0.2 | 11.1 | 92–184 | CH ₃ COCH ₃ P | 19, 26 |
| (CH ₃) ₂ CHCH(CH ₃) ₂ | (y) (f) | 6.9 ± 0.2 | 10.8 | 27–190 | CH ₃ COCH ₃ P | 19, 26 |
| | | 6.8 ± 0.2 | 10.7 | 28–220 | Hg(CH ₃) ₂ P | 33 |
| | (y) | 7.3 ± 0.8 | | 500–560 | CD ₃ CDO T | 29 |
| | | 7.8 ± 0.4 | 11.3 | 166–293 | CD ₃ COCD ₃ P | 26 |
| (C ₂ H ₅) ₂ CH | (y) | 6.8 ± 0.3 | 10.5 | 76–238 | Hg(CH ₃) ₂ P | 41 |
| <i>n</i> -C ₈ H ₁₈ | (y) | | $k = 1.6 \times 10^9$ | 500 | <i>n</i> -C ₈ H ₁₈ T | 40 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range, | Radical source | References |
|---|---------|----------------------------|---|--------------------|--|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| (CH ₃) ₃ CCH ₂ CH(CH ₃) ₂ | | | $k = 0.7 \times 10^9$ | 500 | (CH ₃) ₃ CCH ₂ CH(CH ₃) ₂ T | 40 |
| (CH ₃) ₂ CHCH(CH ₃) CH(CH ₃) ₂ | (y) | 7.9 ± 0.4 | 11.3 | 500 141–332 | (CH ₃) ₂ CHCH(CH ₃) CH(CH ₃) ₂ T CD ₃ COCD ₃ P | 40 26 |
| (CH ₃) ₃ CC(CH ₃) ₃ | (y) | 9.5 ± 0.4 | 11.3 | 162–322 | CD ₃ COCD ₃ P | 26 |
| Cyclo-alkanes | | | | | | |
| cyclo-C ₃ H ₆ | (y) | 10.2 ± 1.0 | 11.1 | 100–250 | Hg(CH ₃) ₂ P | 42 |
| | (y) | 10.3 ± 0.4 | 11.0 | 139–292 | CD ₃ COCD ₃ P | 43 |
| | (y) | 12.9 | 12.18 | 248–404 | CD ₃ COCD ₃ P | 44 |
| cyclo-C ₄ H ₈ | (y) | 9.3 ± 0.4 | 11.4 | 154–307 | CD ₃ COCD ₃ P | 43 |
| | | 10.1 | 12.18 | 262–402 | CD ₃ COCD ₃ P | 45 |
| cyclo-C ₅ H ₁₀ | (i) | . | $k = 5.7 \times 10^7$ | 303 | Hg(CH ₃) ₂ T | 46 |
| | (y) | 8.3 ± 0.2 | 11.4 | 66–296 | CD ₃ COCD ₃ P | 43 |
| | (y) | 9.1 | 12.24 | 250–402 | CD ₃ COCD ₃ P | 44 |
| | (y) | 9.0 | 12.10 | 160–500 | CD ₃ COCD ₃ P | 47 |
| cyclo-C ₆ H ₁₂ | | 8.3 ± 0.2 | 11.3 | 65–189 | CH ₃ COCH ₃ P | 43 |
| | | 8.3 ± 0.3 | 11.2 | 81–220 | Hg(CH ₃) ₂ P | 33 |
| | (y) | 9.5 | 12.47 | 254–481 | CD ₃ COCD ₃ P | 48 |
| cyclo-C ₇ H ₁₄ | (y) | 8.9 | 12.49 | | | 48 |
| Alkenes | | | | | | |
| CH ₂ :CH ₂ | (y) | 10.0 ± 0.4 | 11.3 | 188–340 | CD ₃ COCD ₃ P | 49 |
| | (y) (f) | 7.3 ± 1.0 | | 500–560 | CD ₃ CDO T | 29 |
| CH ₃ CH:CH ₂ | | 8.2 | 11.04 | 107–168 | D.T.B.P | 50 |
| | (y) | 7.7 ± 0.4 | 10.8 | 163–304 | CD ₃ COCD ₃ P | 49 |
| | (y) (f) | 8.0 ± 0.3 | | 500–560 | CD ₃ CDO T | 29, 51 |
| CH ₃ CH:CHCH ₃ | (y) | 7.7 ± 0.4 | 11.1 | 188–342 | CD ₃ COCD ₃ P | 49 |
| CH ₂ :CHCH ₂ CH ₃ | (y) | 7.6 ± 0.4 | 11.2 | 189–340 | CD ₃ COCD ₃ P | 49 |
| (CH ₃) ₂ C:CH ₂ | (y) | 7.3 ± 0.4 | 10.9 | 168–304 | CD ₃ COCD ₃ P | 49 |
| CH ₃ CH ₂ CH ₂ CH:CH ₂ | (y) | 7.6 ± 0.4 | 11.2 | 188–346 | CD ₃ COCD ₃ P | 49 |
| (CH ₃) ₂ CHCH:CH ₂ | (y) | 7.4 ± 0.4 | 11.3 | 189–346 | CD ₃ COCD ₃ P | 49 |
| (CH ₃) ₂ C:C(CH ₃) ₂ | (y) | 7.8 ± 0.4 | 11.6 | 188–341 | CD ₃ COCD ₃ P | 49 |
| Alkynes | | | | | | |
| HC:CH | (y) | 14.0 | | 200–500 | CD ₃ COCD ₃ P | 52 |
| DC:CD | | 18.4 | | 200–500 | CH ₃ COCH ₃ P | 52 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|---------|----------------------------|---|-------------------------------------|---|------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CH ₃ C:CCH ₃ | (y) | 8.6 ± 0.4 | 11.6 | 213–346 | CD ₃ COCD ₃ P | 43 |
| CH:CCH ₂ CH ₃ | (y) | 9.1 ± 0.4 | 11.9 | 183–347 | CD ₃ COCD ₃ P | 43 |
| Aromatic hydrocarbons | | | | | | |
| C ₆ H ₆ | (y) | 9.2 ± 0.4 | $k = 4.48 \times 10^6$ | 209 | D.T.B.P. T | 53 |
| | | | 10.4 | 183–327 | CD ₃ COCD ₃ P | 43 |
| C ₆ H ₅ CH ₃ | (h) | 7 ± 2 11.0 ± 2 | 10.0 | 103–249 | Hg(CH ₃) ₂ P | 54 |
| | | | | 130–230 | D.T.B.P. P | 55 |
| | | | | 149–250 | Hg(CH ₃) ₂ P | 33 |
| | (j) | 13.03 ± 0.27 | 11.2 | 346–701 | | 56, 57, 58 |
| | (z) | 7.4 ± 0.3 | 12.92 ± 0.08 | 100–253 | CH ₃ COCH ₃ P | 59 |
| | | 9.2 ± 0.3 | 10.47 | 159–270 | D.T.B.P. T | 53 |
| | (y) | 8.3 ± 0.3 | $k = (4.0 \pm 0.8) \times 10^5$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| | | 11.0 | 120–334 | CD ₃ COCD ₃ P | 43 | |
| C ₆ H ₅ CD ₃ | | 10.2 | $k = (2.3 \pm 0.4) \times 10^4$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| | | | 10.7 | 100–300 | | 72 |
| C ₆ H ₅ CD ₃ | | 11.3 | $k = (2.6 \pm 0.2) \times 10^4$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| | | | 11.8 | 102–284 | CH ₃ COCH ₃ P | 72 |
| C ₆ D ₅ CH ₃ | | 9.6 | $k = (3.8 \pm 0.8) \times 10^5$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| | | | 11.6 | 106–284 | CH ₃ COCH ₃ P | 72 |
| C ₆ D ₅ CH ₃ | | | $k = (1.0 \pm 0.2) \times 10^4$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| | | | $k = 5.1 \times 10^3$ | 182 | CH ₃ COCH ₃ P | 72 |
| C ₆ D ₅ CD ₃ | | | $k = (3.7 \pm 0.2) \times 10^4$ | 60 | CH ₃ N ₂ CH ₃ P | 60 |
| C ₆ H ₅ C ₂ H ₅ | (z) | 7.0 ± 0.3 | 10.82 | 85–183 | CH ₃ COCH ₃ P | 59 |
| C ₆ H ₅ CH(CH ₃) ₂ | (z) | 6.4 ± 0.5 | 10.76 | 123–249 | CH ₃ COCH ₃ P | 59 |
| <i>o</i> -C ₆ H ₄ (CH ₃) ₂ | (z) | 7.8 ± 0.3 | 11.13 | 101–205 | CH ₃ COCH ₃ P | 61 |
| <i>m</i> -C ₆ H ₄ (CH ₃) ₂ | (z) | 8.5 ± 0.3 | 11.45 | 100–197 | CH ₃ COCH ₃ P | 61 |
| <i>p</i> -C ₆ H ₄ (CH ₃) ₂ | (z) | 7.4 ± 0.2 | 10.82 | 99–197 | CH ₃ COCH ₃ P | 61 |
| | (k) | | $k = 3.1 \times 10^7$ | 484 | <i>p</i> -C ₆ H ₄ (CH ₃) ₂ T | 62 |
| C ₆ H ₄ (CH ₃) ₂ | (k) (l) | | $k = 6.9 \times 10^9$ | 700 | C ₆ H ₄ (CH ₃) ₂ T | 63 |
| Halogenated aromatics | | | | | | |
| <i>o</i> -CH ₃ C ₆ H ₄ F | (z) | 6.0 ± 0.4 | 9.76 | 330–460 | CH ₃ COCH ₃ P | 64 |
| <i>m</i> -CH ₃ C ₆ H ₄ F | (z) | 7.1 ± 0.2 | 10.32 | 330–460 | CH ₃ COCH ₃ P | 64 |
| <i>p</i> -CH ₃ C ₆ H ₄ F | (z) | 5.7 ± 0.4 | 9.60 | 330–460 | CH ₃ COCH ₃ P | 64 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|---------|---------------------------------|--|-------------------|--|------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| Alcohols | | | | | | |
| CH ₃ OH | | 8.2 ± 0.2 | 10.7 | 103–219 | CH ₃ COCH ₃ | P 43 |
| | | 8.2 ± 0.5 | 10.6 | 100–250 | Hg(CH ₃) ₂ | P 42 |
| | | 8.7 ± 0.2 | 10.76 ± 0.07 | 133–199 | CH ₃ COCH ₃ | P 65 |
| | (y) | 8.4 ± 0.1 | 10.68 ± 0.05 | 125–250 | CD ₃ COCD ₃ | P 73 |
| CH ₃ OH | (m) | 10.4 | 11.38 | 133–199 | CH ₃ COCH ₃ | P 65 |
| | (y) (m) | 8.1 | 10.38 | 125–250 | CD ₃ COCD ₃ | P 73 |
| CD ₃ OH | | 6.4 ± 0.7 | 9.25 ± 0.3 | 133–199 | CH ₃ COCH ₃ | P 65 |
| | (y) | 9.0 ± 0.1 | 10.46 ± 0.03 | 125–245 | CD ₃ COCD ₃ | P 73 |
| CD ₃ OH | | 11.7 ± 0.6 | 11.25 ± 0.28 | 133–199 | CH ₃ COCH ₃ | P 65 |
| | (y) | 9.3 ± 0.1 | 10.18 ± 0.06 | 125–245 | CD ₃ COCD ₃ | P 73 |
| C ₂ H ₅ OH | (y) | 8.7 ± 0.4 | 11.5 | 189–341 | CD ₃ COCD ₃ | P 43 |
| (CH ₃) ₂ CH(OH) | (y) | 7.3 ± 0.4 | 11.0 | 214–347 | CD ₃ COCD ₃ | P 43 |
| Thiols | | | | | | |
| CH ₃ SH | | | $k = 1.8 \times 10^7$ | 30 | CH ₃ COCH ₃ | P 66 |
| CD ₃ SH | | 4.1 ± 0.2 | 11.03 ± 0.15 | 130–200 | CH ₃ COCH ₃ | P 74 |
| CD ₃ SH | | 8.3 ± 0.1 | 10.88 ± 0.05 | 130–200 | CH ₃ COCH ₃ | P 74 |
| C ₂ H ₅ SH | | | $k = 3.5 \times 10^7$ | 30 | CH ₃ COCH ₃ | P 66 |
| (CH ₃) ₂ CHSH | | | $k = 4.1 \times 10^7$ | 30 | CH ₃ COCH ₃ | P 66 |
| (CH ₃) ₃ CSH | | | $k = 5.9 \times 10^7$ | 30 | CH ₃ COCH ₃ | P 66 |
| Amines (primary) | | | | | | |
| CH ₃ NH ₂ | | 7.6 | 10.9 | 125–157 | D.T.B.P. | T 67 |
| | | 7.2 ± 0.2 | 10.59 ± 0.12 | | CH ₃ N ₂ CH ₃ | P 68 |
| | (y) | 8.4 ± 0.4 | 11.3 | 183–340 | CD ₃ COCD ₃ | P 43 |
| CH ₃ NH ₂ | (m) | 8.7 ± 0.7 | 10.99 ± 0.37 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| CH ₃ NH ₂ | (m) | 5.7 ± 0.40 | 9.55 ± 0.22 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| CH ₃ ND ₂ | | 9.00 ± 0.20 | 11.15 ± 0.12 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| CH ₃ ND ₂ | | 7.00 ± 0.30 | 9.61 ± 0.16 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| CD ₃ NH ₂ | | 6.00 ± 0.50 | 9.77 ± 0.28 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| CD ₃ NH ₂ | | 10.10 ± 0.30 | 10.86 ± 0.16 | 120–175 | CH ₃ N ₂ CH ₃ | P 69, 68 |
| C ₂ H ₅ NH ₂ | | 7.1 | 11.2 | 125–157 | D.T.B.P. | T 67 |
| | | 7.3 ± 0.3 | 10.89 ± 0.18 | 110–180 | CH ₃ N ₂ CH ₃ | P 75 |
| CH ₃ CH ₂ NH ₂ | | | $k = 6.3 \times 10^5$ | 150 | CH ₃ N ₂ CH ₃ | P 75 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> (kcal mole ⁻¹) | log ₁₀ <i>A</i> (cm ³ mole ⁻¹ sec ⁻¹) | Temperature range °C | Radical source | References |
|---|-------|--|---|-------------------------|--|------------|
| CH ₃ CH ₂ NH ₂ | | 8.1 | 11.2 | 110–180 | CH ₃ N ₂ CH ₃ P | 75 |
| CH ₃ CH ₂ NH ₂ | | 6.5 | 9.9 | 110–180 | CH ₃ N ₂ CH ₃ P | 75 |
| CH ₃ CH ₂ ND ₂ | | 7.6 ± 0.4 | 10.04 ± 0.21 | 110–180 | CH ₃ N ₂ CH ₃ P | 75 |
| CD ₃ CH ₂ ND ₂ | | | <i>k</i> = 1.3 × 10 ⁵ | 150 | CH ₃ N ₂ CH ₃ P | 75 |
| Amines (secondary) | | | | | | |
| (CH ₃) ₂ NH | | 7.2 | 11.7 | 125–157 | D.T.B.P. P | 67 |
| | | 7.00 ± 0.30 | 11.20 ± 0.13 | 120–180 | CH ₃ N ₂ CH ₃ P | 70 |
| | (y) | 7.2 ± 0.4 | 11.3 | 184–341 | CD ₃ COCD ₃ P | 43 |
| (CH ₃) ₂ NH | | 6.40 ± 0.30 | 10.81 ± 0.21 | 120–175 | CH ₃ N ₂ CH ₃ P | 70 |
| (CH ₃) ₂ ND | | 8.70 ± 0.60 | 11.46 ± 0.28 | 120–175 | CH ₃ N ₂ CH ₃ P | 70 |
| (CH ₃) ₂ ND | | 7.80 ± 0.40 | 10.65 ± 0.22 | 120–175 | CH ₃ N ₂ CH ₃ P | 70 |
| (C ₂ H ₅) ₂ NH | | 7.2 | 11.8 | 125–157 | D.T.B.P. T | 67 |
| | (n) | 5.7 ± 1.0 | 10.7 | 123–260 | CH ₃ COCH ₃ P | 71 |
| ((CH ₃) ₂ CH) ₂ NH | | 7.8 | 12.0 | 125–157 | D.T.B.P. T | 67 |
| Amines (tertiary) | | | | | | |
| (CH ₃) ₃ N | | 8.0 | 11.4 | 132–269 | CH ₃ COCH ₃ P | 71 |
| | | 8.9 ± 0.1 | 11.9 ± 0.1 | 123–232 | CH ₃ COCH ₃ P | 76 |
| | (y) | 8.8 ± 0.4 | 11.8 | 193–302 | CD ₃ COCD ₃ P | 43 |
| (C ₂ H ₅) ₃ N | (n) | 5.3 ± 1.0 | 10.4 | 123–260 | CH ₃ COCH ₃ P | 71 |
| Imines | | | | | | |
| (CH ₂) ₂ NH | | 4.8 ± 0.3 | 10.7 | 125–157 | D.T.B.P. T | 77 |
| | (o) | 5.28 | 10.76 | 100–216 | CH ₃ COCH ₃ P | 78 |
| | | 4.77 ± 0.44 | 10.29 ± 0.23 | 110–175 | CH ₃ N ₂ CH ₃ P | 79, 80 |
| | (y) | 5.4 | 10.94 | 120–260 | CD ₃ COCD ₃ P | 78 |
| (CH ₂) ₂ NH | (m) | 10.10 ± 2.80 | 11.44 ± 1.48 | 110–175 | CH ₃ N ₂ CH ₃ P | 79, 80 |
| (CH ₂) ₂ NH | | 4.57 ± 0.10 | 10.17 ± 0.05 | 110–175 | CH ₃ N ₂ CH ₃ P | 79, 80 |
| (CH ₂) ₂ ND | | 6.34 ± 0.21 | 10.17 ± 0.13 | 110–175 | CH ₃ N ₂ CH ₃ P | 79, 80 |
| (CH ₂) ₂ NC(CH ₃) ₃ | | 6.6 | 9.6 | 125–157 | D.T.B.P. T | 67 |
| CH ₃ CH:NC(CH ₃) ₃ | | 7.8 | 11.1 | 125–157 | D.T.B.P. T | 67 |
| Amides, azines, etc. | | | | | | |
| HCONH ₂ | | 6.6 | 10.5 | 170–247 | CH ₃ COCH ₃ P | 81 |
| HCONHCH ₃ | | 7.6 | 10.9 | 161–287 | CH ₃ COCH ₃ P | 81 |
| HCON(CH ₃) ₂ | | 8.3 | 11.4 | 120–298 | CH ₃ COCH ₃ P | 81 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | References |
|---|---------|-----------------------------------|--|-------------------|--|------------|
| | | (<i>kcal mole⁻¹</i>) | (<i>cm³ mole⁻¹ sec⁻¹</i>) | °C | | |
| CH ₃ CONH ₂ | (q) | 9.2 ± 0.3 | 10.8 | 144–224 | CH ₃ CONH ₂ P | 82 |
| CH ₃ CON(CH ₃) ₂ | | 8.3 ± 0.2 | 11.3 ± 0.1 | 106–232 | CH ₃ COCH ₃ P | 76 |
| (CH ₃) ₂ NCON(CH ₃) ₂ | | 7.3 ± 0.2 | 11.0 ± 0.1 | 108–235 | CH ₃ COCH ₃ P | 76 |
| CH ₃ ONH ₂ | | 4.53 ± 0.25 | 10.70 ± 0.15 | 70–190 | CH ₃ N ₂ CH ₃ P | 83 |
| CH ₃ OND ₂ | | 5.88 ± 0.22 | 10.55 ± 0.12 | 70–190 | CH ₃ N ₂ CH ₃ P | 83 |
| (CH ₃) ₂ N.NH ₂ | | 5.80 ± 0.20 | 11.34 ± 0.12 | 110–180 | CH ₃ N ₂ CH ₃ P | 68 |
| (CH ₃) ₂ N.NH ₂ | (r) | 8.5 | 11.6 | 110–180 | CH ₃ N ₂ CH ₃ P | 68 |
| CH ₃ CH:NN:CHCH ₃ | | 6.1 | 10.5 | 125–157 | D.T.B.P. T | 67 |
| CH ₃ N ₂ CH ₃ | | 7.6 ± 0.3 | 11.1 | 60–182 | CH ₃ N ₂ CH ₃ P | 84 |
| | | 7.3 | 10.49 | 25–150 | CH ₃ N ₂ CH ₃ P | 85 |
| | | 6.86 ± 0.18 | 10.37 | –47–50 | CH ₃ N ₂ CH ₃ P | 86 |
| | | 8.4 ± 0.3 | 11.40 | 80–180 | CH ₃ N ₂ CH ₃ P | 87 |
| | (s) | 7.83 ± 0.08 | 10.97 ± 0.04 | 70–190 | CH ₃ N ₂ CH ₃ P | 83 |
| | | 8.7 | 11.47 | 50–180 | CH ₃ N ₂ CH ₃ P | 89 |
| CH ₃ N:N(O)CH ₃ | | 6 ± 2 | | 27–121 | CH ₃ N:N(O)CH ₃ P | 90 |
| Aldehydes | | | | | | |
| HCHO | | | $k = 4.7 \times 10^7$ | 120 | CH ₃ COCH ₃ /O ₂ P | 91 |
| | | 6.2 ± 0.3 | 11.06 | 80–180 | CH ₃ N ₂ CH ₃ P | 87 |
| | | 6.6 | 11.25 | 111–146 | D.T.B.P. T | 37 |
| DCDO | | 7.9 ± 0.3 | 11.15 | 80–180 | CH ₃ N ₂ CH ₃ P | 87 |
| CH ₃ CHO | | 7.9 ± 0.3 | 12.15 | 124–156 | D.T.B.P. T | 92 |
| | | 6.8 | 11.5 | 91–165 | CH ₃ N ₂ CH ₃ P | 93 |
| | | 8.5 | 12.4 | 133–291 | CH ₃ CHO P | 94 |
| | | 8.7 | | 109–345 | CH ₃ CHO P | 95 |
| | | 7.6 ± 0.2 | 11.9 ± 0.1 | 119–175 | D.T.B.P. T | 96 |
| | | 6.8 | 11.50 | 25–250 | CH ₃ N ₂ CH ₃ P | 97 |
| | (y) (f) | 6.5 ± 0.3 | | 500–560 | CD ₃ CDO T | 29 |
| CH ₃ CDO | | 7.9 | 11.8 | 27–158 | CH ₃ N ₂ CH ₃ P | 93 |
| C ₂ H ₅ CHO | | 7.5 | 11.9 | 122–156 | D.T.B.P. T | 99, 92 |
| CH ₂ :CHCH ₂ CHO | | 10.9 | 13.3 | 119–175 | D.T.B.P. T | 96 |
| <i>n</i> -C ₃ H ₇ CHO | | 7.3 ± 0.3 | 11.8 ± 0.2 | 119–175 | D.T.B.P. T | 96 |
| <i>n</i> -C ₃ F ₇ CHO | | 5.55 ± 0.20 | 11.20 | 27–306 | CH ₃ COCH ₃ / C ₃ F ₇ CHO P | 98 |
| iso-C ₃ H ₇ CHO | | 8.7 ± 0.3 | 12.6 ± 0.2 | 119–175 | D.T.B.P. T | 96 |
| <i>n</i> -C ₄ H ₉ CHO | | 8.0 ± 0.3 | 12.1 ± 0.2 | 119–175 | D.T.B.P. T | 96 |
| iso-C ₄ H ₉ CHO | | 8.4 ± 0.3 | 12.3 ± 0.2 | 119–175 | D.T.B.P. T | 96 |

Metathetical Reactions of Methyl Radicals—Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References | |
|---|-------------------------------------|----------------------------|---|-------------------|---|-------------------------------------|----|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | | |
| sec-C ₄ H ₉ CHO | | 10.4 ± 0.3 | 13.1 ± 0.3 | 119–175 | D.T.B.P. T | 96 | |
| <i>t</i> -C ₄ H ₉ CHO | | 10.2 ± 0.3 | 13.0 ± 0.3 | 119–175 | D.T.B.P. T | 96 | |
| Ketones | | | | | | | |
| CH ₃ COCH ₃ | (t) | 9.7 ± 0.1 | 11.6 | 121–300 | CH ₃ COCH ₃ P | 100 | |
| | (h) | 9.6 ± 0.4 | 11.5 | 100–250 | CH ₃ COCH ₃ P | 101 | |
| | | 9.7 ± 0.2 | 11.8 | 125–220 | CH ₃ COCH ₃ and Hg(CH ₃) ₂ P | 102 | |
| | | 9.8 | 11.59 | 271–439 | CH ₃ COCH ₃ P | 103 | |
| | | 9.5 ± 1.5 | 11.5 | 127–175 | D.T.B.P. T | 104 | |
| | | 9.5 ± 0.3 | 11.8 | 130–155 | D.T.B.P. T | 105 | |
| | | 9.8 ± 0.4 | 11.60 | 27–412 | CH ₃ COCH ₃ P | 13 | |
| | | 9.56 | 11.43 | 132–292 | CH ₃ COCH ₃ P | 2, 15 | |
| | CD ₃ COCD ₂ H | (y) (u) | 9.95 ± 0.15 | 10.74 | 120–250 | CD ₃ COCD ₃ P | 73 |
| | | (y) (u) | 9.48 | 9.87 | 125–200 | CD ₃ COCD ₃ P | 74 |
| CD ₃ COCD ₃ | (y) | 10.3 ± 0.2 | 11.6 | 138–292 | CD ₃ COCD ₃ P | 100 | |
| | (y) | 10.6 ± 0.3 | 11.8 | 130–290 | CD ₃ COCD ₃ P | 1 | |
| | (y) | 11.6 ± 0.3 | 11.8 | 135–290 | CD ₃ COCD ₃ P | 3 | |
| | (y) | 11.6 | 12.07 | 150–250 | CD ₃ COCD ₃ P | 30 | |
| | (y) | 11.44 ± 0.05 | 11.66 ± 0.03 | 120–250 | CD ₃ COCD ₃ P | 73 | |
| | (y) | 11.29 ± 0.03 | 11.57 ± 0.01 | 125–200 | CD ₃ COCD ₃ P | 74 | |
| | (y) | 10.9 ± 1.0 | 11.5 | 130–200 | CD ₃ COCD ₃ P | 110 | |
| CF ₃ COCH ₃ | | 8.9 | 12.0 | 25–113 | CF ₃ COCH ₃ P | 109 | |
| CH ₃ COC ₂ H ₅ | | 7.4 | 10.6 | 79–190 | CH ₃ COC ₂ H ₅ P | 166 | |
| C ₂ H ₅ COC ₂ H ₅ | | 8.0 ± 0.2 | 11.8 | 130–155 | D.T.B.P. T | 105 | |
| | | 7.0 ± 0.1 | 11.2 | 26–134 | CH ₃ N ₂ CH ₃ P | 85 | |
| | (y) | | $k = 2.9 \times 10^7$ | 141 | CD ₃ COCD ₃ P | 111 | |
| cyclo-C ₃ H ₅ COCH ₃ | | 9.6 ± 1 | | 60–170 | cyclo-C ₃ H ₅ COCH ₃ P | 112 | |
| C ₆ H ₅ COCH ₃ | | 7.4 | 10.7 | 273–407 | C ₆ H ₅ COCH ₃ P | 113 | |
| CH ₃ COCOCH ₃ | | 7.1 ± 0.2 | | 28–200 | CH ₃ COCOCH ₃ P | 114 | |
| | | 8.5 | 11.3 | 140–198 | CH ₃ N ₂ CH ₃ P | 116 | |
| | | 7.7 | | 28–200 | CH ₃ COCOCH ₃ P | 115 | |
| Esters | | | | | | | |
| HCOOCH ₃ | | 9.0 | 10.9 | 77–230 | CH ₃ COCH ₃ P | 117 | |
| | | 9.8 | 11.3 | | | 117 | |
| HCOOC ₂ H ₅ | | 8.2 | 10.5 | 77–230 | CH ₃ COCH ₃ P | 117 | |
| HCOOCH ₂ CH ₂ CH ₃ | | 7.3 | 10.1 | 74–178 | CH ₃ COCH ₃ P | 118 | |
| HCOOCH(CH ₃) ₂ | | 8.9 | 10.9 | 94–181 | CH ₃ COCH ₃ P | 118 | |
| HCOOCH ₂ CH ₂ CH ₂ CH ₃ | | 8.2 | 10.6 | 75–186 | CH ₃ COCH ₃ P | 119 | |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|---|---------|---------------------------------|--|-------------------|--|------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| CH ₃ COOCH ₃ | | 10 ± 0.5 | 11.2 | 63–216 | CH ₃ COOCH ₃ P | 120 |
| CH ₃ COOCD ₃ | | 10 ± 0.5 | 11.4 | 151–340 | CH ₃ COCH ₃ P | 121 |
| | | 10 ± 0.5 | 11.4 | 145–350 | CH ₃ COOCD ₃ P | 122 |
| CH ₃ COOCD ₃ | | 14 ± 1 | 11.8 | 151–274 | CH ₃ COCH ₃ P | 121 |
| C ₂ H ₅ COOC ₂ H ₅ | | 8.2 | 11.3 | 72–344 | C ₂ H ₅ COOC ₂ H ₅ P | 123 |
| CH ₃ OCOOCH ₃ | | 8.9 ± 0.7 | 10.29 ± 0.34 | 122–253 | CH ₃ COCH ₃ P | 106 |
| Acids and acid anhydrides | | | | | | |
| CH ₃ COOD | | 10.2 | 11.1 | 105–285 | CH ₃ COOD P | 124 |
| (CH ₃ CO) ₂ O | | 9.6 | 11.3 | 107–196 | (CH ₃ CO) ₂ O P | 125 |
| Ethers and epoxides | | | | | | |
| CH ₃ OCH ₃ | | 9.5 ± 0.2 | 11.5 | 108–198 | CH ₃ COCH ₃ P | 43 |
| | | 8.4 ± 1.5 | 11.0 | 100–250 | Hg(CH ₃) ₂ P | 42 |
| | | 10 ± 2 | 11.8 | 199–292 | CH ₃ OCH ₃ – H | 126 |
| | (y) (f) | 10.8 ± 1.0 | | 500–560 | CD ₃ CDO T | 29 |
| (C ₂ H ₅) ₂ O | | 9.75 ± 0.5 | 12.14 | 145–179 | D.T.B.P. T | 127 |
| [(CH ₃) ₂ CH] ₂ O | (y) | 7.3 ± 0.4 | 11.1 | 179–339 | CD ₃ COCD ₃ P | 43 |
| | | 9.6 ± 2 | 11.0 | 100–200 | Hg(CH ₃) ₂ P | 42 |
| Peroxides | | | | | | |
| CH ₃ OOCH ₃ | | 10.0 | 12.56 | 124–185 | CH ₃ OOCH ₃ T | 128 |
| ((CH ₃)CHO) ₂ | | | $k = 4.7 \times 10^5$ | 26 | ((CH ₃) ₂ CHO) ₂ P | 129 |
| | | | $k = 5.3 \times 10^6$ | 77 | ((CH ₃) ₂ CHO) ₂ P | 129 |
| ((CH ₃) ₂ CO) ₂ | | 11.7 ± 0.3 | 12.4 | 130–155 | D.T.B.P. T | 105 |
| | | 14.5 ± 2.5 | | 103–145 | D.T.B.P. T | 37 |
| Nitriles | | | | | | |
| CH ₃ CN | (y) | 10.0 ± 0.5 | 11.5 | 100–290 | CD ₃ COCD ₃ P | 130 |
| C ₂ H ₅ CN | (y) | 8.5 ± 0.5 | 11.5 | 133–297 | CD ₃ COCD ₃ P | 130 |
| Metal alkyls | | | | | | |
| Hg(CH ₃) ₂ | (h) | 10.8 | 11.7 | 25–250 | Hg(CH ₃) ₂ P | 131 |
| | | 9.0 ± 0.5 | 11.3 | 25–250 | Hg(CH ₃) ₂ P | 131 |
| | | 10.8 ± 0.3 | 11.7 | 28–251 | Hg(CH ₃) ₂ P | 33 |
| | | 10.2 ± 1.0 | 10.97 | 160–238 | CH ₃ COCH ₃ P | 110 |
| | | | $k = 4.8 \times 10^7$ | 407 | Hg(CH ₃) ₂ T | 132 |
| | (y) | 10.0 ± 1.0 | 11.22 | 125–202 | CD ₃ COCD ₃ P | 110 |
| Cd(CH ₃) ₂ | | 14 ± 2 | 12.8 | 200–275 | Cd(CH ₃) ₂ P | 5 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References | |
|--|-------|---------------------------------|--|-------------------|--|------------|----------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | | |
| Halogenated alkanes | | | | | | | |
| CFH ₃ | (v) | 8.7 ± 0.3 | 11.6 | 125–211 | CH ₃ COCH ₃ | P | 133, 135 |
| | (y) | 11.8 | 11.2 | 193–331 | CD ₃ COCD ₃ | P | 134 |
| CF ₂ H ₂ | (v) | 6.2 ± 0.3 | 10.5 | 129–192 | CH ₃ COCH ₃ | P | 133, 135 |
| | (y) | 10.4 | 11.0 | 122–301 | CD ₃ COCD ₃ | P | 134 |
| CF ₃ H | (y) | 9.8 | 9.8 | 142–293 | CD ₃ COCD ₃ | P | 136, 137 |
| | | 10.4 | 10.0 | 144–309 | CD ₃ COCD ₃ | P | 138 |
| C ₂ F ₅ H | (y) | 9.7 | 10.8 | 135–318 | CD ₃ COCD ₃ | P | 138 |
| C ₃ F ₇ H | (y) | 9.3 | 10.6 | 129–323 | CD ₃ COCD ₃ | P | 138 |
| CH ₃ Cl | (v) | 9.4 ± 0.3 | 11.8 | 127–207 | CH ₃ COCH ₃ | P | 133, 135 |
| CH ₂ Cl ₂ | (v) | 7.2 ± 0.3 | 11.3 | 129–211 | CH ₃ COCH ₃ | P | 133, 135 |
| CHCl ₃ | (v) | 5.8 ± 0.3 | 10.8 | 132–203 | CH ₃ COCH ₃ | P | 133, 135 |
| | | 6.8 | | 30 | CH ₃ COCH ₃ | P | 139 |
| CH ₃ Br | (v) | 10.1 ± 0.3 | 12.5 | 121–208 | CH ₃ COCH ₃ | P | 133, 135 |
| CH ₂ Br ₂ | (v) | 8.7 ± 0.3 | 12.2 | 126–177 | CH ₃ COCH ₃ | P | 133, 135 |
| Hydrides of nitrogen, oxygen, and sulphur | | | | | | | |
| NH ₃ | | 9.8 ± 0.9 | 11.00 ± 0.42 | 110–180 | CH ₃ N ₂ CH ₃ | P | 88, 68 |
| | | 10.0 ± 0.2 | 10.9 ± 0.1 | 144–308 | CH ₃ COCH ₃ | P | 76 |
| | (y) | 10.0 ± 0.4 | 10.8 | 180–339 | CD ₃ COCD ₃ | P | 43 |
| ND ₃ | | 10.9 ± 0.9 | 11.00 ± 0.42 | 110–180 | CH ₃ N ₂ CH ₃ | P | 88, 68 |
| N ₂ H ₄ | | 5.00 ± 0.1 | 11.00 ± 0.05 | 110–180 | CH ₃ N ₂ CH ₃ | P | 88, 68 |
| H ₂ O | (b) | 24.7 | 13.57 | | | | 8 |
| H ₂ S | | 2.6 | 11.4 | 50–140 | CH ₃ COCH ₃ | P | 140 |
| | (w) | 3.6 | 12.1 | 200–360 | CH ₃ CHO | P | 141 |
| Hydrogen halides | | | | | | | |
| HCl | | 2.3 ± 1 | 11.6 | 28–150 | CH ₃ COCH ₃ | P | 142 |
| | (e) | 4.51 | 12.1 | | | | 143, 144 |
| HI | (x) | 2.30 | 12.50 | 260–316 | CH ₃ I/HI | T | 145 |
| FLUORINE ATOM TRANSFER | | | | | | | |
| SF ₆ | | 14.1 | 13.3 | 157–168 | D.T.B.P., T | | 169 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|--|-------|---------------------------------|---|---------------------------|---|--------------------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| CHLORINE ATOM TRANSFER | | | | | | |
| CCl ₄ | | 12.9 ± 0.7 | 13.4 | 90–145 | D.T.B.P. T | 146, 147 |
| CCl ₃ CN | | 10.4 ± 1.0 | 12.9 | 90–145 | D.T.B.P. T | 147 |
| C ₂ Cl ₆ | | 10.1 ± 0.9 | 11.8 | 90–145 | D.T.B.P. T | 146, 147 |
| C ₆ H ₅ CCl ₃ | | 7.6 ± 0.8 | 10.3 | 90–145 | D.T.B.P. T | 147 |
| CCl ₃ COCCl ₃ | | 9.7 ± 0.8 | 12.6 | 90–145 | D.T.B.P. T | 146, 147 |
| BROMINE ATOM TRANSFER | | | | | | |
| CF ₃ Br | | 12.5 ± 1.0 | 13.3 | 90–145 | D.T.B.P. T | 147 |
| CF ₂ Br ₂ | | 6.4 ± 1.0 | 11.0 | 90–145 | D.T.B.P. T | 147 |
| CCl ₃ Br | | 7.1 ± 0.9 | 13.2 | 90–145 | D.T.B.P. T | 147 |
| CCl ₂ Br ₂ | | 7.6 ± 1.1 | 13.8 | 90–145 | D.T.B.P. T | 147 |
| CBr ₄ | | 7.9 ± 1.1 | 14.2 | 90–145 | D.T.B.P. T | 147 |
| IODINE ATOM TRANSFER | | | | | | |
| I ₂ | (aa) | 1.50 | $k = 2 \times 10^{12} - 2 \times 10^{13}$ $k = 2.0 \times 10^{16}$ 13.0 $k = 1.1 \times 10^{13}$ | 20 50 260–316 60 | CH ₃ I/O ₂ P CH ₃ I P CH ₃ I/HI T CH ₃ I F.P. | 148 149 145 165 |
| CF ₃ I | | 7.5 ± 1.0 | 13.8 | 90–145 | D.T.B.P. T | 147 |
| OXYGEN ATOM TRANSFER | | | | | | |
| O ₂ | (bb) | | $k = 5.3 \times 10^8$ | 498 | CH ₃ COCH ₃ /O ₂ T | 150 |
| NO ₂ | | 5.7 | 12.75 | 1150–1590 | CH ₃ NO ₂ ignition | 151 |
| GROUP-TRANSFER REACTIONS | | | | | | |
| ¹⁴ CH ₃ + CH ₃ COCH ₃ = ¹⁴ CH ₃ C ¹⁴ OCH ₃ + CH ₃ | (ee) | | $k = 5.2 \times 10^6$ | 350 | CH ₃ COCH ₃ P | 16 |
| CH ₃ + CF ₃ COCF ₃ = CH ₃ COCF ₃ + CF ₃ | | 5.7 ± 1.5 | 9.3 | 163–245 | CH ₃ N ₂ CH ₃ P, T | 171 |
| | | 6.2 ± 1.0 | 11.4 | 85–210 | CF ₃ COCF ₃ P | 172 |
| | | 7.0 ± 1.0 | 11.7 | 48–240 | CH ₃ COCH ₃ / CF ₃ COCF ₃ P | 173 |

Metathetical Reactions of Methyl Radicals – Continued

| Reactants | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|-----------------------|---------------------------------|-------------------|------------------|-----------|
| | | ($kcal\ mole^{-1}$) | ($cm^3\ mole^{-1}\ sec^{-1}$) | $^{\circ}C$ | | |
| $CH_3 + C_2F_5COC_2F_5$ $= CH_3COC_2F_5 + C_2F_5$ | | 7 ± 1 | | 184–240 | $(C_2F_5)_2CO$ P | 174 |
| $CD_3 + CH_3HgCH_3$ $= CD_3HgCH_3 + CH_3$ | | | $k = 8.9 \times 10^5$ | 180 | CD_3COCD_3 P | 175, 176 |
| $CH_3 + CD_3SH = CH_3SH$ $+ CD_3$ | | 7.6 | 10.73 | 130–200 | CH_3COCH_3 P | 74 |
| $CH_3 + CF_3COCH_3$ $= C_2H_6 + CF_3CO$ | | 14 | | 150–350 | CF_3COCH_3 P | 109 |
| $CH_3 + CH_3COCOCH_3$ $= CH_3COCH_3$ $+ CH_3CO$ | | 5.6 | 10.7 | 100–200 | $CH_3COCOCH_3$ P | 114 |
| | | 6.6 | | 100–200 | $CH_3COCOCH_3$ P | 115 |
| $CH_3 + CH_3CH:CHCHO$ $= CH_3CH:CHCH_3$ $+ CHO$ | | 7.45 ± 1.30 | 11.8 ± 0.4 | 120–250 | CH_3COCH_3 P | 177 |
| $CH_3 + CH_3HgCH_3$ $= C_2H_6 + (Hg + CH_3)$ | | 1.0 | 7 | 175–220 | CH_3HgCH_3 P | 131, 178 |

Ratios of Rate Constants (Methyl Radicals)

(i) Reactions with halogens and hydrogen halides.

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|--|------------------------------------|-------------------------|------------------------------|--|----------------------|
| | | ($kcal\ mole^{-1}$) | | | $^{\circ}C$ | | |
| (1) $CH_3 + HBr = CH_4 + Br$ (2) $CH_3 + Br_2 = CH_3Br + Br$ | | 2.0 | -0.7 | | 37–210 | CH_3I/HBr P | 167, 168 |
| (1) $CH_3 + HI = CH_4 + I$ (2) $CH_3 + I_2 = CH_3I + I$ | | | | 0.15 0.13 ± 0.03 | 280 270–310 | CH_3I/HI T CH_3I/HI T | 152, 154 153, 154 |
| | | 1.3 ± 0.5 0.8 ± 1.0 0.75 | -0.38 -0.50 ± 0.40 -0.64 | | 126–295 260–316 83–192 | CH_3COCH_3 P CH_3I/HI T CH_3I/HI P | 155 145 164 |
| (1) $CH_3 + HBr = CH_4 + Br$ (2) $CH_3 + I_2 = CH_3I + I$ | | 0.8 ± 0.3 1.37 ± 0.26 0.95 | -0.30 ± 0.13 -0.88 | | 60 80–164 83–192 | $CH_3I/HBr/I_2$ P CH_3COCH_3 P CH_3I/HBr P | 163 156 164 |
| (1) $CH_3 + HBr = CH_4 + Br$ (2) $CH_3 + HI = CH_4 + I$ | (cc) | 1.1 ± 0.4 | | 0.15 | 25 | CH_3I/HBr P | 157 |

Ratios of Rate Constants (Methyl Radicals)–Continued

(i) Reactions with halogens and hydrogen halides.—Continued

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|-----------------------------------|---------------------|-----------|-------------------|--|-----------|
| (1) $\text{CH}_3 + \text{CH}_3\text{I} = \text{CH}_4 + \text{CH}_2\text{I}$ (2) $\text{CH}_3 + \text{HI} = \text{CH}_4 + \text{I}$ | | (<i>kcal mole⁻¹</i>) | | 0.03 | °C 270–310 | $\text{CH}_3\text{I}/\text{HI}$ T | 153, 154 |
| (1) $\text{CH}_3 + \text{HCl} = \text{CH}_4 + \text{Cl}$ (2) $\text{CH}_3 + \text{I}_2 = \text{CH}_3\text{I} + \text{I}$ | (dd) | 2.4 | –1.46 | | 98–160 | $\text{CH}_3\text{I}/\text{HCl}$ P | 164 |
| (1) $\text{CH}_3 + \text{I}_2 = \text{CH}_3\text{I} + \text{I}$ (2) $\text{CH}_3 + \text{NO} = \text{CH}_3\text{NO}$ | | | | 6 | 25 | $\text{CH}_3\text{I}/\text{NO}$ P | 158 |
| (1) $\text{CH}_3 + \text{CH}_4 = \text{CH}_4 + \text{CH}_3$ (2) $\text{CH}_3 + \text{I}_2 = \text{CH}_3\text{I} + \text{I}$ | | 6.5 ± 2 | –3.8 | | 30–252 | $\text{CH}_3\text{I}/\text{CH}_4/\text{I}_2$ P | 161 |

(ii) Miscellaneous reactions.

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|-----------------------------------|---------------------|---|-------------------|---|------------|
| (1) $\text{CH}_3 + \text{CH}_3\text{CD}_3 = \text{CH}_3\text{D} + \text{CH}_3\text{CD}_2$ (2) $\text{CH}_3 + \text{CH}_3\text{CD}_3 = \text{CH}_4 + \text{CH}_2\text{CD}_3$ | | (<i>kcal mole⁻¹</i>) | | | °C 328–424 | $\text{CH}_3\text{N}_2\text{CH}_3$ T | 159, 27 |
| (1) $\text{CH}_3 + \text{CH}_3\text{CD}_2\text{CH}_3 = \text{CH}_3\text{D} + \text{CH}_3\text{CDCH}_3$ (2) $\text{CH}_3 + \text{CH}_3\text{CD}_2\text{CH}_3 = \text{CH}_4 + \text{CH}_2\text{CD}_2\text{CH}_3$ | | 0.6 ± 0.04 | –0.32 | | 312–421 | $\text{CH}_3\text{N}_2\text{CH}_3$ T | 159, 27 |
| (1) $\text{CH}_3 + (\text{CH}_3)_3\text{CD} = \text{CH}_3\text{D} + (\text{CH}_3)_3\text{C}$ (2) $\text{CH}_3 + (\text{CH}_3)_3\text{CD} = \text{CH}_4 + \text{CH}_2(\text{CH}_3)_2\text{CD}$ | | –1.7 | –0.66 | | 305–359 | $\text{CH}_3\text{N}_2\text{CH}_3$ T | 159, 27 |
| (1) $\text{CD}_3 + \text{CD}_3\text{COCD}_2\text{CH}_2\text{CH}_3 = \text{CD}_4 + [\text{C}_5\text{D}_4\text{H}_5\text{O}]$ (2) $\text{CD}_3 + \text{CD}_3\text{COCD}_2\text{CH}_2\text{CH}_3 = \text{CD}_3\text{H} + [\text{C}_5\text{D}_5\text{H}_4\text{O}]$ | | –2.3 | –0.51 | | 35–144 | $\text{CD}_3\text{COCD}_2\text{CH}_2\text{CH}_3$ P, T | 36 |
| (1) $\text{CH}_3 + \text{CH}_3\text{OCH}_2\text{CH}_3 = \text{CH}_4 + [\text{C}_3\text{H}_7\text{O}]$ (2) $\text{CH}_3 + \text{CH}_3\text{OCH}_3 = \text{CH}_4 + \text{CH}_2\text{OCH}_3$ | | 0.7 | | | 525 | CH_3OCH_3 T | 160 |
| (1) $\text{CH}_3 + \text{HCHO} = \text{CH}_4 + \text{HCO}$ (2) $\text{CH}_3 + \text{CH}_3\text{OCH}_3 = \text{CH}_4 + \text{CH}_2\text{OCH}_3$ | | | | 4 to 12 | 507 | CH_3OCH_3 T | 9 |
| (1) $\text{CH}_3 + \text{O}_2 = \text{HCHO} + \text{OH}$ (2) $\text{CH}_3 + \text{O}_2 + \text{M} = \text{CH}_3\text{O}_2 + \text{M}$ | | | | 1.0×10^{-8} $k_1/k_2 [\text{M}] = 0.49$ | 25 162 | $\text{CH}_3\text{I}/\text{O}_2$ P $\text{CH}_3\text{N}_2\text{CH}_3/\text{O}_2$ P | 170 162 |

Notes

- (a) When corrected, these activation energies are 10.5 and 12.2 kcal respectively (see ref. 11).
(b) This value was used (without a source being quoted), in the range 1000–1500 °C.
(c) Use of the more recent activation energy for $\text{CH}_3 + \text{CH}_3\text{CHO}$ gives $E = 12.0$ kcal (see ref. 11).
(d) For a re-analysis of these results see reference 15.
(e) Calculated from the reverse reaction.
(f) Assuming an activation energy of 9.2 kcal/mole for the reaction $\text{CD}_3 + \text{CD}_3\text{CDO} = \text{CD}_4 + \text{CD}_3\text{CO}$.
(g) Assuming $k = 10^{12.3} \exp(-13,600/RT)$ for the reaction $\text{CH}_3 + \text{C}_2\text{D}_6 = \text{CH}_3\text{D} + \text{C}_2\text{D}_5$.
(h) The quantities were not calculated in this manner by the original authors.
(i) Assuming $k = 10^{11.7} \exp(-10,800/RT)$ for the reaction $\text{CH}_3 + \text{CH}_3\text{HgCH}_3 = \text{CH}_4 + \text{CH}_3\text{HgCH}_2$.
(j) Pyrolysis of a series of methyl metal derivatives.
(k) Assuming $k = 10^{11.8} \exp(-12,000/RT)$ for the reaction $\text{CH}_3 + \text{D}_2 = \text{CH}_3\text{D} + \text{D}$.
(l) No distinction made between *o*, *m*, and *p* xylenes.
(m) Calculated assuming a zero secondary isotope effect.
(n) These are not the values quoted by the authors, but corrected (see ref. 67) for arithmetical errors.
(o) No significant quantities of CH_3D were formed, when acetone was photolyzed in presence of $(\text{CD}_2)_2\text{NH}$.
(p) These values are doubtful.

- (q) No significant quantities of CH_3D formed during the photolysis of CH_3COND_2 .
(r) Calculated assuming the parameters for abstraction from the $-\text{NH}_2$ group, are as for those in hydrazine.
(s) See also references 69, 70, 75, 79, 80, and 88.
(t) Several other studies are in excellent agreement with those listed. See references 1, 23, 59, 65, 74, 106, 107, and 108.
(u) "A" factor doubtful.
(v) The results for CCl_4 , CH_2Cl_2 , and CH_3Cl are invalid; the others are to be regarded with some suspicion (see ref. 135).
(w) Assuming $k = 10^{11.9} \exp(-7600/RT)$ for the reaction $\text{CH}_3 + \text{CH}_3\text{CHO} = \text{CH}_4 + \text{CH}_3\text{CO}$.
(x) The authors consider the activation energy to be ~ 1 kcal too high.
(y) The attacking radical is CD_3 .
(z) The variations reported within these series may be spurious as variations in E are compensated by variations in A .
(aa) The authors consider the activation energy to be ~ 1 kcal too high.
(bb) The products of this reaction are HCHO and OH .
(cc) Activation energy difference calculated assuming identical A factors.
(dd) "Hot" methyl radicals may have played an important part in this system.
(ee) Assuming $k = 10^{11.6} \exp(-9800/RT)$ for the reaction $\text{CH}_3 + \text{CH}_3\text{COCH}_3 = \text{CH}_4 + \text{CH}_2\text{COCH}_3$.

References (Methyl Radical Reactions)

1. Majury and Steacie, *Disc. Faraday Soc.* **14**, 45 (1953).
2. Majury and Steacie, *Can. J. Chem.* **30**, 800 (1952).
3. Whittle and Steacie, *J. Chem. Phys.* **21**, 993 (1953).
4. Phibbs and Darwent, *Trans. Faraday Soc.* **45**, 541 (1949).
5. Anderson and Taylor, *J. Phys. Chem.* **56**, 498 (1952).
6. Davison and Burton, *J. Am. Chem. Soc.* **74**, 2307 (1952).
7. Gesser and Steacie, *Can. J. Chem.* **34**, 113 (1956).
8. Intezarova, Kondratiev, and Mukhoyan, *Kinetika i Kataliz*, **5**, 585 (1964).
9. Benson and Jain, *J. Chem. Phys.* **31**, 1008 (1959).
10. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
11. Wijnen and Steacie, *Disc. Faraday Soc.* **14**, 118 (1953).
12. Rebbert and Steacie, *Can. J. Chem.* **32**, 113 (1954).
13. Chanmugam and Burton, *J. Am. Chem. Soc.* **78**, 509 (1956).
14. McNesby, Gordon, and Smith, *J. Am. Chem. Soc.* **78**, 1287 (1956).
15. Pritchard and Pritchard, *Can. J. Chem.* **41**, 3042 (1963).
16. Dainton, Ivin, and Wilkinson, *Trans. Faraday Soc.* **55**, 929 (1959).
17. Creak, Dainton, and Ivin, *Trans. Faraday Soc.* **58**, 326 (1962).
18. Trotman-Dickenson, "Gas Kinetics," p. 199 (Butterworths, 1955).
19. Trotman-Dickenson and Steacie, *J. Phys. Chem.* **55**, 908 (1951).
20. McNesby and Gordon, *J. Am. Chem. Soc.* **76**, 4196 (1954).
21. Wijnen, *J. Chem. Phys.* **23**, 1357 (1955).
22. Dainton, Ivin, and Wilkinson, *Trans. Faraday Soc.* **53**, 1204 (1957).
23. Dainton and McElcheran, *Trans. Faraday Soc.* **51**, 657 (1955).
24. Brodskii, Kalinenko, and Lavrovskii, *J. Chem. Soc.* **4443** (1960).
25. Blackmore and Hinshelwood, *Proc. Roy. Soc.* **271A**, 34 (1963).
26. Trotman-Dickenson, Birchard, and Steacie, *J. Chem. Phys.* **19**, 163 (1951).
27. McNesby, *J. Phys. Chem.* **64**, 1671 (1960).
28. McNesby and Gordon, *J. Am. Chem. Soc.* **77**, 4719 (1955).
29. Rice and Varnerin, *J. Am. Chem. Soc.* **77**, 221 (1955).
30. Jackson, McNesby, and Darwent, *J. Chem. Phys.* **37**, 1610 (1962).
31. Jackson and McNesby, *J. Am. Chem. Soc.* **83**, 4891 (1961).
32. Gomer, *J. Am. Chem. Soc.* **72**, 201 (1950).
33. Rebbert and Steacie, *J. Chem. Phys.* **21**, 1723 (1953).
34. Smith and Taylor, *J. Chem. Phys.* **7**, 390 (1939).
35. Jones and Steacie, *Can. J. Chem.* **31**, 505 (1953).
36. McNesby and Gordon, *J. Am. Chem. Soc.* **78**, 3570 (1956).
37. Blake and Kutschke, *Can. J. Chem.* **37**, 1462 (1959).
38. Blake, Henderson, and Kutschke, *Can. J. Chem.* **39**, 1920 (1961).
39. Rice and Vanderslice, *J. Am. Chem. Soc.* **80**, 291 (1958).
40. Blackmore and Hinshelwood, *Proc. Roy. Soc.* **268A**, 36 (1962).
41. Rebbert and Steacie, *Can. J. Chem.* **32**, 40 (1954).
42. Phibbs and Darwent, *Can. J. Res.* **28B**, 395 (1950).
43. Trotman-Dickenson and Steacie, *J. Chem. Phys.* **19**, 329 (1951).
44. McNesby and Gordon, *J. Am. Chem. Soc.* **79**, 825 (1957).
45. Gordon, Smith, and Drew, *J. Chem. Phys.* **36**, 824 (1962).
46. Russell and Bernstein, *J. Chem. Phys.* **30**, 607 (1959).
47. Gordon, *Can. J. Chem.* **43**, 570 (1965).
48. Gordon and Smith, *J. Phys. Chem.* **66**, 521 (1962).
49. Trotman-Dickenson and Steacie, *J. Chem. Phys.* **19**, 169 (1951).
50. Miyoshi and Brinton, *J. Chem. Phys.* **36**, 3019 (1962).
51. Rice and Varnerin, *J. Am. Chem. Soc.* **76**, 2629 (1954).
52. Drew and Gordon, *J. Chem. Phys.* **31**, 1417 (1959).
53. Mulcahy, Williams, and Wilmhurst, *Austral. J. Chem.* **17**, 1329 (1964).
54. Taylor and Smith, *J. Chem. Phys.* **8**, 543 (1940).
55. Szwarc and Roberts, *Trans. Faraday Soc.* **46**, 625 (1950).
56. Price and Trotman-Dickenson, *J. Chem. Soc.* **4205** (1958).
57. Price and Trotman-Dickenson, *Trans. Faraday Soc.* **53**, 939 (1957).
58. Price and Trotman-Dickenson, *Trans. Faraday Soc.* **53**, 1208 (1957).
59. Burkley and Rebbert, *J. Phys. Chem.* **67**, 168 (1963).
60. Cher, *J. Phys. Chem.* **68**, 1316 (1964).
61. Sanders and Rebbert, *J. Phys. Chem.* **67**, 170 (1963).
62. Burr and Strong, *J. Chem. Phys.* **43**, 1432 (1965).
63. Burr and Strong, *J. Am. Chem. Soc.* **86**, 5065 (1964).
64. Wunderlich and Rebbert, *J. Phys. Chem.* **67**, 1382 (1963).
65. Harrison and Shannon, *Can. J. Chem.* **41**, 2455 (1963).
66. Kerr and Trotman-Dickenson, *J. Chem. Soc.* **3322** (1957).

67. Brinton, *Can. J. Chem.* **38**, 1339 (1960).
68. Gray and Thynne, 10th Int. Comb. Symp., p. 435, Comb. Inst., Pittsburgh (1965).
69. Gray and Thynne, *Trans. Faraday Soc.* **59**, 2275 (1963).
70. Gray, Jones, and Thynne, *Trans. Faraday Soc.* **61**, 474 (1965).
71. Kozak and Gesser, *J. Chem. Soc.* 448 (1960).
72. Cher, Hollingsworth, and Sicilio, *J. Phys. Chem.* **70**, 877 (1966).
73. Shaw and Thynne, *Trans. Faraday Soc.* **62**, 104 (1966).
74. Greig and Thynne, *Trans. Faraday Soc.* **62**, 379 (1966).
75. Gray and Jones, *Trans. Faraday Soc.* **62**, 112 (1966).
76. Edwards, Kerr, Lloyd, and Trotman-Dickenson, *J. Chem. Soc.* 621 (1966).
77. Brinton and Volman, *J. Chem. Phys.* **20**, 25 (1952).
78. Klemm, *Can. J. Chem.* **43**, 2633 (1965).
79. Gray and Jones, *Trans. Faraday Soc.* **61**, 2161 (1965).
80. Gray and Jones, *Can. J. Chem.* **43**, 3485 (1965).
81. Cafferata, Kerr, and Trotman-Dickenson, *J. Chem. Soc.* 1386 (1965).
82. Spall and Steacie, *Proc. Roy. Soc.* **239A**, 1 (1957).
83. Thynne, *Trans. Faraday Soc.* **60**, 2207 (1964).
84. Jones and Steacie, *J. Chem. Phys.* **21**, 1018 (1953).
85. Ausloos and Steacie, *Can. J. Chem.* **32**, 593 (1954).
86. Toby, *J. Am. Chem. Soc.* **82**, 3822 (1960).
87. Toby and Kutschke, *Can. J. Chem.* **37**, 672 (1959).
88. Gray and Thynne, *Trans. Faraday Soc.* **60**, 1047 (1964).
89. Toby and Nimoy, *J. Phys. Chem.* **70**, 867 (1966).
90. Gowenlock, *Can. J. Chem.* **42**, 1936 (1964).
91. Hoare, *Trans. Faraday Soc.* **49**, 1292 (1953).
92. Brinton and Volman, *J. Chem. Phys.* **20**, 1053 (1952).
93. Ausloos and Steacie, *Can. J. Chem.* **33**, 31 (1955).
94. Dodd, *Can. J. Chem.* **33**, 699 (1955).
95. Calvert, Pitts, and Thompson, *J. Am. Chem. Soc.* **78**, 4239 (1956).
96. Birrell and Trotman-Dickenson, *J. Chem. Soc.* 2059 (1960).
97. Kerr and Calvert, *J. Phys. Chem.* **69**, 1022 (1965).
98. Pritchard, Hsia, and Miller, *J. Am. Chem. Soc.* **85**, 1568 (1963).
99. Volman and Brinton, *J. Chem. Phys.* **22**, 929 (1954).
100. Trotman-Dickenson and Steacie, *J. Chem. Phys.* **18**, 1097 (1950).
101. Saunders and Taylor, *J. Chem. Phys.* **9**, 616 (1941).
102. Gomer and Kistiakowsky, *J. Chem. Phys.* **19**, 85 (1951).
103. Mandelcorn and Steacie, *Can. J. Chem.* **32**, 331 (1954).
104. Jacquiss, Roberts, and Szwarc, *J. Am. Chem. Soc.* **74**, 6005 (1952).
105. Pritchard, Pritchard, and Trotman-Dickenson, *J. Chem. Soc.* 1425 (1954).
106. Thynne and Gray, *Trans. Faraday Soc.* **58**, 2403 (1962).
107. March and Polanyi, *Proc. Roy. Soc.* **273A**, 360 (1963).
108. Brinton, *J. Am. Chem. Soc.* **83**, 1541 (1961).
109. Sieger and Calvert, *J. Am. Chem. Soc.* **76**, 5197 (1954).
110. Oswin, Rebbert, and Steacie, *Can. J. Chem.* **33**, 472 (1955).
111. Wijnen, *J. Chem. Phys.* **22**, 1631 (1954).
112. Pitts and Norman, *J. Am. Chem. Soc.* **76**, 4815 (1954).
113. Duncan and Trotman-Dickenson, *J. Chem. Soc.* 4672 (1962).
114. Blacet and Bell, *Disc. Faraday Soc.* **14**, 70 (1953).
115. Bell and Blacet, *J. Am. Chem. Soc.* **76**, 5332 (1954).
116. Ausloos and Steacie, *Can. J. Chem.* **33**, 39 (1955).
117. Thynne, *Trans. Faraday Soc.* **58**, 676 (1962).
118. Thynne, *Trans. Faraday Soc.* **58**, 1394 (1962).
119. Thynne, *Trans. Faraday Soc.* **58**, 1533 (1962).
120. Wijnen, *J. Chem. Phys.* **27**, 710 (1957).
121. Wijnen, *Can. J. Chem.* **36**, 176 (1958).
122. Wijnen, *J. Chem. Phys.* **28**, 939 (1958).
123. Wijnen, *J. Am. Chem. Soc.* **80**, 2394 (1958).
124. Ausloos and Steacie, *Can. J. Chem.* **33**, 1530 (1955).
125. Ausloos, *Can. J. Chem.* **34**, 1709 (1956).
126. Marcus, Darwent, and Steacie, *J. Chem. Phys.* **16**, 987 (1948).
127. Long and Skirrow, *Trans. Faraday Soc.* **58**, 1403 (1962).
128. Thynne and Gray, *Trans. Faraday Soc.* **59**, 1149 (1963).
129. McMillan, *J. Am. Chem. Soc.* **83**, 3018 (1961).
130. Wijnen, *J. Chem. Phys.* **22**, 1074 (1954).
131. Gomer and Noyes, *J. Am. Chem. Soc.* **71**, 3390 (1949).
132. Kallend and Purnell, *Trans. Faraday Soc.* **60**, 103 (1964).
133. Raal and Steacie, *J. Chem. Phys.* **20**, 578 (1952).
134. Pritchard, Bryant, and Thommarson, *J. Phys. Chem.* **69**, 664 (1965).
135. Ridge and Steacie, *Can. J. Chem.* **33**, 396 (1955).
136. Pritchard, Pritchard, Schiff, and Trotman-Dickenson, *Trans. Faraday Soc.* **52**, 849 (1956).
137. Pritchard, Pritchard, Schiff, and Trotman-Dickenson, *Chem. and Ind.* 896 (1955).
138. Pritchard and Thommarson, *J. Phys. Chem.* **68**, 568 (1964).
139. Cvetanovic, Raal, and Steacie, *Can. J. Chem.* **31**, 171 (1953).
140. Imai and Toyama, *Bull. Chem. Soc. Japan* **33**, 652 (1960).
141. Imai and Toyama, *Bull. Chem. Soc. Japan* **33**, 1120 (1960).
142. Cvetanovic and Steacie, *Can. J. Chem.* **31**, 158 (1953).
143. Pritchard, Pyke, and Trotman-Dickenson, *J. Am. Chem. Soc.* **77**, 2629 (1955).
144. Pritchard, Pyke, and Trotman-Dickenson, *J. Am. Chem. Soc.* **76**, 1201 (1954).
145. Flowers and Benson, *J. Chem. Phys.* **38**, 882 (1963).
146. Tomkinson, Galvin, and Pritchard, *J. Phys. Chem.* **68**, 541 (1964).
147. Tomkinson and Pritchard, *J. Phys. Chem.* **70**, 1579 (1966).
148. Christie, *Proc. Roy. Soc.* **244A**, 411 (1958).
149. Souffie, Williams, and Hamill, *J. Am. Chem. Soc.* **78**, 917 (1956).
150. Barnard and Honeyman, *Proc. Roy. Soc.* **279A**, 244 (1964).
151. Hiraoka and Hardwick, *J. Chem. Phys.* **39**, 2361 (1963).
152. Benson and O'Neal, *J. Chem. Phys.* **34**, 514 (1961).
153. Sullivan, *J. Phys. Chem.* **65**, 722 (1961).
154. Ogg, *J. Am. Chem. Soc.* **56**, 526 (1934).
155. O'Neal and Benson, *J. Chem. Phys.* **36**, 2196 (1962).
156. Fettis and Trotman-Dickenson, *J. Chem. Soc.* 3037 (1961).
157. Farren, Gilbert, Linnett and Read, *Trans. Faraday Soc.* **60**, 740 (1964).
158. Christie and Frost, *Trans. Faraday Soc.* **61**, 468 (1965).
159. Rice and Vanderslice, *J. Am. Chem. Soc.* **80**, 291 (1958).
160. Anderson and Benson, *J. Chem. Phys.* **36**, 2320 (1962).
161. Willard and Harris, *J. Am. Chem. Soc.* **76**, 4678 (1954).
162. Wenger and Kutschke, *Can. J. Chem.* **37**, 1546 (1959).
163. Anderson and Kistiakowsky, *J. Chem. Phys.* **11**, 6 (1943).
164. Williams and Ogg, *J. Chem. Phys.* **15**, 696 (1947).
165. Davidson and Carrington, *J. Am. Chem. Soc.* **74**, 6277 (1952).
166. Ausloos and Steacie, *Can. J. Chem.* **33**, 1062 (1955).
167. Kistiakowsky and Van Artsdalen, *J. Chem. Phys.* **12**, 469 (1944).
168. Benson and Buss, *J. Chem. Phys.* **28**, 301 (1958).
169. Batt and Cruickshank, *J. Phys. Chem.* **70**, 723 (1966).
170. Heicklen and Johnston, *J. Am. Chem. Soc.* **84**, 4030 (1962).
171. Pritchard and Steacie, *Can. J. Chem.* **35**, 1216 (1957).
172. Alcock and Whittle, *Trans. Faraday Soc.* **61**, 244 (1965).
173. Giles and Whittle, *Trans. Faraday Soc.* **61**, 1425 (1965).
174. Price and Kutschke, *Can. J. Chem.* **38**, 2128 (1960).
175. Rebbert and Ausloos, *J. Am. Chem. Soc.* **86**, 2068 (1964).
176. Rebbert and Ausloos, *J. Am. Chem. Soc.* **85**, 3086 (1963).
177. Allen and Pitts, *J. Phys. Chem.* **70**, 169 (1966).
178. Trotman-Dickenson, "Gas Kinetics," p. 237 et seq. (Butterworths, London, 1955).

Difluoromethylene Radicals

| Reaction | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|-------|--|---|----------------------------|---|-----------|
| $\text{CF}_2 + \text{O}_2 = \text{CO} + 2\text{F} + \text{O}$ | | (<i>kcal mole⁻¹</i>) 14.98 | (<i>cm³ mole⁻¹ sec⁻¹</i>) 12.30 | ^{°C} 1307-2117 | $\text{C}_2\text{F}_4/\text{O}_2$ S.T.* | 1 |

References

1. Modica and La Graff, J. Chem. Phys. **43**, 3383 (1965).

Trifluoromethyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|--|-------|----------------------------|---|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | (kcal mole ⁻¹) | (mole ⁻¹ cc. sec ⁻¹) | °C | | |
| H ₂ | (a) | 9.5 ± 0.7 | 11.86 | 59-158 | CF ₃ COCF ₃ | 1 |
| | | 8.8 | 11.60 | 219-346 | CF ₃ N ₂ CF ₃ | 2 |
| | | 25.1 | 11.89 | 832-1011 | | 3 |
| HD | | 10.5 ± 1.5 | 11.78 | 102-174 | CF ₃ COCF ₃ | 1 |
| HD | | 10.2 ± 1.5 | 11.35 | 102-174 | CF ₃ COCF ₃ | 1 |
| D ₂ | | 10.2 ± 0.7 | 11.45 | 86-196 | CF ₃ COCF ₃ | 1 |
| | | 9.7 | 11.44 | 220-346 | CF ₃ N ₂ CF ₃ | 2 |
| Alkanes | | | | | | |
| CH ₄ | | 10.3 ± 0.5 | 11.70 | 122-251 | CF ₃ COCF ₃ | 4 |
| | | 10.6 | 11.73 | 170-310 | CF ₃ CHO | 5 |
| | | 9.5 ± 2 | 11.2 | 30-350 | CF ₃ COCF ₃ | 6 |
| | | 11.0 ± 0.1 | 11.98 | 84-261 | CF ₃ COCF ₃ | 7 |
| | | 11.3 ± 0.5 | 11.96 ± 0.22 | 153-295 | CF ₃ COCF ₃ | 8 |
| | | 11.0 | 11.98 | | | |
| CHD ₃ | | 10.5 ± 0.3 | 11.04 | 55-354 | CF ₃ COCF ₃ | 9 |
| CHD ₃ | | 12.75 ± 0.3 | 11.33 | 55-354 | CF ₃ COCF ₃ | 9 |
| CD ₄ | | 12.1 ± 2.1 | 11.18 | 106-287 | CF ₃ COCF ₃ | 7 |
| C ₂ H ₆ | | 7.5 ± 0.5 | 11.68 | 81-216 | CF ₃ COCF ₃ | 4 |
| | | 7.5 | 11.63 | 220-350 | CF ₃ N ₂ CF ₃ | 2 |
| C ₃ H ₈ | | 6.2 | 11.67 | 200-310 | CF ₃ N ₂ CF ₃ | 2 |
| | | 6.5 ± 0.5 | 11.75 | 27-119 | CF ₃ COCF ₃ | 10 |
| <i>n</i> -C ₄ H ₁₀ | | 5.1 ± 0.3 | 11.15 | 29-93 | CF ₃ COCF ₃ | 10 |
| | | 5.5 ± 1.0 | 10.9 | 30-350 | CF ₃ COCF ₃ | 6 |
| | | 5.3 | 11.46 | 200-310 | CF ₃ N ₂ CF ₃ | 2 |
| iso-C ₄ H ₁₀ | | 4.7 ± 0.3 | 11.15 | 28-84 | CF ₃ COCF ₃ | 10 |
| | | 4.7 | 11.17 | 170-240 | CF ₃ CHO | 5 |
| (CH ₃) ₃ CD | | 7.4 ± 1.0 | 12.48 | 62-208 | CF ₃ COCF ₃ | 7 |
| neo-C ₅ H ₁₂ | | 7.6 | 11.76 | 250-320 | CF ₃ N ₂ CF ₃ | 2 |
| CH ₃ C(CH ₃) ₂ CH ₂ CH ₃ | | 1.7 | 10.22 | 250-300 | CF ₃ N ₂ CF ₃ | 2 |
| Cyclo-Alkanes | | | | | | |
| cyclo-C ₅ H ₁₀ | | 4.7 | 11.54 | 210-300 | CF ₃ N ₂ CF ₃ | 2 |
| cyclo-C ₆ H ₁₂ | | 5.0 ± 0.2 | 11.44 | 19-91 | CF ₃ COCF ₃ | 11 |

Trifluoromethyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|-------|------------------------------------|--|-------------------|---|-----------|
| | | (<i>kcal mole</i> ⁻¹) | (<i>mole</i> ⁻¹ <i>cc. sec</i> ⁻¹) | °C | | |
| Aromatic hydrocarbons | | | | | | |
| C ₆ H ₆ | (b) | 6.3 ± 0.20 | 11.35 | 23–86 | CF ₃ COCF ₃ | 11 |
| | | 7.7 | 11.44 | | CF ₃ N ₂ CF ₃ | 2 |
| C ₆ H ₅ CH ₃ | (b) | 6.0 | 11.33 | 27–110 | CF ₃ N ₂ CF ₃ | 2 |
| | | 5.91 ± 0.30 | 11.59 ± 0.20 | | CF ₃ COCF ₃ | 12 |
| | | 5.37 ± 0.26 | 11.10 | | CF ₃ COCF ₃ | 11 |
| C ₆ H ₅ CD ₃ | | 5.8 ± 0.4 | 10.13 | 22–72 | CF ₃ COCF ₃ | 13 |
| <i>o</i> -C ₆ H ₄ (CH ₃) ₂ | | 5.55 ± 0.54 | 11.68 | 37–68 | CF ₃ COCF ₃ | 11 |
| Halogenated Aromatics | | | | | | |
| C ₆ H ₅ Cl | | 5.5 ± 0.3 | 10.56 | 20–118 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ Br | | 5.9 ± 0.5 | 11.53 | 23–95 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ I | | 4.5 ± 0.1 | 10.86 | 75–150 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ CH ₂ Cl | | 8.0 ± 0.3 | 12.46 | 67–151 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ CCl ₃ | | 8.8 ± 0.6 | 12.55 | 104–171 | CF ₃ COCF ₃ | 14 |
| C ₆ F ₅ CH ₃ | | 6.38 | 10.75 | | CF ₃ COCF ₃ | 15 |
| Aldehydes | | | | | | |
| CH ₃ CHO | | 4.2 | 10.84 | 170–250 | CF ₃ CHO | 5 |
| CF ₃ CHO | | 8.4 | 11.73 | 150–400 | CF ₃ CHO | 5 |
| Ketones | | | | | | |
| CH ₃ COCH ₃ | | 8.0 | 11.51 | 250–320 | CF ₃ N ₂ CF ₃ | 2 |
| | | 6.9 ± 0.1 | 10.77 | 29–169 | CF ₃ COCF ₃ | 16 |
| | | 8.27 ± 0.17 | 11.52 ± 0.08 | 85–240 | CF ₃ COCF ₃ | 17, 16 |
| CF ₃ COCH ₃ | | 6.6 | 11.3 | 25–350 | CF ₃ COCH ₃ | 18 |
| C ₆ H ₅ COCF ₃ | | 7.2 ± 0.5 | 9.7 | 158–304 | C ₆ H ₅ COCF ₃ | 19 |
| Halogenated Methanes | | | | | | |
| CH ₃ Cl | | 10.6 ± 0.2 | 12.11 ± 0.08 | 144–400 | CF ₃ COCF ₃ | 8 |
| CH ₂ Cl ₂ | | 7.6 ± 0.1 | 11.19 ± 0.08 | 85–204 | CF ₃ COCF ₃ | 20 |
| CHCl ₃ | | 6.6 ± 0.1 | 11.04 ± 0.06 | 94–296 | CF ₃ COCF ₃ | 20 |
| | | 5.3 ± 0.1 | 10.16 ± 0.07 | 35–338 | CF ₃ N ₂ CF ₃ | 21 |
| | | 6.3 ± 0.6 | 11.98 | 118–251 | CF ₃ COCF ₃ | 7 |

Trifluoromethyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|-----------------------------------|---|-------------------|-----------------------------------|-----------|
| | | (<i>kcal mole⁻¹</i>) | (<i>mole⁻¹ cc. sec⁻¹</i>) | °C | | |
| CDCl ₃ | | 9.0 ± 0.4 | 11.78 | 65–264 | CF ₃ COCF ₃ | 7 |
| CH ₃ Br | (d) | 10.9 ± 0.3 | 12.19 ± 0.14 | 150–260 | CF ₃ COCF ₃ | 8 |
| | (e) | 9.9 ± 0.1 | 11.63 ± 0.06 | 150–260 | CF ₃ COCF ₃ | 8 |
| CH ₃ I | | 7.5 ± 0.3 | 10.63 ± 0.16 | 55–210 | CF ₃ COCF ₃ | 8 |
| Hydrogen halides and hydrogen sulphide | | | | | | |
| HCl | | 5.1 ± 0.5 | 11.05 | 20–205 | CF ₃ COCF ₃ | 26 |
| HBr | | 2.9 ± 0.5 | 11.78 | 26 | CF ₃ COCF ₃ | 26 |
| | | | $k = 3.0 \times 10^8$ | | CF ₃ COCF ₃ | 22 |
| H ₂ S | | 3.88 ± 0.26 | 11.65 ± 0.16 | 63–100 | CF ₃ COCF ₃ | 27 |

CHLORINE ATOM TRANSFER (i)

| | | | | | | |
|--|-----|------------|--------------|---------|--|----|
| Cl ₂ | | 3.6 ± 0.5 | 12.89 | 126–235 | CF ₃ COCF ₃ | 26 |
| CH ₃ Cl | | ≥ 17 | | | CF ₃ COCF ₃ | 20 |
| CH ₂ Cl ₂ | | 11.8 ± 0.7 | 11.49 ± 0.1 | 306–449 | CF ₃ COCF ₃ | 20 |
| CHCl ₃ | (f) | 12.0 ± 0.2 | 12.08 ± 0.1 | 95–296 | CF ₃ COCF ₃ | 20 |
| | | 11.0 ± 0.4 | 11.33 ± 0.2 | 35–338 | CF ₃ N ₂ CF ₃ | 21 |
| CCl ₄ | | 9.3 ± 0.4 | 11.79 ± 0.2 | 122–288 | CF ₃ N ₂ CF ₃ | 21 |
| | | 10.4 ± 0.1 | 12.57 ± 0.04 | 96–240 | CF ₃ COCF ₃ | 20 |
| C ₆ H ₅ Cl | | ≥ 13 | | 20–118 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ CH ₂ Cl | | ≥ 13 | | 67–151 | CF ₃ COCF ₃ | 14 |
| C ₆ H ₅ CCl ₃ | | 9.7 ± 0.5 | 12.49 | 104–171 | CF ₃ COCF ₃ | 14 |

BROMINE ATOM TRANSFER

| | | | | | | |
|----------------------------------|-----|-----------|--------------|----------|-----------------------------------|----|
| Br ₂ | | 0.7 ± 0.5 | 12.36 | 178–327 | CF ₃ COCF ₃ | 26 |
| CH ₃ Br | (d) | 8.4 ± 0.1 | 10.83 ± 0.06 | 150–260 | CF ₃ COCF ₃ | 8 |
| | (e) | 8.1 ± 0.2 | 10.41 ± 0.10 | 150–260 | CF ₃ COCF ₃ | 8 |
| C ₆ H ₅ Br | (g) | 1.7 ± 1.1 | 7.91 | 23–95 | CF ₃ COCF ₃ | 14 |
| C ₆ F ₅ Br | (g) | 0.3 ± 0.6 | 7.38 | 39–107 | CF ₃ COCF ₃ | 14 |
| HBr | (a) | 84.2 | 10.71 | 832–1011 | | 23 |

Trifluoromethyl Radicals—Continued

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|-----------|-------|---------------------------------|---|-------------------|----------------|-----------|
| | | <i>(kcal mole⁻¹)</i> | <i>(mole⁻¹ cc. sec⁻¹)</i> | °C | | |

IODINE ATOM TRANSFER

| | | | | | | |
|---------------------------------|--|------------|-------------|--------|-----------------------------------|----|
| I ₂ | | 0.0 ± 0.5 | 12.42 | | CF ₃ COCF ₃ | 26 |
| CH ₃ I | | 3.3 ± 0.15 | 9.59 ± 0.08 | 55–210 | CF ₃ COCF ₃ | 8 |
| C ₆ H ₅ I | | 3.1 ± 0.6 | 10.33 | 75–150 | CF ₃ COCF ₃ | 14 |

OXYGEN ATOM TRANSFER

| | | | | | | |
|------------------|-----|------|-------|---------|-----------------------------------|----|
| N ₂ O | (h) | 24.0 | 13.15 | 316–375 | CF ₃ COCF ₃ | 24 |
|------------------|-----|------|-------|---------|-----------------------------------|----|

Ratios of Rate Constants (Trifluoromethyl Radicals)

| Reaction | Notes | E ₁ – E ₂ | log ₁₀ A ₁ /A ₂ | Temperature range | Radical source | Reference |
|---|-------|---|--|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | °C | | |
| (1) CF ₃ + Br ₂ = CF ₃ Br + Br | | –0.52 ± 1.88 | 0.93 ± 0.60 | 361–431 | CF ₃ H thermal bromination | 22 |
| (2) CF ₃ + HBr = CF ₃ H + Br | | –2.17 ± 0.16 | 0.58 ± 0.08 | 55–334 | CF ₃ COCF ₃ | 25 |
| (1) CF ₃ + I ₂ = CF ₃ I + I | | | | | | |
| (2) CF ₃ + HBr = CF ₃ H + Br | | –2.98 ± 0.12 | 0.64 ± 0.06 | 85–230 | CF ₃ COCF ₃ | 25 |
| (1) CF ₃ + CHD ₃ = CF ₃ H + CD ₃ | | | | | | |
| (2) CF ₃ + CHD ₃ = CF ₃ D + CHD ₂ | | –2.2 | –0.29 | 55–354 | CF ₃ COCF ₃ | 9 |
| (1) CF ₃ + CH ₂ D ₂ = CF ₃ H + CHD ₂ | | | | | | |
| (2) CF ₃ + CH ₂ D ₂ = CF ₃ D + CH ₂ D | | ln k ₁ /k ₂ = ln 1.400 + 0.42 × 10 ⁶ /T ² | | 727–1067 | CF ₃ N ₂ CF ₃ | 28 |

Notes

- (a) CF₃Br inhibition of H₂/O₂ shock tube ignition.
- (b) Values less reliable than others quoted by the same authors.
- (c) Results combined with data from reference 16.
- (d) At low (25mm Hg) CH₃Br pressures.
- (e) At high (180mm Hg) CH₃Br pressures.
- (f) This result is preferred to that obtained with CF₃N₂CF₃.
- (g) "A" factor unreasonable.
- (h) Result described as "semi-quantitative."
- (i) For data involving fluorine atom transfer, see reference 29.

References

1. Ayscough and Polanyi, *Trans. Faraday Soc.* **52**, 960 (1956).
2. Pritchard, Pritchard, Schiff, and Trotman-Dickenson, *Trans. Faraday Soc.* **52**, 849 (1956).
3. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
4. Ayscough, Polanyi, and Steacie, *Can. J. Chem.* **33**, 743 (1955).
5. Dodd and Smith, *J. Chem. Soc.* 1465 (1957).
6. Pritchard, Pritchard, and Trotman-Dickenson, *Chem. and Ind.* 564 (1955).
7. Carmichael and Johnston, *J. Chem. Phys.* **41**, 1975 (1964).

8. Alcock and Whittle, *Trans. Faraday Soc.* **61**, 244 (1965).
9. Sharp and Johnston, *J. Chem. Phys.* **37**, 1541 (1962).
10. Ayscough and Steacie, *Can. J. Chem.* **34**, 103 (1956).
11. Charles and Whittle, *Trans. Faraday Soc.* **52**, 849 (1956).
12. Holmes and Kutschke, *Trans. Faraday Soc.* **58**, 333 (1962).
13. Charles, Pearson, and Whittle, *Trans. Faraday Soc.* **57**, 1356 (1961).
14. Giles and Whittle, *Trans. Faraday Soc.* **62**, 128 (1966).
15. Charles, Pearson, and Whittle, *Trans. Faraday Soc.* **59**, 1156 (1963).
16. Pritchard and Dacey, *Can. J. Chem.* **38**, 182 (1960).
17. Giles and Whittle, *Trans. Faraday Soc.* **61**, 1425 (1965).
18. Sieger and Calvert, *J. Am. Chem. Soc.* **76**, 5197 (1954).
19. Smith and Calvert, *J. Am. Chem. Soc.* **78**, 2345 (1956).
20. Alcock and Whittle, *Trans. Faraday Soc.* **62**, 134 (1966).
21. Alcock and Whittle, *Trans. Faraday Soc.* **62**, 664 (1966).
22. Corbett, Tarr and Whittle, *Trans. Faraday Soc.* **59**, 1609 (1963).
23. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
24. Bell and Kutschke, *Can. J. Chem.* **42**, 2713 (1964).
25. Tucker and Whittle, *Trans. Faraday Soc.* **61**, 866 (1965).
26. Amphlett and Whittle, *Trans. Faraday Soc.* **62**, 1662 (1966).
27. Arthur and Bell, *Can. J. Chem.* **44**, 1445 (1966).
28. Johnston and Tschuikow-Roux, *J. Chem. Phys.* **36**, 463 (1962).
29. Stewart and Cady, *J. Am. Chem. Soc.* **77**, 6110 (1955).

Metathetical Reactions of Halogenated Methyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|--------------------|--|-----------|
| (i) Reactions of CFH ₂ Radicals | | | | | | |
| Hydrogen atom transfer | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| CFH ₂ COCFH ₂ | (a) | 8.0 ± 0.1 | 10.71 | 100–300 | CFH ₂ COCFH ₂ P | 1 |
| (ii) Reactions of CF ₂ Cl Radicals | | | | | | |
| Hydrogen atom transfer | | | | | | |
| cyclo-C ₅ H ₁₀ | (b) | 5.3 ± 0.4 | | 50–227 | CF ₂ ClCOCF ₂ Cl P | 2 |
| Chlorine atom transfer | | | | | | |
| CF ₂ ClCOCF ₂ Cl | (b) | 4 ± 1 | | 20–180 | CF ₂ ClCOCF ₂ Cl P | 3 |
| | (b) | ~ 3 | | 20–184 | CF ₂ ClCOCF ₂ Cl P | 4 |
| (iii) Reactions of CCl ₃ Radicals | | | | | | |
| Hydrogen atom transfer | | | | | | |
| H ₂ | (c) | 11.3 ± 0.5 | 14.86 | 248–302 | CCl ₃ Br P | 5 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (d) | 11.2 | 10.1 | 146–257 | CCl ₃ Br P | 6, 7 |
| CH ₃ CH ₂ CH ₂ CH ₃ | (d) | 7.5 | 9.9 | 146–257 | CCl ₃ Br P | 6, 7 |
| HBr | (d) | | $k = 2.76 \times 10^{11}$ | 190 | CCl ₃ Br P | 6 |
| Chlorine atom transfer | | | | | | |
| Cl ₂ | (d) | 5.3 | 12.86 | 70–155 | CHCl ₃ /Cl ₂ P | 8, 9 |
| CCl ₃ Br | | 18.6 ± 1 17.4 ± 1 | | 225–294 248–302 | CCl ₃ Br P CCl ₃ Br P | 10 5 |
| CCl ₃ COCCL ₃ | | 7 ± 1 | | 107–252 | CCl ₃ COCCL ₃ P | 11 |

Ratios of Rate Constants (Trichloromethyl Radicals)

| Reactants | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|---------------------|-------------------|---------------------------|-----------|
| | | <i>(kcal mole⁻¹)</i> | | °C | | |
| (1) $\text{CCl}_3 + n\text{-C}_4\text{H}_{10}$ = $\text{sec-C}_4\text{H}_9 + \text{CCl}_3\text{H}$ | | -3.08 ± 0.46 | 0.43 ± 0.25 | 197-294 | CCl_3Br P | 7 |
| (2) $\text{CCl}_3 + n\text{-C}_4\text{H}_{10}$ = $p\text{-C}_4\text{H}_9 + \text{CCl}_3\text{H}$ | | | | | | |
| (1) $\text{CCl}_3 + i\text{-C}_4\text{H}_{10}$ = $(\text{CH}_3)_3\text{C} + \text{CCl}_3\text{H}$ | | -2.40 | 0.30 | 132-189 | CCl_3Br P | 7 |
| (2) $\text{CCl}_3 + n\text{-C}_4\text{H}_{10}$ = $\text{sec-C}_4\text{H}_9 + \text{CCl}_3\text{H}$ | | | | | | |

Notes

(a) Calculated assuming $k = 10^{14} \text{ cm}^3 \text{ mole}^{-1} \text{ sec}^{-1}$ for the combination of CFH_2 radicals.

(b) Calculated assuming $k = 10^{14} \text{ cm}^3 \text{ mole}^{-1} \text{ sec}^{-1}$ for the combination of CF_2Cl radicals.

(c) Calculated assuming $k = 10^{10.7} \text{ cm}^3 \text{ mole}^{-1} \text{ sec}^{-1}$ for the combination of CCl_3 radicals.

(d) Calculated assuming $k = 10^{11.80} \text{ cm}^3 \text{ mole}^{-1} \text{ sec}^{-1}$ for the combination of CCl_3 radicals.

References

1. Pritchard, Venugopalan, and Graham, *J. Phys. Chem.* **68**, 1786 (1964).
2. Majer, Phillips, and Robb, *Trans. Faraday Soc.* **61**, 122 (1965).

3. Bowles, Majer, and Robb, *Trans. Faraday Soc.* **58**, 1541 (1962).

4. Bowles, Majer, and Robb, *Trans. Faraday Soc.* **58**, 2394 (1962).

5. Hauteclouque, *Compt. Rend.* **257**, 131 (1963).

6. Tedder and Watson, *Trans. Faraday Soc.* **62**, 1215 (1966).

7. McGrath and Tedder, *Bull. Soc. Chim. Belges* **71**, 772 (1962).

8. Chiltz, Martens, and Mahieu, *Nature* **180**, 1068 (1957).

9. Chiltz, Mahieu, and Martens, *Bull. Soc. Chim. Belges* **67**, 33 (1958).

10. Hauteclouque, *Compt. Rend.* **256**, 2601 (1963).

11. Hauteclouque, *Compt. Rend.* **254**, 3671 (1960).

Review

1. Chiltz, Goldfinger, Huybrechts, Martens, and Verbecke, *Chem. Rev.* **63**, 355 (1963).

Relative Rate Constants (Formyl Radicals)

| Reactants | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | References |
|--|-------|---|---------------------|--|-------------------|---|------------|
| (1) $\text{HCO} + \text{NO}_2 = \text{HNO}_2 + \text{CO}$ (2) $\text{HCO} + \text{NO}_2 = \text{HCO}_2 + \text{NO}$ | | (<i>kcal mole⁻¹</i>) 0.65 | 0.5 | | °C 100–220 | $\text{CH}_2\text{O}/\text{NO}_2$ T | 1, 2 |
| (1) $\text{HCO} + \text{M} = \text{CO} + \text{H} + \text{M}$ (2) $\text{HCO} + \text{O}_2 = \text{CO} + \text{HO}_2$ | | | | 2.7×10^{-9} $\text{cm}^{-3} \text{mole}$ | 36 | $\text{CH}_3\text{COCH}_3/\text{O}_2$ P | 3 |

References

1. Shaw, J. Chem. Soc. 1517 (1964).
2. Pollard and Wyatt, Trans. Faraday Soc. **45**, 760 (1949).
3. Pearson, J. Phys. Chem. **67**, 1686 (1963).

Metathetical Reactions of Methoxy Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | References |
|-------------------------------------|---------|----------------------------|---|-------------------|--------------------------------------|------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Alkanes | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CH ₄ | (a) | 11.0 | 11.8 | 125–250 | | 15 |
| C ₂ H ₆ | (b) | 7.1 | 11.5 | 200–400 | CH ₃ OOCH ₃ T | 1, 2, 3 |
| C ₃ H ₈ | (b) | 5.2 | 11.3 | 200–400 | CH ₃ OOCH ₃ T | 1, 2, 3 |
| n-C ₄ H ₁₀ | (c) (b) | 2.9 | 10.5 | 200–400 | CH ₃ OOCH ₃ T | 1, 2, 3 |
| iso-C ₄ H ₁₀ | (b) | 4.1 | 11.0 | 190–260 | CH ₃ OOCH ₃ T | 1, 2, 3 |
| (CH ₃) ₄ C | (b) | 7.3 | 11.8 | 200–300 | CH ₃ OOCH ₃ T | 1, 2 |
| Cyclo-Alkanes | | | | | | |
| cyclo-C ₃ H ₆ | (b) | 9.7 | 12.2 | 200–400 | CH ₃ OOCH ₃ T | 1, 2 |
| Esters | | | | | | |
| HCOOCH ₃ | | 8.2 | 12.2 | 124–185 | CH ₃ OOCH ₃ T | 4, 5 |
| CH ₃ COOCH ₃ | | ~ 4.5 | | 63–216 | CH ₃ COOCH ₃ P | 6, 14 |
| CH ₃ COOCD ₃ | (d) | ~ 5 | | 30–201 | CH ₃ COOCD ₃ P | 7, 14 |

Ratios of Rate Constants (Methoxy Radicals)

| Reactants | Notes | E ₁ – E ₂ | log ₁₀ A ₁ /A ₂ | Temperature range | Radical source | References |
|---|-------|---|--|-------------------|--|------------|
| | | (kcal mole ⁻¹) | | °C | | |
| (1) CH ₃ O + CH ₃ OH = CH ₃ OH + CH ₂ OH (2) CH ₃ O + HCHO = CH ₃ OH + HCO | | 3.0–4.3 | | 155–180 | CH ₃ OOCH ₃ P | 8, 12 |
| (1) CH ₃ O + HCHO = CH ₃ OH + HCO (2) 2CH ₃ O = CH ₃ OH + HCHO | (f) | $k_1/k_2^{1/2} = 10^{3.68} \exp(-3000/RT)$ | | 50–135 | D.T.B.P. T, P | 9 |
| (1) CH ₃ O + O ₂ = CH ₂ O + HO ₂ (2) 2CH ₃ O = CH ₃ OH + CH ₂ O | (f) | $k_1/k_2^{1/2} = 0.59$ | | room temp. | CH ₃ I/O ₂ P | 10 |
| (1) CH ₃ + CH ₂ (OCH ₃) ₂ = CH ₃ OH + [C ₃ H ₇ O ₂] (2) CH ₃ O = H + CO + H ₂ | | $k_1/k_2 = 1.14 \times 10^5 \text{ cm}^3 \text{ mole}^{-1}$ | | 459 | CH ₂ (OCH ₃) ₂ T | 11 |

Notes

- (a) Calculated from the back reaction.
- (b) The A factors were deduced by a semi-empirical method by Berces and Trotman-Dickenson (ref. 1). The original work contained an arithmetical mistake that resulted in an overestimate of $\log A$ of 0.3 units.
- (c) This value seems likely to be incorrect.
- (d) The reactant radical in this case is CD_3O .
- (e) For further data on methoxy radicals see reference 13.
- (f) Units are $\text{cm}^{3/2} \text{mole}^{-1/2} \text{sec}^{-1/2}$.

References

1. Berces and Trotman-Dickenson, *J. Chem. Soc.* 348 (1961).
2. Shaw and Trotman-Dickenson, *J. Chem. Soc.* 3210 (1960).
3. Shaw and Trotman-Dickenson, *Proc. Chem. Soc.* 61 (1959).
4. Thynne and Gray, *Trans. Faraday Soc.* **59**, 1149 (1963).

5. Thynne and Gray, *Proc. Chem. Soc.* 295 (1962).
6. Wijnen, *J. Chem. Phys.* **27**, 710 (1957).
7. Wijnen, *J. Chem. Phys.* **28**, 939 (1958).
8. Takezaki and Takeuchi, *J. Chem. Phys.* **22**, 1527 (1954).
9. Hoare and Wellington, *8th. Int. Comb. Symp.*, p. 472 (Williams and Wilkins, 1962).
10. Heicklen and Johnston, *J. Am. Chem. Soc.* **84**, 4030 (1962).
11. Molera, Fernandez-Biarge, Centeno, and Arevalo, *J. Chem. Soc.* 2311 (1963).
12. Takezaki, Miyazaki, and Nakahara, *J. Chem. Phys.* **25**, 536 (1956).
13. Dever and Calvert, *J. Am. Chem. Soc.* **84**, 1362 (1962).
14. Wijnen, *J. Chem. Phys.* **28**, 271 (1958).
15. Shaw and Thynne, *Trans. Faraday Soc.* **62**, 104 (1966).

Reviews

1. Gray and Williams, *Chem. Rev.* **59**, 239 (1959).

Metathetical Reactions of Methylthio Radicals

| Reactants | Note | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|------------------------|------|------------------------------------|---|--------------------|--|------------------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| CH ₄ | (a) | (kcal mole ⁻¹) 18.2 | (cm ³ mole ⁻¹ sec ⁻¹) 12.1 | °C 130-200 | | 1 |
| CH ₃ CHO | | 6.8 4.8 | 11.9 12.0 | 189-396 400-440 | CH ₃ CHO/CH ₃ SH CH ₃ CHO/CH ₃ SH | P T 2 3 |

Notes

(a) Calculated from the reverse reactions.

References

1. Greig and Thynne, *Trans. Faraday Soc.* **62**, 379 (1966).
2. Birrell, Smith, Trotman-Dickenson, and Wilkie, *J. Chem. Soc.* 2807 (1957).
3. Imai and Toyama, *Bull. Chem. Soc. Japan* **33**, 1408 (1960).

Reactions of Chloroformyl Radicals

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|--|-------|--|---|-------------------|--|------------|
| $\text{COCl} + \text{Cl}_2 = \text{COCl}_2 + \text{Cl}$ | (a) | (<i>kcal mole</i> ⁻¹) 2.96 | (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) 12.4 | °C 25-55 | CO/Cl ₂ P | 1 5 |
| | | 3.1 | 10.8 | 15-450 | | |
| $\text{COCl} + \text{NOCl} = \text{Cl}_2 + \text{CO} + \text{NO}$ (or $\text{COCl}_2 + \text{NO}$) | | 1.14 | 13.68 | 25-55 | CO/Cl ₂ /NOCl P | 2 |
| $\text{COCl} + \text{O}_2 = \text{CO}_2 + \text{ClO}$ | | 3.3 | 10.9 | 20-200 | Cl ₂ /O ₂ /CO P | 3, 4 |

Notes

(a) Review of literature data.

References

1. Burns and Dainton, *Trans. Faraday Soc.* **48**, 39 (1952).
2. Burns and Dainton, *Trans. Faraday Soc.* **48**, 52 (1952).
3. Rollefson, *J. Am. Chem. Soc.* **55**, 148 (1933).
4. Trotman-Dickenson, "Gas Kinetics," p. 259 et seq. (Butterworths, 1955).
5. Bodenstein, Brenschede, and Schumacher, *Z. Phys. Chem.* **40B**, 121 (1938).

Metathetical Reactions of Cyano Radicals

| Reactions | Notes | E | $\log_{10} A$ | Temperature range | Radical source | References |
|---------------------------------|-------|--|--|--------------------------|--|------------|
| $H_2 + CN = HCN + H$ | | (<i>kcal mole⁻¹</i>) ~ 7 | (<i>cm³ mole⁻¹ sec⁻¹</i>) | °C | D.F. | 1 |
| $ClCN + CN = C_2N_2 + Cl$ | | 6.0 | 13.05 | 1727-2527 | ClCN ... S.T. | 2 |
| $C_2N_2 + CN = \text{products}$ | | 2.1 | 10.86 | 28-174 | C_2N_2 F.P. | 3 |
| $O_2 + CN = NCO + O$ | | | $k = 4.6 \times 10^{12}$ $k = 5.5 \times 10^{12}$ | room temp. room temp. | C_2N_2/O_2 F.P. C_2N_2/O_2 F.P. | 5 3 |

Ratios of Rate Constants

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | References |
|--|-------|--|---------------------|-----------|-------------------|----------------|------------|
| (1) $CH_4 + CN = CH_3 + HCN$ (2) $C_2H_6 + CN = C_2H_5 + HCN$ | | (<i>kcal mole⁻¹</i>) 3.7 ± 0.2 | 1.31 | | °C 30-150 | ICN P | 4 |
| (1) $C_2H_6 + CN = C_2H_5 + HCN$ (2) $C_3H_8 + CN = n-C_3H_7 + HCN$ | | -0.1 ± 0.2 | 0.00 | | 30-150 | ICN P | 4 |
| (1) $C_2H_6 + CN = C_2H_5 + HCN$ (2) $C_3H_8 + CN = i-C_3H_7 + HCN$ | | -0.1 ± 0.3 | 0.04 | | 30-150 | ICN P | 4 |
| (1) $C_3H_8 + CN = n-C_3H_7 + HCN$ (2) $C_3H_8 + CN = i-C_3H_7 + HCN$ | | -0.1 ± 0.3 | 0.08 | | 30-150 | ICN P | 4 |

References

1. Hartel and Polanyi, *Z. Phys. Chem.* **B11**, 97 (1930).
2. Schofield, Tsang, and Bauer, *J. Chem. Phys.* **42**, 2132 (1965).
3. Paul and Dalby, *J. Chem. Phys.* **37**, 592 (1962).
4. Goy, Shaw, and Pritchard, *J. Phys. Chem.* **69**, 1504 (1965).
5. Basco, *Proc. Roy. Soc.* **283A**, 302 (1965).

Ratios of Rate Constants (Ethyne Radicals)

(i) Reactions of the type (1) $C_2H + RH = C_2H_2 + R$

(2) $C_2H + BrC_2H = C_4H_2 + Br.$

| RH | Notes | k_1/k_2 | Temperature | Radical source | References |
|--|-------|-----------|-------------|----------------------|------------|
| Alkanes | | | | | |
| | | | °C | | |
| CH ₄ | | 0.020 | 27 | BrC ₂ H P | 1 |
| C ₂ H ₆ | | 0.40 | 27 | BrC ₂ H P | 1 |
| <i>n</i> -C ₄ H ₁₀ | | 1.05 | 27 | BrC ₂ H P | 1 |
| iso-C ₄ H ₁₀ | | 1.25 | 27 | BrC ₂ H P | 1 |
| (CH ₃) ₄ C | | 1.1 | 27 | BrC ₂ H P | 1 |
| (CH ₃) ₃ C.C(CH ₃) ₃ | | 1.2 | 27 | BrC ₂ H P | 1 |
| Cyclo-alkanes | | | | | |
| cyclo-C ₃ H ₆ | | 0.21 | 27 | BrC ₂ H P | 1 |
| cyclo-C ₄ H ₈ | | 1.35 | 27 | BrC ₂ H P | 1 |
| spiro-C ₅ H ₈ | | 0.64 | 27 | BrC ₂ H P | 1 |
| cyclo-C ₅ H ₁₀ | | 2.1 | 27 | BrC ₂ H P | 1 |
| cyclo-C ₆ H ₁₂ | | 3.2 | 27 | BrC ₂ H P | 1 |
| cyclo-C ₆ D ₁₂ | | 2.3 | 27 | BrC ₂ H P | 1 |
| Halogenated alkanes | | | | | |
| C ₂ H ₅ Cl | | 0.14 | 27 | BrC ₂ H P | 1 |

(ii) Reactions of the type (1) $C_2H + RH = C_2H_2 + R$

(2) $C_2H + CH_2:CHCH_2CH_3 = C_2H_2 + C_4H_7$

| RH | Notes | k_1/k_2 | Temperature | Radical source | Reference |
|--|-------|-----------|-------------|----------------------|-----------|
| Alkenes | | | | | |
| | | | °C | | |
| CH ₂ :CHCH ₃ | | 0.63 | 27 | BrC ₂ H P | 2 |
| cis-CH ₃ CH:CHCH ₃ | | 1.04 | 27 | BrC ₂ H P | 2 |
| trans-CH ₃ CH:CHCH ₃ | | 1.06 | 27 | BrC ₂ H P | 2 |
| CH ₂ :C(CH ₃) ₂ | | 0.90 | 27 | BrC ₂ H P | 2 |
| CH ₂ :CHCH ₂ CH ₂ CH ₃ | | 1.7 | 27 | BrC ₂ H P | 2 |
| CH ₂ :CHCH(CH ₃) ₂ | | 1.22 | 27 | BrC ₂ H P | 2 |

References

1. Tarr, Strausz, and Gunning, *Trans. Faraday Soc.* **61**, 1946 (1965).

2. Tarr, Strausz, and Gunning, *Trans. Faraday Soc.* **62**, 1221 (1966).

Metathetical Reactions of Ethyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|----------------------------|---|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| H ₂ | (a) | 11.5 ± 1 | | | | 1 |
| | | 4.05 | 12.19 | 752-917 | H ₂ /O ₂ ignition | S.T. |
| | (b) | 11.3 ± 0.5 | 11.51 | 83-322 | (C ₂ D ₅) ₂ CO | P |
| | | 15.9 | 12.8 | | | 3 |
| D ₂ | | 13.3 ± 0.5 | 12.0 | 54-287 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 1 |
| Alkanes | | | | | | |
| <i>n</i> -C ₄ H ₁₀ | | 15.2 | 11.77 | 420-530 | <i>n</i> -C ₄ H ₁₀ | T |
| | (b) | 10.4 | 11.03 | 106-325 | (C ₂ D ₅) ₂ CO | P |
| | | | | | | 4 |
| <i>i</i> -C ₄ H ₁₀ | (b) | 8.9 | 10.62 | 87-319 | (C ₂ D ₅) ₂ CO | P |
| | | | | | | 5 |
| neo-C ₅ H ₁₂ | (b) | 12.6 | 11.25 | 86-324 | (C ₂ D ₅) ₂ CO | P |
| | | | | | | 5 |
| cyclo-C ₆ H ₁₂ | (b) | 10.4 ± 0.5 | 11.42 ± 0.2 | 99-293 | (C ₂ D ₅) ₂ CO | P |
| | | | | | | 3 |
| <i>n</i> -C ₆ H ₁₄ | (b) | 10.1 ± 0.5 | 11.14 ± 0.2 | 87-252 | (C ₂ D ₅) ₂ CO | P |
| | | | | | | 3 |
| <i>n</i> -C ₇ H ₁₆ | | 10.6 ± 0.4 | 11.69 ± 0.2 | 124-200 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 6 |
| Alkenes | | | | | | |
| 1,5-C ₆ H ₁₀ | | 6.3 ± 0.4 | 10.1 ± 0.2 | 71-175 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 7 |
| 1-C ₇ H ₁₄ | | 8.3 ± 0.5 | 11.19 ± 0.3 | 85-200 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 6 |
| 1-C ₈ H ₁₆ | | 8.3 ± 0.2 | 11.19 ± 0.3 | 85-180 | (C ₂ H ₅) ₂ CO | P |
| | | 7.5 ± 0.5 | 10.6 ± 0.3 | 66-152 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 6 |
| | | | | | | 9 |
| trans 4-C ₈ H ₁₆ | | 8.7 ± 1.0 | 11.5 ± 0.6 | 85-165 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 6 |
| (CH ₃) ₂ C:CHCH: C(CH ₃) ₂ | | 7.6 ± 0.4 | 11.1 ± 0.2 | 53-147 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 9 |
| Cyclo alkenes | | | | | | |
| cyclohexadiene,-1,3. | | 5.4 ± 0.5 | 10.6 ± 0.3 | 25-160 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 9, 8 |
| cyclohexadiene,-1,4. | | 5.8 ± 0.1 | 11.3 ± 0.1 | 50-140 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 10 |
| cyclohexene | | 8.2 ± 0.5 | 11.5 ± 0.3 | 23-250 | (C ₂ H ₅) ₂ CO | P |
| | | 7.5 ± 0.4 | 10.2 ± 0.2 | 40-210 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 6 |
| | | | | | | 10 |
| cycloheptatriene | | 6.5 ± 0.5 | 10.9 ± 0.3 | 50-130 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 10 |
| cyclooctatetraene | | 8.6 ± 1.2 | 11.6 ± 0.7 | 50-130 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 10 |
| cyclooctadiene,-1,5. | | 6.8 ± 0.7 | 10.9 ± 0.4 | 60-130 | (C ₂ H ₅) ₂ CO | P |
| | | | | | | 10 |

Metathetical Reactions of Ethyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|----------------------------|---|-------------------|--|-----------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| Alkynes | | | | | | |
| <i>i</i> -C ₇ H ₁₂ | | 7.6 ± 0.2 | 10.9 ± 0.1 | 23–180 | (C ₂ H ₅) ₂ CO P | 6 |
| Aldehydes | | | | | | |
| C ₂ H ₅ CHO | | 7.6 ± 1.0 | 11.5 | 134–156 | D.T.B.P. T | 11 |
| | | 5.9 | 10.8 | 91–315 | C ₂ H ₅ CHO P | 12 |
| | | 6.3 | | 100–175 | C ₂ H ₅ CHO P | 12, 13 |
| Ketones | | | | | | |
| CH ₃ COC ₂ H ₅ | | 8.0 ± 0.1 | 11.0 | 79–234 | CH ₃ COC ₂ H ₅ P | 16 |
| (C ₂ H ₅) ₂ CO | | 7.4 | 10.9 | 25–225 | (C ₂ H ₅) ₂ CO P | 14, 21 |
| | | 7.6 | 11.0 | 41–158 | (C ₂ H ₅) ₂ N ₂ P | 15 |
| | | 7.8 ± 0.2 | 11.1 ± 0.1 | 50–215 | (C ₂ H ₅) ₂ CO P | 6 |
| | | 8.9 | 11.7 | 75–170 | (C ₂ H ₅) ₂ CO P | 17 |
| (CH ₃ CD ₂) ₂ CO | (c) | 11.7 | 11.4 | 24–365 | (CH ₃ CD ₂) ₂ CO P | 18 |
| (CH ₃ CD ₂) ₂ CO | (c) | 8.7 | 11.6 | 24–365 | (CH ₃ CD ₂) ₂ CO P | 18, 19 |
| | | 9.2 ± 0.4 | 11.1 ± 0.1 | 24–365 | (CH ₃ CD ₂) ₂ CO P | 19 |
| (C ₂ D ₅) ₂ CO | (b) | 9.6 ± 0.4 | 11.3 | 160–314 | (C ₂ D ₅) ₂ CO P | 19 |
| | | 9.0 ± 0.5 | 10.95 | 50–324 | (C ₂ D ₅) ₂ CO P | 5 |
| C ₂ F ₅ COC ₂ H ₅ | | 6.8 | 10.85 | 100–250 | C ₂ F ₅ COC ₂ H ₅ P | 37 |
| C ₃ F ₇ COC ₂ H ₅ | | 7.2 | 10.9 | 80–362 | C ₃ F ₇ COC ₂ H ₅ P | 20 |
| Esters | | | | | | |
| HCOOC ₂ H ₅ | | 7.8 | 10.6 | 77–230 | CH ₃ COCH ₃ P | 17 |
| C ₂ H ₅ COOC ₂ H ₅ | | 9.8 | 11.5 | 108–344 | C ₂ H ₅ COOC ₂ H ₅ P | 22 |
| C ₂ H ₅ COOCH ₂ CH: CH ₂ | | 5.8 ± 1.4 | 9.7 ± 0.8 | 79–160 | (C ₂ H ₅) ₂ CO P | 23 |
| Azo-compounds | | | | | | |
| (C ₂ H ₅) ₂ N ₂ | | 7.5 | 10.9 | 74–178 | (C ₂ H ₅) ₂ N ₂ P | 24 |
| | | 8.0 ± 0.2 | 11.4 | 27–175 | (C ₂ H ₅) ₂ N ₂ P | 25 |
| Metal alkyl | | | | | | |
| (C ₂ H ₅) ₂ Hg | | 6.2 | 10.3 | 75–200 | (C ₂ H ₅) ₂ Hg P | 26 |
| Hydrogen halide | | | | | | |
| HI | | 1.1 | 11.92 | 263–303 | HI/C ₂ H ₅ I T | 27 |

Metathetical Reactions of Ethyl Radicals – Continued

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|------------------------|-------|----------------------------|---|-------------------|---|-----------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| CHLORINE ATOM TRANSFER | | | | | | |
| Cl ₂ | | 1.0 | 13.1 | | | 28 |
| IODINE ATOM TRANSFER | | | | | | |
| I ₂ | (d) | 0.2 | 12.50 | 263–303 | HI/C ₂ H ₅ I T | 27 |
| OXYGEN ATOM TRANSFER | | | | | | |
| N ₂ O | (e) | 31.0 | 17.8 | 553–588 | C ₂ H ₆ /N ₂ O T | 29 |

Ratios of Rate Constants (Ethyl Radicals)

| Reaction | Notes | <i>E</i> ₁ – <i>E</i> ₂ | $\log_{10} A_1/A_2$ | <i>k</i> ₁ / <i>k</i> ₂ | Temperature range | Radical source | Reference |
|---|-------|---|---------------------|---|---------------------------|--|------------------------|
| | | (kcal mole ⁻¹) | | | °C | | |
| (1) C ₂ H ₅ + HBr = C ₂ H ₆ + Br (2) C ₂ H ₅ + Br ₂ = C ₂ H ₅ Br + Br | | 0 | –0.22 | | 30–90 | C ₂ H ₆ /Br ₂ /HBr P | 30, 36 |
| (1) C ₂ H ₅ + HI = C ₂ H ₆ + I (2) C ₂ H ₅ + I ₂ = C ₂ H ₅ I + I | | 0.90 | –0.58 | 0.15 0.13 ± 0.03 | 260 250–280 263–303 | C ₂ H ₅ I/HI T C ₂ H ₅ I/HI T C ₂ H ₅ I/HI T | 33, 32 31, 32 27 |
| (1) C ₂ H ₅ + HBr = C ₂ H ₆ + Br (2) C ₂ H ₅ + I ₂ = C ₂ H ₅ I + I | | 2.29 ± 0.08 | 0.23 ± 0.04 | | 55–115 | CH ₃ COC ₂ H ₅ P | 34 |
| (1) C ₂ H ₅ + I ₂ = C ₂ H ₅ I + I (2) C ₂ H ₅ + O ₂ = C ₂ H ₅ O ₂ | | | | 13 | 25 | C ₂ H ₅ I/O ₂ P | 35 |
| (1) C ₂ H ₅ + I ₂ = C ₂ H ₅ I + I (2) C ₂ H ₅ + NO = C ₂ H ₅ NO | | | | 7 | 25 | C ₂ H ₅ I/NO P | 35 |

Notes

(a) Estimated from the reaction D₂ + C₂H₅, assuming the difference in activation energies to be equal to the difference of the zero-point energies.

(b) The attacking radical is C₂D₅.

(c) The attacking radical is CH₃CD₂.

(d) Calculated, assuming the activation energy of 0.2 kcal/mole.

(e) This *A* factor seems improbably high.

References

1. Wijnen and Steacie, *J. Chem. Phys.* **20**, 205 (1952).
2. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
3. Boddy and Steacie, *Can. J. Chem.* **39**, 13 (1961).
4. Purnell and Quinn, *Proc. Roy. Soc.* **270A**, 267 (1962).
5. Boddy and Steacie, *Can. J. Chem.* **38**, 1576 (1960).
6. James and Steacie, *Proc. Roy. Soc.* **244A**, 289 (1958).
7. James and Troughton, *Trans. Faraday Soc.* **62**, 145 (1966).

8. Brown and James, Proc. Chem. Soc. 81 (1962).
9. Brown and James, Can. J. Chem. **43**, 1102 (1965).
10. Brown and James, Can. J. Chem. **43**, 660 (1965).
11. Volman and Brinton, J. Chem. Phys. **22**, 929 (1954).
12. Kerr and Trotman-Dickenson, J. Chem. Soc. 1611 (1960).
13. Blacet and Pitts, J. Am. Chem. Soc. **74**, 3382 (1952).
14. Kutschke, Wijnen, and Steacie, J. Am. Chem. Soc. **74**, 714 (1952).
15. Ausloos and Steacie, Can. J. Chem. **32**, 593 (1954).
16. Ausloos and Steacie, Can. J. Chem. **33**, 1062 (1955).
17. Thynne, Trans. Faraday Soc. **58**, 676 (1962).
18. Wijnen and Steacie, Can. J. Chem. **29**, 1092 (1951).
19. James and Steacie, Proc. Roy. Soc. **245A**, 470 (1958).
20. Pritchard and Thommarson, J. Phys. Chem. **69**, 1001 (1965).
21. Brinton and Steacie, Can. J. Chem. **33**, 1840 (1955).
22. Wijnen, J. Am. Chem. Soc. **80**, 2394 (1958).
23. James and Troughton, Trans. Faraday Soc. **62**, 120 (1966).
24. Ausloos and Steacie, Bull. Soc. Chim. Belges. **63**, 87 (1954).
25. Cerfontain and Kutschke, Can. J. Chem. **36**, 344 (1958).
26. Ivin and Steacie, Proc. Roy. Soc. **208A**, 25 (1951).
27. Hartley and Benson, J. Chem. Phys. **39**, 132 (1963).
28. Goldfinger, Huybrechts, Martens, Meyers, and Olbrechts, Trans. Faraday Soc. **61**, 1933 (1965).
29. Kenright and Trenwith, J. Chem. Soc. 2079 (1959).
30. Anderson and Van Artsdalen, J. Chem. Phys. **12**, 479 (1944).
31. Sullivan, J. Phys. Chem. **65**, 722 (1961).
32. Ogg, J. Am. Chem. Soc. **56**, 526 (1934).
33. Benson and O'Neal, J. Chem. Phys. **34**, 514 (1961).
34. Fettis and Trotman-Dickenson, J. Chem. Soc. 3037 (1961).
35. Christie and Frost, Trans. Faraday Soc. **61**, 468 (1965).
36. Benson and Buss, J. Chem. Phys. **28**, 301 (1958).
37. Thommarson and Pritchard, J. Phys. Chem. **70**, 2307 (1966).

Perfluoroethyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| | | <i>(kcal mole⁻¹)</i> | <i>(mole⁻¹ cc sec⁻¹)</i> | °C | | |
| H ₂ | (a) | 11.9 | 12.72 | 137-237 | (C ₂ F ₅) ₂ CO | 1 |
| | | 12.4 ± 0.2 | 13.20 | 146-313 | C ₂ F ₅ CHO | 2 |
| D ₂ | (a) | 12.6 ± 0.2 | 12.45 | 135-339 | C ₂ F ₅ CHO | 2 |
| | | 14.1 ± 0.3 | 13.08 | 154-259 | (C ₂ F ₅) ₂ N ₂ | 3 |
| CH ₄ | | 10.6 | 11.62 | 150-272 | (C ₂ F ₅) ₂ CO | 1 |
| C ₂ H ₆ | | 8.7 ± 0.2 | 12.18 | 84-226 | (C ₂ F ₅) ₂ N ₂ | 3 |
| cyclo-C ₆ H ₁₂ | | 6.0 ± 0.2 | 12.18 | 28-132 | (C ₂ F ₅) ₂ N ₂ | 3 |
| CF ₃ CHO | | 9.7 ± 0.2 | 12.40 | 138-220 | (C ₂ F ₅) ₂ N ₂ | 3 |
| C ₂ F ₅ CHO | | 4.5 ± 0.2 | 10.49 | 27-307 | C ₂ F ₅ CHO | 4 |
| | | 4.9 ± 0.2 | 10.74 | 135-339 | C ₂ F ₅ CHO | 2 |
| CH ₃ COCH ₃ | | 8.4 ± 0.2 | 11.71 | 82-220 | (C ₂ F ₅) ₂ N ₂ | 3 |
| C ₂ F ₅ COC ₂ H ₅ | | 5.6 ± 0.2 | 11.34 | 50-250 | C ₂ F ₅ COC ₂ H ₅ | 5 |

Notes

(a) These results are less reliable than those obtained from perfluoro ketone systems.

References

1. Price and Kutschke, *Can. J. Chem.* **38**, 2128 (1960).
2. Pritchard and Foote, *J. Phys. Chem.* **68**, 1016 (1964).
3. Pritchard, Dacey, Kent, and Simonds, *Can. J. Chem.* **44**, 171 (1966).
4. Pritchard, Miller, and Foote, *Can. J. Chem.* **40**, 1830 (1962).
5. Thommarson and Pritchard, *J. Phys. Chem.* **70**, 2307 (1966).

Metathetical Reactions of Halogenated Ethyl Radicals

| Reactions | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|-------|--------------------------------|--|--------------------|--|-----------|
| $\text{CH}_2\text{CH}_2\text{Cl} + \text{Cl}_2$ $= \text{CH}_2\text{ClCH}_2\text{Cl} + \text{Cl}$ | | (kcal mole^{-1}) 0 | $(\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1})$ | $^{\circ}\text{C}$ | $\text{CH}_2:\text{CH}_2/\text{Cl}_2$ P | 1, 2 |
| $\text{C}_2\text{H}_3\text{Cl}_2 + \text{Cl}_2$ $= \text{CH}_2\text{ClCHCl}_2 + \text{Cl}$ | (c) | 0.92 ± 0.05 | 11.75 ± 0.2 | 25-55 | $\text{CH}_2:\text{CHCl}/\text{Cl}_2$ P | 1, 2, 3 |
| $\text{CHCl}_2\text{CHCl} + \text{Cl}_2$ $= \text{CHCl}_2\text{CHCl}_2 + \text{Cl}$ | | 2.74 ± 0.6 | 11.7 ± 0.3 | 30-65 | <i>cis</i> - $\text{CHCl}:\text{CHCl}/\text{Cl}_2$ P | 4, 3 |
| $\text{C}_2\text{HCl}_4 + \text{Cl}_2$ $= \text{CHCl}_2\text{CCl}_3 + \text{Cl}$ | (b) | 5.1 ± 0.2 | 11.5 ± 0.2 | 80-140 | $\text{CHCl}:\text{CCl}_2/\text{Cl}_2$ P | 5, 3, 12 |
| $\text{C}_2\text{Cl}_5 + \text{Cl}_2 = \text{C}_2\text{Cl}_6 + \text{Cl}$ | (a) | 5.5 | 11.3 | 87-247 | $\text{CCl}_2:\text{CCl}_2/\text{Cl}_2$ P | 6, 7 |
| $\text{CF}_2\text{CF}_2\text{Cl} + \text{Cl}_2$ $= \text{CF}_2\text{ClCF}_2\text{Cl} + \text{Cl}$ | (d) | 0.8 | | 30-60 | $\text{CF}_2:\text{CF}_2/\text{Cl}_2$ P | 13 |
| $\text{C}_2\text{F}_3\text{Cl}_2 + \text{Cl}_2 = \text{C}_2\text{F}_3\text{Cl}_3$ $+ \text{Cl}$ | (e) | 2.3 | | 30-60 | $\text{CF}_2:\text{CFCl}/\text{Cl}_2$ P | 13 |
| $\text{CFCl}_2\text{CFCl} + \text{Cl}_2$ $= \text{CFCl}_2\text{CFCl}_2 + \text{Cl}$ | (f) | 5.35 ± 0.3 | | 30-70 | $\text{CFCl}:\text{CFCl}/\text{Cl}_2$ P | 14 |
| $\text{C}_2\text{F}_4\text{Br} + \text{HBr} = \text{C}_2\text{F}_4\text{Br}_2$ $+ \text{H}$ | | 51.2 | 13.07 | 855-1013 | $\text{H}_2/\text{O}_2/\text{C}_2\text{F}_4\text{Br}_2$ S.T. | 15 |

Ratios of Rate Constants (Halogenated Ethyl Radicals)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | References |
|---|-------|---|---------------------|-----------|-----------------------------|---|------------|
| (1) $\text{C}_2\text{H}_4\text{Br} = \text{C}_2\text{H}_4 + \text{Br}$ (2) $\text{C}_2\text{H}_4\text{Br} + \text{HBr} = \text{C}_2\text{H}_5\text{Br} + \text{Br}$ | | (kcal mole^{-1}) 14 ± 2 | | | $^{\circ}\text{C}$ 25-54 | $\text{C}_2\text{H}_4/\text{HBr}$ γ radiation | 16, 18 |
| (1) $\text{C}_2\text{H}_4\text{Br} = \text{C}_2\text{H}_4 + \text{Br}$ (2) $\text{C}_2\text{H}_4\text{Br} + \text{Br}_2 = \text{C}_2\text{H}_4\text{Br}_2 + \text{Br}$ | | 6.1 | | | 60-80 | $\text{C}_2\text{H}_4/\text{Br}_2$ P | 17 |

Notes

- (a) Data on this reaction can also be found in references 8 to 11.
- (b) The reactant radical is probably $\text{CHCl}_2\text{CCl}_2^{\cdot}$.
- (c) The reactant radical is probably $\text{CH}_2\text{ClCHCl}_2^{\cdot}$.
- (d) Calculated assuming zero activation energy for the combination of $\text{CF}_2\text{CF}_2\text{Cl}$ radicals.
- (e) Calculated assuming zero activation energy for the combination of $\text{C}_2\text{F}_3\text{Cl}_2$ radicals.
- (f) Calculated assuming zero activation energy for the combination of CFCl_2CFCl radicals.

References

1. Dainton, Lomax, and Weston, *Trans. Faraday Soc.* **58**, 308 (1962).
2. Ayscough, Cocker, Dainton, and Hirst, *Trans. Faraday Soc.* **58**, 318 (1962).
3. Ayscough, Cocker, Dainton, Hirst, Lomax, and Weston, *Proc. Chem. Soc.* 244 (1961).
4. Ayscough, Cocker, Dainton, and Hirst, *Trans. Faraday Soc.* **58**, 295 (1962).
5. Dainton, Lomax, and Weston, *Trans. Faraday Soc.* **53**, 460 (1957).
6. Dusoleil, Goldfinger, Maheiu-Van der Auwera, Martens, and Van der Auwera, *Trans. Faraday Soc.* **57**, 2197 (1961).
7. Adam, Goldfinger, and Gosselain, *Bull. Soc. Chim. Belges* **65**, 549 (1956).
8. Adam and Goldfinger, *Bull. Soc. Chim. Belges* **65**, 561 (1956).
9. Adam, Dusoleil, and Goldfinger, *Bull. Soc. Chim. Belges* **65**, 942 (1956).
10. Ackerman, Chiltz, Goldfinger, and Martens, *Bull. Soc. Chim. Belges* **66**, 325 (1957).
11. Chiltz, Maheiu, and Martens, *Bull. Soc. Chim. Belges* **67**, 33 (1958).
12. Chiltz, Dusoleil, Goldfinger, Huybrechts, Mahieu, Martens, and Van der Auwera, *Bull. Soc. Chim. Belges* **68**, 5 (1959).
13. Castellano, Bergamin, and Schumacher, *Z. Phys. Chem.* **27**, 112 (1961).
14. Vallana, Castellano, and Schumacher, *Z. Phys. Chem.* **46**, 294 (1965).
15. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
16. Armstrong and Spinks, *Can. J. Chem.* **37**, 1210 (1959).
17. Schmitz, Schumacher, and Jager, *Z. Phys. Chem.* **51B**, 281 (1942).
18. Oldershaw and Cvetanovic, *J. Chem. Phys.* **41**, 3639 (1964).

Review

1. Chiltz, Goldfinger, Huybrechts, Martens, and Verbecke, *Chem. Rev.* **63**, 355 (1963).

Ratios of Rate Constants (Carbonylcarbene Radicals)

Ratios of the form (1) $\text{CCO} + \text{C}_3\text{O}_2 = \text{Polymer} + n\text{CO}$

(2) $\text{CCO} + \text{C}_n\text{H}_{2n} = \text{C}_{n+1}\text{H}_{2n} + \text{CO}$

Where C_nH_{2n} is an olefine, and $\text{C}_{n+1}\text{H}_{2n}$ is the diene and/or alkyne formed by addition of a carbon atom

| Olefine | Notes | k_1/k_2 | Temperature | Radical source | Reference |
|---|-------|-----------|-------------|--------------------------|-----------|
| C_2H_4 | (a) | 1.4 | 0 | C_3O_2 P | 1 |
| | (a) | 1.26 | 0 | C_3O_2 P | 2 |
| | (b) | 2.79 | 0 | C_3O_2 P | 2 |
| $\text{CH}_3\text{CH}:\text{CH}_2$ | (a) | 3.93 | 0 | C_3O_2 P | 2 |
| | (b) | 9.72 | 0 | C_3O_2 P | 2 |
| $\text{CH}_2:\text{CHCH}_2\text{CH}_3$ | (a) | 13.14 | 0 | C_3O_2 P | 2 |
| | (b) | 38.91 | 0 | C_3O_2 P | 2 |
| $\text{CH}_2:\text{C}(\text{CH}_3)_2$ | (a) | 14.04 | 0 | C_3O_2 P | 2 |
| | (a) | 8.03 | 0 | C_3O_2 P | 2 |
| $\text{cis-CH}_3\text{CH}:\text{CHCH}_3$ | (a) | 8.03 | 0 | C_3O_2 P | 2 |
| $\text{trans-CH}_3\text{CH}:\text{CHCH}_3$ | (a) | 14.33 | 0 | C_3O_2 P | 2 |
| $(\text{CH}_3)_2\text{C}:\text{CHCH}_3$ | (a) | 34.38 | 0 | C_3O_2 P | 2 |
| $(\text{CH}_3)_2\text{C}:\text{C}(\text{CH}_3)_2$ | (a) | 67.20 | 0 | C_3O_2 P | 2 |

Notes

- (a) Photolysis at 2537 Å.
 (b) Photolysis at > 3100 Å.

References

1. Bayes, J. Am. Chem. Soc. **84**, 4077 (1962).
2. Baker, Kerr, and Trotman-Dickenson, J. Chem. Soc. 975A (1966).

Metathetical Reactions of Acetyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|-----------------------------------|-------|---|---|-------------------|-----------------------------|-----------|
| $I_2 + CH_3CO$ $= CH_3COI + I$ | | 0 (<i>kcal mole</i> ⁻¹) | 12.6 (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) | °C 222-268 | CH ₃ COI/HI T | 1 |

Ratios of Rate Constants (Acetyl and Trifluoroacetyl Radicals)

| Reaction | Notes | <i>E</i> ₁ - <i>E</i> ₂ | $\log_{10}A_1/A_2$ | <i>k</i> ₁ / <i>k</i> ₂ | Temperature range | Radical source | Reference |
|--|-------|---|------------------------------------|---|-------------------|--|-----------|
| (1) CH ₃ CO = CH ₃ + CO (2) CH ₃ CO + HBr = CH ₃ CHO + Br | | (kcal mole ⁻¹) ~ 13 | | | °C 100-300 | CH ₃ COCH ₃ P | 2 |
| (1) CH ₃ CO = CH ₃ + CO (2) CH ₃ CO + O ₂ = CH ₃ O + CO ₂ | | ~ 9 | - 4.68 (mole cm ⁻³) | | | CH ₃ COCH ₃ /O ₂ P | 3 |
| (1) CH ₃ CO + HI = CH ₃ CHO + I (2) CH ₃ CO + I ₂ = CH ₃ COI + I | | 1.5 | - 0.47 | | 222-268 | CH ₃ COI/HI T | 1 |
| (1) CF ₃ CO = CF ₃ + CO (2) CF ₃ CO + Br ₂ = CF ₃ COBr + Br | (a) | 6.0 | - 4.7 | | 19-251 | CF ₃ COCF ₃ /Br ₂ P | 4 |

Notes

(a) Values very doubtful.

References

1. O'Neal and Benson, *J. Chem. Phys.* **37**, 540 (1962).
2. Ridge and Steacie, *Can. J. Chem.* **33**, 383 (1955).
3. Cerfontain and Kutschke, *J. Am. Chem. Soc.* **84**, 4017 (1962).
4. Tucker and Whittle, *Trans. Faraday Soc.* **61**, 484 (1965).

Ratios of Rate Constants (Ethoxy Radicals)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|---------------------|----------------|-------------------|-----------------------|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) $C_2H_5O + (C_2H_5)_2CO$ = C_2H_5OH + $C_2H_4COC_2H_5$ | | | | | | | |
| (2) $C_2H_5O + O_2$ = $CH_3CHO + HO_2$ | | | | 0.1 ± 0.05 | 35 | $(C_2H_5)_2CO/O_2$ P | 1 |
| (1) $C_2H_5O + (C_2H_5)_2N_2$ = C_2H_5OH + $C_2H_4N_2C_2H_5$ | | | | 0.6 ± 0.3 | 118 | $(C_2H_5)_2N_2/O_2$ P | 2 |
| (2) $C_2H_5O + O_2$ = $CH_3CHO + HO_2$ | | | | 1.1 ± 0.4 | 152 | $(C_2H_5)_2N_2/O_2$ P | 2 |
| (1) $C_2H_5O = CH_3 + CH_2O$ | | | | | | | |
| (2) $C_2H_5O + C_2H_5COOC_2H_5$ = $C_2H_5OH + C_5H_9O_2$ | | 7.5 ± 1 | | | 29-195 | $C_2H_5COOC_2H_5$ P | 3 |

NOTE: For further data on ethoxy radicals see reference 4.

3. Wijnen, J. Am. Chem. Soc. **82**, 3034 (1960).

4. Heicklen and Johnston, J. Am. Chem. Soc. **84**, 4394 (1962).

References

Review

1. Jolley, J. Am. Chem. Soc. **79**, 1537 (1957).
2. Cerfontain and Kutschke, J. Am. Chem. Soc. **84**, 4017 (1962).

1. Gray and Williams, Chem. Rev. **59**, 239 (1959).

Metathetical Reactions of Allyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|--|-------|---|--|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| cyclo-C ₅ H ₁₀ | | <i>(kcal mole⁻¹)</i> 31.8 ± 3.6 | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C 450–514 | CD ₃ COCD ₃ P | 1 |
| C ₆ H ₅ CH ₃ | (a) | 14 to 17 | | 459–592 | CH ₂ :CHCH ₂ Br T | 2 |
| CH ₂ :CHCH ₂ CH(CH ₃) ₂ | | 12 | | 450–530 | CH ₂ :CHCH ₂ CH (CH ₃) ₂ T | 3 |

Note

(a) Calculated assuming a steric factor in the range 10⁻¹ to 10⁻².

References

1. Gordon, Smith, and McNesby, *J. Am. Chem. Soc.* **81**, 5059 (1959).
2. Szwarc, Ghosh, and Sehon, *J. Chem. Phys.* **18**, 1142 (1950).
3. Taniowski, *J. Chem. Soc.* 7436 (1965).

Metathetical Reactions of *n*-Propyl Radicals ^(a)

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|---|-------|-----------------------------------|---|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| <i>n</i> -C ₃ H ₇ CHO | | (kcal mole ⁻¹) 6.7 | (cm ³ mole ⁻¹ sec ⁻¹) 11.0 | °C 98-361 | <i>n</i> -C ₃ H ₇ CHO P | 1 |
| <i>n</i> -C ₃ H ₇ CHO | | 10.8 | 11.0 | 191-300 | <i>n</i> -C ₃ H ₇ CHO P | 1 |
| (<i>n</i> -C ₃ H ₇) ₂ CO | | 6.5 | 10.4 | 55-161 | (<i>n</i> -C ₃ H ₇) ₂ CO P | 2 |
| HCOOCH ₂ CH ₂ CH ₃ | | 7.6 | 10.9 | 74-178 | CH ₃ COCH ₃ P | 3 |
| (<i>n</i> -C ₃ H ₇) ₂ N ₂ | | 7.9 | 11.3 | 25-291 | (<i>n</i> -C ₃ H ₇) ₂ N ₂ P | 4 |

Ratios of Rate Constants (*n*-Propyl Radicals)

| Reactions | Notes | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|--|-------|--------------------------------|-------------------|--|-----------|
| (1) <i>n</i> -C ₃ H ₇ + HI = C ₃ H ₈ + I | | 0.11 | 290 | <i>n</i> -C ₃ H ₇ I/HI | T |
| (2) <i>n</i> -C ₃ H ₇ + I ₂ = C ₃ H ₇ I + I | | | | | |
| (1) <i>n</i> -C ₃ H ₇ + I ₂ = C ₃ H ₇ I + I | | 11 | 25 | <i>n</i> -C ₃ H ₇ I/NO | P |
| (2) <i>n</i> -C ₃ H ₇ + NO = C ₃ H ₇ NO | | | | | |
| (1) <i>n</i> -C ₃ H ₇ + I ₂ = C ₃ H ₇ I + I | | 22 | 25 | <i>n</i> -C ₃ H ₇ I/O ₂ | P |
| (2) <i>n</i> -C ₃ H ₇ + O ₂ = C ₃ H ₇ O ₂ | | | | | |

Note

(a) The rate constants are based on log *k* (cm³ mole⁻¹ sec⁻¹) = 10^{13.4} for the combination of *n*-propyl radicals.

References

1. Kerr and Trotman-Dickenson, *Trans. Faraday Soc.* **55**, 921 (1959).
2. Masson, *J. Am. Chem. Soc.* **74**, 4731 (1952).
3. Thynne, *Trans. Faraday Soc.* **58**, 1394 (1962).
4. Kerr and Calvert, *J. Am. Chem. Soc.* **83**, 3391 (1961).
5. Benson and O'Neal, *J. Chem. Phys.* **34**, 514 (1961).
6. Ogg, *J. Am. Chem. Soc.* **56**, 526 (1934).
7. Christie and Frost, *Trans. Faraday Soc.* **61**, 468 (1965).

Metathetical Reactions of Isopropyl Radicals ^(a)

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H ₂ | | 12.5 | | 260-320 | CH ₃ CH:CH ₂ + H | 1 |
| Alkenes | | | | | | |
| cyclohexadiene-1,4 | | 6.5 | 11.4 | 75-136 | (<i>i</i> -C ₃ H ₇) ₂ CO P | 2 |
| cyclohexadiene-1,3 | | 7.1 ± 0.7 | 11.5 ± 0.4 | 42-133 | (<i>i</i> -C ₃ H ₇) ₂ CO P | 3 |
| Aldehydes and Ketones | | | | | | |
| <i>i</i> -C ₃ H ₇ CHO | | 9.5 | 10.7 | 283-377 | <i>i</i> -C ₃ H ₇ CHO P | 4 |
| <i>i</i> -C ₃ H ₇ CHO | | 6.3 | 10.8 | 117-354 | <i>i</i> -C ₃ H ₇ CHO P | 4 |
| (<i>i</i> -C ₃ H ₇) ₂ CO | | 8.5 ± 0.1 | 11.1 | 100-400 | (<i>i</i> -C ₃ H ₇) ₂ CO P | 5 |
| [(CH ₃) ₂ CD] ₂ CO | (b) | 9.3 ± 0.3 | 10.6 | 200-400 | [(CH ₃) ₂ CD] ₂ CO P | 6 |
| [(CH ₃) ₂ CD] ₂ CO | (b) | 11.7 ± 1.1 | 11.3 | 300-400 | [(CH ₃) ₂ CD] ₂ CO P | 6 |
| Ester | | | | | | |
| HCOOCH(CH ₃) ₂ | | 6.6 | 9.9 | 94-181 | CH ₃ COCH ₃ P | 7 |
| Azo-compound | | | | | | |
| (<i>i</i> -C ₃ H ₇) ₂ N ₂ | | 6.5 ± 0.5 | 9.9 | 30-120 | (<i>i</i> -C ₃ H ₇) ₂ N ₂ P | 8 |
| | | 6.7 ± 0.4 | 10.0 | 35-127 | (<i>i</i> -C ₃ H ₇) ₂ N ₂ P | 9 |

Ratios of Rate Constants (Isopropyl Radicals)

| Reactions | Notes | E ₁ - E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|--------------------------------|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) <i>i</i> -C ₃ H ₇ + I ₂ = C ₃ H ₇ I + I (2) <i>i</i> -C ₃ H ₇ + NO = C ₃ H ₇ NO | | | | 22 | 25 | <i>i</i> -C ₃ H ₇ I/NO P | 10 |
| (1) <i>i</i> -C ₃ H ₇ + I ₂ = C ₃ H ₇ I + I (2) <i>i</i> -C ₃ H ₇ + O ₂ = C ₃ H ₇ O ₂ | | | | ~ 3 | 25 | <i>i</i> -C ₃ H ₇ I/O ₂ P | 10 |
| (1) <i>i</i> -C ₃ H ₇ + <i>i</i> -C ₃ H ₇ I = C ₃ H ₆ + C ₃ H ₈ + I | (c) | | | 520(3130Å) | 35 | <i>i</i> -C ₃ H ₇ I P | 11 |
| (2) <i>i</i> -C ₃ H ₇ + I ₂ = <i>i</i> -C ₃ H ₇ I + I | (c) | | | 170(2300Å) | 35 | <i>i</i> -C ₃ H ₇ I P | 11 |
| (1) C ₃ H ₇ + O ₂ = C ₃ H ₆ + HO ₂ (2) C ₃ H ₇ + O ₂ = C ₃ H ₇ O ₂ | (d) | 19.0 | 6.6 | | 345-472 | C ₃ H ₈ /O ₂ T | 12 |

Notes

(a) The rate constants are based on $\log k$ ($\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1}$) = $10^{12.9}$ for the combination of isopropyl radicals.

(b) The attacking radical is $(\text{CH}_3)_2\text{CD}$.

(c) The difference in values is due to a hot radical effect. The value at 3130 Å approximates to the correct value for thermally equilibrated isopropyl radicals.

(d) Summary of a large body of literature data. The propyl radical is probably a mixture of *n* and isopropyl.

References

1. Hoey and Le Roy, *Can. J. Chem.* **33**, 580 (1955).
2. James and Suart, *J. Am. Chem. Soc.* **86**, 5424 (1964).
3. James and Suart, *J. Phys. Chem.* **69**, 2362 (1965).
4. Kerr and Trotman-Dickenson, *Trans. Faraday Soc.* **55**, 921 (1959).
5. Heller and Gordon, *J. Phys. Chem.* **60**, 1315 (1956).
6. Heller and Gordon, *J. Phys. Chem.* **62**, 709 (1958).
7. Thynne, *Trans. Faraday Soc.* **58**, 1394 (1962).
8. Durham and Steacie, *Can. J. Chem.* **31**, 377 (1953).
9. Riem and Kutschke, *Can. J. Chem.* **38**, 2332 (1960).
10. Christie and Frost, *Trans. Faraday Soc.* **61**, 468 (1965).
11. McMillan and Noyes, *J. Am. Chem. Soc.* **80**, 2108 (1958).
12. Sattersfield and Reid, *J. Phys. Chem.* **59**, 283 (1955).

Perfluoropropyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|------------------------------------|---|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| | | (<i>kcal mole</i> ⁻¹) | (<i>mole</i> ⁻¹ <i>cm</i> ³ <i>sec</i> ⁻¹) | °C | | |
| H ₂ | (a) | 12.3 ± 0.4 | 12.64 | 125-243 | (<i>n</i> -C ₃ F ₇) ₂ CO | 1 |
| | | 12.1 ± 0.2 | 12.86 | 157-319 | C ₃ F ₇ CHO | 2 |
| D ₂ | (b) | 13.8 ± 0.5 | 12.78 | 85-182 | (<i>n</i> -C ₃ F ₇) ₂ CO | 1 |
| | | 12.9 ± 0.8 | 12.69 | | (<i>n</i> -C ₃ F ₇) ₂ CO | 3 |
| | (a) | 14.0 ± 0.1 | 12.98 | 165-297 | C ₃ F ₇ CHO | 2 |
| CH ₄ | | 9.5 ± 0.5 | 10.99 | 70-166 | (<i>n</i> -C ₃ F ₇) ₂ CO | 3 |
| C ₂ H ₆ | (c) | 9.2 ± 0.5 | 12.24 | 87-196 | (<i>n</i> -C ₃ F ₇) ₂ CO | 3 |
| cyclo-C ₆ H ₁₂ | | 5.2 ± 0.1 | 11.08 | 25-290 | (<i>n</i> -C ₃ F ₇) ₂ CO | 4 |
| C ₃ F ₇ CHO | (c) | 4.0 ± 0.3 | 10.27 | 28-315 | C ₃ F ₇ CHO | 5 |
| | | 5.5 ± 0.2 | 10.98 | 165-297 | C ₃ F ₇ CHO | 2 |
| CH ₃ COCH ₃ | | 7.2 ± 0.4 | 11.83 | 27-306 | C ₃ F ₇ CHO | 6 |
| C ₃ F ₇ COC ₂ H ₅ | | 8.4 ± 0.3 | 11.77 | 80-362 | C ₃ F ₇ COC ₂ H ₅ | 7 |

Notes

(a) These results are less reliable than those obtained from perfluoro ketone systems.

(b) Assuming $k = 10^{10.99} \exp(-9500/RT)$ for the reaction $n\text{-C}_3\text{F}_7 + \text{CH}_4 = \text{C}_3\text{F}_7\text{H} + \text{CH}_3$.

(c) For both these compounds it seems likely that errors have been made in the determination of the activation energies, although the rate constants were probably of the correct magnitude.

References

1. Miller and Steacie, *J. Am. Chem. Soc.* **80**, 6486 (1958).
2. Pritchard and Foote, *J. Phys. Chem.* **68**, 1016 (1964).
3. Giacometti and Steacie, *Can. J. Chem.* **36**, 1493 (1958).
4. Pritchard and Miller, *J. Phys. Chem.* **63**, 2074 (1959).
5. Pritchard, Miller, and Foote, *Can. J. Chem.* **40**, 1830 (1962).
6. Pritchard, Hsia, and Miller, *J. Am. Chem. Soc.* **85**, 1568 (1963).
7. Pritchard and Thommarson, *J. Phys. Chem.* **69**, 1001 (1965).

Metathetical Reactions of Halogenated Propyl Radicals

| Reactions | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|-------|--|---|-------------------|--|-----------|
| $\text{CCl}_3\text{CH}_2\text{CH}_2 + \text{CCl}_3\text{Br}$ $= \text{CCl}_3\text{CH}_2\text{CH}_2\text{Br}$ $+ \text{CCl}_3$ | | 3.4 <i>(kcal mole⁻¹)</i> | 8 <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C 103-193 | $\text{C}_2\text{H}_4/\text{CCl}_3\text{Br}$ P | 1 |
| $\text{ICH}_2\text{CH}_2\text{CH}_2 + \text{I}_2$ $= \text{ICH}_2\text{CH}_2\text{CH}_2\text{I} + \text{I}$ | (a) | 0.5 | 12.5 | | cyclo- $\text{C}_3\text{H}_6/\text{I}_2$ T | 2 |

Note

(a) Estimated from the equilibrium constant and collision theory.

References

1. Tedder and Walton, *Trans. Faraday Soc.* **60**, 1769 (1964).
2. Benson, *J. Chem. Phys.* **34**, 521 (1961).

Ratios of Rate Constants (Isopropoxy Radicals)

| Reaction | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|---------------------|------------|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) $i\text{-C}_3\text{H}_7\text{O} = \text{CH}_3\text{CHO} + \text{CH}_3$ (2) $i\text{-C}_3\text{H}_7\text{O} + (\text{CH}_3)_2\text{CHOCC}_2\text{H}_5 = i\text{-C}_3\text{H}_7\text{OH} + \text{R}$ | | < 6 | | | 28 | $\text{C}_2\text{H}_5\text{COO}\dot{\text{C}}(\text{CH}_3)_2$ P | 1 |
| (1) $i\text{-C}_3\text{H}_7\text{O} = \text{CH}_3\text{CHO} + \text{CH}_3$ (2) $i\text{-C}_3\text{H}_7\text{O} + \text{NO} = \text{CH}_3\text{COCH}_3 + \text{HNO}$ | | 16 | | | 175-200 | $(\text{CH}_3)_2\text{CHONO}$ T | 2 |
| (1) $i\text{-C}_3\text{H}_7\text{O} + \text{CH}_3\text{CHO} = i\text{-C}_3\text{H}_7\text{OH} + \text{CH}_3\text{CO}$ (2) $i\text{-C}_3\text{H}_7\text{O} + (i\text{-C}_3\text{H}_7\text{O})_2 = i\text{-C}_3\text{H}_7\text{OH} + \text{R}$ | | | | 27 ± 4 | 26 | $(i\text{-C}_3\text{H}_7\text{O})_2$ P | 3 |

References

1. Wijnen, J. Am. Chem. Soc. **82**, 1847 (1960).
2. Ferguson and Phillips, J. Chem. Soc. 4416 (1965).
3. McMillan, J. Am. Chem. Soc. **83**, 3018 (1961).

Review

1. Gray and Williams, Chem. Rev. **59**, 239 (1959).

Metathetical Reactions of Butyl Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|--|-------|----------------------------|---|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| (i) n-butyl radicals | (a) | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| <i>n</i> -C ₄ H ₉ CHO | | 5.4 | 10.9 | 61-229 | <i>n</i> -C ₄ H ₉ CHO P | 1 |
| HCOOCH ₂ CH ₂ CH ₂ CH ₃ | | 5.3 | 10.2 | 75-186 | CH ₃ COCH ₃ P | 2 |
| (ii) sec-butyl radicals | (a) | | | | | |
| CH ₃ CH ₂ CH (CH ₃)CHO | | 4.9 | 10.7 | 25-349 | CH ₃ CH ₂ CH (CH ₃)CHO P | 3 |
| (iii) iso-butyl radicals | (a) | | | | | |
| <i>i</i> -C ₄ H ₉ CHO | | 12.7 ± 0.2 | 12.62 ± 0.05 | 178-279 | <i>i</i> -C ₄ H ₉ CHO P | 4 |
| <i>i</i> -C ₄ H ₉ CHO | | 6.5 ± 0.1 | 11.71 ± 0.07 | 117-230 | <i>i</i> -C ₄ H ₉ CHO P | 4 |
| (<i>i</i> -C ₄ H ₉) ₂ CO | | 7.6 | 11.4 | 78-194 | (<i>i</i> -C ₄ H ₉) ₂ CO P | 5 |
| | (c) | 6.8 ± 0.2 | 11.06 ± 0.09 | | | 4 |
| (iv) t-butyl radicals | (b) | | | | | |
| (CH ₃) ₃ CCHO | | 10.0 | 11.2 | 240-386 | (CH ₃) ₃ CCHO P | 6 |
| (CH ₃) ₃ CCHO | (d) | 4.3 | 9.8 | 60-386 | (CH ₃) ₃ CCHO P | 6 |

Ratios of Rate Constants (*t*-Butyl Radical)

| Reaction | Notes | E ₁ - E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|--|--------------------------------|-------------------|--|-----------|
| | | (kcal mole ⁻¹) | | | °C | | |
| (1) <i>t</i> -C ₄ H ₉ + HBr = C ₄ H ₁₀ + Br (2) <i>t</i> -C ₄ H ₉ + Br ₂ = C ₄ H ₉ Br + Br | | 8.0 | 5.11 | | 40-85 | (CH ₃) ₃ CH/Br ₂ T | 7, 8, 9 |
| (1) <i>t</i> -C ₄ H ₉ + HI = C ₄ H ₁₀ + I (2) <i>t</i> -C ₄ H ₉ + I ₂ = C ₄ H ₉ I + I | | | | 0.226 0.234 0.251 | 526 552 583 | (CH ₃) ₃ CH/I ₂ T | 8 |

Notes

- (a) The rate constants are based on log *k* (cm³ mole⁻¹ sec⁻¹) = 10¹⁴ for the combination of *n*-, *sec*-, and *iso*-butyl radicals.
 (b) The rate constants are based on log *k* (cm³ mole⁻¹ sec⁻¹) = 10^{12.5} for the combination of *t*-butyl radicals.
 (c) Recalculation of data from reference 5.
 (d) This value of the activation energy is probably low.

References

- Kerr and Trotman-Dickenson, *J. Chem. Soc.* 1602 (1960).
- Thynne, *Trans. Faraday Soc.* **58**, 1533 (1962).
- Gruver and Calvert, *J. Am. Chem. Soc.* **78**, 5208 (1956).
- Metcalfe and Trotman-Dickenson, *J. Chem. Soc.* 5072 (1960).
- Kraus and Calvert, *J. Am. Chem. Soc.* **79**, 5921 (1957).
- Birrell and Trotman-Dickenson, *J. Chem. Soc.* 4218 (1960).
- Eckstein, Scheraga, and Van Artsdalen, *J. Phys. Chem.* **22**, 28 (1954).
- Teranishi and Benson, *J. Am. Chem. Soc.* **85**, 2887 (1963).
- Benson and Buss, *J. Chem. Phys.* **28**, 301 (1958).

Metathetical Reactions of *t*-Butoxy Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|------------------------------------|-------|-----------------------------------|---|-------------------|----------------|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| (CH ₃) ₃ CH | | 4.0 (kcal mole ⁻¹) | 10.8 (cm ³ mole ⁻¹ sec ⁻¹) | °C 25-79 | D.T.B.P. P | 4 |

Ratios of Rate Constants (*t*-Butoxy Radicals)

| Reaction | Notes | E ₁ - E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|--|-------------------|----------------|-----------|
| (1) (CH ₃) ₃ CO + CH ₂ O = (CH ₃) ₃ COH + HCO | | (kcal mole ⁻¹) | | | °C | | |
| (2) (CH ₃) ₃ CO = CH ₃ + CH ₃ COCH ₃ | | | | (3.8 ± 0.8) × 10 ⁵ cm ³ mole ⁻¹ | 135 | D.T.B.P. T | 1 |
| (1) (CH ₃) ₃ CO + ((CH ₃) ₃ CO) ₂ = (CH ₃) ₃ COH + R | | | | | | | |
| (2) (CH ₃) ₃ CO = CH ₃ + CH ₃ COCH ₃ | | -3 | | | 25-79 | D.T.B.P. P | 2 |
| (1) (CH ₃) ₃ CO + (CH ₂) ₂ NH = (CH ₃) ₃ COH + R | | | | | | | |
| (2) (CH ₃) ₃ CO = CH ₃ + CH ₃ COCH ₃ | | -12 ± 2 | | | 129-154 | D.T.B.P. P | 3 |

References

1. Hoare and Wellington, 8th Int. Comb. Symp., p. 472 (Williams and Wilkins 1962).
2. McMillan and Wijnen, Can. J. Chem. **36**, 1227 (1958).
3. Brinton and Volman, J. Chem. Phys. **20**, 25 (1952).
4. McMillan, J. Am. Chem. Soc. **82**, 2422 (1960).

Review

1. Gray and Williams, Chem. Rev. **59**, 239 (1959).

Metathetical Reactions of Peracid Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|--|-------|---|---|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| (i) Peracetic Radicals CH ₃ CHO | | (kcal mole ⁻¹) k = (8.05 ± 2.40) × 10 ⁶ | (cm ³ mole ⁻¹ sec ⁻¹) | °C 20 | CH ₃ CHO/O ₂ P | 1 |
| (ii) Perpropionic Radicals C ₂ H ₅ CHO | | k = (4.35 ± 0.91) × 10 ⁷ | | 22 | C ₂ H ₅ CHO/O ₂ P | 1 |

Ratio of Rate Constants (Peracid Radicals)

| Reaction | Notes | E ₁ - E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|---|-------|---|--|--------------------------------|-------------------|--|-----------|
| (i) Peracetic Radicals | | (kcal mole ⁻¹) | | | °C | | |
| (1) CH ₃ CO ₃ + CH ₃ CHO = CH ₃ CO ₃ H + CH ₃ CO | | E ₁ - ½E ₂ = 7.2 ± 1.0 | | | 20-30 | CH ₃ CHO/O ₂ P | 2 |
| (2) 2CH ₃ CO ₃ = (CH ₃ CO) ₂ O ₂ + O ₂ | | | | | | | |
| (ii) Perpropionic Radicals | | | | | | | |
| (1) C ₂ H ₅ CO ₃ + C ₂ H ₅ CHO = C ₂ H ₅ CO ₃ H + C ₂ H ₅ CO | | E ₁ - ½E ₂ = 6.75 ± 0.5 | | | 20-47 | C ₂ H ₅ CHO/O ₂ P | 3 |
| (2) 2C ₂ H ₅ CO ₃ = (C ₂ H ₅ CO) ₂ O ₂ + O ₂ | | | | | | | |

References

1. McDowell and Sharples, Can. J. Chem. **36**, 268 (1958).
2. McDowell and Sharples, Can. J. Chem. **36**, 251 (1958).
3. McDowell and Sharples, Can. J. Chem. **36**, 258 (1958).

Metathetical Reactions of Peroxy Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|---------|---------------------------------|--|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| (i) Isobutyl peroxy radicals | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| (CH ₃) ₃ CH | (a) (b) | 16.0 | $4 \times 10^{-2}Z$ | 327-427 | (CH ₃) ₃ CH/O ₂ - H | 1 |
| (ii) Cyclohexenyl peroxy radicals | | | | | | |
| cyclohexene | (a) (b) | 7.0 | $6 \times 10^{-7}Z$ | 152-352 | C ₆ H ₁₀ /O ₂ - H | 1 |
| (iii) Isopropyl benzene peroxy radicals | | | | | | |
| C ₆ H ₅ CH(CH ₃) ₂ | (a) (b) | 7.0 | $2 \times 10^{-6}Z$ | 252-357 | C ₆ H ₅ CH(CH ₃) ₂ - H | 1 |

Notes

- (a) The peroxy radical is the radical formed by oxygen addition to any hydrocarbon radical produced in the primary act.
 (b) "Z" is the collision number.

Reference

1. Burgess and Robb, *Trans. Faraday Soc.* **54**, 1015 (1958).

Metathetical Reactions of Phenyl Radicals

| Reactants | Notes | <i>E</i> | $\log_{10}A^{(a)}$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H ₂ | | 6.5 | 10.97 | 180-350 | (C ₆ H ₅) ₂ Hg P | 1 |
| CH ₄ | | 7.5 | 11.19 | 180-350 | (C ₆ H ₅) ₂ Hg P | 1 |
| | | 11.1 | 11.9 | 277-407 | CH ₃ COC ₆ H ₅ P | 2 |
| cyclo-C ₃ H ₆ | | 8.5 | 11.4 | 310-407 | CH ₃ COC ₆ H ₅ P | 2 |
| <i>i</i> -C ₄ H ₁₀ | | 6.7 | 11.8 | 277-407 | CH ₃ COC ₆ H ₅ P | 2 |
| CH ₃ COC ₆ H ₅ | | 6.2 | 11.6 | 277-407 | CH ₃ COC ₆ H ₅ P | 2 |
| CF ₃ H | | 5.2 | 10.17 | 180-350 | (C ₆ H ₅) ₂ Hg P | 1 |
| GROUP TRANSFER REACTION | | | | | | |
| C ₆ H ₅ + C ₆ H ₅ COCH ₃ = C ₆ H ₅ C ₆ H ₅ + CH ₃ CO | | 6.2 | 9.6 | 277-407 | CH ₃ COC ₆ H ₅ P | 2 |

Note

(a) These values are based on $\log k$ (cm³ mole⁻¹ sec⁻¹) = 10¹⁴ for the combination of phenyl radicals.

References

1. Fielding and Pritchard, *J. Phys. Chem.* **66**, 821 (1962).
2. Duncan and Trotman-Dickenson, *J. Chem. Soc.* 4672 (1962).

Ratios of Rate Constants (Tolyl Radicals)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|---------------------|-----------|-------------------|--|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) $\text{CH}_3\text{C}_6\text{H}_4 + \text{C}_6\text{H}_4(\text{CH}_3)_2$ $= \text{CH}_3\text{C}_6\text{H}_5$ $+ \text{CH}_3\text{C}_6\text{H}_4\text{CH}_2$ | (a) | | | 10.7 | 700 | $(\text{CH}_3)_2\text{C}_6\text{H}_4/\text{D}_2$ | T |
| (2) $\text{CH}_3\text{C}_6\text{H}_4 + \text{D}_2$ $= \text{CH}_3\text{C}_6\text{H}_4\text{D} + \text{D}$ | | | | | | | |
| (1) $p\text{-CH}_3\text{C}_6\text{H}_4 + \text{C}_6\text{H}_4(\text{CH}_3)_2$ $= \text{CH}_3\text{C}_6\text{H}_5$ $+ \text{CH}_3\text{C}_6\text{H}_4\text{CH}_2$ | (b) | | | 24.6 | 484 | $p\text{-(CH}_3)_2\text{C}_6\text{H}_4/\text{D}_2$ | T |
| (2) $p\text{-CH}_3\text{C}_6\text{H}_4 + \text{D}_2$ $= \text{CH}_3\text{C}_6\text{H}_4\text{D} + \text{D}$ | | | | | | | |

Notes

References

- (a) No distinction was made between the *o*, *m*, and *p* xylenes.
 (b) There appears to be a misprint in the original paper. The numbering of the reactions has been reversed.

1. Burr and Strong, J. Am. Chem. Soc. **86**, 5065 (1964).
2. Burr and Strong, J. Chem. Phys. **43**, 1432 (1965).

Hydroxyl Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|----------------------------------|---------|----------------------------|---|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Hydrogen | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| H ₂ | | 11.0 | 14.52 | 105-216 | H ₂ O/ discharge | 1 |
| | (a) | 10.0 | $k = 2.0 \times 10^{10}$ 14.1 | 520 | H ₂ /O ₂ ignition | 2 3 |
| | | 10.0 | 14.40 | | H ₂ O ₂ flames | 4 |
| | (b) | 5.9 ± 1.0 | $k = 4.3 \times 10^9$ 13.80 ± 0.7 | 37 | NO ₂ /H reaction | 5 |
| | | | | 27-1700 | | 6 |
| | (b) (c) | 5.0(5.3) ± 0.6 | $k = (3.5 \pm 0.3) \times 10^9$ 13.13(13.33) | 27 | H ₂ / discharge | 7 |
| | (b) (c) | 5.3(5.5) ± 0.6 | 13.33(13.49) | 27-799 | | 8 |
| | | | $k = 1.1 \times 10^{12}$ | 27-799 | | 9 |
| | | | $k = (3.9 \pm 0.2) \times 10^9$ | 642 | H ₂ /O ₂ flame | 10 |
| | (b) | 5.2 | 13.36 | 27 | NO ₂ /H reaction | 11 |
| | | 6.4 | 14.17 | 27-1677 | | 11 |
| | | | | 687-807 | H ₂ /O ₂ ignition | 12 |
| Alkanes | | | | | | |
| CH ₄ | | 8.3 | 14.38 | | H ₂ O/ discharge | 13 |
| | | 9.0 | 14.54 | 1027-1527 | CH ₄ /O ₂ flame | 14 |
| | | | $k = 2 \times 10^{13}$ | 1377-1567 | CH ₄ /O ₂ flame | 15 |
| | (d) | 6.5 | 14.15 | 930-1530 | CH ₄ /O ₂ flame | 16 |
| | | 7.9 | 14.36 | | | 17 |
| C ₂ H ₂ | | | $k = 2 \times 10^{12}$ | 1400-1700 | C ₂ H ₂ /O ₂ flame | 18 |
| | | 7.3 | 14.44 | 50-237 | H ₂ O/ discharge | 19 |
| C ₂ H ₄ | | | $k = 1 \times 10^{13}$ | 977-1127 | C ₂ H ₄ /O ₂ flame | 20 |
| | | 6.4 | 14.60 | 77-178 | H ₂ O/ discharge | 19 |
| C ₂ H ₆ | | 5.5 | 14.11 | 65-228 | H ₂ O/ discharge | 19 |
| | (e) | | $k = 2.5 \times 10^{13}$ | 1147-1337 | C ₂ H ₆ /H ₂ /O ₂ flame | 34 |
| | | | $k = 2.1 \times 10^{13}$ | 520 | H ₂ /O ₂ ignition | 21 |
| | | | $k = 5 \times 10^{12}$ | 1027-1227 | C ₂ H ₆ /O ₂ flame | 20 |
| C ₃ H ₈ | (e) | | $k = 4.3 \times 10^{13}$ | 520 | H ₂ /O ₂ ignition | 21 |
| n-C ₄ H ₁₀ | | | $k = 5.8 \times 10^{13}$ | 520 | H ₂ /O ₂ ignition | 21 |
| i-C ₄ H ₁₀ | | | $k = 3.2 \times 10^{13}$ | 520 | H ₂ /O ₂ ignition | 21 |
| Aldehydes | | | | | | |
| HCHO | (b) | 0.9 | 14.11 | 73-216 | H ₂ O/ discharge | 22 |
| | | 13.0 | 15.7 | 73-1339 | | 20 |
| CH ₃ CHO | | 4.0 | 13.53 | 53-209 | H ₂ O/ discharge | 22 |
| Acid | | | | | | |
| HNO ₃ | | | $k = 1.0 \times 10^{11}$ | 27 | HNO ₃ flash photolysis | 23 |
| Halogenated Alkane | | | | | | |
| CH ₃ Br | | | $k = 1.5 \times 10^{13}$ | 1527-1727 | CH ₄ /O ₂ flame | 24 |

Hydroxyl Radicals – Continued

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|-----------|-------|----------------------------|---|-------------------|----------------|-----------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |

OXYGEN ATOM TRANSFER

| | | | | | | |
|-----|---------|--------------------|-------------------------------------|-----------|---------------------------------------|----|
| CO | (g) | 7.0 | 13.08 | 70–203 | H ₂ O/ discharge | 1 |
| | (a) | 5.7 | 12.46 | | | 3 |
| | | | $k = 9 \times 10^{11}$ | 1677 | CH ₄ /O ₂ flame | 15 |
| | (d) | 5.7 | 12.46 | | | 17 |
| | (b) | 7.7 | 12.85 | 127–1727 | | 20 |
| | | 4.0 | 12 | 1107–1447 | flame study. | 25 |
| | (b) | 6.2 ± 0.6 | 12.6 ± 0.3 | 107–1677 | | 8 |
| | (b) (c) | $0.5(0.8) \pm 0.6$ | $11.5(11.7) \pm 0.3$ | 200–800 | | 8 |
| | | | $k = (1.15 \pm 0.5) \times 10^{11}$ | 27 | NO ₂ /H reaction | 11 |
| (b) | 0.6 | 11.49 | 27–1677 | | 11 | |

MISCELLANEOUS REACTIONS

| | | | | | | |
|----------------------------------|--|------|-------|---------|---|----|
| OH + F ₂ = HF + F + O | | 18.0 | 15.3 | | F ₂ /H ₂ O flames | 26 |
| | | 18.0 | 13.85 | 497–557 | F ₂ /H ₂ O ignition | 26 |

Hydroxyl Radicals (Rate Constant Ratios)

| Reaction | Notes | E ₁ – E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|--|-------|---------------------------------|--|--------------------------------|-------------------|--|-----------|
| | | (kcal mole ⁻¹) | | | °C | | |
| (1) C ₂ H ₆ + OH = C ₂ H ₅ + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 12 | 540 | H ₂ /O ₂ ignition | 27 |
| (1) C ₃ H ₈ + OH = C ₃ H ₇ + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 27 | 520 | H ₂ /O ₂ ignition | 27 |
| (1) n-C ₄ H ₁₀ + OH = C ₄ H ₉ + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 36 | 520 | H ₂ /O ₂ ignition | 27 |
| (1) i-C ₄ H ₁₀ + OH = C ₄ H ₉ + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 20 | 520 | H ₂ /O ₂ ignition | 27 |
| (1) HCHO + OH = HCO + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 42 | 540 | H ₂ /O ₂ ignition | 27 |
| (1) (C ₂ H ₅) ₄ Si + OH = (C ₂ H ₅) ₃ Si.C ₂ H ₅ + H ₂ O (2) H ₂ + OH = H + H ₂ O | (f) | | | 74 | 520 | H ₂ /O ₂ ignition | 27 |
| (1) H ₂ O ₂ + OH = HO ₂ + H ₂ O (2) H ₂ + OH = H ₂ O + H | | | | 4.8 to 5.7 | 447 | H ₂ /O ₂ ignition | 28 |
| | | | | 7.1 | 500 | H ₂ /O ₂ ignition | 29 |
| | | | | 5.5 | 440 | H ₂ /O ₂ ignition | 30 |
| | | | | 4.3 ± 0.3 | 440 | H ₂ /O ₂ pyrolysis | 27 |
| | | | | 4.7 | 500 | H ₂ combustion | 27 |

Hydroxyl Radicals (Rate Constant Ratios) – Continued

| Reaction | Notes | $E_1 - E_2$ (<i>kcal mole⁻¹</i>) | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range °C | Radical source | Reference |
|---|-------|--|---------------------|----------------------|--------------------------|---|----------------------|
| (1) $H_2 + OH = H + H_2O$ (2) $CO + OH = CO_2 + H$ | | 4.0 ± 0.3 | 1.49 ± 0.7 | 5.0 3.3 | 200–350 520 | H ₂ O photolysis H ₂ /O ₂ ignition CO/H ₂ combustion | 31 27 27 |
| | (b) | 5.16 | 1.98 | | 500–1002 | | 27 |
| | (b) | 4.6 ± 0.3 | 1.87 | | 27–1002 | | 11 |
| (1) $D_2 + OH = D + HDO$ (2) $CO + OH = CO_2 + H$ | | 6.4 ± 0.3 | 2.20 ± 0.09 | | 200–300 | H ₂ O photolysis | 31 |
| (1) $CH_4 + OH = CH_3 + H_2O$ (2) $CO + OH = CO_2 + H$ | | 7.3 | 2.28 | | 400–650 | H ₂ O ₂ pyrolysis | 32 |
| (1) $HCHO + OH = HCO + H_2O$ (2) $CH_4 + OH = CH_3 + H_2O$ | | | | 33 27 22 33 | 525 600 650 500 | H ₂ O ₂ pyrolysis H ₂ O ₂ pyrolysis H ₂ O ₂ pyrolysis CH ₄ combustion | 32 32 32 33 |

Notes

- (a) Used at flame temperatures.
- (b) Critical survey of literature data.
- (c) $\Delta H_f(OH) = 9.33$ (10.0) kcal/mole.
- (d) This value was used by the authors in the region 1000°–1500° but its origin was not stated.
- (e) Measured relative to $OH + H_2 = H_2O + H$, for which $k = 1.6 \times 10^9$ at 520 °C.
- (f) Calculated on the assumption that oxygen atom reactions could be neglected.
- (g) The products of this reaction are $CO_2 + H$.

References

1. Avramenko and Lorentso, *Zhur. Fiz. Khim.* **24**, 207 (1950).
2. Baldwin, *Trans. Faraday Soc.* **52**, 1344 (1956).
3. Kondratiev, 7th. Int. Comb. Symp., p. 41 (Butterworths, London, 1959).
4. Fenimore and Jones, *J. Phys. Chem.* **65**, 993 (1961).
5. Del Greco and Kaufman, *Disc. Faraday Soc.* **33**, 128 (1962).
6. Kaufman and Del Greco, 9th. Int. Comb. Symp., p. 659 (Academic Press, N.Y., 1963).
7. Wise, Ablow, Sancier, and Wood, Project Squid, Semi Annual Report, p. 59 (April, 1964).
8. Dixon-Lewis, Sutton, and Williams, *Trans. Faraday Soc.* **61**, 255 (1965).
9. Dixon-Lewis, Sutton, and Williams, *J. Chem. Soc.* 5724 (1965).
10. Dixon-Lewis, Sutton and Williams, 10th. Int. Comb. Symp. Comb. Inst., p. 495 (Pittsburgh, 1965).
11. Dixon-Lewis, Wilson, and Westenberg, *J. Chem. Phys.* **44**, 2877 (1966).
12. Skinner and Ringrose, *J. Chem. Phys.* **42**, 2190 (1965).
13. Avramenko and Kolesnikova, "Advances in photochemistry," p. 25 (Interscience Publishers, N.Y., 1964).
14. Fenimore and Jones, *J. Phys. Chem.* **65**, 2200 (1961).
15. Westenberg and Fristom, *J. Phys. Chem.* **65**, 591 (1961).
16. Fristom, 9th. Int. Comb. Symp., p. 560 (Academic Press N.Y., 1963).
17. Intezarova, Kondratiev, and Mukhoyan, *Kinetika i Kataliz* **5**, 585 (1964).
18. Fenimore and Jones, *J. Chem. Phys.* **41**, 1887 (1964).
19. Avramenko and Lorentso, *Dokl. Akad. Nauk. S.S.S.R.* **67**, 867 (1949).
20. Westenberg and Fristom, 10th. Int. Comb. Symp. Comb. Inst., p. 473 (Pittsburgh, 1965).
21. Baldwin and Walker, *Trans. Faraday Soc.* **60**, 1236 (1964).
22. Avramenko and Lorentso, *Dokl. Akad. Nauk. S.S.S.R.* **69**, 205 (1949).
23. Husain and Norrish, *Proc. Roy. Soc.* **273A**, 165 (1963).
24. Wilson, 10th. Int. Comb. Symp. Comb. Inst., p. 47 (Pittsburgh, 1965).
25. Jost, Schecker, and Wagner, *Z. Phys. Chem.* **45**, 56 (1965).
26. Nosova, Lovachev, and Vedenev, *Comb. and Flame* **8**, 163 (1964).
27. Baldwin, Jackson, Walker, and Webster, 10th. Int. Comb. Symp. Comb. Inst., p. 473 (Pittsburgh, 1965).
28. Forst and Giguere, *J. Phys. Chem.* **62**, 340 (1958).
29. Baldwin and Mayor, *Trans. Faraday Soc.* **56**, 103 (1960).
30. Baldwin and Doran, *Trans. Faraday Soc.* **57**, 1578 (1961).
31. Ung and Back, *Can. J. Chem.* **42**, 753 (1964).
32. Hoare, *Nature* **194**, 283 (1962).
33. Blundell, Cook, Hoare, and Milne, 10th. Int. Comb. Symp. Comb. Inst., p. 445 (Pittsburgh, 1965).
34. Fenimore and Jones, 9th. Int. Comb. Symp., p. 560 (Academic Press, N.Y., 1963).

Metathetical Reactions of Hydroperoxyl Radicals

| Reactants | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|-------------------------------|-------|------------------------------------|--|-------------------|---|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| H ₂ | | (kcal mole ⁻¹) 24.0 | (cm ³ mole ⁻¹ sec ⁻¹) 11.08 | °C 500 | H ₂ /O ₂ ignition | 1, 2 |
| | | 14.8 ± 2.2 | k = 1.1 × 10 ⁷ | 617-1007 | H ₂ /O ₂ ignition | 3 |
| | | | k = 1.3 × 10 ⁶ | 500 | H ₂ /O ₂ ignition | 4 |
| | | | 34.6 | 16.51 | 687-807 | |
| H ₂ O | (a) | 8.0 | 8.1 | 600-615 | H ₂ /O ₂ ignition | 1, 8 |
| | (b) | 30.0 | 13.26 | 600-615 | H ₂ /O ₂ ignition | 2 |

Ratios of Rate Constants (Hydroperoxyl Radicals)

| Reactions | Notes | E ₁ - E ₂ | log ₁₀ A ₁ /A ₂ | k ₁ /k ₂ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|--------------------------------|-------------------|--|-----------|
| (1) CO + HO ₂ = CO ₂ + OH (2) H ₂ + HO ₂ = H + H ₂ O ₂ | | (kcal mole ⁻¹) | | 9.5 ± 2 | °C 500 | H ₂ /CO/O ₂ ignition | 5 |
| (1) HCHO + HO ₂ = HCO + H ₂ O ₂ (2) CO + HO ₂ = CO ₂ + OH | | | | 340 | 525 | CH ₄ /O ₂ T | 6 |

Notes

- (a) These values must be rejected on thermochemical grounds.
(b) Is a recalculation of data from (a).

References

1. Voevodsky, 7th. Int. Comb. Symp., p. 34 (Butterworths, London, 1959).

2. Kondratiev, Dokl. Akad. Nauk. S.S.S.R. **137**, 120 (1961).
3. Baldwin and Mayor, Trans. Faraday Soc. **56**, 103 (1960).
4. Miyama and Takeyama, J. Chem. Phys. **41**, 2287 (1964).
5. Baldwin, Jackson, Walker, and Webster, 10th Int. Comb. Symp. Comb. Inst., p. 423 (Pittsburgh, 1965).
6. Blundell, Cook, Hoare, and Milne, 10th. Int. Comb. Symp. Comb. Inst., p. 445 (Pittsburgh, 1965).
7. Dixon-Lewis and Williams, Nature **196**, 1309 (1962).
8. Poltorak and Voevodsky, Zhur. Fiz. Khim. **24**, 299 (1950).
9. Skinner and Ringrose, J. Chem. Phys. **42**, 2190 (1965).

Metathetical Reactions of Imino Radicals

| Reactants | Notes | | E | log ₁₀ A | Temperature range | Radical source | Reference |
|------------------------|-------|-----|---------------------------------|--|-------------------|----------------|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | | |
| HNCO | (a) | 3.6 | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C -31-200 | HNCO P | 1 |

Note

(a) The products of this reaction are NH₂ and NCO.

Reference

1. Mui and Back, Can. J. Chem. **41**, 826 (1963).

Metathetical Reactions of Amino Radicals

| Reactants | Notes | E | $\log_{10}A$ | Temperature range | Radical source | References |
|--|-------|---|--|--------------------------|-------------------------------|------------|
| $\text{NH}_2 + \text{N}_2\text{H}_4$ $= \text{NH}_3 + \text{N}_2\text{H}_3$ | | (kcal mole^{-1}) $k = 7.9 \times 10^{10}$ | $(\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1})$ 13.5 | $^{\circ}\text{C}$ 25 | | 1 |
| | | 17 | | 830-1130 | N_2H_4 S.T. | 4 |
| $\text{NH}_2 + \text{O}_2 = \text{NH} + \text{HO}_2$ | | 42.5 ± 1.2 | | 1277-2027 | NH_3/O_2 S.T. | 2 |

Metathetical Reactions of Amino Radicals Ratios of Rate Constants

| Reactants | Notes | $E_1 - E_2$ | $\log_{10}A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|---------------------------|--------------------|----------------------|--------------------|---|-----------|
| (1) $\text{NH}_2 + (\text{CH}_3)_2\text{N} \cdot \text{NH}_2$ $= \text{NH}_3 + (\text{CH}_3)_2\text{N} \cdot \text{NH}$ | | (kcal mole^{-1}) | | | $^{\circ}\text{C}$ | | |
| (2) $\text{NH}_2 + (\text{CH}_3)_2\text{N} \cdot \text{NH}_2$ $= \text{NH}_3 + (\text{CH}_3)(\text{CH}_2)\text{N} \cdot \text{NH}_2$ | | | | 6.6×10^{-3} | 250 | $(\text{CH}_3)_2\text{N} \cdot \text{NH}_2$ T | 3 |

References

1. Diesen, J. Chem. Phys. **39**, 2121 (1963).
2. Takeyama and Miyama, J. Chem. Phys. **42**, 3737 (1965).
3. Cordes, J. Phys. Chem. **65**, 1473 (1961).
4. Michel and Wagner, 10th Int. Comb. Symp. Comb. Inst., p. 353 (Pittsburgh, 1965).

Metathetical Reactions of Difluoroamino Radicals

| Reactions | Notes | E | log ₁₀ A | Temperature range | Radical source | Reference |
|--|-------|----------------------------|---|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | | |
| Alkanes | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| <i>n</i> -C ₄ H ₁₀ | | 22.20 ± 0.08 | 11.83 ± 0.04 | 180–282 | N ₂ F ₄ / <i>n</i> -C ₄ H ₁₀ T | 1 |
| <i>i</i> -C ₄ H ₁₀ | | 18.49 ± 0.10 | 10.49 ± 0.04 | 155–282 | N ₂ F ₄ / <i>i</i> -C ₄ H ₁₀ T | 1 |
| cyclo-C ₅ H ₁₀ | | 19.91 ± 0.11 | 10.93 ± 0.05 | 180–282 | N ₂ F ₄ /cyclo-C ₅ H ₁₀ T | 1 |
| neo-C ₅ H ₁₂ | | 26.68 ± 0.07 | 13.22 ± 0.03 | 180–282 | N ₂ F ₄ /neo-C ₅ H ₁₂ T | 1 |
| Ketone | | | | | | |
| CH ₃ COCH ₃ | | 19.56 ± 0.11 | 10.71 ± 0.05 | 170–280 | N ₂ F ₄ /CH ₃ COCH ₃ T | 2 |
| FLUORINE ATOM TRANSFER | | | | | | |
| F ₂ O | | 22.5 ± 0.7 | 14.01 | 120–170 | N ₂ F ₄ /F ₂ O T | 3 |

References

1. Grzechowiak, Kerr, and Trotman-Dickenson, Chem. Comm. 109 (1965).
2. Grzechowiak, Kerr, and Trotman-Dickenson, J. Chem. Soc. 5080 (1965).
3. Rubinstein, Sicre, and Schumacher, Z. Phys. Chem. **43**, 64 (1964).

Metathetical Reactions of Nitrate Radicals

| Reactions | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|-------|---------------------------|--|--------------------|---|-----------|
| $\text{NO}_3 + \text{NO} = 2\text{NO}_2$ | | (kcal mole^{-1}) | $(\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1})$ | $^{\circ}\text{C}$ | | |
| | | 1.4 ± 2.5 | $k = 6 \times 10^{12}$ | 27 | N_2O_5 S.T. | 1 |
| | | | 13.8 | 27 | NO_2/O_3 P | 2 |
| | | 1.7 | $k = 2.7 \times 10^{12}$ | 200-550 | NO_2 T | 3 |
| | | | 12.62 | 25 | NO_2 F.P. | 4 |
| $\text{NO}_3 + \text{NO}_2 = \text{NO}_2 + \text{NO} + \text{O}_2$ | | 3.9 ± 1.0 | 11.22 | 27-547 | N_2O_5 S.T. | 1, 2 |
| | | 3.2 ± 1.0 | 11.07 ± 0.47 | 200-550 | NO_2 T | 4 |
| $\text{NO}_3 + \text{NOCl} = \text{NO}_2 + \text{NO}_2\text{Cl}$ | (a) | | $k = 7.0 \times 10^7$ | 40 | $\text{NOCl}/\text{O}_3/\text{N}_2\text{O}_5$ T | 6 |
| | | | $k = 2.3 \times 10^7$ | 40 | | 2 |

Ratios of Rate Constants

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|--|-------|---------------------------|---------------------|-----------|--------------------|------------------------------------|-----------|
| (1) $\text{NO}_3 + \text{NO} = 2\text{NO}_2$ (2) $\text{NO}_3 + \text{NO}_2 = \text{NO}_2 + \text{NO} + \text{O}_2$ | | (kcal mole^{-1}) | | | $^{\circ}\text{C}$ | | |
| | (b) | 1.4 | 2.2 | 60 | 434 | NO_2 T | 7 |
| | (c) | 2.3 | 3.2 | | 20-30 | $\text{NO}/\text{N}_2\text{O}_5$ T | 8, 9 |
| | | | | | 20-30 | $\text{NO}/\text{N}_2\text{O}_5$ T | 8, 9 |

Notes

(a) The A factor was assumed, only the rate constant was determined.

(b) At 400 mm total pressure.

(c) At 57 mm total pressure.

References

1. Davidson and Schott, *J. Chem. Phys.* **27**, 317 (1957).
2. Schott and Davidson, *J. Am. Chem. Soc.* **80**, 1841 (1958).
3. Ford, Doyle, and Endow, *J. Chem. Phys.* **32**, 1256 (1960).
4. Ashmore and Burnett, *Trans. Faraday Soc.* **58**, 253 (1962).
5. Husain and Norrish, *Proc. Roy. Soc.* **273A**, 165 (1963).
6. Johnston and Leighton, *J. Am. Chem. Soc.* **75**, 3612 (1953).
7. Ashmore and Levitt, *Research* **9**, S25 (1956).
8. Hisatsune, Crawford, and Ogg, *J. Am. Chem. Soc.* **79**, 4648 (1957).
9. Hisatsune, McHale, Nightingale, Rotenberg, and Crawford, *J. Chem. Phys.* **23**, 2467 (1955).

Metathetical Reactions of Thiyl Radicals

| Reactant | Note | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | References |
|----------|------|----------|---------------|-------------------|----------------|------------|
|----------|------|----------|---------------|-------------------|----------------|------------|

HYDROGEN ATOM TRANSFER

| | | | | | | |
|---------------------------|-----|--|--|---------------|--|------|
| CH_3OCH_3 | (a) | 6.8 <i>(kcal mole⁻¹)</i> | 11 <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C 360-440 | $\text{CH}_3\text{OCH}_3/\text{H}_2\text{S}$ T | 1, 2 |
|---------------------------|-----|--|--|---------------|--|------|

Notes

(a) For further data on systems containing HS radicals, see reference 3.

References

1. Imai and Toyama, Bull. Chem. Soc. Japan **34**, 328 (1961).
2. Anderson and Benson, J. Chem. Phys. **39**, 1677 (1963).
3. Imai, Yoshida, and Toyama, Bull. Chem. Soc. Japan **35**, 752 (1962).

Reactions of Chloromonoxy Radicals

| Reactions | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|-----------|---------------------------|--|--------------------|----------------------------|-----------|
| $\text{ClO} + \text{Cl}_2\text{O} = \text{ClO}_2 + \text{Cl}_2$ | | (kcal mole^{-1}) | $(\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1})$ | $^{\circ}\text{C}$ | Cl_2O F.P. | 1 |
| $\text{ClO} + \text{Cl}_2\text{O} = \text{Cl} + \text{O}_2 + \text{Cl}_2$ | | | $k = 10^8$ | | | |
| $\text{ClO} + \text{NO}_3\text{Cl} = \text{NO}_2 + \text{Cl}_2 + \text{O}_2$ | ~ 17 | | $k = 5.3 \times 10^7$ | 90-130 | NO_3Cl T | 2 |

Ratios of Rate Constants (Chloromonoxy Radicals)

| Reactions | Notes | $E_1 - E_2$ | $\log_{10} A_1/A_2$ | k_1/k_2 | Temperature range | Radical source | Reference |
|---|-------|---------------------------|---------------------|---------------|---------------------------|--|-----------|
| (1) $\text{ClO} + \text{C}_3\text{H}_8 = \text{HOCl} + \text{CH}_3\text{CHCH}_3$ (2) $\text{ClO} + \text{C}_3\text{H}_8 = \text{HOCl} + \text{CH}_3\text{CH}_2\text{CH}_2$ | | (kcal mole^{-1}) | | 7.2 ± 0.5 | $^{\circ}\text{C}$ 100 | $\text{C}_3\text{H}_8/\text{Cl}_2\text{O}$ T | 3 |

References

1. Edgecombe, Norrish, and Thrush, Proc. Roy. Soc. **243A**, 24 (1957).
2. Cafferata, Sicre, and Schumacher, Z. Phys. Chem. **29**, 188 (1961).
3. Phillips and Shaw, Proc. Chem. Soc. 294 (1962).

Reactions of Borine Radicals

| Reactants | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|--|--|-------------------|---|-----------------|
| $\text{BH}_3 + \text{B}_2\text{H}_6 = \text{B}_3\text{H}_7 + \text{H}_2$ | | (<i>kcal mole</i> ⁻¹) 11.5 | (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) 11.06 | °C | | 1, 9 |
| $\text{BH}_3 + \text{B}_2\text{D}_6 = \text{BH}_3\text{BD}_3 + \text{BD}_3$ | | 6.0 7.8 | 14.2 13.45 | 24-44 | $\text{B}_2\text{H}_6/\text{B}_2\text{D}_6$ T | 4, 5, 6, 3 1 |
| $\text{BH}_3 + \text{BH}_3\text{CO} = \text{B}_2\text{H}_6 + \text{CO}$ | | 7.0 | 11.4 | 0-30 | BH_3CO T | 7, 4, 3 |
| $\text{BH}_3 + \text{H}_2\text{O} = \text{BH}_2\text{OH} + \text{H}_2$ | | 6 | 11.28 | 51 | $\text{B}_2\text{H}_6/\text{H}_2\text{O}$ T | 8, 9 |
| $\text{BH}_3 + \text{HD} = \text{BH}_2\text{D} + \text{H}_2$ | | 7 | 11.6 | | | 9 |

Ratios of Rate Constants Involving Borine Radicals

| Reaction | Notes | <i>E</i> ₁ - <i>E</i> ₂ | $\log_{10} A_1/A_2$ | <i>k</i> ₁ / <i>k</i> ₂ | Temperature range | Radical source | Reference |
|--|-------|---|---------------------|---|-------------------|-------------------------------------|-----------|
| (1) $\text{BH}_3 + \text{B}_2\text{D}_6 = \text{BD}_3\text{BH}_3 + \text{BD}_3$ (2) $\text{BH}_3 + \text{D}_2 = \text{BH}_2\text{D} + \text{HD}$ | | (<i>kcal mole</i> ⁻¹) | | 0.25 | °C 55 | $\text{B}_2\text{H}_6/\text{D}_2$ T | 2, 3 |
| (1) $\text{BH}_3 + \text{O}_2 = \text{BH}_2\text{OH} + \text{O}$ (2) $\text{BH}_3 + \text{O}_2 + \text{M} = \text{HBO}_2 + \text{H}_2 + \text{M}$ | | 2.0 | | | 150-200 | $\text{B}_2\text{H}_6/\text{O}_2$ T | 1 |

References

1. Roth and Bauer, 5th Int. Comb. Symp., p. 710 (Reinhold, N.Y., 1955).
2. Marcus, J. Chem. Phys. **23**, 1107 (1955).
3. Garabedian and Benson, J. Am. Chem. Soc. **86**, 176 (1964).
4. Bauer, Shepp, and McCoy, J. Am. Chem. Soc. **75**, 1003 (1953).
5. Trotman-Dickenson, "Gas Kinetics," p. 237 et seq. (Butterworths, London, 1955).
6. Maybury and Koski, J. Chem. Phys. **21**, 742 (1953).
7. Burg, J. Am. Chem. Soc. **74**, 3482 (1952).
8. Weiss and Shapiro, J. Am. Chem. Soc. **75**, 1221 (1953).
9. Bauer, J. Am. Chem. Soc. **78**, 5775 (1956).

Metathetical Reactions of F₃SO and FSO₃ Radicals

| Reactions | Notes | <i>E</i> | log ₁₀ <i>A</i> | Temperature range | Radical source | Reference |
|--|-------|--|---|-------------------|--|-----------|
| F ₃ SO + F ₂ = F ₄ SO + F | | (kcal mole ⁻¹) 10.0 ± 2 | (cm ³ mole ⁻¹ sec ⁻¹) | °C 5-20 | F ₂ /F ₂ SO P | 1 |
| FSO ₃ + F ₂ = F ₂ SO ₃ + F | | 14.7 ± 1.0 | 8.68 | 230-250 | F ₂ /F ₂ S ₂ O ₆ T | 2 |

References

1. Castellano and Schumacher, Z. Phys. Chem. N.F. **40**, 51 (1964).

2. Castellano and Schumacher, Z. Phys. Chem. N.F. **44**, 57 (1965).

Radical Disproportionation Reactions

The reactions are classified in the same order as the tables of radical reactions. Thus the first set of reactions are those involving atoms, followed by reactions of radicals containing one carbon atom, etc.

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | $\log_{10} k$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|--|------------------------|--|--------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H + OH = H ₂ + O | | 5.8 ± 1.5 7.4 | 12.76 ± 1.0 12.74 | | 25-2000 687-807 | H ₂ /O ₂ S.T. | 4 12 |
| H + HO ₂ = 2OH | | | | 14.86 13.84 | 25 500 | H ₂ /O ₂ - H | 5 6 |
| H + HNO = H ₂ + NO | | | | > 9.78 > 10.48 12.78 ± 0.25 | 20 -47 1300-1700 | H ₂ /discharge H ₂ /discharge H ₂ /O ₂ /N ₂ flame | 1 2 3 |
| H + C ₂ F ₄ Br = C ₂ F ₄ + HBr | | 11.5 | 16.3 | | 855-1013 | H ₂ /O ₂ /C ₂ F ₄ Br ₂ S.T. | 7 |
| Cl + C ₂ HCl ₄ = C ₂ HCl ₃ + Cl ₂ | | | | 13.85 | 224 | CHCl:CCl ₂ /Cl ₂ P | 25, 24 |
| Cl + C ₂ Cl ₅ = C ₂ Cl ₄ + Cl ₂ | | 1.2 | 15.6 | | | | 24 |
| Cl + COCl = CO + Cl ₂ | (a) | 0.8 2.4 | 14.6 13.70 | | 25-55 15-450 | CO/Cl ₂ P | 33, 24 34 |
| Cl + Cl ₃ = 2Cl ₂ | | | | ≤ 14.23 | 20 | Cl ₂ /discharge | 19, 20 |
| I + NOI = NO + I ₂ | | | | 12.60 | 60 | NO/I ₂ F.P. | 30 |
| 2CN = C ₂ + N ₂ | | 96 | | | 3177-4527 | C ₂ N ₂ S.T. | 14, 15 |
| 2C ₂ H ₃ Cl ₂ = C ₂ H ₃ Cl + C ₂ H ₃ Cl ₃ | | 0.3 | 12.9 | | 25-55 | CH ₂ :CHCl/Cl ₂ P | 29 |
| 2CHCl ₂ CHCl = CHCl:CHCl + CHCl ₂ CHCl ₂ | | 0.5 ± 0.5 | 13.47 ± 0.3 | | 30-65 | <i>cis</i> -CHCl:CHCl/Cl ₂ P | 31 |
| 2C ₂ Cl ₅ = C ₂ Cl ₄ + C ₂ Cl ₆ | | 0.08 | 11.66 | | 87-247 | C ₂ Cl ₄ /Cl ₂ P | 32 |
| 2CH ₃ CO ₃ = (CH ₃ CO) ₂ O ₂ + O ₂ | | | | 13.95 | 20 | CH ₃ CHO/O ₂ P | 27 |
| 2C ₂ H ₅ CO ₃ = (C ₂ H ₅ CO) ₂ O ₂ + O ₂ | | | | 13.43 | 22 | CH ₃ CH ₂ CHO/O ₂ P | 28 |
| 2OH = H ₂ + O ₂ | | 48.6 | 12.85 | | 687-807 | H ₂ /O ₂ S.T. | 12 |
| 2OH = H ₂ O + O | (b) | | | 12.18 12.08 | 37-107 27 | H ₂ /discharge H ₂ /discharge | 10 11 |
| | (b) | 1.0 ± 0.5 3.8 | 12.88 ± 0.3 14.87 | | 25-2000 687-807 | H ₂ /O ₂ S.T. H ₂ /O ₂ S.T. | 4 12 |
| | (b) | | | 12.19 | 27 | NO ₂ /H reaction | 13 |
| OH + HNO = H ₂ O + NO (or H ₂ + NO ₂) | | | | 13.95 | 1300-1700 | H ₂ /O ₂ /NO flame | 3 |
| 2HO ₂ = H ₂ O ₂ + O ₂ | | | | 12.25 13.81 | room temp. 25 | H ₂ /O ₂ - H | 6 5 |
| 2NH ₂ = NH ₃ + NH | | | | 13.40 11.66 | 1627-2127 | NH ₂ NH ₂ S.T. NH ₃ F.P. | 8 9 |
| 2N ₂ H ₃ = 2NH ₃ + N ₂ | | | | ≥ 12.48 | 150 | H ₂ /discharge | 18 |

Radical Disproportionation Reactions—Continued

The reactions are classified in the same order as the tables of radical reactions. Thus the first set of reactions are those involving atoms, followed by reactions of radicals containing one carbon atom, etc.

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | $\log_{10} k$ | Temperature range | Radical source | Reference |
|--|-------|----------------------------|---|---|-------------------|---|--------------|
| | | (kcal mole ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | (cm ³ mole ⁻¹ sec ⁻¹) | °C | | |
| 2HNO = H ₂ O + N ₂ O | | | | > 7.48 ~ 8.9 | 27 190 | NO/H reaction (CH ₃) ₂ CHNO T | 16 17 |
| 2NO ₃ = 2NO ₂ + O ₂ | | 7.7 ± 1.0 | 12.42 | | 280–830 | N ₂ O ₅ S.T. | 26 |
| 2ClO = Cl ₂ + O ₂ | | | | 10.38 10.93 | 20 | Cl ₂ O F.P. Cl ₂ O F.P. | 21 22, 23 |

Ratios of Rate Constants (Radical Disproportionation Reactions)

| Reactions | Notes | Rate constants ratios | Temperature range | Radical source | Reference |
|--|-------|--|-------------------|--|-----------|
| | | cm ³ , mole. sec. units | °C | | |
| (1) H + HO ₂ = 2OH (2) 2HO ₂ = H ₂ O ₂ + O ₂ | | $k_1^2/k_2 = 5.1 \times 10^{14}$ | 500 | H ₂ /O ₂ ignition | 35, 36 |
| (1) 2HS = H ₂ + S ₂ (2) 2HS = H ₂ S + S | | $k_1/k_2 = 0.15$ | room temp. | H ₂ S P | 37 |
| (1) 2HNO = H ₂ O + N ₂ O (2) 2HNO = 2NO + H ₂ | | $k_1/k_2 = 8$ | 25 | C ₂ H ₅ OH/NO H | 38 |
| (1) C ₂ H ₅ + C ₃ H ₉ O ₂ = C ₂ H ₄ + C ₂ H ₅ COOC ₂ H ₅ (2) C ₂ H ₅ + C ₃ H ₉ O ₂ = C ₂ H ₆ + C ₅ H ₈ O ₂ | | $k_1/k_2 = 0.1$ | 29 | C ₂ H ₅ COOC ₂ H ₅ P | 39 |
| (1) 2C ₂ H ₅ O = C ₂ H ₅ OH + CH ₃ CHO (2) C ₂ H ₅ O = CH ₃ + CH ₂ O | | $k_1/k_2^2 = 1.2 \times 10^{10}$ $k_1/k_2^2 = (3 \pm 1) \times 10^{12}$ | 152 30 | C ₂ H ₅ COOC ₂ H ₅ P C ₂ H ₅ COOC ₂ H ₅ P | 40 40 |
| (1) 2(CH ₃) ₃ CO = (CH ₃) ₃ COH + (CH ₃) ₂ CCH ₂ O (2) (CH ₃) ₃ CO = CH ₃ COCH ₃ + CH ₃ | | $k_1/k_2^2 \leq 3.0 \times 10^8$ | 25 | ((CH ₃) ₃ CO) ₂ P | 41 |

Notes

- (a) Review of literature data.
(b) Value based on E.S.R. measurements of the radical concentration.

References

- Clyne and Thrush, *Trans. Faraday Soc.* **57**, 1305 (1961).
- Clyne and Thrush, *Disc. Faraday Soc.* **33**, 139 (1962).
- Bulewicz and Sugden, *Proc. Roy. Soc.* **277A**, 143 (1964).
- Kaufman and Del Greco, 9th Int. Comb. Symp., p. 659 (Academic Press, N.Y., 1963).
- Burgess and Robb, *Chem. Soc. Special Publ. No. 9* 167 (1958).
- Dixon-Lewis and Williams, *Nature* **196**, 1309 (1962).
- Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
- Diesen, *J. Chem. Phys.* **39**, 2121 (1963).
- Salzman and Bair, *J. Chem. Phys.* **41**, 3654 (1964).
- Del Greco and Kaufman, *Disc. Faraday Soc.* **33**, 128 (1962).
- Westenberg and De Haas, *J. Chem. Phys.* **43**, 1550 (1965).
- Skinner and Ringrose, *J. Chem. Phys.* **42**, 2190 (1965).
- Dixon-Lewis, Wilson, and Westenberg, *J. Chem. Phys.* **44**, 2877 (1966).
- Fairbairn, *Proc. Roy. Soc.* **267A**, 88 (1962).
- Paterson and Greene, *J. Chem. Phys.* **36**, 1146 (1962).
- Clyne, 10th Int. Comb. Symp., p. 311 (Comb. Inst., Pittsburgh, 1965).

17. Ferguson and Phillips, *J. Chem. Soc.* 4416 (1965).
18. Schiavello and Volpi, *J. Chem. Phys.* **37**, 1510 (1962).
19. Hutton and Wright, *Trans. Faraday Soc.* **61**, 78 (1965).
20. Hutton, *Nature* **203**, 835 (1964).
21. Edgecombe, Norrish, and Thrush, *Proc. Roy. Soc.* **243A**, 24 (1957).
22. Lipscomb, Norrish, and Porter, *Nature* **174**, 785 (1954).
23. Porter and Wright, *Disc. Faraday Soc.* **14**, 23 (1953).
24. Goldfinger, Jeunehomme, and Martens, *J. Chem. Phys.* **29**, 456 (1958).
25. Huybrechts, Meyers, and Verbecke, *Trans. Faraday Soc.* **58**, 1128 (1962).
26. Schott and Davidson, *J. Am. Chem. Soc.* **80**, 1841 (1958).
27. McDowell and Sharples, *Can. J. Chem.* **36**, 268 (1958).
28. McDowell and Sharples, *Can. J. Chem.* **36**, 1227 (1958).
29. Dainton, Lomax, and Weston, *Trans. Faraday Soc.* **58**, 308 (1962).
30. Ayscough, Cocker, Dainton, and Hirst, *Trans. Faraday Soc.* **58**, 295 (1962).
31. Dusoleil, Goldfinger, Mahieu-van der Auwera, Martens, and van der Auwera, *Trans. Faraday Soc.* **57**, 2197 (1961).
32. Burns and Dainton, *Trans. Faraday Soc.* **48**, 39 (1952).
33. Bodenstein, Brenschede, and Schumacher, *Z. Phys. Chem.* **40B**, 121 (1938).
34. Baldwin and Mayor, *Trans. Faraday Soc.* **56**, 103 (1960).
35. Baldwin and Mayor, *Trans. Faraday Soc.* **56**, 80 (1960).
36. Darwent and Roberts, *Proc. Roy. Soc.* **216A**, 344 (1953).
37. Knight and Gunning, *Can. J. Chem.* **39**, 2466 (1961).
38. Wijnen, *J. Am. Chem. Soc.* **82**, 3034 (1960).
39. Wijnen, *J. Am. Chem. Soc.* **80**, 2394 (1958).
40. McMillan and Wijnen, *Can. J. Chem.* **36**, 1227 (1958).




Disproportionation/Combination Ratios

Index

| (a) Hydrogen Atom Transfer | Page |
|--|------|
| Reactions of | |
| 1. Hydrogen atoms, H..... | 109 |
| 2. Methyl, CH ₃ | 109 |
| 3. Methoxy, CH ₃ O..... | 110 |
| 4. Halogenated methyl, CCl ₃ and CF ₂ H..... | 110 |
| 5. Vinyl, CH ₂ CH..... | 111 |
| 6. Ethyl, C ₂ H ₅ | 111 |
| 7. Ethoxy, C ₂ H ₅ O..... | 112 |
| 8. Hydroxy ethyl, CH ₃ CH(OH)..... | 112 |
| 9. Halogenated ethyls, C ₂ F ₅ , C ₂ H ₄ Cl..... | 112 |
| 10. Allyl, CH ₂ CHCH ₂ | 113 |
| 11. <i>n</i> -Propyl, CH ₃ CH ₂ CH ₂ | 113 |
| 12. <i>iso</i> -Propyl, CH ₃ CHCH ₃ | 113 |
| 13. Propoxy, CH ₃ CH ₂ CO..... | 114 |
| 14. <i>n</i> -Perfluoropropyl, CF ₃ CF ₂ CF ₂ | 114 |
| 15. Butyls, CH ₃ CH ₂ CH ₂ CH ₂ , CH ₃ CHCH ₂ CH ₃ , (CH ₃) ₂ CHCH ₂ , CH ₃) ₃ C..... | 114 |
| 16. Monochlorobutyl, C ₄ H ₈ Cl..... | 115 |
| 17. Pentyl, <i>n</i> -C ₅ H ₁₁ , cyclo-C ₅ H ₉ | 115 |
| 18. Hexyl, cyclo-C ₆ H ₁₁ , C ₆ H ₁₃ | 115 |
| 19. Amino, NH ₂ | 115 |
| 20. Nitric oxide, NO..... | 115 |
| 21. Iodine atom, I..... | 115 |
| (b) Chlorine Atom Transfer | |
| 1. Halogenated methyl, CF ₂ Cl..... | 115 |
| 2. Halogenated ethyl, C ₂ H ₄ Cl, C ₂ F ₂ Cl ₃ | 116 |
| 3. Monochlorobutyl, C ₄ H ₈ Cl..... | 116 |
| (c) Oxygen Atom Transfer | |
| 1. Methyl, CH ₃ | 116 |

Disproportionation/Combination Ratios

The following convention is used throughout this table, e.g. for ethyl and *n*-propyl radicals, the possible reactions are:



k_2/k_1 is listed as $\text{C}_2\text{H}_5/n\text{-C}_3\text{H}_7$ and k_3/k_1 as $n\text{-C}_3\text{H}_7/\text{C}_2\text{H}_5$.

The table is constructed with reference to the radical receiving the transferred atom. In the above example, k_2/k_1 is in the section on ethyl radicals and k_3/k_1 , with the *n*-propyl radicals. The radicals are further classified according to the number of carbon atoms they contain. Thus the table starts with H atom reactions, followed by C_1 reactions (methyl radicals and halogen and oxygen containing derivatives of methyl radicals,) then by C_2 reactions, etc.

| Reactants | Notes | $k_{\text{dis}}/k_{\text{comb.}}$ | Temperature range | Radical source | Reference |
|---|-------|-----------------------------------|-------------------|--|-----------|
| HYDROGEN ATOM TRANSFER | | | | | |
| Hydrogen atom | | | | | |
| H/ C_2H_5 | | 0.05 ± 0.03 | °C 26–203 | $(\text{C}_2\text{H}_5)_2\text{CO} + \text{H}_2$ – H | 1 |
| D/ C_2H_5 | (a) | 0.21 ± 0.07 | 26–203 | $(\text{C}_2\text{H}_5)_2\text{CO} + \text{D}_2$ – H | 1 |
| D/ CH_3CD_2 | | 0.06 ± 0.03 | 26–203 | $(\text{CH}_3\text{CD}_2)_2\text{CO} + \text{D}_2$ – H | 1 |
| D/ <i>i</i> - C_3H_7 | | 0.2 | 85 | | 2 |
| H/ C_5H_{11} | | 0.099 | 25 | iso- C_5H_{12} – H | 3 |
| H/ C_6H_{13} | | 0.12 | 25 | <i>n</i> - C_6H_{14} – H | 4 |
| Methyl | | | | | |
| $\text{CH}_3/\text{CH}_2\cdot\text{CH}$ | | 2.3 | 175 | $\text{CH}_2\cdot\text{CHCHO}$ and $\text{HCOOCH}\cdot\text{CH}_2$ + CH_3 | 5 |
| $\text{CH}_3/\text{C}_2\text{H}_5$ | | 0.04 ± 0.02 | 78–130 | $\text{CH}_3\text{COC}_2\text{H}_5$ P | 6 |
| | | 0.055 | 77–230 | $\text{HCOOC}_2\text{H}_5 + \text{CH}_3$ | 7 |
| | | 0.039 ± 0.007 | 110–166 | $(\text{CH}_3)_2\text{CO} + (\text{C}_2\text{H}_5)_2\text{CO}$ P | 8 |
| | | 0.039 | | | |
| $\text{CH}_3/\text{CH}_3\text{CD}_2$ | | 0.06 ± 0.01 | 90 | $(\text{CH}_3)_2\text{CO}$ + $(\text{CH}_3\text{CD}_2)_2\text{CO}$ P | 9 |
| $\text{CD}_3/\text{C}_2\text{H}_5$ | | < 0.08 | 27 | $(\text{CD}_3)_2\text{CO} + (\text{C}_2\text{H}_5)_2\text{CO}$ P | 10 |
| $\text{CH}_3/n\text{-C}_3\text{H}_7$ | | 0.095 ± 0.01 | 74–178 | $\text{HCOOCH}_2\text{CH}_2\text{CH}_3$ + CH_3 | 11 |
| | | 0.025 ± 0.004 | 139–173 | <i>n</i> - $\text{C}_3\text{H}_7\text{CHO}$ + CH_3 | 12 |
| | | 0.041 ± 0.01 | 118–144 | $(\text{CH}_3)_2\text{CO}$ + $(n\text{-C}_3\text{H}_7)_2\text{N}_2$ P | 8 |
| | (b) | 0.14 | | | 13 |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | <i>k</i> _{dis} / <i>k</i> _{comb.} | Temperature range | Radical source | Reference |
|---|-------|---|-------------------|---|-----------|
| | | | °C | | |
| CD ₃ /CH ₃ CH ₂ CD ₂ | | 0.05 | 25–28 | CD ₃ COCD ₂ CH ₂ CH ₃ P | 14 |
| CH ₃ / <i>i</i> -C ₃ H ₇ | | 0.216 ± 0.032 | 80–150 | (CH ₃) ₂ CO + CH ₃ COCH(CH ₃) ₂ P | 15 |
| | | 0.195 0.21 | 94–181 | HCOOCH(CH ₃) ₂ + CH ₃ | 11 |
| CD ₃ /CH ₃ CDCH ₃ | | 0.17 ± 0.03 | 53–116 | CD ₃ COCD(CH ₃) ₂ P | 16 |
| CH ₃ / <i>n</i> -C ₄ H ₉ | | 0.15 | 70–195 | HCOOC ₄ H ₉ + CH ₃ | 13, 103 |
| CH ₃ / <i>sec</i> -C ₄ H ₉ | | 0.30 | –103–+25 | <i>cis</i> -CH ₃ CH:CHCH ₃ + H | 17 |
| | | 0.07 | 107–168 | CH ₃ CH:CH ₂ + CH ₃ | 18 |
| CH ₃ / <i>t</i> -C ₄ H ₉ | (c) | 0.85 ± 0.1 0.88 | 20–60 25–79 | ((CH ₃) ₃ CO) ₂ P ((CH ₃) ₃ CO) ₂ P | 19 20 |
| | (d) | 0.699 ± 0.037 0.806 0.80 | 80–188 | ((CH ₃) ₃ C) ₂ CO P | 15 21 |
| CH ₃ / <i>n</i> -C ₅ H ₁₁ | | 0.1 | 63 | CH ₃ COC ₅ H ₁₁ P | 22 |
| CD ₃ /cyclo-C ₅ H ₉ | | 0.31 | 132–218 | (CD ₃) ₂ CO P | 23 |
| CH ₃ /CDO | | 2.5 | 80–180 | (CH ₃) ₂ N ₂ P | 24 |
| CH ₃ /CH ₃ O | (a) | 1.25 1.51 ± 0.2 | 29 124–185 | CH ₃ COOCH ₃ P CH ₃ OOCH ₃ T | 25 26 |
| CH ₃ /CD ₃ O | | 1.4 ± 0.1 | 30–201 | CH ₃ COOCD ₃ P | 27 |
| CD ₃ /CD ₃ O | | 1.8 | 30–201 | CH ₃ COOCD ₃ P | 27 |
| CH ₃ /CH ₃ CO | | 0.06 0.05 | 29 30 | CH ₃ COOCH ₃ P CH ₃ COOCD ₃ P | 25 104 |
| | | ≅ 0.095 | 55 | CH ₃ CO.COCH ₃ –H | 28 |
| CD ₃ /CD ₃ CO | | 0.014 to 0.055 | 25 | CD ₃ COCD ₂ CH ₂ CH ₃ P | 14 |
| CH ₃ /CH ₃ COCH ₂ | | 0.25 ± 0.25 | 184–285 | CH ₃ COCH ₃ P | 29 |
| CH ₃ /(CH ₃) ₂ CHO | | 3.4 | 26 | ((CH ₃) ₂ CHO) ₂ P | 30 |
| Methoxy | | | | | |
| CH ₃ O/CH ₃ O | | ≧ 60 9.3 ± 0.6 | 25 room temp. | (CH ₃) ₂ N ₂ + O ₂ P CH ₃ I + O ₂ P | 31 32 |
| Halogenated Methyls | | | | | |
| CCl ₃ /C ₂ H ₅ | | 0.24 ± 0.04 0.22 ± 0.03 | 25 0–58 | (C ₂ H ₅) ₂ CO + CCl ₄ P C ₂ H ₄ + CCl ₄ P | 33 34 |
| CCl ₃ /C ₂ H ₄ Cl | | 0.11 ± 0.02 0.14 ± 0.03 | 26 0–58 | C ₂ H ₄ + CCl ₄ P C ₂ H ₄ + CCl ₄ P | 105 34 |
| CCl ₃ /C ₂ H ₄ COC ₂ H ₅ | | 0.9 ± 0.1 | 25 | (C ₂ H ₅) ₂ CO + CCl ₄ P | 33 |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | <i>k</i> _{dis} / <i>k</i> _{comb.} | Temperature range | Radical source | Reference |
|---|-------|---|-------------------|---|-----------|
| | | | °C | | |
| CF ₂ H/CF ₂ H | | 0.19 | 75–119 | CF ₂ ClH – H | 35 |
| Vinyl | | | | | |
| CH ₂ :CH/CH ₂ :CH | | 1.1 | 175 | CH ₂ :CHCHO and HCOOCH:CH ₂ + CH ₃ | 5 |
| | (e) | 0.5 | 272–301 | | 5 |
| | | 0.02 | 50 | (CH ₂ :CH) ₂ Hg P | 36 |
| CH ₂ :CH/C ₂ H ₅ | | 0.03 | 161–175 | (C ₂ H ₅) ₂ CO P | 37 |
| Ethyl | | | | | |
| C ₂ H ₅ /CH ₂ :CH | | 0.12 | 161–175 | (C ₂ H ₅) ₂ CO P | 37 |
| C ₂ H ₅ /C ₂ H ₅ | | 0.36 | 75 | (C ₂ H ₅) ₂ Hg P | 38 |
| | | 0.46 | 200 | (C ₂ H ₅) ₂ Hg P | 38 |
| | | 0.11 | 25 | (C ₂ H ₅) ₂ CO P | 39 |
| | | 0.085 | 101 | (C ₂ H ₅) ₂ CO P | 39 |
| | | 0.17 | 25 | C ₂ H ₄ + H | 40 |
| | | 0.22–0.61 | 42 | C ₂ H ₄ + H | 41 |
| | | 0.47 | 200 | C ₂ H ₄ + H | 41 |
| | | 0.10 | 25 | C ₂ H ₅ CHO P | 42 |
| | | 0.13 | 350 | (C ₂ H ₅) ₂ Hg – T | 43 |
| | | 0.13 | 122 | C ₂ H ₅ CHO + CH ₃ | 44 |
| | | 0.15 | 156 | C ₂ H ₅ CHO + CH ₃ | 44 |
| | | 0.13 ± 0.02 | 26–74 | (C ₂ H ₅) ₂ N ₂ P | 45 |
| | | 0.12 | 100–200 | (C ₂ H ₅) ₂ CO P | 46 |
| | | 0.15 ± 0.01 | 25 | C ₂ H ₄ + H | 47 |
| | | 0.136 ± 0.02 | 50–215 | (C ₂ H ₅) ₂ CO P | 48 |
| | | 0.12 ± 0.01 | 27–118 | (C ₂ H ₅) ₂ N ₂ P | 49 |
| | | 0.15 | 50–315 | C ₂ H ₅ CHO P | 50 |
| | | 0.14 | 77–230 | HCOOC ₂ H ₅ + CH ₃ | 7 |
| | (f) | { 0.16 | – 65 | | |
| | | { 0.13 | 0 | (C ₂ H ₅) ₂ N ₂ P | 51, 52 |
| | | { 0.11 ₉ | 40 | | |
| | | 0.130 ± 0.007 | 115–150 | (C ₂ H ₅) ₂ CO P | 8 |
| | | 0.11 ± 0.01 | 22 | (C ₂ H ₅) ₂ N ₂ F.P. | 53 |
| | | 0.123 ± 0.008 | 25–200 | (C ₂ H ₅) ₂ CO P | 54 |
| | | 0.134 ± 0.003 | 79–141 | (C ₂ H ₅) ₂ CO P | 55 |
| | | 0.11 | room temp. | CH ₂ :CH ₂ + H | 56 |
| | | 0.137 | 71–175 | (C ₂ H ₅) ₂ CO P | 106 |
| | | 0.135 | | | |
| CH ₃ CD ₂ /CH ₃ CD ₂ | (g) | 0.1 | 24–180 | (CH ₃ CD ₂) ₂ CO P | 57 |
| C ₂ D ₅ /C ₂ D ₅ | | 0.098 ₅ ± 0.008 | 50–197 | (C ₂ D ₅) ₂ CO P | 58 |
| C ₂ H ₅ /CH ₂ CHCH ₂ | | 0.05 | 134–175 | (C ₂ H ₅) ₂ CO P | 106 |
| C ₂ H ₅ / <i>n</i> -C ₃ H ₇ | | 0.14 | 117 | (C ₂ H ₅) ₂ CO + HCOO <i>n</i> -C ₃ H ₇ P | 59 |
| | | 0.081 ± 0.01 | 61–129 | (C ₂ H ₅) ₂ CO + (<i>n</i> -C ₃ H ₇) ₂ N ₂ P | 8 |
| | | 0.06 ± 0.04 | – 78–+24 | C ₃ H ₈ – H | 60 |
| | | 0.08 | | | |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | <i>k</i> _{dis} / <i>k</i> _{comb.} | Temperature Range | Radical Source | Reference |
|--|-------|---|-------------------|---|-----------|
| | | | °C | | |
| C ₂ H ₅ / <i>i</i> -C ₃ H ₇ | | 0.2 | room temp. | CH ₂ :CH ₂ + CH ₃ CH:CH ₂ + H | 107 |
| | | 0.21 ± 0.02 | 25 | C ₃ H ₈ - H | 61 |
| | | 0.43 ± 0.03 | 34-144 | HCOO <i>i</i> -C ₃ H ₇ + (C ₂ H ₅) ₂ CO P | 59 |
| | | 0.43 | 48-112 | C ₂ H ₅ CHO + <i>i</i> -C ₃ H ₇ CHO P | 62 |
| | | 0.43 | | | |
| C ₂ H ₅ /sec-C ₄ H ₉ | | 0.23 ± 0.01 | 25 | <i>n</i> -C ₄ H ₁₀ - H | 61 |
| C ₂ H ₅ / <i>i</i> -C ₄ H ₉ | | 0.02 ± 0.02 | 0 | <i>i</i> -C ₄ H ₁₀ - H | 60 |
| C ₂ H ₅ / <i>t</i> -C ₄ H ₉ | | 0.3 | room temp. | CH ₂ :CH ₂ + (CH ₃) ₂ C:CH ₂ + H | 107 |
| | | 0.53 | 73-80 | (C ₂ H ₅) ₂ CO + ((CH ₃) ₃ C) ₂ CO P | 62 |
| | | 0.54 ± 0.01 | 25 | <i>i</i> -C ₄ H ₁₀ - H | 61 |
| | | 0.54 | | | |
| C ₂ H ₅ /cyclo-C ₅ H ₉ | | 0.27 ± 0.05 | 25 | cyclo-C ₅ H ₁₀ - H | 61 |
| C ₂ H ₅ / <i>n</i> -C ₅ H ₁₁ | | 0.08 ± 0.04 | 0 | <i>n</i> -C ₅ H ₁₂ - H | 60 |
| C ₂ H ₅ /sec-C ₅ H ₁₁ | (h) | 0.25 ± 0.01 | 25 | <i>n</i> -C ₅ H ₁₂ - H | 61 |
| C ₂ H ₅ / <i>t</i> -C ₅ H ₁₁ | | 0.60 ± 0.01 | 25 | (CH ₃) ₂ CHCH ₂ CH ₃ - H | 61 |
| C ₂ H ₅ /cyclo-C ₆ H ₇ | (i) | 0.38 ± 0.03 | 50-120 | (C ₂ H ₅) ₂ CO P | 64 |
| C ₂ H ₅ /((CH ₃) ₃ CCHCH ₃) | | 0.20 ± 0.02 | 25 | (CH ₃) ₃ CCH ₂ CH ₃ - H | 61 |
| C ₂ H ₅ /sec-hexyl | (j) | 0.27 ± 0.01 | 25 | <i>n</i> -C ₆ H ₁₄ - H | 61 |
| C ₂ H ₅ /((CH ₃) ₂ CC ₃ H ₇) | | 0.74 ± 0.03 | 25 | (CH ₃) ₃ CHCH ₂ CH ₂ CH ₃ - H | 61 |
| C ₂ H ₅ /((CH ₃) ₂ CCH(CH ₃) ₂) | | 0.72 ± 0.01 | 25 | (CH ₃) ₂ CHCH(CH ₃) ₂ - H | 61 |
| C ₂ H ₅ /CH ₃ C(C ₂ H ₅) ₂ | | 0.8 | 25 | CH ₃ CH(C ₂ H ₅) ₂ - H | 61 |
| C ₂ H ₅ /CH ₃ CH ₂ O | | 1.3 ± 0.2 | 15-29 | C ₂ H ₅ COOC ₂ H ₅ P | 65 |
| Ethoxy | | | | | |
| CH ₃ CH ₂ O/C ₂ H ₅ | | 2.3 ± 0.3 | 29 | C ₂ H ₅ COOC ₂ H ₅ P | 65 |
| CH ₃ CH ₂ O/CH ₃ CH ₂ O | | 12 ± 2 | 25 | C ₂ H ₅ I + O ₂ P | 66 |
| Hydroxy ethyl | | | | | |
| CH ₃ CH(OH)/CH ₃ CH(OH) | (k) | 0.3 | 25 | C ₂ H ₅ OH - H | 67 |
| Halogenated ethyl | | | | | |
| C ₂ F ₅ /C ₂ H ₅ | | 0.56 | 21-240 | C ₂ F ₅ COC ₂ H ₅ P | 112 |
| C ₂ H ₄ Cl/C ₂ H ₅ | | 0.22 | 0-58 | C ₂ H ₄ + CCl ₄ P | 34 |

Disproportionation/Combination Ratios—Continued

| Reactants | Notes | <i>k</i> _{dis} / <i>k</i> _{comb.} | Temperature Range | Radical Source | Reference |
|--|---|---|-------------------|---|--|
| | | | °C | | |
| C ₂ H ₄ Cl/C ₂ H ₄ Cl | | ≤ 0.05 | 0–58 | C ₂ H ₄ + CCl ₄ P | 34 |
| | | ≤ 0.1 | 26 | C ₂ H ₄ + CCl ₄ P | 105 |
| | (l) | 0.36 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |
| | (m) | 24 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |
| | | | | | |
| C ₂ H ₄ Cl/C ₄ H ₈ Cl | (n) | 0.40 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |
| Allyl | | | | | |
| CH ₂ CHCH ₂ /C ₂ H ₅ | | 0.13 | 134–175 | (C ₂ H ₅) ₂ CO P | 106 |
| <i>n</i>-Propyl | | | | | |
| <i>n</i> -C ₃ H ₇ /C ₂ H ₅ | | 0.058 ± 0.01 | 61–129 | (C ₂ H ₅) ₂ CO + (<i>n</i> -C ₃ H ₇) ₂ N ₂ P | 8 |
| <i>n</i> -C ₃ H ₇ / <i>n</i> -C ₃ H ₇ | | 0.21 | 113 | (<i>n</i> -C ₃ H ₇) ₂ CO P | 69 |
| | | 0.1 | 25 | <i>n</i> -C ₃ H ₇ CHO P | 75 |
| | | 0.3 | 30–108 | (<i>n</i> -C ₃ H ₇) ₂ Hg P | 63 |
| | | 0.125 ± 0.01 | 100–150 | (<i>n</i> -C ₃ H ₇) ₂ CO P | 108 |
| | | 0.16 | 25–191 | <i>n</i> -C ₃ H ₇ CHO P | 70 |
| | | 0.157 | 25–130 | (<i>n</i> -C ₃ H ₇) ₂ N ₂ P | 71 |
| | | 0.14 | 74–178 | HCOOCH ₂ CH ₂ CH ₃ + CH ₃ | 11 |
| | | 0.141 ± 0.015 | 18–150 | (<i>n</i> -C ₃ H ₇) ₂ N ₂ P | 8 |
| | | 0.14 | | | |
| | | | | | |
| C ₂ H ₅ CD ₂ /C ₂ H ₅ CD ₂ | | 0.15 | 25–28 | CH ₃ CH ₂ CD ₂ COCD ₃ P | 14 |
| Iso-Propyl | | | | | |
| iso-C ₃ H ₇ /C ₂ H ₅ | | 0.19 | 48–112 | C ₂ H ₅ CHO + <i>i</i> -C ₃ H ₇ CHO P | 62 |
| | | 0.2 | room temp. | CH ₂ :CH ₂ + CH ₃ CH:CH ₂ + H | 107 |
| | | 0.07 | 34–144 | (C ₂ H ₅) ₂ CO P | 59 |
| | | 0.19 | | | |
| iso-C ₃ H ₇ /iso-C ₃ H ₇ | | 1.5 | 30 | CH ₃ CH:CH ₂ + H | 72 |
| | | 2. | 200 | CH ₃ CH:CH ₂ + H | 72 |
| | | 1. | 270 | (<i>i</i> -C ₃ H ₇) ₂ Hg T | 73 |
| | | 0.53 | 30 | (<i>i</i> -C ₃ H ₇) ₂ N ₂ P | 74 |
| | | 0.5 | 25 | <i>i</i> -C ₃ H ₇ CHO P | 75 |
| | | 0.77 | 25 | CH ₃ CH:CH ₂ + H | 76 |
| | | 0.48 | 320 | CH ₃ CH:CH ₂ + H | 76 |
| | | 0.6 | 200 | (<i>i</i> -C ₃ H ₇) ₂ CO P | 77 |
| | | 0.65 | 20–261 | <i>i</i> -C ₃ H ₇ CHO P | 109 |
| | | 0.5 ± 0.05 | room temp. | CH ₃ CH:CH ₂ + H | 56 |
| | | 0.54 | 61–127 | ((CH ₃) ₂ CH) ₂ N ₂ P | 78 |
| | | 0.65 | 94–181 | HCOO <i>i</i> -C ₃ H ₇ + CH ₃ | 11 |
| | | 0.58 ± 0.04 | 75–136 | ((CH ₃) ₂ CH) ₂ CO P | 79 |
| | | 0.65 | | | |
| | (CH ₃) ₂ CD/(CH ₃) ₂ CD | | 0.63 ± 0.04 | 25–125 | ((CH ₃) ₂ CD) ₂ CO P |
| | | 0.67 | 53–116 | (CH ₃) ₂ CDCOCD ₃ P | 16 |
| iso-C ₃ H ₇ /sec-C ₄ H ₉ | | 0.4 | room temp. | CH ₃ CH:CH ₂ + CH ₃ CH:CHCH ₃ + H | 107 |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | <i>k</i> _{dis} / <i>k</i> _{comb.} | Temperature Range | Radical Source | Reference |
|---|-------|---|-------------------|--|-----------|
| | | | °C | | |
| iso-C ₃ H ₇ / <i>t</i> -C ₄ H ₉ | | 0.5 | room temp. | CH ₃ CH:CH ₂ + (CH ₃) ₂ C:CH ₂ + H | 107 |
| | | 0.70 | 72–82 | (<i>t</i> -C ₄ H ₉) ₂ CO + (<i>i</i> -C ₃ H ₇) ₂ CO P | 62 |
| iso-C ₃ H ₇ /cyclo-C ₆ H ₇ | (i) | 0.52 ± 0.09 | 75–136 | (<i>i</i> -C ₃ H ₇) ₂ CO P | 79 |
| Propoxy | | | | | |
| C ₂ H ₅ CO/C ₂ H ₅ | | 0.4 ± 0.1 | 24 | (C ₂ H ₅) ₂ CO FP | 81 |
| <i>n</i>-Perfluoropropyl | | | | | |
| <i>n</i> -C ₃ F ₇ /C ₂ H ₅ | | 0.40 | 87–196 | (<i>n</i> -C ₃ F ₇) ₂ CO P | 82 |
| | (o) | 0.021 exp (2.2/RT) | 80–256 | <i>n</i> -C ₃ F ₇ COC ₂ H ₅ P | 83 |
| Butyls | | | | | |
| <i>n</i> -C ₄ H ₉ / <i>n</i> -C ₄ H ₉ | | 5. | 140 | (<i>n</i> -C ₄ H ₉) ₂ Hg P | 84 |
| | | 0.94 ± 0.05 | 70–195 | HCOO <i>n</i> -C ₄ H ₉ + CH ₃ | 13 |
| | | 0.95 ± 0.05 | 75–186 | HCOO <i>n</i> -C ₄ H ₉ + CH ₃ | 103 |
| | (p) | 10 ^{0.6} exp (–1.3/RT) | 61–229 | <i>n</i> -C ₄ H ₉ CHO P | 15 |
| | | 0.95 | | | |
| sec-C ₄ H ₉ / <i>i</i> -C ₃ H ₇ | (r) | 0.4 | room temp. | CH ₃ CH:CH ₂ + CH ₃ CH:CHCH ₃ + H | 107 |
| sec-C ₄ H ₉ /sec-C ₄ H ₉ | (q) | 1.64 | 25 | CH ₃ CH ₂ CH(CH ₃)CHO P | 85 |
| | | 2.27 | 100 | (CH ₃ CH ₂ CH(CH ₃)) ₂ CO P | 86 |
| | | 1.5 | room temp. | CH ₃ CH:CHCH ₃ + H | 110 |
| | | 0.95 | –103–+25 | cis-CH ₃ CH:CHCH ₃ + H | 17 |
| | | ≤ 1.2 | 107–168 | CH ₃ CH:CH ₂ + CH ₃ | 87 |
| | | 1.5 | 24 | CH ₃ CH ₂ CH:CH ₂ + H | 84 |
| | | 3.5 | 220 | CH ₃ CH ₂ CH:CH ₂ + H | 84 |
| | | 0.63 ± 0.08 | –78–+25 | trans-CH ₃ CH:CHCH ₃ + H | 88 |
| sec-C ₄ H ₉ / <i>t</i> -C ₄ H ₉ | (r) | 0.23 | room temp. | CH ₃ CH:CHCH ₃ + (CH ₃) ₂ C:CH ₂ + H | 107 |
| iso-C ₄ H ₉ /iso-C ₄ H ₉ | | 0.42 | 78–109 | ((CH ₃) ₂ CHCH ₂) ₂ CO P | 86 |
| | | 0.17 | 26–124 | (CH ₃) ₂ CHCH ₂ CHO P | 89 |
| | | 0.17 | | | |
| <i>t</i> -C ₄ H ₉ /C ₂ H ₅ | (r) | 1.7 | room temp. | CH ₂ :CH ₂ + (CH ₃) ₂ C:CH ₂ + H | 107 |
| | | 0.31 | 73–80 | (<i>t</i> -C ₄ H ₉) ₂ CO + (C ₂ H ₅) ₂ CO P | 62 |
| <i>t</i> -C ₄ H ₉ /iso-C ₃ H ₇ | (r) | 1.2 | room temp. | CH ₃ CH:CH ₂ + (CH ₃) ₂ C:CH ₂ + H | 107 |
| | | 0.67 | 72–82 | (<i>t</i> -C ₄ H ₉) ₂ CO + (<i>i</i> -C ₃ H ₇) ₂ CO P | 62 |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | $k_{dis}/k_{comb.}$ | Temperature Range | Radical Source | Reference |
|---|-------|---------------------|-------------------|---|-----------|
| | | | °C | | |
| $t\text{-C}_4\text{H}_9/\text{sec-}\text{C}_4\text{H}_9$ | (r) | 2.0 | room temp. | $\text{CH}_3\text{CH}:\text{CHCH}_3$ + $(\text{CH}_3)_2\text{C}:\text{CH}_2$ + H | 107 |
| $t\text{-C}_4\text{H}_9/t\text{-C}_4\text{H}_9$ | | 4.5 | 23 | $(\text{CH}_3)_2\text{C}:\text{CH}_2$ + H | 84 |
| | | 6.5 | 300 | $(\text{CH}_3)_2\text{C}:\text{CH}_2$ + H | 84 |
| | | 4.59 | 117–322 | $(t\text{-C}_4\text{H}_9)_2\text{CO}$ P | 86 |
| | | 7.4 | 20–60 | $(t\text{-C}_4\text{H}_9\text{O})_2$ P | 90 |
| | | 4.38 | 27–230 | $t\text{-C}_4\text{H}_9\text{CHO}$ P | 91 |
| | | 3.19 | 50–80 | $(t\text{-C}_4\text{H}_9)_2\text{CO}$ and $t\text{-C}_4\text{H}_9\text{CHO}$ P | 62 |
| | | 2.2 ± 0.3 | room temp. | $(\text{CH}_3)_2\text{C}:\text{CH}_2$ + H | 107 |
| | | 3.2 | | | |
| Monochlorobutyl | | | | | |
| $\text{C}_4\text{H}_8\text{Cl}/\text{C}_2\text{H}_4\text{Cl}$ | | 0.07 | 23 | $\text{CH}_2:\text{CH}_2 + \text{COCl}_2$ P | 68 |
| Pentyls | | | | | |
| $n\text{-C}_5\text{H}_{11}/n\text{-C}_5\text{H}_{11}$ | | 0.2 | 63 | $n\text{-C}_5\text{H}_{11}\text{COCH}_3$ P | 22 |
| $\text{C}_5\text{H}_{11}/\text{C}_5\text{H}_{11}$ | | 2.45 | 25 | iso- C_5H_{12} – H | 3 |
| | | ≥ 0.66 | 21 | $n\text{-C}_5\text{H}_{12}$ – H | 92 |
| $\text{cyclo-}\text{C}_5\text{H}_9/\text{cyclo-}\text{C}_5\text{H}_9$ | | 0.19 | 29 | cyclo- C_5H_{10} – H | 93 |
| | | 1.0 | 24 | cyclo- C_5H_{10} – H | 94 |
| | | 1.0 | 26–250 | cyclo- C_5H_{10} – H | 95 |
| | | 1.0 | | | |
| Hexyls | | | | | |
| $\text{cyclo-}\text{C}_6\text{H}_{11}/\text{cyclo-}\text{C}_6\text{H}_{11}$ | | 0.46 | 29 | cyclo- C_6H_{12} – H | 111 |
| $\text{cyclo-}\text{C}_6\text{H}_{11}/\text{cyclo-}\text{C}_6\text{H}_{11}$ | (s) | 0.42 | 29 | | 93 |
| $\text{C}_6\text{H}_{13}/\text{C}_6\text{H}_{13}$ | | 0.97 ± 0.08 | 25 | $n\text{-C}_6\text{H}_{14}$ – H | 4 |
| Amino | | | | | |
| NH_2/NH_2 | | 0.18 | no indication | NH_3 F.P. | 97 |
| Nitric Oxide | | | | | |
| $\text{NO}/\text{CH}_3\text{O}$ | (t) | 0.5 | 174 | | 96 |
| $\text{NO}/\text{CH}_3\text{CH}_2\text{O}$ | | 0.28 | 95–135 | $(\text{C}_2\text{H}_5\text{O})_2$ P | 96 |
| $\text{NO}/(\text{CH}_3)_2\text{CHO}$ | (u) | { 0.15 0.175 | 26 79 | $(i\text{-C}_3\text{H}_7\text{O})_2$ P | 30 30 |
| Iodine atom | | | | | |
| $\text{I}/\text{C}_2\text{H}_5$ | | 0.33 ± 0.03 | 28 | $(\text{C}_2\text{H}_5)_2\text{CO} + \text{C}_2\text{H}_5\text{I}$ P | 98 |

CHLORINE ATOM TRANSFER

| | | | | | |
|---|--|-------|--------|-----------------------------|----|
| Halogenated Methyl | | | | | |
| $\text{CF}_2\text{Cl}/\text{CF}_2\text{Cl}$ | | ≤ 0.5 | 75–119 | CF_2ClH – H | 99 |

Disproportionation/Combination Ratios – Continued

| Reactants | Notes | <i>k</i> dis/ <i>k</i> comb. | Temperature Range | Radical Source | Reference |
|--|---------|------------------------------|-------------------|---|-----------|
| °C | | | | | |
| Halogenated Ethyl | | | | | |
| C ₂ H ₄ Cl/C ₂ H ₄ Cl | (m) | ≤ 0.1 | 26 | C ₂ H ₄ + CCl ₄ P | 105 |
| | | 5.8 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |
| C ₂ F ₂ Cl ₃ /C ₂ F ₂ Cl ₃ | | ≤ 0.13 | 30–70 | CF ₂ :CCl ₂ + Cl ₂ P | 100 |
| C ₂ H ₄ Cl/C ₄ H ₈ Cl | (n) (v) | 0.5 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |
| Monochlorobutyl | | | | | |
| C ₄ H ₈ Cl/C ₂ H ₄ Cl | (n) (w) | 4.0 | 23 | C ₂ H ₄ + COCl ₂ P | 68 |

OXYGEN ATOM TRANSFER

| | | | | | |
|----------------------------------|---------|-----|-------|---------------------------------------|-----|
| Methyl | | | | | |
| CH ₃ /NO ₂ | | 2.0 | 55–90 | CH ₃ CHO + NO ₂ | 101 |
| | (x) (y) | 2.4 | 25 | CH ₃ NO ₂ P | 102 |
| | (x) (y) | 1.8 | 105 | CH ₃ NO ₂ P | 102 |

Notes

- (a) Authors consider this value to be too high.
- (b) Unpublished work by Thynne.
- (c) Author believes the radicals to be vibrationally excited.
- (d) A recalculation of data from Kerr and Trotman-Dickenson, *J. Chem. Soc.* 1609 (1960).
- (e) Calculated from data of Le Roy and Tickner, *J. Chem. Phys.* 19, 1247 (1951).
- (f) The temperature dependence gives either *Edis-Combs* = -0.3 kcal/mole or *k*dis/*k*comb = $T^{-0.7}$.
- (g) The ethylene formed was more than 90 percent C₂H₂D₂.
- (h) *Sec*-pentyl is a mixture of the radicals CH₃CH₂CHCH₂CH₃ and CH₃CHCH₂CH₂CH₃.
- (i) *Cyclo*-C₆H₇ is the cyclohexadienyl-1,4 radical.
- (j) *Sec*-hexyl is a mixture of the radicals CH₃CH-C₄H₉ and C₂H₅CHC₃H₇.
- (k) This value is probably low.
- (l) Combination product is 1,4-C₄H₈Cl₂.
- (m) Combination product is 1,3-C₄H₈Cl₂.
- (n) Combination product is 1,6-C₆H₁₂Cl₂.
- (o) Alternatively *k*dis/*k*comb = 0.33 with a large scatter.
- (p) Temperature coefficient very doubtful.
- (q) In the range 25–243 °C, the results fit the expression *k*dis/*k*comb = 0.715 exp (500/RT).
- (r) Very doubtful.
- (s) *Cyclo*-C₆H₁₁ is the methylcyclopentanyl radical.
- (t) Unpublished work by Phillips.
- (u) The ratio could be 0.16 within the experimental error of both results.
- (v) Products of the disproportionation reaction are C₄H₈ and 1,2-C₂H₄Cl₂.
- (w) Products of the disproportionation reaction are C₂H₄ and 1,4-C₄H₈Cl₂.
- (x) The "disproportionation reaction" is the sum of the rate constants for two reactions viz. (i) CH₃ + NO₂ = CH₃ONO, and (ii) CH₃ + NO₂ = CH₃O + NO.
- (y) Combination product is CH₃NO₂.

References

- (1) Heller and Gordon, *J. Chem. Phys.* 36, 2648 (1962).
- (2) Heller and Gordon, *J. Phys. Chem.* 64, 390 (1960).
- (3) Kuntz, *J. Phys. Chem.* 69, 2291 (1965).
- (4) Kuntz, *J. Phys. Chem.* 69, 4396 (1965).
- (5) Weir, *J. Chem. Soc.* 6870 (1965).
- (6) Ausloos and Steacie, *Can. J. Chem.* 33, 1062 (1955).
- (7) Thynne, *Trans. Faraday Soc.* 58, 676 (1962).
- (8) Grotewold and Kerr, *J. Chem. Soc.* 4337 (1963).
- (9) Heller, *J. Chem. Phys.* 28, 1255 (1958).
- (10) Wijnen, *J. Chem. Phys.* 22, 1631 (1954).
- (11) Thynne, *Trans. Faraday Soc.* 58, 1394 (1962).
- (12) Grotewold and Kerr, *J. Chem. Soc.* 4342 (1963).
- (13) Thynne, *Proc. Chem. Soc.* 18 (1961).
- (14) Ausloos and Murad, *J. Am. Chem. Soc.* 80, 5929 (1958).
- (15) Kerr and Trotman-Dickenson, *J. Chem. Soc.* 1609 (1960).
- (16) Heller, *J. Chem. Phys.* 35, 1711 (1961).
- (17) Rabinovitch and Diesen, *J. Chem. Phys.* 30, 735 (1959).
- (18) Miyoshi and Brinton, *J. Chem. Phys.* 36, 3019 (1962).
- (19) Frey, *Proc. Chem. Soc.* 385 (1959).
- (20) McMillan, *J. Am. Chem. Soc.* 82, 2422 (1960).
- (21) Blake, Henderson, and Kutschke, *Can. J. Chem.* 39, 519 (1961).
- (22) Wijnen, *J. Am. Chem. Soc.* 83, 3752 (1961).
- (23) Gordon, *Can. J. Chem.* 43, 570 (1965).
- (24) Toby and Kutschke, *Can. J. Chem.* 37, 672 (1959).
- (25) Wijnen, *J. Chem. Phys.* 27, 710 (1957).
- (26) Thynne and Gray, *Trans. Faraday Soc.* 59, 1149 (1963).
- (27) Wijnen, *J. Chem. Phys.* 28, 939 (1958).
- (28) Harrison and Lossing, *Can. J. Chem.* 37, 1478 (1959).
- (29) Darwent, Allard, Hartman, and Lange, *J. Phys. Chem.* 64, 1847 (1960).
- (30) McMillan, *J. Am. Chem. Soc.* 83, 3018 (1961).
- (31) Dever and Calvert, *J. Am. Chem. Soc.* 84, 1362 (1962).
- (32) Heicklen and Johnston, *J. Am. Chem. Soc.* 84, 4030 (1962).
- (33) Gregory and Wijnen, *J. Chem. Phys.* 38, 2925 (1963).
- (34) Rocquitte and Wijnen, *J. Am. Chem. Soc.* 85, 2053 (1963).
- (35) Bellas, Strausz, and Gunning, *Can. J. Chem.* 43, 1022 (1965).
- (36) Sherwood and Gunning, *J. Phys. Chem.* 69, 2323 (1965).

- (37) James and Troughton, *Trans. Faraday Soc.* **62**, 145 (1966).
- (38) Ivin and Steacie, *Proc. Roy. Soc.* **208A**, 25 (1951).
- (39) Kutschke, Wijnen, and Steacie, *J. Am. Chem. Soc.* **74**, 714 (1952).
- (40) Moore and Taylor, *J. Chem. Phys.* **8**, 504 (1940).
- (41) Le Roy and Khan, *J. Chem. Phys.* **15**, 816 (1947).
- (42) Blacet and Pitts, *J. Am. Chem. Soc.* **74**, 3382 (1952).
- (43) Gowenlock, Polanyi, and Warhurst, *Proc. Roy. Soc.* **218A**, 269 (1953).
- (44) Volman and Brinton, *J. Chem. Phys.* **22**, 929 (1954).
- (45) Ausloos and Steacie, *Bull. Soc. Chim. Belges.* **63**, 87 (1954).
- (46) Brinton and Steacie, *Can. J. Chem.* **33**, 1840 (1955).
- (47) Smith, Beatty, Pindar, and Le Roy, *Can. J. Chem.* **33**, 821 (1955).
- (48) James and Steacie, *Proc. Roy. Soc.* **244A**, 289 (1958).
- (49) Cerfontain and Kutschke, *Can. J. Chem.* **36**, 344 (1958).
- (50) Kerr and Trotman-Dickenson, *J. Chem. Soc.* 1611 (1960).
- (51) Dixon, Stefani, and Szwarc, *J. Am. Chem. Soc.* **85**, 2551 (1963).
- (52) Matsuoka, Dixon, Stefani, and Szwarc, *Proc. Chem. Soc.* 304 (1962).
- (53) Dingley and Calvert, *J. Am. Chem. Soc.* **85**, 856 (1963).
- (54) Brown and James, *Can. J. Chem.* **43**, 660 (1965).
- (55) James and Troughton, *Trans. Faraday Soc.* **62**, 120 (1966).
- (56) Boddy and Robb, *Proc. Roy. Soc.* **249A**, 518 (1959).
- (57) Wijnen and Steacie, *Can. J. Chem.* **29**, 1092 (1951).
- (58) Boddy and Steacie, *Can. J. Chem.* **38**, 1576 (1960).
- (59) Thynne, *Proc. Chem. Soc.* 68 (1961).
- (60) Holroyd and Pierce, *J. Phys. Chem.* **68**, 1392 (1964).
- (61) Holroyd and Klein, *J. Phys. Chem.* **67**, 2273 (1963).
- (62) Dominguez, Kerr, and Trotman-Dickenson, *J. Chem. Soc.* 3357 (1962).
- (63) Caule and Steacie, *Can. J. Chem.* **29**, 103 (1951).
- (64) Brown and James, *Can. J. Chem.* **43**, 660 (1965).
- (65) Wijnen, *J. Am. Chem. Soc.* **82**, 3034 (1960).
- (66) Heicklen and Johnston, *J. Am. Chem. Soc.* **84**, 4030 (1962).
- (67) Knight and Gunning, *Can. J. Chem.* **39**, 2466 (1961).
- (68) Heicklen, *J. Am. Chem. Soc.* **87**, 445 (1965).
- (69) Masson, *J. Am. Chem. Soc.* **74**, 4731 (1952).
- (70) Kerr and Trotman-Dickenson, *Trans. Faraday Soc.* **55**, 572 (1959).
- (71) Kerr and Calvert, *J. Am. Chem. Soc.* **83**, 3391 (1961).
- (72) Moore, *J. Chem. Phys.* **16**, 916 (1948).
- (73) Chilton and Gowenlock, *Trans. Faraday Soc.* **49**, 1451 (1953).
- (74) Durham and Steacie, *Can. J. Chem.* **31**, 377 (1953).
- (75) Blacet and Calvert, *J. Am. Chem. Soc.* **73**, 661 (1951).
- (76) Hoey and Le Roy, *Can. J. Chem.* **33**, 580 (1955).
- (77) Heller and Gordon, *J. Phys. Chem.* **60**, 1315 (1956).
- (78) Riem and Kutschke, *Can. J. Chem.* **38**, 2332 (1960).
- (79) James and Suart, *J. Am. Chem. Soc.* **86**, 5424 (1964).
- (80) Heller and Gordon, *J. Phys. Chem.* **62**, 709 (1958).
- (81) Fischer and Mains, *J. Phys. Chem.* **68**, 188 (1964).
- (82) Giacometti and Steacie, *Can. J. Chem.* **36**, 1493 (1958).
- (83) Pritchard and Thommarson, *J. Phys. Chem.* **69**, 1001 (1965).
- (84) Moore and Wall, *J. Chem. Phys.* **17**, 1325 (1949).
- (85) Gruver and Calvert, *J. Am. Chem. Soc.* **78**, 5208 (1956).
- (86) Kraus and Calvert, *J. Am. Chem. Soc.* **79**, 5921 (1957).
- (87) Miyoshi and Brinton, *J. Chem. Phys.* **36**, 3019 (1962).
- (88) Kubin, Rabinovitch, and Harrington, *J. Chem. Phys.* **37**, 937 (1962).
- (89) Metcalfe and Trotman-Dickenson, *J. Chem. Soc.* 5072 (1960).
- (90) Frey, *Proc. Chem. Soc.* 385 (1959).
- (91) Birrell and Trotman-Dickenson, *J. Chem. Soc.* 4218 (1960).
- (92) Back, *Trans. Faraday Soc.* **54**, 512 (1958).
- (93) Beck, Kniebes, and Gunning, *J. Chem. Phys.* **22**, 678 (1954).
- (94) Stock and Gunning, *Can. J. Chem.* **38**, 2295 (1960).
- (95) Gunning and Stock, *Can. J. Chem.* **42**, 357 (1964).
- (96) Arden, Phillips, and Shaw, *J. Chem. Soc.* 5126 (1964).
- (97) Salzman and Bair, *J. Chem. Phys.* **41**, 3654 (1964).
- (98) Guercione and Wijnen, *J. Chem. Phys.* **38**, 1 (1963).
- (99) Bellas, Strausz, and Gunning, *Can. J. Chem.* **43**, 1022 (1965).
- (100) Vallana, Castellano, and Schumacher, *Z. Phys. Chem.* **46**, 294 (1965).
- (101) Phillips and Shaw, 10th Int. Comb. Symp., p. 423, Combustion Institute, Pittsburgh (1965).
- (102) Rebbert and Slagg, *Bull. Soc. Chim. Belges* **71**, 709 (1962).
- (103) Thynne, *Trans. Faraday Soc.* **58**, 1533 (1962).
- (104) Wijnen, *J. Chem. Phys.* **28**, 271 (1958).
- (105) Roquette and Wijnen, *J. Chem. Phys.* **38**, 4 (1963).
- (106) James and Troughton, *Trans. Faraday Soc.* **62**, 145 (1966).
- (107) Boddy and Robb, *Proc. Roy. Soc.* **249A**, 547 (1959).
- (108) Whiteway and Masson, *J. Chem. Phys.* **25**, 233 (1956).
- (109) Kerr and Trotman-Dickenson, *Trans. Faraday Soc.* **55**, 921 (1959).
- (110) Boddy and Robb, *Proc. Roy. Soc.* **249A**, 532 (1959).
- (111) Beck, Kniebes, and Gunning, *J. Chem. Phys.* **22**, 672 (1954).
- (112) Thommarson and Pritchard, *J. Phys. Chem.* **70**, 2307 (1966).

Molecule-Molecule Reactions (Hydrogen and Hydrogen Halides)

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|---------|---------------------------------|--|------------------------------|--|------------------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| H ₂ + O ₂ = 2OH | | 45.0 67.0 | (<i>k</i> = 84.0) 14.4 | 485 687-807 | H ₂ /O ₂ ignition H ₂ /O ₂ S.T. | 1 2 |
| H ₂ + I ₂ = 2HI | (a) | 39.0 41.0 | 14.1 15.09 | 10-235 598-774 | HI T H ₂ /I ₂ T | 3 9, 10 |
| | | | <i>k</i> = 6.60 ± 0.09 | 394 | H ₂ /I ₂ T | 11 |
| | | 41.24 ± 0.25 | 14.35 ± 0.07 | 105-360 | H ₂ /I ₂ T | 12 |
| D ₂ + I ₂ = 2DI | (b) (c) | 41.47 ± 0.20 | 14.14 ± 0.07 | 360-465 | D ₂ /I ₂ T | 13 |
| H ₂ + HI = H ₂ ^o + HI | | 44 ± 3 ≥ 47 | 13.7 | 420-480 420-480 | H ₂ /I ₂ /HI T H ₂ /I ₂ /HI T | 14 17 |
| H ₂ + NO ₂ = NO + H ₂ O | | 18 | 9.3 | 384-434 | H ₂ /O ₂ /NO ₂ T | 15 |
| HCl + NO ₂ = HNO ₂ + Cl | | 23.4 | 11.6 | 100-420 | HCl/NO ₂ T | 23 |
| HBr + NO ₂ = HNO ₂ + Br | | 13.0 | 11.0 | 180-310 | HBr/NO ₂ T | 23 |
| HBr + C ₂ F ₄ = C ₂ F ₄ Br + H | | 54.6 | 16.96 | 858-1013 | H ₂ /O ₂ /C ₂ F ₄ Br ₂ S.T. | 22 |
| 2HI = H ₂ + I ₂ | (d) (a) | 44.0 49.2 44.20 ± 0.25 | 13.9 15.56 13.54 ± 0.07 | 10-235 597-774 105-360 | HI T HI T H ₂ /I ₂ T | 3 9, 10 12 |
| HI + CH ₃ I = CH ₄ + I ₂ | (e) | 33.4 | 14.3 | 270-320 | CH ₃ I/HI T | 19 |
| DI + CH ₃ I = CH ₃ D + I ₂ | | | <i>k</i> = 1.54 | 250 | CH ₃ I/HI/DI T | 20 |
| HI + C ₂ H ₅ I = C ₂ H ₆ + I ₂ | (e) | 29.8 | 13.7 | 250-300 | C ₂ H ₅ I/HI T | 19 |
| HI + <i>n</i> -C ₃ H ₇ I = C ₃ H ₈ + I ₂ | (e) | 19.2 | 14.1 | 260-300 | <i>n</i> -C ₃ H ₇ I/HI T | 19 |

Ratios of Rate Constants

| Reactions | Notes | <i>E</i> ₁ - <i>E</i> ₂ | $\log_{10} A_1/A_2$ | <i>k</i> ₁ / <i>k</i> ₂ | Temperature range | Radical source | Reference |
|--|-------|---|---------------------|---|-------------------|---|-----------|
| | | <i>(kcal mole⁻¹)</i> | | | °C | | |
| (1) HBr + CH ₂ N ₂ = CH ₃ Br + N ₂ (2) HCl + CH ₂ N ₂ = CH ₃ Cl + N ₂ | | -1.5 | -0.091 | | -78-27 | CH ₂ N ₂ /HBr/HCl T | 21 |
| (1) HI + CH ₂ N ₂ = CH ₃ I + N ₂ (2) HBr + CH ₂ N ₂ = CH ₃ Br + N ₂ | | | | 4 ± 2 | 27 | CH ₂ N ₂ /HI/HBr T | 21 |
| (1) HI + CH ₂ N ₂ = CH ₃ I + N ₂ (2) HCl + CH ₂ N ₂ = CH ₃ Cl + N ₂ | | | | 50 ± 15 | 27 | CH ₂ N ₂ /HI/HCl T | 21 |

Notes

(a) These values summarise a considerable body of early data (references 4 to 8).

(b) For data on the reaction $\text{HD} + \text{I}_2 = \text{HI} + \text{DI}$, see reference 16.

(c) For a summary of early data see reference 3.

(d) For data on the reaction $2\text{DI} = \text{D}_2 + \text{I}_2$ see references 13 and 18.

(e) The A factor values originally reported (reference 19) are too large by a factor of ten (reference 20).

References

1. Semenov, *Acta Physicochimica U.S.S.R.* **20**, 291 (1945).
2. Skinner and Ringrose, *J. Chem. Phys.* **42**, 2190 (1965).
3. Trotman-Dickenson, "Gas Kinetics," p. 255 et seq. (Butterworths, London, 1955).
4. Bodenstein, *Z. Phys. Chem.* **13**, 56 (1894).
5. Bodenstein, *Z. Phys. Chem.* **22**, 1 (1897).
6. Bodenstein, *Z. Phys. Chem.* **29**, 295 (1899).
7. Kassel, *Proc. Nat. Acad. Sci. Wash.* **16**, 358 (1930).
8. Kistiakowsky, *J. Am. Chem. Soc.* **50**, 2315 (1928).
9. Graven, *J. Am. Chem. Soc.* **78**, 3297 (1956).
10. Horie, Ishii, and Amano, *J. Phys. Chem.* **68**, 1264 (1964).
11. Sullivan, *J. Chem. Phys.* **36**, 1925 (1962).
12. Sullivan, *J. Chem. Phys.* **30**, 1292 (1959).
13. Sullivan, *J. Chem. Phys.* **39**, 3001 (1963).
14. Steiner, *Trans. Faraday Soc.* **36**, 1111 (1940).
15. Crist and Wertz, *J. Chem. Phys.* **7**, 719 (1939).
16. Magee, *J. Am. Chem. Soc.* **79**, 5375 (1957).
17. Sullivan, *J. Chem. Phys.* **30**, 1577 (1959).
18. Taylor and Crist, *J. Am. Chem. Soc.* **63**, 1377 (1941).
19. Ogg, *J. Am. Chem. Soc.* **56**, 526 (1934).
20. Newton, *J. Chem. Phys.* **18**, 797 (1950).
21. Hassler and Setser, *J. Am. Chem. Soc.* **87**, 3793 (1965).
22. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
23. Rosser and Wise, *J. Phys. Chem.* **64**, 602 (1960).

Molecule-Molecule Reactions (Hydrocarbons and Hydrocarbon Derivatives)

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--|-------|------------------------------------|---|-------------------|--|-----------|
| Alkanes | | | | | | |
| $\text{CH}_4 + \text{O}_3 = \text{products}$ | | (<i>kcal mole</i> ⁻¹) | (<i>cm</i> ³ <i>mole</i> ⁻¹ <i>sec</i> ⁻¹) | °C | | |
| | | 15.35 | 11.21 | 35-64 | $\text{CH}_4/\text{O}_3/\text{O}_2$ T | 1 |
| | | 13.90 | 11.15 | 35-64 | CH_4/O_3 T | 1 |
| $\text{C}_2\text{H}_6 + \text{C}_2\text{H}_4 = 2\text{C}_2\text{H}_5$ | | 68.9 | 17.6 | 752-917 | $\text{H}_2/\text{O}_2/\text{C}_2\text{H}_4$ S.T. | 2 |
| Alkenes and aromatics | | | | | | |
| $\text{CH}_2:\text{CHCH}:\text{CH}_2$ + $\text{C}_2\text{N}_2 = \text{H}_2 + 2$ - cyano pyridine | | 31.6 | 12.20 | 325-450 | $\text{C}_2\text{N}_2/\text{C}_4\text{H}_6$ T | 3 |
| $\text{CH}_2:\text{CHCH}:\text{CH}_2$ + $\text{CF}_3\text{CN} = \text{H}_2 + 2$ - trifluoromethyl pyridine | | 21.5 | 9.32 | 350-520 | $\text{CF}_3\text{CN}/\text{C}_4\text{H}_6$ T | 4 |
| $\text{CH}_2:\text{CHCH}:\text{CH}_2$ + $\text{CF}_3\text{CF}_2\text{CN} = \text{H}_2$ + 2-pentafluoro- ethyl pyridine | | 25.0 | 10.42 | 340-460 | $\text{C}_2\text{F}_5\text{CN}/\text{C}_4\text{H}_6$ T | 5 |
| $\text{CH}_2:\text{CHCH}:\text{CH}_2$ + $\text{CF}_3\text{CF}_2\text{CF}_2\text{CN}$ = $\text{H}_2 + 2$ -perfluoro- propyl pyridine | | 25.8 | 10.51 | 340-460 | $\text{C}_3\text{F}_7\text{CN}/\text{C}_4\text{H}_6$ T | 5 |
| $2\text{C}_6\text{H}_6 = \text{C}_{12}\text{H}_{10} + \text{H}_2$ | | 40 | 14.0 | 900-1210 | C_6H_6 T | 6 |
| Alkynes | | | | | | |
| $2\text{C}_2\text{H}_2 = \text{products}$ | (b) | 50.2 | 16.57 | 352-472 | C_2H_2 T | 7 |
| $2\text{C}_2\text{H}_2 = \text{C}_4\text{H}_2 + \text{H}_2$ | (b) | ~ 30 | | | C_2H_2 S.T. | 8 |
| $2\text{C}_2\text{H}_2 = \text{C}_4\text{H}_2 + \text{H}_2$ | (b) | 36.4 ± 3 | 12.89 | 1127-2227 | C_2H_2 S.T. | 9 |
| $2\text{C}_2\text{H}_2 = \text{C}_2\text{H}_2^* + \text{C}_2\text{H}_2$ | (b) | 50 | 14.04 | 1227-2227 | C_2H_2 S.T. | 10 |
| $2\text{C}_2\text{H}_2 = \text{C}_4\text{H}_3 + \text{H}$ | (b) | 38.8 ± 7.5 | 13.47 ± 0.78 | 1630-1922 | C_2H_2 S.T. | 11 |
| | (a) | 41.6 ± 0.6 | 13.77 ± 0.11 | 347-2177 | | 11 |
| Hydrocarbon derivatives | | | | | | |
| $\text{CH}_3\text{CHO} + \text{O}_2$ = $\text{HO}_2 + \text{CH}_3\text{CO}$ | | 29 ± 2 | | 320-380 | $\text{CH}_3\text{CHO}/\text{O}_2$ T | 12 |
| $\text{CH}_3\text{I} + \text{I}^* - \text{I}$ = $\text{CH}_3\text{I}^* + \text{I}_2$ | (c) | 9.0 | 6.40 | 60-140 | $\text{CH}_3\text{I}/\text{I}_2$ T | 13 |
| $\text{CH}_3\text{OCH}_3 + \text{NO}$ = HNO + CH_2OCH_3 | | 43.4 | 14.0 | 500-600 | $\text{CH}_3\text{OCH}_3/\text{NO}$ T | 14 |

Notes

- (a) Critical review of literature data.
- (b) These values are included in the review estimate (a), the products of the reaction being considered to be $C_4H_3 + H$.
- (c) These values may be low because of complicating heterogeneous reactions.

References

1. Dilleuth, Skidmore, and Schubert, *J. Phys. Chem.* **64**, 1496 (1960).
2. Skinner and Ringrose, *J. Chem. Phys.* **43**, 4129 (1965).
3. Hawkins and Janz, *J. Am. Chem. Soc.* **74**, 1790 (1952).
4. Jarvie and Janz, *J. Phys. Chem.* **60**, 1430 (1956).
5. Monahan and Janz, *J. Phys. Chem.* **69**, 1070 (1965).
6. Hou and Palmer, *J. Phys. Chem.* **69**, 863 (1965).
7. Silcocks, *Proc. Roy. Soc.* **242A**, 411 (1957).
8. Greene, Taylor, and Patterson, *J. Phys. Chem.* **62**, 238 (1958).
9. Aten and Greene, *Comb and Flame* **5**, 55 (1961).
10. Palmer and Dormish, *J. Phys. Chem.* **68**, 1553 (1964).
11. Gay, Kistiakowsky, Michael, and Niki, *J. Chem. Phys.* **43**, 1720 (1965).
12. Sokolova, Markevich, and Nalbandyan, *Russ. J. Phys. Chem. (Eng. Trans.)* **35**, 415 (1961).
13. Schmied and Fink, *J. Chem. Phys.* **27**, 1034 (1957).
14. McKenney, Wojciechowski, and Laidler, *Can. J. Chem.* **41**, 1993 (1963).

Molecule-Molecule Reactions Involving Nitric Oxide

| Reaction | Notes | E | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|-----------------------|--|-------------------|------------------------------|-----------|
| | | ($kcal\ mole^{-1}$) | ($cm^3\ mole^{-1}\ sec^{-1}$) | $^{\circ}C$ | | |
| $NO + CH_3OCH_3 = HNO + CH_2OCH_3$ | | 43.4 | 14.0 | 500-600 | CH_3OCH_3/NO T | 1 |
| $NO + CCl_3CHO = HNO + CCl_3CO$ | | 37.1 | 13.5 | 356-421 | CCl_3CHO/NO T | 20, 21 |
| $NO + F_2 = ONF + F$ | | 1.5 ± 1.0 | 11.78 | -78-+27 | NO_2/F_2 D.F. | 2 |
| $NO + XeF_4 = ONF + XeF_3$ | | 7 to 9 | | 27-77 | XeF_4/NO T | 3 |
| $NO + XeF_2 = ONF + XeF$ | | 10 to 12 | | 27-77 | XeF_4/NO T | 3 |
| $NO + Cl_2 = NOCl + Cl$ | | | $k = (2.7 \pm 1.0) \times 10^2$ $k = (8.3 \pm 3.0) \times 10^2$ $k = (10.0 \pm 4.0) \times 10^2$ | 158 178 198 | } $NOCl$ T | 4 |
| | (b) | 20.3 | 12.6 | 25-55 | | |
| $NO + NO_2Cl = NO_2 + NOCl$ | | 6.9 | 11.92 | 1-71 | NO_2Cl/NO T | 19 |
| $NO + O_3 = NO_2 + O_2$ | | 2.5 ± 0.3 | 11.90 | -75--43 | NO/O_3 T | 5 |
| | | | $k = 3 \times 10^{10}$ | 29 | NO/O_3 T | 6 |
| | (a) | 2.55 ± 0.22 | 12.08 | -28-+78 | NO/O_3 | 7 |
| | | | $k = 7.6 \times 10^9$ | 60 | NO/O_3 T | 8 |
| | | | $k = 1.5 \times 10^{10}$ | 25 | NO_2/O_2 discharge | 9 |
| | | | $k = (1.26 \pm 0.30) \times 10^{10}$ | 25 | O_3/N_2 discharge | 10 |
| | | 2.46 ± 0.15 | 11.76 | -57-+49 | NO/O_3 T | 11 |
| $NO + NO = N_2 + O_2$ | | 63.8 | 12.42 | 1062-1257 | NO T | 12, 13 |
| | | 63.1 | 13.49 | 1400-1800 | NO T | 14 |
| | | 57.0 | 12.91 | 2730-4030 | NO S.T. | 15 |
| $NO + N_2O = N_2 + NO_2$ | | 50.0 | 14.3 | 1227-1927 | NO/N_2O S.T. | 16 |
| | | 50.0 | 14.4 | 700-760 | | 17 |
| $CH_3CH_2O^{15}NO + ^{14}NO = CH_3CH_2O^{14}NO + ^{15}NO$ | | 21.4 ± 0.5 | 11.48 | 130-150 | $CH_3CH_2O^{15}NO/^{14}NO$ T | 22 |
| $CF_3CH_2O^{15}NO + ^{14}NO = CF_3CH_2O^{14}NO + ^{15}NO$ | | 23.7 ± 0.5 | 12.78 | 130-150 | $CF_3CH_2O^{15}NO/^{14}NO$ T | 22 |

Notes

- (a) Reaction carried out using supersonic nozzle technique.
- (b) Estimated from the reverse reaction.

References

1. McKenney, Wojciechowski, and Laidler, *Can. J. Chem.* **41**, 1993 (1963).
2. Rapp and Johnston, *J. Chem. Phys.* **33**, 695 (1960).
3. Johnston and Woolfolk, *J. Chem. Phys.* **41**, 269 (1964).
4. Ashmore and Spencer, *Trans. Faraday Soc.* **55**, 1868 (1959).
5. Johnston and Crosby, *J. Chem. Phys.* **22**, 689 (1954).
6. Ford, Doyle, and Endow, *J. Chem. Phys.* **26**, 1337 (1957).
7. Marte, Tschuikow-Roux, and Ford, *J. Chem. Phys.* **39**, 3227 (1963).
8. Borok, *Zhur. Fiz. Khim.* **35**, 2275 (1961).
9. Phillips and Schiff, *J. Chem. Phys.* **37**, 924 (1962).
10. Phillips and Schiff, *J. Chem. Phys.* **36**, 1509 (1962).
11. Clyne, Thrush, and Wayne, *Trans. Faraday Soc.* **60**, 359 (1964).
12. Kaufman and Kelso, *J. Chem. Phys.* **23**, 1702 (1955).
13. Kaufman and Decker, 7th Int. Comb. Symp., p. 57 (Butterworths, London, 1959).
14. Yuan, Slaughter, Koerner, and Daniels, *J. Phys. Chem.* **63**, 952 (1959).
15. Freedman and Daiber, *J. Chem. Phys.* **34**, 1271 (1961).
16. Fishburne and Edse, *J. Chem. Phys.* **41**, 1297 (1964).
17. Kaufman and Kelso, *J. Chem. Phys.* **23**, 602 (1955).
18. Ashmore and Chanmugam, *Trans. Faraday Soc.* **49**, 270 (1953).
19. Freiling, Johnston, and Ogg, *J. Chem. Phys.* **20**, 327 (1952).
20. Verhoek, *Trans. Faraday Soc.* **31**, 1521 (1935).
21. Trotman-Dickenson, "Gas Kinetics," p. 212 (Butterworths, London, 1955).
22. Kuhn and Günthard, *Helv. Chim. Acta.* **43**, 607 (1960).

Nitrogen Dioxide (Molecule-Molecule Reactions)

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|---|-------|---------------------------------|--|-------------------|---|-----------|
| | | <i>(kcal mole⁻¹)</i> | <i>(cm³ mole⁻¹ sec⁻¹)</i> | °C | | |
| NO ₂ + H ₂ = NO + H ₂ O | | 18. | 9.3 | 384-434 | H ₂ /O ₂ /NO ₂ T | 19 |
| NO ₂ + HCHO = HNO ₂ + HCO | | 15.1 | 10.1 | 118-160 | HCHO/NO ₂ T | 20 |
| NO ₂ + CHOCHO = HNO ₂ + COCHO | | 19.8 | 11.9 | 160-210 | CHOCHO/NO ₂ T | 21 |
| NO ₂ + CH ₃ CHO = HNO ₂ + CH ₃ CO | | 16.0 | 12.9 | 118-143 | CH ₃ CHO/NO ₂ T | 22 |
| NO ₂ + NH ₃ = HNO ₂ + NH ₂ | | 27.5 | 12.7 | 327-527 | NH ₃ /NO ₂ T | 1 |
| NO ₂ + HCl = HNO ₂ + Cl | | 23.4 | 11.6 | 100-420 | HCl/NO ₂ T | 2 |
| NO ₂ + HBr = HNO ₂ + Br | | 13.0 | 11.0 | 180-310 | HBr/NO ₂ T | 2 |
| NO ₂ + F ₂ = NO ₂ F + F | | 10.5 | 12.2 | 28-70 | NO ₂ /F ₂ T | 7 |
| NO ₂ + F ₂ O = NO ₂ F + FO | | 14.48 | 11.11 | 60-80 | F ₂ O/NO ₂ T | 18 |
| NO ₂ + Cl ₂ O = NO ₂ Cl + OCl | | 11.6 | 10.64 | 20-60 | NO ₂ /Cl ₂ O T | 3 |
| NO ₂ + ClO ₂ = NO ₃ + OCl | | 11.5 | 10.68 | | | 3 |
| NO ₂ + NOCl = NO + NO ₂ Cl | | 10.0 | 10.28 | -20-+8 | NOCl/ClO ₂ T | 4 |
| NO ₂ + CO = CO ₂ + NO | | 31.6 | 13.08 | 267-454 | CO/NO ₂ T | 5 |
| | | 27.8 | 11.68 | 225-290 | CO/NO ₂ T | 6 |
| NO ₂ + O ₃ = NO ₃ + O ₂ | | 7.0 | 12.77 | 13-29 | O ₃ /NO ₂ T | 8 |
| | | | $k = 1.96 \times 10^7$ | room temp. | O ₃ /NO ₂ T | 9 |
| 2NO ₂ = NO + NO ₃ | | | $k = 2.6 \times 10^4$ | 434 | NO ₂ T | 10 |
| | (a) | 23.90 ± 0.60 | 11.89 ± 0.25 | 200-430 | NO ₂ T | 11 |
| | | 23. | | | | 12 |
| 2NO ₂ = 2NO + O ₂ | (b) | 27.10 | 12.69 | 320-380 | NO ₂ T | 13, 11 |
| | | 26.90 | 12.60 | 357-747 | NO ₂ T | 14, 11 |
| | | 26.90 ± 0.10 | 12.60 ± 0.04 | 200-550 | NO ₂ T | 11, 10 |
| 2NO ₂ = products | (c) | 25.6 | 11.8 | 319-383 | | 15 |
| | (c) | 25.0 ± 5.0 | 13.40 | 1127-2027 | NO ₂ S.T. | 16 |
| | (c) | 25.7 | 12.95 | 1227-1827 | NO ₂ S.T. | 17 |

Notes

- (a) Estimated from the reverse reaction.
- (b) These values were recalculated from the original data (reference 11).
- (c) These values represent the sums of the contributions from $2\text{NO}_2 = 2\text{NO} + \text{O}_2$ and $2\text{NO}_2 = \text{NO} + \text{NO}_3$.

References

1. Rosser and Wise, *J. Chem. Phys.* **25**, 1078 (1956).
2. Rosser and Wise, *J. Phys. Chem.* **64**, 602 (1960).
3. Martin, Meise, and Engelmann, *Z. Phys. Chem.* **24**, 285 (1960).
4. Martin and Kohnlein, *Z. Phys. Chem.* **17**, 375 (1958).
5. Johnston, Bonner, and Wilson, *J. Chem. Phys.* **26**, 1002 (1957).
6. Brown and Crist, *J. Chem. Phys.* **9**, 840 (1941).
7. Perrine and Johnston, *J. Chem. Phys.* **21**, 2202 (1953).
8. Johnston and Yost, *J. Chem. Phys.* **17**, 386 (1949).
9. Ford, Doyle, and Endow, *J. Chem. Phys.* **26**, 1336 (1957).
10. Ashmore and Levitt, *Research* **9**, s25 (1956).
11. Ashmore and Burnett, *Trans. Faraday Soc.* **58**, 253 (1962).
12. Schott and Davidson, *J. Am. Chem. Soc.* **80**, 1841 (1958).
13. Bodenstein and Ramstetter, *Z. Phys. Chem.* **100**, 106 (1922).
14. Rosser and Wise, *J. Chem. Phys.* **24**, 493 (1956).
15. Kassel, "The Kinetics of Homogeneous Gas Reactions," p. 156 (Chemical Catalog Co., N.Y., 1932).
16. Huffman and Davidson, *J. Am. Chem. Soc.* **81**, 2311 (1959).
17. Fishburne, Bergbauer, and Edse, *J. Chem. Phys.* **43**, 1847 (1965).
18. De Staricco, Sicre, and Schumacher, *Z. Phys. Chem.* **39**, 337 (1963).
19. Crist and Wertz, *J. Chem. Phys.* **7**, 719 (1939).
20. Pollard and Wyatt, *Trans. Faraday Soc.* **45**, 760 (1949).
21. Thomas, *Trans. Faraday Soc.* **49**, 630 (1953).
22. McDowell and Thomas, *Trans. Faraday Soc.* **46**, 1030 (1950).

Molecule-Molecule Reactions (Inorganic Molecules)

| Reactions | Notes | <i>E</i> | $\log_{10} A$ | Temperature range | Radical source | Reference |
|--------------------------------------|-------|---|---|-------------------|--|------------|
| $F_2 + CO = FCO + F$ | | <i>(kcal mole⁻¹)</i> 13.5 ± 1.2 | <i>(cm³ mole⁻¹ sec⁻¹)</i> 11.67 | °C 15-45 | F ₂ /CO/O ₂ T | 1, 2 |
| $F_2 + ClO_2 = FClO_2 + F$ | | 8.5 ± 0.4 | 10.6 | -46--26 | F ₂ /ClO ₂ T | 3 |
| | | 8.0 ± 0.5 | 10.11 | -46--26 | F ₂ /ClO ₂ T | 4 |
| $Cl_2 + O_3 = ClO + ClO_2$ | | 26.0 ± 1 | | 35-60 | Cl ₂ /O ₃ T | 5 |
| $2ClCN = Cl_2 + C_2N_2$ | | 60. | 13. | 1727-2527 | ClCN S.T. | 6 |
| $2NOCl = Cl_2 + 2NO$ | | 23.4 | 12.5 | 150-250 | | 7 |
| | | 23.6 ± 0.7 | 12.7 ± 0.3 | 150-411 | NOCl T | 8 |
| $2NOI = I_2 + 2NO$ | | | $k \leq 4 \times 10^{10}$ | 60 | NO/I ₂ F.P. | 15 |
| $2O_3 = 3O_2$ | | 18.8 | 12.65 | 70-100 | O ₃ T | 9 |
| $CO + O_2 = CO_2 + O$ | | 51.0 ± 7.0 | 12.54 ± 0.22 | 2127-2727 | CO/O ₂ S.T. | 10 |
| $CO + B_2H_6 = BH_3 + BH_3CO$ | (a) | 17.65 | 11.32 | 0-60 | | 11 |
| $BF_3 + BCl_3 = BF_2Cl + BCl_2$ | | | $k \approx 5 \times 10^3$ | 23 | BF ₃ /BCl ₃ T | 12 |
| $PH_3 + B_2H_6 = PH_3BH_3(g) + BH_3$ | | 11.4 | 9.5 | -24-0 | PH ₃ /B ₂ H ₆ T | 14, 13, 11 |

Notes

(a) Critical review of literature data.

References

1. Heras, Arvia, Aymonino, and Schumacher, *Anales Asoc. Quim. Argentina* **50**, 120 (1962).
2. Heras, Arvia, Aymonino, and Schumacher, *Z. Phys. Chem. N.F.* **28**, 250 (1961).
3. Aymonino, Sicre, and Schumacher, *J. Chem. Phys.* **22**, 756 (1954).
4. Aymonino, Sicre, and Schumacher, *Anales Asoc. Quim. Argentina* **43**, 26 (1955).
5. Schumacher, *Z. Phys. Chem.* **13**, 353 (1957).
6. Schofield, Tsang, and Bauer, *J. Chem. Phys.* **42**, 2132 (1965).
7. Ashmore and Spencer, *Trans. Faraday Soc.* **55**, 1868 (1959).
8. Ashmore and Spencer, *Trans. Faraday Soc.* **58**, 1801 (1962).
9. Pshezhetskii, Morozov, Kamenetskaya, Sirystskaya, and Bribova, *Zhur. Fiz. Khim.* **33**, 2306 (1959).
10. Sulzmann, Myers, and Bartle, *J. Phys. Chem.* **42**, 3969 (1965).
11. Garabedian and Benson, *J. Am. Chem. Soc.* **86**, 176 (1964).
12. Nightingale and Crawford, *J. Chem. Phys.* **22**, 1468 (1954).
13. Bauer, *J. Am. Chem. Soc.* **78**, 5775 (1956).
14. Brumberger and Marcus, *J. Chem. Phys.* **24**, 741 (1956).
15. Porter, Szabo, and Townsend, *Proc. Roy. Soc.* **270A**, 493 (1962).

Termolecular Reactions

| Reactions | Notes | <i>E</i> | $\log_{10}A$ | Temperature range | Radical source | Reference |
|---|-------|------------------------------------|---|-------------------|--|-----------|
| | | (<i>kcal mole</i> ⁻¹) | (<i>cm</i> ^{6 mole} ^{-2 sec} ⁻¹) | °C | | |
| 2NO + O ₂ = 2NO ₂ | (a) | | $k = (7.45 \pm 0.22) \times 10^9$ | 25 | NO/O ₂ T | 1 |
| | | | $k = 6.0 \times 10^9$ | 25 | NO/O ₂ T | 2 |
| | | | $k = 7.10 \times 10^9$ | 25 | NO/O ₂ T | 3 |
| 2NO + Cl ₂ = 2NOCl | (b) | 4.2 | 10.4 | | | 8, 9 |
| | | 6.0 | 11.4 | 100-250 | NO/Cl ₂ T | 10 |
| 2NO + Br ₂ = 2NOBr | | 1.6 | 10.6 | -8-15 | NO/Br ₂ T | 11, 8 |
| 2NO + H ₂ = products | (c) | 47 | 18.7 | 700-825 | NO/H ₂ T | 12, 8 |
| NO + NO ₂ + O ₂ = NO ₂ + NO ₃ | (d) | | $k = 6.58 \times 10^7$ | 25 | NOCl/NO ₂ /O ₂ T | 13 |
| | | | $k = 3.0 \times 10^7$ | | | 15 |
| | | -0.96 | 7.32 | 200-430 | NO ₂ T | 14 |
| 2NO ₂ + O ₂ = 2NO ₃ | (e) | 25 | ($k = 7 \times 10^{-12}$) | 27 | | 15 |
| H + O ₂ + H ₂ = H ₂ O + OH | | | $k = 2 \times 10^{11}$ | 830-1630 | H ₂ /O ₂ S.T. | 16 |
| CF ₃ NO + 2NO = CF ₃ + N ₂ + NO ₃ | | | $k = 5.6 \times 10^5$ | 24 | CF ₃ I/NO P | 17 |
| CH ₃ NO + 2NO = CH ₃ + N ₂ + NO ₃ | (f) | -1.8 | 6.41 | 25-70 | CH ₃ I/NO P | 18 |
| C ₂ H ₅ NO + 2NO = C ₂ H ₅ + N ₂ + NO ₃ | (f) | -2.9 | 6.28 | 25-70 | C ₂ H ₅ I/NO P | 18 |
| CH ₃ CH ₂ CH ₂ NO + 2NO = CH ₃ CH ₂ CH ₂ + N ₂ + NO ₃ | (f) | -7 | 3.7 | 25-45 | C ₃ H ₇ I/NO P | 18 |
| (CH ₃) ₂ CHNO + 2NO = (CH ₃) ₂ CH + N ₂ + NO ₃ | (f) | -4 | 5.3 | 25-45 | (CH ₃) ₂ CHI/NO P | 18 |
| (CH ₃) ₃ CNO + 2NO = (CH ₃) ₃ C + N ₂ + NO ₃ | (f) | -7.5 | | 25-70 | (CH ₃) ₃ CI/NO P | 18 |

Notes

(a) Further data on this reaction can be found in references 4 to 7.

(b) Critical survey of literature data. The Arrhenius plot is curved, and the values given are valid only at low temperatures.

(c) It is unlikely that the nitric oxide-hydrogen reaction is a simple termolecular process.

(d) Temperature independent. Estimated from the reverse reaction.

(e) Estimated from the reverse reaction.

(f) These values are for an overall reaction which could be RNO + NO ⇌ R(NO)₂; R(NO)₂ + NO = R + N₂ + NO₃.

References

1. Brown and Crist, *J. Chem. Phys.* **9**, 840 (1941).
2. Smith, *J. Am. Chem. Soc.* **65**, 74 (1943).
3. Johnston and Slentz, *J. Am. Chem. Soc.* **73**, 2948 (1951).
4. Bodenstein and Lindner, *Z. Phys. Chem.* **100**, 87 (1922).

5. Bodenstein and Wachenheim, *Z. Electrochem.* **24**, 183 (1918).

6. Matthes, Miss Dissertation Berlin, 1933, quoted by Schumacher "Chemische Gasreaktion," p. 314 (Steinhopff, Dresden, 1938).

7. Briner, Pfeiffer, and Malet, *J. Chim. Phys.* **21**, 25 (1924).

8. Trotman-Dickenson, "Gas Kinetics," p. 265 (Butterworths, London, 1955).

9. Welinsky and Taylor, *J. Chem. Phys.* **6**, 466 (1938).

10. Ashmore and Spencer, *Trans. Faraday Soc.* **55**, 1868 (1959).

11. Krauss, *Z. Phys. Chem.* **175A**, 295 (1936).

12. Hinshelwood and Mitchell, *J. Chem. Soc.* 378 (1936).

13. Ray and Ogg, *J. Chem. Phys.* **29**, 984 (1957).

14. Ashmore and Burnett, *Trans. Faraday Soc.* **58**, 253 (1962).

15. Schott and Davidson, *J. Am. Chem. Soc.* **80**, 1841 (1958).

16. Belles and Lauver, *J. Chem. Phys.* **40**, 415 (1964).

17. Heicklen, *J. Phys. Chem.* **70**, 112 (1966).

18. Christie, Foot, and Voisey, *Trans. Faraday Soc.* **61**, 674 (1965).

Inversion (Group Transfer) Reactions

| Reactions | Notes | <i>E</i> | log ₁₀ <i>A</i> | Temperature range | Radical source | Reference |
|---|-------|---|---|-------------------|--|-------------------|
| $^{14}\text{CH}_3 + \text{CH}_3\text{COCH}_3$ = $^{14}\text{CH}_3\text{COCH}_3 + \text{CH}_3$ | (a) | (kcal mole ⁻¹) $k = 5.2 \times 10^6$ | (cm ³ mole ⁻¹ sec ⁻¹) | °C 350 | CH ₃ COCH ₃ P | 1 |
| $\text{CH}_3 + \text{CF}_3\text{COCF}_3$ = $\text{CH}_3\text{COCF}_3 + \text{CF}_3$ | | 5.7 ± 1.5 | 9.3 | 163-245 | CH ₃ N ₂ CH ₃ P and T | 2 |
| | | 6.2 ± 1.0 | 11.4 | 85-210 | CF ₃ COCF ₃ P | 3 |
| | | 7.0 ± 1.0 | 11.7 | 48-240 | CH ₃ COCH ₃ / CF ₃ COCF ₃ P | 4 |
| $\text{CH}_3 + \text{C}_2\text{F}_5\text{COC}_2\text{F}_5$ = $\text{CH}_3\text{COC}_2\text{F}_5 + \text{C}_2\text{F}_5$ | | 7 ± 1 | | 184-240 | (C ₂ F ₅) ₂ CO P | 5 |
| $\text{CD}_3 + \text{CH}_3\text{HgCH}_3$ = $\text{CD}_3\text{HgCH}_3 + \text{CH}_3$ | | | $k = 8.9 \times 10^5$ | 180 | CD ₃ COCD ₃ P | 6, 7 |
| $\text{CH}_3 + \text{CD}_3\text{SH} = \text{CH}_3\text{SH}$ + CD ₃ | | 7.6 | 10.73 | 130-200 | CH ₃ COCH ₃ P | 8 |
| $\text{CH}_3 + \text{CF}_3\text{COCH}_3$ = $\text{C}_2\text{H}_6 + \text{CF}_3\text{CO}$ | | 14. | | 150-350 | CF ₃ COCH ₃ P | 9 |
| $\text{CH}_3 + \text{CH}_3\text{COCOCH}_3$ = CH_3COCH_3 + CH ₃ CO | | 5.6 | 10.7 | 100-200 | CH ₃ COCOCH ₃ P | 10 |
| | | 6.6 | | 100-200 | CH ₃ COCOCH ₃ P | 11 |
| $\text{CH}_3 + \text{CH}_3\text{CH:CHCHO}$ = $\text{CH}_3\text{CH:CHCH}_3$ + CHO | | 7.45 ± 1.30 | 11.8 ± 0.4 | 120-250 | CH ₃ COCH ₃ P | 12 |
| $\text{CH}_3 + \text{CH}_3\text{HgCH}_3$ = $\text{C}_2\text{H}_6 + (\text{Hg} + \text{CH}_3)$ | | 1.0 | 7 | 175-220 | (CH ₃) ₂ Hg P | 13, 14 |
| $\text{C}_6\text{H}_5 + \text{C}_6\text{H}_5\text{COCH}_3$ = $\text{C}_6\text{H}_5\text{C}_6\text{H}_5 + \text{CH}_3\text{CO}$ | (b) | 6.2 | 9.6 | 277-407 | C ₆ H ₅ COCH ₃ P | 15 |
| $\text{CF}_3\text{CH}_2\text{O}^{15}\text{NO} + ^{14}\text{NO}$ = $\text{CF}_3\text{CH}_2\text{O}^{14}\text{NO}$ + ¹⁵ NO | | 23.7 ± 0.5 | 12.78 | 130-150 | CF ₃ CH ₂ O ¹⁵ NO/ ¹⁴ NO T | 16 |
| $\text{CH}_3\text{CH}_2\text{O}^{15}\text{NO} + ^{14}\text{NO}$ = $\text{CH}_3\text{CH}_2\text{O}^{14}\text{NO}$ + ¹⁵ NO | | 21.4 ± 0.5 | 11.48 | 130-150 | CH ₃ CH ₂ O ¹⁵ NO/ ¹⁴ NO T | 16 |
| $\text{BH}_3 + \text{B}_2\text{D}_6 = \text{BH}_3\text{BD}_3$ + BD ₃ | | 6.0 | 14.2 | 24-44 | B ₂ H ₆ /B ₂ D ₆ T | 17, 14, 21, 22 |
| | | 7.8 | 13.45 | | | 18 |
| $\text{BH}_3 + \text{BH}_3\text{CO} = \text{B}_2\text{H}_6$ + CO | | 7.0 | 11.4 | 0-30 | BH ₃ CO T | 19, 17, 22 |
| $\text{PH}_3 + \text{B}_2\text{H}_6 = \text{BH}_3\text{PH}_3$ + BH ₃ | | 11.4 ± 2 | 9.5 | -24-0 | | 20, 22 |

Notes

- (a) Assuming $k = 10^{11.6} \exp(-9800/RT)$ for the reaction
 $\text{CH}_3 + \text{CH}_3\text{COCH}_3 = \text{CH}_4 + \text{CH}_2\text{COCH}_3$.
(b) C_6H_5 refers to the phenyl radical.

References

1. Dainton, Ivin, and Wilkinson, *Trans. Faraday Soc.* **55**, 929 (1959).
2. Pritchard and Steacie, *Can. J. Chem.* **35**, 1216 (1957).
3. Alcock and Whittle, *Trans. Faraday Soc.* **61**, 244 (1965).
4. Giles and Whittle, *Trans. Faraday Soc.* **61**, 1425 (1965).
5. Price and Kutschke, *Can. J. Chem.* **38**, 2128 (1960).
6. Rebbert and Ausloos, *J. Am. Chem. Soc.* **86**, 2068 (1964).
7. Rebbert and Ausloos, *J. Am. Chem. Soc.* **85**, 3086 (1963).
8. Greig and Thynne, *Trans. Faraday Soc.* **62**, 379 (1966).
9. Sieger and Calvert, *J. Am. Chem. Soc.* **76**, 5197 (1954).
10. Blacet and Bell, *Disc. Faraday Soc.* **14**, 70 (1953).
11. Bell and Blacet, *J. Am. Chem. Soc.* **76**, 5332 (1954).
12. Allen and Pitts, *J. Phys. Chem.* **70**, 1691 (1966).
13. Gomer and Noyes, *J. Am. Chem. Soc.* **71**, 3390 (1949).
14. Trotman-Dickenson, "Gas Kinetics," p. 237 et seq. (Butterworths, London, 1955).
15. Duncan and Trotman-Dickenson, *J. Chem. Soc.* 4672 (1962).
16. Kuhn and Gunthard, *Helv. Chim. Acta.* **43**, 607 (1960).
17. Bauer, Shepp, and McCoy, *J. Am. Chem. Soc.* **75**, 1003 (1953).
18. Roth and Bauer, 5th Int. Comb. Symp., p. 710 (Reinhold, New York, 1955).
19. Burg, *J. Am. Chem. Soc.* **74**, 3482 (1952).
20. Brumberger and Marcus, *J. Chem. Phys.* **24**, 741 (1956).
21. Maybury and Koski, *J. Chem. Phys.* **21**, 742 (1953).
22. Garabedian and Benson, *J. Am. Chem. Soc.* **86**, 176 (1964).

(cut here)

Announcement of New Publications on Standard Reference Data

Superintendent of Documents,
Government Printing Office,
Washington, D.C. 20402

Dear Sir:

Please add my name to the announcement list of new publications to be issued in the series: National Standard Reference Data Series—National Bureau of Standards.

Name _____

Company _____

Address _____

City _____ State _____ Zip Code _____

(Notification Key N337)



