

Published in final edited form as:

Nat Genet. 2009 March ; 41(3): 354–358. doi:10.1038/ng.306.

TAC3 and TACR3 Mutations in Familial Hypogonadotropic Hypogonadism Reveal a Key Role for Neurokinin B in the Central Control of Reproduction

A.Kemal Topaloglu^{#1}, Frank Reimann^{#2}, Metin Guclu³, Ayse Serap Yalin⁴, L. Damla Kotan⁵, Keith M Porter⁶, Ayse Serin⁵, Neslihan O Mungan¹, Joshua R Cook⁶, Sazi Imamoglu³, N. Sema Akalin⁴, Bilgin Yuksel¹, Stephen O'Rahilly⁶, and Robert K Semple⁶

¹Cukurova University, Faculty of Medicine, Department of Pediatric Endocrinology and Metabolism, Balcali, Adana, TURKEY

²Cambridge Institute for Medical Research and Department of Clinical Biochemistry, University of Cambridge, Wellcome Trust/MRC Building, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 0XY, UK

³Uludag University, Faculty of Medicine, Department of Endocrinology and Metabolism, Bursa, TURKEY

⁴Marmara University, Faculty of Medicine, Department of Endocrinology and Metabolism, Istanbul, TURKEY

⁵Cukurova University, Faculty of Medicine, Department of Forensic Medicine, Balcali, Adana, TURKEY

⁶University of Cambridge Metabolic Research Laboratories, Institute of Metabolic Science, Addenbrooke's Hospital, Hills Road, Cambridge CB2 0QQ, UK

[#] These authors contributed equally to this work.

Abstract

The timely secretion of gonadal sex steroids is essential for the initiation of puberty, the post-pubertal maintenance of secondary sexual characteristics and the normal perinatal development of male external genitalia. Normal gonadal steroid production requires the actions of the pituitary-derived gonatrophins, LH and FSH. We report four human pedigrees with severe congenital gonadotrophin deficiency and pubertal failure in which all affected individuals are homozygous

Reprints and permissions information is available at npg.nature.com/reprintsandpermissions.

Correspondence and requests for materials should be addressed to A. Kemal Topaloglu, MD, Cukurova University, Faculty of Medicine, Department of Pediatric Endocrinology and Metabolism, Balcali, Adana 01330 TURKEY, Phone: +90322 3387717, Fax: +90322 3387717, ktopaloglu@cu.edu.tr or so104@medschl.cam.ac.uk.

Author Contributions

This study was designed by A.K.T.; Patient identification and phenotypic assessment was undertaken by A.K.T., M.G, A.S.Y, N.O.M, S.I., S.A., and B.Y.; DNA extraction, autozygosity mapping, and DNA sequencing were performed by AKT, LDK, and AS; Mutagenesis, cloning and functional studies were undertaken by F.R., K.M.P., R.K.S. and J.R.C.; The manuscript was written by A.K.T., R.K.S., F.R. and S.O.R..

List of URLs. AutoSNPa is available at <http://dna.leeds.ac.uk/autosnpa/>. The PRINTS-S database of protein family fingerprints is at <http://www.bioinf.man.ac.uk/dbbrowser/PRINTS>.

Supplementary Information is linked to the online version of this manuscript at www.nature.com/nature

for loss-of-function mutations in *TAC3* (encoding Neurokinin B) or its receptor *TACR3* (encoding NK3R). Neurokinin B, a member of the substance P-related tachykinin family, is known to be highly expressed in hypothalamic neurons that also express kisspeptin¹, a recently identified regulator of gonadotropin-releasing hormone secretion². These findings implicate Neurokinin B as a critical central regulator of human gonadal function and suggest novel approaches to the pharmacological control of human reproduction and sex hormone-related diseases.

The hallmark of sexual maturity is secretion from the pituitary gland of the gonadotrophins luteinising hormone (LH) and follicle stimulating hormone (FSH) which act together, in gonadal tissue, to drive both sex hormone secretion and gametogenesis³. The production of LH and FSH from pituitary gonadotrope cells is largely controlled by the pulsatile delivery of gonadotrophin releasing hormone (GnRH) from a functionally interconnected group of secretory neurons whose cell bodies are located predominantly in the medial basal hypothalamus in humans³. This complex central neuroendocrine regulatory unit is susceptible to negative feedback regulation from the gonads by sex steroids³, however although other physiological regulators have been proposed, there is considerable uncertainty about the relative importance and species specificity of many of these inputs⁴. The gonadal endocrine system is active *in utero* and for the first few months of life before being centrally suppressed. The mechanisms subserving this neural suppression and the rekindling of the GnRH pulse generating system at the beginning of puberty are, however, an enduring enigma⁵.

Recently, through a combination of human and murine genetics, the hypothalamically-expressed peptide kisspeptin and its receptor GPR54 have been established to be necessary for normal GnRH secretion and thus gonadotrope function^{6,7}. Human genetics has been a powerful contributor to the discovery of critical molecular elements of the central control of reproduction identifying a number of molecules which are essential for the embryonic migration of GnRH neurons to the hypothalamus from the olfactory placode (*KALI*, *NELF*, *FGFR1*, *PROK1*, *PROKR2*). Humans with genetic defects in those molecules generally present clinically with failure to enter puberty and low circulating gonadotropin levels (hypogonadotropic hypogonadism (HH)) associated with impairment of olfactory sensation (MIM-308700,147950). Mutations either in the GnRH receptor or in GPR54 have been reported in humans with familial HH but normal olfaction (so called normosmic idiopathic HH (nIHH; MIM-146110)). However these genetic defects explain only a minority of cases of familial nIHH⁸ implying the existence of other, as yet unknown, key positive regulators of the central control of the reproductive axis.

Nine consanguineous Turkish families with multiple members affected by nIHH were subjected to genome-wide SNP analysis, and within each family regions of homozygosity common to all affected, but not found in unaffected individuals, were identified⁹. In three families (Fig 1A) homozygosity at a locus on the long arm of chromosome 4 cosegregated with nIHH. Alignment of all 6 affected members of the three families refined the critical interval to a 2.74 MB genomic region extending from 102.20 to 104.94 MB encompassing 20 known or predicted genes (Fig 1B). Of those genes *TACR3* (NM-001059) was prioritized for further analysis. *TACR3* encodes NK3R, the receptor for Neurokinin B, a tachykinin

peptide which is known to be highly expressed in hypothalamic neurons that also express kisspeptin¹. Synthetic NK3R agonists have also been reported to modulate reproductive function but the site, and indeed direction, of such actions have been unclear^{10,11}. Homozygous non-synonymous mutations in the coding sequence of *TACR3* were found in all affected subjects (and no unaffected subjects) in all three families (supplementary note and figure). In family 1 all affected individuals were homozygous for a G to A transversion at cDNA nucleotide 278. This mutation leads to the substitution of glycine at residue 93 for aspartic acid (G93D). G93 lies in the first transmembrane domain of the G protein-coupled NK3R (Fig 1C), and, although it is not highly conserved (Fig 1D), introduction of a negatively charged side change in this very hydrophobic domain would likely be highly thermodynamically unfavourable. Indeed, only 1 of 739 members of the rhodopsin receptor superfamily has a charged side chain at this position. In families 2 and 3 a C to T transition at cDNA nucleotide 1057 was found, producing a proline to serine change at residue 353 (P353S), which lies within the sixth transmembrane domain (Fig 1C). P353 is exquisitely conserved (Fig 1D): 735 out of 739 members of the rhodopsin receptor superfamily have a proline at this position. Neither of these mutations were found in 100 Turkish controls, in the 6 other families from the original genome wide SNP-analysis, nor 50 patients with nHH from a variety of ethnic groups.

In one multiplex family in which *TACR3* mutations had been excluded (Fig 2A) 2 autozygous regions were identified which were common to the two affected family members but not to their unaffected sister. Although these regions - from 125.7 to 130.4 MB on chromosome 4 and 54.5 to 57.6 MB on chromosome 12 - encompassed 21 and 102 known or predicted genes respectively, the only obvious biologically plausible candidate was *TAC3* (NM-0013251), on chromosome 12 (Fig 2A), encoding neurokinin B, the preferred ligand for the neurokinin-3 receptor. On sequencing *TAC3* we identified a homozygous T to C transversion at cDNA nucleotide 269 in both affected siblings, leading to substitution of methionine by threonine at residue 90 (M90T) of the prohormone (Fig 2B). The parents and unaffected sister were heterozygous for the mutation, which was not detected in 100 ethnically-matched controls. Met90 is the C terminal amino acid of the mature neurokinin B decapeptide, and falls within the canonical tachykinin motif, Phe-X-Gly-Leu-Met-NH₂, which is universally conserved among known tachykinins¹² (e.g. Fig 2C). *In vivo* NKB is known to be post-translationally modified by C-terminal amidation, which has been shown to be necessary for full activity of the closely related neurokinin A peptide¹³.

NK3R activation is linked to Gq proteins and its action results in an increase in intracellular calcium. To evaluate the functional consequences of the *TACR3* mutations we expressed wild type and mutant *TACR3* in HEK293 cells and assessed the ability of neurokinin B (NKB), to stimulate calcium flux. In transiently transfected cells, identified on the basis of coexpressed GFP, NKB elicited a robust increase in intracellular calcium with an EC₅₀ of 6.6±1.3 nM, while no response to NKB was seen in untransfected cells (Fig 3). Both G93D and P353S mutant receptors showed severely attenuated ability to respond to stimulation, with EC₅₀s for NKB of 290±146 nM and 4.0±2.1 µM respectively. Thus, both the missense *TACR3* mutations found in all affected subjects in Families 1 through 3 show unequivocal evidence of impaired receptor signaling.

To assess the activity of the M90T mutant NKB found in Family 4, the mature mutant decapeptide was synthesized in both C terminal-amidated and non-amidated form, and the effects of each peptide on wild type receptor were assessed in the same heterologous expression system. The amidated mutant peptide (predicted to be the form that would occur in nature) had severely reduced activity, being around 10 fold less potent than amidated wild type NKB at stimulating the NK3R (EC_{50} -M90T = 52 ± 6 nM; Fig 4). As expected, the non-amidated mutant NKB had little detectable ability to elicit calcium flux (data not shown).

The phenotypes of subjects homozygous for either *TAC3* or *TACR3* mutations provide strong evidence that NKB, acting through its receptor NK3R, is essential for the central hormonal control of human reproduction (supplementary note and supplementary table 1). All subjects who are in the second decade of life or older show unequivocal evidence of failure of pubertal progression with low circulating sex steroid levels and pre-pubertal levels of circulating gonadotropins. Furthermore, all affected males have micropenis, strongly suggesting failure of normal intrauterine and perinatal activation of the reproductive axis¹⁴. The only homozygote (out of 9) in any of the four families who does not yet exhibit a clear phenotype is a 5 year-old female in whom HH would not be expected to have become overt.

NK3R is widely expressed, particularly in the central nervous system¹⁵, and previous studies in rodents utilising pharmacological or genetic manipulation have suggested potential roles in a wide range of CNS functions¹⁶⁻¹⁷. Additionally, polymorphisms in the *TACR3* locus have been weakly genetically associated with a range of human neurobehavioural phenotypes¹⁸. In this regard, it is notable that patients with homozygous loss of function mutations in *TACR3* lack any other clear neurobehavioural or other phenotypic abnormalities outside the reproductive axis. However, both affected subjects with *TAC3* mutations do have mild learning disability (supplementary note) suggesting that NKB action, possibly through receptors other than NK3R, may have effects on higher cognitive function.

The precise mechanism whereby NKB and the NK3R exert their effects on the control of reproduction remains to be established. It is most likely, however, that this effect is exerted at the level of hypothalamic GnRH release. NK3R is expressed on rodent GnRH-expressing neurons^{19,20}. Axons of neurons expressing NKB are closely anatomically apposed to those of GnRH neurons within the median eminence of the hypothalamus^{19,21}, and NKB immunoreactive varicosities have been reported to be in direct contact with GnRH immunoreactive axons²². NKB expression is highest in the arcuate nucleus, where it colocalises with estrogen receptor alpha (Era) and dynorphin²³, implicated in progesterone feedback to GnRH secretion²⁴. Nevertheless NK3R and NKB are also expressed in peripheral organs including ovaries and uterus²⁵, so additional peripheral effects cannot be formally excluded. However there is a paucity of pharmacological or murine genetic evidence directly implicating NKB or NK3R in gonadal function. Mice with genetic ablation of *Tac2* (the murine orthologue of *TAC3*) have not been described and studies of *Tacr3*^{-/-} mice have not reported a reproductive phenotype^{17,26,27}. Notably, central administration of a potent NK3R agonist to ovariectomized, oestrogen-primed rats suppressed LH production rather than stimulating it¹⁰. It thus remains possible that there is true divergence in the reproductive role of NKB/NK3R between humans and rodents.

In summary, we have identified loss-of-function mutations in either neurokinin B or its receptor in four out of nine multiplex families affected by hypogonadotrophic hypogonadism. These findings establish that NKB action *via* the NK3R is necessary for the central neuroendocrine control of human reproduction. These families represents the first examples of inherited defects of tachykinin signalling in any human disorder and suggest that the NKB/NK3R system may provide a novel avenue for the pharmacological manipulation of human fertility and the treatment of sex steroid-related diseases, such as cancer of the breast and prostate.

Methods Summary

The ethics committee of the Cukurova University Faculty of Medicine approved this study, and full informed consent was obtained for each participant. nIHH was defined by conventional criteria (supplementary methods). Normal karyotypes were confirmed and mutations in *GNRHR*, *GNRH1*, *GPR54*, *KISS1*, *KAL1*, *NELF*, *PROK2*, *PROK2R* and *FGFR1* genes were ruled out by sequencing. Whole genome SNP analysis used 250K NspI Affymetrix SNP microarrays (Affymetrix, CA, USA) and data were analyzed using AutoSNPa software⁹. PCR-amplified exons and splice junctions of *TAC3* or *TACR3* were sequenced on an ABI PRISM 3130 autosequencer (Applied Biosystems, Foster City, CA). For primer sequences and annealing temperatures see supplementary Table 2.

Wild type or mutant *TACR3* in pIRES2-AcGFP1 (Clontech) were transfected into HEK293A cells (QBiogene) using polyethylenimine (Sigma) before seeding cells into poly-L-lysine-coated glass-bottomed dishes 16 hours later. At 24 hours cells were loaded with Fura2-AM for 30 min. Experiments were performed on an inverted fluorescence microscope (Eclipse TE2000, Nikon, UK) with a 40x oil-immersion objective. Fura2 was excited at 340, 360 and 380 nm, and GFP at 475 nm, using a 75W xenon arc lamp and a monochromator (Cairn Research, Faversham, UK) and MetaFluor software (Molecular Devices, UK). Emission, filtered at 510/80 and 535/25 for Fura2 and GFP respectively, was recorded with a QuantEM CCD camera (Photometrics, Roper Scientific, UK). NKB (Sigma) or custom made mutant NKB with or without C terminal amidation (New England Peptides) were perfused in bath solution at ~1 ml/min with a chamber volume of ~0.2 ml. 340/380 nm ratio was calculated from images taken at 100 ms excitation at each wavelength at 0.5 Hz. Free Ca^{2+} -concentrations were calculated for individual cells after background subtraction using the equation of Grynkiewicz et al (1985) assuming a K_D of 224 nM. Minimal and maximal signals were recorded in the presence of 5 mM ionomycin in 5 mM EGTA/0 mM Ca^{2+} and 5 mM Ca^{2+} respectively at the end of the experiment. Further detail is given in the supplementary methods.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by grants to A.K.T., N.O.M., and B.Y. from The Scientific and Technological Research Council of Turkey (TÜBİTAK: The Support Programme for Scientific and Technological Research Projects (1001),

Grant no: 106S276), by the Wellcome Trust (R.K.S.: Intermediate Clinical Fellowship 080952/Z/06/Z; S.O.: Programme Grant 078986/Z/06/Z; F.R.: Senior Basic Science Fellowship 084210/Z/07/Z) and the U.K. NIHR Cambridge Biomedical Research Centre. F.R. has also received support from St. John's College, Cambridge.

References

1. Goodman RL, et al. *Endocrinology*. 2007; 148(12):5752. [PubMed: 17823266]
2. Gianetti E, Seminara S. *Reproduction*. 2008; 136(3):295. [PubMed: 18515314]
3. Plant TM. *J Neuroendocrinol*. 2008; 20(6):719. [PubMed: 18601694]
4. Herbison, AE. *Knobil and Neill's Physiology of Reproduction*. Neill, JD., editor. Elsevier; Amsterdam: 2006. p. 1415-1482.
5. Herbison AE. *Hormone research*. 2007; 68(Suppl 5):75. [PubMed: 18174715]
6. Seminara SB, et al. *N Engl J Med*. 2003; 349(17):1614. [PubMed: 14573733]
7. de Roux N, et al. *Proc Natl Acad Sci U S A*. 2003; 100(19):10972. [PubMed: 12944565]
8. Crowley WF Jr, Pitteloud N, Seminara S. *Trans Am Clin Climatol Assoc*. 2008; 119:29. [PubMed: 18596868]
9. Carr IM, et al. *Hum Mutat*. 2006; 27(10):1041. [PubMed: 16941472]
10. Sandoval-Guzman T, Rance NE. *Brain Res*. 2004; 1026(2):307. [PubMed: 15488494]
11. Rance NE. *Peptides*. 2008
12. Almeida TA, et al. *Curr Med Chem*. 2004; 11(15):2045. [PubMed: 15279567]
13. Patacchini R, et al. *J Pharmacol Exp Ther*. 1993; 264(1):17. [PubMed: 8380857]
14. Grumbach MM. *J Clin Endocrinol Metab*. 2005; 90(5):3122. [PubMed: 15728198]
15. Pinto FM, et al. *Eur J Pharmacol*. 2004; 494(2-3):233. [PubMed: 15212980]
16. Hasenohrl RU, et al. *Neuropeptides*. 2000; 34(5):272. [PubMed: 11049731]
17. Siuciak JA, et al. *Psychopharmacology (Berl)*. 2007; 194(2):185. [PubMed: 17558564]
18. Saito S, et al. *Neuroreport*. 2008; 19(4):471. [PubMed: 18287949] Foroud T, et al. *Alcohol Clin Exp Res*. 2008; 32(6):1023. [PubMed: 18422838]
19. Krajewski SJ, et al. *J Comp Neurol*. 2005; 489(3):372. [PubMed: 16025449]
20. Todman MG, Han SK, Herbison AE. *Neuroscience*. 2005; 132(3):703. [PubMed: 15837132]
21. Goubillon ML, et al. *Endocrinology*. 2000; 141(11):4218. [PubMed: 11089556]
22. Ciofi P, Leroy D, Tramu G. *Neuroscience*. 2006; 141(4):1731. [PubMed: 16809008]
23. Burke MC, Letts PA, Krajewski SJ, Rance NE. *J Comp Neurol*. 2006; 498(5):712. [PubMed: 16917850]
24. Goodman RL, et al. *Endocrinology*. 2004; 145(6):2959. [PubMed: 14988383]
25. Patak E, et al. *Br J Pharmacol*. 2003; 139(3):523. [PubMed: 12788812] Loffler S, et al. *Regul Pept*. 2004; 122(2):131. [PubMed: 15380930]
26. Nordquist RE, et al. *Psychopharmacology (Berl)*. 2008; 198(2):211. [PubMed: 18351324]
27. Kung TT, et al. *Pharmacol Res*. 2004; 50(6):611. [PubMed: 15501700]
28. Rosenfield, RL. *Pediatric endocrinology: A practical clinical guide*. Radovick, S.; MacGillivray, MH., editors. Humana Press; 2003. p. 451




Figure 1.

A. Families used to identify critical interval on chromosome 4 harbouring genetic defect causing nHH **B.** Critical region on chromosome 4 showing homozygosity (black) in all affected individuals, with known or predicted genes in region indicated. **C.** Location of pathogenic mutation in NK3R **D.** evolutionary conservation of mutated residues.




Figure 2.

A. Family used to identify region on chromosome 12 harbouring genetic defect causing nIHH with detail of subset of genes in critical interval **B.** Location of pathogenic mutation in TAC3. Shaded amino acids represent the tachykinin signature motif. **C.** Conservation of mutated residue among selected paralogues and orthologues.




Figure 3.

Both missense mutations in NK3R result in loss of receptor function. **A)** Bright field (left) and GFP-fluorescent only (right) identifying TACR3 transfected HEK293 cells. **B)** Pseudocolour images of 340/380 nm ratio for the same optical field as in (A) before (left) and during (right) the application of 100 nM wt-NKB. Note that only GFP-positive cells respond. **C)** Representative Ca^{2+} -responses of 3-5 wt- or mutant TACR3 transfected cells to application of different NKB doses as indicated. **D)** NKB dose responses for wt (black squares, $n=93-206$), G93D- (red squares, $n=112-145$) and P353S-mutant (green squares,

n=156) TACR3. Peak $[Ca^{2+}]$ -responses recorded as in (C) in at least 5 dishes/construct were averaged over 30s and the data was fit with the logistic function $[Ca^{2+}] = (A1-A2)/(1 + ([NKB]/EC50)) + A2$ using origin software (Microcal). The resulting fit parameters were: wt-TACR3: $A1=45 \pm 6$ nM, $A2=227 \pm 6$ nM, $EC50=6.6 \pm 1.3$ nM; TACR3-G93D: $A1=45 \pm 2$ nM, $A2=91 \pm 5$ nM, $EC50=290 \pm 146$ nM; TACR3-P353S: $A1=47 \pm 3$ nM, $A2=75$ nM (fixed), $EC50=4.0 \pm 2.1$ μ M. Error bars represent 1SE.




Figure 4.

The missense mutations in NKB results in severe loss of agonist function at the NK3R **A)** Representative Ca^{2+} -responses of 3 wt-TACR3 transfected cells to application of different NKB and NKB-M90T doses as indicated. **B)** NKB dose responses for NKB (black squares), and NKB-M90T (red circles). Peak $[Ca^{2+}]$ -responses recorded as in **(A)** in 5 dishes ($n=113$ cells) were averaged over 30s and the data was fit with the logistic function $[Ca^{2+}] = (A1 - A2)/(1 + ([NKB]/EC50)) + A2$ using origin software (Microcal). The resulting fit parameters were: NKB: $A1=40 \pm 8$ nM, $A2=242 \pm 11$ nM, $EC50=5.8 \pm 1.5$ nM; NKB-M90T: $A1=38 \pm 4$ nM, $A2=242$ nM (fixed), $EC50=52 \pm 6$ nM. Error bars represent 1SE; responses to similar doses of the two peptides were compared using a paired t-test; i $p=3 \times 10^{-12}$, ii $p=3 \times 10^{-20}$, iii $p=6 \times 10^{-7}$.