

Tansley Review No. 111

Possible roles of zinc in protecting plant cells from damage by reactive oxygen species

ISMAIL CAKMAK

Department of Soil Science and Plant Nutrition, Faculty of Agriculture, University of Cukurova, 01330 Adana, Turkey (fax +90 322 3386747; e-mail cakmak@mail.cu.edu.tr)

Received 20 August 1999; accepted 6 January 2000

Dedicated to the memory of the late Prof. Dr Drs. h. c. Horst Marschner.

CONTENTS

SUMMARY	185	1. <i>Impairments in membrane structure</i>	193
I. INTRODUCTION	186	2. <i>Phospholipids and –SH groups</i>	195
II. EFFECT OF ZINC ON PRODUCTION OF REACTIVE OXYGEN SPECIES	186	3. <i>Alterations in ion absorption</i>	195
1. <i>Superoxide-generating NADPH oxidase</i>	186	(a) <i>Membrane-bound ATPases</i>	195
2. <i>Zinc deficiency potentiates iron-mediated free radical production</i>	189	(b) <i>Nutrient uptake</i>	197
(a) <i>Iron accumulation in zinc-deficient plants</i>	189	(c) <i>Changes in activity of ion channels</i>	197
(b) <i>Iron-induced production of free radicals</i>	189	IV. DETOXIFICATION OF REACTIVE OXYGEN SPECIES	198
3. <i>Zinc deficiency-enhanced photooxidation</i>	191	1. <i>Superoxide dismutases</i>	198
(a) <i>Decrease in photosynthesis</i>	191	2. <i>H₂O₂-scavenging enzymes</i>	198
(b) <i>Light-induced leaf chlorosis</i>	192	V. INVOLVEMENT OF ZINC IN PLANT STRESS TOLERANCE	199
(c) <i>Decrease in indole-3-acetic acid</i>	192	VI. CONCLUSIONS	199
III. MEMBRANE DAMAGE BY REACTIVE OXYGEN SPECIES	193	ACKNOWLEDGEMENTS	200
		REFERENCES	200

SUMMARY

Zinc deficiency is one of the most widespread micronutrient deficiencies in plants and causes severe reductions in crop production. There are a number of physiological impairments in Zn-deficient cells causing inhibition of the growth, differentiation and development of plants. Increasing evidence indicates that oxidative damage to critical cell compounds resulting from attack by reactive O₂ species (ROS) is the basis of disturbances in plant growth caused by Zn deficiency. Zinc interferes with membrane-bound NADPH oxidase producing ROS. In Zn-deficient plants the iron concentration increases, which potentiates the production of free radicals. The Zn nutritional status of plants influences photooxidative damage to chloroplasts, catalysed by ROS. Zinc-deficient leaves are highly light-sensitive, rapidly becoming chlorotic and necrotic when exposed to high light intensity. Zinc plays critical roles in the defence system of cells against ROS, and thus represents an excellent protective agent against the oxidation of several vital cell components such as membrane lipids and proteins, chlorophyll, SH-containing enzymes and DNA. The cysteine, histidine and glutamate or aspartate residues represent the most critical Zn-binding sites in enzymes, DNA-binding proteins (Zn-finger proteins) and membrane proteins. In addition, animal studies have shown that Zn is involved in inhibition of apoptosis (programmed cell death) which is preceded by DNA and membrane damage through reactions with ROS.

Key words: iron accumulation, membrane damage, NADPH oxidase, photooxidation, stress tolerance, superoxide dismutase, superoxide radical, zinc.

I. INTRODUCTION

Zinc deficiency is a common micronutrient deficiency in plants growing in different climatic regions of the world, particularly in arid and semi-arid regions where alkaline soils predominate (Welch *et al.*, 1991; Takkar & Walker, 1993; White & Zasoski, 1999). In an extensive soil survey, Sillanpää (1990) found approx. 50% of the soil samples collected in 25 countries to contain a deficient level of plant-available Zn. Consequently, severe depressions in plant growth and yield have been reported for many countries, for example Australia (Graham *et al.*, 1992), Turkey (Cakmak *et al.*, 1996a, 1999) and India (Takkar *et al.*, 1989).

Several physiological processes are impaired in plants suffering from Zn deficiency. Zinc deficiency causes rapid inhibition of plant growth and development, and thus of final yield. Zinc plays a fundamental role in several critical cellular functions such as protein metabolism, gene expression, structural and functional integrity of biomembranes, photosynthetic C metabolism and IAA metabolism (Marschner, 1995). Compared with other micronutrients, Zn exists in biological systems in high concentrations, particularly in biomembranes. According to Williams (1988), concentrations of Zn in animal systems range from $<10^{-9}$ M in cytoplasm to $>10^{-3}$ M in some membrane vesicles. There are many binding sites for Zn within the membranes, particularly in the interior sites of membranes. As shown with brush border membranes, the maximal binding capacity of membranes is around 400 nmol Zn²⁺ per mg protein (Prasad *et al.*, 1996). In roots of wheat seedlings, cytoplasmic concentration of total Zn has been estimated at approx. 0.4 mM (Santa Maria & Cogliatti, 1988). Most of the critical functions of Zn in cells are related to its ability to form tetrahedral coordination bonds in different vital cell constituents. Cysteine, histidine and aspartate or glutamate are major cellular ligands of Zn that form tetrahedral coordinations (Williams, 1988; Vallee & Auld, 1990; Vallee & Falchuk, 1993). These ligands, especially cysteine and histidine, bind to Zn with a greater affinity and with more stability than to Fe (Berg & Shi, 1996). Thus free radical formation, via reactions between Fe and cysteine and histidine residues, is blocked in the presence of adequate Zn (Searle & Tomasi, 1982; Girotti *et al.*, 1985; Bray & Bettger, 1990).

An increase in the level of reactive O₂ species (ROS) and a decrease in detoxification mechanisms may be the major reasons for impairment of various cellular functions in Zn-deficient plants. Zinc is required for detoxification of ROS including O₂^{•-} (superoxide radical) and H₂O₂ (hydrogen peroxide). Because of its fundamental role in the activation and expression of genes (Klug & Rhodes, 1987; Vallee & Falchuk, 1993), Zn might be involved in oxidative

stress-induced expression of genes encoding anti-oxidative defence enzymes such as H₂O₂-scavenging ascorbate peroxidase and glutathione reductase (Gressel & Galun, 1994; Allen, 1995; Alscher *et al.*, 1997). Several studies have reported that low amounts of Zn in plant cells may enhance production of O₂^{•-} during photosynthetic electron transport (Marschner & Cakmak, 1989; Cakmak *et al.*, 1995; Cakmak & Engels, 1999) and induce O₂^{•-}-generating membrane-bound NADPH oxidase (Cakmak & Marschner, 1988a; Pinton *et al.*, 1994).

Evidence from animal studies has shown that Zn interacts with the binding of Fe to membranes and thus restricts Fe-induced production of highly toxic hydroxyl radicals (OH[•]), a reaction which is referred to as site-specific OH[•] production (Girotti *et al.*, 1985; Powell *et al.*, 1994). Also, apoptosis (programmed cell death) is initiated by ROS (Greenberg *et al.*, 1994; Sarafian & Bredesen, 1994; Dangl *et al.*, 1996), and is stimulated by low concentrations of soluble Zn in the cytosol (Treves *et al.*, 1994; Zalewski *et al.*, 1994). These results imply that an adequate Zn supply is critically important for protection of cells against damaging attacks by ROS.

This review emphasizes the roles played by Zn in the generation and detoxification of ROS and in the protection and maintenance of cellular integrity in plants. Additionally, attention is given to animal studies dealing with the inhibitory effects of Zn on the production of ROS in animal cells. These subjects have been reviewed briefly by Brown *et al.* (1993) and Marschner (1995). For more detailed information on the role of Zn in the protection of cellular components from harmful reactions with toxic O₂ species in animal cells, readers are referred to the excellent reviews by Willson (1988) and Bray & Bettger (1990).

II. EFFECT OF ZINC ON PRODUCTION OF REACTIVE OXYGEN SPECIES

1. Superoxide-generating NADPH oxidase

Production of the superoxide radical (O₂^{•-}) in plant cells is an unavoidable process that is catalysed by a one-electron reduction of molecular O₂ (Elstner, 1982; Asada & Takahashi, 1987; Asada, 1994). Further reactions of O₂^{•-} are responsible for the generation of even more toxic O₂ species such as the hydroxyl radical (OH[•]) and singlet oxygen (¹O₂). Production of O₂^{•-} and its derivatives can be induced when plants are exposed to various biotic and abiotic stresses. In most cases, induced production of O₂^{•-} is catalysed by NAD(P)H-oxidizing enzyme systems localized in different cell compartments, such as cell walls (Gross *et al.*, 1977; Cakmak *et al.*, 1987); plasma membranes (Pinton *et al.*, 1994; Doke *et al.*, 1996; Murphy & Auh, 1996); cytosol and micro-




Fig. 1. Changes with plant age in the levels of $O_2^{\cdot-}$ in microsomal membrane fraction isolated from roots of cotton plants grown with (+Zn, 10^{-6} M $ZnSO_4$) and without (-Zn) Zn supply in nutrient solution. Levels of $O_2^{\cdot-}$ were determined by electron spin resonance (ESR) spectroscopy measurement of the 1,2-dihydroxybenzene-3,5-disulphonic acid (Tiron) semiquinone radical which is formed from the oxidation of Tiron by $O_2^{\cdot-}$ (McRae *et al.*, 1982). Increases in amplitude of the Tiron ESR signal indicate greater production of $O_2^{\cdot-}$ (Cakmak, 1988; Cakmak & Marschner, 1988a).

somes (Cakmak & Marschner, 1988a,b); peroxisomes (Del Rio *et al.*, 1998; Lopez-Huertas *et al.*, 1999); and mitochondria (I. Cakmak, unpublished). The production of $O_2^{\cdot-}$ and its derivatives from NAD(P)H oxidases is associated with different cellular functions, for example lignification of cell walls (Gross *et al.*, 1977), resistance against pathogen infection (Mehdy *et al.*, 1996; Bolwell & Wojtaszek, 1997), and reduction and uptake of iron by roots (Cakmak *et al.*, 1987).

Toxic O_2 species produced by NADPH-dependent oxidases are also involved in damage to several critical cell constituents (Svingen *et al.*, 1979; Jabs *et al.*, 1996; Wojtaszek, 1997). Many enzymatic and non-enzymatic lipid peroxidation processes show a high dependency on NADPH as a basic electron donor, particularly in the presence of redox active transition metals such as Fe. Supplementation of NADPH, together with ferric iron, to isolated cell compartments from animal cells (such as microsomes and mitochondria) results in a rapid peroxidation of lipids (Minotti, 1990; Glinn *et al.*, 1991). Moreover, the role of soybean lipoxygenase in oxidative cell stress is possibly related to the ability of this enzyme to produce $O_2^{\cdot-}$ by oxidizing NADPH (Roy *et al.*, 1994).

The activity of $O_2^{\cdot-}$ -generating NADPH oxidases is greatly influenced by Zn deficiency in animal and plant cells. Zinc exerts a strong inhibitory effect on the generation of $O_2^{\cdot-}$ by NADPH oxidase. In Zn-deficient animal cells, NADPH-dependent production of $O_2^{\cdot-}$ was much higher than in Zn-sufficient cells, and this was considered as a major reason for Zn deficiency-induced membrane damage (Chvapil, 1979; Hammermüller *et al.*, 1987). Hammermüller *et al.* (1987) reported that Zn deficiency results in a two- to three-fold increase in NADPH-dependent H_2O_2 production in rat micro-

somes. However, under the same conditions Cu deficiency did not affect NADPH-dependent H_2O_2 production. In agreement with these results, Burke & Fenton (1985) reported a 10-fold increase in NADPH-dependent lipid peroxidation in Zn-deficient rat cells. Similarly, in plant cells Zn interferes with NADPH-dependent $O_2^{\cdot-}$ generation. In microsomal membrane and cytosolic fractions isolated from cotton roots, levels of $O_2^{\cdot-}$ measured by electron spin resonance (ESR) spectroscopy were higher in Zn-deficient than in Zn-sufficient plants. Levels of $O_2^{\cdot-}$ obtained by ESR spectroscopy are determined using Tiron as a spin probe for $O_2^{\cdot-}$ (McRae *et al.*, 1982). The amplitude of the Tiron-ESR signal in the microsomal membrane fraction from cotton roots is shown in Fig. 1, and reveals a progressive increase in signal with the severity of Zn deficiency, but the signal was not affected in Zn-sufficient plants. The changes in amount of $O_2^{\cdot-}$ measured directly by ESR spectroscopy were positively correlated with the activity of an $O_2^{\cdot-}$ -generating NADPH oxidase in the same microsomal membranes and cytosolic fractions (Cakmak & Marschner, 1988a,b). This result indicated a role of $O_2^{\cdot-}$ -generating NADPH oxidase in the level of $O_2^{\cdot-}$ measured by ESR spectroscopy. Activity of $O_2^{\cdot-}$ -generating NADPH oxidase in Zn-deficient roots is very sensitive to Zn; resupply of Zn to Zn-deficient plants rapidly decreased the generation of $O_2^{\cdot-}$ in microsomal membranes (Cakmak & Marschner, 1988b). The activity of $O_2^{\cdot-}$ -generating NADPH oxidase determined in Zn-deficient plants was unstable and highly sensitive to heating or storage on ice, and had an optimum pH of 7.6 (Cakmak *et al.*, 1988a). As in the $O_2^{\cdot-}$ -generating NADPH oxidases described in different animal and plant cells (Bolwell & Wojtaszek, 1997), NADPH oxidase in Zn-deficient root cells also shows a high dependency on

Table 1. NAD(P)H oxidation and $O_2^{\cdot-}$ production in plasma membrane vesicles isolated from roots of bean plants grown without ($-Zn$) and with ($+Zn = 2 \times 10^{-6} M$) Zn supply

	+Zn	-Zn	$\pm Zn$
	(nmol mg ⁻¹ protein min ⁻¹)		
NAD(P)H oxidation			
NADH	3.3 (100)	8.4 (264)	6.4 (194)
NADPH	2.5 (100)	6.6 (264)	4.3 (172)
$O_2^{\cdot-}$ production			
NADH	4.3 (100)	9.1 (212)	6.9 (160)
NADPH	2.8 (100)	8.2 (293)	4.8 (171)

$\pm Zn$ indicates re-supply of $4 \times 10^{-6} M$ Zn to deficient plants 24 h before vesicle isolation. Data are means of three independent preparations of membrane vesicles. Standard deviation did not exceed 5% of the means. Values in parentheses indicate activities presented as a percentage of those in Zn sufficient ($+Zn$) plants (Pinton *et al.*, 1994).

a flavin (as an electron carrier) for a maximal activity (Cakmak & Marschner, 1988a; Pinton *et al.*, 1994).

Based on these results, it is possible that Zn-deficient cells have enhanced activity of a plasma membrane-bound $O_2^{\cdot-}$ -generating NADPH oxidase. This assumption was tested using isolated plasma membrane vesicles from roots of Zn-deficient and Zn-sufficient bean plants (Pinton *et al.*, 1994). As with the results obtained with microsomal membrane fractions, the activity of NADPH-dependent $O_2^{\cdot-}$ -generating oxidase was strongly increased by Zn deficiency in plasma membrane vesicles (Table 1). Pinton *et al.* (1994) also showed that Zn deficiency-induced $O_2^{\cdot-}$ generation was greater when NADPH instead of NADH was used as electron donor (Table 1). The $O_2^{\cdot-}$ -generating NADPH oxidase activity of Zn-deficient membranes is very similar to that induced in plant cells or animal phagocytotic cells in response to pathogenic infection (Doke *et al.*, 1996; Bolwell & Wojtaszek, 1997). Increases in plasma membrane-bound NADPH oxidase activity as a result of pathogen invasion are associated with high production of $O_2^{\cdot-}$ and other harmful O_2 species, and are produced in order to kill the pathogens and activate defence-related genes in host cells (Mehdy *et al.*, 1996). It is not known if Zn interferes with pathogen-induced NADPH-dependent $O_2^{\cdot-}$ generation in plant cells. In animal systems, $O_2^{\cdot-}$ -generating NADPH oxidase of phagocytic cells was strongly decreased by high Zn concentrations (Suzuki *et al.*, 1985). In contrast, magnesium was required for optimal activity (Cross *et al.*, 1999).

Enhanced production of ROS by NADPH-dependent oxidases is a common phenomenon occurring during apoptosis (Dangl *et al.*, 1996; Jabs *et al.*, 1996; Wojtaszek, 1997). Reactive O_2 species causing membrane and DNA damage have been

shown to be major mediators of apoptosis (Sarafian & Bredesen, 1994; Ridgley *et al.*, 1999). There is increasing evidence in the animal literature demonstrating an important role for Zn in inhibiting apoptosis. Treves *et al.* (1994) reported that Zn is the most abundant intracellular metal and functions as a key intracellular regulator of apoptosis. A decrease in intracellular Zn^{2+} concentration triggers apoptotic cell death by inducing DNA degradation. Similarly, Zalewski *et al.* (1993, 1994) reported that the concentration of readily soluble Zn compounds in the cytosol plays a determining role in preventing DNA fragmentation, and thus apoptosis. According to Parat *et al.* (1997), inhibitory action of Zn on apoptosis seems to be related to its role in protecting cell membranes and DNA from damaging attack by toxic O_2 species. In view of the inhibitory effect of Zn on NADPH oxidase, it is logical to assume that prevention of apoptosis by Zn can be attributed to a reduced activity of NADPH oxidases involved in ROS-mediated apoptotic cell death. This area is an interesting research topic for further study. Recently, a zinc-finger protein encoded by the *Arabidopsis LSD1* gene was described as being involved in the repression of superoxide-dependent cell death in plants (Dietrich *et al.*, 1997).

The enhancing effect of Zn deficiency on $O_2^{\cdot-}$ -generating NADPH oxidase can be ascribed to the dual effects of Zn on enzyme activity. First, Zn interferes with enzymatic oxidation of NADPH either by a specific binding of Zn to NADPH (Chvapil, 1979) or by an alteration of the redox properties of the NADPH-oxidizing enzyme complex (Jeffery, 1983). Accordingly, Zn could decrease the NADPH oxidation capacity of microsomal and plasma membrane fractions (Cakmak & Marschner, 1988b; Table 1). Secondly, the inhibitory effect of Zn on NADPH-dependent $O_2^{\cdot-}$ -generating oxidase

can also be related to greater activity of $O_2^{\cdot-}$ -scavenging superoxide dismutases (SODs) in Zn-sufficient cells (section IV).

2. Zinc deficiency potentiates iron-mediated free radical production


(a) *Iron accumulation in zinc-deficient plants.* In many monocotyledonous and dicotyledonous crop species grown either in soil or solution culture, Zn deficiency results in a high accumulation of Fe in roots and shoots. In a nutrient solution experiment with nine plant species, Rahimi & Bussler (1979) reported an excessive accumulation of Fe in leaves of Zn-deficient plants. Iron concentration of leaves ranged between 180 and 800 mg kg⁻¹ d. wt for Zn-deficient plants, and between 75 and 300 mg kg⁻¹ d. wt for Zn-sufficient plants. In a further nutrient solution study with 10 different wheat cultivars, shoot concentrations of Fe in Zn-deficient compared to Zn-sufficient plants were, on average, twofold higher at the beginning of Zn deficiency and four-fold higher under severe Zn deficiency (Cakmak *et al.*, 1996b). Such high accumulation of Fe was also shown in maize and wheat plants grown on Zn-deficient soils (Warnock, 1970; I. Cakmak, unpublished). Leaves of maize plants grown in a Zn-deficient calcareous soil contained 573 mg Fe kg⁻¹ d. wt, but when plants were supplied with adequate Zn the leaves contained only 80 mg Fe kg⁻¹ d. wt (Warnock, 1970). Jackson *et al.* (1967) found up to 1580 mg Fe kg⁻¹ d. wt in leaves of Zn-deficient maize plants grown under field conditions. Application of Zn to maize plants markedly reduced Fe concentration in leaves. Ambler *et al.* (1970) made similar observations on soybean plants growing in nutrition solution. They demonstrated a substantial decrease in Fe concentration in xylem exudate with increasing concentration of Zn from a deficient to an adequate level in nutrient solution.

Accumulation of Fe in Zn-deficient plants is particularly pronounced under high light intensity. In bean plants grown in nutrient solution, increases in light intensity from 80 to 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ enhanced Fe concentration from 204 to 242 mg kg⁻¹ d. wt for Zn-sufficient plants, and from 269 to 1203 mg kg⁻¹ d. wt for Zn-deficient plants (Cakmak, 1988). Such dramatic changes in Fe concentration of Zn-deficient leaves were not found for Mn concentration. It is possible that Zn deficiency has a specific stimulating effect on root uptake and root-to-shoot translocation of Fe. Excessive accumulation of Fe in Zn-deficient plants found in various growth-chamber, glasshouse and field experiments could not be explained by a 'concentration effect' (i.e. by decreased biomass production) resulting from Zn deficiency-induced growth inhibition (Jackson *et al.*, 1967; Warnock, 1970; Brown & Jones, 1977;



Cakmak *et al.*, 1996b). However in some nutrient solution experiments, Zn deficiency either did not affect or even decreased Fe uptake and translocation, although Zn-deficient plants had higher Fe concentrations in shoots (Walter *et al.*, 1994; Rengel *et al.*, 1998a). As discussed below, such high levels of accumulation of Fe in plant tissues can be responsible for excessive production of toxic O_2 species and extensive cellular damage.

(b) *Iron-induced production of free radicals.* High levels of accumulation of Fe in plant or animal cells are responsible for the initiation of severe oxidative stress because they produce ROS by various cellular reactions (Halliwell & Gutteridge, 1984; Price & Hendry, 1991; Hendry, 1993; Becanne *et al.*, 1998). Interestingly, the role of Fe in generating ROS and causing cellular damage in Zn-deficient plants has not been studied, and has only rarely been discussed in the literature.

The powerful oxidant OH^{\cdot} is produced by the Fe-catalysed Haber–Weiss reaction. This reaction involves reduction of Fe(III) by $O_2^{\cdot-}$ to Fe(II) (reaction 1), and reoxidation of Fe(II) by H_2O_2 to OH^{\cdot} (reaction 2), which is known as the Fenton reaction. The requirement of Fe to cause oxidative cell damage is not only dependent on OH^{\cdot} production by the Fenton reaction. There are also certain Fe species which are highly toxic and produced by the reaction of Fe^{2+} with O_2 , for example, the ferryl ($Fe^{2+}O$), perferryl ($Fe^{2+}O_2$) species, or an unknown $Fe^{2+} \dots Fe^{3+}$ complex (Girrotti, 1985; Minotti, 1990; Tang *et al.*, 1997). It has recently been reported that in peroxidation of membrane lipids some Fe^{2+}/O_2 species are more effective than the OH^{\cdot} produced by the Fenton reaction (Qian & Buettner, 1999).



Most of the Fe in plant cells is present in the ferric form (FeIII) and it is stored in phytoferritin in leaves (Marschner, 1995). Iron is also chelated with low-molecular-weight compounds such as citrate, ADP and amino acids. Such ferric compounds can be reduced with $O_2^{\cdot-}$ (reaction 1) or several physiological reductants such as ascorbic acid. The reduced ferrous (FeII) compounds can then be oxidized, causing the production of H_2O_2 (reaction 3) or $O_2^{\cdot-}$ (reaction 4) (Halliwell & Gutteridge, 1984; Toyokuni, 1996).





Increased Fe accumulation (such as that found for Zn-deficient plants) has also been reported for plants subjected to various stress conditions such as root

anoxia, drought and light (Price & Hendry, 1989, 1991; Hendry, 1993). Under these stress conditions accumulation of Fe is associated with enhanced lipid peroxidation and chlorophyll damage. In the case of drought stress, chloroplast membranes enhanced their production of $O_2^{\cdot-}$ in response to Fe accumulation, and Fe-catalysed formation of O_2 radicals has been considered as a major factor contributing to drought damage in plant cells (Price & Hendry, 1989, 1991).

Plants grown under flooded conditions accumulate high levels of Fe (Yamaguchi, 1989; Sahrawat *et al.*, 1996). During exposure of flooded plants to O_2 , oxidation of Fe(II) leads to production of $O_2^{\cdot-}$ which is suggested as a cause of flooding damage to plants (Hendry & Brocklebank, 1985; Hendry, 1993; Neue *et al.*, 1998). Thongbai *et al.* (1999) studied the generation of free radicals in rice plants growing in an Fe-toxic soil and showed that post-anoxic injury in rice plants is related to Fe-induced production of free radicals measured by electron paramagnetic resonance spectroscopy. Such reactions would be more distinct in Zn-deficient plants because of the high accumulation of Fe in these plants under flooded conditions (e.g. in rice, Forno *et al.*, 1975; Neue *et al.*, 1998), and because of their reduced ability to defend against free radical reactions (section IV).



Iron can also stimulate free radical reactions by binding to critical cell constituents such as proteins, phospholipids and DNA. As shown in animal cells, redox cycling of Fe (also Cu) at or near the binding sites of cell constituents causes production of OH^{\cdot} via the Haber–Weiss reaction (reactions 5–6). This reaction has been called site-specific OH^{\cdot} production, and leads to site-specific damage in cells (Girotti, 1985; Thomas *et al.*, 1986; Chevion, 1988). As a result of its extremely short half-life and poor diffusion, the level of OH^{\cdot} should be particularly high at the Fe-binding sites in cells, such as $-SH$ groups and phosphate groups of cell membranes (reactions 5 and 6).







In most cases, Fe and Zn compete for the binding sites on proteins and phospholipids. The ability of Zn to compete with Fe at binding sites on the membrane has been suggested by Girotti *et al.* (1985) and Bray & Bettger (1990) as an explanation for the increased lipid peroxidation under Zn deficiency. They suggest that the existence of adequate amounts of Zn in the vicinity of Fe-binding sites blocks binding of Fe and thus inhibits Fe-catalysed site-specific generation of OH^{\cdot} . In support of this theory, Kunimoto *et al.* (1981) observed that membrane lipids with negative surface charges (i.e. Fe-binding sites) are more rapidly oxidized by the addition of Fe than membrane lipids carrying a positive charge. This result emphasizes the importance of Fe-binding sites of membranes in the generation of free radicals and peroxidative cell damage. Hydroxyl radicals (OH^{\cdot}) produced at Fe-binding sites on membranes can rapidly initiate peroxidation of phospholipids, causing site-specific damage.

Willson (1988) also suggested that protection of biomembranes and maintenance of cellular integrity by Zn are predominantly controlled by binding of Zn to $-SH$ -containing compounds, particularly in membrane proteins. As given schematically in reactions 7 and 8 (Willson, 1988), under Zn-deficient conditions membrane $-SH$ groups can be occupied by Fe, with a concomitant generation of $O_2^{\cdot-}$ and membrane damage (reaction 7). By competing with Fe, or other redox active metals such as Cu(II), for binding to $-SH$ groups of membrane proteins, Zn inhibits metal-induced generation of $O_2^{\cdot-}$ and the related membrane damage (Willson, 1988). When bound to $-SH$ groups or other critical Fe-binding sites of membranes, Zn, as a redox-inactive element in biological systems, cannot undergo a cyclic reduction and reoxidation to produce free radicals (reaction 8).

Evidence for the cellular importance of the binding of Zn to $-SH$ groups was shown *in vitro* by Searle & Tomasi (1982) in a model study. They compared Zn and Fe for their role in the production of the hydroxyl radical (OH^{\cdot}) in the presence of cysteine. When incubated with $FeSO_4$, cysteine (RS^-) complexes with Fe^{2+} (reaction 9), and this Fe^{2+} -cysteine complex is oxidized producing a ferric complex and



$O_2^{\cdot-}$ (reaction 10). The superoxide radical produced reacts with a new Fe^{2+} -cysteine complex to give rise to H_2O_2 (reaction 11). Hydrogen peroxide, which can also be produced by non-enzymatic dismutation of $O_2^{\cdot-}$ (reaction 12), is reduced by Fe^{2+} -cysteine complex to OH^{\cdot} , as in the Fenton reaction (reaction 13).



In the presence of adequate amounts of $ZnSO_4$, Zn^{2+} ions bind to cysteine, blocking binding of Fe to cysteine and thus preventing formation of $Fe^{2+}(RS^-)_2$ and ROS during the redox cycling of Fe-cysteine complex. Zinc is catalytically inert with respect to reactions that produce free radicals when bound to cysteine or other relevant ligands. According to Berg & Shi (1996), specific binding of Zn to tetrahedral sites of a large number of vital cell compounds, and the lack of redox activity of Zn, may be the bases for many of its crucial physiological functions in cells. Accordingly, it has been found that inhibition by Zn of peroxidative damage in animal cells is related to displacement of redox active metals (e.g. Fe and Cu) from cellular binding sites and, consequently, its ability to inhibit the site-specific production of ROS (Girotti *et al.*, 1985; Powell *et al.*, 1994).

Zinc protects DNA-binding proteins (Zn-finger proteins) from reacting with Fe. In these proteins, Zn is tetrahedrally coordinated with cysteinyl-SH groups and imidazole histinyl groups (Vallee & Auld, 1990; Rhodes & Klug, 1993). Under conditions of high-Fe and low-Zn concentrations, Zn bound to the finger protein can be exchanged by Fe(II), and Fe-substituted finger protein, along with H_2O_2 , can induce generation of the potent oxidant OH^{\cdot} (Conte *et al.*, 1996). According to Conte *et al.* (1996), because of the very close proximity of the Zn-finger protein to DNA, Fe-substituted Zn-finger proteins may cause extensive DNA damage via OH^{\cdot} . DNA damage and impairments in gene expression, as described in animal cells, can also be expected in Zn-deficient plant cells containing high Fe concentrations. Further studies are needed on this effect in plants.

3. Zinc deficiency-enhanced photooxidation

(a) *Decrease in photosynthesis.* Chloroplasts are the major cell compartments producing $O_2^{\cdot-}$ and $O_2^{\cdot-}$ -

derived toxic O_2 species such as 1O_2 and OH^{\cdot} (Asada & Takahashi, 1987; Elstner & Osswald, 1994). During photosynthetic electron transport, part of the reducing equivalents generated are transferred to molecular O_2 with the concomitant production of $O_2^{\cdot-}$ in the chloroplastic stroma, which is called the Mehler reaction. It is estimated that under non-stressed conditions up to 20–25% of total non-cyclic photosynthetic electron transport is consumed by the Mehler reaction at light saturation (Robinson, 1988; Osmond & Grace, 1995; Lovelock & Winter, 1996). Production of $O_2^{\cdot-}$ during photosynthetic electron transport can be intensified when plants are exposed to high light intensity combined with an environmental stress factor that restricts photosynthetic CO_2 fixation, such as chilling, drought stress and mineral nutrient deficiencies (Asada *et al.*, 1977; Elstner *et al.*, 1988; Cakmak & Marschner, 1992; Osmond & Grace, 1995; Biehler & Fock, 1996; Polle, 1996; Fryer *et al.*, 1998; Heiser *et al.*, 1998; Cakmak & Engels, 1999). Such conditions lead to limited $NADP^+$ availability for acceptance of electrons from photosystem I, thus intensifying electron flow to O_2 from ferredoxin with a concomitant photooxidative damage to thylakoid constituents by ROS. Under these conditions, conversion of the absorbed light energy into chemical energy can be impaired, causing the transfer of excess energy from the excited triplet chlorophyll state to ground state O_2 to form singlet oxygen (1O_2) (Elstner, 1982; Demming-Adams & Adams, 1992). Singlet oxygen produced in thylakoid membranes is directly responsible for the damage to proteins of the photosystem II reaction centres, especially D1 polypeptide (Hideg *et al.* 1994; Huner *et al.*, 1998).

Photooxidative damage in Zn-deficient leaves can be expected as a result of impaired photosynthetic CO_2 fixation and reduced activity of superoxide dismutase (SOD). In cauliflower, reduction in photosynthesis induced by Zn deficiency is associated with a decrease in intercellular CO_2 concentration and stomatal conductance (Sharma *et al.*, 1994). Stomatal limitation is considered an important reason for reduced photosynthesis in Zn-deficient plants. Sharma *et al.* (1995) reported a significant role of Zn in the regulation of the stomatal aperture. This role of Zn was ascribed to maintenance of a high K concentration in guard cells. A decrease in carbonic anhydrase activity due to Zn deficiency is well known, and may be a factor contributing to reduced photosynthesis (Ohki, 1976; Rengel, 1995a; Cakmak & Engels, 1999). Using two wheat cultivars differing in sensitivity to Zn deficiency, Fischer *et al.* (1997) showed that for both Zn-deficient and Zn-sufficient conditions, a Zn-deficiency-resistant cultivar exhibited a higher rate of net photosynthesis than a sensitive cultivar. The higher level of photosynthesis in Zn-deficiency-resistant cultivars was related to higher carbonic

anhydrase activity, because irrespective of Zn supply the resistant cultivar had an inherently higher carbonic anhydrase activity than the sensitive cultivar (Rengel, 1995a).

Inhibition of photosynthesis in Zn-deficient plants can also be a consequence of a Zn-deficiency-induced reduction in phloem sap sink demand. Severe inhibition of meristem growth is a typical response of plants to Zn deficiency, which is possibly caused by reduced levels of IAA and protein synthesis (Kitagishi *et al.*, 1987; Cakmak *et al.*, 1989). Additionally, in Zn-deficient plants there is an enhanced accumulation of carbohydrates, especially in phloem sap source leaves, possibly resulting from either impaired phloem export of carbohydrates or decreased sink demand (Sharma *et al.*, 1982; Marschner & Cakmak, 1989; Marschner *et al.*, 1996). Inhibition of photosynthesis by accumulation of carbohydrates in leaves has been well documented (Krapp *et al.*, 1991) and may be a core problem in Zn-deficiency-dependent decreases in photosynthetic CO₂ fixation.

(b) *Light-induced leaf chlorosis.* Irrespective of the underlying causes, impairments in photosynthetic CO₂ fixation in Zn-deficient plants may cause a corresponding impairment in utilization of electrons and absorbed light energy for CO₂ fixation. These conditions in Zn-deficient plants may accentuate photogeneration of ROS and photooxidative damage to chloroplasts, as shown schematically in Fig. 2.

Photooxidative damage of chloroplast pigments is particularly pronounced in leaves showing impaired phloem sap sucrose export from source to sink tissues. Inhibition of sucrose phloem transport as a result of pathogen infection of sieve tubes (Osswald & Elstner, 1986); Mg or K deficiency (Cakmak, 1994; Cakmak *et al.*, 1994); petiole phloem destruction (Wood *et al.*, 1986); or high apoplastic invertase activity (Von Schaewen *et al.*, 1990) is associated with a rapid development of leaf chlorosis, particularly under high light intensity (Dickinson *et al.*, 1991; Cakmak & Marschner, 1992). Similarly, enhancements in chlorosis and necrosis due to increased light intensity are very typical in Zn-deficient source leaves, reflected in a massive accumulation of sucrose and starch (Marschner & Cakmak, 1989) causing a high potential for photooxidative damage of chloroplast constituents (Fig. 2). In accordance with this suggestion, enhancements in light intensity from 80 to 490 $\mu\text{mol m}^{-2} \text{s}^{-1}$ markedly depressed shoot elongation and stimulated appearance of leaf chlorosis under Zn deficiency, but not at adequate Zn supply (Cakmak, 1988; Fig. 3a,b). Also, partial shading of Zn-deficient leaves prevented or strongly delayed appearance of chlorosis in the shaded areas (Fig. 3c). Similar results have also been shown for citrus (Cakmak *et al.*, 1995) and rice plants (Obata *et al.*, 1997). In citrus orchards

in California, Zn deficiency chlorosis has been reported to be more severe on branches directly exposed to sunlight (Chapman, 1966). Similarly, in citrus orchards in southern Turkey we observed that under Zn-deficient conditions, Zn deficiency symptoms are very slight in the north-facing parts, while south-facing parts of trees show very severe Zn deficiency chlorosis (Cakmak *et al.*, 1995; Fig. 3d). The enhancements in severity of chlorosis and necrosis under high light intensity were not caused by lower Zn concentrations in leaves. Shaded (green) and non-shaded (chlorotic/necrotic) parts of leaves had about the same Zn concentration (Marschner & Cakmak, 1989; Cakmak *et al.*, 1995). These observations, and results with light-induced chlorosis under Zn deficiency, indicate that increased severity of leaf chlorosis under high light intensity in Zn-deficient conditions is a consequence of photooxidative damage to chloroplast pigments catalysed by ROS. Similarly, in Mg- or K-deficient plants, increases in light intensity markedly stimulated the development of leaf chlorosis and necrosis. The increase in leaf chlorosis under Mg or K deficiency was associated with severe inhibition of phloem transport of sucrose from source leaves, and thus a massive accumulation of carbohydrates in leaves (Marschner & Cakmak, 1989; Cakmak & Marschner, 1992; Cakmak *et al.*, 1994). With the severity of leaf chlorosis in Mg- or K-deficient plants, activities of O₂^{•-}- and H₂O₂-scavenging enzymes showed a substantial increase, indicating enhanced production of ROS in chloroplasts (Cakmak & Marschner, 1992; Cakmak, 1994). Such distinct effects of high light levels on development of leaf chlorosis are very specific for Zn-, Mg- and K-deficient plants, and do not occur in P-deficient plants where activities of antioxidative enzymes are not enhanced, sucrose export is not affected, and carbohydrates do not accumulate in leaves despite severe decreases in shoot growth (Cakmak, 1994; Cakmak *et al.*, 1994).

Photooxidative damage of the chloroplast constituents under Zn deficiency can also be aggravated by reduced activity of enzymes scavenging O₂^{•-} and H₂O₂ in chloroplasts (section IV). As the photooxidative damage is typical for plants exposed to drought, chilling and airborne pollutants such as ozone and SO₂ (Asada, 1994; Foyer *et al.*, 1994; Alscher *et al.*, 1997), the Zn nutritional status of plants under such conditions should become more important. Therefore improvement of the Zn nutritional status of plants can be of crucial importance in protecting plants from such environmental stresses (section V).

(c) *Decrease in indole-3-acetic acid.* In addition to leaf chlorosis and necrosis, decrease in leaf and fruit size and inhibition of shoot elongation are further characteristic morphological changes occurring in Zn-deficient plants. It is widely accepted that these

morphological changes in Zn-deficient plants are attributable to decreased levels of the growth hormone indole-3-acetic acid (IAA) (Skoog, 1940; Marschner, 1995). Zinc-deficiency-dependent decrease in the concentration of IAA (Cakmak *et al.*, 1989; Domingo *et al.*, 1990) is more distinct when plants are grown under higher light intensity (Skoog, 1940; Cakmak, 1988; Römheld & Marschner, 1991). In *in vitro* studies, light-induced oxidative degradation of IAA has been shown, especially in the presence of high Fe (Cakmak, 1988; Dunlop & Robacker, 1988). Iron, by catalysing the Haber-Weiss reaction, is responsible for the production OH^\cdot . IAA is extremely sensitive to OH^\cdot , and can be oxidized rapidly upon exposure to high concentrations of OH^\cdot (Brennan & Frenkel, 1983; Cakmak, 1988; Dunlop & Robacker, 1988). Besides non-enzymic oxidation, IAA is also oxidized by H_2O_2 -dependent peroxidases (Schneider & Wightman, 1975; Gazaryan *et al.*, 1996), and $\text{O}_2^{\cdot-}$ and $^1\text{O}_2$ are involved in this degradation pathway (Metodiewa *et al.*, 1992). Higher peroxidase activity, $\text{O}_2^{\cdot-}$ and Fe in Zn-deficient plants may result in enhanced oxidative degradation of IAA. This could be another relevant reason why Zn-deficient plants become very sensitive to ROS when grown under high-light conditions.

III. MEMBRANE DAMAGE BY REACTIVE OXYGEN SPECIES

1. Impairments in membrane structure

One of the well established cellular functions of Zn is its role in maintaining the structural integrity and controlling permeability of biomembranes. In Zn-deficient plant cells, loss of membrane integrity and increased membrane permeability to inorganic ions were first demonstrated by Welch *et al.* (1982). In

experiments with wheat, they showed that Zn deficiency caused greater leakage of ^{32}P and ^{36}Cl from root cells than in Zn-sufficient plants. Increased leakage of ions was also found in plants with adequate levels of Zn in their roots and shoots when grown in an external medium without Zn supply. In view of these results, Welch *et al.* (1982) concluded that Zn has a direct effect on the structural integrity of biomembranes. To fulfil this action, Zn must be continuously present in the external medium during plant growth. Similar observations were also made in animal studies. According to Lindsay *et al.* (1989), an increase in K leakage from Zn-deficient cells can be ascribed to the protective role of Zn at the apoplasmic side of the plasma membrane. Apparently, Zn reacts with negatively charged molecules associated with the plasma membrane, stabilizing the membrane structure (Lindsay *et al.*, 1989).

In studies with cotton, wheat, tomato and apple plants, Cakmak & Marschner (1988c) showed that leakage not only of inorganic ions, but also of organic compounds from roots, increased in response to Zn deficiency. As shown in Table 2, Zn deficiency was associated with a marked increase in leakage of amino acids, carbohydrates and phenolics from root cells. These increases in membrane permeability were rapidly reversed by a resupply of Zn to Zn-deficient plants (Cakmak & Marschner, 1988c; Zhang *et al.*, 1991). The role of Zn in membrane integrity is thought to be independent of the role of Ca, because increases in membrane permeability caused by Zn deficiency could not be reduced by a supply of Ca to the external medium (Cakmak & Marschner, 1988c). The independent roles of Ca and Zn in membrane integrity were confirmed in a model study with phospholipid vesicles by Kaszuba & Hunt (1990). They showed that both Zn and Ca are




Fig. 2. Schematic representation of the photosynthetic electron flow to CO_2 and O_2 and formation of toxic oxygen species in chloroplasmic stroma. +Zn and -Zn refer to adequate and deficient Zn supply, respectively.




Fig. 3. (a) Effect of different light intensities on shoot growth of 20-d-old bean (*Phaseolus vulgaris*) plants in nutrient solution without Zn supply. (b) Primary leaves of 18-d-old Zn-deficient bean plants grown at light intensities of 80 (left), 230 (middle) and 490 (right) $\mu\text{mol m}^{-2} \text{s}^{-1}$. (c) Effect of partial shading on development of interveinal chlorosis in Zn-deficient primary leaf of 20-d-old bean plants. Partial shading of the leaf was started before the appearance of visual symptoms of chlorosis, and continued for 9 d. (d) Direction effect on Zn deficiency chlorosis in grapefruit trees grown at the field station of the University of Cukurova, Turkey. Zinc deficiency symptoms were severe on sun-exposed sides of trees (south-facing parts) and absent or slight in the north- and north-west-facing parts of trees. Zinc concentrations in leaves from different directions were similarly very low, ranging from 7 to 9 mg kg^{-1} d. wt.

independently required for stabilization of vesicles, as these ions bind to different sites on the membrane.

Enhanced leakage of organic compounds and nutrients from roots of Zn-deficient plants may have some ecological implications such as nutrient mobilization and infection of roots by pathogenic and non-pathogenic microorganisms (Marschner, 1995; Rengel *et al.*, 1998b). Organic compounds released from roots can change the biological and chemical activity of the rhizosphere, thereby changing the availability to plant roots of micronutrients such as Zn, Cu, Ni and Mn (Zhang *et al.*, 1991; Welch, 1995; Rengel *et al.*, 1996; Rengel, 1997).

2. Phospholipids and –SH groups

Increases in membrane permeability are normally associated with peroxidation of membrane constituents, especially unsaturated fatty acids (Demopoulos, 1973; Dhindsa *et al.*, 1981; Girotti, 1985). In cotton roots, reducing the Zn supply reduced levels of fatty acids, particularly unsaturated fatty acids (Table 3). Compared to Zn-adequate cotton roots, saturated fatty acids were decreased in Zn-deficient roots by 22%, and unsaturated fatty acids by 31%. The total amount of phospholipids in roots of Zn-deficient plants showed a similar decrease (Table 3). Unsaturated fatty acids are known to be more susceptible to peroxidation than saturated fatty acids because they contain double bonds (Demopoulos, 1973; Girotti, 1985). Peroxidative damage to biomembranes begins with oxidation of unsaturated bonds in fatty acids and proceeds through a series of free radical reactions, resulting in substantial alteration in membrane structure. Disturbances in the membrane architecture by peroxidative damage impair the integrity and functions of the proteins and other membrane-associated structures, causing membranes to become leaky and unstable (Demopoulos, 1973; Van Ginkel & Sevanian, 1994).

As with unsaturated fatty acids, sulfhydryl (–SH) groups in cell membranes are very sensitive to Zn deficiency. The number of –SH groups in the plasma membrane proteins of root cells was decreased by Zn deficiency in barley (Welch & Norvell, 1993) and in wheat, especially in Zn-deficiency-sensitive wheat cultivars (Table 4). Resupply of Zn to Zn-deficient wheat plants for 24 h clearly increased the number of –SH groups (Rengel, 1995b). This effect of resupplied Zn on –SH groups was ascribed to an increase in SOD activity and a decrease in $O_2^{\cdot-}$ -generating NADPH oxidase in Zn-deficient roots. It could also result from Zn protection of –SH groups in membrane proteins from redox reactions with redox-active transition metals such as Fe(III) and Cu(II). Zinc also protects –SH groups in animal cell membranes. Xia *et al.* (1999) have reported that deficiency of Zn in erythrocyte membranes is associated with important decreases in the levels of –SH groups, and these decreases could be rapidly reversed by a resupply of Zn to Zn-deficient membranes. By affecting the redox status of –SH groups, Zn can exert a significant role in ion-channel gating phenomena or in the activity of transporter proteins in the plasma membranes of root cells, thereby altering ion uptake and efflux rates (Kochian, 1993; Welch & Norvell, 1993; Welch, 1995).

3. Alterations in ion absorption

(a) *Membrane-bound ATPases.* Impairments in the integrity of cell membranes of Zn-deficient plants can cause alterations in the activity of membrane-bound proton-pumping enzymes and ion channels, and thus affect the uptake of nutrients across the plasma membranes. Limited data exist in the literature concerning the role(s) of Zn in membrane-bound ATPase activity. In plasma membrane-enriched vesicles isolated from bean roots, Zn

Table 2. Concentrations of amino acids, sugars and phenolics in the root exudates of cotton, wheat and apple plants grown with different Zn supplies

Zn supply (M)	Amino acids	Sugars	Phenolics
($\mu\text{g g}^{-1}$ d. wt root per 6 h)			
Cotton			
8×10^{-7} (adequate)	48 ± 11	375 ± 41	117 ± 22
0 (deficient)	165 ± 16	751 ± 273	161 ± 31
Wheat			
8×10^{-7} (adequate)	21 ± 2	315 ± 72	34 ± 6
0 (deficient)	48 ± 3	615 ± 61	80 ± 6
Apple			
5×10^{-7} (adequate)	12 ± 3	275 ± 13	103 ± 38
0 (deficient)	55 ± 13	823 ± 36	350 ± 121

Data represent means \pm SD of 4 independent replications (Cakmak & Marschner, 1988c).

Table 3. Concentrations of fatty acids and total phospholipids in 20-d-old cotton roots supplied with different Zn concentrations

	Zn treatment		
	Zn ₀	Zn ₁	Zn ₂
	(µg g ⁻¹ root f. wt)		
Fatty acids			
Total	1123 ± 48(73)	1335 ± 151(87)	1531 ± 50
Saturated	532 ± 26(73)	613 ± 44(91)	677 ± 44
Unsaturated	591 ± 62(69)	721 ± 114(84)	854 ± 55
Total phospholipids	1530 ± 60(69)	1700 ± 240(76)	2230 ± 111

Zn₂, 8 × 10⁻⁷ M, adequate Zn supply; Zn₁, 2 × 10⁻¹ M; Zn₀, 0. Values in parentheses are percentage of Zn₂ treatment. Data represent means ± SD of 3 independent replications (Cakmak & Marschner, 1988c).

Table 4. Concentrations of reactive sulfhydryl groups in the root-cell plasma membranes of a Zn deficiency-sensitive durum wheat (*cv. Durati*) and Zn deficiency-resistant bread wheat (*cv. Warigal*) grown with sufficient (+Zn) and deficient (-Zn) Zn supply in nutrient solution

Cultivar	Zn supply		
	Deficient	Sufficient	Mean
	(µmol -SH g ⁻¹ root d. wt)		
'Durati' (sensitive)	3.9	5.3	4.6
'Warigal' (resistant)	6.0	7.0	6.5
Mean	4.9	6.2	

Data represent means of 6 replications. The overall means for Zn treatments and the genotypes are given because the interaction effect was not significant ($P \leq 0.50$). The HSD_{0.05} value is 0.7 µmol g⁻¹ root d. wt for both effects (Rengel, 1995b).

deficiency reduced vanadate-sensitive ATP hydrolysis and, more markedly, ATP-dependent proton-pumping activity, resulting in a decrease in pH gradient across the membrane (Pinton *et al.*, 1993). A resupply of Zn to Zn-deficient plants for 24 h evidently restored the proton pumping activity of membrane vesicles (Pinton *et al.* 1993). These effects of Zn substantiate the importance of Zn in maintaining the proton gradient across membranes. Accordingly, in studies with excised maize roots it has been shown that the membrane potential of root cells is higher in the presence than in the absence of Zn, possibly because of the stimulation by Zn of plasma-membrane-bound ATPase activity (Kennedy & Gonsalves, 1987). These authors suggested that Zn²⁺-ATP might be a substrate for plasma membrane ATPase, and higher membrane potential maintained by the presence of Zn benefits the uptake of nutrients. In a further study with the plasma membrane vesicles isolated from corn roots, Kastrop *et al.* (1996) showed that Zn in the form of ZnATP²⁻, but not free Zn²⁺ activated H⁺-ATPase and demonstrated the importance of Zn in regulating proton-pumping activity of plasma membranes and nutrient

uptake by plant roots. Decreases in plasma-membrane-bound ATPase activity due to Zn deficiency were also found in chickpea roots, and were attributed to a modification of the enzyme complex resulting from insufficient membrane Zn (Lawrence *et al.*, 1995).

Disturbances in the activity of membrane-bound ATPases caused by Zn deficiency may be a consequence of free radical attack on enzymes. For example, Na⁺/K⁺-ATPase and Ca-ATPase activities in animal cells are very sensitive to oxidative damage by ROS. Oxidation of -SH groups of enzymes appears to be the major reason for loss of enzyme activity in stressed cells. In studies with animal cells, Kurella *et al.* (1995) reported the existence of a relationship between the decrease in Na⁺/K⁺-ATPase activity and oxidation of -SH groups in the enzyme by free radicals. Oxidation of one or two -SH groups per mole enzyme resulted in a 43% decrease in enzyme activity, and with oxidation of an additional three to four -SH groups per mole enzyme there was a decrease of 70% in enzyme activity. Similarly, in plant cells oxidation of ATPase -SH groups in purified plasma membrane vesicles

from *Elodea canadensis* resulted in a significant loss of ATPase activity (Elzenga *et al.*, 1989).

(b) *Nutrient uptake.* Impaired activity of the ATP-dependent proton gradient across the plasma membrane may result in changes in ion absorption by root cells. Among the mineral nutrients, uptake of P was especially influenced by Zn deficiency. In a number of plant species, Zn deficiency is associated with a marked increase in P uptake by roots (Loneragan *et al.*, 1982; Cakmak & Marschner, 1986; Rengel & Graham, 1995; Parker, 1997). As a consequence, symptoms of Zn deficiency can be mistaken for symptoms of P toxicity. In contrast to P, net uptake of NO_3^- was substantially depressed by Zn deficiency (Marschner & Cakmak, 1986; Cakmak & Marschner, 1990; Norvell & Welch, 1993). Decreases in net NO_3^- uptake are attributed to enhanced net efflux of NO_3^- through the plasma membranes, due to its enhanced membrane permeability under Zn deficiency (Cakmak & Marschner, 1990). Zinc deficiency also alters the uptake of other nutrient elements such as Fe (section II, 2a), B, Mg, K, Mn and Cl, but not consistently (Marschner & Cakmak, 1986; Graham *et al.*, 1987; Norvell & Welch, 1993; Parker, 1997). As shown for uptake of micronutrients and P in different wheat genotypes, the effects of Zn deficiency on uptake of mineral nutrients can be genotype-dependent (Rengel & Graham, 1995).

Zinc sufficiency is also necessary to control the uptake and accumulation of Na in plants (Shukla & Prasad, 1974; Norvell & Welch, 1993) and in algae (Rybol *et al.*, 1992). Increased Na uptake under Zn deficiency is possibly related to impairments in membrane stability and thus a loss of the cell's ability to control Na uptake. Using yeast protoplasts, Kovac *et al.* (1987) showed that Zn has a particular role in maintaining membrane stability and protecting membranes against osmotic stress. In the absence of Zn, plasma membranes of yeast protoplasts lost their stability and became very permeable, leaking intercellular solutes. As suggested by Norvell & Welch (1993), improvement of the Zn nutritional status of plants growing in saline soils of arid and semi-arid regions is important not only to overcome Zn-deficiency-related decreases in plant productivity, but also to protect plants against excessive salinity and resulting injury.

(c) *Changes in activity of ion channels.* Significant progress has been made in understanding the importance of ion channels in ion transport across the membranes. Ion channels are biologically very active proteins that permit rapid and selective transport of ions depending on their opening and closure status (Tester, 1990). Ion channel proteins contain reactive -SH groups, and the activity of the channel is dependent on the redox status of these -SH groups. There is some evidence of involvement

of Zn in regulating the activity of ion channels in animal cells. In frog muscle cells a low concentration of extracellular Zn caused an increase in K efflux via K channels (Spalding *et al.*, 1986). This effect of Zn was attributed to the binding of Zn to histidine residues in the K channel protein. Zinc may also interact with Cl^- channels. In *Xenopus* oocytes, extracellular Zn reduced Cl^- efflux via a Cl^- channel by facilitating the slow gating of the channel (Chen, 1998). Reduction in Cl^- efflux across Cl^- channels, resulting from increasing concentration of extracellular Zn, has been explained by the binding of Zn to channel molecules in the ionic form, ZnOH^+ (Spalding *et al.*, 1990).

In plant and algal cells Zn exerts similar inhibitory roles on efflux of ions via anion channels (Hedrich & Kurkdjian, 1988; Tyerman, 1992). Using isolated vacuoles from sugar beet, Hedrich & Kurkdjian (1988) showed that Zn strongly reduces activity of the Ca-activated, slow vacuole-type anion channels. This blocking action of Zn on channel activity occurs in a few minutes of exposure of vacuoles to a very low Zn concentration ($0.1 \mu\text{M ZnCl}_2$). In wheat and barley seedlings, Welch and co-workers (Welch *et al.*, 1982; Norvell & Welch, 1993; Welch & Norvell, 1993) reported that Zn-deficient root cells enhanced leakage of Cl^- compared with Zn-sufficient plants. It seems likely that Zn may affect the activity of the putative Cl^- channels in root cell membranes by interacting with -SH groups in sulphur-containing amino acid constituents of membrane peptides.

In view of decreases in the amount of -SH groups in the root cell plasma membranes caused by Zn deficiency, Welch & Norvell (1993) suggested that Zn is required for protection of membrane -SH groups against free radicals, thus influencing activity of ion transport proteins as well as ion channels. Alterations in influx and efflux of ions in Zn-deficient plants, compared with Zn-sufficient ones, may at least in part be attributed to the redox state of reactive -SH groups contained in ion channels.

In most cases, unregulated Ca accumulation in the cytosol initiates programmed cell death (Dangl *et al.*, 1996; Levine *et al.*, 1996). Toxic O_2 species are involved in increased concentrations of cytosolic Ca^{2+} because they oxidize critical residues in Ca channel proteins. In animal cells, Ca efflux from Ca-accumulating organelles is enhanced when -SH residues in Ca channels are oxidized by toxic O_2 species (Stoyanovsky *et al.*, 1994; Donoso *et al.*, 1997). Similarly, in Zn-deficient plant cells an enhanced free radical-dependent oxidation of -SH groups and other critical residues of Ca channels might be expected. This could result in an enhanced Ca efflux into the cytosol from Ca-accumulating organelles (e.g. vacuole and endoplasmic reticulum), thereby causing a dramatic increase in

cytosolic Ca concentration. Protection of the critical –SH groups in Ca channels by Zn has been reported in different animal cells (Chiamvimonvat *et al.*, 1995, cited by Xia *et al.*, 1999). Protection of –SH groups by improved Zn nutrition may be provided by reduced activity of $O_2^{\cdot-}$ -generating NADPH oxidase (section II, 1). Hypothetically, maintenance of low cytosolic Ca concentration may be impaired by Zn deficiency, triggering programmed cell death. For plant cells, there are no studies or discussion in the literature related to the death of Zn-deficient cells as a possible consequence of increased cytosolic Ca concentration or DNA degradation. This point needs to be clarified in future studies.

IV. DETOXIFICATION OF REACTIVE OXYGEN SPECIES

1. Superoxide dismutases

The superoxide radical and its derivatives are unavoidable products of normal cell metabolism, and their generation is particularly high during electron transport in chloroplasts and mitochondria (Cakmak *et al.*, 1993; Elstner & Osswald, 1994; Polle, 1996; Alscher *et al.*, 1997). To minimize the destructive effects of ROS, plant cells are equipped with various antioxidants and antioxidative defence enzymes such as superoxide dismutases (SODs), ascorbate peroxidases, ascorbic acid, α -tocopherol and carotenoids (Asada, 1994; Cakmak, 1994; Foyer *et al.*, 1994). By catalysing detoxification of $O_2^{\cdot-}$ to O_2 and H_2O_2 and blocking $O_2^{\cdot-}$ -driven cell damage, SODs are a major component of the antioxidative defence system of plant cells (Fridovich, 1986; Bowler *et al.*, 1994). According to their metal cofactor, SODs are classified into three types containing either Mn (Mn-SOD), Fe (Fe-SOD) or Cu and Zn (CuZn-SOD). In general, Mn-SOD is located in mitochondria, Fe-SOD in chloroplasts, and CuZn-SOD in chloroplasts and the cytosol (Bowler *et al.*, 1994). Manganese-containing SOD is also present in the peroxisomes (Sandali *et al.*, 1987).

Three types of SOD are distinguished according to their sensitivity to cyanide and H_2O_2 . Activity of CuZn-SOD is sensitive to both H_2O_2 and cyanide, while Fe-SOD is not sensitive to cyanide, but H_2O_2 inhibits its activity. Mn-SOD is not affected by H_2O_2 or cyanide (Scandalios, 1993; Bowler *et al.*, 1994). Measurement of these isoforms of SOD is useful for determining the micronutrient status of plants, for example in Zn (Cakmak & Marschner, 1987; Cakmak *et al.*, 1997), Fe (Iturbe-Ormaetxe *et al.*, 1995) and Mn (Garcia *et al.*, 1981; Yu *et al.*, 1998).

In higher plants, CuZn-SOD is the most abundant SOD, while Mn-SOD and Fe-SOD form a smaller proportion of total SOD activity (Jackson *et al.*, 1978; Alscher *et al.*, 1997). In leaves of different

wheat cultivars, deficient supply of Zn decreased total SOD activity and, more distinctly, CuZn-SOD activity, whereas Mn-SOD activity was not affected by Zn deficiency (Cakmak *et al.*, 1997, 1998). Recently, similar results were found by Yu *et al.* (1999) in various wheat genotypes differing in sensitivity to Zn deficiency. They measured both activity and concentration of CuZn-SOD using capillary electrophoresis, and suggested that CuZn-SOD can be used as a tool in assessing the genotypic variation in sensitivity to Zn deficiency. Despite severe decreases in CuZn-SOD activity, there is no compensatory increase in Mn-SOD activity in Zn-deficient plants. Cakmak *et al.* (1997), studying wheat, and Yu & Rengel (1999), working on lupin, found that Zn deficiency markedly reduced CuZn-SOD activity, but did not affect Mn-SOD activity. However, under Mn deficiency decreases in Mn-SOD were accompanied by increases in CuZn-SOD activity (Del Rio *et al.*, 1978; Yu & Rengel, 1999). As in leaves, Zn deficiency lowered SOD activity in roots of cotton plants, and these decreases in SOD activity were paralleled by increases in the amount of $O_2^{\cdot-}$ measured by ESR in the cytosol of root cells (Cakmak & Marschner, 1988a,b). Decreases in CuZn-SOD activity caused by Zn deficiency have also been shown in bean (Cakmak & Marschner, 1993; Wenzel & Mehlhorn, 1995), *Lemna gibba* (Vaughan *et al.*, 1982) and tobacco (Yu *et al.*, 1998).

Irrespective of the Zn nutritional status of plants, CuZn-SOD activity can be decreased when plants are exposed to a strong photooxidative stress or conditions producing high amounts of H_2O_2 . In plant and animal cells, H_2O_2 is effective in inactivating and degrading CuZn-SOD (Strack *et al.*, 1996; Casona *et al.*, 1997). Inhibition of CuZn-SOD activity by H_2O_2 was ascribed to reduction of enzyme-bound Cu^{2+} to Cu^+ by H_2O_2 ; the reduced Cu reacts with H_2O_2 , resulting in production of OH^{\cdot} . This hydroxyl radical was suggested by Hodgson & Fridovich (1975) to be responsible for inactivation of CuZn-SOD. Inactivation of SOD can also occur in plants with high sensitivity to low temperature (Michalski & Kaniuga, 1981). This inactivation was attributed to an inhibitory action of H_2O_2 which is accumulated at low temperatures.

2. H_2O_2 -scavenging enzymes

Higher sensitivity of Zn-deficient plants to oxidative stress conditions cannot be attributed solely to reduced CuZn-SOD activity. Zinc is also indirectly required for high activity of the enzymes involved in H_2O_2 detoxification such as catalase, ascorbate peroxidase and glutathione reductase. In Zn-deficient plants activity of catalase was reduced, although Zn is not a required cofactor for catalase activity. This decrease in catalase activity with Zn deficiency was assumed to be related to inhibition of

catalase by $O_2^{\cdot-}$ (Cakmak & Marschner, 1988a). The catalase enzyme is very sensitive to $O_2^{\cdot-}$ and can be inactivated by increased levels of $O_2^{\cdot-}$ (Fridovich, 1986). Because Zn deficiency strongly reduces protein synthesis (Kitagishi & Obata, 1986; Cakmak *et al.*, 1989; Marschner, 1995), biosynthesis of H_2O_2 -scavenging enzymes can be impaired as a result of Zn deficiency. Accordingly, in bean plants Zn deficiency reduced the activities of glutathione reductase and ascorbate peroxidase in young leaves, although these enzymes do not need Zn for their activity (Cakmak & Marschner, 1993). Consequently, because of the reduced activity of enzymes scavenging $O_2^{\cdot-}$ and H_2O_2 , enhanced production of OH^{\cdot} (via the Haber-Weiss reaction) can be expected in Zn-deficient plants. However, concentrations of OH^{\cdot} in Zn-deficient plants have not been reported.

V. INVOLVEMENT OF ZINC IN PLANT STRESS TOLERANCE

As SODs play critical roles in the oxidative defence systems of all biological tissue (Bowler *et al.*, 1992, 1994; Scandalios, 1993), it can be suggested that plants with reduced SOD activity (i.e. marginal or severe Zn deficiency) should be highly sensitive to oxidative stress factors such as drought, chilling, O_3 , pathogenic infection and salinity. There are results which support this suggestion. In bush bean, Zn deficiency caused a high sensitivity to O_3 toxicity (Wenzel & Mehlhorn, 1995). Ozone is known to exert its deleterious effect on plant cells by producing toxic O_2 species such as $O_2^{\cdot-}$ (Sharma & Davis, 1997). The higher O_3 sensitivity of Zn-deficient bush bean plants correlated well with a reduced activity of CuZn-SOD, prompting Wenzel & Mehlhorn (1995) to suggest that CuZn-SOD plays an important role in plant defence systems against O_3 toxicity. Similarly, overproduction of CuZn-SOD in the cytosol in response to O_3 exposure protected transgenic tobacco plants from O_3 damage (Pitcher & Zilinskas, 1996). Mn-SOD and Fe-SOD were also overexpressed in chloroplasts from plants exposed to O_3 , thereby contributing to plant tolerance against O_3 toxicity (Van Camp *et al.*, 1994, 1996).

Besides protecting against O_3 toxicity, increased activity of CuZn-SOD also enhanced plant resistance to other oxidative stress factors. An NaCl-tolerant cultivar of pea, compared with a sensitive cultivar, had enhanced chloroplastic and cytosolic CuZn-SOD activity when treated with NaCl (Hernandez *et al.*, 1995). In transgenic tobacco plants, salt stress was associated with overexpression of cytosolic and chloroplastic CuZn-SOD enzymes (Van Camp *et al.*, 1996). Drought stress represents another oxidative environmental factor causing enhanced production of $O_2^{\cdot-}$ and H_2O_2 , particularly in chloroplasts (Price & Hendry, 1989, 1991; Smirnov, 1993; Bohnert *et al.*, 1995). In pea plants, decreases in leaf water

potential caused by withholding water resulted in higher cytosolic and chloroplastic CuZn-SOD activity. This was considered an adaptive mechanism for minimizing drought-related cell damage in plants (Mittler & Zilinskas, 1994).

Increased activity of CuZn-SOD and its involvement in stress tolerance has also been shown in plants affected by low temperatures. Chilling stress, in combination with higher light intensity, is responsible for extensive photooxidative damage to chloroplasts by ROS (Wise & Naylor, 1987; Long *et al.*, 1994). Such photooxidative damage was greatly decreased in transgenic tobacco plants that were capable of overexpressing genes encoding the chloroplastic CuZn-SOD enzymes (Sen Gupta *et al.*, 1993a,b). Overexpression of CuZn-SOD genes was also demonstrated in plants exposed to herbicide treatment (Perl *et al.*, 1993), sulphur dioxide (Madamanchi *et al.*, 1994), UV-B radiation (Willekens *et al.*, 1994; Rao *et al.*, 1996) and pathogenic infection (Mittler *et al.*, 1996; Fodor *et al.*, 1997). As shown under field conditions, chilling stress in citrus and drought stress in wheat became more pronounced in plants suffering from Zn deficiency (Cakmak *et al.*, 1995; Ekiz *et al.*, 1998). These results may be due to reduced activity of enzymes scavenging $O_2^{\cdot-}$ and H_2O_2 in Zn-deficient tissues. In wheat plants, decreases in grain yield due to drought stress were more marked when plants were Zn-deficient (Ekiz *et al.*, 1998). By affecting the synthesis and activity of antioxidative enzymes, Zn is an important factor in plant defence systems against destructive O_2 species. Thus improvement of the Zn nutritional status of plants may be of great importance for their survival under oxidative stress conditions (drought, chilling, high light levels, ozone and salinity).

VI. CONCLUSIONS

This review presents several lines of evidence demonstrating that enhanced production of ROS is primarily involved in Zn-deficiency-induced impairments in cellular function and integrity. Zinc performs various important roles in protecting cells from the damaging reactions caused by ROS. As summarized in Fig. 4, Zn is directly or indirectly required for scavenging $O_2^{\cdot-}$ and H_2O_2 , and thus for blocking generation of the powerful oxidant OH^{\cdot} . In Zn-deficient cells, Fe accumulation is substantial and results in a high physiological demand for Zn, particularly at membrane-binding sites for Fe. In the absence of Zn, binding of Fe to membrane constituents or DNA leads to site-specific production of OH^{\cdot} . Zinc also interferes with $O_2^{\cdot-}$ -generating oxidases located in plasma membranes. Zinc is particularly needed within the environment of plasma membranes to maintain their structural and functional integrity. These Zn-deficiency-




Fig. 4. Major physiological changes occurring in Zn-deficient cells and the effect on plant growth.

related disturbances in cellular metabolism are responsible for oxidative damage of membrane proteins, phospholipids, chlorophyll, nucleic acids, SH-containing enzymes and IAA, and thus inhibition of plant growth. Recent reports demonstrate that shoot and root meristematic activities of plants are rapidly blocked under oxidative stress conditions as a result of DNA damage (Reichheld *et al.*, 1999). Very high concentrations of Zn in meristematic plant cells (Kitagishi & Obata, 1986; Hossain *et al.*, 1997) demonstrate the crucial roles played by Zn in highly metabolically active differentiating cells.

ACKNOWLEDGEMENTS

I thank Prof. Dr Ross Welch (Cornell University, US Soil, Plant and Nutrition Laboratory) and Dr Zed Rengel (Western Australia University) for constructive comments on the manuscript and correction of the English text. Financial support from the Alexander von Humboldt Foundation is gratefully acknowledged.

REFERENCES

- Allen RD. 1995. Dissection of oxidative stress tolerance using transgenic plants. *Plant Physiology* **107**: 1049–1054.
- Alscher RG, Donahue JL, Cramer CL. 1997. Reactive oxygen species and antioxidants: relationships in green cells. *Physiologia Plantarum* **100**: 224–233.
- Ambler JE, Brown JC, Gauch HG. 1970. Effect of zinc on translocation of iron in soybean plants. *Plant Physiology* **46**: 320–323.
- Asada K. 1994. Production and action of active oxygen species in photosynthetic tissues. In: Foyer CH, Mullineaux PM, eds. *Causes of photooxidative stress and amelioration of defense systems in plants*. Boca Raton, FL, USA: CRC Press, 77–104.
- Asada K, Takahashi M. 1987. Production and scavenging of active oxygen in photosynthesis. In: Kyle DJ, Osmond CB, Arntzen CJ, eds. *Photoinhibition*. Amsterdam, The Netherlands: Elsevier, 227–287.
- Asada K, Takahashi MA, Tanaka K, Nakano Y. 1977. Formation of active oxygen and its fate in chloroplasts. In: Hayaishi O, Asada K, eds. *Biochemical and medical aspects of active oxygen*. Tokyo, Japan: University Park Press, 45–63.
- Becanne M, Moran JF, Iturbe-Ormaetxe I. 1998. Iron-dependent oxygen free radical generation in plants subjected to environmental stress: toxicity and antioxidant protection. *Plant and Soil* **201**: 137–147.
- Berg JM, Shi Y. 1996. The galvanization of biology: a growing appreciation for the roles of zinc. *Science* **271**: 1081–1085.
- Biehler K, Fock H. 1996. Evidence for the contribution of the Mehler-peroxidase reaction in dissipating excess electrons in drought-stressed wheat. *Plant Physiology* **12**: 265–272.
- Bohnert HJ, Nelson D, Jensen RG. 1995. Adaptations to environmental stress. *Plant Cell* **7**: 1099–1111.
- Bolwell GP, Wojtaszek P. 1997. Mechanisms for the generation of reactive oxygen species in plant defence – a broad perspective. *Physiological and Molecular Plant Pathology* **51**: 347–366.
- Bowler C, Camp WV, Mantogu MV, Inzé D. 1994. Superoxide dismutase in plants. *Critical Reviews in Plant Science* **13**: 199–218.
- Bowler C, Van Montagu M, Inzé D. 1992. Superoxide dismutase and stress tolerance. *Annual Review of Plant Physiology and Plant Molecular Biology* **43**: 83–116.
- Bray TM, Bettger WJ. 1990. The physiological role of zinc as an antioxidant. *Free Radicals in Biology and Medicine* **8**: 281–291.
- Brennan T, Frenkel C. 1983. Non-enzymatic oxidation of indole-3-acetic acid by H₂O₂ and Fe²⁺ ions. *Botanical Gazette* **144**: 32–36.
- Brown JC, Jones WE. 1977. Fitting plants nutritionally to soils. II. Cotton. *Agronomy Journal* **69**: 405–409.
- Brown PH, Cakmak I, Zhang Q. 1993. Form and function of zinc in plants. In: Robson AD, ed. *Zinc in soils and plants*. Dordrecht, The Netherlands: Kluwer Academic, 93–106.
- Burke JP, Fenton MR. 1985. Effects of zinc-deficient diet on lipid peroxidation in liver and tumour subcellular membranes. *Proceedings of the Society for Experimental Biology and Medicine* **179**: 187–191.
- Cakmak I. 1988. *Morphologische und Physiologische Veraenderungen in Zinkmangelpflanzen*. PhD thesis, University of Hohenheim–Stuttgart, Germany.
- Cakmak I. 1994. Activity of ascorbate-dependent H₂O₂-scavenging enzymes and leaf chlorosis are enhanced in magnesium and potassium deficient leaves, but not in phosphorus deficient leaves. *Journal of Experimental Botany* **45**: 1259–1266.
- Cakmak I, Atli M, Kaya R, Evliya H, Marschner H. 1995. Association of high light and zinc deficiency in cold induced leaf chlorosis in grapefruit and mandarin trees. *Journal of Plant Physiology* **146**: 355–360.

- Cakmak I, Engels C. 1999.** Role of mineral nutrients in photosynthesis and yield formation. In: Rengel Z, ed. *Mineral nutrition of crops*. New York, USA: Haworth Press, 141–168.
- Cakmak I, Hengeler C, Marschner H. 1994.** Changes in phloem export of sucrose in leaves in response to phosphorus, potassium and magnesium deficiency in bean plants. *Journal of Experimental Botany* **45**: 1251–1247.
- Cakmak I, Kalayci M, Ekiz H, Braun HJ, Yilmaz A. 1999.** Zinc deficiency as an actual problem in plant and human nutrition in Turkey: a NATO – Science for Stability Project. *Field Crops Research* **60**: 175–188.
- Cakmak I, Marschner H. 1986.** Mechanism of phosphorus induced zinc deficiency in cotton. I. Zinc deficiency-enhanced uptake rate of phosphorus. *Physiologia Plantarum* **68**: 483–490.
- Cakmak I, Marschner H. 1987.** Mechanism of phosphorus induced zinc deficiency in cotton. III. Changes in physiological availability of zinc in plants. *Physiologia Plantarum* **70**: 13–20.
- Cakmak I, Marschner H. 1988a.** Enhanced superoxide radical production in roots of zinc deficient plants. *Journal of Experimental Botany* **39**: 1449–1460.
- Cakmak I, Marschner H. 1988b.** Zinc-dependent changes in ESR signals, NADPH oxidase and plasma membrane permeability in cotton roots. *Physiologia Plantarum* **73**: 182–186.
- Cakmak I, Marschner H. 1988c.** Increase in membrane permeability and exudation in roots of zinc deficient plants. *Journal of Plant Physiology* **132**: 356–361.
- Cakmak I, Marschner H. 1990.** Decrease in nitrate uptake and increase in proton release in zinc deficient cotton, sunflower and buckwheat plants. *Plant and Soil* **129**: 261–268.
- Cakmak I, Marschner H. 1992.** Magnesium deficiency and high light intensity enhance activities of superoxide dismutase, ascorbate peroxidase and glutathione reductase in bean leaves. *Plant Physiology* **98**: 1222–1227.
- Cakmak I, Marschner H. 1993.** Effect of zinc nutritional status on activities of superoxide radical and hydrogen peroxide scavenging enzymes in bean leaves. *Plant and Soil* **155/156**: 127–130.
- Cakmak I, Marschner H, Bangerth F. 1989.** Effect of zinc nutritional status on growth, protein metabolism and levels of indole-3-acetic acid and other phytohormones in bean (*Phaseolus vulgaris* L.). *Journal of Experimental Botany* **40**: 405–412.
- Cakmak I, Oztürk L, Eker S, Torun , Kalfa H I, Yilmaz A. 1997.** Concentration of zinc and activity of copper/zinc-superoxide dismutase in leaves of rye and wheat cultivars differing in sensitivity to zinc deficiency. *Journal of Plant Physiology* **151**: 91–95.
- Cakmak I, Sari N, Marschner H, Kalayci M, Yilmaz A, Eker S, Gulut K. 1996b.** Dry matter production and distribution of zinc in bread and durum wheat genotypes differing in zinc efficiency. *Plant and Soil* **180**: 173–181.
- Cakmak I, Strbac D, Marschner H. 1993.** Activities of hydrogen peroxide-scavenging enzymes in germinating wheat seeds. *Journal of Experimental Botany* **44**: 127–132.
- Cakmak I, Torun B, Erenoglu B, Oztürk L, Marschner H, Kalayci M, Ekiz H, Yilmaz A. 1998.** Morphological and physiological differences in cereals in response to zinc deficiency. *Euphytica* **100**: 349–357.
- Cakmak I, Van Wetering DAM, Marschner H, Bienfait HF. 1987.** Involvement of superoxide radical in extracellular ferric reduction by iron deficient bean roots. *Plant Physiology* **85**: 310–314.
- Cakmak I, Yilmaz A, Ekiz H, Torun B, Erenoglu B, Braun HJ. 1996a.** Zinc deficiency as a critical nutritional problem in wheat production in Central Anatolia. *Plant and Soil* **180**: 165–172.
- Casona LC, Gomez LD, Lascano HR, Gonzalez AC, Trippi VS. 1997.** Inactivation and degradation of CuZn-SOD by active oxygen species in wheat chloroplasts exposed to photo-oxidative stress. *Plant and Cell Physiology* **38**: 433–440.
- Chapman HD. 1966.** Zinc. In: Chapman HD, ed. *Diagnostic criteria for plants and soils*. University of California, Riverside, CA, USA: Division of Agricultural Sciences, 484–499.
- Chen TY. 1998.** Extracellular zinc ion inhibits CIC-0 chloride channels by facilitating slow gating. *Journal of General Physiology* **112**: 715–726.
- Chevion M. 1988.** A site-specific mechanism for free radical induced biological damage: the essential role of redox-active transition metals. *Free Radical Biology and Medicine* **5**: 27–37.
- Chiamvimonvat N, O'Rourke B, Kallen RG, Hofmann F, Flockerzi V, Marban E. 1995.** Functional consequences of sulfhydryl modification in pore-forming subunits of cardiovascular Ca^{2+} and Na^{+} channels. *Circulation Research* **76**: 325–334.
- Chvapil M. 1979.** Zinc and NADPH-oxidation-dependent lipid peroxidation. In: *Oxygen free radicals and tissue damage*. CIBA Foundation Symposium (New Series) No. 65. Amsterdam, The Netherlands: Excerpta Medica, 163–166.
- Conte D, Narindrasorasak S, Sarkar B. 1996.** *In vivo* and *in vitro* iron-replaced zinc finger generates free radicals and causes DNA damage. *Journal of Biological Chemistry* **271**: 5125–5130.
- Cross AR, Erichson R, Ellis BA, Curnutte JT. 1999.** Spontaneous activation of NADPH oxidase in a cell-free system: unexcepted multiple effects of magnesium ion concentrations. *Biochemical Journal* **338**: 229–233.
- Dangl JL, Dietrich RA, Richberg MH. 1996.** Death don't have no mercy: cell death programs in plant-microbe interactions. *Plant Cell* **8**: 1793–1807.
- Del Rio LA, Pastori GM, Palma JM, Sandalio LM, Sevilla F, Corpas FJ, Jiménez A, Lopez-Huertas E, Hernandez JA. 1998.** The activated oxygen role of peroxisomes in senescence. *Plant Physiology* **166**: 1195–1200.
- Del Rio LA, Sevilla F, Gomez M, Yanez J, Lopez-Gorge J. 1978.** Superoxide dismutase: an enzyme system for the study of micronutrient interactions in plants. *Planta* **140**: 221–225.
- Demming-Adam B, Adams WW. 1992.** Photoprotection and other responses of plants to high light stress. *Annual Review of Plant Physiology and Plant Molecular Biology* **43**: 599–626.
- Demopoulos HB. 1973.** The basis of free radical pathology. *Federation Proceedings* **32**: 1859–1861.
- Dhindsa RS, Plumb-Dhindsa P, Thorpe TA. 1981.** Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *Journal of Experimental Botany* **32**: 93–101.
- Dickinson CD, Altabella T, Chrispeels MJ. 1991.** Slow-growth phenotype of transgenic tomato expressing apoplastic invertase. *Plant Physiology* **95**: 420–425.
- Dietrich RA, Richberg MH, Schmidt R, Dean C, Dangl JL. 1997.** A novel zinc finger protein is encoded by the *Arabidopsis* *LSD1* gene and functions as a negative regulator of plant cell death. *Cell* **88**: 685–694.
- Doke N, Miura Y, Sanchez LM, Park HJ, Noritake T, Yoshioka H, Kawaika K. 1996.** The oxidative burst protects plants against pathogen attack: mechanism and role as an emergency signal for plant bio-defense – a review. *Gene* **179**: 45–51.
- Domingo AL, Nayotome Y, Tamai M, Takaki H. 1990.** Indole carboxylic acid in zinc-deficient radish shoots. *Soil Science and Plant Nutrition* **36**: 555–560.
- Donoso P, Rodriguez P, Marambio P. 1997.** Rapid kinetic studies of SH oxidation-induced calcium release from sarcoplasmic reticulum vesicles. *Archives of Biochemistry and Biophysics* **341**: 295–299.
- Dunlop JR, Robacker KK. 1988.** Nutrient salts promote light-induced degradation of indole-3-acetic acid in tissue culture media. *Plant Physiology* **88**: 379–382.
- Elstner EF. 1982.** Oxygen activation and oxygen toxicity. *Annual Review of Plant Physiology* **33**: 73–96.
- Elstner EF, Osswald W. 1994.** Mechanism of oxygen activation during plant stress. *Proceedings of the Royal Society of Edinburgh Section B* **102B**: 131–154.
- Elstner EF, Wagner GA, Schutz W. 1988.** Activated oxygen in green plants in relation to stress conditions. *Current Topics in Plant Biochemistry and Physiology* **7**: 159–187.
- Ekiz H, Bagci SA, Kiral AS, Eker S, Gültekin I, Alkan A, Cakmak I. 1998.** Effects of zinc fertilization and irrigation on grain yield and zinc concentration of various cereals grown in zinc-deficient calcareous soils. *Journal of Plant Nutrition* **21**: 2245–2256.
- Elzenga JTM, Staal M, Prins HBA. 1989.** ATPase activity of isolated plasma membrane vesicles of leaves of *Elodea* as affected by thiol reagents and NADH/NAD⁺ ratio. *Physiologia Plantarum* **76**: 379–385.

- Fischer ES, Thimm O, Rengel Z. 1997.** Zinc nutrition influences the CO₂ gas exchange in wheat. *Photosynthetica* **33**: 505–508.
- Fodor J, Gullner G, Adam AL, Barna B, Komives T, Kiraly Z. 1997.** Local and systemic responses of antioxidants to tobacco mosaic virus infection and salicylic acid in tobacco. *Plant Physiology* **114**: 1443–1451.
- Forno DA, Yoshida S, Asher CJ. 1975.** Zinc deficiency in rice. I. Soil factors associated with the deficiency. *Plant and Soil* **42**: 537–550.
- Foyer CH, Lelandais M, Kunert KJ. 1994.** Photooxidative stress in plants. *Physiologia Plantarum* **92**: 696–717.
- Fridovich I. 1986.** Biological effects of the superoxide radical. *Archives of Biochemistry and Biophysics* **247**: 1–11.
- Fryer MJ, Andrews JR, Oxborough K, Blowers DA, Baker NR. 1998.** Relationship between CO₂ assimilation, photosynthetic electron transport, and active O₂ metabolism in leaves of maize in the field during periods of low temperature. *Plant Physiology* **116**: 571–580.
- Garcia JE, Gomez M, Janes J, Lopez-George J, Del Rio LA. 1981.** Isozyme pattern of the metalloenzyme system superoxide dismutase during growth of peas (*Pisum sativum* L.) under different iron nutrient concentrations. *Zeitschrift für Pflanzenphysiologie* **105**: 21–29.
- Gazaryan IG, Lagrimini LM, Ashby GA, Thorneley RNF. 1996.** Mechanism of indole-3-acetic acid oxidation by plant peroxidases: anaerobic stopped-flow spectrophotometric studies on horseradish and tobacco peroxidases. *Biochemical Journal* **313**: 841–847.
- Girotti AW. 1985.** Mechanisms of lipid peroxidation. *Free Radical Biology and Medicine* **1**: 87–95.
- Girotti AW, Thomas JP, Jordan JE. 1985.** Inhibitory effect of zinc(II) on free radical lipid peroxidation in erythrocyte membranes. *Free Radical Biology and Medicine* **1**: 395–401.
- Glinn M, Ernster L, Lee CP. 1991.** Initiation of lipid peroxidation in submitochondrial particles: effects of respiratory inhibitors. *Archives of Biochemistry and Biophysics* **290**: 57–65.
- Graham RD, Ascher JS, Hynes SC. 1992.** Selecting zinc-efficient cereal genotypes for soils of low zinc status. *Plant and Soil* **146**: 241–250.
- Graham RD, Welch RM, Grunes DL, Carey EE, Norvell WA. 1987.** Effect of zinc deficiency on the accumulation of boron and other mineral nutrients in barley. *Soil Science Society of America Journal* **51**: 652–657.
- Greenberg JT, Guo A, Klessig DF, Ausubel FM. 1994.** Programmed cell death in plants: a pathogen-triggered response activated coordinately with multiple defense functions. *Cell* **77**: 551–563.
- Gressel J, Galun E. 1994.** Genetic controls of photooxidant tolerance. In: Foyer CH, Mullineaux P, eds. *Causes of photooxidative stress and amelioration of defense systems in plants*. Boca Raton, FL, USA: CRC Press, 237–273.
- Gross GG, Janse C, Elstner EF. 1977.** Involvement of malate, monophenols, and the superoxide radical in hydrogen peroxide formation by isolated cell walls from horseradish (*Armoracia lapathifolia* Gilib). *Planta* **136**: 271–276.
- Halliwell B, Gutteridge JMC. 1984.** Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochemical Journal* **219**: 1–14.
- Hammermüller JD, Bray TM, Bettger WJ. 1987.** Effect of zinc and copper deficiency on microsomal NADPH-dependent active oxygen generation in rat lung and liver. *Journal of Nutrition* **117**: 894–901.
- Hedrich R, Kurkdjian A. 1988.** Characterization of an anion-permeable channel from sugar beet vacuoles: effect of inhibitors. *EMBO Journal* **7**: 3661–3666.
- Heiser I, Oswald W, Elstner EF. 1998.** The formation of reactive oxygen species by fungal and bacterial phytotoxins. *Plant Physiology and Biochemistry* **36**: 703–713.
- Hendry GAF. 1993.** Oxygen, free radical process and seed longevity. *Seed Science Research* **3**: 141–153.
- Hendry GAF, Brocklebank KJ. 1985.** Iron-induced oxygen radicals metabolism in waterlogged plants. *New Phytologist* **101**: 199–206.
- Hernandez JA, Olmos E, Corpas FJ, Sevilla F, Del Rio LA. 1995.** Salt-induced oxidative stress in chloroplast of pea plants. *Plant Science* **105**: 151–167.
- Hideg E, Spetea C, Vass I. 1994.** Singlet oxygen and free radical production during acceptor- and donor-side-induced photo-inhibition. Studies with spin trapping EPR spectroscopy. *Biochimica et Biophysica Acta* **1186**: 143–152.
- Hodgson EK, Fridovich I. 1975.** The interaction of bovine erythrocyte superoxide dismutase with hydrogen peroxide. Inactivation of enzyme. *Biochemistry* **14**: 5249–5299.
- Hossain B, Hirata N, Nagatomo Y, Akashi R, Takagi H. 1997.** Internal zinc accumulation is correlated with increased growth in rice suspension culture. *Journal of Plant Growth Regulation* **16**: 239–243.
- Huner NPA, Oquist G, Sarhan F. 1998.** Energy balance and acclimation to light and cold. *Trends in Plant Science* **3**: 224–230.
- Iturbe-Ormaetxe I, Moran JF, Arrese-Igor C, Gogorcena Y, Klucas RV, Becana M. 1995.** Activated oxygen and antioxidant defenses in iron-deficient pea plants. *Plant, Cell & Environment* **18**: 421–429.
- Jabs T, Dietrich A, Dangl JL. 1996.** Initiation of runaway cell death in an *Arabidopsis* mutant by extracellular superoxide. *Science* **273**: 1853–1856.
- Jackson C, Dench J, Moore AL, Halliwell B, Foyer C, Hall DO. 1978.** Subcellular localization and identification of superoxide dismutase in leaves of higher plants. *European Journal of Biochemistry* **91**: 339–344.
- Jackson TL, Hay J, Moore DP. 1967.** The effects of zinc on yield and chemical composition of sweet corn in Willamette Valley. *American Society of Horticultural Sciences* **91**: 462–471.
- Jeffery EH. 1983.** The effect of zinc on NADPH oxidation and monooxygenase activity in rat hepatic microsomes. *Molecular Pharmacology* **23**: 467–473.
- Kastrup BV, Steiger S, Lüttge U, Fischer-Schliebs E. 1996.** Regulatory effects of zinc on corn root plasma membrane H⁺-ATPase. *New Phytologist* **134**: 61–73.
- Kazuba M, Hunt GRA. 1990.** Protection against membrane damage: a ¹H-NMR investigation of the effect of Zn²⁺ and Ca²⁺ on the permeability of phospholipid vesicles. *Journal of Inorganic Biochemistry* **40**: 217–255.
- Kennedy CD, Gonsalves FAN. 1987.** The action of divalent zinc, cadmium, mercury, copper and lead on the trans-root potential and H⁺-efflux of excised roots. *Journal of Experimental Botany* **38**: 800–817.
- Kitagishi K, Obata H. 1986.** Effects of zinc deficiency on the nitrogen metabolism of meristematic tissues of rice plants with reference to protein synthesis. *Soil Science and Plant Nutrition* **32**: 397–405.
- Kitagishi K, Obata H, Kondo T. 1987.** Effect of zinc deficiency on 80S ribosome content of meristematic tissues of rice plant. *Soil Science and Plant Nutrition* **33**: 423–430.
- Klug A, Rhodes D. 1987.** ‘Zinc fingers’: a novel protein motif for nucleic acid recognition. *Trends in Biochemical Sciences* **12**: 464–469.
- Kochian LV. 1993.** Zinc absorption from hydroponic solutions by plant roots. In: Robson AD, ed. *Zinc in soils and plants*. Dordrecht, The Netherlands: Kluwer Academic, 45–57.
- Kovac L, Bohmerova E, Necas O. 1987.** The plasma membrane of yeast protoplasts exposed to hypotonicity becomes porous but does not disintegrate in the presence of protons or polyvalent cations. *Biochimica et Biophysica Acta* **899**: 265–275.
- Krapp A, Quick WP, Stitt M. 1991.** Ribulose-1,5-bisphosphate carboxylase-oxygenase, other Calvin-cycle enzymes, and chlorophyll decrease when glucose is supplied to mature spinach leaves via the transpiration stream. *Planta* **186**: 58–69.
- Kunimoto M, Inoue K, Nojima S. 1981.** Effect of ferrous ion and ascorbate-induced lipid peroxidation on liposomal membranes. *Biochimica et Biophysica Acta* **646**: 169–178.
- Kurella EG, Osipov A, Goldman R, Boldyrev AA, Kagan VE. 1995.** Inhibition of Na⁺/K⁺-ATPase by phenoxyl radicals of etoposide (VP-16): role of sulfhydryl oxidation. *Biochimica et Biophysica Acta* **1232**: 52–58.
- Lawrence K, Bhalla P, Prakash CM. 1995.** Changes in NAD(P)H-dependent redox activities in plasmalemma-enriched vesicles isolated from boron- and zinc-deficient chick pea roots. *Journal of Plant Physiology* **146**: 652–657.
- Levine A, Pennell R, Palmer R, Lamb CJ. 1996.** Calcium-mediated apoptosis in plant hypersensitive response. *Current Biology* **6**: 427–437.
- Lindsay CB, Rodrigues L, Pasternak CA. 1989.** Protection of

- cells against membrane damage by haemolytic agents: divalent cations and protons act at the extracellular side of the plasma membrane. *Biochimica et Biophysica Acta* **983**: 56–64.
- Loneragan JF, Grunes DL, Welch RM, Aduayi E, Tengah A, Lazar VA, Cary EE. 1982.** Phosphorus accumulation and toxicity in leaves in relation to zinc supply. *Soil Science Society of America Journal* **46**: 345–352.
- Long SP, Humphries S, Falkowski PG. 1994.** Photoinhibition of photosynthesis in nature. *Annual Review of Plant Physiology and Plant Molecular Biology* **45**: 633–662.
- Lopez-Huertas E, Corpas FJ, Sandalio LM, Del Rio LA. 1999.** Characterization of membrane polypeptides from pea leaf peroxisomes involved in superoxide radical generation. *Biochemical Journal* **337**: 531–536.
- Lovelock C, Winter K. 1996.** Oxygen-dependent electron transport and protection from photoinhibition in leaves of tropical tree species. *Planta* **198**: 580–587.
- Madamanchi NR, Donahue JI, Cramer CL, Alscher RG, Pedersen K. 1994.** Differential response of Cu, Zn superoxide dismutase in two pea cultivars during a short-term exposure to sulfur dioxide. *Plant Molecular Biology* **26**: 95–103.
- Marschner H. 1995.** *Mineral nutrition of higher plants*, 2nd edn. London, UK: Academic Press.
- Marschner H, Cakmak I. 1986.** Mechanism of phosphorus induced zinc deficiency in cotton. II. Evidence for impaired shoot control of phosphorus uptake and translocation under zinc deficiency. *Physiologia Plantarum* **68**: 491–496.
- Marschner H, Cakmak I. 1989.** High light intensity enhances chlorosis and necrosis in leaves of zinc-, potassium- and magnesium-deficient bean (*Phaseolus vulgaris*) plants. *Journal of Plant Physiology* **134**: 308–315.
- Marschner H, Kirkby EA, Cakmak I. 1996.** Effect of mineral nutritional status on shoot–root partitioning of photo assimilates and cycling of mineral nutrients. *Journal of Experimental Botany* **47**: 1255–1263.
- McRae DG, Baker JE, Thompson JE. 1982.** Evidence for involvement of the superoxide radical in the conversion of 1-aminocyclopropane-1-carboxylic acid to ethylene by pea microsomal membranes. *Plant and Cell Physiology* **23**: 375–383.
- Mehdy MC, Sharma YK, Sathasivan K, Bays NW. 1996.** The role of activated oxygen species in plant disease resistance. *Physiologia Plantarum* **98**: 365–374.
- Metodiewa D, De Melo MP, Escobar JA, Cilento G, Dunford HB. 1992.** Horseradish peroxidase-catalysed aerobic oxidation and peroxidation of indole-3-acetic acid. I. Optical spectra. *Archives of Biochemistry and Biophysics* **296**: 27–33.
- Michalski WP, Kaniuga Z. 1981.** Photosynthetic apparatus of chilling-sensitive plants. X. Relationship between superoxide dismutase activity and photoperoxidation of chloroplast lipids. *Biochimica et Biophysica Acta* **637**: 159–167.
- Minotti G. 1990.** NADPH- and adriamycin-dependent microsomal release of iron and lipid peroxidation. *Archives of Biochemistry and Biophysics* **277**: 268–276.
- Mittler R, Shulaev V, Seskar M, Lam M. 1996.** Inhibition of programmed cell death in tobacco plants during a pathogen-induced hypersensitive response at low oxygen pressure. *Plant Cell* **8**: 1991–2001.
- Mittler R, Zilinskas BA. 1994.** Regulation of pea cytosolic ascorbate peroxidase and other antioxidant enzymes during the progression of drought stress and following recovery from drought. *Plant Journal* **5**: 397–405.
- Murphy TM, Auh CK. 1996.** The superoxide synthase of plasma membrane preparations from cultured rose cells. *Plant Physiology* **110**: 621–629.
- Neue HU, Quijano C, Senadhira D, Setter T. 1998.** Strategies for dealing with micronutrient disorders and salinity in lowland rice systems. *Field Crops Research* **56**: 139–155.
- Norvell AW, Welch RM. 1993.** Growth and nutrient uptake by barley (*Hordeum vulgare* L. cv. Herta): studies using an N-(2-hydroxyethyl) ethylenedinitrilotriacetic acid-buffered nutrient solution technique. I. Zinc ion requirements. *Plant Physiology* **101**: 619–625.
- Obata H, Shimoyama A, Umebayashi M. 1997.** Effect of shading of zinc deficiency symptoms in rice plant. *Soil Science and Plant Nutrition* **43**: 933–936.
- Ohki H. 1976.** Effect of zinc nutrition on photosynthesis and carbonic anhydrase activity in cotton. *Physiologia Plantarum* **38**: 300–304.
- Osmond CB, Grace SC. 1995.** Perspective on photoinhibition and photorespiration in the field: quintessential inefficiencies of the light and dark reactions of photosynthesis? *Journal of Experimental Botany* **46**: 1351–1362.
- Osswald WF, Elstner EF. 1986.** Mechanismen der pathologischen Pigmentbleichung. *Berichte der Deutschen Botanischen Gesellschaft* **99**: 341–356.
- Parat M-O, Richard M-J, Pollet S, Hadjur C, Favier A, Beani J-C. 1997.** Zinc and DNA fragmentation in keratinocyte apoptosis: first inhibitory effect in UVB irradiated cells. *Journal of Photochemistry and Photobiology B: Biology* **37**: 101–106.
- Parker DR. 1997.** Responses of six crop species to solution Zn²⁺ activities buffered with HEDTA. *Soil Science Society of America Journal* **61**: 167–176.
- Perl A, Perl-Treves R, Galili S, Aviv D, Shalgi E, Malkin S, Galun E. 1993.** Enhanced oxidative-stress defense in transgenic potato expressing tomato Cu, Zn superoxide dismutase. *Theoretical and Applied Genetics* **85**: 568–576.
- Pinton R, Cakmak I, Marschner H. 1993.** Effects of zinc deficiency on proton fluxes in plasma membrane-enriched vesicles isolated from bean roots. *Journal of Experimental Botany* **44**: 623–630.
- Pinton R, Cakmak I, Marschner H. 1994.** Zinc deficiency enhanced NAD(P)H-dependent superoxide radical production in plasma membrane vesicles isolated from roots of bean plants. *Journal of Experimental Botany* **45**: 45–50.
- Pitcher LH, Zilinskas BA. 1996.** Overexpression of copper/zinc superoxide dismutase in the cytosol of transgenic tobacco confers partial resistance to ozone-induced foliar necrosis. *Plant Physiology* **110**: 583–588.
- Polle A. 1996.** Mehler reaction: friend or foe in photosynthesis? *Botanica Acta* **109**: 84–89.
- Powell SR, Donna H, Aiuto L, Wapnir RA, Teichberg S, Tortolani AJ. 1994.** Zinc improves postischemic recovery of isolated rat hearts through inhibition of oxidative stress. *American Journal of Physiology* **266**: 2497–2507.
- Prasad R, Kaur D, Kumar V. 1996.** Kinetics characterization of zinc binding to brush border membranes from rat kidney cortex: interaction with cadmium. *Biochimica et Biophysica Acta* **1284**: 69–78.
- Price AH, Hendry AF. 1991.** Iron-catalysed oxygen radical formation and its possible contribution to drought damage in nine native grasses and three cereals. *Plant, Cell & Environment* **14**: 477–484.
- Price AH, Hendry GAF. 1989.** Stress and the role of activated oxygen scavengers and protective enzymes in plants subjected to drought. *Biochemical Society Transactions* **17**: 493–494.
- Qian SY, Buettner GR. 1999.** Iron and dioxygen chemistry is an important route to initiation of biological free radical oxidations: an electron paramagnetic resonance spin trapping study. *Free Radical Biology and Medicine* **26**: 1447–1456.
- Rahimi A, Bussler W. 1979.** Die Entwicklung und der Zn-, Fe- und P-Gehalt höherer Pflanzen in Abhängigkeit vom Zinkangebot. *Zeitschrift für Pflanzenernährung und Bodenkunde* **142**: 15–27.
- Rao M, Paliyath G, Ormrod DP. 1996.** Ultraviolet-B and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiology* **110**: 125–136.
- Reichheld J-P, Vernoux T, Lardon F, Van Montagu M, Inze D. 1999.** Specific checkpoints regulate plant cell cycle progression in response to oxidative stress. *Plant Journal* **17**: 647–656.
- Rengel Z. 1995a.** Carbonic anhydrase activity in leaves of wheat genotypes differing in zinc efficiency. *Journal of Plant Physiology* **147**: 251–256.
- Rengel Z. 1995b.** Sulfhydryl groups in root-cell plasma membranes of wheat genotypes differing in zinc efficiency. *Physiologia Plantarum* **95**: 604–612.
- Rengel Z. 1997.** Root exudation and microflora populations in rhizosphere of crop genotypes differing in tolerance to micronutrient deficiency. *Plant and Soil* **196**: 255–260.
- Rengel Z, Graham RD. 1995.** Wheat genotypes differ in zinc efficiency when grown in chelate-buffered nutrient solution. II. Nutrient uptake. *Plant and Soil* **176**: 317–324.
- Rengel Z, Gutteridge R, Hirsch P, Hornby D. 1996.** Plant genotype, micronutrient fertilization and take-all infection influence bacterial populations in the rhizosphere of wheat. *Plant and Soil* **183**: 269–277.

- Rengel Z, Römheld V, Marschner H. 1998a. Uptake of zinc iron by wheat genotypes differing in tolerance to zinc deficiency. *Journal of Plant Physiology* **152**: 433–438.
- Rengel Z, Ross G, Hirsch P. 1998b. Plant genotype and micronutrients status influence colonization of wheat roots by soil bacteria. *Journal of Plant Nutrition* **21**: 99–113.
- Rhodes D, Klug A. 1993. Zinc fingers. *Scientific American* **268**: 32–39.
- Ridgley EL, Xiong Z, Ruben L. 1999. Reactive oxygen species activate a Ca^{2+} -dependent cell death pathway in the unicellular organism *Trypanosoma brucei brucei*. *Biochemical Journal* **340**: 33–40.
- Robinson JM. 1988. Does O_2 photoreduction occur within chloroplasts *in vivo*? *Physiologia Plantarum* **72**: 666–680.
- Römheld V, Marschner H. 1991. Function of micronutrients in plants. In: Mortvedt JJ, Cox FR, Shuman LM, Welch RM, eds. *Micronutrients in agriculture*. SSSA Book Series No. 4. Madison, WI, USA: Soil Science Society of America, 297–328.
- Roy P, Roy SK, Mitra A, Kulkarni AP. 1994. Superoxide generation by lipoxygenase in the presence of NADH and NADPH. *Biochimica et Biophysica Acta* **1214**: 171–179.
- Rygel J, Arnold WM, Zimmermann U. 1992. Zinc and salinity effects on membrane transport in *Chara connivens*. *Plant, Cell & Environment* **15**: 11–23.
- Sahrawat KL, Mulbah CK, Diatta S, Delaune RD, Patrick WH, Singh BN, Jones MP. 1996. The role of tolerant genotypes and plant nutrients in the management of iron toxicity in lowland rice. *Journal of Agricultural Science* **126**: 143–149.
- Sandalio LM, Palma JM, Del Rio LA. 1987. Localization of manganese superoxide dismutase in peroxisomes isolated from *Pisum sativum* L. *Plant Science* **51**: 1–8.
- Santa Maria GE, Cogliatti DH. 1988. Bidirectional Zn-fluxes and compartmentation in wheat seedling roots. *Journal of Plant Physiology* **132**: 312–315.
- Sarafian TA, Bredesen DE. 1994. Is apoptosis mediated by reactive oxygen species? *Free Radicals Research* **21**: 1–8.
- Scandalios JG. 1993. Oxygen stress and superoxide dismutase. *Plant Physiology* **101**: 7–12.
- Schneider EA, Wightman F. 1974. Metabolism of auxin in higher plants. *Annual Review of Plant Physiology* **25**: 487–513.
- Searle AJF, Tomasi A. 1982. Hydroxyl free radical production in iron-cysteine solutions and protection by zinc. *Journal of Inorganic Biochemistry* **17**: 161–166.
- Sen Gupta A, Heinen J, Holaday AS, Burke JJ, Allen RD. 1993a. Increased resistance to oxidative stress in transgenic plants that over-express chloroplastic Cu/Zn superoxide dismutase. *Proceedings of the National Academy of Sciences, USA* **90**: 1629–1633.
- Sen Gupta A, Webb RP, Holaday AS, Allen RD. 1993b. Over-expression of superoxide dismutase protects plants from oxidative stress. *Plant Physiology* **103**: 1067–1073.
- Sharma CP, Sharma PN, Bisht SS, Nautiyal BD. 1982. Zinc deficiency induces changes in cabbage. In: Scaife A, ed. *Proceedings of the Ninth Plant Nutrition Colloquium*, Commonwealth Agricultural Bureau, Farnham Royal, UK. Wallingford, UK: CAB International, 601–606.
- Sharma PN, Kumar N, Bisht SS. 1994. Effect of zinc deficiency on chlorophyll content, photosynthesis and water relations of cauliflower plants. *Photosynthetica* **30**: 353–359.
- Sharma PN, Tripathi A, Bisht SS. 1995. Zinc requirement for stomatal opening in cauliflower. *Plant Physiology* **107**: 751–756.
- Sharma YK, Davis KR. 1997. The effects of ozone on antioxidant responses in plants. *Free Radicals in Biology and Medicine* **23**: 480–488.
- Shukla UC, Prasad KG. 1974. Ameliorative role of zinc on maize growth under alkali soil condition. *Agronomy Journal* **66**: 804–806.
- Sillanpää M. 1990. *Micronutrients assessment at the country level: an international study*. FAO Soils Bulletin 63. Rome, Italy: Food and Agriculture Organization of the United Nations.
- Skoog F. 1940. Relationship between zinc and auxin in the growth of higher plants. *American Journal of Botany* **27**: 939–950.
- Smirnoff N. 1993. The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytologist* **125**: 27–58.
- Spalding BC, Swift JG, Horowicz P. 1986. Zinc inhibition of potassium efflux in depolarized frog muscle and its modification by external hydrogen ions and diethylpyrocarbonate treatment. *Journal of Membrane Biology* **93**: 157–164.
- Spalding BC, Taber P, Swift JG, Horowicz P. 1990. Zinc inhibition of chloride efflux from skeletal muscle of rana pipiens and its modification by external pH and chloride activity. *Journal of Membrane Biology* **116**: 195–214.
- Stoyanovsky DA, Salama G, Kagan VE. 1994. Ascorbate/iron activates Ca^{2+} -release channels of skeletal sarcoplasmic reticulum vesicles reconstituted in lipid bilayers. *Archives of Biochemistry and Biophysics* **308**: 214–221.
- Strack PR, Waxmann L, Fagan JM. 1996. ATP-stimulated degradation of oxidatively modified superoxide dismutase by cathepsin D in cardiac tissue extracts. *Biochemical and Biophysical Research Communications* **219**: 348–353.
- Suzuki H, Pabst MJ, Johnston RB Jr. 1985. Enhancement by Ca^{2+} or Mg^{2+} of catalytic activity of the superoxide-producing NADPH oxidase in membrane fractions in human neutrophils and monocytes. *Journal of Biological Chemistry* **260**: 3635–3639.
- Svingen BA, Beuge JA, O'Neal FO, Aust SD. 1979. The mechanism of NADPH-dependent lipid peroxidation. The propagation of lipid peroxidation. *Journal of Biological Chemistry* **254**: 5892–5899.
- Takkar PN, Chibba IM, Mehta SK. 1989. *Twenty years of coordinated research of micronutrients in soil and plants (1967–1987)*. Bhopal, India: Indian Institute of Soil Science.
- Takkar PN, Walker CD. 1993. The distribution and correction of zinc deficiency. In: Robson AD, ed. *Zinc in soils and plants*. Dordrecht, The Netherlands: Kluwer Academic, 151–166.
- Tang LX, Yang JL, Shen X. 1997. Effects of additional iron-chelators on Fe^{2+} -initiated lipid peroxidation: evidence to support the Fe^{2+} - Fe^{3+} complex as the initiator. *Journal of Inorganic Biochemistry* **68**: 265–272.
- Tester M. 1990. Plant ion channels. Whole-cell and single-channel studies. *New Phytologist* **114**: 305–340.
- Thomas JP, Bachowski GJ, Girotti AW. 1986. Inhibition of cell membrane lipid peroxidation by cadmium- and zinc-metallothioneins. *Biochimica et Biophysica Acta* **884**: 448–461.
- Thongbai P, Goodman BA, Marschner H. 2000. Free radicals as a result of post-anoxic injury of rices (*Oryza sativa*) in an iron-toxic soil. *Journal of Plant Nutrition*. (In press.)
- Toyokuni S. 1996. Iron-induced carcinogenesis: the role of redox regulation. *Free Radical Biology and Medicine* **20**: 553–566.
- Treves S, Trentini PL, Ascanelli M, Bucci G, di Virgilio F. 1994. Apoptosis is dependent on intracellular zinc and independent of intracellular calcium in lymphocytes. *Experimental Cell Research* **211**: 339–343.
- Tyerman SD. 1992. Anion channels in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **43**: 351–373.
- Vallee BL, Auld DS. 1990. Zinc coordination, function, and structure of zinc enzymes and other proteins. *Biochemistry* **29**: 5647–5659.
- Vallee BL, Falchuk KH. 1993. The biochemical basis of zinc physiology. *Physiological Reviews* **73**: 79–118.
- Van Camp W, Capiou K, Van Montagu M, Inzé D, Slooten L. 1996. Enhancement of oxidative stress tolerance transgenic plants overproducing Fe-superoxide dismutase in chloroplasts. *Plant Physiology* **112**: 1703–1714.
- Van Camp W, Van Montagu M, Inzé D. 1994. Superoxide dismutases. In: Foyer CH, Mulineaux PM, eds. *Causes of photooxidative stress and amelioration of defense systems in plants*. Boca Raton, FL, USA: CRC Press, 318–341.
- Van Ginkel G, Sevanian A. 1994. Lipid peroxidation-induced membrane structural alterations. *Methods in Enzymology* **233**: 273–289.
- Vaughan D, Dekock PC, Ord BG. 1982. The nature and localization of superoxide dismutase on fronds of *Lemna gibba* L. and the effect of copper and zinc deficiency on its activity. *Physiologia Plantarum* **54**: 253–257.
- Von Schaewen A, Stitt M, Schmidt R, Sonnewald U, Willmitzer L. 1990. Expression of yeast-derived invertase in the cell wall of tobacco and *Arabidopsis* plants lead to accumulation of carbohydrate and inhibition of photosynthesis and strongly influences growth and phenotype of transgenic tobacco plants. *EMBO Journal* **9**: 3033–3044.
- Walter A, Römheld V, Marschner H. 1994. Is the release of phytosiderophores in zinc-deficient wheat plants response to impaired iron utilization? *Physiologia Plantarum* **92**: 493–500.

- Warnock RE. 1970.** Micronutrient uptake and mobility within corn plants (*Zea mays* L.) in relation to phosphorus-induced zinc deficiency. *Soil Science Society of American Proceedings* **34**: 765–769.
- Welch RM. 1995.** Micronutrient nutrition of plants. *Critical Reviews in Plant Sciences* **14**: 49–82.
- Welch RM, Allaway WH, House WA, Kubota J. 1991.** Geographic distribution of trace element problems. In: Mortvedt JJ, Cox FR, Shuman LM, Welch RM, eds. *Micronutrients in agriculture*. SSSA Book Series No. 4. Madison, USA: Soil Science Society of America, 31–57.
- Welch RM, Norvell WA. 1993.** Growth and nutrient uptake by barley (*Hordeum vulgare* L. cv. Herta): studies using an *N*-(2-hydroxyethyl) ethylenedinitrioltriacetic acid-buffered nutrient solution technique. II. Role of zinc in the uptake and root leakage of mineral nutrients. *Plant Physiology* **101**: 627–631.
- Welch RM, Webb MJ, Loneragan JF. 1982.** Zinc in membrane function and its role in phosphorus toxicity. In: Scaife A, ed. *Proceedings of the Ninth Plant Nutrition Colloquium*. Warwick, UK. Wallingford, UK: CAB International, 710–715.
- Wenzel AA, Mehlhorn H. 1995.** Zinc deficiency enhances ozone toxicity in bush beans (*Phaseolus vulgaris* L. cv. Saxa). *Journal of Experimental Botany* **46**: 867–872.
- White JG; Zasoski RH. 1999.** Mapping soil micronutrients. *Field Crops Research* **60**: 11–26.
- Willekens H, Van Camp W, Van Montagu M, Inzé D, Langebartels C, Sandermann H. 1994.** Ozone, sulphur dioxide, and ultraviolet B have similar effects on mRNA accumulation of antioxidant genes in *Nicotiana plumbaginifolia* L. *Plant Physiology* **106**: 1007–1014.
- Williams RJP. 1988.** An introduction to the biochemistry of zinc. In: *Zinc in human biology*. London, UK: Springer-Verlag, 15–31.
- Willson RL. 1988.** Zinc and iron in free radical pathology and cellular control. In: Mills CF, ed. *Zinc in human biology*. London, UK: Springer-Verlag, 147–172.
- Wise RR, Naylor AW. 1987.** Chilling-enhanced photooxidation. The peroxidative destruction of lipids during chilling injury to photosynthesis and ultrastructure. *Plant Physiology* **83**: 272–277.
- Wojtaszek P. 1997.** Oxidative burst: an early plant response to pathogen infection. *Biochemical Journal* **322**: 861–892.
- Wood LJ, Murray BJ, Okatan Y, Nooden LD. 1986.** The effect of petiole phloem disruption on starch and mineral distribution in senescing soybean leaves. *American Journal of Botany* **73**: 1377–1383.
- Xia J, Browning JD, O'Dell BL. 1999.** Decreased plasma membrane thiol concentration is associated with increased osmotic fragility of erythrocytes in zinc-deficient rats. *Journal of Nutrition* **129**: 814–819.
- Yamaguchi M. 1989.** Rice bronzing in Nigeria caused by nutrient imbalances and its control by potassium sulfate application. *Plant and Soil* **117**: 275–286.
- Yu Q, Osborne L, Rengel Z. 1998.** Micronutrient deficiency changes activities of superoxide dismutase and ascorbate peroxidase in tobacco plants. *Journal of Plant Nutrition* **21**: 1427–1437.
- Yu Q, Rengel Z. 1999.** Micronutrient deficiency influences plant growth and activities of superoxide dismutases in narrow-leaved lupins. *Annals of Botany* **83**: 175–182.
- Yu Q, Worth C, Rengel Z. 1999.** Using capillary electrophoresis to measure Cu/Zn superoxide dismutase concentration in leaves of wheat genotypes differing in tolerance to zinc deficiency. *Plant Science* **143**: 231–239.
- Zalewski PD, Forbes IJ, Betts WH. 1993.** Correlation of apoptosis with change in intracellular labile Zn(II) using zinquin [(2-methyl-8-*p*-toluenesulphonamide-6-quinolyloxy)-acetic acid], a new specific fluorescent probe for Zn(II). *Biochemical Journal* **296**: 403–408.
- Zalewski PD, Forbes IJ, Seemark RF, Borlinghaus R, Betts WH, Lincoln SF, Ward AD. 1994.** Flux of intercellular labile zinc during apoptosis (gene-regulated cell death) revealed by a specific chemical probe, Zinquin. *Current Biology* **1**: 153–161.
- Zhang F, Römheld V, Marschner H. 1991.** Release of zinc mobilizing root exudates in differential plant species as affected by zinc nutritional status. *Journal of Plant Nutrition* **14**: 675–686.