

6-3-2021

Targeted Treatment of Head and Neck (Pre)Cancer: Preclinical Target Identification and Development of Novel Therapeutic Applications

Anne van Harten

Thomas Jefferson University; Vrije Universiteit Amsterdam

Ruud H. Brakenhoff

Vrije Universiteit Amsterdam

Follow this and additional works at: <https://jdc.jefferson.edu/cbfp>



Part of the [Otolaryngology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

van Harten, Anne and Brakenhoff, Ruud H., "Targeted Treatment of Head and Neck (Pre)Cancer: Preclinical Target Identification and Development of Novel Therapeutic Applications" (2021).

Department of Cancer Biology Faculty Papers. Paper 181.

<https://jdc.jefferson.edu/cbfp/181>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Cancer Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Review

Targeted Treatment of Head and Neck (Pre)Cancer: Preclinical Target Identification and Development of Novel Therapeutic Applications

Anne M. van Harten ^{1,2} and Ruud H. Brakenhoff ^{1,*}

¹ Cancer Center Amsterdam, Otolaryngology-Head and Neck Surgery, Tumor Biology & Immunology Section, Vrije Universiteit Amsterdam, Amsterdam UMC, 1081 HV Amsterdam, The Netherlands; am.vanharten@amsterdamumc.nl or anne.vanharten@jefferson.edu

² Sidney Kimmel Cancer Center, Department of Cancer Biology, Thomas Jefferson University, Philadelphia, PA 19107, USA

* Correspondence: rh.brakenhoff@amsterdamumc.nl; Tel.: +31-(0)20-4440953

Simple Summary: Head and neck squamous cell carcinomas (HNSCCs) develop from mucosal cells in the oral cavity, pharynx and larynx after either prolonged exposure to tobacco and alcohol, or a transforming infection with high-risk human papillomavirus (HPV). HPV-negative HNSCCs develop in a zone of premalignant mucosal cells centimeters in diameter and characterized by tumor-associated genetic changes, also referred to as ‘fields’, which can present as white leukoplakia lesions but are mostly invisible. Patients with HPV-negative HNSCC have an overall 5-years survival rate of 50–60%, despite application of intense treatment protocols, and current treatment regimens seem to have reached their plateau. Recently, immunotherapy using immune checkpoint inhibitors has been introduced, but seems effective in only some patients. Targeted treatments have failed to find their way into the clinic while novel therapies are urgently awaited that could target the tumor as well as the precancerous cells. However, recent data suggest that we are at the dawn of a new era. This review focusses on the preclinical identification of druggable targets for therapy for HPV-negative HNSCC and their preceding precancerous changes.

Abstract: Head and neck squamous cell carcinomas (HNSCC) develop in the mucosal lining of the upper-aerodigestive tract. In carcinogen-induced HNSCC, tumors emerge from premalignant mucosal changes characterized by tumor-associated genetic alterations, also coined as ‘fields’ that are occasionally visible as leukoplakia or erythroplakia lesions but are mostly invisible. Consequently, HNSCC is generally diagnosed *de novo* at more advanced stages in about 70% of new diagnosis. Despite intense multimodality treatment protocols, the overall 5-years survival rate is 50–60% for patients with advanced stage of disease and seems to have reached a plateau. Of notable concern is the lack of further improvement in prognosis despite advances in treatment. This can be attributed to the late clinical presentation, failure of advanced HNSCC to respond to treatment, the deficit of effective targeted therapies to eradicate tumors and precancerous changes, and the lack of suitable markers for screening and personalized therapy. The molecular landscape of head and neck cancer has been elucidated in great detail, but the absence of oncogenic mutations hampers the identification of druggable targets for therapy to improve outcome of HNSCC. Currently, functional genomic approaches are being explored to identify potential therapeutic targets. Identification and validation of essential genes for both HNSCC and oral premalignancies, accompanied with biomarkers for therapy response, are being investigated. Attentive diagnosis and targeted therapy of the preceding oral premalignant (preHNSCC) changes may prevent the development of tumors. As classic oncogene addiction through activating mutations is not a realistic concept for treatment of HNSCC, synthetic lethality and collateral lethality need to be exploited, next to immune therapies. In recent studies it was shown that cell cycle regulation and DNA damage response pathways become significantly altered in HNSCC causing replication stress, which is an avenue that deserves further exploitation as an HNSCC vulnerability for treatment. The focus of this review is to summarize the current literature on the preclinical identification of potential druggable targets for therapy of (pre)HNSCC, emerging



Citation: van Harten, A.M.; Brakenhoff, R.H. Targeted Treatment of Head and Neck (Pre)Cancer: Preclinical Target Identification and Development of Novel Therapeutic Applications. *Cancers* **2021**, *13*, 2774. <https://doi.org/10.3390/cancers13112774>

Academic Editors: Robert Mandic and Boris A. Stuck

Received: 3 May 2021

Accepted: 28 May 2021

Published: 3 June 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

from the variety of gene knockdown and knockout strategies, and the testing of targeted inhibitors. We will conclude with a future perspective on targeted therapy of HNSCC and premalignant changes.

Keywords: HNSCC; targeted treatment; preclinical development

1. Introduction

Head and neck squamous cell carcinoma (HNSCC) ranks in the top ten of all cancers with respect to incidence, and encompasses the tumors that develop in the mucosal lining of the oral cavity, oropharynx, hypopharynx, and larynx. Annually 700,000 new cases are diagnosed worldwide. The incidence is gender biased; HNSCC occurs about two-times more often in males than in females [1]. This bias may be related to the major classical risk factors for HNSCC; carcinogen exposure from tobacco smoke and (excessive) alcohol consumption, with concomitant exposure increasing the risk [2–4]. A second risk factor for HNSCC is infection with a high-risk type human papillomavirus (hrHPV), particularly for tumors arising in the oropharynx [2–8]. Finally, patients with certain hereditary syndromes may predispose to HNSCC development, of which the most prominent example is the genetic instability syndrome Fanconi anemia (FA) [9–13].

The presence or absence of hrHPV analyzed by surrogate marker p16^{Ink4A}-immunostaining, often followed by HPV DNA PCR, is currently used to classify tumors in the oropharynx in HPV-positive and HPV-negative disease [3,4]. These subgroups differ in progression-free and overall survival as well as in clinical and molecular characteristics [4,14–16]. The most recent data from a Dutch cohort of 1204 oropharyngeal squamous cell carcinoma (OPSCC) patients revealed a 5-years overall survival of 79% for the HPV-positive group and of 43% of the HPV-negative group [17]. Consequently, HPV-positive and HPV-negative OPSCC are considered as separate disease entities. The difference in prognosis has led furthermore to treatment de-escalation trials for HPV-positive HNSCC, although this has not yet resulted in a more personalized treatment based on HPV status [18,19].

As mentioned above, the major risk factors for HPV-negative HNSCC are tobacco and alcohol exposure [2,4,20,21]. Cancer risk relates to exposure, generally summarized as pack years and unit years, and particularly the combination of smoking with alcohol exposure increases risk synergistically [22]. However, HPV-negative tumors may also occur occasionally in patients who hardly smoked or consumed alcohol, and the reason for cancer development in these cases is still elusive [2,23]. Patients generally present with tumors *de novo*, but molecular research during the last decades revealed that HPV-negative HNSCC develops in premalignant changes in the mucosal lining of the head and neck region [4,24–29], which, however, are mostly not visible by eye.

The premalignant cells cover large areas of the mucosal lining with dimensions of multiple centimeters in diameter, and contain some of the tumor-associated genetic changes (mutations and copy number aberrations) that are also observed in the majority of HNSCC [4,29–31]. These cells form a contiguous premalignant “field” that is mostly invisible to the naked eye but may be microscopically identified as dysplasia in surgical margins of excised tumor specimen. Fields can also be identified per definition using genetic markers on DNA obtained from surgical margins or cells obtained through oral brushing [29–31]. Only a smaller subgroup of premalignant fields are macroscopically visible as leukoplakia (white lesions) or erythroplakia (red lesions), which occur with a prevalence of 0.1–0.2% or 0.01–0.02%, respectively [4,32–34]. Sometimes multiple independent premalignant fields are found in a patient, harboring different gene mutations or genomic aberrations [4,24,26]. The fields can eventually evolve into a squamous cell carcinoma. As most of these fields are centimeters in diameter but not visible to the naked eye, fields may remain undetected and stay behind when tumors are excised. When premalignant fields stay behind, new tumors could form in these fields, which are clinically diagnosed as local recurrences when they occur within a distance of 2 cm from the original tumor and within a timespan

of less than 3 years, or as a second primary tumor when located more than 2 cm from the original tumor or after 3 years [4,35]. Ideally, to prevent tumors developing in these fields, (targeted) treatments are needed to eradicate these cells before they transform, often again, into a malignancy. The treatment of premalignant fields is challenging as most are not macroscopically visible and their dimensions are covering large mucosal areas. Smaller visible abnormalities can be resected or treated with laser therapy, but often recur after treatment [33]. Visible or invisible, the curative treatment of premalignant fields remains problematic.

2. Current Treatment Protocols of HNSCC

2.1. Surgery and (Chemo-)Radiotherapy

During the last two decades, the therapeutic arsenal to treat cancer has been changing rapidly, and treatment of a variety of malignancies is increasingly based on tumor genetics, thereby aiming to employ personalized strategies with targeted agents, to reduce toxicity and enhance therapeutic efficacy [36,37]. Despite great efforts to uncover new targeted treatments that could find their way into the clinic, the mainstays of HNSCC treatment remain surgery and radiotherapy, the latter with or without concomitant cisplatin-based chemotherapy. Treatment planning is currently still based on site, tumor stage, imaging and post-operative histological findings, but not on genetics or hrHPV presence.

Albeit treatment protocols are generally intense and may cause disfigurements and toxicities in patients, responses remain somewhat disappointing [38]. Of all tumors, approximately 30% are diagnosed at an early stage [2,39], and these are usually treated with single modality treatment that comprise either surgical resection or radiotherapy, depending on the tumor site [40]. Complete cure is often obtained after treatment, and the 5-years survival rate is around 90% [41].

However, the majority of patients (70%) present with advanced stages of disease, with regional lymph node metastasis or even metastases at distant sites [2,39]. These more advanced tumors are treated either with concomitant chemoradiotherapy and when required with surgical salvage, or with upfront surgery combined with post-operative (chemo)radiotherapy. In some centers neoadjuvant (chemo)radiotherapy followed by surgery may be applied [40,42]. Cisplatin has been the primary choice of chemotherapy since 1977 [43], and is combined with concomitant locoregional radiotherapy. For recurrent-metastatic disease and patients unfit for platinum-based therapy, immunotherapy with anti-PD-(L)1 antibodies, anti-EGFR targeting antibody cetuximab, or invasive multidrug chemotherapies are being applied (Figure 1).

Cisplatin is the mostly applied cytotoxic drug in chemotherapy regimen, which is often applied in combination with radiotherapy. The drug is able to form both intra-strand and inter-strand crosslinking bridges between the two complementary DNA strands of both the genomic and mitochondrial DNA [44,45]. This covalent inter-strand crosslink hampers DNA replication as the replication fork is challenged to pass the DNA crosslink. The key pathway to resolve such DNA crosslinks is the FA/BRCA-pathway (Figure 1) [11,46,47]. Cisplatin acts as radiation-sensitizer, but overall responses differ between tumors. A biomarker or biological explanation for response to cisplatin is unknown other than a defective FA/BRCA pathway [48]. Many HNSCC patients suffer from cisplatin-induced toxicity, and consequently they are frequently hospitalized and can often not sustain the full treatment protocol. Lastly, tumors are or may become resistant to cisplatin.

Most HNSCC patients receive irradiation with photons (IR) in approximately two Gray fractions and up to a total dose of 70 Gray [21,49–51]. Radiation (RT) induces the formation of DNA peroxides by water radiolysis in the presence of oxygen, generating reactive oxygen species (ROS) which induce a high number of single strand DNA (ssDNA) breaks (Figure 1) [52,53]. These breaks lead to the stalling of replication forks and G2/M-checkpoint activation. DNA damage is induced by both the stalled replication forks at ssDNA breaks and the formation of double stranded DNA (dsDNA) breaks during replication when ssDNA breaks that have not been repaired turn into dsDNA breaks [54,55].

Radiation-induced dsDNA damage is commonly repaired by non-homologous end joining (NHEJ) and microhomology-mediated end joining (MMEJ) [56–60]. NHEJ and MMEJ are error-prone DNA repair mechanisms, introducing mutations that are potentially lethal. Nonetheless, it has also been described that MMEJ may contribute to IR-resistance [58].




Figure 1. Approved and clinically applied treatment interventions for HNSCC. Early stage HNSCC is treated with either surgery or radiotherapy with photons or protons, while advanced stage HNSCC is treated by either upfront cisplatin-based chemoradiotherapy and salvage surgery when required, or upfront surgical resection with post-operative (chemo-)radiation. For patients unfit to receive cisplatin, the anti-EGFR monoclonal antibody cetuximab can be applied. Furthermore, patients with recurrent or metastatic disease are treated by the EXTREME protocol and with anti-PD-(L)1 antibodies, such as nivolumab and pembrolizumab, although response rates are low and clinical biomarkers for response are still under investigation. Abbreviations: ROS: reactive oxygen species, ICL: inter-strand crosslink, ssDNA break: single strand DNA break, dsDNA break: double strand DNA break, NK cell: natural killer cell, DC: dendritic cell, T cell: T lymphocyte, PD-1: programmed cell death protein 1, TME: tumor microenvironment, EGFR: epithelial growth factor receptor, PD-L1: programmed death-ligand 1, MAPK: mitogen-activated protein kinase.

2.2. Bio-Radiotherapy

In 2007, the FDA approved chimeric monoclonal antibody cetuximab (Erbix) for treatment of HNSCC (Figure 1) [61,62]. Cetuximab targets the membrane protein epithelial growth factor receptor (EGFR). Although the exact mechanism of action is not completely understood in head and neck cancer, it hinders the binding of EGF to the receptor and thereby inhibits downstream signaling pathways such as the Ras-MAPK-ERK pathway [61]. EGFR overexpression is frequently found in HNSCC, which is in some cases associated with amplification of chromosome 7p [4]. However, single agent response rates remain low in clinical trials (13%) and biomarkers that predict therapy outcome with cetuximab in HNSCC are unidentified [62,63]. It has also been suggested that cetuximab mainly acts as an activator of the immune system in HNSCC patients, thereby acting as a bridge between tumor cells expressing EGFR and immune cells such as CD16-positive NK-cells and dendritic cells (reviewed in [64,65]). The lack of biomarkers predicting response, together with the observation that EGFR small molecule inhibitors such as gefitinib are not particularly effective in the treatment of HNSCC, may point towards an immunological response rather than a molecular response to cetuximab treatment [66,67].

2.3. Immune Checkpoint Inhibitors

Evading the immune response is a commonly accepted hallmark of cancer [68], and HNSCC is known to be very immune suppressive [69]. By overexpression or downregulation of certain immune-associated membrane receptors, tumor cells become unrecognizable to the immune system [65,69–71]. One of these immune-suppressive systems is catalyzed by the interaction between PD-1 on the lymphocytes in the tumor microenvironment (TME) and its ligand PD-L1 on the tumor cells. The development of monoclonal antibodies, such

as nivolumab (Opdivo®) and pembrolizumab (Keytruda®) directed against PD-1, have been shown to be an effective immunotherapeutic strategy in HNSCC [69] (Figure 1). These immune checkpoint inhibitors have been FDA approved for HNSCC since 2016 [72] for recurrent and metastatic disease, and will become more prominent in upfront treatment protocols. Tumor-specific PD-L1 expression is hypothesized to be a predictive biomarker for response, but results are inconclusive [73,74]. It is promising that the response rate in HNSCC to PD-1 antibody treatment is around 25% in both HPV-positive and HPV-negative HNSCC [75]. A recent study with a murine oral squamous cell carcinoma model using 4NQO exposure indicated that PD-1 antibody treatment inhibited progression of the premalignant lesions into carcinoma, which was to a lesser extent also observed in PD-L1 knockout mice treated with 4NQO [76]. This study suggests that PD-(L)1 antibody treatment could be effective in a preventive setting, although data from clinical trials will be important to test this hypothesis. It should be noted that antibody infusion is a rather invasive procedure for treating premalignant changes, as is the toxicity profile of these immune checkpoint inhibitors [77].

3. Molecular Landscape of HNSCC

3.1. Copy Number Alterations

The majority of HNSCC tumors are characterized by a high level of genomic instability, in part resulting from frequent inactivation of cell cycle control [78], which is reflected by many copy number alterations. In the vast majority of HNSCC, gains of chromosomal arms 3q, 5p and 7p, and losses of 3p, 4p and 18q are observed, irrespective of HPV status [3,79–84]. Focal amplifications of 3q26/28 are associated with over-expression of *TP63*, *SOX2* and *PIK3CA* [3,82,85] and lymph node metastasis is associated with the loss of 4p in oral cancers [86]. HPV-negative HNSCC often contains copy number (CN) gains of chromosomes 8q, 9q and 11p, and CN losses of 7q, 8p, 9p, 11q and 18p [3,79–81]. Some of the CN alterations in HPV-negative tumors are already present in premalignant cells [3,4,26,87]. Cell cultures obtained from premalignant fields, including leukoplakia and erythroplakia lesions, contain CN losses in 3p, 8p and 9p21, and amplifications of 3q and 8q, which are also often observed in HNSCC [3,26,87,88].

3.2. Cancer Driver Genes in HNSCC

Comprehensive genomic profiling of HNSCC by the TCGA consortium emphasized the large number of tumor suppressor genes that are inactivated by mutations or chromosomal aberrations, and these studies also highlighted the tremendous heterogeneity of HNSCC [3,38,89,90]. Driving oncogene mutations are largely underrepresented in HNSCC [3], hampering the development of targeted treatments that exploit the concept of ‘oncogene addiction’.

Loss of function of tumor suppressor genes *TP53* and *CDKN2A* (p16^{Ink4A}) through gene mutation, methylation, or in the case of *CDKN2A* through focal loss of 9p21, are often found in both premalignant cells and HPV-negative HNSCC, and occur early in oncogenesis [3,26,29]. HPV-positive tumors typically lack mutations in *TP53* and *CDKN2A*, as the viral oncogenes E6 and E7 block the same signaling pathways. The fact that HPV-positive tumors have a favorable prognosis and are typically *TP53* wild type, interferes with the analysis of *TP53* mutations in relation to clinical outcome. Overall survival of HNSCC patients correlates poorly with *TP53* status particularly when stratified for HPV, conflicting with earlier assumptions when HPV status was not considered [3,78,91,92]. Both *CDKN2A* and *TP53*, through its downstream target p21^{Cip1}, are important inhibitors of cyclin-CDK complexes and can cause cell cycle arrest. Loss of function of these genes thus results in diminished G1/S-checkpoint control and deregulated cellular proliferation, which is considered an important hallmark of cancer [68,93]. Besides *TP53* and *CDKN2A*, gene mutations in *FAT1*, *CASP8*, *AJUBA*, *PIK3CA*, *NOTCH1*, *KMT2D*, *NSD1*, *TGFBR2* and *HRAS* are observed in HPV-negative HNSCC [3]. Of note, alterations of *NOTCH1* in cancer can both be oncogenic or imply loss of function depending on the context, but *NOTCH1*

mutations in HNSCC are generally considered as loss of function mutations and therefore *NOTCH1* acts as tumor suppressor [94]. *PIK3CA* mutations are found in both HPV-positive and -negative tumors, but mutational profiles differ. Amino acid substitutions E542K and E545K in the helical domain of *PIK3CA* are predominantly found in HPV-positive HNSCC, while *PIK3CA* mutations in HPV-negative HNSCC are generally found in the kinase domain [85,95,96].

Of the most frequently mutated genes in HNSCC, only those with oncogenic mutations in *PIK3CA* and *HRAS* can be targeted with small molecule inhibitors that are currently in preclinical testing for HNSCC and other malignancies (see Section 4). Unfortunately, only a minority of HPV-negative HNSCC harbor either a *PIK3CA* or *HRAS* mutation, and therefore only a small patient group might benefit from a therapy regimen targeting these mutations [3].

It stands out that most of the frequently mutated genes in HNSCC such as *TP53*, *CDKN2A*, *CCND1*, *HRAS*, *PIK3CA*, *PTEN* and *RB1* are responsible for increased cell proliferation and deregulation of cell cycle control [3]. In addition, mutations in *CASP8* as well as in *TP53*, *HRAS* and *PIK3CA* contribute to evasion of apoptotic cell death, all typical hallmarks of cancer [68]. Since HNSCC is mainly driven by mutations in tumor suppressor genes, these alterations in cell cycle progression and escape of cell death may serve as potential vulnerabilities to the efficient eradication of the tumor cells, as well as premalignant cells, with appropriately selected targeted agents.

4. Classic Approaches to Identifying Targets for Therapy

We define ‘targeted therapy’ as the therapeutic exploitation of a specific cellular vulnerability in the context of the somatic genetic changes in the tumor cells. Generally, three concepts are considered for the application of targeted therapies for different malignancies: oncogene addiction, synthetic lethality, and collateral lethality (Figure 1). We will discuss these principles and their feasibility in the preclinical development of targeted therapies for HNSCC.

4.1. Oncogene Addiction

Activating mutations of oncogenes may lead to the dependency of the cancer cell on this hyper-activation of the subsequent gene or the pathway it acts in [97]. Oncogene addiction forms an excellent vulnerability that can be targeted by specific inhibitors, and will lead to cancer-specific cell death, while untransformed cells may not be affected as they may rely on backup pathways (Figure 2, left panel) [98]. Targeting the *BRAF*^{V600E} mutation in melanoma using tyrosine kinase inhibitor vemurafenib dramatically changed the clinical treatment of melanoma [99]. Despite initial spectacular results, it also became apparent that resistance is often observed. Moreover, the *BRAF*^{V600E} and vemurafenib story has made crystal clear how critical it is that the pathways are understood in the smallest detail to overcome potential problems such as, in this case, the *BRAF*-paradox; inhibition of B-Raf with vemurafenib caused skin tumors in normal skin by inhibitor-induced refitting of the Raf pathway [100]. Despite these considerations, oncogene addition remains a promising concept in cancer therapy.

Unfortunately, the carcinogenesis of HNSCC is mainly driven by loss of tumor suppressor function [3], and therefore exploitation of oncogene addiction as a vulnerability concept is limited, but nonetheless explored. Oncogene *HRAS* is mutated (*HRAS*^{mut}) in about 4% of HNSCC [3]. Two phase II clinical trials (NCT03719690, NCT02383927) are currently being conducted to investigate the efficacy of tipifarnib, a selective inhibitor of farnesyltransferase that was shown to be an effective strategy for *HRAS*^{mut} HNSCC in a preclinical study [101]. A second oncogene, *PIK3CA*, is mutated in 20% of HNSCC, but only a minority of mutations are activating mutations [3,102]. *PIK3CA* acts in the PI3K-AKT-mTOR pathway that stimulates cell survival upon activation. Different mTOR inhibitors (everolimus and temsirolimus) and PI3K inhibitors (PX-866 and buparlisib) have been tested in phase II clinical trials, but response rates were disappointing and severe

grade 3/4 side effects were observed (reviewed in [103]). A phase III clinical trial is currently being conducted (NCT04338399) in which PI3K inhibitor buparlisib is administered with paclitaxel in patients with recurrent or metastatic HNSCC.




Figure 2. Concepts to identify therapeutic vulnerabilities in cancer cells. Left panel: Oncogene addiction. When a cancer cell harbors an activating mutation of an oncogene, a cancer cell may become completely dependent on this protein product, resulting in cell death upon targeted inhibition. Inhibition of the same albeit wild type protein impacts survival of normal cells less or not at all; these cells do not rely on this protein only for survival. Middle panel: Synthetic lethality. Inhibition of either two proteins or pathways with comparable function does not affect cell survival in normal cells. However, when a cancer cell harbors a gene mutation that inactivates a protein in such a pathway, cancer cells may become fully dependent on the other pathway. Inhibition of the remaining functional pathway will induce specific cancer cell death. Right panel: Collateral lethality. When a tumor suppressor gene is lost through loss of the chromosomal region, other neighboring genes located at that region may become dose-limiting and cause increased sensitivity for inhibiting drugs. Likewise, when a gene is lost or partially lost due to copy number losses, inhibition of a paralogue gene may specifically affect cancer cell viability. Abbreviations: WT: wild type, MUT: mutant.

The most obvious molecular candidate for a targeted approach in HNSCC seems to be EGFR. Of note, most HNSCCs show overexpression of EGFR, but never activating point mutations, which may explain the lack of response to EGFR-targeting kinase inhibitors such as gefitinib and erlotinib [4,66], suggesting that HNSCC cells might not be addicted to EGFR. Cetuximab, a human-mouse chimeric antibody directed against EGFR, has been registered for HNSCC, but it remains unclear whether its working mechanism relates to EGFR inhibition or activation of the immune system. Moreover, for HPV-positive disease, treatment with cetuximab is inferior to cisplatin and seems not to increase response rates [18,19].

4.2. Synthetic Lethality

As HNSCC is the main result of tumor suppressor gene inactivation, we have to rely on alternative concepts for targeted therapy approaches, such as synthetic lethality (Figure 2, middle panel). A classic example of synthetic lethality is the vulnerability to PARP inhibition in homologous recombination (HR) deficient tumors [104,105]. Neither the inhibition of PARP alone nor the loss of homology-directed repair impacts cell survival in normal cells. The loss of an HR gene such as *BRCA1* or *BRCA2* induces breast cancer oncogenesis. Additional loss of the second allele causes these tumors to be HR-deficient. When

PARP function is inhibited by PARP inhibitors in these HR-deficient tumors, cell death is induced. In normal HR-proficient cells, HR-directed repair rescues PARP inhibition.

During the past decade, synthetic lethality has been exploited as a therapeutic approach for most malignancies including HNSCC, but so far with limited success. Although synergy is observed when combining two treatments (radiotherapy supplemented with cisplatin, Aurora inhibition combined with Wee1 inhibition, Haspin knockout combined with Aurora inhibitor [106–108], and many others (see Section 7), these do not follow the classical synthetic lethality concepts, as in these combination treatments one drug sensitizes the response to the other. In the case of classical chemoradiotherapy with cisplatin and irradiation, both induce DNA damage. Nonmalignant cells will respond by cell cycle arrest and DNA repair, but these controls are diminished in cancer cells due to genomic alterations, therefore cancer cells suffer from additional intrinsic DNA damage by replication stress from the uncontrolled entry into S-phase. These treatments are obviously to a lesser extent also toxic to nonmalignant cells with proper cell cycle control, creating a small therapeutic window, in the case of radiotherapy enhanced by image-guided planning and intensity modulation.

Classic synthetic lethal interactions have not yet led to clinical implementation as new therapies for HNSCC. However, exploiting rewired pathways such as those that regulate cell cycle control in cancer cells by targeting an additional cell cycle regulating protein also follows the concept of synthetic lethal interactions. Recent studies suggest that loss of cell cycle control by p53 and/or p16^{Ink4A} loss of functions sensitize (pre)HNSCC cells to Wee1, Chk1 and PLK1 inhibition [84,109–112]. Normal cells may stay arrested in G1/G0, while cancer cells are forced to enter S-phase and progress through the cell cycle and have to rely on the rewired control mechanisms that act during DNA replication and G2-M. When these rewired mechanisms are inhibited, the cancer cells die.

4.3. Collateral Lethality

Tumor suppressor genes become inactivated during carcinogenesis through mutations or genomic deletion. In case of genomic deletion, a larger part of the chromosome is generally deleted and neighboring genes may be lost in this process. The collateral loss of these passenger genes can be exploited for therapy since these genes may be involved in cellular processes, while the gene dosage and associated protein expression has now been halved in the cancer cells. Loss of such passenger genes may also be compensated through redundancy by paralogue genes as postulated by Muller et al. [113]. Subsequent interference with the paralogue gene may lead to decreased cell survival (Figure 2, Right panel). To identify redundant paralogue genes after loss of a passenger gene, integrated analysis is required to match genomic copy number losses with vulnerabilities to interference with known paralogues [110,113–115].

We previously reported on indications that collateral lethality may occur in premalignant oral cells and HNSCC tumor cells in a customized screen in multiple cell lines models with 319 siRNAs, preselected from prior HNSCC lethality screens, [110]. Although we were able to identify collateral lethality for some of these hits, no druggable targets were found in this small RNA interference screen. More comprehensive screening methods that integrate copy number alterations with functional data from high-throughput genetic screens need to be applied to pinpoint new therapeutic avenues based on collateral lethality to treat HNSCC.

5. Identification of Essential Genes in (Pre)HNSCC Cells by Descriptive and Functional Genomics

Fundamental understanding of the biology of the tumor cells is required to allow preclinical identification of vulnerabilities of HNSCC, and the selection of biomarkers for clinical stratification. Together these form the initial steps towards better and more personalized treatments for HNSCC. The descriptive genomics data established by consortia such as the TCGA [3], which summarize the driver mutations and genomic changes causing HNSCC in great detail, have revealed a wealth of new molecular information (see below).

The TCGA aimed to identify new driver genes that could be targeted by small molecule inhibitors, but the identification of new therapeutic targets was insignificant due to the absence of oncogene drivers in HNSCC. However, new technological developments have allowed for a more functional genomics approach, and descriptive and functional genomics revealed complimentary insights into the drivers of HNSCC. To obtain unbiased data on essential genes and synergistic targets in tumor cells, (kinome) short hairpin RNA (shRNA) drop-out screens, microRNA expression screens, genome-wide array-based siRNA screens, CRISPR-Cas9 knockout screens and drug library screens are being used to identify essential genes in HNSCC cells and targeted inhibitors.

5.1. *shRNA (Kinome) Drop-Out Library Screens*

Small interference RNAs (siRNAs) are synthetic 20–25 nucleotide double-stranded RNA molecules that are complementary to gene transcripts, and cause degradation of the transcript or inhibition of translation upon binding [116]. Short hairpin RNAs (shRNAs) are the cloned versions of these siRNAs, inserted in plasmid or lentiviral vectors [117]. Both siRNAs and shRNAs can be pooled in so called libraries that target the whole genome or a part of the genome. Functional RNA interference kinome screens, are used to identify targetable kinases. Cells are plated in large culture dishes, infected with a lentiviral library containing the pooled shRNAs, and grown for a prolonged time after resistance-marker selection. DNA sequencing reveals the relative increase or decrease of the shRNA constructs that have a stimulating or inhibiting effect on proliferation, between freshly infected cells (t0) and study endpoint. Further validation of the identified hits is required to ensure validity and essentiality, since either off-target effects are observed with shRNAs that result in false-positive hits, or cells had been infected with multiple shRNAs leading to a specific lethality [118]. The incomplete knockdown of the target genes by shRNAs (and siRNAs) mimics drug inhibition best, but may cause essential genes to be missed in these screens. Despite these considerations, it stands out that in the shRNA screens conducted in HNSCC, many genes regulating the cell cycle, as well as DNA damage response, emerged as essential [119–123].

5.2. *Genome-Wide Array-Based siRNA Screens and microRNA Expression Library*

Pooled shRNA screens are relatively time-efficient compared to array-based siRNA screens that demand large scale experiments with extensive robotics (see below), but amplification of the shRNA libraries may cause the library not to be fully representative. In addition, shRNAs are processed as microRNA genes, and consequently may behave as miRNAs, the small 20–25 base pairs of single stranded RNAs that are naturally expressed in cells and regulate gene expression by targeting multiple RNA transcripts [116,124]. An alternative is array-based siRNA screens with synthetic and optimized RNA molecules, which are not processed but directly act on the transcripts in the cells. These screens are conducted in 96 or 384 well plates and only one siRNA pool (a mix of a few siRNAs complementary to the same transcript) targeting only one single gene is administered per well, together with a pre-optimized lipid transfection reagent. Cells are added and usually 96 h after transfection cell viability can be measured and essential hits can be identified by bioinformatic analysis [125]. Several genome-wide and sub-genome custom library siRNA screens have been conducted in HNSCC cell lines as well as premalignant oral cell lines to identify essential genes [109,110,126–128]. Again, many genes involved in cell cycle regulation, DNA damage response and mitotic spindle regulation have been identified as potential therapeutic candidates for HNSCC. In addition, siRNA screens have been conducted to identify biomarkers for therapy response or combination therapies [129]. To identify the hits that are tumor cell-specific, the siRNAs are also tested in nonmalignant cells. These unbiased screens also reveal tumor-specific hits that are not easily explained, such as splice factors or ribosomal genes [110,127]. Some genes appear to be essential for all cells including normal cells and are generally involved in protein homeostasis, such

as ubiquitin genes, or protein trafficking. However, most genes target tumor cells more effectively than normal cells.

A similar approach to array-based siRNA screens is the use of array-based microRNA expression libraries to identify putative targets for therapy. The overexpression of the microRNAs inhibits the expression of a variety of target genes, causing an effect on cell proliferation or other cellular processes. Expression of microRNAs miR-181a, -326, and -345 have been reported to specifically kill HNSCC tumor cells by decreasing ATM expression [130].

5.3. CRISPR-Cas9 Knockout Screens

The CRISPR-Cas9 genome editing approaches have accelerated functional genomic screens both to identify cancer cell vulnerabilities and elucidate gene function. CRISPR-Cas9 knockout screens, contrary to siRNA and shRNA libraries, enable complete knockout of target genes, which have become a game changer in the field of functional genomics. The Cas9 endonuclease is directed to specific loci in the genome (the PAM sequence) by so called guide RNAs (gRNAs) that have a sequence complementary to the gene of interest. The DNA break induced by Cas9 and the gRNA in the gene of interest is repaired by error-prone NHEJ, which leads to a deletion or insertion and functional knockout of the gene. Screens can either be using a pooled lentiviral library with cloned gRNAs, followed by library sequencing to identify depleted or enriched gRNAs targeting essential genes, or by using an array-based synthetic gRNA approach with viability readout [131]. Several reports have been published that utilized a CRISPR screen approach to uncover biological mechanisms and essential genes for therapy [106,132–135]. Furthermore, as part of the Cancer Dependency Map Project (the Wellcome Sanger Institute and the Broad Institute), several HNSCC cell lines have already been screened and these datasets are publicly available [136–138]. Similarly, to shRNA and siRNA screens, lethal hits can be missed in CRISPR screens, because of unspecific gRNAs or gRNAs that induce functional splice variants in the gene. The latest versions of libraries have been optimized, however, and are very specific. Limitations are that a fully active NHEJ system must be available in cells.

5.4. Drug Library Screens

Many investigators choose a shortcut and directly screen for cancer cell vulnerabilities by using drug libraries. Many commercial drug libraries are available to screen for targeted inhibitors that may impact survival of cell lines *in vitro*. Multiple array-based drug screens have been performed in HNSCC cell lines and primary HNSCC cells in 2D tissue culture, to identify synergistic drug combinations or new targets for therapy [139–144]. These screens prominently identified many cell cycle and DNA damage response inhibitors impacting survival of HNSCC cells. Vulnerability data from drug screens in many cancer cell lines including HNSCC are collected in datasets such as Genomics of Drug Sensitivity in Cancer (GDSC) (The Cancer Genome Project by the Wellcome Sanger Institute and Massachusetts General Hospital Cancer Center) [145]. It would be of interest to obtain survival data of premalignant oral cells with drug libraries to exploit treatments for high-risk premalignant changes. In addition, further research is needed to move these *in vitro* observations into clinical trials. The obvious limitations of drugs screens are that these are biased, in that many inhibitors have off-target effects and inhibit a multitude of proteins, which complicates the understanding of the workings mechanism of a particular drug and hampers insights into the underlying biological process.

5.5. Descriptive Genomics Technologies

Descriptive genomics encompasses large scale DNA analyses, transcriptomics and proteomics, and can be applied on both cultured cells and patient samples. These approaches describe and analyze the molecular landscape of cancer including somatic mutations and copy number alterations, profiles of methylation, gene expression, microRNAs, and proteins, the latter including post-translational modifications such as phosphorylation.

Data are collected in publicly available databases, which allows for computational biology approaches by any research to uncover new tumor vulnerabilities and biomarkers for therapy. Several initiatives by large consortia to perform in depth profiling of a multitude of cancers have been initiated in the last decade, and both large scale genomics data and data on sensitivity to drug libraries are collected, by, for example, The Cancer Genome Atlas Network [3], The Human Protein Atlas [146] and L1000 [147] amongst others (reviewed in [148]). Initiatives must be large scale as the significance of modeled predictions on tumor classification or outcome directly increases with the number of analyzed samples. However, technical aspects such as sequencing depth and data of normal control samples also support the interpretation of these data. Several *in silico* analysis of HNSCC have now been published [142,149–151]. Furthermore, machine learning models are developed to predict HNSCC tumor progression [152], along with other HNSCC data sources as summarized by Willems et al. [153]. Predicted tumor vulnerabilities and potential biomarkers from computational approaches are however hypothesis-generating and these hypotheses will need to be tested by both *in vitro* and *in vivo* preclinical studies before translation into the clinic. To further utilize the abundance of data obtained through genomics, transcriptomics, and proteomics, together with tumor cell vulnerability data obtained through functional genomics approaches such as siRNA, CRISPR or drug library screens, computational platforms have been developed to identify disease- and subgroup-specific putative targets while estimating the efficiency and toxicity based on modeled predictions through *in silico* analysis [148]. Although the data is out there, understanding the underlying biological principles and translating these to new therapeutic avenues is a challenge for the future.

6. Changes in Cell Cycle in HNSCC

Unbiased screening for vulnerabilities in HNSCC cells has uncovered that inhibition of proteins that regulate cell cycle and DNA damage response impact survival of both premalignant oral cells and HNSCC (see above). A recent proteogenomic study using 108 HPV-negative HNSCC patient tumors strengthened this observation by uncovering that important drivers of HNSCC carcinogenesis act in the cell cycle [151]. The cell cycle is compromised in most HPV-negative HNSCC and premalignant cells through loss of function of p53, p16^{Ink4A} (mutations, methylations and focal losses of chromosomal locus 9p21) and frequent amplification of cyclin D1 (Figure 3) [3,26]. In normal cells, cellular growth stimulatory signals (mitogens) induce cyclin D1 expression and consequently CDK4/6 activation, Rb phosphorylation, E2F release and transition from G1- to S-phase. This is counteracted by p16^{Ink4A} induced arrest [154–156]. Cellular stress leads to increased stabilization of p53 through posttranslational modifications of both p53 and its E3 ubiquitin ligase MDM2 which becomes inactive by the modification. If DNA damage occurs, the ATM-Chk2 signaling cascade is activated [155,157–167]. The p53 protein acts as tetrameric stress-induced transcription factor and induces p21^{Cip1} expression, causing the inhibition of the cyclin-CDK complexes, and cells stay in G1-phase or arrest in S-phase and particularly G2-phase to support efficient DNA repair or induce apoptosis [160,161,163,167–169]. Unscheduled S-phase entry induces replication stress and subsequent DNA damage, and consequently cell death by p53 induction. Remarkably, knockout of p53 or p21^{Cip1} reduces the replication stress induced DNA damage, a counterintuitive finding in mouse cells after knockout of all Rb proteins [170]. This observation was made after starvation induced replication stress in engineered cells, which might differ from the real situation in cancer cells. Although the precise mechanisms remain unclear, most tumor cells as well as premalignant cells suffer from replication stress induced DNA damage.

Release from cellular stress inducers results in degradation of p53 by MDM2, and allows restart of cell cycle progression by multiple mechanisms [160,168]. Upon loss of p53, p16^{Ink4A} or both in cancer cells, the progression from G1- to S-phase is especially impacted due to loss of the G1-checkpoint. In addition, DNA damage induced cell cycle arrest is inhibited by the loss of p53 and, together with an altered expression of cyclin D1, HNSCC cells are unperturbed continuing from G1- into S-phase [3,4,171]. The involvement of cyclin

D1 goes beyond complexing with CDK4/6 alone, since it is also involved in regulation of the DNA damage response through BRCA2, RAD51 and p21^{Cip1}, and different chromatin modification pathways (as reviewed in [172]). Together, these frequently occurring genetic alterations result in a rewired cell cycle and deregulated DNA damage response, explaining the tumor vulnerability to regulators of these processes.




Figure 3. Cell cycle control in premalignant mucosal cells and HPV-negative HNSCC is impaired through frequent inactivation of p53 by (point) mutations. P16^{Ink4A} encoded by the *CDKN2A* gene at chromosome 9p21 is frequently inactivated by either mutations or promotor methylation, or focal homozygous losses of chromosome 9p21. Combined with the amplification of the *CCND1* gene which encodes for cell cycle regulating protein cyclin D1, these alterations result in loss of normal cell cycle control and increase tolerance for DNA damage and aneuploidy as often observed in cancer cells.

Many classical cytotoxic chemotherapeutic agents used in cancer therapy exploit alterations in cell cycle regulation in cancer cells. As mentioned above (Figure 1), the crosslinking agent cisplatin remains the first choice of chemotherapy in HNSCC combined with radiotherapy. As reviewed by Williams and Stoeber, cisplatin treatment efficiently affects DNA replication in S-phase and the subsequent G2-phase [173]. Cells arrest in the S- and G2-phase to allow time for DNA repair. Additionally, classically applied chemotherapeutic agents in standard oncological treatment protocols like 5-fluorouracil (5-FU, a thymidylate synthase inhibitor), methotrexate (an inhibitor of dihydrofolate reductase), irinotecan/camptothecin (inhibitors of topomerase I) and the RNR-complex inhibitor and nucleotide analogue gemcitabine, all affect DNA replication and S-phase progression [173,174]. Tubulin-targeting agents docetaxel, paclitaxel and vincristine furthermore interfere with mitotic progression by stabilizing the spindles [175]. Although these chemotherapeutics target both malignant and rapid proliferating untransformed cells and may cause severe toxicities, these therapeutics illustrate that the mechanisms underlying DNA replication and cell cycle progression harbor targetable vulnerabilities in HNSCC and other cancer cells (Table 1).

7. Clinical Perspective

7.1. Targeting the Cell Cycle for Therapy

During the last decade, a novel interest emerged in targeting the cell cycle in HNSCC (Table 1) as well as other cancers [155,176–186]. The therapeutic effects of classical cytotoxic agents and γ -irradiation already point to the efficacy of targeting the rewired cell cycle in cancer cells. To execute more effective targeted treatments protocols for HNSCC in clinical care, the research objective is to make treatments more efficient and with less

collateral damage in nonmalignant cells. The application of functional genomic approaches by mRNA interference with siRNAs and shRNAs, microRNA expression libraries, high-throughput CRISPR-Cas9 screens, and drug library screens enabled the identification of essential genes specifically to HNSCC cells but at a lesser or no extent to nonmalignant cells [121,127,137,187,188]. Several S-phase and DNA damage related genes have been identified as potential targets for treatment of HNSCC as well as high-risk premalignant changes (Table 1). *ATM* mRNA knockdown was reported to be lethal when using microRNA expressing oligonucleotides [187]. The complexity of the ATM protein as well as redundancy in the kinase domain of ATM with other PIKK-family members, complicates the development of specific inhibitors with sufficient bioavailability *in vivo*. Nevertheless, a phase I clinical trial with the new ATM inhibitor AZD0156 is now conducted in solid tumors (NCT02588105) [111,187,189–192]. Similarly, PIKK-family member *ATR* was identified as an essential gene in HNSCC, but small molecule kinase inhibitors has lacked specificity in clinical trials so far. Recent developments uncovered highly potent small molecules against ATR and DNA-PK, and clinical trials are conducted in HNSCC with these inhibitors both as single agent and in combination (Table 1; ATR: NCT04576091, NCT04491942; DNA-PK: NCT04533750, NCT01353625) [48,111,177,193,194].

Interference of Aurora proteins, *FOXM1*, *KIF11* and *PLK1* showed promising results *in vitro* and *in vivo* in HNSCC, but these molecular targets are currently not tested in HNSCC in clinical trials [107,109,127,188,195–204]. Furthermore, therapeutic inhibition of S-phase regulator CDC7, the function of which is essential for origin firing and replication fork formation [205], is currently clinically tested in a phase I trial with CDC7 inhibitor LY3143921 in HPV-negative HNSCC patients (NCT03096054). Although the clinical application of monotherapy with CDK4/6 inhibitors might not be suitable for HNSCC, clinical trials in combination with cetuximab, radiotherapy, PI3K/mTOR inhibition or anti-PD-(L)1 antibodies are currently being conducted (NCT03065062, NCT03024489, NCT04000529) [206–208].

Table 1. Preclinical studies referring to druggable hits in cell cycle control.

Target	Inhibitor or Interference Method		HPV	Ref
ATM	Antisense oligodeoxynucleotides	<i>In vitro</i>	U	[189]
	Nanoparticles with AS-ODNs	<i>In vitro</i>	U	[190]
	Antisense oligodeoxynucleotides	<i>In vitro, in vivo</i>	Positive	[191]
	microRNA expression	<i>In vitro</i>	Negative	[187]
	AZD0156	<i>In vitro</i>	Negative	[209]
ATR	KU-55933 +/- photons +/- protons	<i>In vitro</i>	Both	[210]
	siRNA interference	<i>In vitro</i>	U	[193]
	AZD6738 +/- KU-0060648	<i>In vitro</i>	U	[48]
	AZD6738 +/- Paclitaxel or Cisplatin	<i>In vitro, in vivo</i>	Both	[194]
	AZD6738; VX-970	<i>In vitro</i>	Negative	[209]
AURORA	VE-821 +/- photons +/- protons	<i>In vitro</i>	Both	[210]
	siRNA interference +/- Paclitaxel	<i>In vitro</i>	Negative	[199]
	R763, Alisertib ^a	<i>In vitro</i>	U	[200]
	Alisertib ^a +/- MG132	<i>In vitro, in vivo</i>	Positive	[201]
	Danusertib ^b	<i>In vitro, in vivo</i>	Both	[188]
CDC7	Alisertib ^a +/- Adavosertib ^c	<i>In vitro, in vivo</i>	Negative	[107]
	Alisertib ^a ; Danusertib ⁱ	<i>In vitro</i>	Negative	[209]
	VX-680 +/- Haspin ^{KO} or CHR-3464	<i>In vitro</i>	Negative	[106]
	XL413 +/- Cisplatin and Fluorouracil	<i>In vitro, in vivo</i>	Both	[202]
	CDK4/6	Palbociclib + Cetuximab	Phase I trial	Both
Abemaciclib +/- Torin2 or Everolimus		<i>In vitro, in vivo</i>	U	[207]
Ribociclib +/- RT		<i>In vitro</i>	U	[208]
Palbociclib + Cetuximab		Phase II trial	Negative	[211]

Table 1. Cont.

Target	Inhibitor or Interference Method		HPV	Ref
	Palbociclib + Cisplatin	Phase I trial, <i>in vivo</i> , <i>in vitro</i>	Negative	[212]
	Abemaciclib + Metformin	<i>In vivo</i>	U	[213]
	Palbociclib + Navitoclax	<i>In vitro</i>	Both	[214]
	Palbociclib	<i>In vitro</i>	Negative	[209]
Chk1/2	PF-00477736 +/- RT	<i>In vitro</i>	Positive	[215]
	AZD7762 +/- Cisplatin	<i>In vitro</i>	Negative	[216]
	siRNA interference	<i>In vitro</i>	U	[193]
	MK-8776 +/- Adavosertib ^c	<i>In vitro</i>	U	[217]
	CCT244747 +/- RT +/- Paclitaxel	<i>In vitro</i> , <i>in vivo</i>	Both	[218]
	Prexasertib ^d	Phase I trial	U	[219]
	AZD7762 or Rabusertib ^e or MK8776 + / - RT	<i>In vitro</i>	Positive	[220]
	Prexasertib ^d +/- RT +/- Cetuximab	<i>In vitro</i> , <i>in vivo</i>	Both	[221]
	AZD7762	<i>In vitro</i> , <i>in vivo</i>	Both	[198]
	Prexasertib ^d	Phase I trial	Both	[222]
	siRNA interference; Prexasertib ^d	<i>In vitro</i>	Both	[112]
	siRNA interference	<i>In vitro</i>	Negative	[126]
	siRNA interference; Rabusertib ^e ; Prexasertib ^d ; MK-8776; PF-477736	<i>In vitro</i>	Both	[111]
	PF-00477736 +/- Alpelisib ^h	<i>In vitro</i> , <i>in vivo</i>	U	[223]
	shRNA; MK-8776 +/- Niraparib + / - RT	<i>In vitro</i> , <i>in vivo</i>	Both	[224]
	Prexasertib ^d +/- Cisplatin + / - RT	<i>In vitro</i> , <i>in vivo</i>	Both	[225]
	Prexasertib ^d ; MK8776	<i>In vitro</i>	Negative	[209]
	Prexasertib ^d	<i>In vivo</i>	Both	[226]
DNA-PK	KU-0060648 +/- AZD6738	<i>In vitro</i>	U	[48]
	CC-115	<i>In vitro</i>	Negative	[209]
	KU-57788 or IC87361 +/- Olaparib or Veliparib +/- RT	<i>In vitro</i>	Both	[227]
	NU7441 +/- Olaparib	<i>In vitro</i> , <i>in vivo</i>	Negative	[228]
	KU-57788 +/- photons +/- protons	<i>In vitro</i>	Both	[210]
FOXM1	siRNA interference	<i>In vitro</i>	Positive	[203]
	Thiostrepton	<i>In vitro</i> , <i>in vivo</i>	Positive	[204]
KIF11	Ispinesib	Phase II trial	U	[195]
	siRNA interference; Ispinesib	<i>In vitro</i> , <i>in vivo</i>	Negative	[127]
PLK1	siRNA interference +/- RT	<i>In vitro</i> , <i>in vivo</i>	Negative	[196]
	BI2536	<i>In vitro</i>	U	[197]
	Volasertib ^f	<i>In vitro</i> , <i>in vivo</i>	Both	[198]
	siRNA interference; Volasertib ^f ; GSK461364; Rigosertib ^g ; HMN-214	<i>In vitro</i> , <i>in vivo</i>	Neg, preHN	[109]
	BI2536	<i>In vitro</i>	Negative	[209]
Wee1	Adavosertib ^c	<i>In vitro</i> , <i>in vivo</i>	Both	[121]
	Adavosertib ^c +/- Cisplatin	<i>In vitro</i> , <i>in vivo</i>	Negative	[229]
	Adavosertib ^c +/- Cisplatin	<i>In vitro</i> , <i>in vivo</i>	Positive	[230]
	Adavosertib ^c +/- RT	<i>In vitro</i>	Positive	[220]
	Adavosertib ^c +/- Vorinostat	<i>In vitro</i> , <i>in vivo</i>	Negative	[231]
	Adavosertib ^c	<i>In vitro</i>	Negative	[232]
	Adavosertib ^c	<i>In vitro</i> , <i>in vivo</i>	Both	[198]
	Adavosertib ^c	<i>In vitro</i> , <i>in vivo</i>	U	[233]
	Adavosertib ^c + Cisplatin + Docetaxel	Phase I trial	Both	[234]
	siRNA interference	<i>In vitro</i>	Negative	[126]
	Adavosertib ^c	<i>In vitro</i>	Negative	[209]
	Adavosertib ^c	<i>In vitro</i>	Positive	[235]

Table 1. Cont.

Target	Inhibitor or Interference Method		HPV	Ref
	Adavosertib ^c +/- Alisertib ^a	<i>In vitro, in vivo</i>	Negative	[107]
	siRNA interference; Adavosertib ^e	<i>In vitro</i>	Neg, preHN	[110]
	Adavosertib ^c	<i>In vitro</i>	Positive	[236]
	Adavosertib ^c + Cisplatin	Phase I trial	U	[237]
	shRNA; Adavosertib ^e +/- Niraparib +/- RT	<i>In vitro, in vivo</i>	Both	[224]
<i>Ledgens</i>	^a MLN8237	^f BI6727	RT radiotherapy	
	^b PHA-739358	^g ON-01910	U unknown	
	^c AZD1775/MK-1775	^h BYL-71	preHN premalignant oral cells	
	^d LY2606368	ⁱ AMG900	KO CRISPR-mediated knockout	
	^e LY2603618			

In several reports the susceptibility of HNSCC to Chk1 and Wee1 inhibitors has been published (Table 1) [107,110–112,121,126,193,198,209,215–226,229–237]. Chk1 is mainly involved in S-phase regulation upon stalled replication forks, which makes it a feasible target for cancers with high replication stress such as HNSCC [180,238–242]. Clinical trials with the Chk1 inhibitor SRA737 and the dual Chk1/Chk2 inhibitor prexasertib have been completed for treatment of solid tumors amongst which is HNSCC (NCT01115790, NCT02555644, NCT02797964).

Wee1 is a critical regulator of the G2/M-checkpoint through CDK1 phosphorylation, which inactivates the protein, but also plays a role in S-phase regulation by inactivating CDK2 phosphorylation against a background of replication problems [180,186,241,242]. The Wee1 inhibitor adavosertib is the only compound tested in clinical trials, including several studies in HNSCC, either as monotherapy or in combination with conventional chemotherapy and radiation (NCT04460937, NCT03028766) [177]. A phase I trial with promising results in HNSCC has been published [234].

Premalignant cells have been shown to be specifically vulnerable to PLK1 and Wee1 inhibition, whilst normal oral keratinocyte and fibroblast cells are not affected [109,110]. This strengthens the hypothesis that loss of p53, p16^{Ink4A} and to a lesser extent cyclin D1 amplification as seen in premalignant cells [26], sensitizes cells to these inhibitors, which may support the initiation of clinical trials. Particularly the application as monotherapy for treatment of high-risk premalignant changes characterized by morphological and genetic changes, might contribute to eradication of visible changes such as leukoplakias, but may also lead to less recurrent disease after treatment of the index tumor and improve overall survival rates of HNSCC patients.

Altogether, these preclinical and clinical trial data indicate the feasibility of targeting DNA replication and cell cycle progression in HNSCC. It should be noted, however, that the data in cell lines also point towards tumor-specific differences in response, possibly reflecting inter-tumor heterogeneity of HNSCC [90]. Clinical trials will provide important additional information on efficacy and biomarkers for patient selection in the future, as well as useful combination therapies. For many targets, small molecule inhibitors that show better bio-distribution and target-specificity *in vivo* and alternative therapeutic molecules such as protein degrading (PROTAC) molecules [243] are needed for more successful clinical implementation and increased responses.

7.2. Combining PD-L1 Antibodies with DNA Damaging Agents

Preclinical research has shown that activation of the cGAS/STING pathway after DNA damage enhances immune cell infiltration into the tumor microenvironment (TME) through secretion of pro-inflammatory type I interferon (IFN) and induction of a senescence-associated secretory phenotype (SASP) [244]. More data has been published recently indicating that degree of successful cGAS/STING pathway activation after classical DNA-damaging chemotherapeutics and radiation therapy influences treatment outcome through antitumor immunity (as reviewed in [245]). By inducing DNA damage and activation of T-

cells, two hallmarks of cancer are being exploited, with potentially better responses and less therapy resistance [68,246]. To reduce toxicity as observed with classical chemotherapies and to enhance tumor-specificity, further research should determine whether combination of immune checkpoint inhibitors such as anti-PD-(L)1 antibodies with targeted agents improves HNSCC response rates and survival. As shown in other tumor types (reviewed in [246]), inhibitors targeting molecules such as ATR, Wee1 and Chk1, which induce replication-associated DNA damage and potentially cGAS/STING pathway activation through subsequent cytosolic DNA fragments in HNSCC, are promising candidates for combination treatments in HNSCC. Clinical trials are currently conducted combining ATR and Chk1 inhibitors with anti-PD-(L)1 antibodies in solid tumors including HNSCC (NCT04266912, NCT02264678, NCT04095273, NCT03495323) [246].

8. Conclusions

Survival rates for late stage HNSCC remain disappointing and while protocols for first-line treatment have been optimized in recent decades, this has not fundamentally improved survival since the implementation of cisplatin in 1977. The FDA approval of cetuximab in 2007 and two anti-PD-1 antibodies in 2016 have increased the arsenal, but the response rates still leave much to be desired and the lack of reliable biomarkers for response hamper implementation given the costs and associated toxicities of these new agents. Treatments with targeted agents have hardly been introduced in routine care as activating mutations in oncogenes are scarce in HNSCC, and clinical results so far were disappointing. In the last decade, through unbiased screening of siRNA and shRNA libraries, inhibitor libraries and more recently guide RNA libraries utilizing CRISPR-Cas9 technology, a better understanding of HNSCC vulnerabilities was established. Especially deregulation of the cell cycle in HNSCC has emerged as a candidate to be exploited for clinical use, not only to treat invasive cancers but also high-risk premalignant mucosal cells. In addition, combination therapies with immune checkpoint inhibitors and targeted agents that affect DNA damaging, such as ATR, Chk1 and Wee1 inhibitors, have potential to further increase response rates and survival. Better understanding of genomic alterations and loss of passenger genes will furthermore contribute to the identification of targets through synthetic and collateral lethality, which will expand our pre-clinical toolbox to uncover new HNSCC vulnerabilities in specific genetic backgrounds. Lastly, research should increase the focus on premalignant mucosal cells, and test vulnerability of these cells to new therapeutic compounds. By targeting these lesions together with the index tumor during treatment, recurrent and secondary tumors may be prevented. Furthermore, diagnosis and treatment of high-risk premalignant lesions by targeted inhibitors with their generally mild toxicity profiles could prevent the malignant progression of these precancers into difficult to treat HNSCC tumors.

Author Contributions: Writing—original draft preparation, A.M.v.H. and R.H.B.; writing—review and editing, A.M.v.H. and R.H.B., visualization, A.M.v.H.; supervision, R.H.B. All authors have read and agreed to the published version of the manuscript.

Funding: Funding was provided by the Cancer Center Amsterdam foundation of Amsterdam UMC, the Netherlands.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: The manuscript was partly based on the thesis of the first author, Anne M. van Harten. Figures 1–3 were created with [BioRender.com](https://www.biorender.com) (accessed on 3 May 2021).

Conflicts of Interest: AMH declares no conflict of interest. RHB has a collaboration with InteRNA Technologies BV, AstraZeneca and NLC-Health-Venture-Builder, and research support from GenMab BV, Bristol Meyers Squibb, and Hanarth Foundation. RHB is on the advisory board of Nanobiotix.

References

1. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2018**, *68*, 394–424. [\[CrossRef\]](#)
2. Leemans, C.R.; Snijders, P.J.F.; Brakenhoff, R.H. The molecular landscape of head and neck cancer. *Nat. Rev. Cancer* **2018**, *18*, 269–282. [\[CrossRef\]](#)
3. Lawrence, M.S.; Sougnez, C.; Lichtenstein, L.; Cibulskis, K.; Lander, E.; Gabriel, S.B.; Getz, G.; Ally, A.; Balasundaram, M.; Birol, I.; et al. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature* **2015**, *517*, 576–582. [\[CrossRef\]](#)
4. Johnson, D.E.; Burtneess, B.; Leemans, C.R.; Lui, V.W.Y.; Bauman, J.E.; Grandis, J.R. Head and neck squamous cell carcinoma. *Nat. Rev. Dis. Prim.* **2020**, *6*, 92. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Leemans, C.R.; Braakhuis, B.J.M.; Brakenhoff, R.H. The molecular biology of head and neck cancer. *Nat. Rev. Cancer* **2011**, *11*, 9–22. [\[CrossRef\]](#)
6. Brakenhoff, R.H.; Wagner, S.; Klussmann, J.P. Molecular Patterns and Biology of HPV-Associated HNSCC. In *HPV Integration in Head and Neck Squamous Cell Carcinomas: Cause and Consequence*; Recent Results in Cancer Research; Springer: Berlin/Heidelberg, Germany, 2017; Volume 206, pp. 57–72, ISBN 978-3-319-43578-7.
7. Doorbar, J.; Egawa, N.; Griffin, H.; Kranjec, C.; Murakami, I. Human papillomavirus molecular biology and disease association. *Rev. Med. Virol.* **2015**, *25*, 2–23. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Graham, S.V. The human papillomavirus replication cycle, and its links to cancer progression: A comprehensive review. *Clin. Sci.* **2017**, *131*, 2201–2221. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Furquim, C.P.; Pivovar, A.; Amenábar, J.M.; Bonfim, C.; Torres-Pereira, C.C. Oral cancer in Fanconi anemia: Review of 121 cases. *Crit. Rev. Oncol. Hematol.* **2018**, *125*, 35–40. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Nalepa, G.; Clapp, D.W. Fanconi anaemia and cancer: An intricate relationship. *Nat. Rev. Cancer* **2018**, *18*, 168–185. [\[CrossRef\]](#)
11. Romick-Rosendale, L.E.; Lui, V.W.Y.; Grandis, J.R.; Wells, S.I. The Fanconi anemia pathway: Repairing the link between DNA damage and squamous cell carcinoma. *Mutat. Res. Fundam. Mol. Mech. Mutagen.* **2013**, *743–744*, 78–88. [\[CrossRef\]](#)
12. Van Zeeburg, H.J.T.; Snijders, P.J.F.; Wu, T.; Gluckman, E.; Soulier, J.; Surralles, J.; Castella, M.; Van Der Wal, J.E.; Wennerberg, J.; Califano, J.; et al. Clinical and molecular characteristics of squamous cell carcinomas from Fanconi anemia patients. *J. Natl. Cancer Inst.* **2008**, *100*, 1649–1653. [\[CrossRef\]](#)
13. Van Zeeburg, H.J.T.; Snijders, P.J.F.; Pals, G.; Hermsen, M.A.J.A.; Rooimans, M.A.; Bagby, G.; Soulier, J.; Gluckman, E.; Wennerberg, J.; Leemans, C.R.; et al. Generation and molecular characterization of head and neck squamous cell lines of Fanconi anemia patients. *Cancer Res.* **2005**, *65*, 1271–1276. [\[CrossRef\]](#)
14. Pulte, D.; Brenner, H. Changes in survival in head and neck cancers in the late 20th and early 21st century: A period analysis. *Oncologist* **2010**, *15*, 994–1001. [\[CrossRef\]](#)
15. Barnes, L.; Eveson, J.W.; Reichart, P.; Sidransky, D. *World Health Organization Classification of Tumours. Pathology & Genetics. Head and Neck Tumours*; International Agency for Research on Cancer (IARC): Lyon, France, 2005; ISBN 9283224175.
16. Alabi, O.; O’Neill, J.P. ‘Good cancer gone bad’: A narrative review of HPV oropharyngeal cancer and potential poor outcomes. *Eur. Arch. Oto-Rhino-Laryngol.* **2020**, *277*, 2185–2191. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Nauta, I.H.; Rietbergen, M.M.; Van Bokhoven, A.A.J.D.; Bloemena, E.; Lissenberg-Witte, B.I.; Heideman, D.A.M.; De Jong, R.J.B.; Brakenhoff, R.H.; Leemans, C.R. Evaluation of the eighth TNM classification on p16-positive oropharyngeal squamous cell carcinomas in the Netherlands and the importance of additional HPV DNA testing. *Ann. Oncol.* **2018**, *29*, 1273–1279. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Mehanna, H.; Robinson, M.; Hartley, A.; Kong, A.; Foran, B.; Fulton-Lieuw, T.; Dalby, M.; Mistry, P.; Sen, M.; O’Toole, L.; et al. Radiotherapy plus cisplatin or cetuximab in low-risk human papillomavirus-positive oropharyngeal cancer (De-ESCALaTE HPV): An open-label randomised controlled phase 3 trial. *Lancet* **2019**, *393*, 51–60. [\[CrossRef\]](#)
19. Gillison, M.L.; Trotti, A.M.; Harris, J.; Eisbruch, A.; Harari, P.M.; Adelstein, D.J.; Jordan, R.C.K.; Zhao, W.; Sturgis, E.M.; Burtneess, B.; et al. Radiotherapy plus cetuximab or cisplatin in human papillomavirus-positive oropharyngeal cancer (NRG Oncology RTOG 1016): A randomised, multicentre, non-inferiority trial. *Lancet* **2019**, *393*, 40–50. [\[CrossRef\]](#)
20. Blot, W.J.; McLaughlin, J.K.; Winn, D.M.; Austin, D.F.; Greenberg, R.S.; Preston-Martin, S.; Bernstein, L.; Schoenberg, J.B.; Stemhagen, A.; Fraumeni, J.F. Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res.* **1988**, *48*, 3282–3287.
21. Cramer, J.D.; Burtneess, B.; Le, Q.T.; Ferris, R.L. The changing therapeutic landscape of head and neck cancer. *Nat. Rev. Clin. Oncol.* **2019**, *16*, 669–683. [\[CrossRef\]](#)
22. Osazuwa-Peters, N.; Adjei Boakye, E.; Chen, B.Y.; Tobo, B.B.; Varvares, M.A. Association between head and neck squamous cell carcinoma survival, smoking at diagnosis, and marital status. *JAMA Otolaryngol. Neck Surg.* **2017**, *144*, 43–50. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Smeets, S.J.; Brakenhoff, R.H.; Ylstra, B.; Van Wieringen, W.N.; Van De Wiel, M.A.; Leemans, C.R.; Braakhuis, B.J.M. Genetic classification of oral and oropharyngeal carcinomas identifies subgroups with a different prognosis. *Cell. Oncol.* **2009**, *31*, 291–300. [\[CrossRef\]](#)
24. Slaughter, D.P.; Southwick, H.W.; Smejkal, W. “Field cancerization” in oral stratified squamous epithelium. Clinical implications of multicentric origin. *Cancer* **1953**, *6*, 963–968. [\[CrossRef\]](#)

25. Tabor, M.P.; Brakenhoff, R.H.; Ruijter-Schippers, H.J.; Kummer, J.A.; Leemans, C.R.; Braakhuis, B.J.M. Genetically altered fields as origin of locally recurrent head and neck cancer: A retrospective study. *Clin. Cancer Res.* **2004**, *10*, 3607–3613. [[CrossRef](#)] [[PubMed](#)]
26. De Boer, D.V.; Brink, A.; Buijze, M.; Stigter-van Walsum, M.; Hunter, K.D.; Ylstra, B.; Bloemena, E.; Leemans, C.R.; Brakenhoff, R.H. Establishment and genetic landscape of precancer cell model systems from the head and neck mucosal lining. *Mol. Cancer Res.* **2018**, *17*, 120–130. [[CrossRef](#)] [[PubMed](#)]
27. Braakhuis, B.; Leemans, C.; Brakenhoff, R. Expanding fields of genetically altered cells in head and neck squamous carcinogenesis. *Semin. Cancer Biol.* **2005**, *15*, 113–120. [[CrossRef](#)]
28. Braakhuis, B.J.M.; Tabor, M.P.; Kummer, J.A.; Leemans, C.R.; Brakenhoff, R.H. A genetic explanation of Slaughter’s concept of field cancerization: Evidence and clinical implications. *Cancer Res.* **2003**, *63*, 1727–1730.
29. Califano, J.; Van der Riet, P.; Westra, W.; Nawroz, H.; Clayman, G.; Piantadosi, S.; Corio, R.; Lee, D.; Greenberg, B.; Koch, W.; et al. Genetic progression model for head and neck cancer: Implications for field cancerization. *Cancer Res.* **1996**, *56*, 2488–2492. [[CrossRef](#)]
30. Smetsers, S.E.; Velleuer, E.; Dietrich, R.; Wu, T.; Brink, A.; Buijze, M.; Deeg, D.J.H.; Soulier, J.; Leemans, C.R.; Braakhuis, B.J.M.; et al. Noninvasive molecular screening for oral precancer in Fanconi anemia patients. *Cancer Prev. Res.* **2015**, *8*, 1102–1111. [[CrossRef](#)] [[PubMed](#)]
31. Graveland, A.P.; Golusinski, P.J.; Buijze, M.; Douma, R.; Sons, N.; Kuik, D.J.; Bloemena, E.; Leemans, C.R.; Brakenhoff, R.H.; Braakhuis, B.J.M. Loss of heterozygosity at 9p and p53 immunopositivity in surgical margins predict local relapse in head and neck squamous cell carcinoma. *Int. J. Cancer* **2011**, *128*, 1852–1859. [[CrossRef](#)] [[PubMed](#)]
32. Reichart, P.A.; Philipsen, H.P. Oral erythroplakia—A review. *Oral Oncol.* **2005**, *41*, 551–561. [[CrossRef](#)]
33. Brouns, E.; Baart, J.A.; Karagozoglu, K.H.; Aartman, I.H.A.; Bloemena, E.; Van der Waal, I. Malignant transformation of oral leukoplakia in a well-defined cohort of 144 patients. *Oral Dis.* **2014**, *20*, e19–e24. [[CrossRef](#)]
34. Schepman, K.P.; Van Der Meij, E.H.; Smeele, L.E.; Van Der Waal, I. Malignant transformation of oral leukoplakia: A follow-up study of a hospital-based population of 166 patients with oral leukoplakia from The Netherlands. *Oral Oncol.* **1998**, *34*, 270–275. [[CrossRef](#)]
35. Gleber-Netto, F.O.; Braakhuis, B.J.M.; Triantafyllou, A.; Takes, R.P.; Kelner, N.; Rodrigo, J.P.; Strojan, P.; Vander Poorten, V.; Rapidis, A.D.; Rinaldo, A.; et al. Molecular events in relapsed oral squamous cell carcinoma: Recurrence vs secondary primary tumor. *Oral Oncol.* **2015**, *51*, 738–744. [[CrossRef](#)]
36. Wong, K.M.; Capasso, A.; Eckhardt, S.G. The changing landscape of phase I trials in oncology. *Nat. Rev. Clin. Oncol.* **2016**, *13*, 106–117. [[CrossRef](#)] [[PubMed](#)]
37. Moscow, J.A.; Fojo, T.; Schilsky, R.L. The evidence framework for precision cancer medicine. *Nat. Rev. Clin. Oncol.* **2018**, *15*, 183–192. [[CrossRef](#)]
38. Canning, M.; Guo, G.; Yu, M.; Myint, C.; Groves, M.W.; Byrd, J.K.; Cui, Y. Heterogeneity of the head and neck squamous cell carcinoma immune landscape and its impact on immunotherapy. *Front. Cell Dev. Biol.* **2019**, *7*, 52. [[CrossRef](#)]
39. Kobayashi, K.; Hisamatsu, K.; Suzui, N.; Hara, A.; Tomita, H.; Miyazaki, T. A Review of HPV-related head and neck cancer. *J. Clin. Med.* **2018**, *7*, 241. [[CrossRef](#)] [[PubMed](#)]
40. Koyfman, S.A.; Ismaila, N.; Holsinger, F.C. Management of the neck in squamous cell carcinoma of the oral cavity and oropharynx: ASCO clinical practice guideline summary. *J. Oncol. Pract.* **2019**, *15*, 273–278. [[CrossRef](#)]
41. DeSantis, C.E.; Lin, C.C.; Mariotto, A.B.; Siegel, R.L.; Stein, K.D.; Kramer, J.L.; Alteri, R.; Robbins, A.S.; Jemal, A. Cancer treatment and survivorship statistics, 2014. *CA. Cancer J. Clin.* **2014**, *64*, 252–271. [[CrossRef](#)]
42. Szturcz, P.; Wouters, K.; Kiyota, N.; Tahara, M.; Prabhas, K.; Noronha, V.; Castro, A.; Licitra, L.; Adelstein, D.; Vermorken, J.B. Commentary on “Weekly low-dose versus three-weekly high-dose cisplatin for concurrent chemoradiation in locoregionally advanced non-nasopharyngeal head and neck cancer: A systematic review and meta-analysis of aggregate data”. *Oncologist* **2017**, *22*, 1022–1023. [[CrossRef](#)]
43. Wittes, R.E.; Cvitkovic, E.; Shah, J.; Gerold, F.P.; Strong, E.W. CIS-Dichlorodiammineplatinum(II) in the treatment of epidermoid carcinoma of the head and neck. *Cancer Treat. Rep.* **1977**, *61*, 359–366.
44. Tchounwou, P.B.; Dasari, S.; Noubissi, F.K.; Ray, P.; Kumar, S. Advances in our understanding of the molecular mechanisms of action of cisplatin in cancer therapy. *J. Exp. Pharmacol.* **2021**, *13*, 303–328. [[CrossRef](#)]
45. Dasari, S.; Bernard Tchounwou, P. Cisplatin in cancer therapy: Molecular mechanisms of action. *Eur. J. Pharmacol.* **2014**, *740*, 364–378. [[CrossRef](#)]
46. Lopez-Martinez, D.; Liang, C.-C.; Cohn, M.A. Cellular response to DNA interstrand crosslinks: The Fanconi anemia pathway. *Cell. Mol. Life Sci.* **2016**, *73*, 3097–3114. [[CrossRef](#)]
47. Rodriguez, A.; D’Andrea, A. Fanconi anemia pathway. *Curr. Biol.* **2017**, *27*, R986–R988. [[CrossRef](#)] [[PubMed](#)]
48. Hafi, H.; Dillon, M.T.; Barker, H.E.; Kyula, J.N.; Schick, U.; Paget, J.T.; Smith, H.G.; Pedersen, M.; McLaughlin, M.; Harrington, K.J. Combined ATR and DNA-PK inhibition radiosensitizes tumor cells independently of their p53 status. *Front. Oncol.* **2018**, *8*, 245. [[CrossRef](#)]
49. De Felice, F.; Polimeni, A.; Valentini, V.; Brugnoletti, O.; Cassoni, A.; Greco, A.; De Vincentiis, M.; Tombolini, V. Radiotherapy controversies and prospective in head and neck cancer: A literature-based critical review. *Neoplasia* **2018**, *20*, 227–232. [[CrossRef](#)] [[PubMed](#)]

50. Mazzola, R.; Fiorentino, A.; Ricchetti, F.; Gregucci, F.; Corradini, S.; Alongi, F. An update on radiation therapy in head and neck cancers. *Expert Rev. Anticancer Ther.* **2018**, *18*, 359–364. [[CrossRef](#)] [[PubMed](#)]
51. Maghami, E.; Koyfman, S.A.; Weiss, J. Personalizing postoperative treatment of head and neck cancers. *Am. Soc. Clin. Oncol. Educ. Book. Am. Soc. Clin. Oncol. Annu. Meet.* **2018**, *38*, 515–522. [[CrossRef](#)] [[PubMed](#)]
52. Sonveaux, P. ROS and radiotherapy: More we care. *Oncotarget* **2017**, *8*, 35482–35483. [[CrossRef](#)] [[PubMed](#)]
53. Gray, L.H.; Conger, A.D.; Ebert, M.; Hornsey, S.; Scott, O.C.A. The Concentration of oxygen dissolved in tissues at the time of irradiation as a factor in radiotherapy. *Br. J. Radiol.* **1953**, *26*, 638–648. [[CrossRef](#)]
54. Good, J.S.; Harrington, K.J. The hallmarks of cancer and the radiation oncologist: Updating the 5Rs of radiobiology. *Clin. Oncol.* **2013**, *25*, 569–577. [[CrossRef](#)]
55. Bhattacharya, S.; Asaithamby, A. Repurposing DNA repair factors to eradicate tumor cells upon radiotherapy. *Transl. Cancer Res.* **2017**, *6*, S822–S839. [[CrossRef](#)]
56. Seol, J.-H.; Shim, E.Y.; Lee, S.E. Microhomology-mediated end joining: Good, bad and ugly. *Mutat. Res. Mol. Mech. Mutagen.* **2018**, *809*, 81–87. [[CrossRef](#)]
57. Wang, C.; Lees-Miller, S.P. Detection and repair of ionizing radiation-induced DNA double strand breaks: New developments in nonhomologous end joining. *Int. J. Radiat. Oncol.* **2013**, *86*, 440–449. [[CrossRef](#)] [[PubMed](#)]
58. Dutta, A.; Eckelmann, B.; Adhikari, S.; Mokim Ahmed, K.; Sengupta, S.; Pandey, A.; Hegde, P.M.; Tsai, M.-S.; Tainer, J.A.; Weinfeld, M.; et al. Microhomology-mediated end joining is activated in irradiated human cells due to phosphorylation-dependent formation of the XRCC1 repair complex. *Nucleic Acids Res.* **2017**, *45*, 2585–2599. [[CrossRef](#)]
59. Ferguson, D.O.; Holloman, W.K.; Chen, D.J.; Kurimasa, A.; Li, G.C.; Lehnert, B.E.; Goodwin, E.H.; Chun, J.; Alt, F.W. Recombinational repair of gaps in DNA is asymmetric in *Ustilago maydis* and can be explained by a migrating D-loop model. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 5419–5424. [[CrossRef](#)] [[PubMed](#)]
60. Hanscom, T.; McVey, M. Regulation of error-prone DNA double-strand break repair and its impact on genome evolution. *Cells* **2020**, *9*, 1657. [[CrossRef](#)] [[PubMed](#)]
61. Rivera, F.; García-Castaño, A.; Vega, N.; Vega-Villegas, M.E.; Gutiérrez-Sanz, L. Cetuximab in metastatic or recurrent head and neck cancer: The EXTREME trial. *Expert Rev. Anticancer Ther.* **2009**, *9*, 1421–1428. [[CrossRef](#)]
62. Tian, Y.; Lin, J.; Tian, Y.; Zhang, G.; Zeng, X.; Zheng, R.; Zhang, W.; Yuan, Y. Efficacy and safety of anti-EGFR agents administered concurrently with standard therapies for patients with head and neck squamous cell carcinoma: A systematic review and meta-analysis of randomized controlled trials. *Int. J. Cancer* **2018**, *142*, 2198–2206. [[CrossRef](#)]
63. Vermorken, J.B.; Trigo, J.; Hitt, R.; Koralewski, P.; Diaz-Rubio, E.; Rolland, F.; Knecht, R.; Amellal, N.; Schueler, A.; Baselga, J. Open-label, uncontrolled, multicenter phase II study to evaluate the efficacy and toxicity of cetuximab as a single agent in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck who failed to respond to platinum-based therapy. *J. Clin. Oncol.* **2007**, *25*, 2171–2177. [[CrossRef](#)]
64. Ferris, R.L.; Lenz, H.-J.; Trotta, A.M.; García-Foncillas, J.; Schulten, J.; Audhuy, F.; Merlano, M.; Milano, G. Rationale for combination of therapeutic antibodies targeting tumor cells and immune checkpoint receptors: Harnessing innate and adaptive immunity through IgG1 isotype immune effector stimulation. *Cancer Treat. Rev.* **2018**, *63*, 48–60. [[CrossRef](#)]
65. Ling, D.C.; Bakkenist, C.J.; Ferris, R.L.; Clump, D.A. Role of immunotherapy in head and neck cancer. *Semin. Radiat. Oncol.* **2018**, *28*, 12–16. [[CrossRef](#)]
66. Tang, X.; He, J.; Li, B.; Zheng, Y.; Li, K.; Zou, S.; Chen, L. Efficacy and safety of gefitinib in patients with advanced head and neck squamous cell carcinoma: A meta-analysis of randomized controlled trials. *J. Oncol.* **2019**, *2019*, 1–9. [[CrossRef](#)]
67. Jin, W.J.; Erbe, A.K.; Schwarz, C.N.; Jaquish, A.A.; Anderson, B.R.; Sriramaneni, R.N.; Jagodinsky, J.C.; Bates, A.M.; Clark, P.A.; Le, T.; et al. Tumor-specific antibody, cetuximab, enhances the in situ vaccine effect of radiation in immunologically cold head and neck squamous cell carcinoma. *Front. Immunol.* **2020**, *11*, 1. [[CrossRef](#)] [[PubMed](#)]
68. Hanahan, D.; Weinberg, R.A. Hallmarks of cancer: The next generation. *Cell* **2011**, *144*, 646–674. [[CrossRef](#)] [[PubMed](#)]
69. Forster, M.D.; Devlin, M.-J. Immune checkpoint inhibition in head and neck cancer. *Front. Oncol.* **2018**, *8*, 310. [[CrossRef](#)] [[PubMed](#)]
70. Zolkind, P.; Uppaluri, R. Checkpoint immunotherapy in head and neck cancers. *Cancer Metastasis Rev.* **2017**, *36*, 475–489. [[CrossRef](#)] [[PubMed](#)]
71. Ferris, R.L. Immunology and immunotherapy of head and neck cancer. *J. Clin. Oncol.* **2015**, *33*, 3293–3304. [[CrossRef](#)]
72. Wei, S.C.; Duffy, C.R.; Allison, J.P. Fundamental mechanisms of immune checkpoint blockade therapy. *Cancer Discov.* **2018**, *8*, 1069–1086. [[CrossRef](#)]
73. Saleh, K.; Eid, R.; Haddad, F.G.H.; Khalife-Saleh, N.; Kourie, H.R. New developments in the management of head and neck cancer—Impact of pembrolizumab. *Ther. Clin. Risk Manag.* **2018**, *14*, 295–303. [[CrossRef](#)] [[PubMed](#)]
74. Qiao, X.; Jiang, J.; Pang, X.; Huang, M.; Tang, Y.; Liang, X.; Tang, Y. The evolving landscape of PD-1/PD-L1 pathway in head and neck cancer. *Front. Immunol.* **2020**, *11*, 11. [[CrossRef](#)]
75. Bardhan, K.; Anagnostou, T.; Boussiotis, V.A. The PD1:PD-L1/2 Pathway from discovery to clinical implementation. *Front. Immunol.* **2016**, *7*, 550. [[CrossRef](#)]
76. Monteiro de Oliveira Novaes, J.A.; Hirz, T.; Guijarro, I.; Nilsson, M.; Piseigna, M.A.; Poteete, A.; Barsoumian, H.B.; Fradette, J.J.; Chen, L.N.; Gibbons, D.L.; et al. Targeting of CD40 and PD-L1 pathways inhibits progression of oral premalignant lesions in a carcinogen-induced model of oral squamous cell carcinoma. *Cancer Prev. Res.* **2020**, *14*, 313–324. [[CrossRef](#)]

77. Naidoo, J.; Page, D.B.; Li, B.T.; Connell, L.C.; Schindler, K.; Lacouture, M.E.; Postow, M.A.; Wolchok, J.D. Toxicities of the anti-PD-1 and anti-PD-L1 immune checkpoint antibodies. *Ann. Oncol.* **2015**, *26*, 2375–2391. [[CrossRef](#)]
78. Donehower, L.A.; Soussi, T.; Korkut, A.; Liu, Y.; Schultz, A.; Cardenas, M.; Li, X.; Babur, O.; Hsu, T.-K.; Lichtarge, O.; et al. Integrated analysis of TP53 gene and pathway alterations in the cancer genome atlas. *Cell Rep.* **2019**, *28*, 1370–1384. [[CrossRef](#)]
79. Zhang, Y.; Koneva, L.A.; Virani, S.; Arthur, A.E.; Virani, A.; Hall, P.B.; Warden, C.D.; Carey, T.E.; Chepeha, D.B.; Prince, M.E.; et al. Subtypes of HPV-positive head and neck cancers are associated with HPV characteristics, copy number alterations, PIK3CA mutation, and pathway signatures. *Clin. Cancer Res.* **2016**, *22*, 4735–4745. [[CrossRef](#)] [[PubMed](#)]
80. Cheng, H.; Yang, X.; Si, H.; Saleh, A.D.; Xiao, W.; Coupar, J.; Gollin, S.M.; Ferris, R.L.; Issaeva, N.; Yarbrough, W.G.; et al. Genomic and transcriptomic characterization links cell lines with aggressive head and neck cancers. *Cell Rep.* **2018**, *25*, 1332–1345. [[CrossRef](#)]
81. Marescalco, M.S.; Capizzi, C.; Condorelli, D.F.; Barresi, V. Genome-wide analysis of recurrent copy-number alterations and copy-neutral loss of heterozygosity in head and neck squamous cell carcinoma. *J. Oral Pathol. Med.* **2014**, *43*, 20–27. [[CrossRef](#)] [[PubMed](#)]
82. Davidson, M.A.; Shanks, E.J. 3q26-29 Amplification in head and neck squamous cell carcinoma: A review of established and prospective oncogenes. *FEBS J.* **2017**, *284*, 2705–2731. [[CrossRef](#)] [[PubMed](#)]
83. Gollin, S.M. Cytogenetic alterations and their molecular genetic correlates in head and neck squamous cell carcinoma: A next generation window to the biology of disease. *Genes Chromosom. Cancer* **2014**, *53*, 972–990. [[CrossRef](#)]
84. Van Harten, A.M.; Poell, J.B.; Buijze, M.; Brink, A.; Wells, S.I.; René Leemans, C.; Wolthuis, R.M.F.; Brakenhoff, R.H. Characterization of a head and neck cancer-derived cell line panel confirms the distinct TP53-proficient copy number-silent subclass. *Oral Oncol.* **2019**, *98*, 53–61. [[CrossRef](#)]
85. Taberna, M.; Mena, M.; Pavón, M.A.; Alemany, L.; Gillison, M.L.; Mesía, R. Human papillomavirus-related oropharyngeal cancer. *Ann. Oncol.* **2017**, *28*, 2386–2398. [[CrossRef](#)]
86. Wreesmann, V.B.; Wang, D.; Goberdhan, A.; Prasad, M.; Ngai, I.; Schnaser, E.A.; Sacks, P.G.; Singh, B. Genetic abnormalities associated with nodal metastasis in head and neck cancer. *Head Neck* **2004**, *26*, 10–15. [[CrossRef](#)]
87. Towle, R.; Tsui, I.F.L.; Zhu, Y.; MacLellan, S.; Poh, C.F.; Garnis, C. Recurring DNA copy number gain at chromosome 9p13 plays a role in the activation of multiple candidate oncogenes in progressing oral premalignant lesions. *Cancer Med.* **2014**, *3*, 1170–1184. [[CrossRef](#)] [[PubMed](#)]
88. Van Zeeburg, H.J.T.; Graveland, A.P.; Brink, A.; Nguyen, M.; Leemans, C.R.; Bloemena, E.; Braakhuis, B.J.M.; Brakenhoff, R.H. Generation of precursor cell lines from preneoplastic fields surrounding head and neck cancers. *Head Neck* **2013**, *35*, 568–574. [[CrossRef](#)] [[PubMed](#)]
89. Stransky, N.; Egloff, A.M.; Tward, A.D.; Kostic, A.D.; Cibulskis, K.; Sivachenko, A.; Kryukov, G.V.; Lawrence, M.S.; Sougnez, C.; McKenna, A.; et al. The mutational landscape of head and neck squamous cell carcinoma. *Science* **2011**, *333*, 1157–1160. [[CrossRef](#)] [[PubMed](#)]
90. De Roest, R.H.; Mes, S.; Brink, A.; Poell, J.B.; Van de Wiel, M.A.; Bloemena, E.; Thai, E.; Poli, T.; Leemans, C.R.; Brakenhoff, R.H. Molecular characterization of locally relapsed head and neck cancer after concomitant chemoradiotherapy. *Clin. Cancer Res.* **2019**, *25*, 7256–7265. [[CrossRef](#)] [[PubMed](#)]
91. Masica, D.L.; Li, S.; Douville, C.; Manola, J.; Ferris, R.L.; Burtness, B.; Forastiere, A.A.; Koch, W.M.; Chung, C.H.; Karchin, R. Predicting survival in head and neck squamous cell carcinoma from TP53 mutation. *Hum. Genet.* **2015**, *134*, 497–507. [[CrossRef](#)] [[PubMed](#)]
92. Poeta, M.L.; Forastiere, A.; Benoit, N.; Califano, J.A.; Westra, W.; Sidransky, D.; Koch, W.M.; Saunders, J.; Manola, J.; Goldwasser, M.A.; et al. TP53 mutations and survival in squamous-cell carcinoma of the head and neck. *N. Engl. J. Med.* **2007**, *357*, 2552–2561. [[CrossRef](#)] [[PubMed](#)]
93. Hanahan, D.; Weinberg, R.A. The hallmarks of cancer. *Cell* **2000**, *100*, 57–70. [[CrossRef](#)]
94. Fukusumi, T.; Califano, J.A. The NOTCH Pathway in head and neck squamous cell carcinoma. *J. Dent. Res.* **2018**, *97*, 645–653. [[CrossRef](#)]
95. Henderson, S.; Chakravarthy, A.; Su, X.; Boshoff, C.; Fenton, T.R. APOBEC-mediated cytosine deamination links PIK3CA helical domain mutations to human papillomavirus-driven tumor development. *Cell Rep.* **2014**, *7*, 1833–1841. [[CrossRef](#)]
96. Swanton, C.; Mcgranahan, N.; Starrett, G.J.; Harris, R.S. APOBEC Enzymes: Mutagenic fuel for cancer evolution and heterogeneity. *Cancer Discov.* **2015**, *5*, 704–712. [[CrossRef](#)]
97. Weinstein, I.B. CANCER: Enhanced: Addiction to oncogenes—The Achilles heel of cancer. *Science* **2002**, *297*, 63–64. [[CrossRef](#)] [[PubMed](#)]
98. Pagliarini, R.; Shao, W.; Sellers, W.R. Oncogene addiction: Pathways of therapeutic response, resistance, and road maps toward a cure. *EMBO Rep.* **2015**, *16*, 280–296. [[CrossRef](#)]
99. Chapman, P.B.; Hauschild, A.; Robert, C.; Haanen, J.B.; Ascierto, P.; Larkin, J.; Dummer, R.; Garbe, C.; Testori, A.; Maio, M.; et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N. Engl. J. Med.* **2011**, *364*, 2507–2516. [[CrossRef](#)]
100. Sosman, J.A.; Kim, K.B.; Schuchter, L.; Gonzalez, R.; Pavlick, A.C.; Weber, J.S.; McArthur, G.A.; Hutson, T.E.; Moschos, S.J.; Flaherty, K.T.; et al. Survival in BRAF V600-mutant advanced melanoma treated with vemurafenib. *N. Engl. J. Med.* **2012**, *366*, 707–714. [[CrossRef](#)] [[PubMed](#)]

101. Gilardi, M.; Wang, Z.; Proietto, M.; Chillà, A.; Calleja-Valera, J.L.; Goto, Y.; Vanoni, M.; Janes, M.R.; Mikulski, Z.; Gualberto, A.; et al. Tipifarnib as a precision therapy for HRAS-mutant head and neck squamous cell carcinomas. *Mol. Cancer Ther.* **2020**, *19*, 1784–1796. [[CrossRef](#)] [[PubMed](#)]
102. Lui, V.W.Y.; Hedberg, M.L.; Li, H.; Vangara, B.S.; Pendleton, K.; Zeng, Y.; Lu, Y.; Zhang, Q.; Du, Y.; Gilbert, B.R.; et al. Frequent mutation of the PI3K pathway in head and neck cancer defines predictive biomarkers. *Cancer Discov.* **2013**, *3*, 761–769. [[CrossRef](#)] [[PubMed](#)]
103. Saada-Bouزيد, E.; Le Tourneau, C. Beyond EGFR targeting in SCCHN: Angiogenesis, PI3K, and other molecular targets. *Front. Oncol.* **2019**, *9*, 74. [[CrossRef](#)]
104. Bryant, H.E.; Schultz, N.; Thomas, H.D.; Parker, K.M.; Flower, D.; Lopez, E.; Kyle, S.; Meuth, M.; Curtin, N.J.; Helleday, T. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature* **2005**, *434*, 913–917. [[CrossRef](#)]
105. Farmer, H.; McCabe, N.; Lord, C.J.; Tutt, A.N.J.; Johnson, D.A.; Richardson, T.B.; Santarosa, M.; Dillon, K.J.; Hickson, I.; Knights, C.; et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* **2005**, *434*, 917–921. [[CrossRef](#)]
106. Huang, M.; Feng, X.; Su, D.; Wang, G.; Wang, C.; Tang, M.; Paulucci-Holthausen, A.; Hart, T.; Chen, J. Genome-wide CRISPR screen uncovers a synergistic effect of combining Haspin and Aurora kinase B inhibition. *Oncogene* **2020**, *39*, 4312–4322. [[CrossRef](#)]
107. Lee, J.W.; Parameswaran, J.; Sandoval-Schaefer, T.; Eoh, K.J.; Yang, D.; Zhu, F.; Mehra, R.; Sharma, R.; Gaffney, S.G.; Perry, E.B.; et al. Combined aurora kinase A (AURKA) and WEE1 inhibition demonstrates synergistic antitumor effect in squamous cell carcinoma of the head and neck. *Clin. Cancer Res.* **2019**, *25*, 3430–3442. [[CrossRef](#)]
108. Marcu, L.; Van Doorn, T.; Olver, I. Cisplatin and radiotherapy in the treatment of locally advanced head and neck cancer—a review of their cooperation. *Acta Oncol.* **2003**, *42*, 315–325. [[CrossRef](#)]
109. De Boer, D.V.; Martens-de Kemp, S.R.; Buijze, M.; Stigter-van Walsum, M.; Bloemena, E.; Dietrich, R.; Leemans, C.R.; Beusechem Van, V.W.; Braakhuis, B.J.M.; Brakenhoff, R.H. Targeting PLK1 as a novel chemopreventive approach to eradicate preneoplastic mucosal changes in the head and neck. *Oncotarget* **2017**, *8*, 1–13. [[CrossRef](#)]
110. Van Harten, A.M.; De Boer, D.V.; Martens-de Kemp, S.R.; Buijze, M.; Ganzevles, S.H.; Hunter, K.D.; Leemans, C.R.; Van Beusechem, V.W.; Wolthuis, R.M.F.; De Menezes, R.X.; et al. Chemopreventive targeted treatment of head and neck precancer by Wee1 inhibition. *Sci. Rep.* **2020**, *10*, 1–12. [[CrossRef](#)]
111. Van Harten, A.M.; Buijze, M.; Van der Mast, R.; Rooimans, M.A.; Martens-de Kemp, S.R.; Bachas, C.; Brink, A.; Stigter-van Walsum, M.; Wolthuis, R.M.F.; Brakenhoff, R.H. Targeting the cell cycle in head and neck cancer by Chk1 inhibition: A novel concept of bimodal cell death. *Oncogenesis* **2019**, *8*, 1–16. [[CrossRef](#)]
112. Gadhikar, M.A.; Zhang, J.; Shen, L.; Rao, X.; Wang, J.; Zhao, M.; Kalu, N.N.; Johnson, F.M.; Byers, L.A.; Heymach, J.; et al. CDKN2A/p16 deletion in head and neck cancer cells is associated with cdk2 activation, replication stress, and vulnerability to CHK1 inhibition. *Cancer Res.* **2018**, *78*, 781–797. [[CrossRef](#)]
113. Muller, F.L.; Aquilanti, E.A.; DePinho, R.A. Collateral lethality: A new therapeutic strategy in oncology. *Trends Cancer* **2015**, *1*, 161–173. [[CrossRef](#)]
114. Menezes, R.X.; Boetzer, M.; Sieswerda, M.; Van Ommen, G.J.B.; Boer, J.M. Integrated analysis of DNA copy number and gene expression microarray data using gene sets. *BMC Bioinform.* **2009**, *10*, 203. [[CrossRef](#)]
115. Zhao, D.; DePinho, R.A. Synthetic essentiality: Targeting tumor suppressor deficiencies in cancer. *BioEssays* **2017**, *39*, 1700076. [[CrossRef](#)]
116. Elbashir, S.M.; Harborth, J.; Lendeckel, W.; Yalcin, A.; Weber, K.; Tuschl, T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* **2001**, *411*, 494–498. [[CrossRef](#)]
117. Brummelkamp, T.R.; Bernards, R.; Agami, R. A system for stable expression of short interfering RNAs in mammalian cells. *Science* **2002**, *296*, 550–553. [[CrossRef](#)]
118. Rao, D.D.; Senzer, N.; Cleary, M.A.; Nemunaitis, J. Comparative assessment of siRNA and shRNA off target effects: What is slowing clinical development. *Cancer Gene Ther.* **2009**, *16*, 807–809. [[CrossRef](#)]
119. Hinz, T.K.; Kleczko, E.K.; Singleton, K.R.; Calhoun, J.; Marek, L.A.; Kim, J.; Tan, A.C.; Heasley, L.E. Functional RNAi screens define distinct protein kinase vulnerabilities in EGFR-dependent HNSCC cell lines. *Mol. Pharmacol.* **2019**, *96*, 862–870. [[CrossRef](#)]
120. Fu, G.; Somasundaram, R.T.; Jessa, F.; Srivastava, G.; MacMillan, C.; Witterick, I.; Walfish, P.G.; Ralhan, R. ER maleate is a novel anticancer agent in oral cancer: Implications for cancer therapy. *Oncotarget* **2016**, *7*, 17162–17181. [[CrossRef](#)]
121. Moser, R.; Xu, C.; Kao, M.; Annis, J.; Lerma, L.A.; Schaupp, C.M.; Gurley, K.E.; Jang, I.S.; Biktasova, A.; Yarbrough, W.G.; et al. Functional kinomics identifies candidate therapeutic targets in head and neck cancer. *Clin. Cancer Res.* **2014**, *20*, 4274–4288. [[CrossRef](#)]
122. Yeh, M.-H.; Tsai, T.-C.; Kuo, H.-P.; Chang, N.-W.; Lee, M.-R.; Chung, J.-G.; Tsai, M.-H.; Liu, J.-Y.; Kao, M.-C. Lentiviral short hairpin RNA screen of human kinases and phosphatases to identify potential biomarkers in oral squamous cancer cells. *Int. J. Oncol.* **2011**, *39*, 1221–1231. [[CrossRef](#)]
123. Yamaguchi, K.; Iglesias-Bartolomé, R.; Wang, Z.; Callejas-Valera, J.L.; Amornphimoltham, P.; Molinolo, A.A.; Cohen, E.E.; Califano, J.A.; Lippman, S.M.; Luo, J.; et al. A synthetic-lethality RNAi screen reveals an ERK-mTOR co-targeting pro-apoptotic switch in PIK3CA + oral cancers. *Oncotarget* **2016**, *7*, 10696–10709. [[CrossRef](#)]
124. Boudreau, R.L.; Martins, I.; Davidson, B.L. Artificial MicroRNAs as siRNA shuttles: Improved safety as compared to shRNAs in vitro and in vivo. *Mol. Ther.* **2009**, *17*, 169–175. [[CrossRef](#)]

125. Bachas, C.; Hodzic, J.; Van der Mijn, J.C.; Stoepker, C.; Verheul, H.M.W.; Wolthuis, R.M.F.; Felley-Bosco, E.; Van Wieringen, W.N.; Van Beusechem, V.W.; Brakenhoff, R.H.; et al. Rscreenorm: Normalization of CRISPR and siRNA screen data for more reproducible hit selection. *BMC Bioinform.* **2018**, *19*, 301. [[CrossRef](#)]
126. Xu, C.; Nikolova, O.; Basom, R.S.; Mitchell, R.M.; Shaw, R.; Moser, R.D.; Park, H.; Gurley, K.E.; Kao, M.C.; Green, C.L.; et al. Functional precision medicine identifies novel druggable targets and therapeutic options in head and neck cancer. *Clin. Cancer Res.* **2018**, *24*, 2828–2843. [[CrossRef](#)]
127. Martens-de Kemp, S.R.; Nagel, R.; Stigter-van Walsum, M.; Van der Meulen, I.H.; Van Beusechem, V.W.; Braakhuis, B.J.M.; Brakenhoff, R.H. Functional genetic screens identify genes essential for tumor cell survival in head and neck and lung cancer. *Clin. Cancer Res.* **2013**, *19*, 1994–2003. [[CrossRef](#)]
128. Xu, C.; Wang, P.; Liu, Y.; Zhang, Y.; Fan, W.; Upton, M.P.; Lohavanichbutr, P.; Houck, J.R.; Doody, D.R.; Futran, N.D.; et al. Integrative genomics in combination with rna interference identifies prognostic and functionally relevant gene targets for oral squamous cell carcinoma. *PLoS Genet.* **2013**, *9*, e1003169. [[CrossRef](#)] [[PubMed](#)]
129. Martens-de Kemp, S.R.; Brink, A.; Van der Meulen, I.H.; De Menezes, R.X.; Te Beest, D.E.; Leemans, C.R.; Van Beusechem, V.W.; Braakhuis, B.J.M.; Brakenhoff, R.H. The FA/BRCA pathway identified as the major predictor of cisplatin response in head and neck cancer by functional genomics. *Mol. Cancer Ther.* **2017**, *16*, 540–550. [[CrossRef](#)] [[PubMed](#)]
130. Sangiorgi, B.; De Souza, F.C.; Mota de Souza Lima, I.; dos Santos Schiavinato, J.L.; Corveloni, A.C.; Thomé, C.H.; Araújo Silva, W.; Faça, V.M.; Covas, D.T.; Zago, M.A.; et al. A high-content screening approach to identify MicroRNAs against head and neck cancer cell survival and EMT in an inflammatory microenvironment. *Front. Oncol.* **2019**, *9*, 1100. [[CrossRef](#)] [[PubMed](#)]
131. Doench, J.G. Am I ready for CRISPR? A user's guide to genetic screens. *Nat. Rev. Genet.* **2018**, *19*, 67–80. [[CrossRef](#)]
132. Chai, A.W.Y.; Yee, P.S.; Price, S.; Yee, S.M.; Lee, H.M.; Tiong, V.K.H.; Gonçalves, E.; Behan, F.M.; Bateson, J.; Gilbert, J.; et al. Genome-wide CRISPR screens of oral squamous cell carcinoma reveal fitness genes in the Hippo pathway. *Elife* **2020**, *9*, 1–81. [[CrossRef](#)]
133. Dok, R.; Bamps, M.; Glorieux, M.; Zhao, P.; Sablina, A.; Nuyts, S. Radiosensitization approaches for HPV-positive and HPV-negative head and neck squamous carcinomas. *Int. J. Cancer* **2020**, *146*, 1075–1085. [[CrossRef](#)]
134. Abraham, C.G.; Ludwig, M.P.; Andrysik, Z.; Pandey, A.; Joshi, M.; Galbraith, M.D.; Sullivan, K.D.; Espinosa, J.M. Δ Np63 α suppresses TGFB2 expression and RHOA activity to drive cell proliferation in squamous cell carcinomas. *Cell Rep.* **2018**, *24*, 3224–3236. [[CrossRef](#)]
135. Jamieson, S.M.; Tsai, P.; Kondratyev, M.K.; Budhani, P.; Liu, A.; Senzer, N.N.; Chiorean, E.G.; Jalal, S.I.; Nemunaitis, J.J.; Kee, D.; et al. Evofosfamide for the treatment of human papillomavirus-negative head and neck squamous cell carcinoma. *JCI Insight* **2018**, *3*. [[CrossRef](#)]
136. Meyers, R.M.; Bryan, J.G.; McFarland, J.M.; Weir, B.A.; Sizemore, A.E.; Xu, H.; Dharia, N.V.; Montgomery, P.G.; Cowley, G.S.; Pantel, S.; et al. Computational correction of copy number effect improves specificity of CRISPR–Cas9 essentiality screens in cancer cells. *Nat. Genet.* **2017**, *49*, 1779–1784. [[CrossRef](#)]
137. Behan, F.M.; Iorio, F.; Picco, G.; Gonçalves, E.; Beaver, C.M.; Migliardi, G.; Santos, R.; Rao, Y.; Sassi, F.; Pinnelli, M.; et al. Prioritization of cancer therapeutic targets using CRISPR–Cas9 screens. *Nature* **2019**, *568*, 511–516. [[CrossRef](#)]
138. Lenoir, W.F.; Lim, T.L.; Hart, T. PICKLES: The database of pooled in-vitro CRISPR knockout library essentiality screens. *Nucleic Acids Res.* **2018**, *46*, D776–D780. [[CrossRef](#)]
139. Perry, J.; Ashford, B.; Thind, A.S.; Gauthier, M.-E.; Minaei, E.; Major, G.; Iyer, N.G.; Gupta, R.; Clark, J.; Ranson, M. Comprehensive mutational and phenotypic characterization of new metastatic cutaneous squamous cell carcinoma cell lines reveal novel drug susceptibilities. *Int. J. Mol. Sci.* **2020**, *21*, 9536. [[CrossRef](#)]
140. Bryant, J.; Batis, N.; Franke, A.C.; Clancey, G.; Hartley, M.; Ryan, G.; Brooks, J.; Southam, A.D.; Barnes, N.; Parish, J.; et al. Repurposed quinacrine synergizes with cisplatin, reducing the effective dose required for treatment of head and neck squamous cell carcinoma. *Oncotarget* **2019**, *10*, 5229–5244. [[CrossRef](#)]
141. Okazaki, S.; Shintani, S.; Hirata, Y.; Suina, K.; Semba, T.; Yamasaki, J.; Umene, K.; Ishikawa, M.; Saya, H.; Nagano, O. Synthetic lethality of the ALDH3A1 inhibitor dyclonine and xCT inhibitors in glutathione deficiency-resistant cancer cells. *Oncotarget* **2018**, *9*, 33832–33843. [[CrossRef](#)]
142. Chia, S.; Low, J.-L.; Zhang, X.; Kwang, X.-L.; Chong, F.-T.; Sharma, A.; Bertrand, D.; Toh, S.Y.; Leong, H.-S.; Thangavelu, M.T.; et al. Phenotype-driven precision oncology as a guide for clinical decisions one patient at a time. *Nat. Commun.* **2017**, *8*, 435. [[CrossRef](#)] [[PubMed](#)]
143. Johnston, P.A.; Sen, M.; Hua, Y.; Camarco, D.P.; Shun, T.Y.; Lazo, J.S.; Wilson, G.M.; Resnick, L.O.; LaPorte, M.G.; Wipf, P.; et al. HCS campaign to identify selective inhibitors of IL-6-induced STAT3 pathway activation in head and neck cancer cell lines. *Assay Drug Dev. Technol.* **2015**, *13*, 356–376. [[CrossRef](#)]
144. Srivastava, G.; Matta, A.; Fu, G.; Somasundaram, R.T.; Datti, A.; Walfish, P.G.; Ralhan, R. Anticancer activity of pyrithione zinc in oral cancer cells identified in small molecule screens and xenograft model: Implications for oral cancer therapy. *Mol. Oncol.* **2015**, *9*, 1720–1735. [[CrossRef](#)]
145. Yang, W.; Soares, J.; Greninger, P.; Edelman, E.J.; Lightfoot, H.; Forbes, S.; Bindal, N.; Beare, D.; Smith, J.A.; Thompson, I.R.; et al. Genomics of drug sensitivity in cancer (GDSC): A resource for therapeutic biomarker discovery in cancer cells. *Nucleic Acids Res.* **2013**, *41*, D955–D961. [[CrossRef](#)]

146. Uhlen, M.; Fagerberg, L.; Hallstrom, B.M.; Lindskog, C.; Oksvold, P.; Mardinoglu, A.; Sivertsson, A.; Kampf, C.; Sjostedt, E.; Asplund, A.; et al. Tissue-based map of the human proteome. *Science* **2015**, *347*, 1260419. [[CrossRef](#)] [[PubMed](#)]
147. Subramanian, A.; Narayan, R.; Corsello, S.M.; Peck, D.D.; Natoli, T.E.; Lu, X.; Gould, J.; Davis, J.F.; Tubelli, A.A.; Asiedu, J.K.; et al. A next generation connectivity map: L1000 platform and the first 1,000,000 profiles. *Cell* **2017**, *171*, 1437–1452. [[CrossRef](#)]
148. Paananen, J.; Fortino, V. An omics perspective on drug target discovery platforms. *Brief. Bioinform.* **2020**, *21*, 1937–1953. [[CrossRef](#)] [[PubMed](#)]
149. Keam, B.; Park, J.-Y.; Kim, J.-P.; Kim, G.-D.; Yu, Y.-S.; Cho, S.-H.; Kim, S.; Ahn, H.-K.; Chun, S.-H.; Kwon, J.-H.; et al. Comprehensive analysis of mutation-based and expressed genes-based pathways in head and neck squamous cell carcinoma. *Processes* **2021**, *9*, 792. [[CrossRef](#)]
150. Ran, Q.-C.; Long, S.-R.; Ye, Y.; Xie, C.; XuXiao, Z.-L.; Liu, Y.-S.; Pang, H.-X.; Sunchuri, D.; Teng, N.-C.; Guo, Z.-L. Mining TCGA database for prognostic genes in head and neck squamous cell carcinoma microenvironment. *J. Dent. Sci.* **2021**, *16*, 661–667. [[CrossRef](#)] [[PubMed](#)]
151. Huang, C.; Chen, L.; Savage, S.R.; Eiguez, R.V.; Dou, Y.; Li, Y.; da Veiga Leprevost, F.; Jaehnig, E.J.; Lei, J.T.; Wen, B.; et al. Proteogenomic insights into the biology and treatment of HPV-negative head and neck squamous cell carcinoma. *Cancer Cell* **2021**, *39*, 1–19. [[CrossRef](#)]
152. Cavalieri, S.; De Cecco, L.; Brakenhoff, R.H.; Serafini, M.S.; Canevari, S.; Rossi, S.; Lanfranco, D.; Hoebbers, F.J.P.; Wesseling, F.W.R.; Keek, S.; et al. Development of a multiomics database for personalized prognostic forecasting in head and neck cancer: The Big Data to Decide EU Project. *Head Neck* **2021**, *43*, 601–612. [[CrossRef](#)]
153. Willems, S.M.; Abeln, S.; Feenstra, K.A.; De Bree, R.; Van der Poel, E.F.; Baatenburg de Jong, R.J.; Heringa, J.; Van den Brekel, M.W.M. The potential use of big data in oncology. *Oral Oncol.* **2019**, *98*, 8–12. [[CrossRef](#)]
154. Hume, S.; Dianov, G.L.; Ramadan, K. A unified model for the G1/S cell cycle transition. *Nucleic Acids Res.* **2020**, *48*, 12483–12501. [[CrossRef](#)]
155. Otto, T.; Sicinski, P. Cell cycle proteins as promising targets in cancer therapy. *Nat. Rev. Cancer* **2017**, *17*, 93–115. [[CrossRef](#)]
156. Rubin, S.M.; Sage, J.; Skotheim, J.M. Integrating old and new paradigms of G1/S control. *Mol. Cell* **2020**, *80*, 183–192. [[CrossRef](#)] [[PubMed](#)]
157. Sherr, C.J. Living with or without cyclins and cyclin-dependent kinases. *Genes Dev.* **2004**, *18*, 2699–2711. [[CrossRef](#)] [[PubMed](#)]
158. Shadfan, M.; Lopez-Pajares, V.; Yuan, Z.-M. MDM2 and MDMX: Alone and together in regulation of p53. *Transl. Cancer Res.* **2012**, *1*, 88–89. [[CrossRef](#)]
159. Tian, X.; Huang, B.; Zhang, X.P.; Lu, M.; Liu, F.; Onuchic, J.N.; Wang, W. Modeling the response of a tumor-suppressive network to mitogenic and oncogenic signals. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 5337–5342. [[CrossRef](#)] [[PubMed](#)]
160. Gil, J.; Peters, G. Regulation of the INK4b-ARF-INK4a tumour suppressor locus: All for one or one for all. *Nat. Rev. Mol. Cell Biol.* **2006**, *7*, 667–677. [[CrossRef](#)]
161. Giono, L.E.; Manfredi, J.J. The p53 tumor suppressor participates in multiple cell cycle checkpoints. *J. Cell. Physiol.* **2006**, *209*, 13–20. [[CrossRef](#)] [[PubMed](#)]
162. Smith, J.; Tho, L.M.; Xu, N.; Gillespie, D.A. The ATM-Chk2 and ATR-Chk1 pathways in DNA damage signaling and cancer. *Adv. Cancer Res.* **2010**, *108*, 73–112. [[CrossRef](#)]
163. Shiloh, Y.; Ziv, Y. The ATM protein kinase: Regulating the cellular response to genotoxic stress, and more. *Nat. Rev. Mol. Cell Biol.* **2013**, *14*, 197–210. [[CrossRef](#)]
164. Blackford, A.N.; Jackson, S.P. ATM, ATR, and DNA-PK: The trinity at the heart of the DNA damage response. *Mol. Cell* **2017**, *66*, 801–817. [[CrossRef](#)] [[PubMed](#)]
165. Reinhardt, H.C.; Schumacher, B. The p53 network: Cellular and systemic DNA damage responses in aging and cancer. *Trends Genet.* **2012**, *28*, 128–136. [[CrossRef](#)] [[PubMed](#)]
166. Sherr, C.J.; McCormick, F. The RB and p53 pathways in cancer. *Cancer Cell* **2002**, *2*, 103–112. [[CrossRef](#)]
167. Polager, S.; Ginsberg, D. P53 and E2f: Partners in life and death. *Nat. Rev. Cancer* **2009**, *9*, 738–748. [[CrossRef](#)] [[PubMed](#)]
168. Hydbring, P.; Malumbres, M.; Sicinski, P. Non-canonical functions of cell cycle cyclins and cyclin-dependent kinases. *Nat. Rev. Mol. Cell Biol.* **2016**, *17*, 280–292. [[CrossRef](#)]
169. Massagué, J. G1 cell-cycle control and cancer. *Nature* **2004**, *432*, 298–306. [[CrossRef](#)]
170. Benedict, B.; Van Harn, T.; Dekker, M.; Hermsen, S.; Kucukosmanoglu, A.; Pieters, W.; Delzenne-Goette, E.; Dorsman, J.C.; Petermann, E.; Foijer, F.; et al. Loss of p53 suppresses replication-stress-induced DNA breakage in G1/S checkpoint deficient cells. *Elife* **2018**, *7*. [[CrossRef](#)]
171. Dhingra, V.; Verma, J.; Misra, V.; Srivastav, S.; Hasan, F. Evaluation of cyclin D1 expression in head and neck squamous cell carcinoma. *J. Clin. Diagn. Res.* **2017**, *11*, EC01–EC04. [[CrossRef](#)]
172. Musgrove, E.; Caldon, C.; Barraclough, J.; Stone, A.; Sutherland, R. Cyclin D as a therapeutic target in cancer. *Nat. Rev. Cancer* **2011**, *11*, 558–572. [[CrossRef](#)]
173. Williams, G.H.; Stoerber, K. The cell cycle and cancer. *J. Pathol.* **2012**, *226*, 352–364. [[CrossRef](#)]
174. Kitao, H.; Iimori, M.; Kataoka, Y.; Wakasa, T.; Tokunaga, E.; Saeki, H.; Oki, E.; Maehara, Y. DNA replication stress and cancer chemotherapy. *Cancer Sci.* **2018**, *109*, 264–271. [[CrossRef](#)] [[PubMed](#)]
175. Mody, M.D.; Gill, H.S.; Saba, N.F. The evolving and future role of taxanes in squamous cell carcinomas of the head and neck a review. *JAMA Otolaryngol. Head Neck Surg.* **2016**, *142*, 898–905. [[CrossRef](#)]

176. Yap, T.A.; Plummer, R.; Azad, N.S.; Helleday, T. The DNA damaging revolution: PARP inhibitors and beyond. *Am. Soc. Clin. Oncol. Educ. B.* **2019**, *39*, 185–195. [[CrossRef](#)] [[PubMed](#)]
177. Gorecki, L.; Andrs, M.; Korabecny, J. Clinical candidates targeting the ATR–CHK1–WEE1 axis in cancer. *Cancers* **2021**, *13*, 795. [[CrossRef](#)]
178. Cleary, J.M.; Aguirre, A.J.; Shapiro, G.I.; D’Andrea, A.D. Biomarker-guided development of DNA repair inhibitors. *Mol. Cell* **2020**, *78*, 1070–1085. [[CrossRef](#)]
179. Ingham, M.; Schwartz, G.K. Cell-cycle therapeutics come of age. *J. Clin. Oncol.* **2017**, *35*, 2949–2959. [[CrossRef](#)] [[PubMed](#)]
180. Forment, J.V.; O’Connor, M.J. Targeting the replication stress response in cancer. *Pharmacol. Ther.* **2018**, *188*, 155–167. [[CrossRef](#)]
181. Ubhi, T.; Brown, G.W. Exploiting DNA replication stress for cancer treatment. *Cancer Res.* **2019**, *79*, 1730–1739. [[CrossRef](#)]
182. Bartek, J.; Mistrik, M.; Bartkova, J. Thresholds of replication stress signaling in cancer development and treatment. *Nat. Struct. Mol. Biol.* **2012**, *19*, 5–7. [[CrossRef](#)]
183. Dent, P. Investigational CHK1 inhibitors in early phase clinical trials for the treatment of cancer. *Expert Opin. Investig. Drugs* **2019**, *28*, 1095–1100. [[CrossRef](#)]
184. Dobbstein, M.; Sørensen, C.S. Exploiting replicative stress to treat cancer. *Nat. Rev. Drug Discov.* **2015**, *14*, 405–423. [[CrossRef](#)]
185. Dominguez-Brauer, C.; Thu, K.L.; Mason, J.M.; Blaser, H.; Bray, M.R.; Mak, T.W. Targeting mitosis in cancer: Emerging strategies. *Mol. Cell* **2015**, *60*, 524–536. [[CrossRef](#)]
186. Geenen, J.J.J.; Schellens, J.H.M. Molecular pathways: Targeting the protein kinase Wee1 in cancer. *Clin. Cancer Res.* **2017**, *23*, 4540–4544. [[CrossRef](#)] [[PubMed](#)]
187. Lindenbergh-van der Plas, M.; Martens-de Kemp, S.R.; De Maaker, M.; Van Wieringen, W.N.; Ylstra, B.; Agami, R.; Cerisoli, F.; Leemans, C.R.; Braakhuis, B.J.M.; Brakenhoff, R.H. Identification of lethal microRNAs specific for head and neck cancer. *Clin. Cancer Res.* **2013**, *19*, 5647–5657. [[CrossRef](#)] [[PubMed](#)]
188. Kalu, N.N.; Mazumdar, T.; Peng, S.; Tong, P.; Shen, L.; Wang, J.; Banerjee, U.; Myers, J.N.; Pickering, C.R.; Brunell, D.; et al. Comprehensive pharmacogenomic profiling of human papillomavirus-positive and -negative squamous cell carcinoma identifies sensitivity to aurora kinase inhibition in KMT2D mutants. *Cancer Lett.* **2018**, *431*, 64–72. [[CrossRef](#)] [[PubMed](#)]
189. Zou, J.; Qiao, X.; Ye, H.; Yang, Y.; Zheng, X.; Zhao, H.; Liu, S. Antisense inhibition of ATM gene enhances the radiosensitivity of head and neck squamous cell carcinoma in mice. *J. Exp. Clin. Cancer Res.* **2008**, *27*, 56. [[CrossRef](#)] [[PubMed](#)]
190. Zou, J.; Qiao, X.; Ye, H.; Zhang, Y.; Xian, J.; Zhao, H.; Liu, S. Inhibition of ataxia-telangiectasia mutated by antisense oligonucleotide nanoparticles induces radiosensitization of head and neck squamous-cell carcinoma in mice. *Cancer Biother. Radiopharm.* **2009**, *24*, 339–346. [[CrossRef](#)] [[PubMed](#)]
191. Feng, J.; Zou, J.; Li, L.; Zhao, Y.; Liu, S. Antisense oligodeoxynucleotides targeting ATM strengthen apoptosis of laryngeal squamous cell carcinoma grown in nude mice. *J. Exp. Clin. Cancer Res.* **2011**, *30*, 43. [[CrossRef](#)] [[PubMed](#)]
192. Weber, A.M.; Ryan, A.J. ATM and ATR as therapeutic targets in cancer. *Pharmacol. Ther.* **2015**, *149*, 124–138. [[CrossRef](#)]
193. Sankunni, M.; Parikh, R.A.; Lewis, D.W.; Gooding, W.E.; Saunders, W.S.; Gollin, S.M. Targeted inhibition of ATR or CHEK1 reverses radioresistance in oral squamous cell carcinoma with distal chromosome arm 11q loss. *Genes Chromosom. Cancer* **2014**, *53*, 129–143. [[CrossRef](#)]
194. Leonard, B.C.; Lee, E.D.; Bholra, N.E.; Li, H.; Sogaard, K.K.; Bakkenist, C.J.; Grandis, J.R.; Johnson, D.E. ATR inhibition sensitizes HPV– and HPV+ head and neck squamous cell carcinoma to cisplatin. *Oral Oncol.* **2019**, *95*, 35–42. [[CrossRef](#)]
195. Tang, P.A.; Siu, L.L.; Chen, E.X.; Hotte, S.J.; Chia, S.; Schwarz, J.K.; Pond, G.R.; Johnson, C.; Colevas, A.D.; Synold, T.W.; et al. Phase II study of ispinesib in recurrent or metastatic squamous cell carcinoma of the head and neck. *Investig. N. Drugs* **2008**, *26*, 257–264. [[CrossRef](#)]
196. Gerster, K.; Shi, W.; Ng, B.; Yue, S.; Ito, E.; Waldron, J.; Gilbert, R.; Liu, F.F. Targeting Polo-like kinase 1 enhances radiation efficacy for head-and-neck squamous cell carcinoma. *Int. J. Radiat. Oncol. Biol. Phys.* **2010**, *77*, 253–260. [[CrossRef](#)]
197. Wagenblast, J.; Hirsh, D.; Thron, L.; Arnoldner, C.; Diensthuber, M.; Stöver, T.; Hambek, M. Effects of the Polo-like-kinase-1-inhibitor BI2536 in squamous cell carcinoma cell lines of the head and neck. *Oncol. Lett.* **2012**, *4*, 175–177. [[CrossRef](#)] [[PubMed](#)]
198. Zhang, M.; Singh, R.; Peng, S.; Mazumdar, T.; Sambandam, V.; Shen, L.; Tong, P.; Li, L.; Kalu, N.N.; Pickering, C.R.; et al. Mutations of the LIM protein AJUBA mediate sensitivity of head and neck squamous cell carcinoma to treatment with cell-cycle inhibitors. *Cancer Lett.* **2017**, *392*, 71–82. [[CrossRef](#)] [[PubMed](#)]
199. Mazumdar, A.; Henderson, Y.C.; El-Naggar, A.K.; Sen, S.; Clayman, G.L. Aurora kinase A inhibition and paclitaxel as targeted combination therapy for head and neck squamous cell carcinoma. *Head Neck* **2009**, *31*, 625–634. [[CrossRef](#)]
200. Hoellein, A.; Pickhard, A.; von Keitz, F.; Schoeffmann, S.; Piontek, G.; Rudelius, M.; Baumgart, A.; Wagenpfeil, S.; Peschel, C.; Dechow, T.; et al. Aurora kinase inhibition overcomes cetuximab resistance in squamous cell cancer of the head and neck. *Oncotarget* **2011**, *2*, 599–609. [[CrossRef](#)]
201. Shaikh, M.H.; Idris, A.; Johnson, N.W.; Fallaha, S.; Clarke, D.T.W.; Martin, D.; Morgan, I.M.; Gabrielli, B.; McMillan, N.A.J. Aurora kinases are a novel therapeutic target for HPV-positive head and neck cancers. *Oral Oncol.* **2018**, *86*, 105–112. [[CrossRef](#)]
202. Jin, S.; Ma, H.; Yang, W.; Ju, H.; Wang, L.; Zhang, Z. Cell division cycle 7 is a potential therapeutic target in oral squamous cell carcinoma and is regulated by E2F1. *J. Mol. Med.* **2018**, *96*, 513–525. [[CrossRef](#)] [[PubMed](#)]
203. Chen, W.; Yuan, K.; Tao, Z.-Z.; Xiao, B.-K. Deletion of Forkhead Box M1 transcription factor reduces malignancy in laryngeal squamous carcinoma cells. *Asian Pac. J. Cancer Prev.* **2011**, *12*, 1785–1788.

204. Jiang, L.; Wu, X.; Wang, P.; Wen, T.; Yu, C.; Wei, L.; Chen, H. Targeting FoxM1 by thiostrepton inhibits growth and induces apoptosis of laryngeal squamous cell carcinoma. *J. Cancer Res. Clin. Oncol.* **2015**, *141*, 971–981. [[CrossRef](#)] [[PubMed](#)]
205. Labib, K. How do Cdc7 and cyclin-dependent kinases trigger the initiation of chromosome replication in eukaryotic cells? *Genes Dev.* **2010**, *24*, 1208–1219. [[CrossRef](#)] [[PubMed](#)]
206. Michel, L.; Ley, J.; Wildes, T.M.; Schaffer, A.; Robinson, A.; Chun, S.-E.; Lee, W.; Lewis, J.; Trinkaus, K.; Adkins, D. Phase I trial of palbociclib, a selective cyclin dependent kinase 4/6 inhibitor, in combination with cetuximab in patients with recurrent/metastatic head and neck squamous cell carcinoma. *Oral Oncol.* **2016**, *58*, 41–48. [[CrossRef](#)] [[PubMed](#)]
207. Ku, B.M.; Yi, S.Y.; Koh, J.; Bae, Y.H.; Sun, J.M.; Lee, S.H.; Ahn, J.S.; Park, K.; Ahn, M.J. The CDK4/6 inhibitor LY2835219 has potent activity in combination with mTOR inhibitor in head and neck squamous cell carcinoma. *Oncotarget* **2016**, *7*, 14803–14813. [[CrossRef](#)] [[PubMed](#)]
208. Tai, T.S.; Lin, P.M.; Wu, C.F.; Hung, S.K.; Huang, C.I.; Wang, C.C.; Su, Y.C. CDK4/6 inhibitor LEE011 is a potential radiation-sensitizer in head and neck squamous cell carcinoma: An in vitro study. *Anticancer Res.* **2019**, *39*, 713–720. [[CrossRef](#)]
209. Deneka, A.Y.; Einarson, M.B.; Bennett, J.; Nikonova, A.S.; Elmekawy, M.; Zhou, Y.; Lee, J.W.; Burtness, B.A.; Golemis, E.A. Synthetic lethal targeting of mitotic checkpoints in HPV-negative head and neck cancer. *Cancers* **2020**, *12*, 306. [[CrossRef](#)]
210. Vitti, E.T.; Kacperek, A.; Parsons, J.L. Targeting DNA Double-Strand Break Repair Enhances Radiosensitivity of HPV-Positive and HPV-Negative Head and Neck Squamous Cell Carcinoma to Photons and Protons. *Cancers* **2020**, *12*, 1490. [[CrossRef](#)]
211. Adkins, D.; Ley, J.; Neupane, P.; Worden, F.; Sacco, A.G.; Palka, K.; Grilley-Olson, J.E.; Maggiore, R.; Salama, N.N.; Trinkaus, K.; et al. Palbociclib and cetuximab in platinum-resistant and in cetuximab-resistant human papillomavirus-unrelated head and neck cancer: A multicentre, multigroup, phase 2 trial. *Lancet Oncol.* **2019**, *20*, 1295–1305. [[CrossRef](#)]
212. Robinson, A.M.; Rathore, R.; Redlich, N.J.; Adkins, D.R.; VanArsdale, T.; Van Tine, B.A.; Michel, L.S. Cisplatin exposure causes c-Myc-dependent resistance to CDK4/6 inhibition in HPV-negative head and neck squamous cell carcinoma. *Cell Death Dis.* **2019**, *10*, 867. [[CrossRef](#)] [[PubMed](#)]
213. Hu, Q.; Peng, J.; Jiang, L.; Li, W.; Su, Q.; Zhang, J.; Li, H.; Song, M.; Cheng, B.; Xia, J.; et al. Metformin as a senostatic drug enhances the anticancer efficacy of CDK4/6 inhibitor in head and neck squamous cell carcinoma. *Cell Death Dis.* **2020**, *11*, 925. [[CrossRef](#)]
214. Gadsden, N.J.; Fulcher, C.D.; Li, D.; Shrivastava, N.; Thomas, C.; Segall, J.E.; Prystowsky, M.B.; Schlecht, N.F.; Gavathiotis, E.; Ow, T.J. Palbociclib renders human papilloma virus-negative head and neck squamous cell carcinoma vulnerable to the senolytic agent navitoclax. *Mol. Cancer Res.* **2021**, *19*, 862–873. [[CrossRef](#)]
215. Busch, C.-J.; Kriegs, M.; Laban, S.; Tribius, S.; Knecht, R.; Petersen, C.; Dikomey, E.; Rieckmann, T. HPV-positive HNSCC cell lines but not primary human fibroblasts are radiosensitized by the inhibition of Chk1. *Radiother. Oncol.* **2013**, *108*, 495–499. [[CrossRef](#)] [[PubMed](#)]
216. Gadhikar, M.A.; Sciuto, M.R.; Alves, M.V.O.; Pickering, C.R.; Osman, A.A.; Neskey, D.M.; Zhao, M.; Fitzgerald, A.L.; Myers, J.N.; Frederick, M.J. Chk1/2 inhibition overcomes the cisplatin resistance of head and neck cancer cells secondary to the loss of functional p53. *Mol. Cancer Ther.* **2013**, *12*, 1860–1873. [[CrossRef](#)]
217. Bridges, K.A.; Chen, X.; Liu, H.; Rock, C.; Buchholz, T.A.; Shumway, S.D.; Skinner, H.D.; Meyn, R.E. MK-8776, a novel chk1 kinase inhibitor, radiosensitizes p53-defective human tumor cells. *Oncotarget* **2016**, *7*, 71660–71672. [[CrossRef](#)]
218. Barker, H.E.; Patel, R.; McLaughlin, M.; Schick, U.; Zaidi, S.; Nutting, C.M.; Newbold, K.L.; Bhide, S.; Harrington, K.J. CHK1 inhibition radiosensitizes head and neck cancers to paclitaxel-based chemoradiotherapy. *Mol. Cancer Ther.* **2016**, *15*, 2042–2054. [[CrossRef](#)] [[PubMed](#)]
219. Hong, D.; Infante, J.; Janku, F.; Jones, S.; Nguyen, L.M.; Burris, H.A.; Naing, A.; Bauer, T.M.; Piha-Paul, S.; Johnson, F.M.; et al. Phase I Study of LY2606368, a checkpoint kinase 1 inhibitor, in patients with advanced cancer. *J. Clin. Oncol.* **2016**, *34*, 1764–1771. [[CrossRef](#)] [[PubMed](#)]
220. Busch, C.-J.J.; Kröger, M.S.; Jensen, J.; Kriegs, M.; Gatzemeier, F.; Petersen, C.; Münscher, A.; Rothkamm, K.; Rieckmann, T. G2-checkpoint targeting and radiosensitization of HPV/p16-positive HNSCC cells through the inhibition of Chk1 and Wee1. *Radiother. Oncol.* **2017**, *122*, 260–266. [[CrossRef](#)] [[PubMed](#)]
221. Zeng, L.; Beggs, R.R.; Cooper, T.S.; Weaver, A.N.; Yang, E.S. Combining Chk1/2 inhibition with cetuximab and radiation enhances in vitro and in vivo cytotoxicity in head and neck squamous cell carcinoma. *Mol. Cancer Ther.* **2017**, *16*, 591–600. [[CrossRef](#)]
222. Hong, D.S.; Moore, K.; Patel, M.; Grant, S.C.; Burris, H.A.; William, W.N.; Jones, S.; Meric-Bernstam, F.; Infante, J.; Golden, L.; et al. Evaluation of prexasertib, a checkpoint kinase 1 inhibitor, in a phase Ib Study of patients with squamous cell carcinoma. *Clin. Cancer Res.* **2018**, *24*, 3263–3272. [[CrossRef](#)] [[PubMed](#)]
223. Yang, C.-Y.; Liu, C.-R.; Chang, I.Y.-F.; OuYang, C.-N.; Hsieh, C.-H.; Huang, Y.-L.; Wang, C.-I.; Jan, F.-W.; Wang, W.-L.; Tsai, T.-L.; et al. Cotargeting CHK1 and PI3K Synergistically suppresses tumor growth of oral cavity squamous cell carcinoma in patient-derived xenografts. *Cancers* **2020**, *12*, 1726. [[CrossRef](#)]
224. Molkenline, J.M.; Molkenline, D.P.; Bridges, K.A.; Xie, T.; Yang, L.; Sheth, A.; Heffernan, T.P.; Clump, D.A.; Faust, A.Z.; Ferris, R.L.; et al. Targeting DNA damage response in head and neck cancers through abrogation of cell cycle checkpoints. *Int. J. Radiat. Biol.* **2020**, *2020*, 1–8. [[CrossRef](#)] [[PubMed](#)]
225. Zeng, L.; Nikolaev, A.; Xing, C.; Della Manna, D.L.; Yang, E.S. CHK1/2 inhibitor prexasertib suppresses NOTCH signaling and enhances cytotoxicity of cisplatin and radiation in head and neck squamous cell carcinoma. *Mol. Cancer Ther.* **2020**, *19*, 1279–1288. [[CrossRef](#)] [[PubMed](#)]

226. Chaudhary, R.; Slebos, R.J.C.; Song, F.; McCleary-Sharpe, K.P.; Masannat, J.; Tan, A.C.; Wang, X.; Amaladas, N.; Wu, W.; Hall, G.E.; et al. Effects of checkpoint kinase 1 inhibition by prexasertib on the tumor immune microenvironment of head and neck squamous cell carcinoma. *Mol. Carcinog.* **2021**, *60*, 138–150. [[CrossRef](#)]
227. Lee, T.W.; Wong, W.W.; Dickson, B.D.; Lipert, B.; Cheng, G.J.; Hunter, F.W.; Hay, M.P.; Wilson, W.R. Radiosensitization of head and neck squamous cell carcinoma lines by DNA-PK inhibitors is more effective than PARP-1 inhibition and is enhanced by SLFN11 and hypoxia. *Int. J. Radiat. Biol.* **2019**, *95*, 1597–1612. [[CrossRef](#)] [[PubMed](#)]
228. Zeng, L.; Boggs, D.H.; Xing, C.; Zhang, Z.; Anderson, J.C.; Wajapeyee, N.; Veale, C.; Bredel, M.; Shi, L.Z.; Bonner, J.A.; et al. Combining PARP and DNA-PK inhibitors with irradiation inhibits HPV-negative head and neck cancer squamous carcinoma growth. *Front. Genet.* **2020**, *11*, 1036. [[CrossRef](#)]
229. Osman, A.A.; Monroe, M.M.; Ortega Alves, M.V.; Patel, A.A.; Katsonis, P.; Fitzgerald, A.L.; Neskey, D.M.; Frederick, M.J.; Woo, S.H.; Caulin, C.; et al. Wee-1 kinase inhibition overcomes cisplatin resistance associated with high-risk TP53 mutations in head and neck cancer through mitotic arrest followed by senescence. *Mol. Cancer Ther.* **2015**, *14*, 608–619. [[CrossRef](#)] [[PubMed](#)]
230. Tanaka, N.; Patel, A.A.; Wang, J.; Frederick, M.J.; Kalu, N.N.; Zhao, M.; Fitzgerald, A.L.; Xie, T.X.; Silver, N.L.; Caulin, C.; et al. Wee-1 kinase inhibition sensitizes high-risk HPV+HNSCC to apoptosis accompanied by downregulation of MCL-1 and XIAP antiapoptotic proteins. *Clin. Cancer Res.* **2015**, *21*, 4831–4844. [[CrossRef](#)]
231. Tanaka, N.; Patel, A.A.; Tang, L.; Silver, N.L.; Lindemann, A.; Takahashi, H.; Jaksik, R.; Rao, X.; Kalu, N.N.; Chen, T.C.; et al. Replication stress leading to apoptosis within the S-phase contributes to synergism between vorinostat and AZD1775 in HNSCC harboring high-risk TP53 mutation. *Clin. Cancer Res.* **2017**, *23*, 6541–6554. [[CrossRef](#)]
232. Kao, M.; Green, C.; Sidorova, J.; Méndez, E. Strategies for targeted therapy in head and neck squamous cell carcinoma using WEE1 inhibitor AZD1775. *JAMA Otolaryngol. Neck Surg.* **2017**, *143*, 631. [[CrossRef](#)]
233. Yuan, M.-L.; Li, P.; Xing, Z.-H.; Di, J.-M.; Liu, H.; Yang, A.-K.; Lin, X.-J.; Jiang, Q.-W.; Yang, Y.; Huang, J.-R.; et al. Inhibition of WEE1 suppresses the tumor growth in laryngeal squamous cell carcinoma. *Front. Pharmacol.* **2018**, *9*, 1041. [[CrossRef](#)] [[PubMed](#)]
234. Mendez, E.; Rodriguez, C.P.; Kao, M.C.; Raju, S.; Diab, A.; Harbison, R.A.; Konnick, E.Q.; Mugundu, G.M.; Santana-Davila, R.; Martins, R.; et al. A phase I clinical trial of AZD1775 in combination with neoadjuvant weekly docetaxel and cisplatin before definitive therapy in head and neck squamous cell carcinoma. *Clin. Cancer Res.* **2018**, *24*, 2740–2748. [[CrossRef](#)] [[PubMed](#)]
235. Diab, A.; Kao, M.; Kehrl, K.; Kim, H.Y.; Sidorova, J.; Mendez, E. Multiple defects sensitize p53-deficient head and neck cancer cells to the Wee1 kinase inhibition. *Mol. Cancer Res.* **2019**, *17*, 1115–1128. [[CrossRef](#)] [[PubMed](#)]
236. Diab, A.; Gem, H.; Swanger, J.; Kim, H.Y.; Smith, K.; Zou, G.; Raju, S.; Kao, M.; Fitzgibbon, M.; Loeb, K.R.; et al. FOXM1 drives HPV+ HNSCC sensitivity to WEE1 inhibition. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 28287–28296. [[CrossRef](#)]
237. Kong, A.; Good, J.; Kirkham, A.; Savage, J.; Mant, R.; Llewellyn, L.; Parish, J.; Spruce, R.; Forster, M.; Schipani, S.; et al. Phase I trial of WEE1 inhibition with chemotherapy and radiotherapy as adjuvant treatment, and a window of opportunity trial with cisplatin in patients with head and neck cancer: The WISTERIA trial protocol. *BMJ Open* **2020**, *10*, e033009. [[CrossRef](#)]
238. González Besteiro, M.A.; Gottifredi, V. The fork and the kinase: A DNA replication tale from a CHK1 perspective. *Mutat. Res. Rev. Mutat. Res.* **2015**, *763*, 168–180. [[CrossRef](#)]
239. Chen, T.; Stephens, P.A.; Middleton, F.K.; Curtin, N.J. Targeting the S and G2 checkpoint to treat cancer. *Drug Discov. Today* **2012**, *17*, 194–202. [[CrossRef](#)]
240. Furgason, J.M.; Bahassi, E.M. Targeting DNA repair mechanisms in cancer. *Pharmacol. Ther.* **2013**, *137*, 298–308. [[CrossRef](#)]
241. Goto, H.; Izawa, I.; Li, P.; Inagaki, M. Novel regulation of checkpoint kinase 1: Is checkpoint kinase 1 a good candidate for anti-cancer therapy? *Cancer Sci.* **2012**, *103*, 1195–1200. [[CrossRef](#)]
242. Toledo, L.; Neelsen, K.J.; Lukas, J. Replication catastrophe: When a checkpoint fails because of exhaustion. *Mol. Cell* **2017**, *66*, 735–749. [[CrossRef](#)]
243. Khan, S.; He, Y.; Zhang, X.; Yuan, Y.; Pu, S.; Kong, Q.; Zheng, G.; Zhou, D. PROteolysis TArgeting Chimeras (PROTACs) as emerging anticancer therapeutics. *Oncogene* **2020**, *39*, 4909–4924. [[CrossRef](#)] [[PubMed](#)]
244. Motwani, M.; Pesiridis, S.; Fitzgerald, K.A. DNA sensing by the cGAS–STING pathway in health and disease. *Nat. Rev. Genet.* **2019**, *20*, 657–674. [[CrossRef](#)] [[PubMed](#)]
245. Yum, S.; Li, M.; Chen, Z.J. Old dogs, new trick: Classic cancer therapies activate cGAS. *Cell Res.* **2020**, *30*, 639–648. [[CrossRef](#)] [[PubMed](#)]
246. Reisländer, T.; Groelly, F.J.; Tarsounas, M. DNA damage and cancer immunotherapy: A STING in the tale. *Mol. Cell* **2020**, *80*, 21–28. [[CrossRef](#)] [[PubMed](#)]