

Review

# Targeting Cancer Stem Cells: A Strategy for Effective Eradication of Cancer

Masahiro Shibata <sup>1,2</sup> and Mohammad Obaidul Hoque <sup>1,3,4,\*</sup> 

<sup>1</sup> Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University School of Medicine, Baltimore, MD 21231 USA; m-shibata@med.nagoya-u.ac.jp

<sup>2</sup> Department of Breast and Endocrine Surgery (Surgery II), Nagoya University Graduate School of Medicine, Nagoya 4668550, Japan

<sup>3</sup> Department of Oncology, Johns Hopkins University School of Medicine, Baltimore, MD 21231, USA

<sup>4</sup> Department of Urology, Johns Hopkins University School of Medicine, Baltimore, MD 21231, USA

\* Correspondence: mhoque1@jhmi.edu; Tel.: +1-410-502-8778; Fax: +1-410-614-1411

Received: 11 April 2019; Accepted: 23 May 2019; Published: 27 May 2019



**Abstract:** Cancer stem cells (CSCs) are subpopulations of tumor cells with the ability to self-renew, differentiate, and initiate and maintain tumor growth, and they are considered to be the main drivers of intra- and inter-tumoral heterogeneity. While conventional chemotherapy can eradicate the majority of non-CSC tumor cells, CSCs are often drug-resistant, leading to tumor recurrence and metastasis. The heterogeneity of CSCs is the main challenge in developing CSC-targeting therapy; therefore, we and other investigators have focused on developing novel therapeutic strategies that combine conventional chemotherapy with inhibitors of CSC-regulating pathways. Encouraging preclinical findings have suggested that CSC pathway blockade can indeed enhance cellular sensitivity to non-targeted conventional therapy, and this work has led to several ongoing clinical trials of CSC pathway inhibitors. Our studies in bladder cancer and lung adenocarcinoma have demonstrated a crucial role of YAP1, a transcriptional regulator of genes that promote cell survival and proliferation, in regulating CSC phenotypes. Moreover, using cell lines and patient-derived xenograft models, we showed that inhibition of YAP1 enhances the efficacy of conventional therapies by attenuating CSC stemness features. In this review, we summarize the therapeutic strategies for targeting CSCs in several cancers and discuss the potential and challenges of the approach.

**Keywords:** cancer; cancer stem cell; YAP1; verteporfin; combinational therapy

## 1. Introduction

The aggressive nature of certain solid tumors is thought to result, at least in part, from the expansion of cancer stem cell (CSC) populations with self-renewal and differentiation properties that ultimately lead to intra- and inter-tumoral heterogeneity [1–5]. Indeed, the ratio of CSCs to non-CSCs in a tumor cell population correlates with poor clinical outcome in many cancers [6]. While chemotherapy usually eradicates the bulk population of non-CSC tumor cells, CSCs are relatively chemoresistant and not only become enriched after chemotherapy [7–13] but also enter a transient dormant status that may result in tumor recurrence and metastasis [14,15]. Recent work suggests that CSCs may contribute to therapy resistance by repopulating residual tumors between chemotherapy cycles [14]. Thus, a promising and urgently needed therapeutic strategy for many solid tumors is to develop novel therapies that inhibit specific CSC-generating and -expanding pathways in a given tumor [16,17].

Increasing evidence suggests that blockade of factors and/or pathways necessary for CSC survival and growth can increase the cellular sensitivity to conventional chemotherapy, and several clinical trials aiming to abrogate CSCs are currently ongoing [4]. However, it is important to note that CSCs


are not a single cell type but instead are composed of multiple heterogeneous phenotypes, making it extremely difficult to predict whether a specific CSC-targeting therapy will be effective for an individual patient. Furthermore, there are currently no known universal markers for CSCs, which eliminates the possibility of developing a ‘pan-CSC’-targeting therapy [18,19]. Due to this diversity among CSCs, there is a need to identify CSC-specific markers or regulatory pathways characteristic of a given tumor to enable the development of tailored therapies. In other words, effective targeting of CSCs will require a better understanding of the intra- and inter-tumoral diversities of CSC types and functions.

In this regard, the past several years have seen increasing efforts to develop CSC-targeting therapeutic approaches. One of these is YAP1, a transcriptional regulator and the main effector of the Hippo pathway, which normally regulates cellular proliferation and differentiation but also promotes malignant phenotypes and drug resistance in various cancers [20]. Our group recently demonstrated that YAP1 regulates the generation and expansion of CSCs in urothelial carcinoma of bladder and lung adenocarcinoma and that pharmacological inhibition of YAP1, in combination with chemotherapies, suppresses cancer progression by attenuating cancer stemness features [20–22].

Here, we review current knowledge of some of the potential mechanisms by which CSCs can be targeted and the therapeutic challenges faced in eliminating CSCs in various cancers. We also discuss our recent works on CSCs with a particular focus on YAP1-targeting strategies.

## 2. CSC-Targeting Therapies

Recently, several attempts have been conducted aiming to combine conventional drugs with CSC-targeting agents (Figure 1) [6,19]. However, the heterogeneity and high plasticity of not only CSCs but also many non-CSC tumor cells amplify the challenge of identifying suitable targeting molecules for diverse cancers [23]. Such plasticity enables non-CSCs to transdifferentiate into CSCs that may be resistant to specific CSC-targeting therapies [24]. In addition, many surface markers are also expressed on normal stem cell populations where they may play essential roles in tissue hemostasis and renewal [24]. Therefore, there is an urgent need to identify novel CSC specific molecules and/or pathways for specific cancer types that are essentially absent (or expressed at low levels) in normal stem cells.



**Figure 1.** Combining conventional cytotoxic drugs with cancer stem cell (CSC)-targeting agents. (a) Although chemotherapeutic and molecular-targeted drugs can attack most cancer cells, CSCs can evade these agents, leading to tumor regrowth. (b) Combination therapy with CSC-targeting agents and conventional drugs is predicted to be more effective because it eliminates both CSCs and non-CSC tumor cells.

Numerous studies have described the existence of CSC populations in a variety of cancers, including breast, colorectal, lung, ovary, bladder, and head and neck cancers. Malignant stemness markers and factors differ among cancer types, among the same cancer in different patients, and among different regions within the same tumors; nevertheless, all CSCs share the ability to drive tumorigenesis, metastasis, drug resistance, and immunosuppressive microenvironment [6,25]. These features make CSCs attractive targets for the development of more effective treatments for cancer [6]. To date, attempts have been made to target stemness markers, such as STAT3 and NANOG [6], as well as pathways that regulate malignant stemness, most notably the Wnt/ $\beta$ -catenin, Notch, hedgehog, and JAK-STAT pathways, which promote stemness features in various cancers [26]. We summarize these attempts and current status in Table 1. In the following sections, we describe some of the CSC markers and pathways that are subject to recent efforts to develop therapeutic strategies in several major cancers.

**Table 1.** Therapeutic attempts to target cancer stem cell (CSC).

Target	Cancer Type	Inhibitor	Result	Reference
ALDH1	Breast	Histone deacetylase inhibitor	CD44 <sup>+</sup> CD24 <sup>-/low</sup> cell population was decreased and stemness markers were downregulated.	[27,28]
	NSCLC	Disulfiram	Combination of disulfiram and copper downregulated stemness-related genes.	[29]
			When combined with diethylamino-benzaldehyde resensitized resistant cells to cisplatin.	[30]
	Ovary	Solanum incanum extract	A phase II trial showed prolonged survival when disulfiram was combined with cisplatin and vinorelbine.	[31]
	Ovary	Solanum incanum extract	Notch1 and FoxM1 were downregulated, which resulted in increased chemotherapeutic sensitivities.	[32]
CD44	Breast	Anti CD44 antibody	Nanoparticles with CD44 antibody and gemcitabine specifically targeted CD44 <sup>+</sup> cells.	[33]
CD133	Ovary	Anti CD133 antibody-toxin conjugate	Cellular growth was inhibited and tumor progression was suppressed in a mouse model.	[34]
Hedgehog	Bladder	Cyclopamine	Tumor formation was suppressed via inhibition of GALNT1 that mediates SHH signaling.	[35]
	Lung	GDC-0449	Stemness-related features were suppressed in both NSCLC and small-cell lung cancer cells.	[36]
KLF5	Breast	Metformin	CSC growth was inhibited through suppression of <i>NANOG</i> and <i>FGF-BP1</i> (downstream targets of KLF5).	[37,38]
Notch2 and Notch3	Various cancers Small-cell lung	Tarextumab	Tumorigenesis and cellular growth were suppressed and chemotherapeutic efficacy was increased.	[39]
			A phase Ib trial showed good tolerability and anti-tumor effect.	[40]
PI3K/AKT	Bladder	MyrtoCommulone-A	Several stem cell markers were downregulated and stemness-related features were attenuated.	[41]
	Bladder	Motesanib	Survival-related genes in the PI3K/AKT pathway were decreased and cisplatin sensitivity was enhanced.	[42]
STAT3	Breast	STAT3 inhibitor VII	Combination of STAT3 inhibitor and carboplatin abrogated carboplatin-induced ALDH <sup>+</sup> cell enrichment.	[15]
	Colon	Napabucasin	<i>c-Myc</i> , <i>Nanog</i> and <i>Sox2</i> , were downregulated, which attenuated metastasis in a mouse model.	[43]
			Napabucasin showed prolonged survival in phosphorylated STAT3 positive patients.	[44]
Pancreas	Napabucasin	Cancer relapse and metastasis were blocked in mice.	[43]	
NSCLC	NSCLC	OPB-51602	A phase I trial suggested that NSCLC patients were likely to obtain better response.	[45]
Wnt/ $\beta$ -catenin	Breast	Pyruvium pamoate	CD44 <sup>+</sup> CD24 <sup>-/low</sup> and ALDH <sup>+</sup> cells were suppressed by downregulating <i>NANOG</i> , <i>OCT4</i> , and <i>SOX2</i> .	[46]
	Breast	Resveratrol	Resveratrol, which suppressed Wnt/ $\beta$ -catenin pathway, inhibited CSCs and induced autophagy.	[47]
	Ovary	Imatinib	CSC activity was suppressed when combined with platinum chemotherapy.	[48]
			Phase II clinical trials had only a modest impact on the prognosis of ovarian cancer patients.	[49,50]
YAP1	Bladder	Verteporfin	Combination of YAP1 and COX2 inhibitors with chemotherapy attenuated CSC properties and enhanced chemotherapy response.	[21]
	NSCLC Bladder	Verteporfin	Verteporfin attenuated the resistance to EGFR inhibitors.	[21,51,52]

NSCLC: non-small cell lung carcinoma; SHH: sonic hedgehog signaling molecule.

## 2.1. Breast Cancer

CSCs in breast cancer were first identified as cells with a  $CD44^+CD24^{-/low}$  phenotype [53]. In this pioneering study, putative CSCs and non-CSCs were tested for their ability to form tumors in mice. Transfer of as few as 100  $CD44^+CD24^{-/low}$  cells resulted in tumor formation, whereas the transfer of thousands of other cells with various phenotypes did not [53]. Subsequent studies have identified CD44, CD133, and ALDH as CSC markers in breast cancer [54,55]. In addition, several transcription factors, such as KLF4, NANOG, OCT4, and SOX2, have been reported to be expressed and induce stemness features in breast CSCs [56–58]. Interestingly, while breast cancer sphere-forming cells showed high SOX2 expression, NANOG and OCT4 were not expressed in these cells, suggesting the presence of heterogeneous CSC populations [57]. In addition, several deregulated signaling pathways, including hedgehog, Notch, TGF- $\beta$ /BMP, and Wnt/ $\beta$ -catenin are known to contribute to the enhancement of characteristic CSC phenotypes, such as self-renewal, proliferation, apoptosis evasion, invasion, epithelial-mesenchymal transition (EMT), and drug resistance [59,60].

To date, several approaches to therapeutic targeting of CSC populations have been investigated in breast cancer. The percentage of  $CD44^+CD24^-$  cells is higher in basal-type compared with luminal-type breast cancer [61]; therefore, attempts to target CSCs have mainly focused on the basal subtype. Inhibitors of histone deacetylases, which are important epigenetic modifying enzymes, reduced the abundance of  $CD44^+CD24^{-/low}$ -expressing cells, inhibit the activity of ALDH1, and suppress the expressions of several stemness markers, such as *BMI1*, *NANOG* and *OCT4* [27,28]. Recently, treatment of breast CSCs with interferon- $\beta$  in vitro has been reported to limit stemness, migration, sphere-forming properties, and re-expression of CD24, and promote an epithelial-like morphology [62]. Another study found that the type 2 diabetes drug metformin suppresses CSC growth by targeting KLF5 for degradation and preventing transcription of its downstream target genes, *NANOG* and *FGF-BP1* [37]. Interestingly, metformin has been mentioned as a potential CSC-targeting drug for use as (neo-)adjuvant therapy [38]. The possibility that cytotoxic drugs can selectively be delivered to CSCs is supported by the demonstration that iron oxide magnetic nanoparticles containing anti-CD44 antibody and gemcitabine derivatives can specifically target and kill  $CD44^+$  cells [33].

Several therapeutic agents have been evaluated to target the Wnt/ $\beta$ -catenin signaling pathway, which is an important regulator of CSC characteristics. Resveratrol, a natural polyphenolic compound, reduced the breast CSC population in mice via inhibition of Wnt/ $\beta$ -catenin signaling [47], and a highly potent small molecule antagonist of  $\beta$ -catenin binding to nuclear T-cell factor has been shown to inhibit the growth of breast CSCs and, to a lesser extent, non-CSCs [63]. Pyrvinium pamoate, an anthelmintic drug and inhibitor of the Wnt/ $\beta$ -catenin pathway, prevented the proliferation of breast cancer cells, especially  $CD44^+CD24^{-/low}$  and ALDH $^+$  CSCs, via downregulating NANOG, OCT4, and SOX2 [46]. In a recent study, carboplatin treatment activates STAT3, leading to breast CSCs enrichment, and combination treatment with a STAT3 inhibitor and carboplatin attenuated the stemness-like features, resulting in a more efficient therapeutic response [15].

Micro RNAs (miRNAs) regulate gene expression by destabilizing and/or silencing the translocation of target mRNAs, and many miRNAs with CSC-promoting or -suppressing properties have been investigated as potential therapeutic targets. For example, the tumor suppressor miR-223 is downregulated in  $CD44^+CD24^{-/low}$  triple-negative breast CSCs, and its overexpression resensitized the cells to induction of apoptosis [64]. Inhibition of miR-125a, which regulates TAZ, an effector in the Hippo pathway, led to a significant reduction in the breast CSC pool [65], and miR-34a has also been reported to suppress breast CSC-like characteristics by inhibiting the Notch1 signaling pathway [66].

## 2.2. Colorectal Cancer

CD133 has been identified as a marker of colon CSCs, which make up approximately 2.5% of colorectal cancer tumor cells. Notably,  $CD133^+$  cancer cells rapidly formed tumors after injection into immunodeficient mice, whereas  $CD133^-$  cells did not [67]. Moreover,  $CD133^+$  colorectal cancer cells have also been shown to be resistant to radiotherapy and chemotherapy [68], consistent with a CSC

phenotype. Another known CSC marker, CD44 is enriched on CSC cells with CSC-like properties and may promote their function by forming a positive feedback loop with Ras signaling [69], and CD26<sup>+</sup> colorectal CSCs contribute to tumor initiation by facilitating the EMT [70]. The G protein-coupled receptor LGR5 has also been reported to be a marker for colorectal CSCs during the initial stages of tumorigenesis, and *LGR5* expression levels correlated with aggressive clinicopathological features in colorectal cancer [71–73]. Interestingly, combination targeting of both LGR5<sup>+</sup> cells and differentiated cancer cells prevented tumor resistance and relapse [71,72]. Consistent with their roles in other cancer types, the transcription factors NANOG, OCT4, and SOX2 promote stemness features in colorectal CSCs [68,74].

The transcription factor STAT3 is activated by many signaling pathways involved in the regulation of cell growth and apoptosis. Accordingly, STAT3 is an oncogenic driver and contributes to carcinogenesis by promoting cell survival, angiogenesis and the generation and expansion of CSCs, which leads to drug resistance [75–79]. Although further studies are needed, STAT3 is considered to be a promising CSC target in colon cancer [6]. Other study showed that napabucasin, which inhibits STAT3-driven gene transcription, blocks several key molecules in CSC-related signaling pathways, including *c-Myc*, *Nanog* and *Sox2*, in a colon cancer mouse model, thereby suppressing metastases [43].

In addition to being the subject of numerous preclinical studies, several colorectal CSC targeting therapies have advanced through clinical trials. For example, napabucasin could be safely combined with FOLFIRI (folinic acid, fluorouracil, and irinotecan), a standard chemotherapy regimen for colorectal cancer, and showed encouraging signs of efficacy in phase I and II clinical trials [80]. Subsequently, a phase III trial of napabucasin showed prolonged survival of phosphorylated STAT3 positive patients in advanced colon cancer [44].

### 2.3. Lung Cancer

Multiple markers of lung CSCs have been identified, including CD44, CD133, CD166, and ALDH1 [7,81,82]. However, the intra-tumoral heterogeneity and high plasticity of lung cancer have made it difficult to precisely define which markers drive the CSC properties in a given tumor [82]. Among these molecules, CD44, which regulates adhesion, differentiation, homing and migration, is considered to be a key CSC marker [82]. CD44<sup>+</sup> non-small cell lung cancer (NSCLC) cells were shown to specifically form spheroids and induce tumor initiation via induction of NANOG, OCT4, and SOX2 overexpression [81]. Another study successfully identified lung CSCs as a CD44<sup>+</sup>CD90<sup>+</sup> subpopulation that displayed a mesenchymal morphology and elevated expressions of *NANOG*, *OCT4*, and EMT-related genes [83]. CD133 has also been identified as a lung CSC marker; notably, CD133<sup>+</sup> cells have sphere-forming and tumorigenic capabilities [84], are chemoresistant, and are a biomarker of poor prognosis in lung cancer patients [82]. ALDH1<sup>+</sup> cells also exhibit CSC properties that include the ability to self-renew, initiate tumors, and resist drug therapy [85]. This study also demonstrated that ALDH1<sup>+</sup> cells co-express CD133 in lung cancer patients, and that ALDH1 expression correlated positively with tumor stage and grade and with poor prognosis [85]. One particularly interesting study has explored the expressions of multiple putative stemness markers, such as ALDH1, CD24, CD44, CD133, OCT4, and SOX2, in 371 surgically resected NSCLC specimens [86]. ALDH1, ABCG5, CD44, and SOX2 were shown to be associated with poor differentiation and increased proliferation. In addition, ALDH1, CD44, and SOX2 expressions were more frequently associated with squamous cell carcinoma whereas CD24 and CD166 were more common in adenocarcinoma, which supports the heterogeneity of stemness markers among CSCs of different histologic types [86]. Like other cancer types, lung CSCs have been reported to utilize common signaling pathways, including the Wnt/ $\beta$ -catenin, hedgehog, and Notch pathways [87].

Among the CSC-targeting strategies explored in lung cancer to date, genetic knockdown or pharmacological inhibition of ALDH1 has been shown to attenuate the proliferative and migratory capabilities of NSCLC cells [88]. Similarly, disulfiram, an ALDH1 inhibitor, showed synergistic effects with copper in attenuating lung CSC activity via downregulation of stemness-related genes [29]. In a

study of isolated ALDH1<sup>+</sup> lung cancer cells with stem cell-like characteristics, combination therapy with diethylamino-benzaldehyde and disulfiram was able to resensitize resistant cells to the cytotoxic effects of cisplatin [30]. In a phase IIb trial of NSCLC, disulfiram was well-tolerated and prolonged overall survival when combined with cisplatin and vinorelbine [31]. As in colorectal cancer, STAT3 has been investigated as a CSC-regulatory target in lung cancer. In a phase I trial of the STAT3 inhibitor OPB-51602, EGFR-mutant NSCLC patients were likely to obtain better response among the patients of various cancer types [45]. GDC-0449, an inhibitor of the hedgehog pathway, has also demonstrated efficacy in both NSCLC and small-cell lung cancer via suppression of stemness-related features [36]. Tarextumab, which blocks Notch2 and Notch3 signaling, reduces the abundance of tumor-initiating cells and the growth of lung cancer [39]. A phase Ib trial of tarextumab showed good tolerability and a dose-dependent anti-tumor effect when combined with etoposide and platinum in small-cell lung cancer patients [40].

#### 2.4. Ovarian Cancer

Ovarian cancer is an umbrella term for a broad range of neoplasms of different histopathological types, including serous, endometrioid, clear cell, and mucinous; importantly, these subtypes exhibit different responses to drug treatment [89]. The proportion of CSCs in ovarian cancer varies between 0.1% and 30% depending on the disease type and stage [90]. CD133<sup>+</sup> ovarian cancer cells have been shown to have a more proliferative phenotype than do CD133<sup>-</sup> cells [91]. A recent study showed that CD133<sup>+</sup> CSCs induced CD133<sup>-</sup> cells to undergo the EMT and to display enhanced metastatic capacity via secretion of CCL5 and activation of NF- $\kappa$ B signaling [92]. Another group examined the expression of CD44 on ovarian CSCs by immunohistochemistry and found that its expression was associated with poor prognosis [93]. CD44 knockdown in CSCs suppressed their proliferation, migration/invasion, and sphere-forming abilities and increased their drug sensitivity [93]. ALDH has been reported to be overexpressed in spheroid ovarian CSCs [94], and ALDH1A1 was found to be a predictive biomarker of the response of patients to chemotherapy [95]. Another study also identified an association between ALDH overexpression and poor prognosis and additionally showed that ALDH1A1 overexpression increased the clonogenic potential, expression of DNA repair proteins, and platinum resistance of ovarian cancer cells [96]. ALDH<sup>+</sup>CD133<sup>+</sup> ovarian CSCs appear to have greater growth potential than do ALDH<sup>+</sup>CD133<sup>-</sup> cells; indeed, as few as 11 ALDH<sup>+</sup>CD133<sup>+</sup> cells were required to form tumors in mice [97]. Another study reported that ovarian CSCs were enriched among cells co-expressing ALDH1 and CD44 and that these cells displayed chemoresistance [98]. The authors proposed that high ALDH1 alone may be insufficient to drive the ovarian CSC phenotype, but that ALDH1 and CD44 co-expression may trigger chemoresistance and poor clinical outcome [98].

Many studies have investigated CSC targeting as a therapeutic strategy for ovarian cancer. For example, miR-199a decreased CD44 expression, resulting in reduced tumorigenicity and increased chemotherapeutic sensitivity [99]. In this study, other stemness factors, such as *NANOG*, *OCT4* and *CD133*, were also reduced by miR-199a overexpression [99]. Targeting of CD133<sup>+</sup> cells with an anti-CD133 antibody-toxin conjugate was shown to inhibit the progression of ovarian cancer [34]. ALDH1 overexpression in ovarian CSCs increased taxane and platinum resistance, and conversely, *ALDH1* knockdown overcomes the resistance, enhancing growth inhibition in the presence of chemotherapeutic agents [95]. Treatment of ovarian CSCs with solanum incanum extract inhibited ALDH1 activity and Notch1 and FoxM1 expression, leading to attenuation of stemness features and increased sensitivity to cisplatin and paclitaxel [32]. Imatinib, a small molecule inhibitor of c-KIT (CD117), in combination with platinum cytotoxic drugs inhibited ovarian CSC activity by blockade of the Wnt/ $\beta$ -catenin signaling pathway [48]. However, imatinib had only a modest impact on the prognosis of ovarian cancer patients in phase II clinical trials [49,50]. These unfavorable results may have been due to intra- and/or inter-tumoral heterogeneity among both CSCs and non-CSCs.

## 2.5. Bladder Cancer

The existence of bladder CSCs was first reported in 2009 and has since been confirmed in numerous studies [100]. These cells are thought to be a major reason for the failure of adjuvant treatment and poor therapeutic outcomes of bladder cancer [100,101]. Bladder CSCs can be identified using a number of cell surface markers, including CD44, CD67LR, EMA, and ALDH1A1, and the hedgehog, phosphoinositide 3-kinase (PI3K)/AKT, Wnt/ $\beta$ -catenin, and Notch signaling pathways, which play key roles in the generation, expansion, and the maintenance of stemness, self-renewal, and proliferative potential in these heterogeneous groups of cells [102]. Although agents that target a small number of these pathways and markers are available and some of them are being tested in clinical trials, a clear treatment strategy has not been developed.

One study has shown that basal subtype bladder cancer cells expressing sonic hedgehog signaling molecule (SHH), a vertebrate homolog of *Drosophila* hedgehog, are capable of regenerating all cell types within the urothelium [103], suggesting they are CSCs. Subsequent studies suggested that muscle-invasive bladder carcinoma arises exclusively from SHH-expressing stem cells in the basal urothelium [104] and confirmed the role of SHH in the promotion of EMT, tumorigenicity, and stemness [35,105]. Accordingly, targeting of SHH by cyclopamine was shown to inhibit bladder tumor growth [35].

Available data suggest that targeting of the PI3K/AKT pathway inhibits urothelial tumor growth and metastasis by reducing CSCs and blocking the EMT [106]. The main advances in the area of bladder CSCs involve the PI3K/AKT inhibitor myrtoquomulone-A and the multi-targeted tyrosine kinase inhibitor motesanib. Iskender et al. found that myrtoquomulone-A treatment of the human bladder cancer cell line HTB-9 decreased cell viability, proliferation, and sphere-forming ability and suppressed the expression of common stem cell markers (e.g., CD44, NANOG, OCT4, and SOX2) and other proteins involved in the PI3K/AKT pathway [41]. Similarly, Ho et al. reported that motesanib decreased the expression of survival-related genes in the PI3K/AKT pathway, and combination treatment with motesanib synergistically enhanced the anticancer effect of cisplatin [42].

Elevated activity of the Wnt/ $\beta$ -catenin pathway, which regulates stem cell homeostasis, and increased expression of its components are associated with cancer progression [107]. Among bladder cancer patients, 20 single nucleotide polymorphisms in 40 genes in the Wnt/ $\beta$ -catenin pathway were associated with an increased risk of cancer [108]. A recent study revealed that miR-374a directly targets WNT5A and that low levels of this miRNA were associated with poor prognosis [109]. In this study, miR-374a treatment inhibited the phosphorylation and nuclear translocation of  $\beta$ -catenin, decreasing the expression of cancer stemness-related proteins [109].

The Notch signaling pathway interacts with Wnt and SHH pathways and is involved in the maintenance of stemness features [110,111]. This pathway, which has both oncogenic and tumor-suppressive effects depending on the tissue and cellular context [112], includes four receptors, NOTCH1-4, and five ligands and regulates the transcription of multiple target genes [102]. *NOTCH1* expression is decreased in bladder cancer and its activation reduces cellular proliferation, suggesting that it has a tumor-suppressive role [112]. In contrast, *NOTCH2* acts as an oncogene by promoting cell proliferation, the EMT, and maintenance of stemness; moreover, its inhibition attenuates malignant phenotypes in vitro and in vivo [113]. Due to these contrasting roles and the potential for off-target side effects in other organs, more evidence will be required to support the development of Notch-targeting therapy in cancer [106].

## 2.6. Head and Neck Cancer

As in other cancer types, CD44<sup>+</sup> cells have been reported to possess the unique properties of CSCs in head and neck squamous carcinoma (HNSCC) [114]. In addition, CD24<sup>+</sup>CD44<sup>+</sup> cells were found to be more chemoresistant and tumorigenic compared with CD24<sup>-</sup>CD44<sup>+</sup> cancer cells [115]. CD133 has also been investigated as a CSC marker in HNSCC; consistent with this, CD133<sup>+</sup> cells showed reduced sensitivity to paclitaxel [116,117]. In another study, CD10<sup>+</sup> HNSCC cells expressed OCT3/4,

were relatively resistant to cisplatin and radiation, and had sphere-forming and aggressive tumorigenic properties [118], suggesting that CD10 may be a marker of HNSCC CSCs. In nasopharyngeal carcinoma, the expression levels of NANOG, OCT4, and SOX2 were positively correlated with EMT-associated indicators and worse prognosis [119]. In an oral squamous carcinoma mouse model, Lin28, a key factor for cellular reprogramming and generation of induced pluripotent stem cells, was found to promote stemness properties via inhibition of Let-7 and induction of Oct4 and Sox2 [120]. ALDH1 has also been considered to be a CSC marker in HNSCC; accordingly, increased ALDH1 activity enhances tumorigenesis and decreases the effects of chemoradiotherapy [121]. Moreover, ALDH<sup>+</sup>CD44<sup>+</sup> HNSCC cells are more tumorigenic and have stronger stemness characteristics than cells expressing CD44<sup>+</sup> alone [117].


The oncoprotein BMI1 cooperates with MYC and Snail signaling to suppress cellular senescence and apoptosis and to promote cell proliferation and the EMT [122]. In laryngeal carcinoma, BMI1 was co-expressed with CD133 and found to have proliferation-promoting and anti-apoptotic effects [123]. Examination of specimens from patients with oral lichen planus showed that high BMI1 levels were associated with a higher risk of progressing to oral squamous cell carcinoma [124]. In another study, the expression of BMI1 or ALDH1 was associated with the transformation from oral erythroplakia to cancer [125]. Chemoradiation is a standard therapy for HNSCC, and CSCs generally show resistance to both chemotherapy and radiotherapy via control of cell cycling, efficient DNA repair, elevated free radical scavenging, and microenvironment support [117]. Indeed, in several studies, BMI1 expression was associated with a lack of response to chemoradiotherapy [126,127]. Although there is currently little supportive evidence, it is possible that eliminating CSCs by targeting CSC generating and maintaining pathways may increase the sensitivity of HNSCC to chemotherapy and radiotherapy.

### 3. Targeting CSCs by Inhibiting YAP1

In an attempt to target CSC-regulating effector molecules, we have focused on YAP1, a principal effector of the Hippo signaling pathway that regulates organ growth in normal tissues. Increasing evidence suggests that YAP1 has diverse roles in tumorigenesis and drug resistance [20], and it has been reported to promote malignant features (e.g., cell proliferation, invasion, migration, and anti-apoptosis) [20,128–132] and drug resistance in various cancers, including breast, bladder and lung cancer (Figure 2) [51,52,133]. Recent studies have also shown that YAP1 contributes to the establishment of an immunosuppressive tumor microenvironment [20]. Moreover, there is substantial evidence that YAP1 regulates malignant cell stemness [21]. YAP1 is capable of reprogramming non-CSCs into cells that have CSC-like characteristics [134] and helps cancer cells to maintain their stemness by promoting autophagy [135,136]. In NSCLC, YAP1 directly interacts with OCT4 followed by SOX2 upregulation to facilitate self-renewal and acquisition of endothelial-like properties (vascular mimicry) of CSCs [137]. In ovarian cancer, YAP1 regulates pluripotency and chemoresistance [138]. YAP1 induction concurrently induces two potential stemness markers, EPCAM and keratin 19 in hepatocellular carcinoma [139]. Most recently, YAP1 has been reported to activate SOX9 to confer CSC-like features in esophageal squamous cell carcinoma [140]. In addition, YAP1 signaling is activated by long noncoding RNAs in liver cancer, leading to the induction of self-renewal by CSCs [141]. Importantly, YAP1 interacts with numerous oncogenic signaling pathways, including MAPK, PI3K/mTOR, and Hippo, which enhance tumorigenesis [142]. The interplay between YAP1 and the Wnt/ $\beta$ -catenin pathway is known to be important for the maintenance and expansion of CSCs [142,143]. For example, in the absence of Wnt signaling, YAP1 is restricted to the  $\beta$ -catenin destruction complex, but activation of Wnt signaling induces their release and translocation to the nucleus, where they upregulate transcription of  $\beta$ -catenin and YAP1 target genes [144]. Cancerous tissue is composed of not only cancer cells but also mesenchymal cells, endothelial cells, and immune cells [20]. Interestingly, YAP1 in these cells also contributes to cancer progression. For example, in a stromal cell, YAP1 upregulates transcription of myofibroblast marker, such as *CYR-61* and *CTGF* for activating cancer-associated fibroblast that is required for remodeling cancer stroma [145]. In naïve T-cell, YAP1 promotes polarization to regulatory



T-cell through inducing TGFBR2 for immunosuppressive tumor microenvironment [146]. From these studies, YAP1 is considered an attractive therapeutic target.




**Figure 2.** Tumor-promoting roles of YAP1. YAP1 contributes to cancer progression from multiple aspects, such as tumorigenesis, metastasis, malignant stemness, and immunosuppressive microenvironment.

Verteporfin, a porphyrin derivative, is a photosensitizer used in the photodynamic therapy of macular degeneration [20]. Among several drugs, such as dasatinib and statins, that are capable of blocking YAP1 activities [143,147], verteporfin is known as the most effective suppressor of YAP1 activity and widely used in preclinical research [148,149]. Combination treatment of verteporfin and erlotinib suppressed the tumorigenic properties of erlotinib-resistant NSCLC and bladder cancer cells [21,51]. Moreover, the addition of verteporfin to gefitinib decreased the viability of gefitinib-resistant NSCLC and bladder cancer cells [21,52]. Although several ongoing clinical trials are now evaluating the efficacy of verteporfin as a photosensitizer in different types of cancers, none have declared the intent to evaluate YAP1-inhibitory effects.

Our group recently showed that YAP1 is overexpressed in basal-type bladder cancer that has abundant stem cell phenotype [21]. In this study, high YAP1 expression was associated with increased levels of mesenchymal markers, and YAP1 induced CSC properties, such as sphere-forming and self-renewal abilities, invasiveness and drug resistance via induction of SOX2. We also demonstrated that COX2/PGE2 signaling upregulates SOX2 and that a feedback mechanism regulates SOX2 through these pathways. Moreover, combination treatment with verteporfin and a COX2 inhibitor enhanced chemotherapeutic efficacy through the suppression of CSC properties [21]. Further work will be necessary to determine the potential clinical utility of this combination therapy for targeting CSCs.

We have also demonstrated that YAP1 induces STAT3 phosphorylation in lung adenocarcinoma cells by increasing expression of interleukin-6 (IL-6) (manuscript under review). Both YAP1 and STAT3 have been reported to regulate CSCs and other oncogenic pathways [20,21,79]. Indeed, genetic inhibition of both *YAP1* and *STAT3* decreased the stem cell-like features of lung adenocarcinoma cells, resulting in reduced proliferation and enhanced cisplatin sensitivity. Verteporfin was found to suppress not only YAP1 but also STAT3, a previously reported off-target effect of this compound [150]. Using patient-derived xenograft mice models in which the heterogeneity of the primary human tumor was preserved, we found that simultaneous blockade of YAP1 and STAT3 by verteporfin suppressed CSC activity and tumor growth when combined with chemotherapy. Therefore, verteporfin could be an ideal drug for targeting CSC properties by suppressing both YAP1 and STAT3 (Figure 3).

As noted above, verteporfin has already been used to treat of macular degeneration and known to cause minor side effects, such as blurred vision and headache with rare systemic adverse events, making the path to the clinic easier for this drug than for novel YAP1 and STAT3 inhibitors with unknown safety profiles. Although the available evidence suggests that the combination of conventional therapy plus a YAP1 inhibitor holds great promise for the suppression of CSC expansion, we do not yet know the extent to which the heterogeneity of CSCs will be a confounding factor. Further work will be necessary to understand how different classes of CSCs could be targeted.



**Figure 3.** YAP1 and STAT3; Two oncogenic pathways that promote cancer stemness. YAP1 and STAT3 are independently involved in oncogenic signaling to promote cancer stem cell (CSC) properties. However, YAP1 also promotes IL-6-induced STAT3 phosphorylation and activation. The porphyrin derivative verteporfin inhibits both the YAP1 and STAT3 pathways and may thus be an efficient suppressor of CSC properties.

#### 4. Conclusions

In this review, we have discussed some of the potential mechanisms for targeting CSCs, the therapeutic challenges presented in various cancers, and the development of YAP1-targeting strategies. Considering the complexity and diversity of CSCs and non-CSCs, it will be extremely challenging to completely eradicate all cancer cells. However, the addition of CSC-targeting agents to the repertoire of conventional chemotherapeutic drugs represents a promising strategy to induce synergistic effects against both CSCs and non-CSCs.

**Author Contributions:** M.S. searched the literature and wrote the manuscript. M.O.H. drew conceptual frameworks and revised the manuscript.

**Funding:** This work was funded by NCI R01CA206027 and NCI R01CA208709 (to M.O.H.).

**Acknowledgments:** We thank Anne M. O'Rourke, from Edanz Group ([www.edanzediting.com/ac](http://www.edanzediting.com/ac)) for editing a draft of this manuscript.

**Conflicts of Interest:** The authors declare no conflicts of interest.

#### References

1. Testa, U.; Castelli, G.; Pelosi, E. Lung Cancers: Molecular Characterization, Clonal Heterogeneity and Evolution, and Cancer Stem Cells. *Cancers* **2018**, *10*, 248. [[CrossRef](#)] [[PubMed](#)]
2. Chen, Z.; Fillmore, C.M.; Hammerman, P.S.; Kim, C.F.; Wong, K.K. Non-small-cell lung cancers: A heterogeneous set of diseases. *Nat. Rev. Cancer* **2014**, *14*, 535–546. [[CrossRef](#)] [[PubMed](#)]
3. Qu, H.; Li, R.; Liu, Z.; Zhang, J.; Luo, R. Prognostic value of cancer stem cell marker CD133 expression in non-small cell lung cancer: A systematic review. *Int. J. Clin. Exp. Pathol.* **2013**, *6*, 2644–2650.
4. MacDonagh, L.; Gray, S.G.; Breen, E.; Cuffe, S.; Finn, S.P.; O'Byrne, K.J.; Barr, M.P. Lung cancer stem cells: The root of resistance. *Cancer Lett.* **2016**, *372*, 147–156. [[CrossRef](#)]
5. Singh, A.; Settleman, J. EMT, cancer stem cells and drug resistance: An emerging axis of evil in the war on cancer. *Oncogene* **2010**, *29*, 4741–4751. [[CrossRef](#)] [[PubMed](#)]

6. Desai, A.; Yan, Y.; Gerson, S.L. Concise Reviews: Cancer Stem Cell Targeted Therapies: Toward Clinical Success. *Stem Cells Transl. Med.* **2018**, *8*, 75–81. [[CrossRef](#)]
7. Zakaria, N.; Satar, N.A.; Abu Halim, N.H.; Ngali, S.H.; Yusoff, N.M.; Lin, J.; Yahaya, B.H. Targeting Lung Cancer Stem Cells: Research and Clinical Impacts. *Front. Oncol.* **2017**, *7*, 80. [[CrossRef](#)]
8. Leon, G.; MacDonagh, L.; Finn, S.P.; Cuffe, S.; Barr, M.P. Cancer stem cells in drug resistant lung cancer: Targeting cell surface markers and signaling pathways. *Pharmacol. Ther.* **2016**, *158*, 71–90. [[CrossRef](#)]
9. Suresh, R.; Ali, S.; Ahmad, A.; Philip, P.A.; Sarkar, F.H. The Role of Cancer Stem Cells in Recurrent and Drug-Resistant Lung Cancer. *Adv. Exp. Med. Biol.* **2016**, *890*, 57–74. [[CrossRef](#)]
10. Singh, S.; Chellappan, S. Lung cancer stem cells: Molecular features and therapeutic targets. *Mol. Asp. Med.* **2014**, *39*, 50–60. [[CrossRef](#)]
11. Gottschling, S.; Schnabel, P.A.; Herth, F.J.; Herpel, E. Are we missing the target? Cancer stem cells and drug resistance in non-small cell lung cancer. *Cancer Genom. Proteom.* **2012**, *9*, 275–286.
12. Shibue, T.; Weinberg, R.A. EMT, CSCs, and drug resistance: The mechanistic link and clinical implications. *Nat. Rev. Clin. Oncol.* **2017**, *14*, 611–629. [[CrossRef](#)]
13. Codony-Servat, J.; Rosell, R. Cancer stem cells and immunoresistance: Clinical implications and solutions. *Transl. Lung. Cancer Res.* **2015**, *4*, 689–703. [[CrossRef](#)]
14. Kurtova, A.V.; Xiao, J.; Mo, Q.; Pazhanisamy, S.; Krasnow, R.; Lerner, S.P.; Chen, F.; Roh, T.T.; Lay, E.; Ho, P.L.; et al. Blocking PGE2-induced tumour repopulation abrogates bladder cancer chemoresistance. *Nature* **2015**, *517*, 209–213. [[CrossRef](#)]
15. Lu, H.; Chen, I.; Shimoda, L.A.; Park, Y.; Zhang, C.; Tran, L.; Zhang, H.; Semenza, G.L. Chemotherapy-Induced Ca(2+) Release Stimulates Breast Cancer Stem Cell Enrichment. *Cell Rep.* **2017**, *18*, 1946–1957. [[CrossRef](#)] [[PubMed](#)]
16. Ramos, E.K.; Hoffmann, A.D.; Gerson, S.L.; Liu, H. New Opportunities and Challenges to Defeat Cancer Stem Cells. *Trends Cancer* **2017**, *3*, 780–796. [[CrossRef](#)] [[PubMed](#)]
17. Fiori, M.E.; Villanova, L.; De Maria, R. Cancer stem cells: At the forefront of personalized medicine and immunotherapy. *Curr. Opin. Pharmacol.* **2017**, *35*, 1–11. [[CrossRef](#)] [[PubMed](#)]
18. Aoi, T. Biology of lung cancer: Genetic mutation, epithelial-mesenchymal transition, and cancer stem cells. *Gen. Thorac. Cardiovasc. Surg.* **2016**, *64*, 517–523. [[CrossRef](#)] [[PubMed](#)]
19. Wang, T.; Shigdar, S.; Gantier, M.P.; Hou, Y.; Wang, L.; Li, Y.; Shamaileh, H.A.; Yin, W.; Zhou, S.F.; Zhao, X.; et al. Cancer stem cell targeted therapy: Progress amid controversies. *Oncotarget* **2015**, *6*, 44191–44206. [[CrossRef](#)]
20. Shibata, M.; Ham, K.; Hoque, M.O. A time for YAP1: Tumorigenesis, immunosuppression and targeted therapy. *Int. J. Cancer* **2018**, *143*, 2133–2144. [[CrossRef](#)] [[PubMed](#)]
21. Ooki, A.; Del Carmen Rodriguez Pena, M.; Marchionni, L.; Dinalankara, W.; Begum, A.; Hahn, N.M.; VandenBussche, C.J.; Rasheed, Z.A.; Mao, S.; Netto, G.J.; et al. YAP1 and COX2 Coordinately Regulate Urothelial Cancer Stem-like Cells. *Cancer Res.* **2018**, *78*, 168–181. [[CrossRef](#)] [[PubMed](#)]
22. Ooki, A.; Dinalankara, W.; Marchionni, L.; Tsay, J.J.; Goparaju, C.; Maleki, Z.; Rom, W.N.; Pass, H.I.; Hoque, M.O. Epigenetically regulated PAX6 drives cancer cells toward a stem-like state via GLI-SOX2 signaling axis in lung adenocarcinoma. *Oncogene* **2018**. [[CrossRef](#)] [[PubMed](#)]
23. Yang, Z.; Li, C.; Fan, Z.; Liu, H.; Zhang, X.; Cai, Z.; Xu, L.; Luo, J.; Huang, Y.; He, L.; et al. Single-cell Sequencing Reveals Variants in ARID1A, GPRC5A and MLL2 Driving Self-renewal of Human Bladder Cancer Stem Cells. *Eur. Urol.* **2017**, *71*, 8–12. [[CrossRef](#)] [[PubMed](#)]
24. Zhu, P.; Fan, Z. Cancer stem cells and tumorigenesis. *Biophys. Rep.* **2018**, *4*, 178–188. [[CrossRef](#)] [[PubMed](#)]
25. Fatima, F.; Nawaz, M. Stem cell-derived exosomes: Roles in stromal remodeling, tumor progression, and cancer immunotherapy. *Chin. J. Cancer* **2015**, *34*, 541–553. [[CrossRef](#)] [[PubMed](#)]
26. Ajani, J.A.; Song, S.; Hochster, H.S.; Steinberg, I.B. Cancer stem cells: The promise and the potential. *Semin. Oncol.* **2015**, *42* (Suppl. 1), S3–S17. [[CrossRef](#)]
27. Li, H.Z.; Yi, T.B.; Wu, Z.Y. Suspension culture combined with chemotherapeutic agents for sorting of breast cancer stem cells. *BMC Cancer* **2008**, *8*, 135. [[CrossRef](#)]
28. Schech, A.; Kazi, A.; Yu, S.; Shah, P.; Sabnis, G. Histone Deacetylase Inhibitor Entinostat Inhibits Tumor-Initiating Cells in Triple-Negative Breast Cancer Cells. *Mol. Cancer Ther.* **2015**, *14*, 1848–1857. [[CrossRef](#)]
29. Duan, L.; Shen, H.; Zhao, G.; Yang, R.; Cai, X.; Zhang, L.; Jin, C.; Huang, Y. Inhibitory effect of Disulfiram/copper complex on non-small cell lung cancer cells. *Biochem. Biophys. Res. Commun.* **2014**, *446*, 1010–1016. [[CrossRef](#)]

30. MacDonagh, L.; Gallagher, M.F.; Ffrench, B.; Gasch, C.; Breen, E.; Gray, S.G.; Nicholson, S.; Leonard, N.; Ryan, R.; Young, V.; et al. Targeting the cancer stem cell marker, aldehyde dehydrogenase 1, to circumvent cisplatin resistance in NSCLC. *Oncotarget* **2017**, *8*, 72544–72563. [[CrossRef](#)]
31. Nechushtan, H.; Hamamreh, Y.; Nidal, S.; Gotfried, M.; Baron, A.; Shalev, Y.I.; Nisman, B.; Peretz, T.; Peylan-Ramu, N. A phase IIb trial assessing the addition of disulfiram to chemotherapy for the treatment of metastatic non-small cell lung cancer. *Oncologist* **2015**, *20*, 366–367. [[CrossRef](#)]
32. Wu, Y.H.; Chiu, W.T.; Young, M.J.; Chang, T.H.; Huang, Y.F.; Chou, C.Y. Solanum Incanum Extract Downregulates Aldehyde Dehydrogenase 1-Mediated Stemness and Inhibits Tumor Formation in Ovarian Cancer Cells. *J. Cancer* **2015**, *6*, 1011–1019. [[CrossRef](#)]
33. Aires, A.; Ocampo, S.M.; Simoes, B.M.; Josefa Rodriguez, M.; Cadenas, J.F.; Couleaud, P.; Spence, K.; Latorre, A.; Miranda, R.; Somoza, A.; et al. Multifunctionalized iron oxide nanoparticles for selective drug delivery to CD44-positive cancer cells. *Nanotechnology* **2016**, *27*, 065103. [[CrossRef](#)]
34. Skubitz, A.P.; Taras, E.P.; Boylan, K.L.; Waldron, N.N.; Oh, S.; Panoskaltis-Mortari, A.; Vallera, D.A. Targeting CD133 in an in vivo ovarian cancer model reduces ovarian cancer progression. *Gynecol. Oncol.* **2013**, *130*, 579–587. [[CrossRef](#)] [[PubMed](#)]
35. Li, C.; Du, Y.; Yang, Z.; He, L.; Wang, Y.; Hao, L.; Ding, M.; Yan, R.; Wang, J.; Fan, Z. GALNT1-Mediated Glycosylation and Activation of Sonic Hedgehog Signaling Maintains the Self-Renewal and Tumor-Initiating Capacity of Bladder Cancer Stem Cells. *Cancer Res.* **2016**, *76*, 1273–1283. [[CrossRef](#)]
36. Tian, F.; Mysliwicz, J.; Ellwart, J.; Gamarra, F.; Huber, R.M.; Bergner, A. Effects of the Hedgehog pathway inhibitor GDC-0449 on lung cancer cell lines are mediated by side populations. *Clin. Exp. Med.* **2012**, *12*, 25–30. [[CrossRef](#)] [[PubMed](#)]
37. Shi, P.; Liu, W.; Wang, H.; Li, F.; Zhang, H.; Wu, Y.; Kong, Y.; Zhou, Z.; Wang, C.; Chen, W.; et al. Metformin suppresses triple-negative breast cancer stem cells by targeting KLF5 for degradation. *Cell Discov.* **2017**, *3*, 17010. [[CrossRef](#)] [[PubMed](#)]
38. Barbieri, F.; Thellung, S.; Ratto, A.; Carra, E.; Marini, V.; Fucile, C.; Bajetto, A.; Pattarozzi, A.; Wurth, R.; Gatti, M.; et al. In vitro and in vivo antiproliferative activity of metformin on stem-like cells isolated from spontaneous canine mammary carcinomas: Translational implications for human tumors. *BMC Cancer* **2015**, *15*, 228. [[CrossRef](#)] [[PubMed](#)]
39. Yen, W.C.; Fischer, M.M.; Axelrod, F.; Bond, C.; Cain, J.; Cancilla, B.; Henner, W.R.; Meisner, R.; Sato, A.; Shah, J.; et al. Targeting Notch signaling with a Notch2/Notch3 antagonist (tarextumab) inhibits tumor growth and decreases tumor-initiating cell frequency. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **2015**, *21*, 2084–2095. [[CrossRef](#)] [[PubMed](#)]
40. Lu, H.; Jiang, Z. Advances in antibody therapeutics targeting small-cell lung cancer. *Adv. Clin. Exp. Med.* **2018**, *27*, 1317–1323. [[CrossRef](#)] [[PubMed](#)]
41. Iskender, B.; Izgi, K.; Karaca, H.; Canatan, H. Myrtilone-A treatment decreases pluripotency- and multipotency-associated marker expression in bladder cancer cell line HTB-9. *J. Nat. Med.* **2015**, *69*, 543–554. [[CrossRef](#)]
42. Ho, J.N.; Byun, S.S.; Lee, S.E.; Youn, J.I.; Lee, S. Multikinase inhibitor motesanib enhances the antitumor effect of cisplatin in cisplatin-resistant human bladder cancer cells via apoptosis and the PI3K/Akt pathway. *Oncol. Rep.* **2019**, *41*, 2482–2490. [[CrossRef](#)] [[PubMed](#)]
43. Li, Y.; Rogoff, H.A.; Keates, S.; Gao, Y.; Murikipudi, S.; Mikule, K.; Leggett, D.; Li, W.; Pardee, A.B.; Li, C.J. Suppression of cancer relapse and metastasis by inhibiting cancer stemness. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 1839–1844. [[CrossRef](#)] [[PubMed](#)]
44. Jonker, D.J.; Nott, L.; Yoshino, T.; Gill, S.; Shapiro, J.; Ohtsu, A.; Zalberg, J.; Vickers, M.M.; Wei, A.C.; Gao, Y.; et al. Napabucasin versus placebo in refractory advanced colorectal cancer: A randomised phase 3 trial. *Lancet. Gastroenterol. Hepatol.* **2018**, *3*, 263–270. [[CrossRef](#)]
45. Wong, A.L.; Soo, R.A.; Tan, D.S.; Lee, S.C.; Lim, J.S.; Marban, P.C.; Kong, L.R.; Lee, Y.J.; Wang, L.Z.; Thuya, W.L.; et al. Phase I and biomarker study of OPB-51602, a novel signal transducer and activator of transcription (STAT) 3 inhibitor, in patients with refractory solid malignancies. *Ann. Oncol.* **2015**, *26*, 998–1005. [[CrossRef](#)]
46. Xu, L.; Zhang, L.; Hu, C.; Liang, S.; Fei, X.; Yan, N.; Zhang, Y.; Zhang, F. WNT pathway inhibitor pyrvinium pamoate inhibits the self-renewal and metastasis of breast cancer stem cells. *Int. J. Oncol.* **2016**, *48*, 1175–1186. [[CrossRef](#)]

47. Fu, Y.; Chang, H.; Peng, X.; Bai, Q.; Yi, L.; Zhou, Y.; Zhu, J.; Mi, M. Resveratrol inhibits breast cancer stem-like cells and induces autophagy via suppressing Wnt/beta-catenin signaling pathway. *PLoS ONE* **2014**, *9*, e102535. [[CrossRef](#)]
48. Chau, W.K.; Ip, C.K.; Mak, A.S.; Lai, H.C.; Wong, A.S. c-Kit mediates chemoresistance and tumor-initiating capacity of ovarian cancer cells through activation of Wnt/beta-catenin-ATP-binding cassette G2 signaling. *Oncogene* **2013**, *32*, 2767–2781. [[CrossRef](#)]
49. Coleman, R.L.; Broaddus, R.R.; Bodurka, D.C.; Wolf, J.K.; Burke, T.W.; Kavanagh, J.J.; Levenback, C.F.; Gershenson, D.M. Phase II trial of imatinib mesylate in patients with recurrent platinum- and taxane-resistant epithelial ovarian and primary peritoneal cancers. *Gynecol. Oncol.* **2006**, *101*, 126–131. [[CrossRef](#)]
50. Schilder, R.J.; Sill, M.W.; Lee, R.B.; Shaw, T.J.; Senterman, M.K.; Klein-Szanto, A.J.; Miner, Z.; Vanderhyden, B.C. Phase II evaluation of imatinib mesylate in the treatment of recurrent or persistent epithelial ovarian or primary peritoneal carcinoma: A Gynecologic Oncology Group Study. *J. Clin. Oncol.* **2008**, *26*, 3418–3425. [[CrossRef](#)] [[PubMed](#)]
51. Hsu, P.C.; You, B.; Yang, Y.L.; Zhang, W.Q.; Wang, Y.C.; Xu, Z.; Dai, Y.; Liu, S.; Yang, C.T.; Li, H.; et al. YAP promotes erlotinib resistance in human non-small cell lung cancer cells. *Oncotarget* **2016**, *7*, 51922–51933. [[CrossRef](#)]
52. Lee, J.E.; Park, H.S.; Lee, D.; Yoo, G.; Kim, T.; Jeon, H.; Yeo, M.K.; Lee, C.S.; Moon, J.Y.; Jung, S.S.; et al. Hippo pathway effector YAP inhibition restores the sensitivity of EGFR-TKI in lung adenocarcinoma having primary or acquired EGFR-TKI resistance. *Biochem. Biophys. Res. Commun.* **2016**, *474*, 154–160. [[CrossRef](#)]
53. Al-Hajj, M.; Wicha, M.S.; Benito-Hernandez, A.; Morrison, S.J.; Clarke, M.F. Prospective identification of tumorigenic breast cancer cells. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 3983–3988. [[CrossRef](#)]
54. Gong, W.; Sun, B.; Sun, H.; Zhao, X.; Zhang, D.; Liu, T.; Zhao, N.; Gu, Q.; Dong, X.; Liu, F. Nodal signaling activates the Smad2/3 pathway to regulate stem cell-like properties in breast cancer cells. *Am. J. Cancer Res.* **2017**, *7*, 503–517.
55. Park, S.Y.; Lee, H.E.; Li, H.; Shipitsin, M.; Gelman, R.; Polyak, K. Heterogeneity for stem cell-related markers according to tumor subtype and histologic stage in breast cancer. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **2010**, *16*, 876–887. [[CrossRef](#)]
56. Hu, T.; Liu, S.; Breiter, D.R.; Wang, F.; Tang, Y.; Sun, S. Octamer 4 small interfering RNA results in cancer stem cell-like cell apoptosis. *Cancer Res.* **2008**, *68*, 6533–6540. [[CrossRef](#)]
57. Leis, O.; Eguiara, A.; Lopez-Arribillaga, E.; Alberdi, M.J.; Hernandez-Garcia, S.; Elorriaga, K.; Pandiella, A.; Rezola, R.; Martin, A.G. Sox2 expression in breast tumours and activation in breast cancer stem cells. *Oncogene* **2012**, *31*, 1354–1365. [[CrossRef](#)]
58. Yu, F.; Li, J.; Chen, H.; Fu, J.; Ray, S.; Huang, S.; Zheng, H.; Ai, W. Kruppel-like factor 4 (KLF4) is required for maintenance of breast cancer stem cells and for cell migration and invasion. *Oncogene* **2011**, *30*, 2161–2172. [[CrossRef](#)]
59. Cochrane, C.R.; Szczepny, A.; Watkins, D.N.; Cain, J.E. Hedgehog Signaling in the Maintenance of Cancer Stem Cells. *Cancers* **2015**, *7*, 1554–1585. [[CrossRef](#)]
60. Mamaeva, V.; Niemi, R.; Beck, M.; Ozliseli, E.; Desai, D.; Landor, S.; Gronroos, T.; Kronqvist, P.; Pettersen, I.K.; McCormack, E.; et al. Inhibiting Notch Activity in Breast Cancer Stem Cells by Glucose Functionalized Nanoparticles Carrying gamma-secretase Inhibitors. *Mol. Ther.* **2016**, *24*, 926–936. [[CrossRef](#)]
61. Nilendu, P.; Kumar, A.; Pal, J.K.; Sharma, N.K. Breast cancer stem cells as last soldiers eluding therapeutic burn: A hard nut to crack. *Int. J. Cancer* **2018**, *142*, 7–17. [[CrossRef](#)] [[PubMed](#)]
62. Doherty, M.R.; Cheon, H.; Junk, D.J.; Vinayak, S.; Varadan, V.; Telli, M.L.; Ford, J.M.; Stark, G.R.; Jackson, M.W. Interferon-beta represses cancer stem cell properties in triple-negative breast cancer. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 13792–13797. [[CrossRef](#)]
63. Jang, G.B.; Hong, I.S.; Kim, R.J.; Lee, S.Y.; Park, S.J.; Lee, E.S.; Park, J.H.; Yun, C.H.; Chung, J.U.; Lee, K.J.; et al. Wnt/beta-Catenin Small-Molecule Inhibitor CWP232228 Preferentially Inhibits the Growth of Breast Cancer Stem-like Cells. *Cancer Res.* **2015**, *75*, 1691–1702. [[CrossRef](#)] [[PubMed](#)]
64. Sun, X.; Li, Y.; Zheng, M.; Zuo, W.; Zheng, W. MicroRNA-223 Increases the Sensitivity of Triple-Negative Breast Cancer Stem Cells to TRAIL-Induced Apoptosis by Targeting HAX-1. *PLoS ONE* **2016**, *11*, e0162754. [[CrossRef](#)] [[PubMed](#)]

65. Nandy, S.B.; Arumugam, A.; Subramani, R.; Pedroza, D.; Hernandez, K.; Saltzstein, E.; Lakshmanaswamy, R. MicroRNA-125a influences breast cancer stem cells by targeting leukemia inhibitory factor receptor which regulates the Hippo signaling pathway. *Oncotarget* **2015**, *6*, 17366–17378. [[CrossRef](#)]
66. Kang, L.; Mao, J.; Tao, Y.; Song, B.; Ma, W.; Lu, Y.; Zhao, L.; Li, J.; Yang, B.; Li, L. MicroRNA-34a suppresses the breast cancer stem cell-like characteristics by downregulating Notch1 pathway. *Cancer Sci.* **2015**, *106*, 700–708. [[CrossRef](#)] [[PubMed](#)]
67. Ricci-Vitiani, L.; Lombardi, D.G.; Pilozzi, E.; Biffoni, M.; Todaro, M.; Peschle, C.; De Maria, R. Identification and expansion of human colon-cancer-initiating cells. *Nature* **2007**, *445*, 111–115. [[CrossRef](#)]
68. Zhou, Y.; Xia, L.; Wang, H.; Oyang, L.; Su, M.; Liu, Q.; Lin, J.; Tan, S.; Tian, Y.; Liao, Q.; et al. Cancer stem cells in progression of colorectal cancer. *Oncotarget* **2018**, *9*, 33403–33415. [[CrossRef](#)]
69. Cheng, C.; Yaffe, M.B.; Sharp, P.A. A positive feedback loop couples Ras activation and CD44 alternative splicing. *Genes Dev.* **2006**, *20*, 1715–1720. [[CrossRef](#)]
70. Pang, R.; Law, W.L.; Chu, A.C.; Poon, J.T.; Lam, C.S.; Chow, A.K.; Ng, L.; Cheung, L.W.; Lan, X.R.; Lan, H.Y.; et al. A subpopulation of CD26+ cancer stem cells with metastatic capacity in human colorectal cancer. *Cell Stem Cell* **2010**, *6*, 603–615. [[CrossRef](#)]
71. Schepers, A.G.; Snippert, H.J.; Stange, D.E.; van den Born, M.; van Es, J.H.; van de Wetering, M.; Clevers, H. Lineage tracing reveals Lgr5+ stem cell activity in mouse intestinal adenomas. *Science* **2012**, *337*, 730–735. [[CrossRef](#)]
72. Shimokawa, M.; Ohta, Y.; Nishikori, S.; Matano, M.; Takano, A.; Fujii, M.; Date, S.; Sugimoto, S.; Kanai, T.; Sato, T. Visualization and targeting of LGR5(+) human colon cancer stem cells. *Nature* **2017**, *545*, 187–192. [[CrossRef](#)]
73. Uchida, H.; Yamazaki, K.; Fukuma, M.; Yamada, T.; Hayashida, T.; Hasegawa, H.; Kitajima, M.; Kitagawa, Y.; Sakamoto, M. Overexpression of leucine-rich repeat-containing G protein-coupled receptor 5 in colorectal cancer. *Cancer Sci.* **2010**, *101*, 1731–1737. [[CrossRef](#)]
74. Xu, F.; Dai, C.; Zhang, R.; Zhao, Y.; Peng, S.; Jia, C. Nanog: A potential biomarker for liver metastasis of colorectal cancer. *Dig. Dis. Sci.* **2012**, *57*, 2340–2346. [[CrossRef](#)]
75. Hu, W.; Jin, P.; Liu, W. Periostin Contributes to Cisplatin Resistance in Human Non-Small Cell Lung Cancer A549 Cells via Activation of Stat3 and Akt and Upregulation of Survivin. *Cell. Physiol. Biochem.: Int. J. Exp. Cell. Physiol. Biochem. Pharmacol.* **2016**, *38*, 1199–1208. [[CrossRef](#)]
76. Li, Y.; Du, H.; Qin, Y.; Roberts, J.; Cummings, O.W.; Yan, C. Activation of the signal transducers and activators of the transcription 3 pathway in alveolar epithelial cells induces inflammation and adenocarcinomas in mouse lung. *Cancer Res.* **2007**, *67*, 8494–8503. [[CrossRef](#)] [[PubMed](#)]
77. Shien, K.; Papadimitrakopoulou, V.A.; Ruder, D.; Behrens, C.; Shen, L.; Kalhor, N.; Song, J.; Lee, J.J.; Wang, J.; Tang, X.; et al. JAK1/STAT3 Activation through a Proinflammatory Cytokine Pathway Leads to Resistance to Molecularly Targeted Therapy in Non-Small Cell Lung Cancer. *Mol. Cancer Ther.* **2017**, *16*, 2234–2245. [[CrossRef](#)] [[PubMed](#)]
78. Zhao, M.; Gao, F.H.; Wang, J.Y.; Liu, F.; Yuan, H.H.; Zhang, W.Y.; Jiang, B. JAK2/STAT3 signaling pathway activation mediates tumor angiogenesis by upregulation of VEGF and bFGF in non-small-cell lung cancer. *Lung. Cancer* **2011**, *73*, 366–374. [[CrossRef](#)]
79. Galoczova, M.; Coates, P.; Vojtesek, B. STAT3, stem cells, cancer stem cells and p63. *Cell. Mol. Biol. Lett.* **2018**, *23*, 12. [[CrossRef](#)]
80. Bendell, J.; Hubbard, J.; O’Neil, B.H.; Jonker, D.; Starodub, A.; Peyton, J.; Pitot, H.; Halfdanarson, T.; Nadeau, B.; Zubkus, J.; et al. LBA-003Phase 1b/II study of cancer stemness inhibitor napabucasin in combination with FOLFIRI +/- bevacizumab (bev) in metastatic colorectal cancer (mCRC) patients (pts). *Ann. Oncol.* **2017**, *28*. [[CrossRef](#)]
81. Leung, E.L.; Fiscus, R.R.; Tung, J.W.; Tin, V.P.; Cheng, L.C.; Sihoe, A.D.; Fink, L.M.; Ma, Y.; Wong, M.P. Non-small cell lung cancer cells expressing CD44 are enriched for stem cell-like properties. *PLoS ONE* **2010**, *5*, e14062. [[CrossRef](#)] [[PubMed](#)]
82. Hardavella, G.; George, R.; Sethi, T. Lung cancer stem cells-characteristics, phenotype. *Transl. Lung. Cancer Res.* **2016**, *5*, 272–279. [[CrossRef](#)]
83. Wang, P.; Gao, Q.; Suo, Z.; Munthe, E.; Solberg, S.; Ma, L.; Wang, M.; Westerdaal, N.A.; Kvalheim, G.; Gaudernack, G. Identification and characterization of cells with cancer stem cell properties in human primary lung cancer cell lines. *PLoS ONE* **2013**, *8*, e57020. [[CrossRef](#)] [[PubMed](#)]

84. Eramo, A.; Lotti, F.; Sette, G.; Piloizzi, E.; Biffoni, M.; Di Virgilio, A.; Conticello, C.; Ruco, L.; Peschle, C.; De Maria, R. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ.* **2008**, *15*, 504–514. [[CrossRef](#)]
85. Jiang, F.; Qiu, Q.; Khanna, A.; Todd, N.W.; Deepak, J.; Xing, L.; Wang, H.; Liu, Z.; Su, Y.; Stass, S.A.; et al. Aldehyde dehydrogenase 1 is a tumor stem cell-associated marker in lung cancer. *Mol. Cancer Res.* **2009**, *7*, 330–338. [[CrossRef](#)]
86. Sterlacci, W.; Savic, S.; Fiegl, M.; Obermann, E.; Tzankov, A. Putative stem cell markers in non-small-cell lung cancer: A clinicopathologic characterization. *J. Thorac. Oncol.* **2014**, *9*, 41–49. [[CrossRef](#)] [[PubMed](#)]
87. Gomez-Casal, R.; Bhattacharya, C.; Ganesh, N.; Bailey, L.; Basse, P.; Gibson, M.; Epperly, M.; Levina, V. Non-small cell lung cancer cells survived ionizing radiation treatment display cancer stem cell and epithelial-mesenchymal transition phenotypes. *Mol. Cancer* **2013**, *12*, 94. [[CrossRef](#)]
88. Moreb, J.S.; Baker, H.V.; Chang, L.J.; Amaya, M.; Lopez, M.C.; Ostmark, B.; Chou, W. ALDH isozymes downregulation affects cell growth, cell motility and gene expression in lung cancer cells. *Mol. Cancer* **2008**, *7*, 87. [[CrossRef](#)]
89. Mihanfar, A.; Aghazadeh Attari, J.; Mohebbi, I.; Majidinia, M.; Kaviani, M.; Yousefi, M.; Yousefi, B. Ovarian cancer stem cell: A potential therapeutic target for overcoming multidrug resistance. *J. Cell. Physiol.* **2018**. [[CrossRef](#)]
90. Visvader, J.E.; Lindeman, G.J. Cancer stem cells in solid tumours: Accumulating evidence and unresolved questions. *Nat. Rev. Cancer* **2008**, *8*, 755–768. [[CrossRef](#)] [[PubMed](#)]
91. Ricci, F.; Bernasconi, S.; Porcu, L.; Erba, E.; Panini, N.; Fruscio, R.; Sina, F.; Torri, V.; Broggin, M.; Damia, G. ALDH enzymatic activity and CD133 positivity and response to chemotherapy in ovarian cancer patients. *Am. J. Cancer Res.* **2013**, *3*, 221–229. [[PubMed](#)]
92. Long, H.; Xiang, T.; Qi, W.; Huang, J.; Chen, J.; He, L.; Liang, Z.; Guo, B.; Li, Y.; Xie, R.; et al. CD133+ ovarian cancer stem-like cells promote non-stem cancer cell metastasis via CCL5 induced epithelial-mesenchymal transition. *Oncotarget* **2015**, *6*, 5846–5859. [[CrossRef](#)]
93. Gao, Y.; Foster, R.; Yang, X.; Feng, Y.; Shen, J.K.; Mankin, H.J.; Hornicek, F.J.; Amiji, M.M.; Duan, Z. Up-regulation of CD44 in the development of metastasis, recurrence and drug resistance of ovarian cancer. *Oncotarget* **2015**, *6*, 9313–9326. [[CrossRef](#)] [[PubMed](#)]
94. Liao, J.; Qian, F.; Tchabo, N.; Mhawech-Fauceglia, P.; Beck, A.; Qian, Z.; Wang, X.; Huss, W.J.; Lele, S.B.; Morrison, C.D.; et al. Ovarian cancer spheroid cells with stem cell-like properties contribute to tumor generation, metastasis and chemotherapy resistance through hypoxia-resistant metabolism. *PLoS ONE* **2014**, *9*, e84941. [[CrossRef](#)] [[PubMed](#)]
95. Landen, C.N., Jr.; Goodman, B.; Katre, A.A.; Steg, A.D.; Nick, A.M.; Stone, R.L.; Miller, L.D.; Mejia, P.V.; Jennings, N.B.; Gershenson, D.M.; et al. Targeting aldehyde dehydrogenase cancer stem cells in ovarian cancer. *Mol. Cancer Ther.* **2010**, *9*, 3186–3199. [[CrossRef](#)]
96. Meng, E.; Mitra, A.; Tripathi, K.; Finan, M.A.; Scalici, J.; McClellan, S.; Madeira da Silva, L.; Reed, E.; Shevde, L.A.; Palle, K.; et al. ALDH1A1 maintains ovarian cancer stem cell-like properties by altered regulation of cell cycle checkpoint and DNA repair network signaling. *PLoS ONE* **2014**, *9*, e107142. [[CrossRef](#)] [[PubMed](#)]
97. Silva, I.A.; Bai, S.; McLean, K.; Yang, K.; Griffith, K.; Thomas, D.; Ginestier, C.; Johnston, C.; Kueck, A.; Reynolds, R.K.; et al. Aldehyde dehydrogenase in combination with CD133 defines angiogenic ovarian cancer stem cells that portend poor patient survival. *Cancer Res.* **2011**, *71*, 3991–4001. [[CrossRef](#)]
98. Wang, L.; Mezencev, R.; Bowen, N.J.; Matyunina, L.V.; McDonald, J.F. Isolation and characterization of stem-like cells from a human ovarian cancer cell line. *Mol. Cell. Biochem.* **2012**, *363*, 257–268. [[CrossRef](#)]
99. Cheng, W.; Liu, T.; Wan, X.; Gao, Y.; Wang, H. MicroRNA-199a targets CD44 to suppress the tumorigenicity and multidrug resistance of ovarian cancer-initiating cells. *FEBS J.* **2012**, *279*, 2047–2059. [[CrossRef](#)] [[PubMed](#)]
100. Ohishi, T.; Koga, F.; Migita, T. Bladder Cancer Stem-Like Cells: Their Origin and Therapeutic Perspectives. *Int. J. Mol. Sci.* **2016**, *17*, 43. [[CrossRef](#)] [[PubMed](#)]
101. Hofner, T.; Macher-Goeppinger, S.; Klein, C.; Schillert, A.; Eisen, C.; Wagner, S.; Rigo-Watermeier, T.; Baccelli, I.; Vogel, V.; Trumpp, A.; et al. Expression and prognostic significance of cancer stem cell markers CD24 and CD44 in urothelial bladder cancer xenografts and patients undergoing radical cystectomy. *Urol. Oncol.* **2014**, *32*, 678–686. [[CrossRef](#)] [[PubMed](#)]

102. Li, Y.; Lin, K.; Yang, Z.; Han, N.; Quan, X.; Guo, X.; Li, C. Bladder cancer stem cells: Clonal origin and therapeutic perspectives. *Oncotarget* **2017**, *8*, 66668–66679. [[CrossRef](#)] [[PubMed](#)]
103. Shin, K.; Lee, J.; Guo, N.; Kim, J.; Lim, A.; Qu, L.; Mysorekar, I.U.; Beachy, P.A. Hedgehog/Wnt feedback supports regenerative proliferation of epithelial stem cells in bladder. *Nature* **2011**, *472*, 110–114. [[CrossRef](#)]
104. Shin, K.; Lim, A.; Odegaard, J.I.; Honeycutt, J.D.; Kawano, S.; Hsieh, M.H.; Beachy, P.A. Cellular origin of bladder neoplasia and tissue dynamics of its progression to invasive carcinoma. *Nat. Cell Biol.* **2014**, *16*, 469–478. [[CrossRef](#)] [[PubMed](#)]
105. Islam, S.S.; Mokhtari, R.B.; Noman, A.S.; Uddin, M.; Rahman, M.Z.; Azadi, M.A.; Zlotta, A.; van der Kwast, T.; Yeager, H.; Farhat, W.A. Sonic hedgehog (Shh) signaling promotes tumorigenicity and stemness via activation of epithelial-to-mesenchymal transition (EMT) in bladder cancer. *Mol. Carcinog.* **2016**, *55*, 537–551. [[CrossRef](#)]
106. Fang, D.; Kitamura, H. Cancer stem cells and epithelial-mesenchymal transition in urothelial carcinoma: Possible pathways and potential therapeutic approaches. *Int. J. Urol.* **2018**, *25*, 7–17. [[CrossRef](#)]
107. Moon, R.T.; Kohn, A.D.; De Ferrari, G.V.; Kaykas, A. WNT and beta-catenin signalling: Diseases and therapies. *Nat. Rev. Genet.* **2004**, *5*, 691–701. [[CrossRef](#)]
108. Pierzynski, J.A.; Hildebrandt, M.A.; Kamat, A.M.; Lin, J.; Ye, Y.; Dinney, C.P.; Wu, X. Genetic Variants in the Wnt/beta-Catenin Signaling Pathway as Indicators of Bladder Cancer Risk. *J. Urol.* **2015**, *194*, 1771–1776. [[CrossRef](#)]
109. Chen, X.; Jia, C.; Jin, X.; Gu, X. MicroRNA-374a Inhibits Aggressive Tumor Biological Behavior in Bladder Carcinoma by Suppressing Wnt/beta-Catenin Signaling. *Cell. Physiol. Biochem. Int. J. Exp. Cell. Physiol. Biochem. Pharmacol.* **2018**, *48*, 815–826. [[CrossRef](#)]
110. Lobry, C.; Oh, P.; Aifantis, I. Oncogenic and tumor suppressor functions of Notch in cancer: it's NOTCH what you think. *J. Exp. Med.* **2011**, *208*, 1931–1935. [[CrossRef](#)] [[PubMed](#)]
111. Pannuti, A.; Foreman, K.; Rizzo, P.; Osipo, C.; Golde, T.; Osborne, B.; Miele, L. Targeting Notch to target cancer stem cells. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **2010**, *16*, 3141–3152. [[CrossRef](#)]
112. Goriki, A.; Seiler, R.; Wyatt, A.W.; Contreras-Sanz, A.; Bhat, A.; Matsubara, A.; Hayashi, T.; Black, P.C. Unravelling disparate roles of NOTCH in bladder cancer. *Nat. Rev. Urol.* **2018**, *15*, 345–357. [[CrossRef](#)] [[PubMed](#)]
113. Hayashi, T.; Gust, K.M.; Wyatt, A.W.; Goriki, A.; Jager, W.; Awrey, S.; Li, N.; Oo, H.Z.; Altamirano-Dimas, M.; Buttyan, R.; et al. Not all NOTCH Is Created Equal: The Oncogenic Role of NOTCH2 in Bladder Cancer and Its Implications for Targeted Therapy. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **2016**, *22*, 2981–2992. [[CrossRef](#)] [[PubMed](#)]
114. Prince, M.E.; Sivanandan, R.; Kaczorowski, A.; Wolf, G.T.; Kaplan, M.J.; Dalerba, P.; Weissman, I.L.; Clarke, M.F.; Ailles, L.E. Identification of a subpopulation of cells with cancer stem cell properties in head and neck squamous cell carcinoma. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 973–978. [[CrossRef](#)] [[PubMed](#)]
115. Han, J.; Fujisawa, T.; Husain, S.R.; Puri, R.K. Identification and characterization of cancer stem cells in human head and neck squamous cell carcinoma. *BMC Cancer* **2014**, *14*, 173. [[CrossRef](#)] [[PubMed](#)]
116. Felthaus, O.; Ettl, T.; Gosau, M.; Driemel, O.; Brockhoff, G.; Reck, A.; Zeitler, K.; Hautmann, M.; Reichert, T.E.; Schmalz, G.; et al. Cancer stem cell-like cells from a single cell of oral squamous carcinoma cell lines. *Biochem. Biophys. Res. Commun.* **2011**, *407*, 28–33. [[CrossRef](#)] [[PubMed](#)]
117. Reid, P.A.; Wilson, P.; Li, Y.; Marcu, L.G.; Bezak, E. Current understanding of cancer stem cells: Review of their radiobiology and role in head and neck cancers. *Head Neck* **2017**, *39*, 1920–1932. [[CrossRef](#)] [[PubMed](#)]
118. Fukusumi, T.; Ishii, H.; Konno, M.; Yasui, T.; Nakahara, S.; Takenaka, Y.; Yamamoto, Y.; Nishikawa, S.; Kano, Y.; Ogawa, H.; et al. CD10 as a novel marker of therapeutic resistance and cancer stem cells in head and neck squamous cell carcinoma. *Br. J. Cancer* **2014**, *111*, 506–514. [[CrossRef](#)] [[PubMed](#)]
119. Luo, W.; Li, S.; Peng, B.; Ye, Y.; Deng, X.; Yao, K. Embryonic stem cells markers SOX2, OCT4 and Nanog expression and their correlations with epithelial-mesenchymal transition in nasopharyngeal carcinoma. *PLoS ONE* **2013**, *8*, e56324. [[CrossRef](#)]
120. Chien, C.S.; Wang, M.L.; Chu, P.Y.; Chang, Y.L.; Liu, W.H.; Yu, C.C.; Lan, Y.T.; Huang, P.I.; Lee, Y.Y.; Chen, Y.W.; et al. Lin28B/Let-7 Regulates Expression of Oct4 and Sox2 and Reprograms Oral Squamous Cell Carcinoma Cells to a Stem-like State. *Cancer Res.* **2015**, *75*, 2553–2565. [[CrossRef](#)]
121. Chen, Y.C.; Chen, Y.W.; Hsu, H.S.; Tseng, L.M.; Huang, P.I.; Lu, K.H.; Chen, D.T.; Tai, L.K.; Yung, M.C.; Chang, S.C.; et al. Aldehyde dehydrogenase 1 is a putative marker for cancer stem cells in head and neck squamous cancer. *Biochem. Biophys. Res. Commun.* **2009**, *385*, 307–313. [[CrossRef](#)]



122. Allegra, E.; Trapasso, S.; Pisani, D.; Puzzo, L. The role of BMI1 as a biomarker of cancer stem cells in head and neck cancer: A review. *Oncology* **2014**, *86*, 199–205. [[CrossRef](#)]
123. Chen, H.; Zhou, L.; Dou, T.; Wan, G.; Tang, H.; Tian, J. BMI1'S maintenance of the proliferative capacity of laryngeal cancer stem cells. *Head Neck* **2011**, *33*, 1115–1125. [[CrossRef](#)]
124. Ma, L.; Wang, H.; Yao, H.; Zhu, L.; Liu, W.; Zhou, Z. Bmi1 expression in oral lichen planus and the risk of progression to oral squamous cell carcinoma. *Ann. Diagn. Pathol.* **2013**, *17*, 327–330. [[CrossRef](#)]
125. Feng, J.Q.; Xu, Z.Y.; Shi, L.J.; Wu, L.; Liu, W.; Zhou, Z.T. Expression of cancer stem cell markers ALDH1 and Bmi1 in oral erythroplakia and the risk of oral cancer. *J. Oral Pathol. Med. Off. Publ. Int. Assoc. Oral Pathol. Am. Acad. Oral Pathol.* **2013**, *42*, 148–153. [[CrossRef](#)] [[PubMed](#)]
126. Vormittag, L.; Thurnher, D.; Geleff, S.; Pammer, J.; Heiduschka, G.; Brunner, M.; Grasl, M.; Erovic, B.M. Co-expression of Bmi-1 and podoplanin predicts overall survival in patients with squamous cell carcinoma of the head and neck treated with radio(chemo)therapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2009**, *73*, 913–918. [[CrossRef](#)] [[PubMed](#)]
127. Chen, H.; Zhou, L.; Wan, G.; Dou, T.; Tian, J. BMI1 promotes the progression of laryngeal squamous cell carcinoma. *Oral Oncol.* **2011**, *47*, 472–481. [[CrossRef](#)]
128. Abdelhamed, S.; Ogura, K.; Yokoyama, S.; Saiki, I.; Hayakawa, Y. AKT-STAT3 Pathway as a Downstream Target of EGFR Signaling to Regulate PD-L1 Expression on NSCLC cells. *J. Cancer* **2016**, *7*, 1579–1586. [[CrossRef](#)] [[PubMed](#)]
129. Johnson, R.; Halder, G. The two faces of Hippo: Targeting the Hippo pathway for regenerative medicine and cancer treatment. *Nat. Rev. Drug Discov.* **2014**, *13*, 63–79. [[CrossRef](#)]
130. Nishio, M.; Maehama, T.; Goto, H.; Nakatani, K.; Kato, W.; Omori, H.; Miyachi, Y.; Togashi, H.; Shimono, Y.; Suzuki, A. Hippo vs. Crab: Tissue-specific functions of the mammalian Hippo pathway. *Genes Cells Devoted Mol. Cell. Mech.* **2017**, *22*, 6–31. [[CrossRef](#)] [[PubMed](#)]
131. Pan, D. The hippo signaling pathway in development and cancer. *Dev. Cell* **2010**, *19*, 491–505. [[CrossRef](#)]
132. Staley, B.K.; Irvine, K.D. Hippo signaling in Drosophila: Recent advances and insights. *Dev. Dyn. Off. Publ. Am. Assoc. Anat.* **2012**, *241*, 3–15. [[CrossRef](#)]
133. Zhang, M.; Zeng, J.; Zhao, Z.; Liu, Z. Loss of miR-424-3p, not miR-424-5p, confers chemoresistance through targeting YAP1 in non-small cell lung cancer. *Mol. Carcinog.* **2017**, *56*, 821–832. [[CrossRef](#)] [[PubMed](#)]
134. Zanconato, F.; Cordenonsi, M.; Piccolo, S. YAP/TAZ at the Roots of Cancer. *Cancer Cell* **2016**, *29*, 783–803. [[CrossRef](#)] [[PubMed](#)]
135. Garcia-Prat, L.; Martinez-Vicente, M.; Perdiguero, E.; Ortet, L.; Rodriguez-Ubreva, J.; Rebollo, E.; Ruiz-Bonilla, V.; Gutarra, S.; Ballestar, E.; Serrano, A.L.; et al. Autophagy maintains stemness by preventing senescence. *Nature* **2016**, *529*, 37–42. [[CrossRef](#)]
136. Song, Q.; Mao, B.; Cheng, J.; Gao, Y.; Jiang, K.; Chen, J.; Yuan, Z.; Meng, S. YAP enhances autophagic flux to promote breast cancer cell survival in response to nutrient deprivation. *PLoS ONE* **2015**, *10*, e0120790. [[CrossRef](#)]
137. Bora-Singhal, N.; Nguyen, J.; Schaal, C.; Perumal, D.; Singh, S.; Coppola, D.; Chellappan, S. YAP1 Regulates OCT4 Activity and SOX2 Expression to Facilitate Self-Renewal and Vascular Mimicry of Stem-Like Cells. *Stem Cells* **2015**, *33*, 1705–1718. [[CrossRef](#)]
138. Xia, Y.; Zhang, Y.L.; Yu, C.; Chang, T.; Fan, H.Y. YAP/TEAD co-activator regulated pluripotency and chemoresistance in ovarian cancer initiated cells. *PLoS ONE* **2014**, *9*, e109575. [[CrossRef](#)]
139. Kim, G.J.; Kim, H.; Park, Y.N. Increased expression of Yes-associated protein 1 in hepatocellular carcinoma with stemness and combined hepatocellular-cholangiocarcinoma. *PLoS ONE* **2013**, *8*, e75449. [[CrossRef](#)]
140. Wang, L.; Zhang, Z.; Yu, X.; Huang, X.; Liu, Z.; Chai, Y.; Yang, L.; Wang, Q.; Li, M.; Zhao, J.; et al. Unbalanced YAP-SOX9 circuit drives stemness and malignant progression in esophageal squamous cell carcinoma. *Oncogene* **2019**, *38*, 2042–2055. [[CrossRef](#)]
141. Zhu, P.; Wang, Y.; Wu, J.; Huang, G.; Liu, B.; Ye, B.; Du, Y.; Gao, G.; Tian, Y.; He, L.; et al. LncBRM initiates YAP1 signalling activation to drive self-renewal of liver cancer stem cells. *Nat. Commun.* **2016**, *7*, 13608. [[CrossRef](#)] [[PubMed](#)]
142. Zhang, K.; Qi, H.X.; Hu, Z.M.; Chang, Y.N.; Shi, Z.M.; Han, X.H.; Han, Y.W.; Zhang, R.X.; Zhang, Z.; Chen, T.; et al. YAP and TAZ Take Center Stage in Cancer. *Biochemistry* **2015**, *54*, 6555–6566. [[CrossRef](#)] [[PubMed](#)]
143. Maugeri-Sacca, M.; De Maria, R. Hippo pathway and breast cancer stem cells. *Crit. Rev. Oncol. Hematol.* **2016**, *99*, 115–122. [[CrossRef](#)] [[PubMed](#)]

144. Azzolin, L.; Panciera, T.; Soligo, S.; Enzo, E.; Bicciato, S.; Dupont, S.; Bresolin, S.; Frasson, C.; Basso, G.; Guzzardo, V.; et al. YAP/TAZ incorporation in the beta-catenin destruction complex orchestrates the Wnt response. *Cell* **2014**, *158*, 157–170. [[CrossRef](#)]
145. You, E.; Huh, Y.H.; Kwon, A.; Kim, S.H.; Chae, I.H.; Lee, O.J.; Ryu, J.H.; Park, M.H.; Kim, G.E.; Lee, J.S.; et al. SPIN90 Depletion and Microtubule Acetylation Mediate Stromal Fibroblast Activation in Breast Cancer Progression. *Cancer Res.* **2017**, *77*, 4710–4722. [[CrossRef](#)]
146. Fan, Y.; Gao, Y.; Rao, J.; Wang, K.; Zhang, F.; Zhang, C. YAP-1 Promotes Tregs Differentiation in Hepatocellular Carcinoma by Enhancing TGFBR2 Transcription. *Cell. Physiol. Biochem.* **2017**, *41*, 1189–1198. [[CrossRef](#)]
147. Oku, Y.; Nishiya, N.; Shito, T.; Yamamoto, R.; Yamamoto, Y.; Oyama, C.; Uehara, Y. Small molecules inhibiting the nuclear localization of YAP/TAZ for chemotherapeutics and chemosensitizers against breast cancers. *FEBS Open Bio* **2015**, *5*, 542–549. [[CrossRef](#)]
148. Gibault, F.; Bailly, F.; Corvaisier, M.; Coevoet, M.; Huet, G.; Melnyk, P.; Cotellet, P. Molecular Features of the YAP Inhibitor Verteporfin: Synthesis of Hexasubstituted Dipyrroles as Potential Inhibitors of YAP/TAZ, the Downstream Effectors of the Hippo Pathway. *ChemMedChem* **2017**, *12*, 954–961. [[CrossRef](#)]
149. Liu-Chittenden, Y.; Huang, B.; Shim, J.S.; Chen, Q.; Lee, S.J.; Anders, R.A.; Liu, J.O.; Pan, D. Genetic and pharmacological disruption of the TEAD-YAP complex suppresses the oncogenic activity of YAP. *Genes Dev.* **2012**, *26*, 1300–1305. [[CrossRef](#)] [[PubMed](#)]
150. Zhang, H.; Ramakrishnan, S.K.; Triner, D.; Centofanti, B.; Maitra, D.; Gyorffy, B.; Sebolt-Leopold, J.S.; Dame, M.K.; Varani, J.; Brenner, D.E.; et al. Tumor-selective proteotoxicity of verteporfin inhibits colon cancer progression independently of YAP1. *Sci. Signal.* **2015**, *8*, ra98. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).