

Temperaturejump problem with arbitrary accommodation

S. K. Loyalka, C. E. Siewert, and J. R. Thomas Jr.

Citation: *Physics of Fluids (1958-1988)* **21**, 854 (1978); doi: 10.1063/1.862306

View online: <http://dx.doi.org/10.1063/1.862306>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/pof1/21/5?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[The temperature-jump problem for a variable collision frequency model](#)

Phys. Fluids **14**, 382 (2002); 10.1063/1.1416192

[Temperature slip problem with arbitrary accommodation at the surface](#)

Phys. Fluids **16**, 1162 (1973); 10.1063/1.1694481

[A Nanosecond TemperatureJump Apparatus](#)

Rev. Sci. Instrum. **42**, 1643 (1971); 10.1063/1.1684958

[On the Determination of Thermal Accommodation Coefficients in the TemperatureJump Region](#)

J. Chem. Phys. **46**, 3217 (1967); 10.1063/1.1841192

[Method for Determining Accommodation Coefficients from Data in the TemperatureJump Range without Applying TemperatureJump Theory](#)

J. Chem. Phys. **42**, 1850 (1965); 10.1063/1.1696210

An advertisement for physicist jobs. On the left is a photograph of a man in a dark suit and striped tie, looking surprised with his hand to his ear. To the right, the text 'HAVE YOU HEARD?' is written in large, bold, dark red letters. Below this, in smaller dark red letters, is 'Employers hiring scientists and engineers trust'. Underneath that, 'physicstodayJOBS' is written in a blue, sans-serif font. To the right of the text is a square QR code. At the bottom, the URL 'http://careers.physicstoday.org/post.cfm' is displayed in a small, black, sans-serif font.

HAVE YOU HEARD?

Employers hiring scientists
and engineers trust
physicstodayJOBS

<http://careers.physicstoday.org/post.cfm>

Temperature-jump problem with arbitrary accommodation

S. K. Loyalka

Nuclear Engineering Department, University of Missouri-Columbia, Columbia, Missouri 65201

C. E. Siewert^{a)}

Laboratoire d'Optique Atmospherique, Université des Sciences et Techniques de Lille, France

J. R. Thomas, Jr.

Nuclear Engineering Group, Mechanical Engineering Department, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

(Received 9 August 1977; final manuscript received 10 February 1978)

A concise and accurate result for the temperature-jump coefficient based on the linearized BGK model and arbitrary accommodation is reported. The jump coefficient is expressed as a power series in $(1-\alpha)$, and values of the expansion coefficients are given.

We wish to give a rearranged version of some previous results¹ for the temperature-jump coefficient. As discussed earlier¹ the temperature-jump coefficient can be expressed as

$$\epsilon' = \left(\frac{2-\alpha}{\alpha} \right) \frac{5\sqrt{\pi}}{8} \epsilon_0 + \Delta(\alpha), \quad (1)$$

where α is the accommodation coefficient for the linearized BGK model,

$$\epsilon_0 = \frac{8}{5\sqrt{\pi}} \begin{vmatrix} 1 \\ 0 \end{vmatrix}^t \mathbf{H}_1^{-t} \mathbf{H}_2^t \begin{vmatrix} 1 \\ -\sqrt{\frac{2}{3}} \end{vmatrix}, \quad (2)$$

and

$$\Delta(\alpha) = \frac{(1-\alpha)}{\alpha\sqrt{\pi}} \begin{vmatrix} 1 \\ 0 \end{vmatrix}^t \mathbf{H}_1^{-t} \int_0^\infty \mathbf{H}^{-1}(\eta) \begin{vmatrix} 1 & 0 \\ 0 & \sqrt{\frac{2}{3}} \end{vmatrix} \times \mathbf{A}(\eta) \eta \exp(-\eta^2) d\eta. \quad (3)$$

Here, $\mathbf{H}(\mu)$ is the 2×2 \mathbf{H} matrix introduced by Kriese *et al.*,²

$$\mathbf{H}_B^t = (\pi)^{-1/2} \int_0^\infty \mathbf{H}^t(\mu) \mathbf{Q}^t(\mu) \mathbf{Q}(\mu) \mu^B \exp(-\mu^2) d\mu, \quad (4)$$

$$\mathbf{Q}(\mu) = \begin{vmatrix} \sqrt{\frac{2}{3}}(\mu^2 - \frac{1}{2}) & 1 \\ \sqrt{\frac{2}{3}} & 0 \end{vmatrix}, \quad (5)$$

and the vector $\mathbf{A}(\eta)$ is the solution of the Fredholm equation

$$\mathbf{A}(\eta) = (\alpha - 2) \mathbf{F}(\eta) + (1 - \alpha) (\mathbf{T}\mathbf{A})(\eta), \quad \eta > 0. \quad (6)$$

In addition, we use the superscript t to denote the transpose operation and the superscript $-t$ to denote the transpose-inverse operation. In Eq. (6)

$$\mathbf{F}(\eta) = \eta \exp(-\eta^2) \mathbf{R}(\eta) \mathbf{H}^{-t}(\eta) \mathbf{H}_1^t \begin{vmatrix} \sqrt{\frac{3}{2}} \\ -1 \end{vmatrix}, \quad (7)$$

where

$$\mathbf{R}(\eta) = \frac{1}{N(\eta)} \begin{vmatrix} \sqrt{\frac{3}{2}} N_{22}(\eta) & -N_{12}(\eta) \\ -\sqrt{\frac{3}{2}} N_{12}(\eta) & N_{11}(\eta) \end{vmatrix}. \quad (8)$$

We note that Kriese *et al.*² have given $N_{ij}(\eta)$ and $N(\eta)$ in terms of Dawson's integral. The operator \mathbf{T} in Eq. (6) is such that

$$(\mathbf{T}\mathbf{X})(\eta) = \frac{\eta \exp(-\eta^2)}{\sqrt{\pi}} \mathbf{R}(\eta) \mathbf{H}^{-t}(\eta) \times \int_0^\infty \mathbf{H}^{-1}(\mu) \begin{vmatrix} \sqrt{\frac{3}{2}} & 0 \\ 0 & 1 \end{vmatrix} \mathbf{X}(\mu) \exp(-\mu^2) \mu \frac{d\mu}{\mu + \eta}. \quad (9)$$

We have computed the Hilbert-Schmidt³ norm of the kernel \mathbf{K} of the operator \mathbf{T} , where

$$\mathbf{K}(\eta, \mu) = \frac{\eta \exp(-\eta^2)}{\sqrt{\pi}} \mathbf{R}(\eta) \mathbf{H}^{-t}(\eta) \times \mathbf{H}^{-1}(\mu) \begin{vmatrix} \sqrt{\frac{3}{2}} & 0 \\ 0 & 1 \end{vmatrix} \frac{\mu \exp(-\mu^2)}{\eta + \mu}. \quad (10)$$

We used the definition

$$\|\mathbf{T}\| = \left\{ \int_0^\infty d\eta \int_0^\infty d\mu \operatorname{tr}[\mathbf{K}^t(\mu, \eta) \mathbf{K}(\mu, \eta)] \right\}^{1/2}, \quad (11)$$

where

$$\operatorname{tr}[\mathbf{K}^t \mathbf{K}] = K_{11}^2 + K_{12}^2 + K_{21}^2 + K_{22}^2. \quad (12)$$

We find that $\|\mathbf{T}\| \approx 0.04$; that is, for

$$|(1 - \alpha)| \lesssim (0.04)^{-1},$$

or $\alpha \lesssim 25$, it should be possible to obtain a convergent Neumann-Liouville iterative solution of Eq. (6). In fact for physically interesting cases $\alpha \leq 1$ and in this range, one can expect a very rapidly convergent iterative solution.

Thus, we now express an iterative solution of Eq. (6) as

$$\mathbf{A}(\eta) = (\alpha - 2) \left[\mathbf{F}(\eta) + \sum_{m=1}^{\infty} (1 - \alpha)^m (\mathbf{T}^m \mathbf{F})(\eta) \right]; \quad (13)$$

then, we can write the jump coefficient as

$$\epsilon' = \frac{5\sqrt{\pi}}{8} \left(\frac{2 - \alpha}{\alpha} \right) \left[\epsilon_0 + \sum_{m=1}^{\infty} (1 - \alpha)^m \epsilon_m \right], \quad (14)$$

where

$$\epsilon_m = -\frac{1}{\sqrt{\pi}} \frac{8}{5\sqrt{\pi}} \begin{vmatrix} 1 \\ 0 \end{vmatrix}^t \mathbf{H}^{-t} \mathbf{W}_{m-1} \mathbf{H}_1^t \begin{vmatrix} \sqrt{\frac{2}{3}} \\ -1 \end{vmatrix}, \quad m > 0, \quad (15)$$

$$\mathbf{W}_m = \int_0^{\infty} \mathbf{H}^{-1}(\eta) \begin{vmatrix} 1 & 0 \\ 0 & \sqrt{\frac{2}{3}} \end{vmatrix} (\mathbf{T}^m \mathbf{B})(\eta) \eta \exp(-\eta^2) d\eta, \quad (16)$$

and

$$\mathbf{B}(\eta) = \eta \exp(-\eta^2) \mathbf{R}(\eta) \mathbf{H}^{-t}(\eta). \quad (17)$$

Equation (11) is a convenient result because the ϵ_m do not depend on α . We have evaluated Eqs. (2) and (15) numerically to find the results given in Table I. A Gaussian quadrature scheme was used to evaluate all integrals, and the number of quadrature points was increased until no change in Table I was observed. The final calculation utilized 80 Gauss-Legendre points⁴ in the interval (0, 1), which was mapped onto (0, ∞) according to the transformation $\mu = \mu'/(1 - \mu')$. The accuracy of the present results to the significant digits reported here is verified by the fact that Eqs. (15) give,

$$\sum_{i=0}^{10} \epsilon_i = 1.000000;$$

that is, in the limit $\alpha \rightarrow 0$, from Eq. (11) we get

$$\epsilon' = \frac{5\sqrt{\pi}}{8} \left(\frac{2 - \alpha}{\alpha} \right).$$

TABLE I. Numerical values for ϵ_m as computed from Eq. (15).

m	ϵ_m
0	1.17597
1	-1.60683×10^{-1}
2	-1.37349×10^{-2}
3	-1.38665×10^{-3}
4	-1.44586×10^{-4}
5	-1.52085×10^{-5}
6	-1.60481×10^{-6}
7	-1.69582×10^{-7}
8	-1.79329×10^{-8}
9	-1.89710×10^{-9}
10	-2.00735×10^{-10}

which is an exact result in this limit. Since in view of the expansion in $(1 - \alpha)$ one would expect the maximum numerical error at $\alpha \rightarrow 0$, and since in this limit our result is exact to the number of significant digits quoted, it is clear that the present series expansion should provide exact values of ϵ' to the number of significant digits quoted here.

One of the authors (CES) is grateful to J. Lenoble and the Université des Sciences et Techniques de Lille for their kind hospitality and to the Délégation Générale à la Recherche Scientifique et Technique for partial support of this work. Another (JRT) acknowledges a very helpful discussion with J. Ball.

^{a)}Permanent address: Nuclear Engineering Department, North Carolina State University, Raleigh, N.C. 27607.

¹J. R. Thomas, Jr., *Phys. Fluids* **16**, 1162 (1973).

²J. T. Kriese, T. S. Chang, and C. E. Siewert, *Int. J. Eng. Sci.* **12**, 441 (1974).

³F. Riesz and B. Sz-Nagy, *Functional Analysis* [(Translated from the 2nd French edition by L. Boron) Ungar, New York, 1955].

⁴A. H. Stroud and D. Secrest, *Gaussian Quadrature Formulas* (Prentice-Hall, Englewood Cliffs, N.J., 1966).