

Temperature-Sensitive Transient Receptor Potential Channels in Corneal Tissue Layers and Cells

Stefan Mergler^a Monika Valtink^b Sumioka Takayoshi^c Yuka Okada^c
Masayasu Miyajima^c Shizuya Saika^c Peter S. Reinach^d

^aDepartment of Ophthalmology, Charité – Universitätsmedizin Berlin, Berlin, and ^bInstitute of Anatomy, Faculty of Medicine Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany; ^cDepartment of Ophthalmology, Wakayama Medical University School of Medicine, Wakayama, Japan; ^dDepartment of Ophthalmology and Optometry, Wenzhou Medical University, Wenzhou, PR China

Key Words

Thermo-transient receptor potential channels ·
Electrophysiology · Calcium · Cornea · Inflammation ·
Fibrosis · Wound healing

Abstract

We here provide a brief summary of the characteristics of transient receptor potential channels (TRPs) identified in corneal tissue layers and cells. In general, TRPs are nonselective cation channels which are Ca²⁺ permeable. Most TRPs serve as thermosensitive molecular sensors (thermo-TRPs). Based on their functional importance, the possibilities are described for drug-targeting TRP activity in a clinical setting. TRPs are expressed in various tissues of the eye including both human corneal epithelial and endothelial layers as well as stromal fibroblasts and stromal nerve fibers. TRP vanilloid type 1 (TRPV1) heat receptor, also known as capsaicin receptor, along with TRP melastatin type 8 (TRPM8) cold receptor, which is also known as menthol receptor, are prototypes of the thermo-TRP family. The TRPV1 functional channel is the most investigated TRP channel in these tissues, owing to its contribution to maintaining tissue homeostasis as well as eliciting wound healing responses to injury. Other thermo-TRP family members identified in these tissues are TRPV2, 3

and 4. Finally, there is the TRP ankyrin type 1 (TRPA1) cold receptor. All of these thermo-TRPs can be activated within specific temperature ranges and transduce such inputs into chemical and electrical signals. Although several recent studies have begun to unravel complex roles for thermo-TRPs such as TRPV1 in corneal layers and resident cells, additional studies are needed to further elucidate their roles in health and disease.

© 2014 S. Karger AG, Basel

Introduction

Temperature-sensitive transient receptor potential channels (thermo-TRPs) are a superfamily which is subdivided into 7 groups according to differences in amino acid sequence homology. Specifically, TRPs are subdivided into canonical receptors (TRPC1–6), vanilloid receptors (TRPV1–7), melastatin receptors (TRPM1–8), ankyrin receptor (TRPA1), polycystins (TRPP1–5) and mucolipins (TRPML1–3). Another TRP subfamily is assigned to mechanosensory channels (TRPN), which are not expressed in mammals. TRPV1–4, TRPM8 and TRPA1 belong to the thermo-TRP family. Modulation of thermo-TRP activity by environmental thermal challenges is essential for tissues

Table 1. Characterization of TRP activators, inhibitors and tissue localization

	Selectivity, P _{Ca} :P _{Na}	Activation threshold temperature, °C	Pharmacology	Function	Cornea
TRPV1	4–10	>43	capsaicin, capsazepine	heat sensor osmosensor ¹	epithelium [8, 43], stroma [29], endothelium [9]
TRPV2	1–3	>52	2-APB	heat sensor	endothelium [9]
TRPV3	2.6	30–39	camphor, 2-APB	moderate heat sensor	epithelium [20], endothelium [9]
TRPV4	6–10	24–27	4 α -PDD, GSK1016790A	moderate heat sensor, osmosensor ²	epithelium [19], endothelium [28]
TRPV5	>100	–	low [Ca ²⁺] _i	Ca ²⁺ reabsorption in kidney	–
TRPV6	>100	–	low [Ca ²⁺] _i	–	–
TRPM8	3.3	<23–28	menthol, icilin, eucalyptol	moderate cold sensor	epithelium ³ , nerve fibers [36], stroma ³ , endothelium [24, 26]
TRPA1	0.8	<17	icilin	cold sensor	endothelium [24, 26]

P = Permeability ratio; 2-APB = 2-aminoethoxydiphenyl borate; [Ca²⁺]_i = intracellular Ca²⁺. ¹ Activation by hypertonic challenge. ² Activation by hypotonic challenge. ³ Mergler et al. [unpubl. data].

to offset declines in tissue function that result from thermal exposure [1]. Such changes in different TRP subtypes are elicited for the most part through transient increases in intracellular Ca²⁺ influx as a consequence of channel activation. These rises in turn activate downstream-linked Ca²⁺-sensitive signaling pathways. The magnitude and duration of the Ca²⁺ influx encodes downstream signaling responses that induce either a homeostatic or lethal response. Therefore, identifying which TRP subtypes respond to physiological and pathogenic stresses imposed on corneal tissue can aid in identifying novel drug targets.

This insight can then be utilized to target specific subtypes with drugs or genetic techniques to reverse losses in corneal function which are caused by maladaptive TRP function. At this point, it is increasingly apparent that thermo-TRPs play an important role in eliciting such control. Okada, Reinach and colleagues [2, 3] have recently shown in different studies that inhibition of both TRPV1 and TRPA1 activation after an alkali burn markedly reduces corneal fibrosis inflammation and opacification. On the other hand, TRPV1 activation hastens corneal epithelial healing following epithelial debridement, suggesting that increasing TRPV1 activity in a clinical setting could decrease the likelihood of corneal infection.

Although there are some relevant indications of putative roles of thermo-TRPs in corneal cells, there is scant information about the cellular mechanisms modulating TRP channel activation and deactivation in corneal tissues and cells. This realization has prompted efforts to fill this void, which is needed to develop drugs or gene expression protocols that can selectively modulate TRP behavior without having side effects.

This review describes the functional importance of thermo-TRPs in corneal tissues and cells in maintaining corneal homeostasis. Table 1 provides a summary listing corneal TRP isotype characteristics, functional roles and cell type localization. We also present evidence that TRP malfunction can contribute to the underlying pathophysiology of inappropriate wound healing responses to corneal injury. Finally, the possible translational relevance of the insight gained from this characterization is discussed.

Corneal Epithelium

In the last decade, different types of TRP isotype functional expression were identified on corneal epithelial nerves and also on corneal epithelial cell membranes

(fig. 1) [4]. TRPC4 is a thermally insensitive member of the TRP canonical subfamily which can be activated by intracellular Ca^{2+} store depletion induced by epidermal growth factor (EGF) [5]. TRPC4 activation occurs as a consequence of Ca^{2+} store emptying, which leads to increases in store operated Ca^{2+} channel (SOCC) activity. It is uncertain whether this TRP subtype is the sole component of SOCCs or whether it is a contributor to its function. This uncertainty exists because TRPs are composed of tetrameric subunits, which can form both homomeric and heteromeric configurations. A homomeric TRP is made up of 4 subunits from one subfamily member, whereas a heteromeric TRP counterpart can have subunits derived from different members of a subfamily. These alternatives extend the complexity of delineating the functional roles of TRPs, since the regulation of heteromeric channels can differ from homomeric channels. In any case, in this study, it was shown that SOCC activation leads to capacitative calcium entry, which refills intracellular stores of their depleted Ca^{2+} content [6]. TRPC4 involvement in capacitative calcium entry is apparent, since in TRPC4 siRNA gene-silenced cells, EGF-induced SOCC activation and proliferation were suppressed [5]. TRPC4 expression was also identified in bovine corneal endothelial cells and is also involved in regulating SOCC activity [7].

The vanilloid TRP protein isoform TRPV1 is functionally expressed in both human corneal epithelial cells and corneal endothelial cells [8, 9]. It is activated in nonneural cells by numerous stimuli including heat, voltage, vanilloids, cannabinoids, lipids and protons/cations [10, 11]. In this context, complex regulatory mechanisms are involved in encoding specific responses through Ca^{2+} signaling [12]. Since TRPV1 activity is to some extent voltage dependent [13], membrane depolarization by TRPV1 channels suppressed Ca^{2+} entry through calcium release-activated calcium channel protein 1 (ORAI1), a pore-forming transmembrane protein whose activation elicits SOCC activation in nonexcitable cells (fig. 2). Specifically, ORAI1, which is encoded by the *ORAI1* gene, elicits the calcium release-activated Ca^{2+} current when activated by Ca^{2+} store depletion [14]. The process of ORAI1 channel activation is thought to depend on relocation to the plasma membrane of intracellular store membrane-associated stromal interaction molecule 1 (STIM1). This protein senses the Ca^{2+} filling state of intracellular stores and moves to open SOCCs by interacting with ORAI1 upon intracellular Ca^{2+} store depletion. The contribution by other SOCCs such as TRPC1 still remains to be elucidated. However, the role so far of TRPV1 in connection with ORAI1 has not been investigated in corneal tissues and cells.




Fig. 1. Histological structure of the human cornea. Vertical section of the human cornea. The corneal endothelium, consisting of only one cell layer, is located at the posterior cornea, facing the anterior chamber of the eye. The corneal epithelium is located at the outside of the cornea, facing the air. In between, the corneal stroma is located. HE. $\times 100$.




Fig. 2. Schematic representation of TRPV1 (capsaicin receptor). TRPV1 modulates driving forces for Ca^{2+} entry. It is an ion channel and a polymodal receptor for various inflammatory stimuli, hypertonicity, protons, heat and lipid metabolites. TRPV1 possesses endogenous activators such as endocannabinoids (anandamide, NADA). TRPV1 can also be activated by heat (>43°C) and is permeable for monovalent and divalent cations like other unspecific cation channels. Notably, membrane depolarization ($V_m \uparrow$) by TRPV1 channels results in a reduced Ca^{2+} entry ($\text{Ca}^{2+} \downarrow$) via ORAI. This Ca^{2+} process accordingly modulates TRPV1 function to modify the intracellular Ca^{2+} content.

Another characteristic of TRPs is that they are able to modulate driving forces for Ca^{2+} entry. On the one hand, membrane depolarization by TRPs results in a reduced Ca^{2+} entry through ORAI, as mentioned before. On the other hand, hyperpolarization of the membrane by big potassium or maxi-K channels or small conductance calcium-activated potassium channels results in an increased Ca^{2+} influx [12]. This Ca^{2+} change subsequently modulates TRP and big potassium or maxi-K channel as well as small conductance calcium-activated potassium channel function to fine-tune the intracellular Ca^{2+} concentration. Irrespective of K^+ channel involvement, TRPV1 activation via capsaicin initially leads to increases in intracellular Ca^{2+} levels. Their rises can induce inflammatory cytokine release through mitogen-activated protein kinase (MAPK) signaling in corneal epithelial cells [8]. TRPV1 receptors play a role in mediating corneal epithelial and stromal cell inflammatory mediator infiltration and subsequent hyperalgesia based on results obtained with a mouse corneal wound healing model [3]. In this study, it was shown that in homozygous TRPV1 knockout mice, inflammation and fibrosis were clearly suppressed relative to what occurred in their wild-type counterpart [3]. A similar favorable outcome was obtained in wild-type mice systemically treated with a TRPV1 antagonist subsequent to alkali having burnt their corneas. Therefore, TRPV1 is a potential drug target for improving the outcome of inflammatory/fibrogenic wound healing [3]. Regarding TRPV1-mediated Ca^{2+} signaling, Yang et al. [15] recently showed that activation of a G protein-coupled receptor such as the cannabinoid receptor type 1 (CB1) suppressed TRPV1-induced inflammatory responses to corneal epithelial debridement. A suggested Ca^{2+} signal transduction pathway activated by a CB1 agonist, WIN55,212-2 (WIN), is shown in figure 3. CB1 activation by injury induced a release of endogenous membrane constituents including endocannabinoids, which suppress increases in corneal inflammation and opacification caused by TRPV1 activation. This suppression occurred through interaction between TRPV1 and CB1, which led to declines in TRPV1 phosphorylation status [16]. Whether TRPV1 is directly activated by the CB1 agonist WIN or CB1 instead induces TRPV1 activation through a cytosolic Ca^{2+} change is not yet elucidated. Interestingly, CB1-induced suppression of a TRPV1-induced Ca^{2+} transient was also observed in eye tumor cells [17, 18].

TRPV4 has an osmosensor function in human corneal epithelial cells, since it is activated by exposure to hypotonicity [19]. The phorbol ester 4 α -phorbol 12,13-didecanoate (4 α -PDD) is a selective TRPV4 channel agonist that




Fig. 3. Suggested Ca^{2+} signal transduction pathways activated by the CB1 agonist WIN. TRPV1 can selectively be activated by heat ($>43^\circ\text{C}$) or capsaicin and blocked by capsazepine. The CB1 receptor coupled to $\text{G}_{i/o}$ proteins can be activated by WIN. If TRPV1 is activated by capsaicin, WIN blocks TRPV1, which leads to a reduction of intracellular Ca^{2+} ($\downarrow[\text{Ca}^{2+}]_i$). Probably, WIN may also directly activate TRPV1 by a CB1-independent mechanism.

induced currents in the whole-cell patch-clamp configuration (fig. 4). In these cells, TRPV4 activation elicited regulatory volume decrease behavior, which was validated by showing that in TRPV4 siRNA gene-silenced cells, this response was suppressed [19]. Another TRPV variant, TRPV3, is also expressed in these cells and has roles in thermosensation and cell proliferation regulation [20]. Furthermore, TRPM8 is expressed in these cells, since specific agonists such as menthol or physical cooling induced Ca^{2+} responses which could be blocked by the specific TRPM8 blocker BCTC [unpubl. data].

Taken together, TRPV1, 3 and 4 subtype functional expression suggests that these different subtypes are essential for eliciting homeostatic responses to thermal variations encountered in daily living that otherwise could compromise the maintenance of corneal visual function. Challenges to corneal epithelial function due to changes in tear film osmolarity caused by thermal variation may be compensated for through activation of TRPV1 and TRPV4, which are stimulated by exposure to hyperosmotic and hypoosmotic stresses, respectively. The responses induced by TRPV1 activation may be fine-tuned through complex regulation by G protein-coupled receptors such as CB1.




Fig. 4. Activation of TRPV4 by 4 α -PDD in human corneal epithelial cells. **a** Experimental design (whole-cell configuration). The holding potential (HP) was set to 0 mV to avoid any voltage-dependent ion channel currents. **b** Voltage pulse protocol. **c** Whole-cell currents (not leak subtracted) induced by stimulation voltages between -90 and +90 mV (control) and after extracellular application of 4 α -PDD (2 μ M). 4 α -PDD induced an increase in cation channel outward currents. **d** Effect of extracellular application of 4 α -PDD (2 μ M) summarized in a current/voltage plot (*I*-*V* plot; *n* = 5; \pm SEM). For the current/voltage relation, maximal peak current amplitudes were normalized to each cell membrane capacitance and plotted against the pipette potential (mV) of the electrical stimulation. The upper trace (filled circles) was determined in the presence of 2 μ M 4 α -PDD

and the lower trace (open circles) in the absence of 4 α -PDD (control). The 4 α -PDD-induced outward currents were significantly increased, whereas the inward currents were not influenced (right panel). **e** Time course of currents activated by 4 α -PDD (2 μ M) at +90 mV measured every minute and plotted against time (in minutes; *n* = 5; \pm SEM). Currents were leak subtracted and normalized to the control amplitude (*I*/*I*₀) measured directly before application of 4 α -PDD. The trace (circles) was determined in the presence of 2 μ M 4 α -PDD, which was applied after 2 min. Outward currents (*I*) clearly increased (inset) and typically appeared with some delay because of extracellular Ca²⁺-free solution. The 4 α -PDD-induced outward currents were significantly increased 4 min after the control currents (without 4 α -PDD; right panel).

Corneal Endothelium

The corneal endothelium is a single layer of postmitotic cells arranged in a hexagonal array at the posterior side of the cornea (fig. 1). By eliciting osmotically coupled

fluid flow from the stroma into the anterior chamber, this tissue maintains corneal deturgescence and transparency [21, 22]. For these functions to be maintained, a critical endothelial cell density along with close cell-to-cell apposition is needed for its barrier property to prevent corneal

swelling. Besides voltage-operated Ca^{2+} channels, the non-temperature-sensitive TRPM2 channel is putatively expressed in this tissue [23, 24]. Given that TRPM2 is activated by oxidative stress (H_2O_2), this suggests that corneal endothelial cells can withstand some levels of oxidative stress [25]. In a study, it was shown that a block of voltage-operated Ca^{2+} channels (L-type) could only partially suppress the H_2O_2 -induced Ca^{2+} influx [24]. Furthermore, the cooling agent icilin induced Ca^{2+} increases, suggesting functional thermosensitive TRPM8 expression [26]. This suggests that corneal endothelial cell function can adapt to thermal variations and probably also to oxidative stress through activation of these TRP subtypes. TRPV1–3 functional expression in this cell layer also contributes to thermal sensing [9]. Modulation of their activity by temperature changes underlies essential homeostatic mechanisms, contributing to the support of corneal endothelial function under different ambient conditions. Interestingly, similar observations were made in the immortalized clonal cell populations HCEC-B4G12 and HCEC-H9C1 also under serum-free culture conditions [27]. The osmosensor TRPV4 was also identified in human corneal endothelial cells, because exposure to a hypotonic environment increased the intracellular Ca^{2+} concentration and TRP-like whole-cell currents in these cells [28]. Therefore, the commonality in thermosensitive TRP functional expression in the corneal epithelial and endothelial layers suggests that these tissues are well poised to compensate for losses in function that are caused by exposure to similar thermal stresses. Nevertheless, it is still unclear whether there are differences in their functional expression patterns, since only some of the numerous thermosensitive TRP subtypes have been evaluated.

Corneal Stroma

The stroma is located between the two outer limiting corneal layers and accounts for up to about 90% of the corneal full thickness (fig. 1). It is formed by an extracellular matrix of regularly arranged collagen fibers fixed in a geometric array and interspersed with keratocytes. TRPV1 expression was identified in primary cultured human fibroblasts [29]. Yang et al. [29] suggested that transforming growth factor (TGF)- β_1 -induced fibrosis and inflammation are dependent on TRPV1 transactivation on fibroblasts in an injury severe enough for epithelial-derived TGF- β_1 to gain stromal access. This dependence is evident, since in TRPV1 knockout mice, the corneal wound healing response to an alkali burn has a far more favorable out-

come [3]. These undesirable effects in wild-type mice may result in part from an increased cytokine secretion caused by TGF- β_1 stimulation of its cognate receptor, which mediates phospho-p38 MAPK signaling required for persistent Smad2/3 phosphorylation [29]. The fibrotic response that occurs subsequently to Smad2/3 activation is due to myofibroblast transdifferentiation, which leads to scarring and corneal opacification [30]. This response is dependent on activation of a positive feedback signaling transduction loop wherein p38 MAPK induces TRPV1 activation, leading to prolongation of both phospho-p38 MAPK and Smad2/3 activation through concerted TRPV1 and TGF- β receptor interaction.

Taken together, stromal TRPV1 activation by injury on stromal resident fibroblasts initially mediates innate immune responses, which provide an adaptive advantage against pathogenic infiltration. However, if this response is not self-limiting and instead becomes dysregulated, the resulting proinflammatory cytokine storm can lead to blindness. As the scarring response is dependent on TRPV1-induced prolongation and enhancement of TGF- β receptor signaling, drug inhibition of TRPV1 activation in a clinical setting may reduce fibrosis occurring during wound healing. Very recent studies suggest that keratocytes also express TRPM8, since its functional activity was recently identified in an established human corneal stromal cell line [31; Tuercker, Mergler et al., unpubl. data]. On the other hand, it is evident that TRPV1 activation in vivo has a double-edged sword aspect, since its activation following epithelial debridement instead increases epithelial cell proliferation and migration and hastens wound closure [2]. In an earlier study, it was shown that such promotion of wound healing occurs through the transactivation of EGF receptor [2, 15]. EGF receptor stimulation contributes to increases in cell proliferation and migration, which are essential responses for wound closure.

Corneal Nerve Fibers

The corneal stroma contains nerve fibers that form a dense midstromal and subepithelial plexus with fibers extending into the basal corneal epithelial layer [32]. Various studies demonstrate also an expression of thermosensitive TRPs in noncorneal primary sensory neurons [33]. TRPV1 and TRPM8 are more extensively described in corneal nerve fibers of mice, guinea pigs and humans [34–37]. Specifically, in mouse corneal nerve fibers, the TRPM8 agonist menthol induced TRPM8 activation. This response is also elicited at the same temperature character-

istic of TRPM8 activation and induces increased basal lacrimation, while TRPM8 gene silencing failed to induce this response. A low concentration of menthol increased lacrimation via TRPM8 channels without evoking nociceptive behavior [37]. Physical cooling of the cornea in humans similarly increased lacrimation, whereas warming had an opposite effect [36]. Therefore, it is suggested that TRPM8 expression in corneal nerves provides a cold receptor function on the ocular surface and may also serve as a 'humidity detector'. These studies broaden our understanding of mechanisms that mediate ocular surface homeostasis. They suggest that tear hyposalivation in dry eye syndrome (DES) may be a consequence of impaired TRPM8 function on corneal nerves, since TRPM8 activity contributes to the regulation of basal tear flow in mice.

In conclusion, thermo-TRPs not only play a role in the reflexogenic protection of the ocular surface and anterior eye but also most likely mediate adaptive responses that are essential for maintaining corneal transparency. In this regard, Parra et al. [36] pointed out that prompt responses which are needed to maintain ocular surface hydration during exposure to a desiccating stress may depend on TRPM8 functional expression. Such maintenance may not be solely dependent on TRPM8 expression in corneal nerves but may also depend on TRPV isotype expression in corneal epithelial, stromal and endothelial cells. A possible functional interaction between expression of thermo-TRPs on nerve fibers and neighboring corneal tissues and cells has not been investigated so far.

TRP Drug Targeting in a Clinical Setting

The realization that a variety of different thermosensitive TRP isoforms and a thermoinsensitive counterpart are expressed on the corneal nerves and the cells of the epithelial, endothelial and stromal layers has prompted interest in developing drugs that selectively control their functional expression. This interest stems from studies demonstrating that these TRPs are activated by sensory stimuli and induce responses to environmental stresses that are of adaptive value by reducing declines in corneal function. TRP activation by such challenges can offset declines in epithelial barrier function caused by an anisotonic stress, hasten corneal reepithelialization and reduce pathogenic infiltration, provided the inflammatory response is self-limiting rather than dysregulated. In addition, thermosensitive TRP activation may be essential to offset declines in tear fluid flow needed for sustaining anterior surface ocular health.

Out of the 27 candidate TRP mammalian genes, only 2 different TRPs so far are of potential value for drug development: TRPV1 and TRPM8. TRPV1 has attracted interest because its activation by injury on afferent corneal sensory nerves in some cases elicits debilitating pain. Another reason for targeting TRPV1 is that after a severe penetrating injury, corneal wound healing is inappropriate due to fibrosis and chronic inflammation. In the first case, suppressing TRPV1 activation through agonist downregulation can initially make the pain more severe due to TRPV1 stimulation, which wanes with time. Therefore, efforts are underway to develop novel selective antagonists that block TRPV1-induced nociception. Many candidates have to be screened because it has to be shown that any decline in TRPV1 functional expression does include neither a systemic nor an off-target effect. It is now also apparent that restoring corneal function with a TRPV1 antagonist should be restricted to usage after a severe injury rather than epithelial debridement. This distinction has to be made, since only after an injury that breaches the basement membrane is it possible for TGF- β to gain access to the stroma and induce fibrosis and chronic inflammation by transactivating TRPV1. Otherwise, after subsequent mild injury, it is not beneficial to inhibit TRPV1 activation, since increases in Ca²⁺ influx elicited by this channel are required to stimulate EGF signaling pathways leading to increases in cell proliferation and migration, both of which hasten wound closure and restoration of barrier function. Another reason to pursue the development of selective TRPV1 antagonists is that inhibition of TRPV1 may reduce inflammation that is symptomatic in some DES patients whose tears are hypertonic. These individuals may also benefit in the future from the development of TRPM8 agonists whose activation by temperature lowering can be promoted to enhance basal tear flow and contribute to the maintenance of ocular surface hydration.

Clinical Aspects: Thermography

Thermography is an investigative technique which allows a rapid color-coded display of the temperature across a wide surface by means of infrared detection using thermography devices such as the novel thermography device TG 1000 (Tomey Corp., Nagoya, Japan) or a thermal camera (noncontact thermographic camera, FLIR P 620) [38, 39]. Thermal pictures of ocular inflammation or noninflammatory ocular pathology can be taken to compare them with pictures of healthy ocular physiology [40]. Accordingly, inflammatory ocular conditions can be easily

distinguished from noninflammatory conditions. Thermography can also be utilized in the evaluation of tear film in DES. Notably, the relation between ocular surface temperature (OST) and physical parameters of the anterior eye is important. OST can be specifically assessed by dynamic thermal imaging. Studies using thermal imaging revealed that OST is principally related to the tear film and less related to other corneal characteristics (e.g. corneal thickness, corneal curvature and anterior chamber depth) [41]. So far, however, nothing is known about TRPs in context with thermography. On the other hand, inflammatory processes are associated with an increased temperature of the affected tissue and increased levels of thermo-TRP expression. At this point, older studies demonstrated that administration of cryotherapy caused a decrease in the infiltration of inflammatory cells in alkali-burned corneas (the activity of collagenase was inhibited) [42]. Based on the recent and growing knowledge about the role of TRPs in connection with clinical aspects of eye surface biology (e.g. DES and keratoplasty), especially the TRPV1 subtype could provide an excellent drug target to reduce inflammation and the discomfort experienced by individuals afflicted with DES. Inflammatory inhibition could be achieved by suppressing its channel activity or expression. Based on future characterizations of other TRP functions, novel ligands could be developed to modify TRP function. This could contribute to an efficient targeted therapy program to achieve a better clinical outcome.

Conclusions

Although there is an increasing awareness of the functional importance of thermo-TRP expression in the maintenance of corneal homeostasis during exposure to environmental stresses, it is still not clear whether channelopathies underlie the pathogenesis of some corneal diseases. Such clarification is needed to determine whether they are potential drug targets to reduce the declines in corneal deturgescence associated with disease. Another important area warranting additional investigation is the functional interplay between TRPM8 expressed on nerve fibers and adjacent corneal stromal and epithelial cells. Nevertheless, the insight gained from characterizing TRP functional expression may have an immediate benefit in identifying novel drug targets for improved treatment of DES and reducing sight-compromising fibrosis after a severe injury that breaches the epithelial basement membrane.

Acknowledgements

Stefan Mergler (PhD) is supported by the DFG Me 1706/13-1 and DFG Me 1706/14-1 grants concerning TRP-related research projects. Peter Reinach (PhD) is supported by the Zhejiang provincial government. The authors thank Noushafarin Khajavi (PhD student) and Olaf Strauß (PhD; both Experimental Ophthalmology, Charité – Universitätsmedizin Berlin) for helpful discussions.

References

- Pan Z, Yang H, Reinach PS: Transient receptor potential (TRP) gene superfamily encoding cation channels. *Hum Genomics* 2011;5: 108–116.
- Sumioka T, Okada Y, Reinach PS, Shirai K, Miyajima M, Yamanaka O, Saika S: Impairment of cornea epithelial wound healing in a TRPV1-deficient mouse. *Invest Ophthalmol Vis Sci* 2014;55:3295–3302.
- Okada Y, Reinach PS, Shirai K, Kitano A, Kao WW, Flanders KC, Miyajima M, Liu H, Zhang J, Saika S: TRPV1 involvement in inflammatory tissue fibrosis in mice. *Am J Pathol* 2011;178:2654–2664.
- Murata Y, Masuko S: Peripheral and central distribution of TRPV1, substance P and CGRP of rat corneal neurons. *Brain Res* 2006; 1085:87–94.
- Yang H, Mergler S, Sun X, Wang Z, Lu L, Bonanno JA, Pleyer U, Reinach PS: TRPC4 knockdown suppresses EGF-induced store operated channel activation and growth in human corneal epithelial cells. *J Biol Chem* 2005;280:32230–32237.
- Yang H, Sun X, Wang Z, Ning G, Zhang F, Kong J, Lu L, Reinach PS: EGF stimulates growth by enhancing capacitative calcium entry in corneal epithelial cells. *J Membr Biol* 2003;194:47–58.
- Xie Q, Zhang Y, Cai S, X, Zhai C, Bonanno JA: Expression and functional evaluation of transient receptor potential channel 4 in bovine corneal endothelial cells. *Exp Eye Res* 2005; 81:5–14.
- Zhang F, Yang H, Wang Z, Mergler S, Liu H, Kawakita T, Tachado SD, Pan Z, Capo-Apon-te JE, Pleyer U, Koziel H, Kao WW, Reinach PS: Transient receptor potential vanilloid 1 activation induces inflammatory cytokine release in corneal epithelium through MAPK signaling. *J Cell Physiol* 2007;213:730–739.
- Mergler S, Valtink M, Coulson-Thomas VJ, Lindemann D, Reinach PS, Engelmann K, Pleyer U: TRPV channels mediate temperature-sensing in human corneal endothelial cells. *Exp Eye Res* 2010;90:758–770.
- Fernandes ES, Fernandes MA, Keeble JE: The functions of TRPA1 and TRPV1: moving away from sensory nerves. *Br J Pharmacol* 2012;166:510–521.
- Pingle SC, Matta JA, Ahern GP: Capsaicin receptor: TRPV1 a promiscuous TRP channel. *Handb Exp Pharmacol* 2007;155–171.
- Gees M, Colsoul B, Nilius B: The role of transient receptor potential cation channels in Ca²⁺ signaling. *Cold Spring Harb Perspect Biol* 2010;2:a003962.
- Voets T, Droogmans G, Wissenbach U, Janssens A, Flockerzi V, Nilius B: The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nature* 2004; 430:748–754.

- 14 Feske S, Gwack Y, Prakriya M, Srikanth S, Puppel SH, Tanasa B, Hogan PG, Lewis RS, Daly M, Rao A: A mutation in Orail causes immune deficiency by abrogating CRAC channel function. *Nature* 2006;441:179–185.
- 15 Yang H, Wang Z, Capo-Aponte JE, Zhang F, Pan Z, Reinach PS: Epidermal growth factor receptor transactivation by the cannabinoid receptor (CB1) and transient receptor potential vanilloid 1 (TRPV1) induces differential responses in corneal epithelial cells. *Exp Eye Res* 2010;91:462–471.
- 16 Yang Y, Yang H, Wang Z, Varadaraj K, Kumari SS, Mergler S, Okada Y, Saika S, Kingsley PJ, Marnett LJ, Reinach PS: Cannabinoid receptor 1 suppresses transient receptor potential vanilloid 1-induced inflammatory responses to corneal injury. *Cell Signal* 2013;25:501–511.
- 17 Mergler S, Cheng Y, Skosyrsky S, Garreis F, Pietrzak P, Kociok N, Dwarakanath A, Reinach PS, Kakkassery V: Altered calcium regulation by thermo-sensitive transient receptor potential channels in etoposide-resistant WERI-Rb1 retinoblastoma cells. *Exp Eye Res* 2012;94:157–173.
- 18 Mergler S, Derckx R, Reinach PS, Garreis F, Bohm A, Schmelzer L, Skosyrsky S, Ramesh N, Abdelmessih S, Polat OK, Khajavi N, Riechardt AI: Calcium regulation by temperature-sensitive transient receptor potential channels in human uveal melanoma cells. *Cell Signal* 2013;26:56–69.
- 19 Pan Z, Yang H, Mergler S, Liu H, Tachado SD, Zhang F, Kao WW, Koziel H, Pleyer U, Reinach PS: Dependence of regulatory volume decrease on transient receptor potential vanilloid 4 (TRPV4) expression in human corneal epithelial cells. *Cell Calcium* 2008;44:374–385.
- 20 Yamada T, Ueda T, Ugawa S, Ishida Y, Imayasu M, Koyama S, Shimada S: Functional expression of transient receptor potential vanilloid 3 (TRPV3) in corneal epithelial cells: involvement in thermosensation and wound healing. *Exp Eye Res* 2010;90:121–129.
- 21 Mergler S, Pleyer U: The human corneal endothelium: new insights into electrophysiology and ion channels. *Prog Retin Eye Res* 2007;26:359–378.
- 22 Mergler S, Pleyer U: Physiology of the human corneal endothelium: new insights from electrophysiological investigations (in German). *Klin Monatsbl Augenheilkd* 2011.
- 23 Mergler S, Dannowski H, Bednarz J, Engelmann K, Hartmann C, Pleyer U: Calcium influx induced by activation of receptor tyrosine kinases in SV40-transfected human corneal endothelial cells. *Exp Eye Res* 2003;77:485–495.
- 24 Mergler S, Pleyer U, Reinach P, Bednarz J, Dannowski H, Engelmann K, Hartmann C, Yousif T: EGF suppresses hydrogen peroxide induced Ca^{2+} influx by inhibiting L-type channel activity in cultured human corneal endothelial cells. *Exp Eye Res* 2005;80:285–293.
- 25 Bari MR, Akbar S, Eweida M, Kuhn FJ, Gustafsson AJ, Luckhoff A, Islam MS: H_2O_2 -induced Ca^{2+} influx and its inhibition by N-(p-aminocinnamoyl)anthranilic acid in the β -cells: involvement of TRPM2 channels. *J Cell Mol Med* 2009;13:3260–3267.
- 26 Mergler S, Mertens C, Valtink M, Reinach PS, Szekeley VC, Slavi N, Garreis F, Abdelmessih S, Turker E, Fels G, Pleyer U: Functional significance of thermosensitive transient receptor potential melastatin channel 8 (TRPM8) expression in immortalized human corneal endothelial cells. *Exp Eye Res* 2013;116:337–349.
- 27 Valtink M, Gruschwitz R, Funk RH, Engelmann K: Two clonal cell lines of immortalized human corneal endothelial cells show either differentiated or precursor cell characteristics. *Cells Tissues Organs* 2008;187:286–294.
- 28 Mergler S, Valtink M, Taetz K, Sahlmüller M, Fels G, Reinach PS, Engelmann K, Pleyer U: Characterization of transient receptor potential vanilloid channel 4 (TRPV4) in human corneal endothelial cells. *Exp Eye Res* 2011;93:710–719.
- 29 Yang Y, Yang H, Wang Z, Mergler S, Wolosin JM, Reinach PS: Functional TRPV1 expression in human corneal fibroblasts. *Exp Eye Res* 2013;107:121–129.
- 30 Yang Y, Wang Z, Yang H, Wang L, Gillespie SR, Wolosin JM, Bernstein AM, Reinach PS: TRPV1 potentiates TGF- β induction of corneal myofibroblast development through an oxidative stress-mediated p38-SMAD2 signaling loop. *PLoS One* 2013;8:e77300.
- 31 Zorn-Kruppa M, Tykhonova S, Belge G, Bednarz J, Diehl HA, Engelke M: A human corneal equivalent constructed from SV40-immortalised corneal cell lines. *Altern Lab Anim* 2005;33:37–45.
- 32 Marfurt CF, Cox J, Deek S, Dvorscak L: Anatomy of the human corneal innervation. *Exp Eye Res* 2010;90:478–492.
- 33 Tominaga M, Caterina MJ: Thermosensation and pain. *J Neurobiol* 2004;61:3–12.
- 34 Hirata H, Oshinsky ML: Ocular dryness excites two classes of corneal afferent neurons implicated in basal tearing in rats: involvement of transient receptor potential channels. *J Neurophysiol* 2012;107:1199–1209.
- 35 Madrid R, Donovan-Rodriguez T, Meseguer V, Acosta MC, Belmonte C, Viana F: Contribution of TRPM8 channels to cold transduction in primary sensory neurons and peripheral nerve terminals. *J Neurosci* 2006;26:12512–12525.
- 36 Parra A, Madrid R, Echevarria D, del Olmo S, Morenilla-Palao C, Acosta MC, Gallar J, Dhaka A, Viana F, Belmonte C: Ocular surface wetness is regulated by TRPM8-dependent cold thermoreceptors of the cornea. *Nat Med* 2010;16:1396–1399.
- 37 Robbins A, Kurose M, Winterson BJ, Meng ID: Menthol activation of corneal cool cells induces TRPM8-mediated lacrimation but not nociceptive responses in rodents. *Invest Ophthalmol Vis Sci* 2012;53:7034–7042.
- 38 Morgan PB, Soh MP, Efron N, Tullo AB: Potential applications of ocular thermography. *Optom Vis Sci* 1993;70:568–576.
- 39 Klamann MK, Maier AK, Gonnermann J, Klein JP, Pleyer U: Measurement of dynamic ocular surface temperature in healthy subjects using a new thermography device. *Curr Eye Res* 2012;37:678–683.
- 40 Kawali AA: Thermography in ocular inflammation. *Indian J Radiol Imaging* 2013;23:281–283.
- 41 Purslow C, Wolffsohn J: The relation between physical properties of the anterior eye and ocular surface temperature. *Optom Vis Sci* 2007;84:197–201.
- 42 Toczolowski J: The effect of immunosuppressors and low temperature on the activity of collagenase in the cornea. *Ann Ophthalmol* 1988;20:64–67.
- 43 Mergler S, Garreis F, Sahlmüller M, Reinach PS, Paulsen F, Pleyer U: Thermosensitive transient receptor potential channels in human corneal epithelial cells. *J Cell Physiol* 2011;226:1828–1842.