





## Concise report

## Temporal trends in COVID-19 outcomes in people with rheumatic diseases in Ireland: data from the COVID-19 Global Rheumatology Alliance registry

Richard Conway <sup>1,2</sup>, Elena Nikiphorou<sup>3,4</sup>, Christiana A. Demetriou<sup>5</sup>, Candice Low<sup>6</sup>, Kelly Leamy<sup>7</sup>, John G. Ryan<sup>8</sup>, Ronan Kavanagh<sup>9</sup>, Alexander D. Fraser<sup>10,11</sup>, John J. Carey<sup>12,13</sup>, Paul O'Connell<sup>14,15</sup>, Rachael M. Flood<sup>2,16</sup>, Ronan H. Mullan<sup>2,16</sup>, David J. Kane<sup>2,16</sup>, Nicola Ambrose<sup>17</sup>, Frances Stafford<sup>17</sup>, Philip C. Robinson <sup>18</sup>, Jean W. Liew <sup>19</sup>, Rebecca Grainger <sup>20</sup> and Geraldine M. McCarthy<sup>7</sup> on behalf of the COVID-19 Global Rheumatology Alliance

## Abstract

**Objectives.** Although evidence is accumulating globally, data on outcomes in rheumatic disease and COVID-19 in Ireland are limited. We used data from the COVID-19 Global Rheumatology Alliance (C19-GRA) to describe time-varying COVID-19 outcomes for people with rheumatic disease in Ireland.

**Methods.** Data entered into the C19-GRA provider registry from Ireland between 24 March 2020 and 9 July 2021 were analysed. Differences in the likelihood of hospitalization and mortality according to demographic and clinical variables were investigated using Chi-squared test or Fisher's exact test, as appropriate. Trends in odds of hospitalization and mortality over time were investigated using logistic regression with the time period as a categorical variable.

**Results.** Of 212 cases included, 59.4% were female and median age was 58.0 years (range 13–96). Of the 212 cases, 92 (43%) were hospitalized and 22 (10.4%) died. Increasing age, a diagnosis of gout, ever smoking, glucocorticoid use, having comorbidities and specific comorbidities of cancer, cardiovascular and pulmonary disease were more common in those hospitalized. A diagnosis of inflammatory arthritis, csDMARD and/or b/tsDMARD use were less frequent in those hospitalized. Increasing age, a diagnosis of gout, ever smoking, having comorbidities and specific comorbidities of obesity, cardiovascular and pulmonary disease were more common in those who died. Odds of hospitalization or mortality did not change over time.

**Conclusion.** No temporal trend was observed in either COVID-19-related hospitalization or mortality outcomes for people with rheumatic disease in Ireland.

**Key words:** rheumatic disease, COVID-19, biologics, hospitalization

<sup>1</sup>Department of Rheumatology, St. James's Hospital, <sup>2</sup>Clinical Medicine, Trinity College Dublin, Dublin, Ireland, <sup>3</sup>Department of Rheumatology, <sup>4</sup>Centre for Rheumatic Diseases, King's College London, London, UK, <sup>5</sup>Department of Primary Care and Population Health, University of Nicosia Medical School, Nicosia, Cyprus, <sup>6</sup>Department of Rheumatology, St. Vincent's University Hospital, <sup>7</sup>Department of Rheumatology, Mater Misericordiae Hospital, Dublin, <sup>8</sup>Department of Rheumatology, Cork University Hospital, Wilton, <sup>9</sup>Department of Rheumatology, Galway Clinic, Galway, <sup>10</sup>Department of Rheumatology, University Hospitals Limerick, <sup>11</sup>Graduate Entry Medical School, University of Limerick, Limerick, <sup>12</sup>Department of Rheumatology, Galway University Hospitals, <sup>13</sup>School of Medicine, National University of Ireland Galway, Galway, <sup>14</sup>Department of Rheumatology, Beaumont Hospital, <sup>15</sup>School of Medicine, Royal College of Surgeons in Ireland,

<sup>16</sup>Department of Rheumatology, Tallaght University Hospital, <sup>17</sup>Department of Rheumatology, Blackrock Clinic, Dublin, Ireland, <sup>18</sup>University of Queensland Faculty of Medicine, Brisbane, Australia, <sup>19</sup>Section of Rheumatology, Boston University School of Medicine, Boston, MA, USA and <sup>20</sup>Department of Medicine, University of Otago, Wellington, New Zealand

Submitted 17 December 2021; accepted 1 March 2022

Correspondence to: Richard Conway, Department of Rheumatology, St. James's Hospital, Dublin 8, Ireland.  
E-mail: drrichardconway@gmail.com

### Rheumatology key messages

- The current study includes 212 cases of COVID-19 in people with rheumatic disease reported in Ireland.
- Of the COVID-19 cases reported, 43% were hospitalized and 10% died.
- COVID-19 outcomes for people with rheumatic disease did not show the temporal improvements observed in the general population.

## Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the resulting Coronavirus disease 2019 (COVID-19) continue to present a global health crisis. Significant improvements in COVID-19 outcomes have been demonstrated over time [1–5]. This may relate to accumulating clinical experience leading to an improvement in care [6] along with beneficial effects of several immunomodulating treatments in severe COVID-19 [7–10]. It is unclear whether outcomes for people with rheumatic disease who develop COVID-19 are also improving, with some studies reporting improved outcomes [4, 11]. Because outcomes following SARS-CoV-2 infection may also vary depending on the geography and ethnicity of affected populations, it is important to explore outcomes in different locations and over time [12–16].

We have previously reported predictors of hospitalization in people with rheumatic disease who developed COVID-19 in the first five months of the pandemic in Ireland [17]. In this study we examine temporal trends in COVID-19 outcomes in people with rheumatic disease in Ireland in the first 16 months of the pandemic.

## Methods

### COVID-19 Global Rheumatology Alliance

Data regarding individuals with rheumatic disease with COVID-19 are entered into one of two parallel international data portals hosted in the United States and United Kingdom. Details of the C19-GRA registries have been published previously [18, 19].

### Data collection

Cases in this study were entered into the C19-GRA provider registry from 24 March 2020 to 9 July 2021. Data were collected on baseline rheumatic disease status including demographic and clinical variables such as age, sex, smoking status, rheumatic disease diagnosis, disease activity (as per the physician's global assessment (remission, low, moderate or high/severe)), and comorbidities. Medications were categorized as previously described [17]. DMARDs were grouped as conventional synthetic DMARDs (csDMARDs) or biologic DMARDs (bDMARDs) and targeted synthetic DMARDs (tsDMARDs) [18]. Rheumatic diseases were categorized as (i) inflammatory arthritis (IA); (ii) gout; and (iii) vasculitis, connective tissue diseases and all other diagnoses ('other'). Data collected regarding COVID-19 infection

included method of diagnosis, place of diagnosis, COVID-19 symptoms, and outcomes of COVID-19 disease including hospitalization, ventilation and death.

Demographic and clinical continuous variables were reported as median (IQR) and categorical variables as number and percentage (%). Hospitalization and mortality probability were calculated for each category of the key demographic and clinical variables. Differences in the likelihood of hospitalization and mortality according to demographic and clinical variables were investigated using Chi-squared test or Fisher's exact test, as appropriate.

Trends in the odds of hospitalization and mortality between timing of diagnosis were investigated using logistic regression with hospitalization or mortality as the dependent variable and wave as a categorical predictor. Each case was categorized under a wave using the date of diagnosis as follows: Wave 1 (weeks 10–31), 1 March 2020–1 August 2020; Wave 2 (week 32–47), 2 August 2020–21 November 2020; and Wave 3 (week 48 onwards), 22 November 2020–9 July 2021 [20]. The COVID-19 vaccination programme in Ireland began on 29 December 2020, with 1 million vaccine doses administered by April 2021, and 90% of adults fully vaccinated by September 2021. Appropriate confounders were identified as those demographic and clinical variables that were associated with both the outcome of interest as well as with timing/wave of diagnosis. These confounders were corrected for in each model. The significance of any suspected trend was tested using the 'contrast' post-estimation command.

A sensitivity analysis was performed restricting our results to two institutions employing a rigorous case-finding methodology. These institutions contacted all patients requesting that they notify the rheumatology service if they developed COVID-19 (as part of clinical care), all admissions to the institutions with COVID-19 were screened for rheumatic disease cases, and finally all positive SARS-CoV-2 PCR tests were back-referenced against people attending the rheumatology services using medical record linkage.

All statistical analyses were performed using STATA IC15 (StataCorp. 2017. Stata Statistical Software: Release 15. College Station, TX, USA: StataCorp LLC.) and graphs were prepared using GraphPad Prism version 6 for Windows (GraphPad Software, La Jolla, CA, USA, www.graphpad.com).

This study was approved by the Irish National Research Ethics Committee for COVID-19 (20-NREC-COV-010). The committee waived the need for written informed consent as the data were fully anonymised.

## Results

The 212 cases had a median age of 58.0 years (IQR: 45.5–73.0) and the majority were female (59.4%). Most had IA (136/212 64.2%) with the balance having gout (35/212, 16.5%) and CTD or other diagnoses (50/212, 23.6%). The majority of patients (46.2%) were in remission, with 35.6% of cases having low disease activity. Seventy cases (33%) had no comorbidities; among those with comorbidities, cardiovascular and pulmonary diseases were the most prevalent (42% and 17.5%, respectively). Just under half of cases were hospitalized (92/212, 43%) and 22 (10.4%) died.

### Hospitalization outcome

Demographic and clinical details according to hospitalization are shown in [Table 1](#). Increasing age, a diagnosis of gout, ever smoking, glucocorticoid use, having comorbidities, and specific comorbidities of cancer and cardiovascular and pulmonary disease were more common in those hospitalized. Patients at either high or low disease activity were more likely to be hospitalized. A diagnosis of IA, csDMARD and/or b/tsDMARD use was less frequent in those hospitalized.

### Mortality outcome

Demographic and clinical details according to mortality are shown in [Table 1](#). Increasing age, a diagnosis of gout, ever smoking, having comorbidities and specific comorbidities of obesity, cardiovascular and pulmonary disease were more common in those who died.

### Hospitalization and mortality outcomes according to wave of diagnosis

The majority of people ( $n = 115$ , 54.2%) were diagnosed during the first wave of the pandemic in Ireland, with 8.9% ( $n = 19$ ) in the second wave and 36.3% ( $n = 77$ ) in the third wave. Hospitalization occurred in 49/115 (44.1%) during the first wave, 2/19 (11.1%) during the second wave and 41/77 (54%) during the third wave. Death occurred in 10/115 (8.7%) during the first wave, 1/19 (5.3%) during the second wave and 11/77 (14.3%) during the third wave. A diagnosis of IA, disease activity, cardiovascular disease or obesity was differentially represented in the three waves ([Supplementary Table S1](#), available at *Rheumatology* online). They were thus inserted as confounders, along with age and gender, in the logistic regression model investigating the odds of hospitalisation across waves. Likewise, disease activity, cardiovascular disease and obesity were both added as confounders, along with age and gender, in the respective model for mortality.

Cases diagnosed during wave 2 displayed a decrease in the likelihood of hospitalisation compared with wave 1, whereas cases diagnosed during wave 3 had increased hospitalisation odds compared with wave 1, but all changes were non-significant, as was the

quadratic trend in the odds for hospitalisation between the three waves.

In terms of mortality, cases diagnosed in each successive wave demonstrated again non-statistically significant increased likelihoods of mortality, and a non-statistically increasing linear trend in the odds of mortality across waves ([Fig. 1](#)).

The sensitivity analysis restricting our results to the two institutions with rigorous case-finding methodology, reporting 59% of cases, did not alter the overall results.

## Discussion

This is the largest and most detailed report of people with rheumatic disease and COVID-19 from Ireland. We have previously reported associations of clinical features with hospitalisation [17]. We did not identify any reduction in odds for hospitalisation or mortality over time. This lack of change in odds of hospitalisation and mortality over time was in contrast to other data where clear improvements in COVID-19 outcomes have been demonstrated for non-rheumatic disease patients over time [1, 2, 21]. Likely contributors are better general care measures as well as targeted treatments such as glucocorticoids and tocilizumab [7, 8]. One explanatory hypothesis may be that people with rheumatic diseases, who may be receiving long-term immunomodulating treatments, may be less likely to benefit from the use of these treatments in the setting of subsequent severe COVID-19. Alternatively, our small sample size may have limited ability to demonstrate a change. The roll-out of the COVID-19 vaccine in Ireland occurred in conjunction with the third COVID-19 wave and may have impacted on our findings.


Previous studies in rheumatic disease patients have suggested some improvement in outcomes over time [4, 5, 11]. A cohort study from Boston of 143 rheumatic disease patients demonstrated a reduction in rates of mechanical ventilation, but not of hospitalisation or mortality, over time [11]. An exposure scored matched analysis study of 8540 rheumatic disease patients using the US TriNetX electronic health record database reported a reduction in the risks of hospitalisation, mechanical ventilation and mortality when comparing a late cohort to an early cohort [4]. The findings of these studies, and the current work, must be interpreted with cognizance of the limitations, including selection bias, unmeasured confounders and possible artefact related to the identification of milder cases over time as testing capacity expanded [5].

Our study has several limitations. While this is the largest study of people with rheumatic disease and COVID-19 from Ireland, the statistical power of our study is limited by the low number of cases; we did not undertake multivariable analyses due to this. The C19-GRA is a physician-entered registry and is limited by selection bias with a likely tendency to report more severe cases. There is also the possibility that the reporting of cases varied over time

**TABLE 1** Hospitalisation outcome according to demographic and clinical factors in people with rheumatic disease diagnosed with COVID-19

	All Participants (n = 212)	Not hospitalised (n = 113)	Hospitalised (n = 92)	P- value	Alive (n = 190)	Deceased (n = 22)	P- value
Gender, n (%)							
Female	126 (59.4)	72 (59.0)	50 (41.0)	0.17	114 (90.5)	12 (9.5)	0.62
Male	86 (40.6)	41 (49.4)	42 (50.6)		76 (88.4)	10 (11.6)	
Age (years), n (%)							
18-29	8 (3.8)	6 (85.7)	1 (14.3)	<b>&lt;0.01<sup>a</sup></b>	8 (100.0)	0 (0.0)	<b>&lt;0.01<sup>a</sup></b>
30-49	54 (25.8)	42 (80.8)	10 (19.2)		54 (100.0)	0 (0.0)	
50-65	67 (32.1)	46 (70.8)	19 (29.2)		62 (92.5)	5 (7.5)	
>65	80 (38.3)	17 (21.5)	62 (78.5)		63 (78.8)	17 (21.2)	
Most common rheumatic disease diagnoses <sup>a</sup> , n (%)							
Inflammatory arthritis <sup>b</sup>	136 (64.2)	91 (70.0)	39 (30.0)	<b>&lt;0.01</b>	126 (92.7)	10 (7.3)	0.05
Gout	35 (16.5)	2 (5.7)	33 (94.3)	<b>&lt;0.01</b>	27 (77.1)	8 (22.9)	<b>0.01<sup>a</sup></b>
Connective Tissue Disease and Other <sup>c</sup>	50 (23.6)	23 (47.9)	25 (52.1)	0.25	45 (90.0)	5 (10.0)	0.29
Disease Activity, n (%)							
1	96 (46.2)	34 (36.6)	59 (63.4)	<b>&lt;0.01<sup>a</sup></b>	81 (84.4)	15 (15.6)	0.19 <sup>a</sup>
2	74 (35.6)	54 (75.0)	18 (25.0)		70 (94.6)	4 (5.4)	
3	30 (14.4)	21 (70.0)	9 (30.0)		27 (90.0)	3 (10.0)	
4	5 (2.40)	1 (20.0)	4 (80.0)		5 (100.0)	0 (0.0)	
No comorbidities, n (%)	70 (33.0)	58 (85.3)	10 (14.7)	<b>&lt;0.01</b>	69 (98.6)	1 (1.4)	<b>&lt;0.01</b>
Most common comorbidities, n (%)							
Cancer	10 (4.3)	1 (11.1)	9 (88.9)	<b>0.01<sup>a</sup></b>	7 (77.8)	2 (22.2)	0.24 <sup>a</sup>
Cardiovascular disease <sup>d</sup>	89 (42.0)	23 (26.4)	64 (73.6)	<b>&lt;0.01</b>	70 (78.7)	19 (21.3)	<b>&lt;0.01<sup>a</sup></b>
Pulmonary disease <sup>d</sup>	37 (17.5)	9 (24.3)	28 (75.7)	<b>&lt;0.01</b>	26 (70.3)	11 (29.7)	<b>&lt;0.01<sup>a</sup></b>
Neurological/Neuromuscular/ Psychiatric disease	10 (4.7)	3 (30.0)	7 (70.0)	0.12	8 (80.0)	2 (20.0)	0.28 <sup>a</sup>
Obesity	20 (9.4)	11 (55.0)	9 (45.0)	0.99	14 (70.0)	6 (30.0)	<b>&lt;0.01<sup>a</sup></b>
Smoking status, n (%)							
Never	120 (56.6)	72 (60.5)	47 (39.5)	<b>&lt;0.01</b>	110 (91.7)	10 (8.3)	<b>0.02</b>
Ever	54 (25.5)	15 (28.3)	38 (71.7)		43 (79.6)	11 (20.4)	
Unknown	38 (17.9)						
Medication prior to COVID-19 diagnosis, n (%)							
Steroids	33 (15.6)	11 (34.4)	21 (65.6)	<b>0.01</b>	26 (78.8)	7 (21.2)	0.06
Steroids 10 mg or more	15 (7.1)	4 (28.6)	10 (71.4)	<b>0.04</b>	12 (80.0)	3 (20.0)	0.19
csDMARD monotherapy <sup>e</sup>	61 (28.8)	39 (67.2)	19 (32.8)	<b>0.03</b>	56 (91.8)	5 (8.2)	0.51
b/tsDMARD (monotherapy or in combination with csDMARD) <sup>f</sup>	81 (38.2)	56 (70.0)	24 (30.0)	<b>0.00</b>	75 (92.6)	6 (7.4)	0.27
No complications, n (%)	162 (76.4)	155 (95.7)	7 (4.3)	<b>&lt;0.01</b>	155 (95.7)	7 (4.3)	<b>&lt;0.01</b>
Most common complications, n (%)							
ARDS	10 (4.7)	3 (30.0)	7 (70.0)	<b>&lt;0.01<sup>a</sup></b>	3 (30.0)	7 (70.0)	<b>&lt;0.01<sup>a</sup></b>
Sepsis	9 (4.3)	4 (44.4)	5 (55.6)	<b>&lt;0.01<sup>a</sup></b>	4 (44.4)	5 (55.6)	<b>&lt;0.01<sup>a</sup></b>
Concomitant infection	14 (6.6)	9 (64.3)	5 (35.7)	<b>0.01<sup>a</sup></b>	9 (64.3)	5 (35.7)	<b>&lt;0.01<sup>a</sup></b>
Thromboembolism	11 (5.2)	10 (89.6)	1 (9.1)	<b>1.00<sup>a</sup></b>	10 (90.9)	1 (9.1)	1.00
AKI or renal failure	7 (3.3)	2 (28.6)	5 (71.4)	<b>&lt;0.01<sup>a</sup></b>	2 (28.6)	5 (71.4)	<b>&lt;0.01<sup>a</sup></b>
Deceased, n (%)	22 (10.4)	0 (0.0)	22 (100.0)	<b>&lt;0.01</b>	—	—	

P-value from Pearson's  $\chi^2$  test, unless a=Fisher's Exact test. <sup>a</sup>Patients could be diagnosed with more than one rheumatic diseases. <sup>b</sup>Inflammatory arthritis diagnosis includes: axial spondyloarthritis (including ankylosing spondylitis); PsA; other spondyloarthritis (including reactive arthritis); JIA, oligo; juvenile idiopathic arthritis, poly; systemic JIA; RA; other inflammatory arthritis. <sup>c</sup>Connective tissue disease and other diagnoses include: NCA-associated vasculitis (e.g. GPA, EGPA); other vasculitis including Kawasaki disease; anti-phospholipid antibody syndrome; autoinflammatory syndrome (including TRAPS, CAPS, FMF); Behcet's; Chronic recurrent multifocal osteomyelitis; GCA; IgG4-related disease; inflammatory myopathy (e.g. DM, PM); IBM; mixed connective tissue disease; ocular inflammation; PMR; sarcoidosis; SS; SLE; SSC; undifferentiated CTD; localized scleroderma (morphea); other. <sup>d</sup>Cardiovascular diseases include: cerebrovascular disease; CVD; hypertension; diabetes; and renal disease. Pulmonary diseases include asthma; COPD; and interstitial lung disease. <sup>e</sup>csDMARD monotherapy includes: antimalarials (including HCQ, chloroquine, mepacrine/quinacrine); apremilast; AZA/6-MP; ciclosporin; leflunomide; MTX; MMF/mycophenolic acid; SSZ; tacrolimus; thalidomide/lenalidomide. <sup>f</sup>b/tsDMARD therapy includes: abatacept; belimumab; CD-20 inhibitors (including rituximab, ofatumumab); CYC; IL-1 inhibitors (including anakinra, canakinumab, rilonacept); IL-6 inhibitors (including tocilizumab, sarilumab); IL-12 inhibitors (ustekinumab); IL 23 inhibitors (guselkumab, risankizumab); IL-17 inhibitors (including secukinumab, ixekizumab); JAK inhibitors (including tofacitinib, baricitinib, upadacitinib); TNF-inhibitors (including infliximab, etanercept, adalimumab, golimumab, certolizumab and biosimilars); rituximab within the last 12 months. Bold values indicates the significance of  $P < 0.05$ .

**Fig. 1** Odds for hospitalisation and mortality according to wave of diagnosis

potentially influencing our results. However, restricting our data to the two institutions with rigorous case ascertainment did not alter the overall findings. The C19-GRa is also a case-based registry with no denominator population, therefore inferences cannot be made about the incidence of COVID-19 in people with rheumatic diseases. Additionally, the C19-GRa is by design restricted to people with rheumatic disease and COVID-19; therefore, comparisons cannot be made to people with non-rheumatic disease and COVID-19 nor to rheumatic disease in the absence of COVID-19.

In conclusion, our findings demonstrate no difference in hospitalisation or mortality in patients with rheumatic disease across the first three COVID-19 waves in Ireland.

## Acknowledgements

The authors would like to thank all rheumatology providers who entered data into the registry.

R.C., C.L., K.L., J.G.R., R.K., A.D.F., J.J.C., P.O.C., R.M.F., R.H.M., D.J.K., N.A., F.S. and G.M.M. contributed to data collection, data quality control, data analysis and interpretation. They drafted, and revised, the manuscript critically for important intellectual content and gave final approval of the version to be published. E.N. and C.A.D. contributed to the analysis and interpretation of the data. They drafted, and revised, the manuscript critically for important intellectual content and gave final approval of the version to be published. P.C.R., J.W.L. and R.G. directed the work and contributed to the analysis and interpretation of the data. They drafted, and revised, the manuscript critically for important intellectual content and gave final approval of the version to be published.

**Funding:** No specific funding was received from any bodies in the public, commercial or not-for-profit sectors to carry out the work described in this article.

**Disclosure statement:** The views expressed here are those of the authors and do not necessarily represent the views of the ACR, the EULAR, and the (UK) National Health Service (NHS), the National Institute for Health Research (NIHR) or the (UK) Department of Health. R.C.: Speaker's bureau: Janssen, Roche, Sanofi, Abbvie, Elena. E.N.: Speaker's bureau: AbbVie Eli-Lilly, Gilead, Celtrion, Pfizer, Sanofi. P.C.R.: Speaker's bureau: UCB, Roche, Pfizer, Gilead, Janssen, Novartis, Eli Lilly, Abbvie; Grant/research support from: Abbvie, UCB, Novartis, Janssen, Pfizer. J.W.L.: Grant/research support from: Pfizer. R.G.: Speaker's bureau: Pfizer, Cornerstones, Janssen, Novartis, Abbvie. The other authors have declared no conflicts of interest.

**Ethics:** Irish National Research Ethics Committee for COVID-19 (20-NREC-COV-010).

**Data sharing statement:** Request for access to data from the registry should be made to the Data Access and Sharing Committee of the COVID-19 Global Rheumatology Alliance.

## Data availability statement

The data underlying this article are available on reasonable request to the corresponding author.

## Supplementary data

Supplementary data are available at *Rheumatology* online.

## References

- 1 Anesi GL, Jablonski J, Harhay MO, Atkins JH *et al.* Characteristics, outcomes, and trends of patients with COVID-19-related critical illness at a learning health system in the United States. *Ann Intern Med* 2021;174:613–21.
- 2 Roth GA, Emmons-Bell S, Alger HM *et al.* Trends in patient characteristics and COVID-19 in-hospital mortality in the United States during the COVID-19 pandemic. *JAMA Network Open* 2021;4:e218828.
- 3 Yeates EO, Nahmias J, Chinn J *et al.* Improved outcomes over time for adult COVID-19 patients with acute respiratory distress syndrome or acute respiratory failure. *PLoS One* 2021;16:e0253767.
- 4 Jorge A, D'Silva KM, Cohen A *et al.* Temporal trends in severe COVID-19 outcomes in patients with rheumatic disease: a cohort study. *Lancet Rheumatol* 2021;3:e131–e7.
- 5 Gianfrancesco MA, Robinson PC. Changing COVID-19 outcomes in patients with rheumatic disease—are we really getting better at this? *Lancet Rheumatol* 2021;3:e88–e90.
- 6 Smith V, Devane D, Nichol A, Roche D. Care bundles for improving outcomes in patients with COVID-19 or related conditions in intensive care - a rapid scoping review. *Cochrane Database Syst Rev* 2020;12:Cd013819.
- 7 RECOVERY Collaborative Group, Horby P, Lim WS *et al.* Dexamethasone in hospitalized patients with Covid-19. *New Engl J Med* 2021;384:693–704. <https://pubmed.ncbi.nlm.nih.gov/32678530/>
- 8 RECOVERY Collaborative Group. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet* 2021;397:1637–45. <https://pubmed.ncbi.nlm.nih.gov/33933206/>
- 9 Guimarães PO, Quirk D, Furtado RH *et al.* Tofacitinib in patients hospitalized with Covid-19 pneumonia. *New Engl J Med* 2021;385:406–15.
- 10 Kalil AC, Patterson TF, Mehta AK *et al.* Baricitinib plus remdesivir for hospitalized adults with Covid-19. *New Engl J Med* 2021;384:795–807.
- 11 Serling-Boyd N, D'Silva KM, Hsu TY *et al.* Coronavirus disease 2019 outcomes among patients with rheumatic diseases 6 months into the pandemic. *Ann Rheum Dis* 2021;80:660–6.
- 12 Mazumder A, Arora M, Sra MS, Gupta A *et al.* Geographical variation in case fatality rate and doubling time during the COVID-19 pandemic. *Epidemiol Infect* 2020;148:e163.
- 13 Scirè CA, Carrara G, Zanetti A *et al.* COVID-19 in rheumatic diseases in Italy: first results from the Italian registry of the Italian Society for Rheumatology (CONTROL-19. *Clin Exp Rheumatol* 2020;38:748–53.
- 14 D'Silva KM, Jorge A, Cohen A *et al.* COVID-19 outcomes in patients with systemic autoimmune rheumatic diseases compared to the general population: a US multicenter, comparative cohort study. *Arthritis Rheumatol* 2021;73:914–20.
- 15 FAI2R/SFR/SNFM/IOF/SOFREMIP/CRI/IMIDIATE consortium and contributors. Severity of COVID-19 and survival in patients with rheumatic and inflammatory diseases: data from the French RMD COVID-19 cohort of 694 patients. *Ann Rheum Dis* 2021;80:527–38. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7712850/>
- 16 Pang R, Zhao J, Gan Z *et al.* Evolution of COVID-19 in patients with autoimmune rheumatic diseases. *Aging* 2020;12:23427–35.
- 17 Conway R, Nikiphorou E, Demetriou CA *et al.* Predictors of hospitalization in patients with rheumatic disease and COVID-19 in Ireland: data from the COVID-19 global rheumatology alliance registry. *Rheumatol Adv Pract* 2021;5:rkab031.
- 18 Gianfrancesco M, Hyrich KL, Al-Adely S *et al.* Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: data from the COVID-19 Global Rheumatology Alliance physician-reported registry. *Ann Rheum Dis* 2020;79:859–66.
- 19 Gianfrancesco MA, Hyrich KL, Gossec L *et al.* Rheumatic disease and COVID-19: initial data from the COVID-19 Global Rheumatology Alliance provider registries. *Lancet Rheumatol* 2020;2:e250–3.
- 20 Lima V. The Pandemic One Year on: Trends and Statistics Between Three Waves of the COVID-19 Pandemic in Ireland. Available from: <https://publicpolicy.ie/papers/the-pandemic-one-year-on-trends-and-statistics-between-three-waves-of-the-covid-19-pandemic-in-ireland/> (17 December 2021, date last accessed).
- 21 Conway RP, Byrne DG, O'Riordan DMR *et al.* Emergency medical admissions and COVID-19: impact on 30-day mortality and hospital length of stay. *Ir J Med Sci* 2021;doi:10.1007/s11845-021-02752-7.