

Terpenoids and Sterols from *Hoya multiflora* Blume

Virgilio D. Ebajo Jr.¹, Chien-Chang Shen², Consolacion Y. Ragasa^{1,3*}

¹Chemistry Department, De La Salle University, 2401 Taft Avenue, Manila 1004, Philippines. ²National Research Institute of Chinese Medicine, 155-1, Li-Nong St., Sec. 2, Taipei 112, Taiwan. ³Chemistry Department, De La Salle University Science & Technology Complex Leandro V. Locsin Campus, Binan City, Laguna 4024, Philippines.

ARTICLE INFO

Article history:

Received on: 30/01/2015

Revised on: 12/02/2015

Accepted on: 07/03/2015

Available online: 27/04/2015

Key words:

Hoya multiflora Blume,
Apocynaceae, bauerenol,
lupeol, lupeol acetate, α -
amyryn, α -amyryn acetate, β -
amyryn, β -amyryn acetate,
squalene, lutein, β -sitosterol,
stigmasterol.

ABSTRACT

Chemical investigation of the dichloromethane extracts of *Hoya multiflora* Blume led to the isolation of lupeol (**1a**), α -amyryn (**1b**), β -amyryn (**1c**), lupeol acetate (**2a**), α -amyryn acetate (**2b**), and β -amyryn acetate (**2c**) from the stems; and **1b**, bauerenol (**3**), squalene (**4**), lutein (**5**), β -sitosterol (**6a**), and stigmasterol (**6b**) from the leaves. The structures of **1-6** were identified by comparison of their ¹H and/or ¹³C NMR data with those reported in the literature.

INTRODUCTION

Hoya plants are also called wax plants due to the waxy appearance of their leaves or flowers. There are at least 109 species of *Hoya* found in the Philippines, 88 of these are endemic to the country (Aurigue, 2013). The *Hoya multiflora*, also called shooting-star hoyo is indigenous to the Philippines. This plant was called multiflora due to its multiple flowers (about 40) in the convex umbel (Aurigue, 2013). There are no reported chemical studies and biological activities on *H. multiflora*. However, congeners of the plant have been studied for their chemical constituents. Gas chromatographic analysis on the chemical constituents of *Hoya naumanii* led to the detection of the triterpenes β -amyryn, lupeol and α -amyryn and their 3, 4-*seco*-3-oic acid methyl esters (Baas and Van Berkel, 1991). The isolation of pentacyclic triterpenols δ -amyryn, β -amyryn, lupeol and α -amyryn and their 3, 4-*seco*-3-*nor*-2-ol derivatives (australinolins A–D) from the leaf wax, of, *Hoya australis*, have

been reported (Baas et al., 1992). Moreover, the β -amyryn derivative 5-isopropyl-10 (2-methoxycarbonylethyl)des-A-olean-12-en and the taraxerol derivative 5-isopropyl-10 (2-methoxycarbonylethyl)des-A-olean-14-en were isolated from *Hoya lacunose* (Baas, 1983). The oligosaccharides 6-deoxy-3-, *O*-methyl- β -allopyranosyl (1 \rightarrow 4)- β -cymaropyranosyl (1 \rightarrow 4)- β -cymaronic acid δ -lactone and 6-deoxy-3-*O*-methyl- β -allopyranosyl (1 \rightarrow 4)- β -oleandropyranosyl (1 \rightarrow 4)- β -cymaropyranosyl (1 \rightarrow 4)- β -cymaronic acid δ -lactone and its sodium salt were isolated from *Hoya carmosa* (Yoshikawa et al., 2000). A review on the chemical and pharmacological aspects of *Hoya* species has been provided (Pandey et al., 2006). *Hoya* species yielded pregnanes, lipids, sterols, flavanols, triterpenes, sesquiterpenes and disaccharides. They were reported to exhibit antinematodal activity, hypo sensitization, immunological properties and phytotoxicity; used for the treatment of occupational asthma and sea-squirt asthma and allergies; and employed as antigens and insecticides (Pandey et al., 2006).

This study was conducted as part of our research on the chemical constituents of the genus *Hoya*. We earlier, reported the isolation of lupenone and lupeol from the roots; lupeol, squalene and β -sitosterol from the leaves; and betulin from, the stems of *Hoya mindorensis* Schlechter (Ebajo et al., 2014).

* Corresponding Author

Consolacion Y. Ragasa, Chemistry Department, De La Salle University,
2401 Taft Avenue, Manila 1004, Philippines.
Email: consolacion.ragasa@dlsu.edu.ph




Fig. 1: Chemical Constituents of *Hoya multiflora* lupeol (1a), lupeol acetate (2a), α -amyrin (1b), α -amyrin acetate (2b), β -amyrin (1c), β -amyrin acetate (2a), bauerenol (3), squalene (4), lutein (5), β -sitosterol (6a), and stigmasterol (6b).

We report herein the isolation of lupeol (**1a**), α -amyrin (**1b**), β -amyrin (**1c**), lupeol acetate (**2a**), α -amyrin acetate (**2b**), and β -amyrin acetate (**2c**) from the stems; and **1b**, bauerenol (**3**), squalene (**4**), lutein (**5**), β -sitosterol (**6a**), and stigmasterol (**6b**) from the leaves of *Hoya multiflora* Blume. To the best of our knowledge this is the first report on the isolation of these compounds from *H. multiflora*.

MATERIALS AND METHODS

General Experimental Procedure

NMR spectra were recorded on a Varian VNMRs spectrometer in CDCl_3 at 600 MHz for ^1H NMR and 150 MHz for ^{13}C NMR spectra. Column chromatography was performed, with silica gel 60 (70-230 mesh). Thin layer chromatography, was

performed with plastic backed plates coated with silica gel F₂₅₄ and the plates were visualized by spraying with vanillin/ H_2SO_4 solution followed by warming.

Sample Collection

Hoya multiflora Blume was collected from a garden in Pangasinan, Philippines in September 2013. Voucher specimens were authenticated at the Botany Division of the Philippine National Museum.

General Isolation Procedure

The air-dried leaves (23.0 g), and stems (86.5 g) of *H. multiflora* were ground in a blender, soaked in CH_2Cl_2 for three days and then filtered. The filtrates were concentrated under vacuum to afford crude extracts of leaves (2.0 g), and stems (1.0 g)

which were each chromatographed by gradient elution with CH₂Cl₂, followed by increasing amounts of acetone at 10% increment by volume as eluents. A glass column 12 inches in height and 0.5 inch internal diameter was used for the fractionation of crude extracts. Two milliliter fractions were collected. Fractions with spots of the same *R_f* values were combined and rechromatographed in appropriate solvent systems until TLC pure isolates were obtained. Rechromatography and final purifications were conducted using Pasteur pipettes as columns. One milliliter fractions were collected.

Isolation of Chemical Constituents of the Stems

The CH₂Cl₂ fraction from the chromatography of the crude extract was rechromatographed using 1% EtOAc in petroleum ether to afford a mixture of **2a-2c** (9 mg) after washing with petroleum ether. The 20% acetone in CH₂Cl₂ fraction was rechromatographed using 5% EtOAc in petroleum ether, to afford a mixture of **1a-1c** (4 mg) after washing with petroleum ether.

Isolation of Chemical Constituents of the Leaves

The CH₂Cl₂ fraction from the chromatography of the crude extract was rechromatographed (3 ×) using petroleum ether to afford **4** (5 mg). The 30% acetone in CH₂Cl₂ fraction was rechromatographed (4×) using 20% EtOAc in petroleum ether to afford a mixture of **1b** and **3** (15 mg) after washing with petroleum ether. The 40% acetone in CH₂Cl₂ fraction was rechromatographed (2×) using CH₃CN:Et₂O:CH₂Cl₂ (0.5:0.5:9, v/v) afford a mixture of **6a** and **6b** (4 mg) after washing with petroleum ether. The 60% acetone in CH₂Cl₂ fraction was rechromatographed using CH₃CN:Et₂O:CH₂Cl₂ (1:1:8, v/v) afford **5** (3 mg) after washing with Et₂O.

Lupeol (1a)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ 4.66 (d, *J* = 2.4 Hz, H-29b), 4.55 (d, *J* = 2.4 Hz, H-29a), 3.18 (H-3), 0.98 (s, H₃-23), 0.78 (s, H₃-24), 0.83 (s, H₃-25), 0.95 (s, H₃-26), 1.05 (s, H₃-27), 0.90 (s, H₃-28), 1.66 (s, H₃-30).

α-Amyrin (1b)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ 3.15 (H-3), 0.67 (H-5), 5.12 (H-12), 0.95 (s, H₃-23), 0.76 (s, H₃-24), 0.75 (s, H₃-25), 0.89 (s, H₃-26), 1.01 (s, H₃-27), 0.95 (s, H₃-28), 0.85 (d, *J* = 6.0 Hz, H₃-29), 0.75 (d, *J* = 7.0 Hz, H₃-30).

β-Amyrin (1c)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ 3.15 (H-3), 0.67 (H-5), 5.16 (H-12), 0.77 (s, H₃-23), 0.90 (s, H₃-24), 0.74 (s, H₃-25), 0.93 (s, H₃-26), 1.16 (s, H₃-27), 1.07 (s, H₃-28), 0.86 (s, H₃-29), 0.79 (s, H₃-30).

Lupeol acetate (2a)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): δ 38.44 (C-1), 27.42 (C-2), 80.94 (C-3), 38.44 (C-4), 55.36 (C-5), 17.99

(C-6), 34.19 (C-7), 55.41 (C-8), 50.32 (C-9), 37.06 (C-10), 20.92 (C-11), 25.07 (C-12), 38.02 (C-13), 42.05 (C-14), 27.42 (C-15), 35.55 (C-16), 47.63 (C-17), 48.27 (C-18), 47.99 (C-19), 150.96 (C-20), 29.81 (C-21), 40.01 (C-22), 28.07 (C-23), 15.72 (C-24), 15.96 (C-25), 16.17 (C-26), 14.49 (C-27), 18.22, 18.25 (C-28, C-30), 109.34 (C-29), 171.01, 21.39 (OAc).

α-Amyrin acetate (2b)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): δ 38.44 (C-1), 27.59 (C-2), 80.96 (C-3), 37.69 (C-4), 55.24 (C-5), 18.22 (C-6), 32.85 (C-7), 40.01 (C-8), 47.63 (C-9), 36.77 (C-10), 23.35 (C-11), 124.30 (C-12), 139.61 (C-13), 42.05 (C-14), 26.58 (C-15), 28.05 (C-16), 33.73 (C-17), 59.04 (C-18), 39.63 (C-19), 39.59 (C-20), 31.23 (C-21), 41.52 (C-22), 28.73 (C-23), 16.85 (C-24), 15.72 (C-25), 16.73 (C-26), 23.21 (C-27), 28.73 (C-28), 17.50 (C-29), 21.39 (C-30), 171.01, 21.32 (OAc).

β-Amyrin acetate (2c)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): δ 39.59 (C-1), 27.93 (C-2), 80.8 (C-3), 39.59 (C-4), 55.24 (C-5), 18.19 (C-6), 33.73 (C-7), 38.37 (C-8), 47.4 (C-9), 35.55 (C-10), 23.59 (C-11), 121.62 (C-12), 145.20 (C-13), 42.05 (C-14), 28.73 (C-15), 27.93 (C-16), 32.85 (C-17), 59.0 (C-18), 40.01 (C-19), 41.52 (C-20), 31.23 (C-21), 42.01 (C-22), 29.69 (C-23), 15.72 (C-24), 15.72 (C-25), 16.84 (C-26), 23.59 (C-27), 28.73 (C-28), 17.50 (C-29), 21.39 (C-30), 170.8, 21.32 (OAc).

Baurenol (3)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): δ 36.87 (C-1), 27.69 (C-2), 79.05 (C-3), 38.88 (C-4), 50.41 (C-5), 24.15 (C-6), 116.43 (C-7), 145.35 (C-8), 48.22 (C-9), 35.33 (C-10), 16.85 (C-11), 32.42 (C-12), 37.69 (C-13), 41.52 (C-14), 28.87 (C-15), 37.69 (C-16), 32.04 (C-17), 54.88 (C-18), 35.20 (C-19), 32.04 (C-20), 29.67 (C-21), 31.52 (C-22), 27.53 (C-23), 14.66 (C-24), 12.98 (C-25), 23.65 (C-26), 22.66 (C-27), 39.65 (C-28), 25.63 (C-29), 22.55 (C-30).

Squalene (4)

Colorless oil. ¹³C NMR (150 MHz, CDCl₃): δ 25.69 (C-1), 131.24 (C-2), 124.31 (C-3), 26.66 (C-4), 39.74 (C-5), 134.89 (C-6), 124.41 (C-7), 26.77 (C-8), 39.76 (C-9), 135.10 (C-10), 124.31 (C-11), 28.28 (C-12), 17.68 (C-13), 16.04 (C-14), 16.00 (C-15).

Lutein (5)

Orange crystals. ¹H NMR (600 MHz, CDCl₃): δ 1.05 (s, 2 ring A CH₃), 0.83 (s, ring B CH₃), 0.98 (s, ring B CH₃), 1.60 (allylic CH₃), 1.71 (allylic CH₃), 1.89 (allylic CH₃), 1.951 (allylic CH₃), 1.94 (2 allylic CH₃), 1.45, 1.75 (CH₂), 1.35, 1.85 (CH₂), 2.35, 2.00 (allylic CH₂), 2.38 (allylic CH), 4.23 (br s, CHOH), 3.98 (m, CHOH), 5.52 (br s, =CH), 5.41 (dd, *J* = 9.6, 15.0 Hz, =CH), 6.56-6.65, 6.33 (dd, *J* = 15.0, 3.0 Hz), 6.23 (br d, *J* = 9.6 Hz), 6.09-6.14 (=CH).

β -Sitosterol (6a)

Colorless solid. ^1H NMR (600 MHz, CDCl_3): δ 3.50 (m, H-3), 2.26, 2.21 (H₂-4), 5.33 (dd, $J = 1.8, 4.8$ Hz, H-6), 0.66 (s, CH₃-18), 0.99 (s, CH₃-19), 0.90 (d, $J = 6.6$ Hz, CH₃-21), 0.79 (d, $J = 6.6$ Hz, CH₃-26), 0.82 (d, $J = 6.6$ Hz, CH₃-27), 0.86 (t, $J = 7.2$ Hz, CH₃-29).

Stigmasterol (6b)

Colorless solid. ^1H NMR (600 MHz, CDCl_3): δ 3.50 (m, H-3), 5.33 (dd, $J = 1.8, 4.8$ Hz, H-6), 0.68 (s, CH₃-18), 0.99 (s, CH₃-19), 1.01 (d, $J = 6.6$ Hz, CH₃-21), 5.13 (dd, $J = 8.4, 15.0$ Hz, H-22), 5.00 (dd, $J = 9.0, 15.0$ Hz, H-23), 0.84 (d, $J = 6.6$ Hz, CH₃-26), 0.83 (d, $J = 6.6$ Hz, CH₃-27), 0.80 (t, $J = 6.6$ Hz, CH₃-29).

RESULTS AND DISCUSSION

Silica gel chromatography of the dichloromethane extracts of *Hoya multiflora* Blume afforded a mixture of lupeol (**1a**) (Ragasa *et al.*, 2014a), α -amyrin (**1b**) (Ragasa *et al.*, 2014a) and β -amyrin (**1c**) (Ragasa *et al.*, 2014a) in about 2:1:0.3 ratio and another mixture of lupeol acetate (**2a**) (Tsai *et al.*, 2012), α -amyrin acetate (**2b**) (Ragasa *et al.*, 2014b) and β -amyrin acetate (**2c**) (Feleke and Brehane, 2005) in about 3:1:0.3 ratio from the stems; and a mixture of **1b** and bauerenol (**3**) (Raga *et al.*, 2013a) in about 1:2.5 ratio, squalene (**4**) (Ragasa *et al.*, 2014c), lutein (**5**) (Ragasa *et al.*, 2014d), and another mixture of β -sitosterol (**6a**) (Ragasa *et al.*, 2014e) and stigmasterol (**6b**) (Ragasa *et al.*, 2014e) in about 2:1 ratio from the leaves. The ratio of about 2:1:0.3 for the mixture of **1a**, **1b** and **1c** was deduced from integrations of the ^1H NMR resonances for the olefinic protons of **1a** at δ 4.55 (d, $J = 2.4$ Hz) and 4.66 (d, $J = 2.4$ Hz), **1b** at δ 5.10 (t, $J = 3.6$ Hz) and **1c** at δ 5.16 (t, $J = 3.6$ Hz). The integrations of the ^1H NMR resonances for the olefinic protons of **2a** at δ 4.55 (d, $J = 2.4$ Hz) and 4.66 (d, $J = 2.4$ Hz), **2b** at δ 5.11 (t, $J = 3.6$ Hz) and **2c** at δ 5.16 (t, $J = 3.6$ Hz) indicated that the ratio of **2a**, **2b** and **2c** is about 3:1:0.3. The 1:2.5 ratio of the mixture of **1b** and **3** was determined from the integrations of the ^1H NMR resonances for the olefinic protons of **1b** at δ 5.15 (t, $J = 3.6$ Hz) and **3** at δ 5.39 (dd, $J = 3.0, 7.2$ Hz, H-7). The integrations of the ^1H NMR resonances for the olefinic protons of **6a** at δ 5.33 (H-6) and **6b** at δ 5.33 (H-6), 5.13 (dd, $J = 8.4, 15.0$ Hz, H-22) and 5.00 (dd, $J = 8.4, 15.0$ Hz, H-23) suggested that the ratio of **6a** and **6b** is about 2:1. The structures of **1-6** were identified by comparison of their ^1H and/or ^{13}C NMR data with literature data.

Although no biological activity tests were conducted on the isolated compounds (**1-6**), literature search revealed that these have diverse bioactivities as follows. Lupeol (**1a**) exhibited antiurolithiatic and diuretic activity (Vidya *et al.*, 2002). It prevented the formation of vesical calculi and reduced the size of the preformed stones in rats (Anand *et al.*, 1994). It also showed antifungal activity against *Fusarium oxysporum* and *Penicillium notatum* (Manzano *et al.*, 2013). Lupeol significantly reduced the 451Lu tumor growth in athymic nude mice (Saleem *et al.*, 2008), inhibited the proliferation of MDA-MB-231 human breast cancer

cells in a dose dependent manner (Lambertini *et al.*, 2005), and induced growth inhibition and apoptosis in hepatocellular carcinoma SMMC7721 cells by down-regulation of the death receptor 3 (DR3) expression (Zhang *et al.*, 2009). Lupeol and lupeol acetate (**2a**) have shown hypotensive activity (Saleem *et al.*, 2003), while **1a** also exhibited antidyslipidemic activity in hamster at 100 mg/Kg body weight (Reddy *et al.*, 2009). It exhibited potent anti-inflammatory activity in an allergic airway inflammation model by a significant reduction in eosinophils infiltration and in Th2-associated cytokines levels that trigger the immune responses in asthma (Vasconcelos *et al.*, 2008). A review on the biological activities of lupeol has been provided (Gallo and Sarachine, 2009). β -Amyrin (**1c**) and α -amyrin (**1b**) were reported to possess anti-inflammatory (Recio *et al.*, 1995; Madeiros *et al.*, 2007; Okoye *et al.*, 2014) and analgesic (Otuki *et al.*, 2005; Soldi *et al.*, 2008) properties. β -Amyrin showed antifungal activity against *A. rabiei* with an MIC value of 0.0156 mg/mL (Jabeen *et al.*, 2011). α -Amyrin was proposed as a possible biomarker for the fungal resistance of grape-vine leaves (*Vitis vinifera*) (Batovska *et al.*, 2008). The mixture of **1b** and **1c** effectively reduced the elevated plasma glucose levels during the oral glucose tolerance test (OGTT). Furthermore, the mixture of **1b** and **1c** at 100 mg/kg significantly decreased the VLDL and LDL cholesterol and increased the HDL cholesterol (Santos *et al.*, 2012). A review on the sources and biological activities of **1b** and **1c** has been provided (Vasquez *et al.*, 2012). The anti-inflammatory effect of lupeol acetate (**2a**) involves the opioid system, as indicated by the complete blockade of the opioid antagonist naloxone (Lucetti *et al.*, 2010). α -Amyrin acetate (**2b**) at 100 mg/kg showed significant ($p < 0.05$) inhibition of egg albumen-induced paw edema with 40 % inhibition at the 5th hour. β -Amyrin acetate (**2c**) and **2b** isolated from the *Alstonia boonei* stem bark exhibited profound anti-inflammatory activity (Okoye *et al.*, 2014). Triterpenes **2b** and **2c** were also reported to exhibit sedative, anxiolytic and anticonvulsant properties (Aragao *et al.*, 2009). A mixture of bauerenol (**3**), **1b** and **1c** obtained from *Ardisia* species exhibited angio-suppressive effects on duck chorioallantoic membrane (CAM) (Raga *et al.*, 2013b); restricted inter-capillary length and reduced branch point with 100% CAM viability and embryo survivability and promoted intense expression of the von Willebrand factor (F8) (Raga *et al.*, 2013c); was found toxic to *A. salina* nauplii after 48h of exposure and showed teratologic manifestations on *Danio rerio* embryos (Raga *et al.*, 2014a); and exhibited analgesic property in the acetic acid writhing test and hot plate assay (Raga *et al.*, 2014b). Another study reported that a mixture of bauerenol, α -amyrin and β -amyrin from *Carmona retusa* exhibited 51% analgesic activity and showed 20% anti-inflammatory activity at dosage of 100 mg/kg mouse, while of 250 mg/kg mouse showed a 29% anti-diarrheal activity (Villasenor *et al.*, 2004). Squalene (**4**) was reported to significantly suppress colonic ACF formation and crypt multiplicity**y which strengthened the hypothesis that it possesses chemopreventive activity against colon carcinogenesis (Rao *et al.*, 1998).

It showed cardioprotective effect which is related to inhibition of lipid accumulation by its hypolipidemic properties and/or its antioxidant properties (Farvin *et al.*, 2006). A recent study reported that tocotrienols, carotenoids, squalene and coenzyme Q10 have anti-proliferative effects on breast cancer cells (Loganathan *et al.*, 2013). The preventive and therapeutic potential of squalene containing compounds on tumor promotion and regression have been reported (Desai *et al.*, 1996). A recent review on the bioactivities of squalene has been provided (Ronco and De Stéfani, 2013).

Dietary lutein (**5**), especially at 0.002%, inhibited tumor growth by selectively modulating apoptosis, and by inhibiting angiogenesis (Chew *et al.*, 2003). Another study reported that the chemopreventive properties of all-*trans* retinoic acid and lutein may be attributed to their differential effects on apoptosis pathways in normal *versus* transformed mammary cells (Sumantran *et al.*, 2000). Moreover, very low amounts of dietary lutein (0.002%) can efficiently decrease mammary tumor development and growth in mice (Park *et al.*, 1998). Another study reported that lutein and zeaxanthin reduces the risk of age related macular degeneration (SanGiovanni *et al.*, 2007).

β -Sitosterol (**6a**) was observed to have growth inhibitory effects on human breast MCF-7 and MDA-MB-231 adenocarcinoma cells (Awad *et al.*, 2007). It was shown to be effective for the treatment of benign prostatic hyperplasia (Jayaprakasha *et al.*, 2007). It was also reported to attenuate β -catenin and PCNA expression, as well as quench radical *in-vitro*, making it a potential anticancer drug for colon carcinogenesis (Baskar *et al.*, 2010). It can inhibit the expression of NPC1L1 in the enterocytes to reduce intestinal cholesterol uptake (Jesch *et al.*, 2009). It was reported to induce apoptosis mediated by the activation of ERK and the down regulation of Akt in MCA-102 murine fibrosarcoma cells (Moon *et al.*, 2007).

Stigmasterol (**6b**) shows therapeutic efficacy against Ehrlich ascites carcinoma bearing mice while conferring protection against cancer induced altered physiological conditions (Ghosh *et al.*, 2011). It lowers plasma cholesterol levels, inhibits intestinal cholesterol and plant sterol absorption, and suppresses hepatic cholesterol and classic bile acid synthesis in Wistar as well as WKY rats (Batta *et al.*, 2006). Other studies reported that stigmasterol showed cytostatic activity against Hep-2 and McCoy cells (Gómez *et al.*, 2001), markedly inhibited tumour promotion in two stage carcinogenesis experiments (Kasahara *et al.*, 1994), exhibited antimutagenic (Lim *et al.*, 2005), topical anti-inflammatory (Garcia *et al.*, 1999), anti-osteoarthritic (Gabay *et al.*, 2010) and antioxidant (Panda *et al.*, 2009) activities.

CONCLUSION

Hoya multiflora is a Philippine indigenous ornamental plant with no reported chemical studies and biological activities. This study reports on the terpenoids and sterols with known diverse biological activities which were isolated from the leaves and stems of the plant. Most of these compounds (**1a-1c**, **3-6**) were

reported to exhibit cytotoxic and anticancer properties, while **2a-2c** were reported to possess anti-inflammatory activity.

ACKNOWLEDGEMENT

A research grant from the De La Salle University Science Foundation through the University Research Coordination Office is gratefully acknowledged.

REFERENCES

- Anand R, Patnaik GK, Kulshreshtha DK, Dhawan N. Antirolithiatic activity of lupeol, the active constituent of *Cratevanuriala*. *Phytother Res* 1994; 8 (7):417-421.
- Aurigue FB. A Collection of Philippine Hoyas and Their Culture, Philippine Council for Agriculture, Aquatic and Natural Resources Research and Development (PCAARRD). Department of science and technology (DOST), 2013, 195 pages.
- Aragão GF, Carneiro LMV, Junior APF, Bandeira PN, Lemos ILG, Viana GSdB. Evidence for excitatory and inhibitory amino acids participation in the neuropharmacological activity of alpha- and beta-amyrin acetate. *The Open Pharmacol J* 2009; 3:9-16.
- Awad AB, Chinnman M, Fink CS, Bradford PG. β -Sitosterol activates Fas signaling in human breast cancer cells. *Phytomed* 2007; 14:747-754.
- Baas WJ, Van Berkel IEM, Versluis C, Heerma W, Kreyenbroek MN. Ring-A fissioned 3, 4-seco-3-nor-triterpene-2-aldehydes and related pentacyclic triterpenoids from the leaf wax of *Hoya australis*. *Phytochem* 1992; 31 (6):2073-2076.
- Baas WJ, Van Berkel IEM. 3, 4-Seco-triterpenoid acids and other constituents of the leaf wax of *Hoya naumanii*. *Phytochem* 1991; 30 (5):1625-1628.
- Baas WJ. Dihydronyctanthic acid methyl ester and other 3, 4-seco-pentacyclic triterpenoids from *Hoya lacunosa*. *Phytochem*, 1983; 22 (12):2809-2812.
- Baskar AA, Ignacimuthu S, Paulraj G, Numair K. Chemopreventive potential of β -sitosterol in experimental colon cancer model - an *in vitro* and *in vivo* study. *BMC Comp Alt Med* 2010; 10:24.
- Batovska DI, Todorova IT, Nedelcheva DV, Parushev SP, Atanassov AJ, Hvarleva TD, Djakova GJ, Bankova VS. Preliminary study on biomarkers for the fungal resistance in *Vitis vinifera* leaves, *J Plant Physiol* 2008; 165:791-795.
- Batta AK, Xu G, Honda A, Miyazaki T, Salen G. Stigmasterol reduces plasma cholesterol levels and inhibits hepatic synthesis and intestinal absorption in the rat. *Metabolism* 2006; 55 (3):292-299.
- Chew BP, Brown CM, Park JS, Mixer PF. Dietary lutein inhibits mouse mammary tumor growth by regulating angiogenesis and apoptosis. *Anticancer Res* 2003; 23 (4):3333-3339.
- Desai KN, Wei H, Lamartiniere CA. The preventive and therapeutic potential of the squalene-containing compound, Roindex, on tumor promotion and regression. *Cancer Lett* 1996; 101:93-96.
- Ebajo Jr V, Shen C-C, Ragasa CY. Triterpenes and sterol from *Hoya mindorensis*. *Der Pharma Chemica*. 2014, 6 (4):321-325.
- Farvin KHS, Anandan R, Hari S, Kumar S, Shing KS, Mathew S, Sankar TV, Nair PGV. Cardioprotective effect of squalene on lipid profile in isoprenaline-induced myocardial infarction in rats. *J Med Food* 2006; 9 (4):531-536.
- Feleke S, Brehane A. Triterpene compounds from the latex of *Ficus sur* I. *Bull Chem Soc Ethiop* 2005; 19 (2):307-310.
- Gabay O, Sanchez C, Salvat C, Chevy F, Breton M, Nourissat G. Stigmasterol: a phytosterol with potential anti-osteoarthritic properties. *Osteoarthritis Cartilage* 2010; 18 (1):106-116.
- Gallo MBC, Sarachine MJ. Biological activities of lupeol. *Intl J Biomed Pharm Sci* 2009; 3 (1):46-66.
- García MD, Sáenz MT, Gómez MA, Fernández MA. Topical anti-inflammatory activity of phytosterols isolated from *Eryngium*

foetidum on chronic and acute inflammation models. *Phytother Res* 1999; 13 (1):78–80.

Ghosh T, Maity TK, Singh J. Evaluation of antitumor activity of stigmaterol, a constituent isolated from *Bacopa monnieri* Linn aerial parts against ehrlich ascites carcinoma in mice. *Orient Pharm Exp Med* 2011; 11:41–49.

Gómez MA, García MD, Sáenz MT. Cytostatic activity of *Achillea ageratum* L. *Phytother Res* 2001; 15 (7):633–634.

Jabeen K, Javaid A, Ahmad E, Athar M. Antifungal compounds from *Melia azederach* leaves for management of *Ascochyta rabiei*, the cause of chickpea blight. *Nat Prod Res* 2011; 25 (3):264–276.

Jayaprakasha GK, Mandadi KK, Poulouse SM, Jadegoud Y, Gowda GA, Patil BS. Inhibition of colon cancer growth and antioxidant activity of bioactive compounds from *Poncirus trifoliata* (L.) Raf. *Bioorg Med Chem*, 2007; 15:4923–4932.

Jesch ED, Seo JM, Carr TP, Lee JY. Sitosterol reduces messenger RNA and protein expression levels of Niemann-Pick C1-like 1 in FHs 74 Int cells. *Nutr Res* 2009; 29 (12):859–66.

Kasahara Y, Kumaki K, Katagiri S, Yasukawa K, Yamanouchi S, Takido M. Carthami flos extract and its component, stigmaterol, inhibit tumour promotion in mouse skin two-stage carcinogenesis. *Phytother Res* 1994; 8 (6):327–331.

Lambertini E, Lampronti I, Penolazzi L, Khan MTH, Ather A, Giorgi G, Gambari R, Piva R. Expression of estrogen receptor gene in breast cancer cells treated with transcription factor decoy is modulated by Bangladeshi natural plant extracts. *Oncology Research* 2005; 14: 69–79

Lim J-C, Park JH, Budesinsky M, Kasal A, Han Y-H, Koo B-S, Lee S-I, Lee D-U. Antimutagenic constituents from the thorns of *Gleditsia sinensis*. *Chem Pharm Bull* 2005; 53 (5):561–564.

Loganathan R, Selvaduray KR, Nesaretnam K, Radhakrishnan A. Differential and antagonistic effects of palm tocotrienols and other phytonutrients (carotenoids, squalene and coenzyme Q10) on breast cancer cells *in vitro*. *J Oil Palm Res* 2013; 25:208–215.

Lucetti DL, Lucetti ECP, Bandeira MAM, Veras HNH, Silva AH, Leal LKAM, Lopes AA, Alves VCC, Silva GS, Brito GA, Viana OB. Anti-inflammatory effects and possible mechanism of action of lupeol acetate isolated from *Himatanthus drasticus* (Mart.) Plumel. *J Inflamm* 2010; 7:60.

Madeiras R, Otuki MF, Avellar MC, Calixto JB. Mechanisms underlying the inhibitory actions of the pentacyclic triterpene-amyrin in the mouse skin inflammation induced by phorbol ester 12-O-tetradecanoylphorbol-13-acetate. *Eur J Pharmacol* 2007;55 (9):227–235.

Manzano PI, Miranda M, Abreu-Payrol J, Silva M, Sterner O, Peralta EL. Pentacyclic triterpenoids with antimicrobial activity from the leaves of *Vernonanthura patens* (Asteraceae). *Emir J Food Agric* 2013; 25 (7):539–543

Moon DO, Kyeong Jun L, Yung HC, Gi-Young K. Moon DO, Kyeong Jun L, Yung HC, Gi-Young K. *Int Immunopharmacol* 2007; 7:1044–1053.

Okoye NN, Ajaghaku DL, Okeke HN, Ildogwe EE, Nworu CS, Okoye FBC. beta-Amyrin and alpha-amyrin acetate isolated from the stem bark of *Alstonia boonei* display profound anti-inflammatory activity. *Pharm Biol* 2014; 52 (11):1478–1486.

Otuki C, Ferreira J, Lima F, Meyre-Silva C, Malheiros A, Muller L, Cani G, Santos A, Yunes R, Calixto J. Antinociceptive properties of a mixture of α -amyrin and β -amyrin triterpenes: evidence for participation of protein kinase C and protein kinase A pathways. *J Pharmacol Exp Therapeutics* 2005; 31 (1):310–318.

Panda S, Jafri M, Kar A, Meheta BK. Thyroid inhibitory, anti-peroxidative and hypoglycemic effects of stigmaterol, isolated from *Butea monosperma*. *Fitoter* 2009; 80 (2):123–126.

Pandey SC, Singh SS, Ghosh AC, Deepak D, Khare AJ. *Med Arom Plant Sci* 2004; 26 (4):775–783.

Park JS, Chew BP, Wong TS. Dietary lutein from marigold extract inhibits mammary tumor development in BALB/c mice. *J Nutr* 1998; 128 (10):1650–1656.

Raga DD, Herrera AA, Ragasa CY. Angio-suppressive triterpenoids from *Ardisia cf. elliptica* (subgenus Tinus) on duck (*Anas*

platyrhynchus L.) chorioallantoic membrane. *Chin J Nat Med* 2013a; 11 (2):128–138.

Raga DD, Herrera AA, Shen C-C, Ragasa CY. Triterpenes from *Ardisia cf. elliptica* (subgenus Tinus) limit vascular density and promote von Willebrand factor expression on duck chorioallantoic membrane. *Pharm Chem J* 2013b; 47 (1):44–53.

Raga DD, Herrera AA, Dinah Espineli, Shen C-C, Ragasa CY. Triterpenes from *Ardisia squamulosa* C. Presl (Myrsinaceae) limit angiogenesis and the expression of Von Willebrand factor in duck chorioallantoic membrane. *J Chem Pharm Res*, 2013c; 5 (10):230–239.

Raga DD, Herrera AA, Alimboyoguen AB, Shen C-C, Ragasa CY. Effects of triterpenes from *Ardisia cf. elliptica* (subgenus Tinus) and sterols from *Ardisia pyramidalis* Cav Pers on *Artemia salina* and *Danio rerio* toxicity and caudal fin regeneration. *J Chem Pharm Res* 2014a; 6 (3):1014–1022.

Raga DD, Herrera AA, Shen C-C, Ragasa CY. Analgesic triterpenes from *Ardisia cf. elliptica* (subgenus Tinus) (Myrsinaceae). *Der Pharma Chemica* 2014b; 6 (4):153–161.

Ragasa CY, Caro JL, Shen C-C. Triterpenes and sterol from *Artocarpus ovatus*. *J Appl Pharm Sci* aawaaw2014a; 4 (10):7–11.

Ragasa CY, Ng VAS, De Los Reyes MM, Mandia EH, Shen C-C. Triterpenes and a coumarin derivative from *Kibatalia gitingensis* (Elm.) Woodson. *Der Pharma Chemica* 2014b; 6 (5):360–364.

Ragasa CY, Ng VAS, Ebajo Jr V, De Los Reyes MM, Mandia EH, Shen C-C. Chemical constituents of *Wrightia pubescens* (R.Br.). *Der Pharmacia Lettre* 2014c; 6 (6):14–19.

Ragasa CY, Torres OB, Mandia EH, Shen C-C. Chemical constituents of *Terminalia microcarpa*. *Der Pharmacia Lettre* 2014d, 6 (6):439–442.

Ragasa CY, Caro JL, Lirio LG, Shen CC. Chemical constituents of *Coix lacryma-jobi*. *Res J Pharm Biol Chem Sci* 2014e, 5 (6):344–348.

Rao CV, Mark HLN, Reddy RS. Chemopreventive effect of squalene on colon cancer. *Carcinogenesis* 1998; 19:287–290.

Recio MC, Giner RM, Manez S, Rios JL. Structural requirements for the anti-inflammatory activity of natural triterpenoids. *Planta Med* 1995; 61 (2):181–185.

Reddy KP, Singh AB, Puri A, Srivastava AK, Narender T. Synthesis of novel triterpenoid (lupeol) derivatives and their *in vivo* antihyperglycemic and antidiabetic activity. *Bioorg Med Chem Lett* 2009; 19:4463–4466.

Ronco AL, De Stéfani E. Squalene: a multi-task link in the crossroads of cancer and aging. *Functional Foods in Health and Disease* 2013; 3:462–476.

Saleem M, Maddodi N, Zaid MA, Khan N, Hafeez B, Asim M, Suh Y, Yun J, Setaluri V, Mukhtar H. Lupeol inhibits growth of highly aggressive human metastatic melanoma cells *in vitro* and *in vivo* by inducing apoptosis. *Cancer Therapy: Preclinical* 2008; 14:2119–2127.

Saleem R, Ahmad SI, Ahmed M, Faizi Z, Zikr-ur-Rehman S, Ali M, Faizi S. Hypotensive activity and toxicology of constituents from *Bombax ceiba* stem bark. *Biol Pharm Bull* 2003; 26: 41–46.

SanGiovanni JP, Chew EY, Clemons TE. The relationship of dietary carotenoid and vitamin A, E, and C intake with age-related macular degeneration in a case-control study: AREDS Report No. 22. *Arch Ophthalmol* 2007; 125 (9):1225–1232.

Santos FA, Frota JT, Arruda BR, de Melo TS, de Carvalho AA, da Silva A, Brito GAdC, Chaves MH, Rao VS. Antihyperglycemic and hypolipidemic effects of α , β -amyrin, a triterpenoid mixture, from *Protium heptaphyllum* in mice. *Lipids in Health and Disease* 2012; 11:98.

Soldi C, Pizzolatti G, Luiz A, Marcon R, Meotti F, Miotob L, Santos A. Synthetic derivatives of the α - and β -amyrin triterpenes and their antinociceptive properties. *Bioorg Med Chem* 2008; 16 (6):3377–3386.

Sumantran VN, Zhang R, Lee DS, Wicha MS. Differential regulation of apoptosis in normal *versus* transformed mammary epithelium by lutein and retinoic acid. *Cancer Epidemiol Biomarkers Prev* 2000; 9:257–263.

Tsai P-W, de Castro-Cruz K, Shen C-C, Ragasa CY. Chemical constituents of *Ficus odorata*. *Pharm Chem J* 2012b; 46 (4):225–227.

Vasconcelos JF, Teixeira MM, Barbosa-Filho JM, Lúcio ASSC, Almeida JRGS, Queiroz LP, Ribeiro-dos-Santos R, Soares MBP. The triterpenoid lupeol attenuates allergic airway inflammation in a murine model. *Intl Immunopharmacol* 2008; 8:1216-1221.

Vázquez LH, Palazon J, Navarro-Ocaña A. (2012). The pentacyclic triterpenes, α , β -amyryns: A review of sources and biological activities, *Phytochemicals - A Global Perspective of Their Role in Nutrition and Health*, Rao V (Ed.), ISBN: 978-953-51-0296-0, InTech, Available from: <http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-their-role-in-nutrition-and-health/the-pentacyclic-triterpenes-amyryns-a-review-of-sources-and-biological-activities>.

Vidya L, Leni M, Varalakshmi P. Evaluation of the effect of triterpenes on urinary risk factors of stone formation in pyridoxine hyperoxaluric rats. *Phytother Res* 2002; 16 (6):514-518.

Villasenor IM, Canlas AP, Faustino KM, Plana KG. Evaluation of the bioactivity of triterpene mixture isolated from *Carmona retusa* (Vahl.) Masam leaves. *J Ethnopharmacol* 2004; 92 (1):53-56.

Yoshikawa K, Nishino H, Arihara S, Chang HC, Wang JD, Oligosaccharides from *Hoya carmosa*. *J Nat Prod* 2000; 63 (1):146-148.

Zhang L, Zhang Y, Zhang L, Yang X, Lv Z. Lupeol, a dietary triterpene, inhibited growth, and induced apoptosis through down-regulation of DR3 in SMMC7721 cells. *Cancer Investigation* 2009; 27:163-170.

How to cite this article:

Ebajo Jr. V., Shen C-C., Ragasa C.Y. Terpenoids and Sterols from *Hoya multiflora* Blume. *J App Pharm Sci*, 2015; 5 (04): 033-039