

REVIEW

Open Access

TET proteins and the control of cytosine demethylation in cancer

Laurianne Scourzic^{1,2,3}, Enguerran Mouly^{1,2,3} and Olivier A Bernard^{1,2,3*}

Abstract

The discovery that ten-eleven translocation (TET) proteins are α -ketoglutarate-dependent dioxygenases involved in the conversion of 5-methylcytosines (5-mC) to 5-hydroxymethylcytosine (5-hmC), 5-formylcytosine and 5-carboxycytosine has revealed new pathways in the cytosine methylation and demethylation process. The description of inactivating mutations in *TET2* suggests that cellular transformation is in part caused by the deregulation of this 5-mC conversion. The direct and indirect deregulation of methylation control through mutations in DNA methyltransferase and isocitrate dehydrogenase (IDH) genes, respectively, along with the importance of cytosine methylation in the control of normal and malignant cellular differentiation have provided a conceptual framework for understanding the early steps in cancer development. Here, we review recent advances in our understanding of the cytosine methylation cycle and its implication in cellular transformation, with an emphasis on TET enzymes and 5-hmC. Ongoing clinical trials targeting the activity of mutated IDH enzymes provide a proof of principle that DNA methylation is targetable, and will trigger further therapeutic applications aimed at controlling both early and late stages of cancer development.

Introduction

DNA methylation on carbon 5 of cytosine is one of the best-studied epigenetic marks in mammals and is known to play crucial roles in cellular processes, including gene expression regulation and control of differentiation. However, variations in DNA methylation appear during normal differentiation and aging, and may contribute to

tumorigenesis. The processes of DNA methylation and demethylation as well as enzymes involved in these epigenetic mechanisms have been outlined recently but still need further characterization. Concomitantly, direct and indirect deregulation of methylation control has been uncovered in human malignancies from both hematopoietic and non-hematopoietic origins.

Ten-eleven translocation (TET) enzymes are 2-oxoglutarate, oxygen- and iron-dependent dioxygenases able to catalyze the oxidation of 5-methylcytosine (5-mC) into 5-hydroxymethylcytosine (5-hmC) [1,2]. They have been identified as key players in cytosine demethylation and in the control of cellular differentiation and transformation. Acquired point mutations and deletion events targeting TET genes are frequently observed in human cancers. These mutations affect TET2 and to some extent TET3 and result in partial or total inactivation of the gene. Metabolic perturbations resulting from mutations in genes encoding isocitrate dehydrogenase (IDH), fumarate hydratase (FH) or succinate dehydrogenase (SDH) also inhibit the TET enzymes and, in turn, DNA demethylation. Deregulation of DNA methylation may also be achieved directly through mutations in genes encoding DNA methyltransferase (DNMT) [3,4]. We are now starting to understand the control of TET protein activity, their DNA targeting, and their crosstalk with other epigenetic marks. For example, several proteins that interact with TET proteins (such as O-linked β -D-N acetylglucosamine transferase (OGT)) and with methylated and oxidized cytosines have been identified, highlighting their function in the regulation of chromatin structure. Following the implementation of specific detection methods, much has recently been learned regarding the quantity and location of the oxidized cytosine forms, mainly in embryonic stem (ES) cells, and we are now on the verge of a more complete understanding of their functions.

In this review, we discuss the established and emerging roles of TET enzymes and their functions in cytosine demethylation, with an emphasis on methylcytosine and its oxidized forms in normal tissues. We assess the roles of

* Correspondence: olivier.bernard@inserm.fr

¹Institut National de la Santé et de la Recherche Médicale (INSERM), Unité 1170, équipe labellisée Ligue Contre le Cancer, 94805 Villejuif, France

²Institut Gustave Roussy, 94805 Villejuif, France

Full list of author information is available at the end of the article

TET enzymes in hematological cancers and solid tumors, focusing on mutations involved in TET inactivation. Finally, we discuss the potential translational applications.

The cytosine methylation cycle

5-mC results from the transfer of a methyl group to cytosine within a CpG dinucleotide, mediated by DNMT enzymes encoded by five genes. DNMT1 is mainly responsible for the maintenance of genomic DNA methylation patterns (that is, after DNA replication), whereas DNMT2 (or tRNA cytosine-5-methyltransferase) is an RNA methyltransferase. DNMT3A and DNMT3B are mainly responsible for *de novo* DNA methylation [5]. However, all three enzymes may contribute to both maintenance and *de novo* DNA methylation [6]. The catalytically inactive DNMT3L interacts with these enzymes and the histone 3 tail to stimulate DNA methylation [7]. Furthermore, DNMT3A has recently been identified to be involved in crosstalk with epigenetic marks independently of DNMT3L [8].

Although DNA methylation has long been recognized, and cytosine methylation by DNMT3A and DNMT3B has been shown to be reversible *in vitro* [9], the mechanism of DNA demethylation was unclear until the functional analyses of the TET family proteins [1,2]. Due to its poor recognition of 5-hmC, which results from TET activity, DNMT1 is not able to perform the methylation of the neo-synthesized DNA strand (maintenance methylation). So the methylation information is lost in dividing cells, in a so-called passive manner (Figure 1). The three enzymes of the TET family (TET1, TET2 and TET3) are able to further oxidize 5-hmC into 5-formylcytosine (5-fC) and then 5-carboxycytosine (5-caC) [10,11]. Thymidine DNA glycosylase (TDG) is then able to remove 5-fC and 5-caC, triggering base-excision repair (BER) activity and the reintroduction of unmethylated cytosine [11-13]. The existence of decarboxylases that convert 5-caC to unmethylated cytosine is hypothetical. It has been suggested that the deamination of 5-hmC into 5-hydroxymethyluracil (5-hmU) occurs via activation-induced deaminase (AID) and apolipoprotein B mRNA editing enzyme (APOBEC), followed by TDG and BER mechanisms [14]. However, this remains controversial because 5-hmU residues may also originate from TET-mediated oxidation of thymine [15]. In addition, the activity of recombinant AID decreases with the size of the cytosine C5 electron cloud and does not show any activity on 5-hmC *in vitro* [16,17]. Indeed, AID exhibits its strongest activity against unmodified cytosine. Thymine resulting from deamination of 5-mC is not easily recognized by DNA repair machinery and is considered mutagenic. These branches of the cycle need to be further investigated in a cell- and tissue-dependent context. Regardless, TET proteins as well as several other

proteins (Table 1) are essential players in the demethylation of 5-mC.


DNA methylation distribution and localization

CpGs represent between 1% and 4% of genomic mammalian DNA and approximately 75% of them are methylated. Most CpGs are located in repetitive DNA elements, indicating that cytosine methylation is used as a defense mechanism against transposons and other parasitic elements to maintain the function and stability of the genome [18]. CpGs are also concentrated in CpG islands, which are mainly unmethylated and are associated with DNA accessibility and gene transcription. These CpG islands are usually found close to gene promoters, and their methylation status is strongly correlated with the transcription state of the genes nearby. Methylation can also be identified within gene bodies. It preferentially occurs in a CxG context (where x can be T, A or C) in ES cells and this intragenic methylation is mainly associated with highly expressed genes [19]. Promoter and gene body methylation are being extensively investigated to elucidate specific mechanisms and factors responsible for gene expression modulation. Recently, DNMT3B was reported to be involved in the remethylation of gene body-associated genes following treatment of a colon cancer cell line with DNMT inhibitors [20].

DNA hydroxymethylation distribution and localization

5-hmC was first identified in 1952 in bacteriophage T4 [21], and again 20 years later in the mammalian genome, in which it was found to constitute 0% to 17% of the total number of cytosine bases of brain-extracted DNA in mouse, rat and frog [22]. More recently, 5-hmC was estimated to constitute 0.6% of nucleotides in Purkinje cells, 0.2% in granule cells [1] and 0.03% in mouse ES cells [2]. However, the classical analyses of bisulfite-treated DNA do not discriminate between 5-mC and 5-hmC. Discrepancies among published studies may be due to different methodologies and analytical processes [23-26]. These studies nevertheless provide a general picture of the genome-wide distribution of modified cytosines in ES cells and other tissues.

The distribution of 5-hmC differs in several organs and tissues in mouse [27] and human [28]. The 5-hmC content also varies during development and cell differentiation [29]. For example, pluripotency correlates with high levels of 5-hmC, as observed in the inner cell mass, in multipotent adult stem cells as well as in progenitor cells. Embryonic or induced pluripotent stem cells also show a high 5-hmC level. Among differentiated cells, neuronal cells retain a high 5-hmC content [30,31]. In the blastocyst stage, erasure of DNA methylation and hydroxymethylation marks is followed by their re-establishment



by TET proteins and subsequent *de novo* methylation by DNMT3A and DNMT3B. In the totipotent zygote, the paternal pronucleus shows high levels of 5-hmC [32,33] caused by genome-wide hydroxylation of 5-mC [34,35], in contrast to the low level of 5-hmC in the maternal pronucleus. This phenomenon is linked to TET3 translocation from the cytoplasm to the paternal pronucleus immediately after fertilization [36]. In addition, the maternal factor PGC7 protects 5-mC from Tet3-mediated conversion to 5-hmC by binding to maternal chromatin containing dimethylated histone H3 lysine 9 [37]. The maternal genome undergoes progressive 5-mC demethylation upon cell division. Genome-wide mapping of 5-hmC with

specific antibodies or chemical labeling has enabled the design of 5-hmC distribution maps in mouse and human ES cells, as well as in neurological tissues. These studies have revealed that the 5-hmC mark is not uniformly distributed in the genome and is abundant in gene-rich euchromatin regions, particularly at promoters, exons and transcription start sites of genes expressed at low levels [38]. 5-hmC is mainly enriched in low CpG content regions and in bivalent gene promoters, characterized by both transcriptional permissive trimethylated histone H3 lysine 4 (H3K4me3) and repressive trimethylated histone H3 lysine 27 (H3K27me3) marks. Furthermore, *TET2*-mutated diffuse large B-cell lymphomas

Table 1 Functions and expression of human and murine proteins involved in the cytosine methylation/demethylation cycle

Proteins	Functions	Expression levels during development and in embryonic stem cells	Expression levels in adult tissues	Expression levels in adult hematopoiesis
TET1	Oxidization of 5-mC	High in mES cells, low in oocytes and zygotes [2,30,31]	Variable expression [56]	Low [83]
TET2	Oxidization of 5-mC	High in mES cells, low in oocytes and zygotes [2,30,31]	Widely expressed [56]	High [83]
TET3	Oxidization of 5-mC	Low in mES cells, high in oocytes and zygotes [2,30,31]	Limited expression in colon, stomach, adrenal glands and peripheral blood cells [56]	Low [83]
DNMT1	Methylation maintenance during DNA replication	High in early embryonic stages [129]	Ubiquitous [130,131]	Uniform but low in neutrophils [132]
DNMT3A	<i>De novo</i> methylation	High in early embryonic stages [129]	Ubiquitous [130,131]	Uniform but high in T lymphocytes and neutrophils [132]
DNMT3B	<i>De novo</i> methylation	High in later embryonic stages and differentiated cells [129]	Low levels except testis, thyroid and bone marrow [130,131]	Low expression except in human CD34+ cells [132]
AID	Cytidine deamination	High in immature B cells from fetal bone marrow and liver [133]	High in lymph nodes and moderated in spleen and bone marrow [134]	Mainly in activated mature B cells [134]
TDG	Glycosylation and deamination	Ubiquitous from ED 7.5 to 13.5 in mouse, high in central and peripheral nervous system, thymus, lung, liver, kidney, adrenal glands and intestine at ED 14.5 [135]	Mouse aorta [98]	Not reported
IDH1	Isocitrate decarboxylation of citric acid cycle	Not reported	Cytoplasm [92,103]	Not reported
IDH2	Isocitrate decarboxylation of citric acid cycle	Not reported	Mitochondria [92,103]	Not reported
FH	Hydration of fumarate of citric acid cycle	Not reported	Mitochondria [136]	Mature erythrocytes [136]
SDH	Oxidation of succinate of citric acid cycle	Not reported	Mitochondria [136]	Not reported

References listed in this table are based on mouse model studies, except for [56,103,83,130-133]. 5-mC, 5-methylcytosine; AID, activation-induced deaminase; DNMT, DNA methyltransferase; ED, embryonic day; FH, fumarate hydratase; IDH, isocitrate dehydrogenase; mES, murine embryonic stem; SDH, succinate dehydrogenase; TDG, thymine DNA glycosylase; TET, ten-eleven translocation.

have been associated with a hypermethylation signature on gene promoters identified as bivalent in human ES cells [39]. More recently, 5-hmC was identified in intergenic regions in human ES cells. More specifically, 5-hmC was found in regions comprising *cis*-regulatory elements, such as active enhancers, with co-localization of the histone modification marks monomethylated histone H3 lysine 4 (H3K4me1) and acetylated histone H3 lysine 27 (H3K27ac), and transcription factor binding sites for pluripotency factors, such as OCT4 and NANOG, or the insulator binding protein CTCF [40,41].

Function of oxidized cytosine forms

TET proteins participate in the regulation of gene transcription through the controlled generation of 5-hmC, 5-fC and 5-caC and their subsequent recognition by protein complexes involved in modulating chromatin structure or DNA repair [42-46].

Promoter methylation is associated with the repression of gene expression in somatic cells. It is not clear yet

whether specific readers of methylated DNA, such as methyl-CpG binding protein 2 (MeCP2), or methyl-binding domain (MBD) proteins are recruited to the methylated DNA and prevent the binding of transcription factors, or if they participate directly in the establishment of compact chromatin and gene repression. MBD1, MBD2 and MBD4 preferentially bind methylated DNA, in contrast to MBD3, MBD5 and MBD6 that prefer to bind to non-methylated DNA [47], although MBD5 and MBD6 associate with heterochromatin [48]. MBD4, possessing a DNA glycosylase domain, is also involved in BER following deamination events and is able to interact with mismatch repair proteins [49,50]. Methyl-CpG binding proteins were thought to be unable to bind 5-hmC [51] until recently [43], although the ability of MBD3 to specifically bind 5-hmC [45] is still controversial [44]. The DNA damage control proteins UHRF1 and UHRF2 also have 5-mC binding ability through their SET- and RING-associated domains. Additionally, UHRF1 is able to bind hemimethylated DNA and recruit DNMT1

[52,53]; it has recently been proposed that it may also be able to bind both methylated and hydroxymethylated DNA [42]. However, these 5-mC and 5-hmC readers are rarely found to be mutated in cancer (Table 2).

In ES cells, the distributions of 5-fC and 5-caC resemble those of 5-hmC, with a preference for enhancers,

and bivalent and silent promoters. Analyses of proteins interacting with cytosine-oxidized forms have identified glycosylase and DNA repair proteins interacting with 5-fC at a higher level compared with other cytosine forms, suggesting that 5-fC may trigger repair-associated removal [44].

Table 2 Somatic mutations affecting TET genes in cancer

Genes	Mutations in solid tumors	Mutations in hematological malignancies
TET genes		
<i>TET1</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], lung [106,138,139] and bladder cancer [140]	Rare in AML [141] and CLL [142]
<i>TET2</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], melanoma and lung cancer [106,138,139]	Frequent in various cancers (see Table 3)
<i>TET3</i>	Rare in endometrioid carcinoma and colorectal carcinomas [105,137]	Rare in CLL [142], PTCL [90] and T-ALL [143]
Epigenetic regulators of TET genes		
Methyltransferases		
<i>DNMT1</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], and lung cancer [106,138,139]	Rare in AML [141]
<i>DNMT3A</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], lung cancer [106,138,139] and 2% in non-small cell lung cancer [144]	AML [128], MDS [145] and T-cell lymphomas [90,91]
Deaminase and glycosylase		
<i>AID</i>	Rare in glioblastoma and medulloblastoma [146], endometrioid carcinoma ^a , colon cancer [105,137] and lung cancer [106,138,139]	T-ALL [147]
<i>TDG</i>	Rare in endometrioid carcinoma ^a , rare in glioblastoma ^a , colon cancer [104,136], lung [106,138,139] and thyroid cancer [148]	Not reported
Histone crosstalk regulators		
<i>OGT</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], lung cancer [106,138,139] and breast cancer [115,149]	DLBCL [150] and CLL [142]
<i>IDAX</i>	Rare in breast cancer ^a , glioblastoma ^a , endometrioid carcinoma ^a , kidney ^a , colon cancer [105,137], lung cancer [106,138,139] and 1.2% in mouth and pharynx carcinoma [117]	Not reported
5-mC and 5-hmC readers		
<i>MBD1</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], lung cancer [106,138,139], breast cancer [149] and melanoma [109]	Rare in ALL [151]
<i>MBD4</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137], lung cancer [106,138,139], breast cancer [115] and melanoma [109]	Rare in AML [141]
<i>UHRF1</i>	Rare in endometrioid carcinoma ^a , colorectal carcinomas [105,137] and lung cancer [106,138,139]	Rare in B-ALL [152]
Other genes affecting TET functions		
Metabolic enzymes		
<i>IDH1</i>	Rare in paragangliomas [153], frequent in chondrosarcomas [154], thyroid [155,156], prostate [157] and central nervous system cancers [102,158,159]	Frequent in AML [88], MDS [160], DLBCL [161] and B-ALL [162]
<i>IDH2</i>	Rare in endometrioid carcinoma ^a and colorectal carcinomas [105,137], frequent in chondrosarcomas [154] and central nervous system cancers [102,158,159]	Frequent in AML [88], MDS [160] and AITL [89]
<i>FH</i>	Renal cell carcinoma [163] and paragangliomas [104]	Not reported
<i>SDH</i>	Renal cell carcinoma [164] and paragangliomas [104]	Not reported

^aCOSMIC database. Some mutations listed in this table have not been confirmed as somatic mutations. 5-mC, 5-methylcytosine; 5-hmC, 5-hydroxymethylcytosine; AID, activation-induced deaminase; AML, acute myeloid leukemia; AITL, angioimmunoblastic T-cell lymphoma; B-ALL, B-cell acute lymphoblastic leukemia; CLL, chronic lymphocytic leukemia; DLBCL, diffuse large B-cell lymphoma; DNMT, DNA methyltransferase; FH, fumarate hydratase; IDAX, Inhibition of the Dvl and Axin complex; IDH, isocitrate dehydrogenase; MBD, methyl-binding domain; MDS, myelodysplastic syndrome; OGT, O-linked β -D-N acetylglucosamine transferase; PTCL, peripheral T-cell lymphoma; SDH, succinate dehydrogenase; T-ALL, T-cell acute lymphoblastic leukemia; TDG, thymine DNA glycosylase; TET, ten-eleven translocation; UHRF, ubiquitin-like with PHD and ring finger domains.

TET proteins

TET1 was first identified as a rare fusion partner of the mixed lineage leukemia gene, resulting from the chromosomal translocation t(10;11)(q22;23) in acute leukemia [2,54-57]. The difference between TET proteins relies on their structure (Figure 2) but also on their distinct expression patterns: TET2 is more highly expressed in the hematopoietic system than TET3 and TET1. It is currently thought that the common and main function of TET proteins is to establish or maintain protective boundaries to prevent unwanted methylation of non-methylated regions [58]. Each TET protein may also have specific functions: for example, TET1 oxidizes 5-mC to 5-hmC, and TET2 and TET3 stimulate the removal of 5-hmC [59]. In ES cells, TET2 may preferentially act on gene bodies, and TET1 at promoters [60]. The role of TET-mediated cytosine oxidation at distal enhancers is currently being thoroughly investigated. Super enhancers (enhancer clusters) that produce enhancer-transcribed RNAs in mouse ES cells have recently been associated with H3K27ac, TET1 and a decrease in DNA methylation level at pluripotency-dedicated loci [61]. Also, a specific role for TET2 in the control of enhancer activity has been suggested in the context of murine ES cell differentiation [62]. This mechanism remains to be investigated in the context of cancer, and more specifically in hematological disorders.

Interaction with other proteins

The stability and activity of TET proteins are regulated in several ways. Vitamin C has been reported as a cofactor that enhances the activity of TET enzymes [63,64]. The Dvl-binding protein inhibition of the Dvl and Axin complex (IDAX) can recruit TET2 to unmethylated DNA via the CXXC domain, and at the same time is able to induce its proteolytic degradation by caspase activation [65]. Other proteins interact with TET proteins, such as early B-cell factor 1 [66], or modulate their subcellular localization, such as AID [67], but it is not yet


clear whether they affect TET stability and function. This is also the case for OGT, which can associate with TET proteins [68-70] but appears to differently affect the three proteins. For instance, OGT has been described to trigger the export of TET3 from the nucleus and thus impair its activity [71]. A better understanding of multiple TET functions will arise from the identification of TET partners in normal and cancerous cellular contexts.

Crosstalk with other epigenetic mechanisms

In addition to transcriptional regulation through the readers of 5-hmC, 5-fC and 5-caC, another level of transcriptional regulation mediated by TETs comes from the interplay between DNA and histone modifiers. TET1 has been shown to interact with histone deacetylases through the transcriptional corepressor SIN3 transcription regulator family member A, thereby promoting transcriptional repression [72]. TET proteins can recruit OGT enzymes to chromatin, which catalyzes the addition of O-linked β -D-N acetylglucosamine to serine and threonine within histones and other proteins. TET proteins also interact indirectly with the complex proteins associated with SET1 (COMPASS) complex, which is responsible for mono-, di- and trimethylation of histone 3 lysine 4 and is associated with active transcription. This interaction occurs through the OGT-mediated glycosylation of the COMPASS subunit host cell factor 1. The COMPASS complex of proteins is involved in the regulation of master genes, such as *HOX*, during development, balanced by the action of the polycomb repressive complex (PRC), which catalyzes the repressive mark H3K27me3. In addition, TET1 shares target genes with PRC2 in ES cells [73]. In conclusion, TET proteins also serve as platforms for other epigenetic activities [74].

Other TET functions

The TET family is conserved during evolution. *Drosophila*, for example, has one homologous gene, whose function remains undetermined because of the particular DNA



methylation pattern of flies [75]. Additional TET functions might be uncovered in the future, and a recent report indicates that mammalian TET proteins may catalyze the formation of 5-hydroxymethylcytidine *in vitro*, suggesting a role in RNA modification [76]. Recently, TET triple knockout mouse ES cells were generated using the CRISPR/Cas9 system, suggesting a novel function of these proteins in telomere length regulation [77]. Indeed, triple knockout ES cells have an increased telomere length associated with a higher frequency of telomere-sister chromatid exchange. Although TET proteins seem to be involved in telomere shortening, their precise roles need to be further investigated in the context of both normal and cancerous cells.

TET and cancer

Here, we discuss the role of TET proteins in cancer focusing on TET2 mutations and activity impairment, first in hematopoietic malignancies and then in solid tumors.

TET in hematopoietic malignancies

TET mutations

Inactivation of *TET2* by genomic deletions or mutations has been reported in a wide range of adult hematological malignancies, including acute myeloid leukemia (AML), myelodysplastic syndrome (MDS) and myeloproliferative neoplasms (MPN) [78–80], as well as in lymphoid malignancies [39,81] (Table 3). In myeloid malignancies, *TET2* mutations are associated with a decrease in 5-hmC levels and an increase in 5-mC levels with respect to *TET2*-wild-type samples [82–84]. Many *TET2* acquired missense mutations have been described. Mutations that target the evolutionarily conserved catalytic domain of the protein are predicted to impair its function. Other missense mutations, occurring, for example, in the amino-terminal part of the protein, may also affect its function in an as yet uncharacterized manner. *TET2* mutations are observed on only one of the two gene copies, indicating that partial inactivation of the protein may contribute to cellular transformation [78]. There are marked differences between the three *TET* genes in terms of their expression levels. *TET2*, for example, has a higher expression level in hematological cells than *TET1* or *TET3*. *TET3* expression levels are higher than *TET1* levels in hematopoietic progenitor cells. Mutations in *TET3* have also been described but are much less frequent, probably because of its lower expression in hematopoiesis. Regarding *TET1*, most of the currently described mutations are missense mutations, whose functional consequences have not been established.

Associations with other mutations

Mouse and human studies have shown that the loss of *TET2* endows cells with a growth advantage over

wild-type cells, but does not lead to full transformation. Although this is not always the case, *TET2* mutation frequently occurs before the *JAK2*^{V617F} mutation in the development of MPN [78,85], suggesting that *TET2* mutation may occur very early in cancer development. *TET2* mutations also occur in early progenitors in MDS. Acquired *TET2* mutations are also observed in lymphoma, both B- and T-cell types, and particularly in angioimmunoblastic T-cell lymphoma (AITL). In both T- and B-cell lymphomas, *TET2* mutations have been identified in multipotent progenitors [86] that are able to participate in both myeloid and lymphoid differentiation. Together, these observations indicate that *TET2* loss predisposes but does not trigger cellular transformation. The tumor phenotype depends on cooperating mutations, such as *JAK2* or *KIT* mutations for MPN [87].

In AML, *TET2* mutations occur with other major mutations, particularly internal tandem duplication of *FLT3*, as well as mutations in *RAS*, *NPM1* and *DNMT3A*. Mutations in *TET2*, *IDH1* and *IDH2* are, however, mutually exclusive [88]. The situation is markedly different in AITL. Here, *TET2* mutations are closely associated with *DNMT3A* mutations [86] and, even more intriguing, do occur together with *IDH2* mutations [89–91].

TET and IDH mutations

IDH mutant proteins can inhibit *TET2* activity. The *IDH* genes encode enzymes of the citric acid cycle that convert isocitrate into α -ketoglutarate (α KG) in a nicotinamide adenine dinucleotide phosphate-dependent manner. A variety of human cancers, including AML [92–94], show recurrent missense mutations in *IDH1* and *IDH2* that endow the mutant protein with the ability to synthesize 2-hydroxyglutarate (2HG) from α KG (Table 2). 2HG is a competitive inhibitor of α KG and may inhibit all α KG-dependent dioxygenases, including EGLN prolyl hydroxylases, Jumanji C histone demethylases and TET proteins. In AML, *TET2* and *IDH* mutations are mutually exclusive, suggesting that they target the same pathway [84]. Consistent with this, *TET2*- and *IDH*-mutated primary AML samples show comparable DNA methylation profiles [84,95].

Other examples of TET2 activity targeting in myeloid malignancies

A recent report indicates that mutations in the *WT1* gene are exclusive from *TET2*, *IDH1* and *IDH2* mutations and impair *TET2* activity in human AML. The *WT1* gene encodes a zinc finger transcription factor and is mutated in approximately 8% of patients. Similar to patients with mutations in *IDH1*, *IDH2* and *TET2*, samples from patients with *WT1*-mutated primary AML show decreased 5-hmC levels and changes in 5-hmC localization. This study indicates the involvement of

Table 3 Prevalence of *TET1*, *TET2* and *TET3* mutations in hematological malignancies and solid tumors

Cancer	<i>TET1</i> mutation prevalence (%)	<i>TET2</i> mutation prevalence (%)	<i>TET3</i> mutation prevalence (%)
Myeloid malignancies			
MDS	Not reported	6-26 [78,79,165-169]	Not reported
MDS/MPN			
CMML (adult)	Not reported	20-58 [78,169-174]	Not reported
MPD			
PV	Not reported	6-16 [78,169,170]	Not reported
ET	Not reported	4-5 [78,169,170]	Not reported
MF	Not reported	2-17 [78,169,170]	Not reported
CML	Not reported	2-4 [175,176]	Not reported
AML			
<i>De novo</i> (adult)	Rare [141]	12-27 [169,170,177-181]	Not reported
<i>De novo</i> (pediatric)	Not reported	2-4 [182,183]	Not reported
Secondary AML	Not reported	17-32 [80,85,184,185]	Not reported
Mastocytosis	Not reported	20-29 [87,186]	Not reported
BPDCN	Not reported	25-54 [187-189]	Not reported
Lymphoid malignancies			
B-cell lymphoma			
DLBCL	Not reported	6-12 [39,81]	Not reported
MCL	Not reported	0-4 [81,190]	Not reported
Follicular lymphoma	Not reported	2 [81]	Not reported
CLL	Rare [142]	Not reported	Rare [142]
T cell lymphoma			
AITL	Not reported	33-83 [81,86,91,191,192]	Not reported
PTCL and PTCL, NOS	Not reported	20-49 [81,86,91,191,192]	Rare [90]
T-ALL	Not reported	Not reported	Rare [143]
Solid tumors from			
Endometrium	9*	7*	4*
Breast	Rare [106]	Rare [115]	Rare*
Central nervous system	Rare [193]	Rare*	Rare*
Kidney	Rare*	Rare [113]	Rare*
Large intestine	7 [105,137]	4 [105,137]	5 [105,137]
Liver	Rare [194]	Rare*	Rare [195]
Lung	5 [106,114,138]	2 [115,138,139]	Rare [138,139]
Ovary	Rare*	Rare*	Rare [112]
Pancreas	Rare [196,197]	Rare*	Rare*
Prostate	Rare [106,198]	Rare [198,199]	Rare [199]
Skin	Rare [109]	1 [109]	Rare [116]
Stomach	4 [200,201]	Not reported	Rare*
Urinary tract	4 [140]	4*	Rare*

AITL, angioimmunoblastic T cell lymphoma; AML, acute myeloid leukemia; BPDCN, blastic plasmacytoid dendritic cell neoplasm; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; CMML, chronic myelomonocytic leukemia; DLBCL, diffuse large B cell lymphoma; ET, essential thrombocytosis; FL, follicular lymphoma; MCL, mantle cell lymphoma; MDS, myelodysplastic syndrome; MF, myelofibrosis; MDS/MPN, myelodysplastic syndrome/myeloproliferative neoplasm; MPD, myeloproliferative disorder; PV, polycythemia vera; PTCL, peripheral T cell lymphoma; PTCL,NOS, peripheral T cell lymphoma not otherwise specified; T-ALL, T-cell acute lymphoblastic leukemia; TET, Ten eleven translocation. *COSMIC database. Some mutations listed in this table have not been confirmed as somatic mutations.

WT1 in the regulation of hydroxymethylation and provides an example of TET2 function impairment without *TET2* mutations [96].

Mouse models have shown that microRNAs (miRNAs) miR26a and miR29a are able to regulate TET expression by targeting their 3' untranslated regions (UTRs) [97,98]. Other miRNAs, such as miR125b, miR29b, miR29c, miR101 and miR7, have also been implicated in TET regulation using a 3' UTR human and mouse reporter screen [99]. Recently, miR22 has been shown to be responsible for the downregulation of all three TET genes [100]. Indeed, conditional expression of miR22 in a transgenic mouse model led to reduced levels of 5-hmC, amplification of the hematopoietic stem/progenitor compartment, and development of hematopoietic malignancies. miR22 is highly expressed in more than half of adult MDS and AML samples, providing another example that TET2 activity can be knocked down in the absence of a somatic mutation.

5-hmC and TET in solid tumors

Deregulation of cytosine hydroxymethylation by TET activity

Abnormal patterns of cytosine methylation have been observed in some solid tumors, including melanoma. The melanoma epigenome widely lacks 5-hmC, in association with tumor progression and downregulation of the *TET* family genes [101]. However, somatic *TET* mutations are exceedingly rare in this cancer, suggesting that another mechanism is affecting TET activity. Considering that TET enzymes are dependent on α KG, alteration in genes participating in its production may contribute to the inhibition of TET activity. Accordingly, *IDH1* or *IDH2* mutations are described in 10% of melanomas. These data support a role for deregulation of DNA methylation control during tumor progression rather than during the initial phases.

IDH mutations were first observed in human gliomas [102]. The *IDH*-mutated samples exhibited a hypermethylation phenotype, due to the inactivation of TET proteins by 2HG [103]. In paragangliomas, inactivating mutations in the *SDHx* and *FH* genes, encoding citric acid cycle enzymes (Table 2), result in the accumulation of succinate or fumarate, respectively, and competitive inhibition of α KG-dependent dioxygenases, similar to 2HG [104]. *SDH* mutations induce a hypermethylation phenotype compared to tumors with wild-type *SDH*, and are associated with transcriptional silencing. This argues for a driver role for demethylation deregulation in the development of these tumors.

TET mutations

TET mutations are rare in solid tumors [105–117]. In many instances, acquired mutations are missense mutations

whose functional consequences on TET protein activity are uncertain. A survey of *TET2* mutations in the COSMIC database showed more deleterious mutations in hematological malignancies than in solid tumors (29.8% versus 7.3% for frameshift mutations and 28.1% versus 10.3% for nonsense mutations). Conversely, there are fewer potentially benign mutations in hematological malignancies than in solid tumors (0.25% versus 17.6% in solid tumors for coding-silent mutations and 26.5% versus 63.1% for missense mutations). The dominant expression of *TET2* (with respect to *TET1* and *TET3*) in hematopoiesis results in a strong effect of *TET2* deficiency on 5-hmC levels. Aside from the potential specific functions of *TET2*, because expression of the three *TET* genes is equivalent in other tissues, the consequences of *TET2* deficiency on global cytosine (hydroxy)methylation is expected to be less important than in hematopoietic tissues. *IDH*, *SDH* and *FH* mutations, which result in the inhibition of virtually all α KG-dependent dioxygenases, including all three TET proteins, would therefore more strongly impact DNA methylation control than a single *TET* gene mutation.

Implications for disease

Studies of *TET2* deficiencies in tumor development have revealed the importance of DNA methylation in cellular processes as well as in the progressive development of adult type hematological malignancies.

In terms of potential clinical applicability, it appears difficult to specifically and directly target these TET dioxygenases for cancer treatment because they are inactivated in cancer. Indeed, recent efforts have focused on indirect correction of TET function and 5-hmC deregulation in cancer.

TET inactivation induces a methylation imbalance, including hypermethylation of tumor suppressor genes in malignant clones. These genes may be targeted by hypomethylating agents already used in clinical studies, such as 5-azacitidine and decitabine [118,119]. The global hypomethylation effect of these drugs, which remains nonspecific, seems to be accompanied by local hypermethylation, whose long-term consequences are unknown [20]. The molecular mechanisms of action of these drugs need to be further investigated, and extensive clinical trials are needed to prove their efficacy and to identify biomarkers of clinical responses.

In *IDH1*- or *IDH2*-mutated cancers, the oncometabolite 2HG acts as a biomarker of compromised enzyme activity [120]. This led to the development of *IDH2* inhibitors, now tested in clinical trials [121]. Similarly, *FH* and *SDH* inhibitors could be developed to prevent the overall effect of metabolic TET inactivation in cancer. The activities of TET as well as DNMT enzymes are regulated, in part, by the concentrations of their required cofactors. Thus, the metabolic state of the cell is an


antitumor target, by preventing the activity of the mutated protein but also by manipulating agonist or antagonist functions. In addition to the detection of *TET2* mutations that pre-date full-blown malignancies, recent studies have highlighted preleukemic phases in AML that are associated with mutations in other genes affecting DNA methylation, such as *DNMT3A*, *IDH1* and *IDH2* [122], and in genes involved in chromatin structure, such as *SMC1A* (structural maintenance of chromosome 1A) [123]. These observations suggest that manipulating the control of chromatin structure may be efficient for the treatment of both early and late phases of disease.

Conclusions and future directions

DNA methylation patterns are markedly abnormal in malignant cells in comparison with normal tissues. Abnormal

methylation has been postulated to inactivate tumor suppressor genes through cytosine methylation and to activate oncogenes through cytosine hydroxymethylation and demethylation (Figure 3). An unexpected number of oxidized cytosine forms have been uncovered, whose specific functions need to be investigated. Specific techniques allowing their thorough investigation at the nucleotide level are under development and will enable us to investigate the specific functions of these cytosines in normal cells. This is a requirement for understanding their roles in cellular transformation, because mutations detected in cancer can inactivate or impair DNA methylation (for example, *DNMT3A* mutations) or DNA demethylation (for example, *TET2* or *IDH* mutations).

Although cytosine methylation is associated with gene repression, the exact mechanisms leading from



methylation to gene repression are still elusive, as are the crosstalk with other epigenetic modifications, the factors involved in chromatin modification and the regulation of their activities. DNA methylation and demethylation result from the regulation of different enzymatic activities, which compete with each other for DNA access.

This process is complex enough to appear stochastic, with a slow drift in methylation profiles that is tissue specific as well as age related. This drift leads to cellular heterogeneity and, with respect to methylation and gene repression, allows for cell selection. It is likely that *TET2* deficiency increases cellular heterogeneity and facilitates the selection of fitter cells. We now require a complete understanding of the protein complexes involved in cytosine methylation and demethylation, including the exact role of each of the mammalian TET proteins and their regulatory signals, in order to target these processes.

Work with human samples and murine models has shown that *TET2* deficiency does not trigger full-blown malignancies but predisposes to the development of tumors such as MPN, MDS and lymphoma. The different *Tet2* knockout mouse models exhibit similar phenotypes but also present subtle differences that might be due to the loss of different regions of the gene. So far, it has not been possible to correlate clinical phenotypes with *TET2* mutations (for example, regarding their precise location or heterozygosity). Other questions relate to the dependency of malignant cells on the initial *TET2* mutation, and why some patients with *TET2* mutations remain healthy while others develop a myeloid or a lymphoid malignancy [123–126]. Addressing such questions is important, not only with regard to mutations in genes involved in the control of DNA methylation (such as *TET2* or *DNMT3A*), but also for mutation in genes controlling other functions that predate and may predispose to the development of adult malignancies [125–127].

Abbreviations

2HG: 2-hydroxyglutarate; 5-caC: 5-carboxycytosine; 5-fC: 5-formylcytosine; 5-hmC: 5-hydroxymethylcytosine; 5-hmU: 5-hydroxymethyluracil; 5-mC: 5-methylcytosine; αKG: α-ketoglutarate; AID: Activation-induced deaminase; ATL: Angioimmunoblastic T-cell lymphoma; AML: Acute myeloid leukemia; BER: Base-excision repair; COSMIC: Catalogue of somatic mutations in cancer; DNMT: DNA methyltransferase; ES: Embryonic stem; FH: Fumarate hydratase; IDH: Isocitrate dehydrogenase; MBD: Methyl-binding domain; MBP: Methyl-CpG binding; MDS: Myelodysplastic syndrome; miRNA: microRNA; MPN: Myeloproliferative neoplasms; OGT: O-linked β-D-N acetylglucosamine transferase; PRC2: Polycomb repressive complex 2; SDH: Succinate dehydrogenase; TDG: Thymidine DNA glycosylase; TET: Ten-eleven translocation; UTR: Untranslated region.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LS and OAB drafted the manuscript. All authors participated and agreed to the final version.

Acknowledgements

Work in the laboratory was supported by grants from Institut National de la Santé Et de la Recherche Médicale (INSERM), Institut Gustave Roussy, Institut National du Cancer (INCa) - INCa-DGOS-INSERM 6043, Fondation pour la recherche Médicale (FRM) and Association Laurette Fugain. LS is supported by fellowships from Cancéropôle Ile de France and Fondation ARC. We thank Philippe Dessen for help with COSMIC data.

Author details

¹Institut National de la Santé et de la Recherche Médicale (INSERM), Unité 1170, équipe labellisée Ligue Contre le Cancer, 94805 Villejuif, France.

²Institut Gustave Roussy, 94805 Villejuif, France. ³University Paris 11 Sud, 91405 Orsay, France.

Published online: 29 January 2015

References

- Kriaucionis S, Heintz N. The nuclear DNA base 5-hydroxymethylcytosine is present in Purkinje neurons and the brain. *Science*. 2009;324:929–30.
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, et al. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science*. 2009;324:930–5.
- Russler-Germain DA, Spencer DH, Young MA, Lamprecht TL, Miller CA, Fulton R, et al. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell*. 2014;25:442–54.
- Heyn H, Vidal E, Sayols S, Sanchez-Mut JV, Moran S, Medina I, et al. Whole-genome bisulfite DNA sequencing of a DNMT3B mutant patient. *Epigenetics*. 2012;7:542–50.
- Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell*. 1999;99:247–57.
- Arand J, Spieler D, Karius T, Branco MR, Meilinger D, Meissner A, et al. In vivo control of CpG and non-CpG DNA methylation by DNA methyltransferases. *PLoS Genet*. 2012;8:e1002750.
- Ooi SK, Qiu C, Bernstein E, Li K, Jia D, Yang Z, et al. DNMT3L connects unmethylated lysine 4 of histone H3 to de novo methylation of DNA. *Nature*. 2007;448:714–7.
- Guo X, Wang L, Li J, Ding Z, Xiao J, Yin X, et al. Structural insight into autoinhibition and histone H3-induced activation of DNMT3A. *Nature*. 2014. doi:10.1038/nature13899.
- Chen CC, Wang KY, Shen CK. The mammalian de novo DNA methyltransferases DNMT3A and DNMT3B are also DNA 5-hydroxymethylcytosine dehydroxymethylases. *J Biol Chem*. 2012;287:33116–21.
- Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, et al. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science*. 2011;333:1300–3.
- He YF, Li BZ, Li Z, Liu P, Wang Y, Tang Q, et al. Tet-mediated formation of 5-carboxylcytosine and its excision by TDG in mammalian DNA. *Science*. 2011;333:1303–7.
- Maiti A, Drohat AC. Thymine DNA glycosylase can rapidly excise 5-formylcytosine and 5-carboxylcytosine: potential implications for active demethylation of CpG sites. *J Biol Chem*. 2011;286:35334–8.
- Zhang L, Lu X, Lu J, Liang H, Dai Q, Xu GL, et al. Thymine DNA glycosylase specifically recognizes 5-carboxylcytosine-modified DNA. *Nat Chem Biol*. 2012;8:328–30.
- Guo JU, Su Y, Zhong C, Ming GL, Song H. Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain. *Cell*. 2011;145:423–34.
- Pfaffeneder T, Spada F, Wagner M, Brandmayr C, Laube SK, Eisen D, et al. Tet oxidizes thymine to 5-hydroxymethyluracil in mouse embryonic stem cell DNA. *Nat Chem Biol*. 2014;10:574–81.
- Nabel CS, Jia H, Ye Y, Shen L, Goldschmidt HL, Stivers JT, et al. AID/APOBEC deaminases disfavor modified cytosines implicated in DNA demethylation. *Nat Chem Biol*. 2012;8:751–8.
- Rangam G, Schmitz KM, Cobb AJ, Petersen-Mahrt SK. AID enzymatic activity is inversely proportional to the size of cytosine C5 orbital cloud. *PLoS One*. 2012;7:e43279.

18. Ehrlich M, Gama-Sosa MA, Huang LH, Midgett RM, Kuo KC, McCune RA, et al. Amount and distribution of 5-methylcytosine in human DNA from different types of tissues of cells. *Nucleic Acids Res.* 1982;10:2709–21.
19. Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, et al. Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature.* 2009;462:315–22.
20. Yang X, Han H, De Carvalho DD, Lay FD, Jones PA, Liang G. Gene body methylation can alter gene expression and is a therapeutic target in cancer. *Cancer Cell.* 2014;26:577–90.
21. Wyatt GR, Cohen SS. A new pyrimidine base from bacteriophage nucleic acids. *Nature.* 1952;170:1072–3.
22. Penn NW, Suwalski R, O'Riley C, Bojanowski K, Yura R. The presence of 5-hydroxymethylcytosine in animal deoxyribonucleic acid. *Biochem J.* 1972;126:781–90.
23. Booth MJ, Branco MR, Ficz G, Oxley D, Krueger F, Reik W, et al. Quantitative sequencing of 5-methylcytosine and 5-hydroxymethylcytosine at single-base resolution. *Science.* 2012;336:934–7.
24. Booth MJ, Marsico G, Bachman M, Beraldi D, Balasubramanian S. Quantitative sequencing of 5-formylcytosine in DNA at single-base resolution. *Nat Chem.* 2014;6:435–40.
25. Mohn F, Weber M, Schubeler D, Roloff TC. Methylated DNA immunoprecipitation (MeDIP). *Methods Mol Biol.* 2009;507:55–64.
26. Flusberg BA, Webster DR, Lee JH, Travers KJ, Olivares EC, Clark TA, et al. Direct detection of DNA methylation during single-molecule, real-time sequencing. *Nat Methods.* 2010;7:461–5.
27. Globisch D, Munzel M, Muller M, Michalak S, Wagner M, Koch S, et al. Tissue distribution of 5-hydroxymethylcytosine and search for active demethylation intermediates. *PLoS One.* 2010;5:e15367.
28. Li W, Liu M. Distribution of 5-hydroxymethylcytosine in different human tissues. *J Nucleic Acids.* 2011;2011:870726.
29. Ruzov A, Tsenkina Y, Serio A, Dudnakova T, Fletcher J, Bai Y, et al. Lineage-specific distribution of high levels of genomic 5-hydroxymethylcytosine in mammalian development. *Cell Res.* 2011;21:1332–42.
30. Ito S, D'Alessio AC, Taranova OV, Hong K, Sowers LC, Zhang Y. Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature.* 2010;466:1129–33.
31. Koh KP, Yabuuchi A, Rao S, Huang Y, Cunneiff K, Nardone J, et al. Tet1 and Tet2 regulate 5-hydroxymethylcytosine production and cell lineage specification in mouse embryonic stem cells. *Cell Stem Cell.* 2011;8:200–13.
32. Iqbal K, Jin SG, Pfeifer GP, Szabo PE. Reprogramming of the paternal genome upon fertilization involves genome-wide oxidation of 5-methylcytosine. *Proc Natl Acad Sci U S A.* 2011;108:3642–7.
33. Wossidlo M, Nakamura T, Lepikhov K, Marques CJ, Zakhartchenko V, Boiani M, et al. 5-Hydroxymethylcytosine in the mammalian zygote is linked with epigenetic reprogramming. *Nat Commun.* 2011;2:241.
34. Mayer W, Niveleau A, Walter J, Fundele R, Haaf T. Demethylation of the zygotic paternal genome. *Nature.* 2000;403:501–2.
35. Oswald J, Engemann S, Lane N, Mayer W, Olek A, Fundele R, et al. Active demethylation of the paternal genome in the mouse zygote. *Curr Biol.* 2000;10:475–8.
36. Gu TP, Guo F, Yang H, Wu HP, Xu GF, Liu W, et al. The role of Tet3 DNA dioxygenase in epigenetic reprogramming by oocytes. *Nature.* 2011;477:606–10.
37. Nakamura T, Liu YJ, Nakashima H, Umehara H, Inoue K, Matoba S, et al. PGC7 binds histone H3K9me2 to protect against conversion of 5mC to 5hmC in early embryos. *Nature.* 2012;486:415–9.
38. Yu M, Hon GC, Szulwach KE, Song CX, Zhang L, Kim A, et al. Base-resolution analysis of 5-hydroxymethylcytosine in the mammalian genome. *Cell.* 2012;149:1368–80.
39. Asmar F, Punj V, Christensen J, Pedersen MT, Pedersen A, Nielsen AB, et al. Genome-wide profiling identifies a DNA methylation signature that associates with TET2 mutations in diffuse large B-cell lymphoma. *Haematologica.* 2013;98:1912–20.
40. Stroud H, Feng S, Morey Kinney S, Pradhan S, Jacobsen SE. 5-Hydroxymethylcytosine is associated with enhancers and gene bodies in human embryonic stem cells. *Genome Biol.* 2011;12:R54.
41. Szulwach KE, Li X, Li Y, Song CX, Han JW, Kim S, et al. Integrating 5-hydroxymethylcytosine into the epigenomic landscape of human embryonic stem cells. *PLoS Genet.* 2011;7:e1002154.
42. Frauer C, Hoffmann T, Bultmann S, Casa V, Cardoso MC, Antes I, et al. Recognition of 5-hydroxymethylcytosine by the Uhrf1 SRA domain. *PLoS One.* 2011;6:e21306.
43. Mellen M, Ayata P, Dewell S, Kriaucionis S, Heintz N. MeCP2 binds to 5hmC enriched within active genes and accessible chromatin in the nervous system. *Cell.* 2012;151:1417–30.
44. Spruijt CG, Gnerlich F, Smits AH, Pfaffeneder T, Jansen PW, Bauer C, et al. Dynamic readers for 5-(hydroxy)methylcytosine and its oxidized derivatives. *Cell.* 2013;152:1146–59.
45. Yildirim O, Li R, Hung JH, Chen PB, Dong X, Ee LS, et al. Mbd3/NURD complex regulates expression of 5-hydroxymethylcytosine marked genes in embryonic stem cells. *Cell.* 2011;147:1498–510.
46. Iurlaro M, Ficz G, Oxley D, Raiber EA, Bachman M, Booth MJ, et al. A screen for hydroxymethylcytosine and formylcytosine binding proteins suggests functions in transcription and chromatin regulation. *Genome Biol.* 2013;14:R119.
47. Hendrich B, Bird A. Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Mol Cell Biol.* 1998;18:6538–47.
48. Laget S, Joulie M, Le Masson F, Sasai N, Christians E, Pradhan S, et al. The human proteins MBD5 and MBD6 associate with heterochromatin but they do not bind methylated DNA. *PLoS One.* 2010;5:e11982.
49. Bellacosa A, Cicchillitti L, Schepis F, Riccio A, Yeung AT, Matsumoto Y, et al. MED1, a novel human methyl-CpG-binding endonuclease, interacts with DNA mismatch repair protein MLH1. *Proc Natl Acad Sci U S A.* 1999;96:3969–74.
50. Hendrich B, Hardeland U, Ng HH, Jiricny J, Bird A. The thymine glycosylase MBD4 can bind to the product of deamination at methylated CpG sites. *Nature.* 1999;401:301–4.
51. Valinluck V, Tsai HH, Rogstad DK, Burdzy A, Bird A, Sowers LC. Oxidative damage to methyl-CpG sequences inhibits the binding of the methyl-CpG binding domain (MBD) of methyl-CpG binding protein 2 (MeCP2). *Nucleic Acids Res.* 2004;32:4100–8.
52. Bostick M, Kim JK, Esteve PO, Clark A, Pradhan S, Jacobsen SE. UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science.* 2007;317:1760–4.
53. Sharif J, Muto M, Takebayashi S, Suetake I, Iwamatsu A, Endo TA, et al. The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature.* 2007;450:908–12.
54. Burmeister T, Meyer C, Schwartz S, Hofmann J, Molkenin M, Kowarz E, et al. The MLL recombinome of adult CD10-negative B-cell precursor acute lymphoblastic leukemia: results from the GMALL study group. *Blood.* 2009;113:4011–5.
55. Ittel A, Jeandrier E, Helias C, Perrusson N, Humbrecht C, Liou B, et al. First description of the t(10;11)(q22;q23)/MLL-TET1 translocation in a T-cell lymphoblastic lymphoma, with subsequent lineage switch to acute myelomonocytic myeloid leukemia. *Haematologica.* 2013;98:e166–8.
56. Lorsbach RB, Moore J, Mathew S, Raimondi SC, Mukatira ST, Downing JR. TET1, a member of a novel protein family, is fused to MLL in acute myeloid leukemia containing the t(10;11)(q22;q23). *Leukemia.* 2003;17:637–41.
57. Ono R, Taki T, Taketani T, Taniwaki M, Kobayashi H, Hayashi Y. LCX, leukemia-associated protein with a CXXC domain, is fused to MLL in acute myeloid leukemia with trilineage dysplasia having t(10;11)(q22;q23). *Cancer Res.* 2002;62:4075–80.
58. Jeong M, Sun D, Luo M, Huang Y, Challen GA, Rodriguez B, et al. Large conserved domains of low DNA methylation maintained by Dnmt3a. *Nat Genet.* 2014;46:17–23.
59. Putiri EL, Tiedemann RL, Thompson JJ, Liu C, Ho T, Choi JH, et al. Distinct and overlapping control of 5-methylcytosine and 5-hydroxymethylcytosine by the TET proteins in human cancer cells. *Genome Biol.* 2014;15:R81.
60. Huang Y, Chavez L, Chang X, Wang X, Pastor WA, Kang J, et al. Distinct roles of the methylcytosine oxidases Tet1 and Tet2 in mouse embryonic stem cells. *Proc Natl Acad Sci U S A.* 2014;111:1361–6.
61. Pulakanti K, Pinello L, Stelloh C, Blinka S, Allred J, Milanovich S, et al. Enhancer transcribed RNAs arise from hypomethylated, Tet-occupied genomic regions. *Epigenetics.* 2013;8:1303–20.
62. Hon GC, Song CX, Du T, Jin F, Selvaraj S, Lee AY, et al. 5mC oxidation by Tet2 modulates enhancer activity and timing of transcriptome reprogramming during differentiation. *Mol Cell.* 2014;56:286–97.
63. Yin R, Mao SQ, Zhao B, Chong Z, Yang Y, Zhao C, et al. Ascorbic acid enhances Tet-mediated 5-methylcytosine oxidation and promotes DNA demethylation in mammals. *J Am Chem Soc.* 2013;135:10396–403.

64. Minor EA, Court BL, Young JL, Wang G. Ascorbate induces ten-eleven translocation (Tet) methylcytosine dioxygenase-mediated generation of 5-hydroxymethylcytosine. *J Biol Chem*. 2013;288:13669–74.
65. Ko M, An J, Bandukwala HS, Chavez L, Aijo T, Pastor WA, et al. Modulation of TET2 expression and 5-methylcytosine oxidation by the CXXC domain protein IDAX. *Nature*. 2013;497:122–6.
66. Guilhamon P, Eskandarpour M, Halai D, Wilson GA, Feber A, Teschendorff AE, et al. Meta-analysis of IDH-mutant cancers identifies EBF1 as an interaction partner for TET2. *Nat Commun*. 2013;4:2166.
67. Arioka Y, Watanabe A, Saito K, Yamada Y. Activation-induced cytidine deaminase alters the subcellular localization of Tet family proteins. *PLoS One*. 2012;7:e45031.
68. Chen Q, Chen Y, Bian C, Fujiki R, Yu X. TET2 promotes histone O-GlcNAcylation during gene transcription. *Nature*. 2013;493:561–4.
69. Deplus R, Delatte B, Schwinn MK, Defrance M, Mendez J, Murphy N, et al. TET2 and TET3 regulate GlcNAcylation and H3K4 methylation through OGT and SET1/COMPASS. *EMBO J*. 2013;32:645–55.
70. Vella P, Scelfo A, Jammula S, Chiacchiera F, Williams K, Cuomo A, et al. Tet proteins connect the O-linked N-acetylglucosamine transferase Ogt to chromatin in embryonic stem cells. *Mol Cell*. 2013;49:645–56.
71. Zhang Q, Liu X, Gao W, Li P, Hou J, Li J, et al. Differential regulation of the ten-eleven translocation (TET) family of dioxygenases by O-linked beta-N-acetylglucosamine transferase (OGT). *J Biol Chem*. 2014;289:5986–96.
72. Williams K, Christensen J, Helin K. DNA methylation: TET proteins-guardians of CpG islands? *EMBO Rep*. 2012;13:28–35.
73. Wu H, D'Alessio AC, Ito S, Xia K, Wang Z, Cui K, et al. Dual functions of Tet1 in transcriptional regulation in mouse embryonic stem cells. *Nature*. 2011;473:389–93.
74. Hanover JA, Krause MW, Love DC. Bittersweet memories: linking metabolism to epigenetics through O-GlcNAcylation. *Nat Rev Mol Cell Biol*. 2012;13:312–21.
75. Takayama S, Dhahbi J, Roberts A, Mao G, Heo SJ, Pachter L, et al. Genome methylation in *D. melanogaster* is found at specific short motifs and is independent of DNMT2 activity. *Genome Res*. 2014;24:821–30.
76. Fu L, Guerrero CR, Zhong N, Amato N, Liu Y, Liu S, et al. Tet-mediated formation of 5-hydroxymethylcytosine in RNA. *J Am Chem Soc*. 2014;136:11582–5.
77. Lu F, Liu Y, Jiang L, Yamaguchi S, Zhang Y. Role of Tet proteins in enhancer activity and telomere elongation. *Genes Dev*. 2014;28:2103–19.
78. Delhommeau F, Dupont S, Della Valle V, James C, Trannoy S, Masse A, et al. Mutation in TET2 in myeloid cancers. *N Engl J Med*. 2009;360:2289–301.
79. Langemeijer SM, Kuiper RP, Berends M, Knops R, Aslanyan MG, Massop M, et al. Acquired mutations in TET2 are common in myelodysplastic syndromes. *Nat Genet*. 2009;41:838–42.
80. Tefferi A, Pardanani A, Lim KH, Abdel-Wahab O, Lasho TL, Patel J, et al. TET2 mutations and their clinical correlates in polycythemia vera, essential thrombocythemia and myelofibrosis. *Leukemia*. 2009;23:905–11.
81. Quivoron C, Couronne L, Della Valle V, Lopez CK, Plo I, Wagner-Ballon O, et al. TET2 inactivation results in pleiotropic hematopoietic abnormalities in mouse and is a recurrent event during human lymphomagenesis. *Cancer Cell*. 2011;20:25–38.
82. Pronier E, Almire C, Mokrani H, Vasanthakumar A, Simon A, da Costa Reis Monte Mor B, et al. Inhibition of TET2-mediated conversion of 5-methylcytosine to 5-hydroxymethylcytosine disturbs erythroid and granulomonocytic differentiation of human hematopoietic progenitors. *Blood*. 2011;118:2551–5.
83. Ko M, Huang Y, Jankowska AM, Pape UJ, Tahiliani M, Bandukwala HS, et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature*. 2010;468:839–43.
84. Figueroa ME, Abdel-Wahab O, Lu C, Ward PS, Patel J, Shih A, et al. Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell*. 2010;18:553–67.
85. Couronne L, Lippert E, Andrieux J, Kosmider O, Radford-Weiss I, Penhner D, et al. Analyses of TET2 mutations in post-myeloproliferative neoplasm acute myeloid leukemias. *Leukemia*. 2010;24:201–3.
86. Couronne L, Bastard C, Bernard OA. TET2 and DNMT3A mutations in human T-cell lymphoma. *N Engl J Med*. 2012;366:95–6.
87. Soucie E, Hanssens K, Mercher T, Georgin-Lavialle S, Damaj G, Livideanu C, et al. In aggressive forms of mastocytosis, TET2 loss cooperates with c-KITD816V to transform mast cells. *Blood*. 2012;120:4846–9.
88. Mardis ER, Ding L, Dooling DJ, Larson DE, McLellan MD, Chen K, et al. Recurring mutations found by sequencing an acute myeloid leukemia genome. *N Engl J Med*. 2009;361:1058–66.
89. Cairns RA, Iqbal J, Lemonnier F, Kucuk C, de Leval L, Jais JP, et al. IDH2 mutations are frequent in angioimmunoblastic T-cell lymphoma. *Blood*. 2012;119:1901–3.
90. Palomero T, Couronne L, Khiabanian H, Kim MY, Ambesi-Impimbato A, Perez-Garcia A, et al. Recurrent mutations in epigenetic regulators, RHOA and FYN kinase in peripheral T cell lymphomas. *Nat Genet*. 2014;46:166–70.
91. Sakata-Yanagimoto M, Enami T, Yoshida K, Shiraishi Y, Ishii R, Miyake Y, et al. Somatic RHOA mutation in angioimmunoblastic T cell lymphoma. *Nat Genet*. 2014;46:171–5.
92. Losman JA, Kaelin Jr WG. What a difference a hydroxyl makes: mutant IDH, (R)-2-hydroxyglutarate, and cancer. *Genes Dev*. 2013;27:836–52.
93. McKenney AS, Levine RL. Isocitrate dehydrogenase mutations in leukemia. *J Clin Invest*. 2013;123:3672–7.
94. Lu C, Thompson CB. Metabolic regulation of epigenetics. *Cell Metab*. 2012;16:9–17.
95. Xu W, Yang H, Liu Y, Yang Y, Wang P, Kim SH, et al. Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of alpha-ketoglutarate-dependent dioxygenases. *Cancer Cell*. 2011;19:17–30.
96. Rampal R, Alkalini A, Madzo J, Vasanthakumar A, Pronier E, Patel J, et al. DNA hydroxymethylation profiling reveals that WT1 mutations result in loss of TET2 function in acute myeloid leukemia. *Cell Rep*. 2014;9:1841–55.
97. Fu X, Jin L, Wang X, Luo A, Hu J, Zheng X, et al. MicroRNA-26a targets ten eleven translocation enzymes and is regulated during pancreatic cell differentiation. *Proc Natl Acad Sci U S A*. 2013;110:17892–7.
98. Zhang P, Huang B, Xu X, Sessa WC. Ten-eleven translocation (Tet) and thymine DNA glycosylase (TDG), components of the demethylation pathway, are direct targets of miRNA-29a. *Biochem Biophys Res Commun*. 2013;437:368–73.
99. Cheng J, Guo S, Chen S, Mastriano SJ, Liu C, D'Alessio AC, et al. An extensive network of TET2-targeting MicroRNAs regulates malignant hematopoiesis. *Cell Rep*. 2013;5:471–81.
100. Song SJ, Ito K, Ala U, Kats L, Webster K, Sun SM, et al. The oncogenic microRNA miR-22 targets the TET2 tumor suppressor to promote hematopoietic stem cell self-renewal and transformation. *Cell Stem Cell*. 2013;13:87–101.
101. Lian CG, Xu Y, Ceol C, Wu F, Larson A, Dresser K, et al. Loss of 5-hydroxymethylcytosine is an epigenetic hallmark of melanoma. *Cell*. 2012;150:1135–46.
102. Yan H, Parsons DW, Jin G, McLendon R, Rasheed BA, Yuan W, et al. IDH1 and IDH2 mutations in gliomas. *N Engl J Med*. 2009;360:765–73.
103. Lu C, Ward PS, Kapoor GS, Rohle D, Turcan S, Abdel-Wahab O, et al. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature*. 2012;483:474–8.
104. Letouze E, Martinelli C, Lorient C, Burnichon N, Abermil N, Ottolenghi C, et al. SDH mutations establish a hypermethylator phenotype in paraganglioma. *Cancer Cell*. 2013;23:739–52.
105. Cancer Genome Atlas N. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012;487:330–7.
106. Kan Z, Jaiswal BS, Stinson J, Janakiraman V, Bhatt D, Stern HM, et al. Diverse somatic mutation patterns and pathway alterations in human cancers. *Nature*. 2010;466:869–73.
107. Dulak AM, Stojanov P, Peng S, Lawrence MS, Fox C, Stewart C, et al. Exome and whole-genome sequencing of esophageal adenocarcinoma identifies recurrent driver events and mutational complexity. *Nat Genet*. 2013;45:478–86.
108. Nikolaev SI, Rimoldi D, Iseli C, Valsesia A, Robyr D, Gehrig C, et al. Exome sequencing identifies recurrent somatic MAP2K1 and MAP2K2 mutations in melanoma. *Nat Genet*. 2012;44:133–9.
109. Krauthammer M, Kong Y, Ha BH, Evans P, Bacchiocchi A, McCusker JP, et al. Exome sequencing identifies recurrent somatic RAC1 mutations in melanoma. *Nat Genet*. 2012;44:1006–14.
110. Tarpey PS, Behjati S, Cooke SL, Van Loo P, Wedge DC, Pillay N, et al. Frequent mutation of the major cartilage collagen gene COL2A1 in chondrosarcoma. *Nat Genet*. 2013;45:923–6.
111. Pugh TJ, Morozova O, Attiyeh EF, Asgharzadeh S, Wei JS, Auclair D, et al. The genetic landscape of high-risk neuroblastoma. *Nat Genet*. 2013;45:279–84.

112. Cancer Genome Atlas Research N. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011;474:609–15.
113. Sato Y, Yoshizato T, Shiraiishi Y, Maekawa S, Okuno Y, Kamura T, et al. Integrated molecular analysis of clear-cell renal cell carcinoma. *Nat Genet*. 2013;45:860–7.
114. Peifer M, Fernandez-Cuesta L, Sos ML, George J, Seidel D, Kasper LH, et al. Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat Genet*. 2012;44:1104–10.
115. Stephens PJ, Tarpey PS, Davies H, Van Loo P, Greenman C, Wedge DC, et al. The landscape of cancer genes and mutational processes in breast cancer. *Nature*. 2012;486:400–4.
116. Berger MF, Hodis E, Heffernan TP, Deribe YL, Lawrence MS, Protopopov A, et al. Melanoma genome sequencing reveals frequent PREX2 mutations. *Nature*. 2012;485:502–6.
117. Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, et al. The mutational landscape of head and neck squamous cell carcinoma. *Science*. 2011;333:1157–60.
118. Fenaux P, Mufti GJ, Hellstrom-Lindberg E, Santini V, Gattermann N, Germing U, et al. Azacitidine prolongs overall survival compared with conventional care regimens in elderly patients with low bone marrow blast count acute myeloid leukemia. *J Clin Oncol*. 2010;28:562–9.
119. Fenaux P, Mufti GJ, Hellstrom-Lindberg E, Santini V, Finelli C, Giagounidis A, et al. Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: a randomised, open-label, phase III study. *Lancet Oncol*. 2009;10:223–32.
120. Janin M, Mylonas E, Saada V, Micol JB, Renneville A, Quivron C, et al. Serum 2-hydroxyglutarate production in IDH1- and IDH2-mutated de novo acute myeloid leukemia: a study by the Acute Leukemia French Association group. *J Clin Oncol*. 2014;32:297–305.
121. Wang F, Travins J, DeLaBarre B, Penard-Lacronique V, Schalm S, Hansen E, et al. Targeted inhibition of mutant IDH2 in leukemia cells induces cellular differentiation. *Science*. 2013;340:622–6.
122. Shlush LI, Zandi S, Mitchell A, Chen WC, Brandwein JM, Gupta V, et al. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature*. 2014;506:328–33.
123. Jan M, Snyder TM, Corces-Zimmerman MR, Vyas P, Weissman IL, Quake SR, et al. Clonal evolution of preleukemic hematopoietic stem cells precedes human acute myeloid leukemia. *Sci Transl Med*. 2012;4:149ra18.
124. Busque L, Patel JP, Figueroa ME, Vasanthakumar A, Provost S, Hamilou Z, et al. Recurrent somatic TET2 mutations in normal elderly individuals with clonal hematopoiesis. *Nat Genet*. 2012;44:1179–81.
125. Genovese G, Kahler AK, Handsaker RE, Lindberg J, Rose SA, Bakhoum SF, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med*. 2014;371:2477–87.
126. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371:2488–98.
127. Xie M, Lu C, Wang J, McLellan MD, Johnson KJ, Wendl MC, et al. Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med*. 2014;20:1472–8.
128. Ley TJ, Ding L, Walter MJ, McLellan MD, Lamprecht T, Larson DE, et al. DNMT3A mutations in acute myeloid leukemia. *N Engl J Med*. 2010;363:2424–33.
129. La Salle S, Trasler JM. Dynamic expression of DNMT3a and DNMT3b isoforms during male germ cell development in the mouse. *Dev Biol*. 2006;296:71–82.
130. Robertson KD, Uzvolgyi E, Liang G, Talmadge C, Sumegi J, Gonzales FA, et al. The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Res*. 1999;27:2291–8.
131. Xie S, Wang Z, Okano M, Nogami M, Li Y, He WW, et al. Cloning, expression and chromosome locations of the human DNMT3 gene family. *Gene*. 1999;236:87–95.
132. Mizuno S, Chijiwa T, Okamura T, Akashi K, Fukumaki Y, Niho Y, et al. Expression of DNA methyltransferases DNMT1, 3A, and 3B in normal hematopoiesis and in acute and chronic myelogenous leukemia. *Blood*. 2001;97:1172–9.
133. Meyers G, Ng YS, Bannock JM, Lavoie A, Walter JE, Notarangelo LD, et al. Activation-induced cytidine deaminase (AID) is required for B-cell tolerance in humans. *Proc Natl Acad Sci U S A*. 2011;108:11554–9.
134. Muramatsu M, Sankaranand VS, Anant S, Sugai M, Kinoshita K, Davidson NO, et al. Specific expression of activation-induced cytidine deaminase (AID), a novel member of the RNA-editing deaminase family in germinal center B cells. *J Biol Chem*. 1999;274:18470–6.
135. Niederreither K, Harbers M, Chambon P, Dolle P. Expression of T:G mismatch-specific thymidine-DNA glycosylase and DNA methyl transferase genes during development and tumorigenesis. *Oncogene*. 1998;17:1577–85.
136. Tolley E, Craig I. Presence of two forms of fumarase (fumarate hydratase E.C. 4.2.1.2) in mammalian cells: immunological characterization and genetic analysis in somatic cell hybrids. Confirmation of the assignment of a gene necessary for the enzyme expression to human chromosome 1. *Biochem Genet*. 1975;13:867–83.
137. Seshagiri S, Stawiski EW, Durinck S, Modrusan Z, Storm EE, Conboy CB, et al. Recurrent R-spondin fusions in colon cancer. *Nature*. 2012;488:660–4.
138. Imielinski M, Berger AH, Hammerman PS, Hernandez B, Pugh TJ, Hodis E, et al. Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell*. 2012;150:1107–20.
139. Seo JS, Ju YS, Lee WC, Shin JY, Lee JK, Bleazard T, et al. The transcriptional landscape and mutational profile of lung adenocarcinoma. *Genome Res*. 2012;22:2109–19.
140. Gui Y, Guo G, Huang Y, Hu X, Tang A, Gao S, et al. Frequent mutations of chromatin remodeling genes in transitional cell carcinoma of the bladder. *Nat Genet*. 2011;43:875–8.
141. Dolnik A, Engelmann JC, Scharfenberger-Schmeer M, Mauch J, Kelkenberg-Schade S, Haldemann B, et al. Commonly altered genomic regions in acute myeloid leukemia are enriched for somatic mutations involved in chromatin remodeling and splicing. *Blood*. 2012;120:e83–92.
142. Quesada V, Conde L, Villamor N, Ordóñez GR, Jares P, Bassaganyas L, et al. Exome sequencing identifies recurrent mutations of the splicing factor SF3B1 gene in chronic lymphocytic leukemia. *Nat Genet*. 2012;44:47–52.
143. Kalender Atak Z, De Keersmaecker K, Gianfelici V, Geerdens E, Vandepoel R, Pauwels D, et al. High accuracy mutation detection in leukemia on a selected panel of cancer genes. *PLoS One*. 2012;7:e38463.
144. Kim MS, Kim YR, Yoo NJ, Lee SH. Mutational analysis of DNMT3A gene in acute leukemias and common solid cancers. *APMIS*. 2013;121:85–94.
145. Walter MJ, Ding L, Shen D, Shao J, Grillo M, McLellan M, et al. Recurrent DNMT3A mutations in patients with myelodysplastic syndromes. *Leukemia*. 2011;25:1153–8.
146. Pugh TJ, Weeraratne SD, Archer TC, Pomeranz Krummel DA, Auclair D, Boichicchio J, et al. Medulloblastoma exome sequencing uncovers subtype-specific somatic mutations. *Nature*. 2012;488:106–10.
147. Zhang J, Ding L, Holmfeldt L, Wu G, Heatley SL, Payne-Turner D, et al. The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature*. 2012;481:157–63.
148. Agrawal N, Jiao Y, Sausen M, Leary R, Bettgeowda C, Roberts NJ, et al. Exomic sequencing of medullary thyroid cancer reveals dominant and mutually exclusive oncogenic mutations in RET and RAS. *J Clin Endocrinol Metab*. 2013;98:E364–9.
149. Shah SP, Roth A, Goya R, Oloumi A, Ha G, Zhao Y, et al. The clonal and mutational evolution spectrum of primary triple-negative breast cancers. *Nature*. 2012;486:395–9.
150. Morin RD, Mendez-Lago M, Mungall AJ, Goya R, Mungall KL, Corbett RD, et al. Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma. *Nature*. 2011;476:298–303.
151. Chen C, Bartenhagen C, Gombert M, Okpanyi V, Binder V, Rottgers S, et al. Next-generation-sequencing-based risk stratification and identification of new genes involved in structural and sequence variations in near haploid lymphoblastic leukemia. *Genes Chromosomes Cancer*. 2013;52:564–79.
152. Roberts KG, Morin RD, Zhang J, Hirst M, Zhao Y, Su X, et al. Genetic alterations activating kinase and cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*. 2012;22:153–66.
153. Gaal J, Burnichon N, Korpershoek E, Roncelin I, Bertherat J, Plouin PF, et al. Isocitrate dehydrogenase mutations are rare in pheochromocytomas and paragangliomas. *J Clin Endocrinol Metab*. 2010;95:1274–8.
154. Amary MF, Bacci K, Maggiani F, Damato S, Halai D, Berisha F, et al. IDH1 and IDH2 mutations are frequent events in central chondrosarcoma and central and periosteal chondromas but not in other mesenchymal tumours. *J Pathol*. 2011;224:334–43.
155. Hemery JP, Bastos AU, Cerutti JM. Identification of several novel non-p.R132 IDH1 variants in thyroid carcinomas. *Eur J Endocrinol*. 2010;163:747–55.

156. Murugan AK, Bojdani E, Xing M. Identification and functional characterization of isocitrate dehydrogenase 1 (IDH1) mutations in thyroid cancer. *Biochem Biophys Res Commun*. 2010;393:555–9.
157. Ghiam AF, Cairns RA, Thoms J, Dal Pra A, Ahmed O, Meng A, et al. IDH mutation status in prostate cancer. *Oncogene*. 2012;31:3826.
158. Parsons DW, Jones S, Zhang X, Lin JC, Leary RJ, Angenendt P, et al. An integrated genomic analysis of human glioblastoma multiforme. *Science*. 2008;321:1807–12.
159. Balss J, Meyer J, Mueller W, Korshunov A, Hartmann C, von Deimling A. Analysis of the IDH1 codon 132 mutation in brain tumors. *Acta Neuropathol*. 2008;116:597–602.
160. Kosmider O, Gelsi-Boyer V, Slama L, Dreyfus F, Beyne-Rauzy O, Quesnel B, et al. Mutations of IDH1 and IDH2 genes in early and accelerated phases of myelodysplastic syndromes and MDS/myeloproliferative neoplasms. *Leukemia*. 2010;24:1094–6.
161. Zhang J, Grubor V, Love CL, Banerjee A, Richards KL, Mieczkowski PA, et al. Genetic heterogeneity of diffuse large B-cell lymphoma. *Proc Natl Acad Sci U S A*. 2013;110:1398–403.
162. Kang MR, Kim MS, Oh JE, Kim YR, Song SY, Seo SI, et al. Mutational analysis of IDH1 codon 132 in glioblastomas and other common cancers. *Int J Cancer*. 2009;125:353–5.
163. Tomlinson IP, Alam NA, Rowan AJ, Barclay E, Jaeger EE, Kelsell D, et al. Germline mutations in FH predispose to dominantly inherited uterine fibroids, skin leiomyomata and papillary renal cell cancer. *Nat Genet*. 2002;30:406–10.
164. Ricketts C, Woodward ER, Killick P, Morris MR, Astuti D, Latif F, et al. Germline SDHB mutations and familial renal cell carcinoma. *J Natl Cancer Inst*. 2008;100:1260–2.
165. Bejar R, Stevenson K, Abdel-Wahab O, Galili N, Nilsson B, Garcia-Manero G, et al. Clinical effect of point mutations in myelodysplastic syndromes. *N Engl J Med*. 2011;364:2496–506.
166. Jankowska AM, Szpurka H, Tiu RV, Makishima H, Aftab M, Huh J, et al. Loss of heterozygosity 4q24 and TET2 mutations associated with myelodysplastic/myeloproliferative neoplasms. *Blood*. 2009;113:6403–10.
167. Kosmider O, Gelsi-Boyer V, Cheok M, Grabar S, Della-Valle V, Picard F, et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood*. 2009;114:3285–91.
168. Smith AE, Mohamedali AM, Kulasekararaj A, Lim Z, Gaken J, Lea NC, et al. Next-generation sequencing of the TET2 gene in 355 MDS and CMML patients reveals low-abundance mutant clones with early origins, but indicates no definite prognostic value. *Blood*. 2010;116:3923–32.
169. Tefferi A, Lim KH, Abdel-Wahab O, Lasho TL, Patel J, Patnaik MM, et al. Detection of mutant TET2 in myeloid malignancies other than myeloproliferative neoplasms: CMML, MDS, MDS/MPN and AML. *Leukemia*. 2009;23:1343–5.
170. Abdel-Wahab O, Mullally A, Hedvat C, Garcia-Manero G, Patel J, Wadleigh M, et al. Genetic characterization of TET1, TET2, and TET3 alterations in myeloid malignancies. *Blood*. 2009;114:144–7.
171. Grossmann V, Kohlmann A, Eder C, Haferlach C, Kern W, Cross NC, et al. Molecular profiling of chronic myelomonocytic leukemia reveals diverse mutations in >80% of patients with TET2 and EZH2 being of high prognostic relevance. *Leukemia*. 2011;25:877–9.
172. Itzykson R, Kosmider O, Renneville A, Gelsi-Boyer V, Meggendorfer M, Morabito M, et al. Prognostic score including gene mutations in chronic myelomonocytic leukemia. *J Clin Oncol*. 2013;31:2428–36.
173. Kohlmann A, Grossmann V, Klein HU, Schindela S, Weiss T, Kazak B, et al. Next-generation sequencing technology reveals a characteristic pattern of molecular mutations in 72.8% of chronic myelomonocytic leukemia by detecting frequent alterations in TET2, CBL, RAS, and RUNX1. *J Clin Oncol*. 2010;28:3858–65.
174. Kosmider O, Gelsi-Boyer V, Ciudad M, Racœur C, Jooste V, Vey N, et al. TET2 gene mutation is a frequent and adverse event in chronic myelomonocytic leukemia. *Haematologica*. 2009;94:1676–81.
175. Roche-Lestienne C, Marceau A, Labis E, Nibourel O, Coiteux V, Guilhot J, et al. Mutation analysis of TET2, IDH1, IDH2 and ASXL1 in chronic myeloid leukemia. *Leukemia*. 2011;25:1661–4.
176. Makishima H, Jankowska AM, McDevitt MA, O'Keefe C, Dujardin S, Cazzolli H, et al. CBL, CBLB, TET2, ASXL1, and IDH1/2 mutations and additional chromosomal aberrations constitute molecular events in chronic myelogenous leukemia. *Blood*. 2011;117:e198–206.
177. Shen Y, Zhu YM, Fan X, Shi JY, Wang QR, Yan XJ, et al. Gene mutation patterns and their prognostic impact in a cohort of 1185 patients with acute myeloid leukemia. *Blood*. 2011;118:5593–603.
178. Nibourel O, Kosmider O, Cheok M, Boissel N, Renneville A, Philippe N, et al. Incidence and prognostic value of TET2 alterations in de novo acute myeloid leukemia achieving complete remission. *Blood*. 2010;116:1132–5.
179. Metzeler KH, Maharry K, Radmacher MD, Mrozek K, Margeson D, Becker H, et al. TET2 mutations improve the new European LeukemiaNet risk classification of acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol*. 2011;29:1373–81.
180. Chou WC, Chou SC, Liu CY, Chen CY, Hou HA, Kuo YY, et al. TET2 mutation is an unfavorable prognostic factor in acute myeloid leukemia patients with intermediate-risk cytogenetics. *Blood*. 2011;118:3803–10.
181. Weissmann S, Alpermann T, Grossmann V, Kowarsch A, Nadarajah N, Eder C, et al. Landscape of TET2 mutations in acute myeloid leukemia. *Leukemia*. 2012;26:934–42.
182. Liang DC, Liu HC, Yang CP, Jaing TH, Hung JJ, Yeh TC, et al. Cooperating gene mutations in childhood acute myeloid leukemia with special reference on mutations of ASXL1, TET2, IDH1, IDH2, and DNMT3A. *Blood*. 2013;121:2988–95.
183. Langemeijer SM, Jansen JH, Hooijer J, van Hoogen P, Stevens-Linders E, Massop M, et al. TET2 mutations in childhood leukemia. *Leukemia*. 2011;25:189–92.
184. Kosmider O, Delabesse E, de Mas VM, Cornillet-Lefebvre P, Blanchet O, Delmer A, et al. TET2 mutations in secondary acute myeloid leukemias: a French retrospective study. *Haematologica*. 2011;96:1059–63.
185. Konstantin N, Bultmann S, Szewagierczak A, Dufour A, Ksienzyk B, Schneider F, et al. Genomic 5-hydroxymethylcytosine levels correlate with TET2 mutations and a distinct global gene expression pattern in secondary acute myeloid leukemia. *Leukemia*. 2011;25:1649–52.
186. Tefferi A, Levine RL, Lim KH, Abdel-Wahab O, Lasho TL, Patel J, et al. Frequent TET2 mutations in systemic mastocytosis: clinical, KITD816V and FIP1L1-PDGFRα correlates. *Leukemia*. 2009;23:900–4.
187. Alayed K, Patel KP, Konoplev S, Singh RR, Routbort MJ, Reddy N, et al. TET2 mutations, myelodysplastic features, and a distinct immunoprofile characterize blastic plasmacytoid dendritic cell neoplasm in the bone marrow. *Am J Hematol*. 2013;88:1055–61.
188. Jardin F, Ruminy P, Parmentier F, Troussard X, Vaida I, Stamatoullas A, et al. TET2 and TP53 mutations are frequently observed in blastic plasmacytoid dendritic cell neoplasm. *Br J Haematol*. 2011;153:413–6.
189. Menezes J, Acquadro F, Wiseman M, Gomez-Lopez G, Salgado RN, Talavera-Casanas JG, et al. Exome sequencing reveals novel and recurrent mutations with clinical impact in blastic plasmacytoid dendritic cell neoplasm. *Leukemia*. 2014;28:823–9.
190. Meissner B, Kridel R, Lim RS, Rogic S, Tse K, Scott DW, et al. The E3 ubiquitin ligase UBR5 is recurrently mutated in mantle cell lymphoma. *Blood*. 2013;121:3161–4.
191. Cools J. RHOA mutations in peripheral T cell lymphoma. *Nat Genet*. 2014;46:320–1.
192. Lemonnier F, Couronne L, Parrens M, Jais JP, Travert M, Lamant L, et al. Recurrent TET2 mutations in peripheral T-cell lymphomas correlate with TFH-like features and adverse clinical parameters. *Blood*. 2012;120:1466–9.
193. Jones DT, Jager N, Kool M, Zichner T, Hutter B, Sultan M, et al. Dissecting the genomic complexity underlying medulloblastoma. *Nature*. 2012;488:100–5.
194. Guichard C, Amadio G, Imbeaud S, Ladeiro Y, Pelletier L, Maad IB, et al. Integrated analysis of somatic mutations and focal copy-number changes identifies key genes and pathways in hepatocellular carcinoma. *Nat Genet*. 2012;44:694–8.
195. Kan Z, Zheng H, Liu X, Li S, Barber TD, Gong Z, et al. Whole-genome sequencing identifies recurrent mutations in hepatocellular carcinoma. *Genome Res*. 2013;23:1422–33.
196. Barber LJ, Rosa JM, Kozarewa I, Fenwick K, Assiotis I, Mitsopoulos C, et al. Comprehensive genomic analysis of a BRCA2 deficient human pancreatic cancer. *PLoS One*. 2011;6:e21639.
197. Biankin AV, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, et al. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature*. 2012;491:399–405.
198. Lindberg J, Mills IG, Klevebring D, Liu W, Neiman M, Xu J, et al. The mitochondrial and autosomal mutation landscapes of prostate cancer. *Eur Urol*. 2013;63:702–8.

199. Grasso CS, Wu YM, Robinson DR, Cao X, Dhanasekaran SM, Khan AP, et al. The mutational landscape of lethal castration-resistant prostate cancer. *Nature*. 2012;487:239–43.
200. Wang K, Kan J, Yuen ST, Shi ST, Chu KM, Law S, et al. Exome sequencing identifies frequent mutation of ARID1A in molecular subtypes of gastric cancer. *Nat Genet*. 2011;43:1219–23.
201. Zang ZJ, Cutcutache I, Poon SL, Zhang SL, McPherson JR, Tao J, et al. Exome sequencing of gastric adenocarcinoma identifies recurrent somatic mutations in cell adhesion and chromatin remodeling genes. *Nat Genet*. 2012;44:570–4.