

Q1 Tethered derivatives of D-glucose and pentacyclic triterpenes for homo/heterobivalent inhibition of glycogen phosphorylase

Keguang Cheng, Jun Liu, Hongbin Sun,* Éva Bokor, Katalin Czifrák, Bálint Kónya, Marietta Tóth, Tibor Docsa, Pál Gergely and László Somsák*

Q2 Low micromolar inhibitors (IC_{50} 40–70 μM) were found among the heterobivalent compounds studied, while homobivalent derivatives proved inactive in assays against rabbit muscle glycogen phosphorylase a or b.

Please check this proof carefully. **Our staff will not read it in detail after you have returned it.** Translation errors between word-processor files and typesetting systems can occur so the whole proof needs to be read. Please pay particular attention to: tabulated material; equations; numerical data; figures and graphics; and references. If you have not already indicated the corresponding author(s) please mark their name(s) with an asterisk. Please e-mail a list of corrections or the PDF with electronic notes attached -- do not change the text within the PDF file or send a revised manuscript.

Please bear in mind that minor layout improvements, e.g. in line breaking, table widths and graphic placement, are routinely applied to the final version.

Please note that, in the typefaces we use, an italic vee looks like this: ν , and a Greek nu looks like this: ν .

We will publish articles on the web as soon as possible after receiving your corrections; **no late corrections will be made.**

Please return your **final** corrections, where possible within **48 hours** of receipt, by e-mail to: njc@rsc.org

Reprints—Electronic (PDF) reprints will be provided free of charge to the corresponding author. Enquiries about purchasing paper reprints should be addressed via: <http://www.rsc.org/Publishing/ReSource/PaperReprints/>. Costs for reprints are below:

Reprint costs		
No of pages	Cost for 50 copies	Cost for each additional 50 copies
2-4	£190	£120
5-8	£315	£230
9-20	£630	£500
21-40	£1155	£915
>40	£1785	£1525
<i>Cost for including cover of journal issue:</i> £55 per 50 copies		

Queries are marked on your proof like this **Q1**, **Q2**, etc. and for your convenience line numbers are indicated like this 5, 10, 15, ...

Query reference	Query	Remarks
Q1	For your information: You can cite this article before you receive notification of the page numbers by using the following format: (authors), (journal), (year), DOI: 10.1039/(paper number).	
Q2	Could you please check that the contents entry text does not extend below the horizontal line when typeset? If it does, please suggest modifications that would shorten it by a sufficient amount. Many thanks.	
Q3	Should this be "benzimidazole" rather than "benzimidazol" ?	
Q4	Should this be "cynamic acid" rather than "cynnamic acid" ?	
Q5	Please check that the changes to this part of the sentence have not affected the meaning. Thank you.	
Q6	There is an isolated plus-minus symbol in the middle of Scheme 4. Can it be deleted?	
Q7	Would you like to provide captions for Charts 1 or 2, to aid the reader? In particular, what do MA, UA and OA mean in Chart 2?	
Q8	Chemical names of cpds 19--22: These names contain an unpaired square bracket,]. Is part of the name missing?	
Q9	Chemical names of cpds 31--38: These names contain an unusual numbering in the middle, namely "1H-1,2". Is part of the name missing?	
Q10	Chemical names of cpds 48--51: These names contain ",2]butane". Is this correct?	

Tethered derivatives of D-glucose and pentacyclic triterpenes for homo/heterobivalent inhibition of glycogen phosphorylase†

Keguang Cheng,^a Jun Liu,^b Hongbin Sun,^{*a} Éva Bokor,^c Katalin Czifrák,^c Bálint Kónya,^c Marietta Tóth,^c Tibor Docsa,^d Pál Gergely^e and László Somsák^{*c}

Received (in Montpellier, France) 27th October 2009, Accepted 25th January 2010

First published as an Advance Article on the web

DOI: 10.1039/b9nj00602h

Propargyl esters of the C-28 carboxylic acids of pentacyclic triterpenes (oleanolic, ursolic, and maslinic acids) were coupled with 2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyl azide as well as *N*-(ω -azido-[C-2, C-6, and C-11]alkanoyl)- β -D-glucopyranosylamines under conditions of copper(i)-catalyzed azide-alkyne cycloaddition (CuAAC) to give tethered D-glucose-triterpene heteroconjugates. The *O*-acetyl protecting groups were removed by base-catalyzed hydrolysis. *N*-(ω -Azido-[C-2, C-6, C-11, and C-16]alkanoyl)- β -D-glucopyranosylamines were also tethered by 1,7-octadiyne under CuAAC conditions to furnish D-glucose homoconjugates. *O*-Deacetylation was carried out by the Zemplén protocol. The new compounds were assayed against rabbit muscle glycogen phosphorylase (RMGP) a or b enzymes. Some of the heteroconjugates inhibited the enzyme in the low micromolar range (IC₅₀ values 40–70 μ M), while the homoconjugates proved inefficient as inhibitors.

Introduction

Type 2 diabetes mellitus has become a widespread disease afflicting a very large proportion of the population all over the world.^{1–3} The diseased state is associated with disorders in glucose metabolism by the liver and periphery resulting in elevated blood glucose levels which, in turn, are responsible for fatal long-term complications.^{1,4} An ideal anti-diabetic agent should be capable of lowering blood glucose in both fed and fasted states. Control of the hepatic glycogen metabolism is one of the key events through which insulin maintains blood glucose homeostasis. Among other means for influencing glucose production in the liver, inhibition of glycogen phosphorylase (GP), the rate-limiting enzyme of glycogen degradation, has been regarded as a promising therapeutic approach to the treatment of type 2 diabetes.^{5,6} Some GP inhibitors have shown efficacy in lowering blood glucose in animal models and clinical trials.^{7,8} In the liver and muscle isoforms of GP enzymes, six binding sites have been identified by X-ray crystallographic studies of enzyme-inhibitor complexes: the catalytic, the inhibitor, the allosteric,

the glycogen storage, and the new allosteric sites,^{6,9} as well as the recently discovered benzimidazol site.¹⁰

Among the large variety of compounds tested as GP inhibitors, the most populated class is that of D-glucose derivatives,^{11,12} which bind primarily to the catalytic site of the enzyme, as proven by several X-ray crystallographic investigations.⁹ These glucose analogue inhibitors of GP are characterized by maintaining an intact hexopyranoid sugar ring with the full OH substitution pattern of D-glucose configuration, thus resembling the non-reducing end of the natural substrate glycogen. The modifications are located at the anomeric centre as spirocycles, as well as β -NHCOR, β -NHCONHCOR, and β -C-heterocyclic substituents, just to mention the most efficient ones.^{5,6}

Pentacyclic triterpenes like **1–3** and related compounds have been reported to represent a new class of glycogen phosphorylase inhibitors.^{13–15} X-Ray crystallographic studies revealed the molecular basis of their inhibitory effect, demonstrating that pentacyclic triterpenes such as asiatic and maslinic acids bind to GP at the allosteric site.¹⁶ Oleanolic acid (**1**, OA), ursolic acid (**2**, UA) and maslinic acid (**3**, MA) have recently attracted much attention due to their broad biological activities such as protection of the liver against toxic injury, anti-inflammation, anti-HIV, antitumor, antioxidation, anti-hyperglycemia and cardiovascular activities.¹⁷

Inhibitors having the potential to bind to more than one site of an enzyme may be significantly more efficient than those with a single binding group (for some tentatively selected examples of bi- or trivalent enzyme inhibitors see ref. 18–23). This principle is well known in the interactions of multivalent carbohydrate derivatives with various proteins, and is frequently called the glycoside cluster effect in that field.²⁴ Trivalent glucose analogues have very recently been tested for GP inhibition to show a slightly better effect than

^a Center for Drug Discovery, College of Pharmacy, China Pharmaceutical University, 24 TongjiXiang, Nanjing 210009, China

^b Jiangsu Center for Drug Screening, China Pharmaceutical University, 24 TongjiXiang, Nanjing 210009, China

^c Department of Organic Chemistry, University of Debrecen, POB 20, H-4010 Debrecen, Hungary

^d Cell Biology and Signaling Research Group of the Hungarian Academy of Sciences at the Department of Medical Chemistry, Medical and Health Science Centre, University of Debrecen, Egyetem tér 1, H-4032 Debrecen, Hungary

^e Department of Medical Chemistry, Medical and Health Science Centre, University of Debrecen, Egyetem tér 1, H-4032 Debrecen, Hungary

† Electronic supplementary information (ESI) available: Copies of ¹H NMR and ¹³C NMR spectra. See DOI: 10.1039/b9nj00602h

1 that of derivatives with a single sugar unit.²⁵ Homobivalent
 Q4 indolcarboxamide²⁶ as well as cynammic acid^{27,28} derivatives
 proved very efficient inhibitors of GP.

5 With these preliminaries in mind, we envisaged conjugation
 Q5 of triterpenes and D-glucose in such a way that both could
 bind to the site to which they bind on their own, thus
 providing the first potentially heterobivalent inhibitors of
 GP. The recently reported triterpene–glucose conjugates were
 10 not capable of this because the sugar parts were attached to
 the triterpene *via* the C-6 position.²⁹ Furthermore, some new
 bivalent glucose derivatives are also reported.

Results and discussion

15 Syntheses

The new triterpene glycoconjugates were designed to include
 oleanolic, ursolic, and maslinic acids (1–3) on one hand and
 N-acyl-β-D-glucopyranosylamines on the other, by connecting
 them *via* linker chains of different length. The Cu(I)-catalyzed
 20 azide–alkyne cycloaddition³⁰ (CuAAC) was chosen as the
 linking methodology. The syntheses are summarized in
 Schemes 1–5.

Direct esterification of oleanolic acid 1, ursolic acid 2, and
 25 maslinic acid 3 with propargyl bromide (Scheme 1) afforded
 alkynes 4, 5, and 6, respectively, in excellent yields.

N-Acyl-β-D-glucopyranosylamines with a terminal azide
 group were synthesized from per-O-acetylated-β-D-glucopy-
 30 ranosyl azide³² 7 (Scheme 2). ω-Bromoalkanoyl derivatives
 8–11 were obtained by a ‘Staudinger reaction’ of 7 with PMe₃,
 resulting in an intermediate phosphinimine which, without
 being isolated, was reacted³³ with the corresponding
 ω-bromoalkanoic acid. Subsequent substitution with NaN₃
 in DMF gave compounds 12–15, respectively. Practically each


synthetic step furnished the corresponding product in very
 good yield.

To perform the CuAAC, an alkyne 4–6 and an azide 7 or
 12–14 each were dissolved in CH₂Cl₂–H₂O, followed by the
 addition of a catalytic amount of sodium L-ascorbate and
 5 CuSO₄·5H₂O. The ‘click reactions’ proceeded very well at
 room temperature to afford β-D-glucopyranosyl-1,2,3-triazoles
 16–18 (Scheme 3) and the tethered compounds 22–30
 (Scheme 4) in good to excellent yields. The O-acetyl groups
 were cleaved with 4 N NaOH/MeOH to give the corresponding
 10 deprotected compounds 19–21 (Scheme 3) and 31–37,
 respectively (Scheme 4). During deprotection of 28 the desired
 compound was not obtained; instead compound 38 could be
 isolated as a result of cleavage of the glucosylamide bond.

For bivalent glucose derivatives the N-(ω-azidoalkanoyl)-β-
 15 D-glucopyranosylamines 12–15 were reacted with 1,7-octa-
 diyne (43) under CuAAC conditions (Scheme 5). The reactions
 proceeded smoothly to give good to excellent yields of the
 coupled derivatives 44–47, which were deprotected under
 Zemplén conditions to give compounds 48–51 in similarly
 20 good yields. In order to make comparisons with monovalent
 glucose derivatives, azides 12 and 13 were deprotected by the
 Zemplén protocol to 39 and 41, respectively, which were
 further reduced to the ω-amino compounds 40 and 42.


Glycogen phosphorylase inhibition

The above-synthesized derivatives were evaluated in enzyme
 inhibition assays described previously^{35,36} against rabbit
 muscle glycogen phosphorylase a (RMGP_a) or b (RMGP_b)
 which shared considerable sequence similarity with human
 liver GP (Schemes 1 and 3–5, and Charts 1 and 2. As we found
 30 previously,³⁶ inhibitions of a and b forms of GP showed
 acceptable similarity.




	<i>IC</i> ₅₀ [μM]	R ¹	R ²	R ³	Yield [%]	<i>IC</i> ₅₀ [μM]
1 (oleanolic acid, OA)	14 ¹⁴	H	H	CH ₃	4	405 ± 30
2 (ursolic acid, UA)	9 ¹³	H	CH ₃	H	5	no inh.
3 (maslinic acid, MA)	28 ³¹	OH	H	CH ₃	6	no inh.

Scheme 1 Reagents and conditions: (a) K₂CO₃, propargyl bromide, DMF, rt. Inhibition of RMGP_a (*IC*₅₀ [μM], values are means of three experiments).




	n	Yield [%]
7	1	12 (63%)
8 not isolated	5	13 (85%)
9 (93%)	10	14 (88%)
10 (64%)	15	15 (81%)
11 (76%)		


Scheme 2 Reagents and conditions: (a) PMe₃, Br–(CH₂)_n–COOH, CH₂Cl₂, rt; (b) NaN₃, DMF, rt.



Scheme 3 Reagents and conditions: (a) CuSO₄, sodium L-ascorbate, CH₂Cl₂-H₂O, rt; (b) NaOH, MeOH, rt. Inhibition of RMGPα (*IC₅₀* [μM], values are means of three experiments).



Scheme 4 Reagents and conditions: (a) CuSO₄, sodium L-ascorbate, CH₂Cl₂-H₂O, rt; (b) NaOH, MeOH, rt. Inhibition of RMGPα (*IC₅₀* [μM], values are means of three experiments).



15 **Scheme 5** Reagents and conditions: (a) cat. NaOMe, MeOH rt.; (b) RANEY[®]-Ni, H₂, MeOH, 70 °C; (c) CuSO₄, L-ascorbic acid, CH₂Cl₂-H₂O, rt. Inhibition of RMGPb. 15

20 Inhibition by compounds tested against RMGPb

20
25
30

R	IC_{50} [μM]
1	H 14 ¹⁴
52	Et no inh. ¹⁴
53	Allyl no inh. ¹⁴
54	Bn 461 ¹⁴

55 1.14²⁹

20 The assay results showed that propargylation of the C-28 carboxyl depressed the GPa enzyme inhibitory activity (compare 1–3 to 4–6 in Scheme 1). This observation is similar to the effect of other esterifications of OA (52–54, Chart 1) resulting in a significant loss of activity.¹⁴

25 In the sugar-coupled series (Schemes 3 and 4) the activities of OA derivatives were generally better than those of the derivatives of UA and MA (19 vs. 20 and 21; 22 vs. 25 and 28; 23 vs. 26 and 29; 32 vs. 34 and 36). Deprotection of the sugar part in the 1- β -D-glucopyranosyl-1,2,3-triazole series (Scheme 3) gave better inhibitors with OA (16 vs. 19) and MA (18 vs. 21), while no significant change was observed with UA (17 vs. 20). Appending the sugar to the triazole via the C-6 position as in 55 (Chart 1) gave a very good inhibitor, although the *O*-peracetylated analogue had no activity at all.²⁹

35 Inhibition by compounds tested against RMGPb

35
40
45

Compound	K_i [μM]
56	14 ³⁷ 26 ³⁹
57	180 ³⁸

Chart 1

35 The effect of the length of the linker between the sugar and the triterpene parts was studied in the ω -triazolylalkanoyl-amide series (Scheme 4): with OA (22 vs. 23 and 24) and MA (28 vs. 29 and 30) derivatives the one-carbon linkage was significantly better than the longer ones, while among the UA compounds an opposite effect (27 vs. 25 and 26) was observed. Removal of the *O*-acetyl protecting groups in the ω -triazolylalkanoyl-amide series (Scheme 4, 31–37) brought about no obvious difference. Comparison of 19 with the hydroxy-methyl-triazole 56 shows that the presence of the OA moiety makes the inhibition somewhat better. However, 56 binds to the catalytic site,³⁷ while 19 can be expected to occupy the allosteric site.²⁹ Thus, the comparable inhibitory activities may




Chart 2 ni = no inhibition.

1 not be directly relevant, except in the as-yet unproven case of a
dual binding mode which could be expected to occur between
two enzyme dimers.²⁵ Similar considerations may apply to a
comparison of **31** and **57**.

5 In cases of bivalent glucose derivatives **48–51** (Scheme 5) no
inhibition could be observed. Study of analogous monovalent
compounds revealed that with an azide as endgroup (**39**, **41**)
the inhibitory activity was moderate and decreased with the
length of the linker. Bivalent compound **48** can also be
10 compared with the monovalent triazole **57**³⁸ (Chart 1) to
show that the dimeric structures seem to be too large to
occupy the catalytic site, and no other interactions exist with
the enzyme. In the presence of amine endgroups (**40**, **42**) the
inhibition was much weaker, and with the longer linker chain
15 no effect was detected.

Conclusions

Copper(i)-catalyzed azide–alkyne cycloaddition – ‘click
chemistry’ – proved suitable for the synthesis of conjugates
20 of pentacyclic triterpenes and D-glucose derivatives as new,
potentially heterobivalent inhibitors of glycogen phosphorylase.
Compounds **17** (IC₅₀ = 51 μM), **19** (IC₅₀ = 26 μM), **20**
(IC₅₀ = 45 μM), **22** (IC₅₀ = 68 μM), **27** (IC₅₀ = 65 μM) and
25 (IC₅₀ = 78 μM) were the most potent inhibitors of
RMGPa. Homobivalent glucose derivatives proved inefficient
in RMGPb inhibition assays. The monovalent analogues of
both triterpenes and glucose derivatives proved generally more
efficient than the bivalent compounds.

Experimental

General methods

35 All commercially available solvents and reagents were used
without further purification. Melting points were measured on
a RY-1 or on a Kofler hot-stage melting point apparatus.
Column chromatography was carried out on E. Merck Silica
Gel 60 (230–400 mesh), on silica gel (200–300 mesh, Qindao
Ocean Chemical Company, China), or Kieselgel 60 (Merck,
40 particle size 0.063–0.200 mm). IR spectra were recorded on
Shimadzu FTIR-8400S spectrometer. ¹H- and ¹³C-NMR spectra
were measured on Bruker AV-300 (300/75 MHz for ¹H/¹³C),
Bruker 360 (360/90 MHz for ¹H/¹³C) or Avance DRX 500
(500/125 MHz for ¹H/¹³C) spectrometers. Chemical shifts are
45 reported as values from an internal tetramethylsilane standard.
TLC was performed on DC-Alurolle Kieselgel 60 F254 (Merck),
and the plates were visualised under UV light and by gentle
heating. Mass spectral data were obtained on Agilent 1100
LC/DAD/MSD or Q-ToF Micro MS/MS spectrometers. Optical
50 rotations were measured using a Perkin-Elmer 141 or a Perkin-
Elmer 241 polarimeters at rt. PMe₃ (1 M solution in toluene) and
1,7-octadiyne were purchased from Sigma-Aldrich.

Syntheses

55 General procedure I for the propargylation of oleanolic acid, ursolic acid or maslinic acid

To a solution of a carboxylic acid (**1** or **2** or **3**, 2.2 mmol) in
DMF (5 mL), was added propargyl bromide (2.4 mmol) and

K₂CO₃ (4.4 mmol). The reaction mixture was stirred at rt
1 for 18 h, then concentrated. The residue was diluted with
EtOAc (50 mL), washed successively with 1 N HCl, water,
satd. aq. NaHCO₃, water and brine, dried (MgSO₄), filtered
and concentrated. The residue was purified by column
5 chromatography.

Propargyl 3β-hydroxyolean-12-en-28-oate (4)²⁹. Prepared
from **1** (1 g, 2.2 mmol) and propargyl bromide (0.27 mL,
2.4 mmol) according to General procedure I. The residue was
10 purified by column chromatography (EtOAc–hexane, 1 : 6).
Yield: 1.05 g, 97%, white solid, mp 121–122 °C; R_f = 0.33
(EtOAc–hexane, 1 : 4); [α]_D = +67.9 (c = 0.50, CH₂Cl₂). IR
(KBr, cm⁻¹): 3308, 2945, 2866, 1731, 1157, 1032, 739; ¹H
NMR (300 MHz, CDCl₃): δ 0.74, 0.77, 0.92, 0.98, 1.13 (5 s,
15 each 3H, 5 × CH₃), 0.90 (s, 6H, 2 × CH₃), 0.71–2.04 (m, 22H),
2.41 (t, 1H, J = 2.6 Hz, CH), 2.87 (dd, 1H, J = 4.1, 9.5 Hz,
H-18), 3.21 (dd, 1H, J = 5.1, 10.7 Hz, H-3), 4.56 (dd, 1H, J =
2.6, 15.4 Hz, CO₂CH₂), 4.68 (dd, 1H, J = 2.6, 15.4 Hz,
CO₂CH₂), 5.30 (t, 1H, J = 3.5 Hz, H-12); ¹³C NMR
20 (75 MHz, CDCl₃): δ 15.3, 15.6, 17.1, 18.3, 23.0, 23.4, 23.6,
25.8, 27.2, 27.7, 28.1, 30.7, 32.2, 32.8, 33.1, 33.8, 37.0, 38.5,
38.8, 39.4, 41.3, 41.7, 45.9, 46.8, 47.6, 51.6, 55.2, 74.4, 78.1,
79.0, 122.63, 143.4, 176.8. ESI-MS (positive mode) m/z: 517.3
[M + Na]⁺.

Propargyl 3β-hydroxyurs-12-en-28-oate (5). Prepared from **2**
(2.0 g, 4.4 mmol) and propargyl bromide (0.54 mL, 4.8 mmol)
according to General procedure I. The residue was purified by
column chromatography (EtOAc–hexane, 1 : 5). Yield: 2.0 g,
93%, white solid, mp 129–131 °C; R_f = 0.48 (EtOAc–hexane,
30 1 : 5). IR (KBr, cm⁻¹): 3309, 2927, 2871, 1729, 1454, 1383,
1221, 1167, 1139, 1106, 1032, 996, 757, 667; ¹H NMR
(300 MHz, CDCl₃): δ 0.76, 0.77, 0.91, 0.95, 0.98, 1.08
(6 s, each 3H, 6 × CH₃), 0.87 (d, 3H, J = 6.4 Hz, CH₃),
35 0.75–2.10 (m, 22H), 2.26 (d, 1H, J = 11.3 Hz, H-18), 2.41
(t, 1H, J = 2.4 Hz, CH), 3.20 (dd, 1H, J = 5.1, 10.7 Hz, H-3),
4.57 and 4.65 (dd, each 1H, J = 2.5, 15.6 Hz, COOCH₂), 5.27
(1H, t, J = 3.6 Hz, H-12); ¹³C NMR (75 MHz, CDCl₃): δ 5.5,
15.6, 17.0, 17.2, 18.3, 21.1, 23.3, 23.5, 24.6, 27.3, 28.0, 28.1,
40 30.6, 33.1, 36.4, 37.0, 38.67, 38.75, 38.8, 39.1, 39.6, 42.1, 47.6,
48.2, 51.6, 52.8, 55.3, 74.3, 78.1, 79.0, 125.9, 137.8, 176.6.
ESI-MS (positive mode) m/z: 495.4 [M + H]⁺.

Propargyl 2α,3β-dihydroxyolean-12-en-28-oate (6). Prepared
from **3** (1.4 g, 3.0 mmol) and propargyl bromide (0.37 mL,
3.3 mmol) according to General procedure I. The residue was
purified by column chromatography (EtOAc–hexane, 1 : 5).
Yield: 1.3 g, 87%, white solid, mp 233–234 °C; R_f = 0.69
(EtOAc–hexane, 1 : 5). IR (KBr, cm⁻¹): 3394, 3309, 2946,
1729, 1463, 1388, 1364, 1259, 1217, 1157, 1121, 1049, 1033,
995, 758, 669, 633; ¹H NMR (300 MHz, CDCl₃): δ 0.74, 0.82,
50 0.90, 0.92, 0.98, 1.03, 1.13 (7 s, each 3H, 7 × CH₃), 0.75–2.01
(m, 20H), 2.41 (t, 1H, J = 2.4 Hz, CH), 2.87 (dd, 1H, J = 4.3,
13.7 Hz, H-18), 3.01 (d, 1H, J = 9.5 Hz, H-3), 3.65–3.73
(m, 1H, H-2), 4.57 and 4.69 (dd, each 1H, J = 2.4, 15.6 Hz,
55 COOCH₂), 5.31 (t, 1H, J = 3.5 Hz, H-12); ¹³C NMR
(75 MHz, CDCl₃): δ 16.6, 16.7, 17.2, 18.4, 23.0, 23.5, 23.6,
25.9, 27.7, 28.6, 30.7, 32.2, 32.7, 33.1, 33.9, 38.3, 39.2,
39.5, 41.3, 41.8, 45.9, 46.5, 46.8, 47.6, 51.6, 55.4, 69.0, 74.4,

1 78.1, 84.0, 122.5, 143.5, 176.8. ESI-MS (positive mode) m/z :
533.4 [M + Na]⁺.

5 **General procedure II for the preparation of *N*-(ω -bromoalkanyl)-
2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamines (9–11)**

2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyl azide (**7**, 0.10 g,
0.27 mmol) was dissolved in dry CH₂Cl₂ (3 mL). To the
solution Me₃P (1.1 equiv. of a 1 M solution in toluene) was
added in one portion. The mixture was stirred at rt. until
nitrogen evolution had ceased and TLC (EtOAc–hexane, 1 : 1)
had indicated complete transformation of the azide. This
solution was then reacted with an ω -bromoalkanoic acid
(1.1 equiv., as indicated with the particular compounds) till
the disappearance of the iminophosphorane (TLC, EtOAc–
hexane, 1 : 1). Then, it was diluted with CH₂Cl₂ (5 mL) and
washed with satd. aq. NaHCO₃ solution (2 × 5 mL). The
organic phase was dried over MgSO₄ and the solvent was
removed under diminished pressure. The crude product was
purified by column chromatography.

***N*-(6-Bromohexanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyrano-
sylamine (**9**).** Prepared from **7** (0.50 g 1.34 mmol) according to
General procedure II. The residue was purified by column
chromatography (EtOAc–hexane, 1 : 1). Yield: 0.65 g, 93%,
colourless oil, R_f = 0.35 (EtOAc–hexane, 1 : 1); $[\alpha]_D^{25}$ = +22
(c = 0.59, CHCl₃); ¹H NMR (360 MHz, CDCl₃): δ (ppm)
1.42–1.47 (m, 2H, CH₂), 1.60–1.63 (m, 2H, CH₂), 1.83–1.90
(m, 2H, CH₂), 2.02, 2.04, 2.06, 2.08 (4s, 12H, 4 × OCOCH₃),
2.21–2.27 (m, 2H, CH₂), 3.39–3.43 (m, 2H, CH₂), 3.86 (ddd,
1H, J = 1.2, 2.6, 10.6 Hz, H-5), 4.08 (dd, 1H, J = 1.2,
11.9 Hz, H-6b), 4.32 (dd, 1H, J = 2.6, 11.9 Hz, H-6a), 4.93,
5.06, 5.29, 5.32 (4 pseudo t, 4H, J = 9.2, 10.6 Hz in each, H-1,
H-2, H-3, H-4), 6.63 (d, 1H, J = 9.2 Hz, NH); ¹³C NMR
(90 MHz, CDCl₃): δ (ppm) 20.3 (3), 20.5 (4 × OCOCH₃), 23.9,
27.3, 32.0, 33.3, 35.9 (5 × CH₂), 61.5 (C-6), 67.9, 70.4, 72.5,
73.2 (C-2, C-3, C-4, C-5), 77.7 (C-1), 169.3, 169.6, 170.4,
170.5 (4 × OCOCH₃), 172.8 (NHCO). Anal. calcd. for
C₂₀H₃₀BrNO₁₀ (524.37): C 45.81, H 5.77, N 2.67. Found: C
45.64, H 5.94, N 2.59.

***N*-(11-Bromoundecanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopy-
ranosylamine (**10**).** Prepared from **7** (5.0 g 13.4 mmol)
according to General procedure II. The residue was purified
by column chromatography (EtOAc–hexane, 1 : 1). Yield:
5.13 g, 64%, white crystalline product, mp 61–63 °C; $[\alpha]_D^{25}$ =
+15 (c = 0.38, CHCl₃); ¹H NMR (360 MHz, CDCl₃): δ (ppm)
1.28 (bs, 10H, 5 × CH₂), 1.40–1.45 (m, 2H, CH₂), 1.57–1.50
(m, 2H, CH₂), 1.80–1.86 (m, 2H, CH₂), 2.02, 2.04, 2.06, 2.08
(4 s, 12H, 4 × OCOCH₃), 2.17–2.21 (m, 2H, CH₂), 3.38–3.43
(m, 2H, CH₂), 3.85 (ddd, 1H, J = 1.2, 2.6, 10.6 Hz, H-5), 4.07
(dd, 1H, J = 1.2, 11.9 Hz, H-6b), 4.32 (dd, 1H, J = 2.6, 11.9
Hz, H-6a), 4.93, 5.06, 5.27, 5.32 (4 pseudo t, 4H, J = 9.2, 10.6
Hz in each, H-1, H-2, H-3, H-4), 6.51 (d, 1H, J = 9.2 Hz,
NH); ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.4(3), 20.5
(4 × OCOCH₃), 24.8, 25.0, 27.9, 28.4, 28.8, 29.0, 29.1, 32.5,
33.7, 36.3 (10 × CH₂), 61.5 (C-6), 67.9, 70.4, 72.5, 73.3
(C-2, C-3, C-4, C-5), 77.8 (C-1), 169.3, 169.6, 170.4, 170.6
(4 × OCOCH₃), 173.3 (NHCO). Anal. calcd. for C₂₅H₄₀BrNO₁₀

(594.50): C 50.51, H 6.78, N 2.36. Found: C 50.64, H 6.91,
N 2.49.

***N*-(16-Bromohexadecanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopy-
ranosylamine (**11**).** Prepared from **7** (0.50 g 1.34 mmol)
according to General procedure II. The residue was purified
by column chromatography (EtOAc–hexane, 1 : 1). Yield:
0.68 g, 76%, white crystalline product, mp 91–93 °C; $[\alpha]_D^{25}$ =
+10 (c = 0.20, CHCl₃); ¹H NMR (360 MHz, CDCl₃): δ (ppm)
1.28 (m, 18H, 9 × CH₂), 1.36–1.39 (m, 2H, CH₂), 1.48–1.52
(m, 2H, CH₂), 1.57–1.61 (m, 2H, CH₂), 1.84–1.88 (m, 2H,
CH₂), 2.03, 2.04, 2.06, 2.08 (4 s, 12H, 4 × OCOCH₃),
2.27–2.31 (m, 2H, CH₂), 3.36–3.40 (m, 2H, CH₂), 3.87
(ddd, 1H, J = 1.2, 2.6, 10.6 Hz, H-5), 4.06 (dd, 1H, J =
1.2, 11.9 Hz, H-6b), 4.35 (dd, 1H, J = 2.6, 11.9 Hz, H-6a),
4.98, 5.22, 5.27, 5.32 (4 pseudo t, 4H, J = 9.2, 10.6 Hz in each,
H-1, H-2, H-3, H-4), 6.50 (d, 1H, J = 9.2 Hz, NH); ¹³C NMR
(90 MHz, CDCl₃): δ (ppm) 20.2 (3), 20.3 (4 × OCOCH₃), 23.0,
23.4, 24.8, 25.0, 26.0, 26.3, 28.1, 28.8, 29.0, 29.3, 29.4, 32.0,
33.8, 34.5, 36.2, (16 × CH₂), 61.7 (C-6), 68.1, 71.2, 72.5, 74.3
(C-2, C-3, C-4, C-5), 76.8 (C-1), 169.3, 169.6, 170.4, 170.6
(4 × OCOCH₃), 172.3 (NHCO). Anal. calcd. for
C₃₀H₅₀BrNO₁₀ (664.64): C 54.22, H 7.58, N 2.11. Found: C
54.11, H 7.72, N 2.29.

**General procedure III for the preparation of *N*-(ω -azidoalkanyl)-
2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamines (12–15)**

An *N*-(ω -bromoalkanyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyra-
nosylamine (**9–11**) was dissolved in dry DMSO (15 mL/mmol).
To the solution NaN₃ (2 equiv.) was added in one portion.
The mixture was stirred at rt until the disappearance of the starting
bromide (TLC EtOAc–hexane 1 : 1). The solution was diluted
with water (150 mL), washed with Et₂O (5 × 25 mL) and water
(1 × 25 mL), dried over MgSO₄ and the solvent was removed
under diminished pressure. The crude product was purified by
column chromatography or crystallisation.

***N*-Azidoacetyl-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamine
(**12**).** To a solution of azide **7** (0.20 g, 0.54 mmol) in dry
CH₂Cl₂ (3 mL), PMe₃ (0.54 mL of a 1 M solution in toluene)
was added in one portion. The mixture was stirred at rt until
nitrogen evolution had ceased, and TLC (EtOAc–hexane, 1 : 1)
had indicated complete transformation of **7** (approx. 15 min).
This soln was then reacted with bromoacetic acid (0.082 g,
0.59 mmol). When TLC (EtOAc–hexane, 1 : 1) showed no
more change (conversions were incomplete), the solvent was
evaporated, and the pale yellow oil was dissolved in CH₂Cl₂
(30 mL) and extracted with satd. aq. NaHCO₃ solution
(2 × 30 mL). The organic phase was dried over MgSO₄,
concentrated under diminished pressure, then the residue
(0.20 g) was dissolved in dry DMF (3 mL) and NaN₃ (0.056 g,
0.82 mmol) was added. The reaction mixture was stirred at rt
for 2 h (TLC, EtOAc–hexane, 1 : 1). The mixture was then
diluted with water (20 mL) and extracted with Et₂O (5 × 30 mL).
The combined organic phase was dried over MgSO₄ and the
solvent was removed under diminished pressure. The obtained
syrup was purified by column chromatography (EtOAc–
hexane, 4 : 6) to give **12** as white crystals: 0.15 g, 63%, calcd
for **7**; mp 150–152 °C, $[\alpha]_D^{25}$ = +39 (c = 0.21, CHCl₃),

1 (lit.⁴⁰ $[\alpha]_{\text{D}} +4.1$ ($c = 1$, CHCl_3); ^1H NMR (360 MHz, CDCl_3): δ (ppm) 2.03, 2.04, 2.07, 2.09 (4s, 12H, $4 \times \text{OCOCH}_3$), 3.85 (ddd, 1H, $J = 2.6, 4.0, 9.2$ Hz, H-5), 3.93–4.11 (m, 3H, H-6b, CH_2), 4.30 (dd, 1H $J = 2.6, 13.2$ Hz, H-6a), 4.99, 5.08, 5.24, 5.33 (4 pseudo t, 4H, $J = 9.2, 10.6$ Hz in each, H-1, H-2, H-3, H-4), 7.19 (d, 1H, $J = 9.2$ Hz, NH); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 20.6, 20.5 ($4 \times \text{OCOCH}_3$), 52.4 (CH_2), 61.5 (C-6) 67.9, 70.3, 72.5, 73.6 (C-2, C-3, C-4, C-5), 78.0 (C-1), 167.5, 169.5, 169.8, 170.5 ($4 \times \text{OCOCH}_3$), 170.8 (NHCO). Anal. calcd. for $\text{C}_{16}\text{H}_{22}\text{N}_4\text{O}_{10}$ (430.37): C, 44.65; H, 5.15; N, 13.02; Found: C, 44.53; H, 5.22; N, 13.12.

***N*-(6-Azidohexanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamine (13).** Prepared from **9** (2.00 g, 3.81 mmol) according to General procedure III. Yield: 1.57 g, 85%, white crystalline product, mp 135–137 °C; $[\alpha]_{\text{D}} = +37$ ($c = 0.34$, CHCl_3); ^1H NMR (360 MHz, CDCl_3): δ (ppm) 1.36–1.44 (m, 2H, CH_2), 1.57–1.68 (m, 4H, $2 \times \text{CH}_2$), 2.02, 2.04, 2.05, 2.08 (4 s, 12H, $4 \times \text{OCOCH}_3$), 2.18–2.24 (m, 2H, CH_2), 3.24–3.29 (m, 2H, CH_2), 3.84 (ddd, 1H, $J = 1.2, 2.6, 10.6$ Hz, H-5), 4.08 (dd, 1H, $J = 1.1, 11.9$ Hz, H-6b), 4.31 (dd, 1H, $J = 11.9, 4.0$ Hz, H-6a), 4.92, 5.06, 5.28, 5.31 (4 pseudo t, 4H, $J = 9.2, 10.6$ Hz in each, H-1, H-2, H-3, H-4), 6.57 (d, 1H, $J = 9.2$ Hz, NH); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 20.4 (3), 20.5 ($4 \times \text{OCOCH}_3$), 24.4, 26.0, 28.4, 36.1, 51.0 ($5 \times \text{CH}_2$), 61.5 (C-6), 68.0, 70.4, 73.4, 72.6 (C-2, C-3, C-4, C-5), 77.9 (C-1), 169.5, 169.7, 170.5, 170.8 ($4 \times \text{OCOCH}_3$), 173.0 (NHCO). Anal. calcd. for $\text{C}_{20}\text{H}_{30}\text{N}_4\text{O}_{10}$ (486.48): C 49.38, H 6.22, N 11.52. Found: C 49.46, H 6.14, N 11.59.

***N*-(11-Azidoundecanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamine (14).** Prepared from **10** (1.00 g, 1.69 mmol) according to General procedure III. Yield: 0.83 g, 88%, white crystalline product, mp 68–70 °C; $[\alpha]_{\text{D}} = +13$ ($c = 0.22$, CHCl_3); ^1H NMR (360 MHz, CDCl_3): δ (ppm) 1.25–1.31 (m, 14H, $7 \times \text{CH}_2$), 1.46–1.50 (m, 2H, CH_2), 2.02, 2.04, 2.05, 2.08 (4 s, 12H, $4 \times \text{OCOCH}_3$), 2.20–2.25 (m, 2H, CH_2), 2.52–2.56 (m, 2H, CH_2), 3.17 (ddd, 1H, $J = 1.2, 2.6, 10.6$ Hz, H-5), 3.31 (dd, 1H, $J = 1.1, 11.9$ Hz, H-6b), 3.53 (dd, 1H, $J = 4.0, 11.9$ Hz, H-6a), 4.22, 4.29, 4.56, 4.61 (4 pseudo t, 4H, $J = 9.2, 10.6$ Hz in each, H-1, H-2, H-3, H-4), 7.49 (d, 1H, $J = 9.2$ Hz, NH); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 20.4 (3), 20.5 ($4 \times \text{OCOCH}_3$), 25.0, 26.4, 27.9, 28.0, 28.2, 28.3, 30.1, 32.4, 35.0, 50.2 ($10 \times \text{CH}_2$), 60.9 (C-6), 67.1, 69.6, 72.2, 72.4 (C-2, C-3, C-4, C-5), 77.9 (C-1), 168.5, 168.6, 168.7, 169.3 ($4 \times \text{OCOCH}_3$), 173.0 (NHCO). Anal. calcd. for $\text{C}_{25}\text{H}_{40}\text{N}_4\text{O}_{10}$ (556.62): C 53.95, H 7.24, N 10.07. Found: C 53.76, H 7.04, N 10.19.

***N*-(16-Azidohexadecanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamine (15).** Prepared from **11** (0.4 g, 0.60 mmol) according to General procedure III. Yield: 0.30 g, 81%, white crystalline product, mp 92–94 °C; $[\alpha]_{\text{D}} = +16$ ($c = 0.22$, CHCl_3); ^1H NMR (360 MHz, CDCl_3): δ (ppm) 1.24–1.28 (m, 18H, $9 \times \text{CH}_2$), 1.36–1.39 (m, 2H, CH_2), 1.47–1.50 (m, 2H, CH_2), 1.56–1.61 (m, 2H, CH_2), 1.84–1.87 (m, 2H, CH_2), 2.02, 2.04, 2.07, 2.08 (4s, 12H, $4 \times \text{OCOCH}_3$), 2.27–2.31 (m, 2H, CH_2), 3.38–3.41 (m, 2H, CH_2), 3.21 (ddd, 1H, $J = 1.2, 2.6, 10.6$ Hz, H-5), 3.29 (1H, dd, $J = 1.1, 11.9$ Hz, H-6b),

3.52 (dd, 1H, $J = 4.0, 11.9$ Hz, H-6a), 4.20, 4.25, 4.71, 4.60 (4 pseudo t, 4H, $J = 9.2, 10.6$ Hz, in each, H-1, H-2, H-3, H-4), 7.48 (d, 1H, $J = 9.2$ Hz, NH); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 20.4 (3), 20.5 ($4 \times \text{OCOCH}_3$), 23.3, 23.4, 3 24.8, 26.0, 27.1, 27.6, 28.2, 28.5, 29.0, 29.7, 30.4, 31.9, 33.8, 36.5, 49.2 ($15 \times \text{CH}_2$), 61.5 (C-6), 68.3, 70.0, 72.1, 72.5 (C-2, C-3, C-4, C-5), 78.7 (C-1), 168.5, 168.6, 168.7, 169.3 ($4 \times \text{OCOCH}_3$), 172.1 (NHCO). Anal. calcd. for $\text{C}_{30}\text{H}_{50}\text{N}_4\text{O}_{10}$ (626.75): C 57.49, H 8.04, N 8.94. Found: C 57.23, H 8.24, N 8.82.

General procedures IV for the CuAAC ‘click’ reaction

(a). To a solution of an alkyne (0.27 mmol) and an azide (0.27 mmol) in CH_2Cl_2 (2 mL) and H_2O (2 mL), was added $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (0.32 mmol) and Na-L-ascorbate (0.64 mmol). The resulting solution was stirred for 12 h at rt. The reaction mixture was diluted with H_2O (10 mL), then extracted with CH_2Cl_2 (3×10 mL). The combined organic layer was dried over MgSO_4 , filtered, and concentrated. The residue was purified by column chromatography.

(b). An *N*-(ω -azidoalkanoyl)-2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylamine (0.46 mmol) was added to a 1 : 1 (v/v) mixture of CH_2Cl_2 and water (10 mL/mmol). To the solution 1,7-octadiyne (**43**, 1.0 equiv.), $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (5 mol%) and L-ascorbic acid (15 mol%) were added. The mixture was heated at 50 °C until complete transformation of the starting azide (TLC, EtOAc). The solution was diluted with water (25 mL) and extracted with CH_2Cl_2 (5×12 mL). The organic phase was dried over MgSO_4 and the solvent was removed under diminished pressure. The crude product was purified by column chromatography or crystallisation.

General procedure V for *O*-deacetylation

To a solution or suspension of an *O*-peracetylated compound (0.081 mmol) in MeOH (3 mL) was added 4 N aq. NaOH (0.4 mL), then stirred at rt for 1 h, and neutralized with 1 N HCl (1.8 mL). The mixture was concentrated *in vacuo* and the residue was taken up in EtOAc (50 mL), washed successively with 1 N HCl (3×15 mL), water (3×15 mL) and brine (3×15 mL), dried (Na_2SO_4), filtered and concentrated. The residue was purified by column chromatography.

[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyolean-12-en-28-oate (16). Prepared from **4** (0.13 g, 0.27 mmol) and **7** (0.10 g, 0.27 mmol) according to General procedure IVa. The residue was purified by column chromatography (EtOAc–hexane, 1 : 1). Yield: 0.13 g, 57%, white solid, mp 124–125 °C. $R_f = 0.13$ (EtOAc–hexane, 1 : 2). IR (KBr, cm^{-1}): 2947, 2869, 1757, 1460, 1369, 1227, 1036, 758; ^1H NMR (300 MHz, CDCl_3): δ 0.66, 0.78, 0.89, 0.90, 0.91, 0.99, 1.13 (7 s, each 3H, $7 \times \text{CH}_3$), 0.67–2.09 (m, 22H), 1.86, 2.03, 2.07, 2.09 (4 s, each 3H, $4 \times \text{OCOCH}_3$), 2.86 (dd, 1H, $J = 3.8, 13.9$ Hz, H-18), 3.18–3.23 (m, 1H, H-3), 3.96–4.02 (m, 1H, H-5-Glc), 4.11–4.16 (m, 1H, H-6a-Glc), 4.33 (dd, 1H, $J = 4.8, 12.6$ Hz, H-6b-Glc), 5.17 (s, 2H, COOCH_2), 5.21–5.27 (m, 1H, H-2-Glc), 5.31 (t, 1H, $J = 3.3$ Hz, H-12), 5.39–5.43 (m, 2H, overlapping, H-3-Glc and H-4-Glc), 5.84–5.87 (m, 1H, H-1-Glc), 7.81 (s, 1H, NCH); ^{13}C NMR (75 MHz, CDCl_3):

1 δ 15.3, 15.6, 16.9, 18.4, 20.1, 20.5, 20.6, 23.0, 23.4, 23.6, 25.9,
27.2, 27.7, 28.1, 29.7, 30.7, 32.2, 32.7, 33.1, 33.9, 37.1, 38.5,
38.8, 39.3, 41.4, 41.8, 45.9, 46.7, 47.6, 55.3, 57.4, 61.5, 67.7,
70.3, 72.6, 75.3, 79.0, 85.9, 122.0, 122.5, 143.7, 144.0, 168.6,
5 169.3, 169.9, 170.4, 177.4. ESI-MS (positive mode) m/z : 890.8
[M + Na]⁺.

10 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyl)-1*H*-1,2,3-triazol-
4-ylmethyl] 3 β -hydroxyurs-12-en-28-oate (17).** Prepared from **5**
(0.13 g, 0.27 mmol) and **7** (0.10 g, 0.27 mmol) according to
General procedure IVa. The residue was purified by column
chromatography (EtOAc–hexane, 1:2). Yield: 0.22 g, 96%,
white solid, mp 120–122 °C. R_f = 0.20 (EtOAc–hexane, 1:2);
IR (KBr, cm⁻¹): 2932, 2872, 1757, 1456, 1376, 1228, 1104; ¹H
15 NMR (300 MHz, CDCl₃): δ 0.72, 0.81, 0.94, 0.97, 1.01, 1.11
(6 s, each 3H, 6 \times CH₃), 0.89 (d, 3H, J = 6.3 Hz, CH₃),
0.72–2.13 (m, 22H), 1.89, 2.06, 2.10, 2.13 (4 s, each 3H, 4 \times
OCOCH₃), 2.25 (d, 1H, J = 10.8 Hz, H-18), 3.25 (dd, 1H, J =
4.3, 10.7 Hz, H-3), 4.00–4.04 (m, 1H, H-5-Glc), 4.14–4.19
20 (m, 1H, H-6a-Glc), 4.35 (dd, 1H, J = 4.5, 12.7 Hz, H-6b-Glc),
5.17 and 5.18 (2 d, each 1H, J = 12.8 Hz, COOCH₂),
5.23–5.30 (m, 2H, overlapping, H-12 and H-2-Glc),
5.40–5.48 (m, 2H, overlapping, H-3-Glc and H-4-Glc), 5.90
25 (d, 1H, J = 9.2 Hz, H-1-Glc), 7.82 (s, 1H, NCH); ¹³C NMR
(75 MHz, CDCl₃): δ 15.5, 15.6, 19.96, 17.0, 18.4, 20.1, 20.5,
20.6, 21.1, 23.3, 23.6, 24.2, 27.3, 28.1, 28.2, 30.6, 33.0, 36.4,
37.0, 38.7, 38.8, 38.82, 39.1, 39.6, 42.1, 47.6, 48.1, 52.9, 55.3,
57.3, 61.5, 67.8, 70.4, 72.7, 75.3, 79.1, 86.9, 122.0, 125.8, 138.1,
144.1, 168.6, 169.3, 169.9, 177.3. ESI-MS (positive mode) m/z :
30 890.8 [M + Na]⁺.

35 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyl)-1*H*-1,2,3-triazol-
4-ylmethyl] 2 α ,3 β -dihydroxyolean-12-en-28-oate (18).** Prepared
from **6** (0.14 g, 0.27 mmol) and **7** (0.10 g, 0.27 mmol) according
to General procedure IVa. The residue was purified by column
chromatography (EtOAc–hexane, 1:1). Yield: 0.20 g, 84%,
white solid, mp 158–160 °C, R_f = 0.17 (EtOAc–hexane, 1:1);
IR (KBr, cm⁻¹): 3395, 2948, 1758, 1460, 1368, 1227, 1101,
1037, 758; ¹H NMR (300 MHz, CDCl₃): δ 0.69, 0.85, 0.92,
0.94, 1.00, 1.06, 1.15 (7 s, each 3H, 7 \times CH₃), 0.69–2.12
40 (m, 20H), 1.88, 2.06, 2.10, 2.12 (4 s, each 3H, 4 \times OCOCH₃),
2.89 (dd, 1H, J = 3.1, 13.2 Hz, H-18), 3.05 (d, 1H, J = 9.5 Hz,
H-3 α), 3.68–3.76 (m, 1H, H-2 β), 4.00–4.04 (m, 1H, H-5-Glc),
4.14–4.18 (m, 1H, H-6a-Glc), 4.35 (dd, 1H, J = 4.8, 12.6 Hz,
H-6b-Glc), 5.19 and 5.20 (2 d, each 1H, J = 12.9 Hz,
45 COOCH₂), 5.24–5.30 (m, 1H, H-2-Glc), 5.34 (brs, 1H,
H-12), 5.43–5.46 (m, 2H, H-3-Glc and H-4-Glc), 5.90
(d, 1H, J = 9.2 Hz, H-1-Glc), 7.84 (s, 1H, NCH); ¹³C NMR
(75 MHz, CDCl₃): δ 16.9, 17.0, 17.2, 18.6, 20.3, 20.7, 20.9,
50 23.3, 23.7, 23.9, 26.1, 27.9, 28.8, 30.9, 32.4, 32.9, 33.3, 34.1,
38.6, 39.3, 39.4, 39.7, 41.6, 42.0, 46.1, 46.7, 47.0, 47.9, 55.6,
57.6, 61.8, 68.0, 69.2, 70.6, 72.9, 75.5, 77.4, 84.2, 86.1, 122.2,
122.6, 144.0, 144.3, 168.9, 169.5, 170.1, 170.6, 177.7. ESI-MS
(positive mode) m/z : 906.4 [M + Na]⁺.

Q8 **1-(β -D-Glucopyranosyl)-1*H*-1,2] 3 β -hydroxyolean-12-en-28-
oate (19).** Prepared from **16** (0.07 g, 0.08 mmol) according to
General procedure V. The residue was purified by column
chromatography (EtOAc). Yield: 0.05 g, 91%, white solid,
mp 178–180 °C, R_f = 0.13 (EtOAc); $[\alpha]_D$ = +45 (c = 0.05,

MeOH); IR (KBr, cm⁻¹): 3424, 2942, 1712, 1636, 1052, 1033, 1016, 772; ¹H NMR (300 MHz, C₅D₅N): δ 0.82, 0.85, 0.89, 0.93, 1.01, 1.17, 1.21 (7 s, each 3H, 7 \times CH₃), 0.82–1.92 (m, 22H), 3.08–3.11 (m, 1H, H-18), 3.39–3.44 (m, 1H, H-3), 4.20–4.22 (m, 1H, H-5-Glc), 4.27–4.41 (m, 3H, overlapping, H-6a-Glc, H-6b-Glc, H-4-Glc), 4.50 (m, 1H, H-2-Glc), 4.79 (t, 1H, J = 8.9, 8.9 Hz, H-3-Glc), 5.40 (s, 1H, H-12), 5.47–5.57 (m, 2H, COOCH₂), 6.35 (d, 1H, J = 9.2 Hz, H-1-Glc), 8.64 (s, 1H, NCH); ¹³C NMR (75 MHz, C₅D₅N): δ 15.7, 16.6, 17.4, 18.8, 23.4, 23.6, 23.8, 26.0, 28.1, 28.8, 30.8, 32.7, 33.1, 33.2, 34.0, 37.4, 39.0, 39.4, 39.8, 41.9, 42.1, 46.1, 47.0, 48.1, 55.9, 58.2, 62.4, 71.1, 73.9, 78.2, 79.1, 82.0, 89.6, 123.0, 124.1, 143.5, 144.0, 177.4. ESI-MS (positive mode) m/z : 744.3 [M + HCOO]⁺; HRMS (MALDI) m/z = C₃₉H₆₁N₃O₈ [M + Na]⁺ calcd. 722.4356, found 722.4371.

**1-(β -D-Glucopyranosyl)-1*H*-1,2] 3 β -hydroxyurs-12-en-28-
oate (20).** Prepared from **17** (0.14 g, 0.17 mmol) according to
General procedure V. The residue was purified by column
chromatography (EtOAc). Yield: 0.11 g, 92%, white solid, mp
160–162 °C, R_f = 0.16 (EtOAc); $[\alpha]_D$ = +23 (c = 0.1,
MeOH); IR (KBr, cm⁻¹): 3381, 2923, 2869, 1723, 1454,
1136, 1098, 1051, 1032, 1016, 772; ¹H NMR (300 MHz,
C₅D₅N): δ 0.81 (d, 3H, J = 4.7 Hz, CH₃), 0.83, 0.85, 0.89,
0.97, 1.06, 1.17 (6 s, each 3H, 6 \times CH₃), 2.34 (d, 1H, J = 11.2 Hz,
H-18), 0.75–1.94 (m, 22H), 3.38 (m, 1H, H-3), 4.09–4.17
25 (m, 1H, H-5-Glc), 4.22–4.36 (m, 3H, H-4-Glc and H-6-Glc),
4.46 (d, 1H, J = 10.8 Hz, H-2-Glc), 4.75 (t, 1H, J = 8.8, 8.8
Hz, H-3-Glc), 5.33 (brs, 1H, H-12), 5.44 (s, 2H, COOCH₂),
6.33 (d, 1H, J = 9.2 Hz, H-1-Glc), 8.58 (s, 1H, NCH); ¹³C
30 NMR (75 MHz, C₅D₅N): δ 15.8, 16.6, 17.3, 18.8, 19.1, 21.2,
23.7, 23.8, 24.6, 28.2, 28.8, 30.8, 33.5, 36.8, 37.3, 39.1,
39.2, 39.3, 39.4, 40.0, 42.4, 48.1, 48.4, 53.4, 55.9, 58.1, 62.4,
71.1, 73.9, 78.2, 79.1, 81.9, 89.6, 124.1, 126.2, 129.3, 138.6,
143.5, 177.1. ESI-MS (positive mode) m/z : 744.5 [M +
35 HCOO]⁺; HRMS (MALDI) m/z = C₃₉H₆₁N₃O₈ [M +
Na]⁺ calcd. 722.4356, found 722.4365.

**1-(β -D-Glucopyranosyl)-1*H*-1,2] 2 α ,3 β -dihydroxyolean-12-
en-28-oate (21).** Prepared from **18** (0.13 g, 0.15 mmol) according
40 to General procedure V. The residue was purified by column
chromatography (MeOH–CH₂Cl₂, 1:15). Yield: 0.09 g, 85%,
white solid, mp 205–207 °C, R_f = 0.18 (MeOH–CH₂Cl₂,
1:15); $[\alpha]_D$ = +27 (c = 0.07, MeOH). IR (KBr, cm⁻¹):
45 3461, 2945, 2864, 1720, 1642, 1457, 1051, 1031, 1017, 772, 667;
¹H NMR (300 MHz, C₅D₅N): δ 0.82, 0.85, 0.88, 1.03, 1.07,
1.15, 1.25 (7 s, each 3H, 7 \times CH₃), 0.82–2.26 (m, 20H),
3.07–3.10 (m, 1H, H-18), 3.29 (d, 1H, J = 9.3 Hz, H-3 α),
4.09 (m, 1H, H-2 β), 4.23–4.31 (m, 1H, H-5-Glc), 4.34–4.42
50 (m, 3H, overlapping, H-4-Glc and H-6a-Glc, and H-6b-Glc),
4.52 (d, 1H, J = 11.0 Hz, H-2-Glc), 4.80 (t, 1H, J = 8.9, 8.9
Hz, H-3-Glc), 5.37 (s, 1H, H-12), 5.52 (s, 2H, COOCH₂), 6.38
(d, 1H, J = 9.2 Hz, H-1-Glc), 8.65 (s, 1H, NCH); ¹³C NMR
(75 MHz, C₅D₅N): δ 17.0, 17.5, 17.7, 18.9, 23.4, 23.6, 23.9,
26.0, 28.1, 29.3, 30.7, 32.7, 33.1, 33.2, 33.9, 38.6, 39.8, 41.9,
42.1, 46.1, 47.0, 47.8, 48.1, 55.9, 58.2, 62.4, 68.6, 71.1, 73.9,
79.1, 81.9, 83.9, 89.6, 122.9, 124.1, 143.5, 144.0, 177.4. ESI-MS
(positive mode) m/z : 716.4 [M + H]⁺; HRMS (MALDI)
55 m/z = C₃₉H₆₁N₃O₉ [M + Na]⁺ calcd. 738.4306, found 738.4320.

1 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosylaminocarbonyl-**
methyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyolean-12-en-
28-oate (22). Prepared from **4** (0.11 g, 0.23 mmol) and **12**
2 residue was purified by column chromatography (EtOAc-
3 hexane, 1 : 1). Yield: 0.18 g, 85%, white solid, mp 146–148 °C,
 $R_f = 0.21$ (EtOAc–hexane, 1 : 1); IR (KBr, cm^{-1}): 2947, 2872,
4 1755, 1552, 1463, 1374, 1230, 1175, 1159, 1045, 759, 667; ^1H
NMR (300 MHz, CDCl_3): δ 0.57, 0.78, 0.98, 1.11 (4 s, each
5 3H, $4 \times \text{CH}_3$), 0.89 (s, 9H, $3 \times \text{CH}_3$), 0.57–2.10 (m, 22H), 2.01,
6 2.02, 2.03, 2.08 (4 s, each 3H, $4 \times \text{OCOCH}_3$), 2.82–2.85 (1H, m,
H-18), 3.20 (dd, 1H, $J = 5.0, 10.4$ Hz, H-3), 3.78–3.83 (m, 1H,
H-5-Glc), 4.07 (dd, 1H, $J = 2.1, 12.6$ Hz, H-6a-Glc), 4.28
7 (dd, 1H, $J = 4.3, 12.5$ Hz, H-6b-Glc), 4.90 (t, 1H, $J = 9.6, 9.6$ Hz,
H-4-Glc), 4.94–5.32 (m, 8H, overlapping, NCH_2 , COOCH_2 ,
8 H-1-Glc, H-2-Glc, H-3-Glc, H-12), 6.78 (d, 1H, $J = 8.7$ Hz,
OH), 7.69 (s, 1H, NCH); ^{13}C NMR (75 MHz, CDCl_3): δ 15.4,
9 15.6, 16.7, 18.3, 20.5, 20.7, 23.0, 23.4, 23.6, 26.0, 27.2, 27.6,
10 28.1, 30.6, 32.3, 32.7, 33.0, 33.8, 37.0, 38.4, 38.7, 39.3, 41.3,
11 41.7, 45.8, 46.7, 47.5, 52.5, 55.2, 57.3, 61.5, 68.0, 70.3, 72.4,
12 73.8, 78.5, 79.0, 122.5, 125.4, 143.5, 143.9, 165.4, 169.4, 169.8,
13 170.5, 171.0, 177.6. ESI-MS (positive mode) m/z : 947.6
14 $[\text{M} + \text{Na}]^+$. HRMS (MALDI) $m/z = \text{C}_{45}\text{H}_{72}\text{N}_4\text{O}_{10}$
15 $[\text{M} + \text{Na}]^+$ calcd. 852.0638, found 851.5158.

16 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosylaminocarbonyl-**
pentyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyolean-12-en-
28-oate (23). Prepared from **4** (0.10 g, 0.21 mmol) and **13** (0.10 g,
0.21 mmol) according to General procedure IVa. The residue
17 was purified by column chromatography (EtOAc–hexane,
1 : 1). Yield: 0.16 g, 80%, white solid, mp 99–100 °C, $R_f =$
18 0.12 (EtOAc–hexane, 1 : 1); IR (KBr, cm^{-1}): 2947, 2866, 1755,
19 1537, 1463, 1367, 1229, 1176, 1039, 757, 667; ^1H NMR
(300 MHz, CDCl_3): δ 0.55, 0.77, 0.89, 0.97, 1.11 (5 s, each
20 3H, $5 \times \text{CH}_3$), 0.88 (s, 6H, $2 \times \text{CH}_3$), 0.55–2.07 (m, 28H), 2.02,
21 2.03, 2.04, 2.07 (4 s, each 3H, $4 \times \text{OCOCH}_3$), 2.14–2.18
(m, 2H, CH_2CO), 2.82–2.85 (m, 1H, H-18), 3.20 (dd, 1H,
22 $J = 4.9, 10.6$ Hz, H-3), 3.79–3.84 (m, 1H, H-5-Glc), 4.07
(dd, 1H, $J = 2.1, 12.5$ Hz, H-6a-Glc), 4.28–4.34 (m, 3H,
23 overlapping, NCH_2 and H-6b-Glc), 4.90 (t, 1H, $J = 9.7,$
9.7 Hz, H-4-Glc), 5.06 (t, 1H, $J = 9.7, 9.7$ Hz, H-3-Glc), 5.20
24 (s, 2H, COOCH_2), 5.24–5.34 (m, 3H, overlapping, H-12, H-1-
Glc and H-12'), 6.23 (d, 1H, $J = 9.2$ Hz, NH), 7.56 (s, 1H,
25 NCH); ^{13}C NMR (75 MHz, CDCl_3): δ 15.3, 15.6, 16.7, 18.3,
26 20.6, 20.66, 20.7, 22.9, 23.4, 23.6, 24.2, 25.8, 25.9, 27.2, 27.6,
27 28.1, 29.9, 30.6, 32.3, 32.7, 33.0, 33.8, 36.0, 37.0, 38.4, 38.7,
28 39.3, 41.3, 41.7, 45.8, 46.7, 47.5, 49.9, 55.2, 57.4, 61.6, 68.1,
29 70.7, 72.6, 73.6, 77.2, 78.2, 79.0, 122.4, 123.9, 143.1, 143.6,
30 169.5, 169.8, 170.6, 171.1, 172.7, 177.8. ESI-MS (positive
31 mode) m/z : 1003.5 $[\text{M} + \text{Na}]^+$.

32 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosylaminocarbonyl-**
decyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyolean-12-en-
28-oate (24). Prepared from **4** (0.09 g, 0.18 mmol) and **14**
33 (0.10 g, 0.18 mmol) according to General procedure IVa.
The residue was purified by column chromatography (EtOAc–
34 hexane, 1 : 1). Yield: 0.12 g, 64%, white solid, mp 90–92 °C,
 $R_f = 0.22$ (EtOAc–hexane, 1 : 1); IR (KBr, cm^{-1}): 3359, 2922,
35 2860, 1744, 1693, 1364, 1220, 1049, 1032, 772; ^1H NMR

(300 MHz, CDCl_3): δ 0.52, 0.77, 0.87, 0.88, 0.90, 0.98, 1.11
1 (7 s, each 3H, $7 \times \text{CH}_3$), 0.52–2.07 (m, 38H), 2.02, 2.03, 2.04,
2.07 (4 s, each 3H, $4 \times \text{OCOCH}_3$), 2.13–2.23 (m, 2H, CH_2CO),
2.82 (m, 1H, H-18), 3.20 (dd, 1H, $J = 5.0, 10.9$ Hz, H-3),
3.80–3.85 (m, 1H, H-5-Glc), 4.07 (dd, 1H, $J = 1.9, 12.5$ Hz,
5 H-6a-Glc), 4.28–4.34 (m, 3H, overlapping, H-6b-Glc and
 NCH_2), 4.91 (pseudo t, 1H, $J = 9.6, 9.7$ Hz, H-4-Glc), 5.06
(pseudo t, 1H, $J = 9.6, 9.7$ Hz, H-3-Glc), 5.18 (s, 2H,
 COOCH_2), 5.22–5.34 (m, 3H, overlapping, H-12 and H-1-
Glc, H-2-Glc), 6.30 (d, 1H, $J = 9.3$ Hz, NH), 7.56 (s, 1H, s,
10 NCH); ^{13}C NMR (75 MHz, CDCl_3): δ 15.4, 15.6, 16.8, 18.3,
20.5, 20.6, 20.7, 23.0, 23.4, 23.6, 25.1, 25.8, 26.5, 27.2, 27.7,
28.1, 29.0, 29.1, 29.2, 29.3, 29.33, 30.2, 30.7, 32.4, 32.7, 33.0,
33.9, 36.6, 37.1, 38.5, 38.8, 39.4, 41.4, 41.7, 45.9, 46.8, 47.6,
50.4, 55.2, 57.5, 61.7, 68.3, 70.7, 72.8, 73.6, 77.2, 78.2, 79.0,
15 122.4, 123.9, 143.1, 169.5, 170.0, 170.5, 171.0, 173.3, 177.8.
ESI-MS (positive mode) m/z : 1073.9 $[\text{M} + \text{Na}]^+$.

16 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosylaminocarbonyl-**
methyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyurs-12-en-
28-oate (25). Prepared from **5** (0.11 g, 0.23 mmol) and **12**
17 (0.10 g, 0.23 mmol) according to General procedure IVa.
The residue was purified by column chromatography (EtOAc–
18 hexane, 1 : 1). Yield: 0.024 g, 11%, white solid, mp 116–118 °C,
 $R_f = 0.21$ (EtOAc–hexane, 1 : 1); $[\alpha]_D = +29$ ($c = 0.05,$
19 CHCl_3). IR (KBr, cm^{-1}): 3340, 2947, 2871, 1756, 1454, 1377,
1230, 1046, 1033, 997, 758; ^1H NMR (300 MHz, CDCl_3): δ
0.60, 0.78, 0.91, 0.93, 0.98, 1.06 (6 s, each 3H, $6 \times \text{CH}_3$), 0.83
20 (d, 3H, $J = 6.4$ Hz, CH_3), 0.60–2.01 (m, 22H), 2.01, 2.03, 2.08
(s, 12 H, $4 \times \text{OCOCH}_3$), 2.20 (d, 1H, $J = 11.3$ Hz, H-18), 3.22
21 (dd, 1H, $J = 4.9, 10.8$ Hz, H-3), 3.78–3.83 (m, 1H, H-5-Glc),
4.08 (dd, 1H, $J = 1.9, 12.7$ Hz, H-6a-Glc), 4.28 (dd, 1H, $J =$
22 4.3, 12.6 Hz, H-6b-Glc), 4.87 (pseudo t, 1H, $J = 9.5, 9.6$ Hz,
H-4-Glc), 4.99–5.09 (m, 3H, overlapping, NCH_2CO and H-2-
Glc), 5.16 (d, 1H, $J = 12.7$ Hz, COOCH_2), 5.19 (d, 1H, $J =$
23 12.7 Hz, COOCH_2), 5.22–5.23 (m, 2H, overlapping, H-1-Glc
and H-12), 5.29 (pseudo t, 1H, $J = 9.5, 9.6$ Hz, H-3-Glc), 6.75
24 (d, 1H, $J = 8.7$ Hz, NH), 7.68 (s, 1H, NCH); ^{13}C NMR
(75 MHz, CDCl_3): δ 15.5, 15.6, 16.9, 17.0, 18.3, 20.5, 20.7,
21.1, 23.3, 23.5, 24.2, 27.3, 28.0, 28.2, 30.6, 33.0, 36.6, 37.0,
25 38.7, 38.76, 38.84, 39.1, 39.6, 42.1, 47.6, 48.2, 52.6, 52.9, 55.3,
57.2, 61.6, 68.1, 70.4, 72.5, 73.9, 78.5, 79.1, 125.3, 125.8, 138.0,
144.0, 165.3, 169.4, 169.8, 170.5, 171.0, 177.4. ESI-MS
26 (positive mode) m/z : 947.0 $[\text{M} + \text{Na}]^+$.

27 **[1-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosylaminocarbonyl-**
pentyl)-1*H*-1,2,3-triazol-4-ylmethyl] 3 β -hydroxyurs-12-en-
28-oate (26). Prepared from **5** (0.10 g, 0.21 mmol) and **13**
28 (0.10 g, 0.21 mmol) according to General procedure IVa.
The residue was purified by column chromatography (EtOAc–
29 hexane, 1 : 1). Yield: 0.10 g, 50%, white solid, mp 102–104 °C,
 $R_f = 0.56$ (EtOAc–hexane, 2 : 1); IR (KBr, cm^{-1}): 3352, 2940,
30 2870, 1754, 1534, 1455, 1377, 1228, 1043, 756, 666; ^1H NMR
(300 MHz, CDCl_3): δ 0.59, 0.77, 0.90, 0.93, 0.98, 1.06 (6 s, each
31 3H, $6 \times \text{CH}_3$), 0.83 (d, $J = 6.4$ Hz, 3H, CH_3), 0.59–2.00
32 (m, 28H), 2.02, 2.03, 2.04, 2.08 (4 s, each 3H, $4 \times \text{OCOCH}_3$),
2.14–2.23 (m, 3H, overlapping, H-18 and CH_2CON), 3.21
33 (dd, 1H, $J = 4.0, 10.1$ Hz, H-3), 3.79–3.84 (m, 1H, H-5-
Glc), 4.07 (dd, 1H, $J = 2.1, 12.7$ Hz, H-6a-Glc), 4.29–4.34
34 (m, 3H, overlapping, H-6b-Glc and NCH_2), 4.90 (t, 1H, $J = 9.7,$
9.7 Hz, H-4-Glc), 5.06 (t, 1H, $J = 9.7, 9.7$ Hz, H-3-Glc), 5.20
35 (s, 2H, COOCH_2), 5.24–5.34 (m, 3H, overlapping, H-12, H-1-
Glc and H-12'), 6.23 (d, 1H, $J = 9.2$ Hz, NH), 7.56 (s, 1H,
NCH); ^{13}C NMR (75 MHz, CDCl_3): δ 15.3, 15.6, 16.7, 18.3,
20.6, 20.66, 20.7, 22.9, 23.4, 23.6, 24.2, 25.8, 25.9, 27.2, 27.6,
28.1, 29.9, 30.6, 32.3, 32.7, 33.0, 33.8, 36.0, 37.0, 38.4, 38.7,
39.3, 41.3, 41.7, 45.8, 46.7, 47.5, 49.9, 55.2, 57.4, 61.6, 68.1,
70.7, 72.6, 73.6, 77.2, 78.2, 79.0, 122.4, 123.9, 143.1, 143.6,
169.5, 169.8, 170.6, 171.1, 172.7, 177.8. ESI-MS (positive
mode) m/z : 1003.5 $[\text{M} + \text{Na}]^+$.

Q9 [3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonylmethyl)-1H-1,2] β 3-hydroxyolean-12-en-28-oate (31). Prepared from **22** (0.13 g, 0.14 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH-CH₂Cl₂, 1:15). Yield: 0.055 g, 51%, white solid, mp 197–199 °C, R_f = 0.09 (MeOH-CH₂Cl₂, 1:10); $[\alpha]_D^{25}$ = +37 (c = 0.11, MeOH); IR (KBr, cm⁻¹): 3407, 2943, 2860, 1705, 1556, 1389, 1161, 1059, 1032, 1018, 772; ¹H NMR (300 MHz, C₅D₅N): δ 0.80, 0.90, 1.03, 1.19, 1.24 (5 s, each 3H, 5 \times CH₃), 0.88 (s, 6H, 2 \times CH₃), 0.80–1.96 (m, 22H), 3.13 (d, 1H, J = 10.3 Hz, H-18), 3.43 (t, 1H, J = 7.8 Hz, H-3), 4.02–4.11 (m, 2H, overlapping, H-5-Glc and H-6a-Glc), 4.25–4.27 (m, 2H, overlapping, H-4-Glc and H-2-Glc), 4.37 (dd, 1H, J = 4.3, 11.8 Hz, H-6b-Glc), 4.49 (d, 1H, J = 11.5 Hz, H-1'), 5.43 (s, 1H, H-12), 5.48 (s, 2H, NCH₂CO), 5.62 (s, 2H, COOCH₂), 5.97 (t, 1H, J = 8.9, 8.9 Hz, H-3-Glc), 8.40 (s, 1H, NCH), 10.60 (d, 1H, J = 8.8 Hz, NH); ¹³C NMR (75 MHz, C₅D₅N): δ 15.6, 16.5, 17.1, 18.7, 23.3, 23.6, 23.7, 26.0, 28.0, 28.1, 28.7, 29.9, 30.7, 32.7, 33.05, 33.1, 33.9, 37.3, 38.9, 39.3, 39.7, 41.9, 42.0, 46.1, 46.9, 48.0, 52.7, 55.8, 58.0, 62.4, 71.5, 74.5, 78.1, 79.5, 80.3, 81.5, 123.1, 123.8, 126.7, 135.8, 143.5, 144.0, 150.2, 166.9, 177.3. ESI-MS (negative mode) m/z : 755.5 [M - H]⁺; HRMS (MALDI) m/z = C₄₁H₆₄N₄O₉ [M + Na]⁺ calcd. 779.4571, found 779.4594.

[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonylpentyl)-1H-1,2] β 3-hydroxyolean-12-en-28-oate (32). Prepared from **23** (0.11 g, 0.11 mmol) according to the general procedure V. The residue was purified by column chromatography (MeOH-CH₂Cl₂, 1:15). Yield: 0.076 g, 83%, white solid, mp 164–166 °C, R_f = 0.25 (MeOH-CH₂Cl₂, 1:10); $[\alpha]_D^{25}$ = +42 (c = 0.06, MeOH); IR (KBr, cm⁻¹): 3367, 2939, 2864, 1725, 1663, 1382, 1053, 1032, 1013, 773; ¹H NMR (300 MHz, C₅D₅N): δ 0.81, 0.94, 1.05, 1.20, 1.24 (5 s, each 3H, 5 \times CH₃), 0.89 (s, 6H, 2 CH₃), 0.81–1.93 (m, 28H), 2.41 (t, 2H, J = 7.3 Hz, CH₂CO), 3.15 (d, 1H, J = 13.3 Hz, H-18), 3.45 (brs, 1H, H-3), 4.05–4.13 (m, 2H, H-5-Glc and H-6a-Glc), 4.23–4.30 (m, 4H, H-2-Glc, H-4-Glc and NCH₂), 4.37 (dd, 1H, J = 4.6, 11.8 Hz, H-6b-Glc), 4.48 (d, 1H, J = 11.8 Hz, H-1-Glc), 5.43 (s, 1H, H-12), 5.53 and 5.54 (2 d, each 1H, J = 12.6 Hz, COOCH₂), 6.01 (t, 1H, J = 9.0, 9.0 Hz, H-3-Glc), 8.13 (s, 1H, NCH), 9.62 (d, 1H, J = 9.1 Hz, NH); ¹³C NMR (75 MHz, C₅D₅N): δ 15.6, 16.5, 17.2, 18.8, 23.3, 23.6, 23.8, 25.1, 26.0, 26.4, 28.0, 28.1, 28.8, 30.3, 30.7, 32.8, 33.1, 33.2, 33.9, 36.3, 37.3, 39.0, 39.4, 39.7, 41.9, 42.0, 46.1, 47.0, 48.0, 50.0, 55.8, 58.1, 62.7, 71.8, 74.6, 78.1, 79.7, 80.1, 81.3, 123.0, 124.6, 143.4, 144.0, 173.6, 177.4. ESI-MS (negative mode) m/z : 811.4 [M - H]⁺; HRMS (MALDI) m/z = C₄₅H₇₂N₄O₉ [M + Na]⁺ calcd. 835.5197, found 835.5203.

[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonyldecyl)-1H-1,2] β 3-hydroxyolean-12-en-28-oate (33). Prepared from **24** (0.083 g, 0.08 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH-CH₂Cl₂, 1:20). Yield: 0.068 g, 97%, white solid, mp 195–196 °C, R_f = 0.27 (MeOH-CH₂Cl₂, 1:15); $[\alpha]_D^{25}$ = +31 (c = 0.1, MeOH); IR (KBr, cm⁻¹): 3392, 2928, 1727, 1463, 1386, 1158, 1123, 1050, 1032, 1012, 756, 697; ¹H NMR (300 MHz, C₅D₅N): δ 0.59, 0.86, 1.04, 1.10 (4 s, each 3H,

4 \times CH₃), 0.74 (s, 9H, 3 \times CH₃), 0.59–1.95 (m, 38H), 2.36 (t, 2H, J = 7.3 Hz, CH₂CON), 2.94 (dd, 1H, J = 3.6, 10.0 Hz, H-18), 3.28 (t, 1H, J = 8.0 Hz, H-3), 3.92–3.93 (m, 2H, overlapping, H-5-Glc and H-6a-Glc), 4.04–4.10 (m, 3H, m, overlapping, NCH₂ and H-6b-Glc), 4.26–4.38 (m, 3H, m, overlapping, H-2-Glc, H-3-Glc, and H-4-Glc), 5.26 (s, 1H, H-12), 5.42 (d, 1H, J = 14.1 Hz, COOCH₂), 5.46 (d, 1H, J = 14.1 Hz, COOCH₂), 5.63 (d, 1H, J = 8.6 Hz, H-1-Glc), 8.18 (s, 1H, NCH), 9.67 (d, 1H, J = 8.8 Hz, NH); ¹³C NMR (75 MHz, C₅D₅N): δ 15.7, 16.6, 17.3, 18.8, 23.4, 23.7, 23.8, 25.9, 26.0, 26.7, 28.1, 28.8, 29.2, 29.6, 29.64, 29.7, 30.6, 30.8, 32.8, 33.1, 33.2, 34.0, 36.9, 37.4, 39.0, 39.4, 39.8, 41.9, 42.1, 46.1, 47.0, 48.0, 50.3, 55.8, 58.1, 62.7, 71.8, 74.5, 78.1, 79.6, 80.0, 81.4, 122.8, 124.7, 143.4, 144.0, 174.1, 177.4. ESI-MS (positive mode) m/z : 917.5 [M + Cl]⁺; HRMS (MALDI) m/z = C₅₀H₈₂N₄O₉ [M + Na]⁺ calcd. 905.5980, found 905.6013.

[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonylpentyl)-1H-1,2] β 3-hydroxyurs-12-en-28-oate (34). Prepared from **26** (0.065 g, 0.07 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH-CH₂Cl₂, 1:20). Yield: 0.053 g, 98%, white solid, mp 214–216 °C, R_f = 0.67 (MeOH-CH₂Cl₂, 1:15); $[\alpha]_D^{25}$ = +26 (c = 0.1, MeOH). IR (KBr, cm⁻¹): 3386, 2924, 2869, 1724, 1657, 1456, 1392, 1049, 1032, 1017; ¹H NMR (300 MHz, C₅D₅N): δ 0.83, 0.91, 0.94, 1.04, 1.13, 1.23 (6 s, each 3H, 6 \times CH₃), 0.88 (d, 3H, J = 3.9 Hz, CH₃), 0.83–2.00 (m, 28H), 2.39–2.47 (m, 3H, overlapping, H-18 and CH₂CON), 3.44 (dd, 1H, J = 6.1, 9.8 Hz, H-3), 4.04 (m, 1H, H-5-Glc), 4.16 (t, 1H, J = 9.8, 9.8 Hz, H-4-Glc), 4.24–4.45 (m, 6H, overlapping, NCH₂, H-2-Glc, H-3-Glc and H-6a-Glc, H-6b-Glc), 5.40 (s, 1H, H-12), 5.50 (s, 2H, COOCH₂), 5.96 (1H, overlapping, H-1'), 8.13 (s, 1H, NCH), 9.69 (d, 1H, J = 8.2 Hz, NH); ¹³C NMR (75 MHz, C₅D₅N): δ 15.8, 16.6, 17.3, 18.8, 21.2, 23.7, 23.8, 24.6, 25.1, 26.4, 28.1, 28.4, 28.8, 30.0, 30.3, 30.8, 33.5, 36.4, 37.0, 37.3, 39.1, 39.2, 39.3, 39.4, 40.0, 42.4, 48.0, 48.4, 50.1, 53.4, 55.8, 58.0, 62.7, 71.8, 74.5, 78.2, 79.6, 80.0, 81.3, 124.6, 126.2, 138.6, 143.3, 173.7, 177.1. ESI-MS (positive mode) m/z : 847.5 [M + Cl]⁺. HRMS (MALDI) m/z = C₄₅H₇₂N₄O₉ [M + Na]⁺ calcd. 835.5197, found 835.5206.

[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonyldecyl)-1H-1,2] β 3-hydroxyurs-12-en-28-oate (35). Prepared from **27** (0.10 g, 0.1 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH-CH₂Cl₂, 1:12). Yield: 0.078 g, 93%, white solid, mp 199–200 °C, R_f = 0.12 (MeOH-CH₂Cl₂, 1:15); $[\alpha]_D^{25}$ = +32 (c = 0.06, MeOH); IR (KBr, cm⁻¹): 3409, 2924, 2861, 1726, 1662, 1453, 1382, 1052, 1032, 1013. ¹H NMR (300 MHz, C₅D₅N): δ 0.81, 0.91, 0.94, 1.03, 1.13, 1.23 (6 s, each 3H, 6 \times CH₃), 0.88 (d, 3H, J = 3.6 Hz, CH₃), 0.81–1.99 (m, 38H), 2.43–2.49 (m, 3H, overlapping, H-18 and CH₂CON), 3.44 (dd, 1H, J = 6.8, 9.7 Hz, H-3), 4.04 (m, 1H, H-5-Glc), 4.14 (t, 1H, J = 8.8, 8.8 Hz, H-4-Glc), 4.21–4.33 (m, 2H, NCH₂), 4.35–4.47 (m, 3H, overlapping, H-3-Glc, H-2-Glc and H-1-Glc), 5.39 (brs, 1H, H-12), 5.51 and 5.52 (2 d, each 1H, J = 12.5 Hz, COOCH₂), 8.19 (s, 1H, NCH),

1 9.62 (d, 1H, $J = 9.0$ Hz, NH); ^{13}C NMR (75 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 15.8, 16.6, 17.3, 17.4, 18.8, 21.2, 23.7, 23.8, 24.6, 26.0, 26.8, 28.2, 28.5, 28.9, 29.3, 29.6, 29.67, 29.7, 30.0, 30.6, 30.8, 33.5, 36.9, 37.0, 37.4, 39.2, 39.3, 39.4, 40.0, 42.4, 48.0, 48.4, 50.3, 53.4, 55.9, 58.1, 62.8, 71.9, 74.6, 78.2, 79.7, 80.0, 81.4, 124.7, 126.2, 143.4, 174.1, 177.2. ESI-MS (positive mode) m/z : 927.5 $[\text{M} + \text{HCOO}]^+$; HRMS (MALDI) $m/z = \text{C}_{50}\text{H}_{82}\text{N}_4\text{O}_9$ $[\text{M} + \text{Na}]^+$ calcd. 905.5980, found 905.6004.

10 **[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonyl-pentyl)-1H-1,2] 2 α ,3 β -dihydroxyolean-12-en-28-oate (36).** Prepared from **29** (0.10 g, 0.1 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH– CH_2Cl_2 , 1:10). Yield: 0.075 g, 93%, white solid, mp 180–182 °C, $R_f = 0.09$ (MeOH– CH_2Cl_2 , 1:15); $[\alpha]_D = +23$ ($c = 0.05$, MeOH); IR (KBr, cm^{-1}): 3369, 2944, 2873, 1725, 1664, 1546, 1461, 1260, 1159, 1122, 1049, 1033; ^1H NMR (300 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 0.79, 0.90, 1.02, 1.09, 1.16 (5 s, each 3H, $5 \times \text{CH}_3$), 0.87 (s, 6H, $2 \times \text{CH}_3$), 0.79–2.30 (m, 26H), 2.40 (t, 2H, $J = 7.3$ Hz, CH_2CON), 3.11 (dd, 1H, $J = 4.2$, 13.5 Hz, H-18), 3.37 (d, 1H, $J = 9.3$ Hz, H-3 α), 4.10 (m, 1H, H-2 β), 4.15–4.49 (m, 6H, overlapping, H-1-Glc, H-2-Glc, H-4-Glc, H-5-Glc and H-6-Glc), 5.38 (s, 1H, H-12), 5.52 (s, 2H, COOCH_2), 5.99 (t, 1H, $J = 9.0$, 9.0 Hz, H-3-Glc), 8.11 (s, 1H, NCH), 9.62 (d, 1H, $J = 9.0$ Hz, NH); ^{13}C NMR (75 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 16.9, 17.3, 17.7, 18.9, 23.4, 23.7, 23.9, 25.1, 26.0, 26.5, 28.0, 29.3, 30.0, 30.4, 36.4, 38.6, 39.8, 41.9, 42.1, 46.1, 47.0, 47.8, 48.1, 50.0, 55.9, 58.2, 62.8, 68.6, 71.8, 74.6, 79.7, 80.0, 81.3, 83.8, 122.9, 124.6, 143.3, 144.0, 173.6, 177.4. ESI-MS (positive mode) m/z : 873.5 $[\text{M} + \text{HCOO}]^+$; HRMS (MALDI) $m/z = \text{C}_{45}\text{H}_{72}\text{N}_4\text{O}_{10}$ $[\text{M} + \text{Na}]^+$ calcd. 851.5146, found 851.5158.

35 **[3-Triazol-4-ylmethyl,1-(β -D-glucopyranosylaminocarbonyl-decyl)-1H-1,2] 2 α ,3 β -dihydroxyolean-12-en-28-oate (37).** Prepared from **30** (0.11 g, 0.1 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH– CH_2Cl_2 , 1:15). Yield: 0.065 g, 72%, white solid, mp 163–165 °C, $R_f = 0.18$ (MeOH– CH_2Cl_2 , 1:15); $[\alpha]_D = +33$ ($c = 0.07$, MeOH), IR (KBr, cm^{-1}): 3377, 2923, 2861, 1725, 1053, 1032, 1015, 772; ^1H NMR (300 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 0.79, 1.04, 1.10, 1.18 (4 s, each 3H, $4 \times \text{CH}_3$), 0.89 (s, 9H, $3 \times \text{CH}_3$), 0.79–2.30 (m, 36H), 2.48 (t, 2H, $J = 7.5$ Hz, CH_2CON), 3.12 (dd, 1H, $J = 3.8$, 9.9 Hz, H-18), 3.38 (d, 1H, $J = 9.2$ Hz, H-3 α), 4.05 (m, 1H, H-2 β), 4.07–4.51 (m, 8H, overlapping, H-1-Glc, H-2-Glc, H-4-Glc, H-5-Glc, H-6-Glc and NCH_2), 5.40 (s, 1H, H-12), 5.54 and 5.57 (2 d, each 1H, $J = 12.6$ Hz, COOCH_2), 6.03 (t, 1H, $J = 9.0$, 9.0 Hz, H-3-Glc), 8.12 (s, 1H, NCH), 9.61 (d, 1H, $J = 8.9$ Hz, NH); ^{13}C NMR (75 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 17.0, 17.3, 17.7, 18.9, 23.4, 23.9, 25.95, 26.0, 26.8, 28.0, 29.3, 29.4, 29.6, 29.66, 29.7, 30.0, 30.6, 30.8, 32.8, 33.1, 34.0, 36.9, 38.6, 39.9, 41.9, 42.1, 46.1, 47.0, 47.8, 48.1, 50.3, 55.9, 58.2, 62.8, 68.6, 71.9, 74.7, 79.8, 80.1, 81.4, 83.9, 122.9, 124.7, 143.4, 144.1, 174.0, 177.4. ESI-MS (positive mode) m/z : 943.6 $[\text{M} + \text{HCOO}]^+$; HRMS (MALDI) $m/z = \text{C}_{50}\text{H}_{82}\text{N}_4\text{O}_{10}$ $[\text{M} + \text{Na}]^+$ calcd. 921.5929, found 921.5937.

55 **2-[2,3-Triazol-1-yl,3 β -dihydroxyolean-12-en-28-carboxyloxy-methyl)-1H-1,4-(2 α) acetic acid (38).** Prepared from **28** (0.084 g,

0.09 mmol) according to General procedure V. The residue was purified by column chromatography (MeOH– CH_2Cl_2 , 1:10). Yield: 0.05 g, 91%, white solid, mp 225–227 °C, $R_f = 0.06$ (MeOH– CH_2Cl_2 , 1:15); $[\alpha]_D = +82$ ($c = 0.06$, MeOH); IR (KBr, cm^{-1}): 3403, 2940, 2862, 1724, 1450, 1386, 1229, 1159, 1050, 1033; ^1H NMR (300 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 0.84, 1.02, 1.07, 1.13, 1.17 (5 s, each 3H, $5 \times \text{CH}_3$), 0.87 (s, 6H, $2 \times \text{CH}_3$), 0.84–2.22 (m, 20H), 3.12 (dd, 1H, $J = 3.7$, 13.5 Hz, H-18), 3.37 (d, 1H, $J = 9.4$ Hz, H-3 α), 4.06–4.14 (m, 1H, H-2 β), 5.40 (s, 1H, H-12), 5.54 (s, 2H, NCH_2CO), 5.64 (s, 2H, COOCH_2), 5.67 (s, 2H, COOCH_2), 8.45 (s, 1H, NCH); ^{13}C NMR (75 MHz, $\text{C}_5\text{D}_5\text{N}$): δ 16.9, 17.3, 17.7, 18.9, 23.4, 23.7, 23.9, 26.0, 28.1, 29.3, 30.0, 30.8, 32.7, 33.1, 34.0, 38.6, 39.8, 39.9, 41.9, 42.1, 46.1, 47.0, 47.8, 48.1, 52.3, 55.9, 58.2, 68.6, 83.9, 122.6, 126.3, 143.5, 144.0, 177.4. ESI-MS (positive mode) m/z : 634.9 $[\text{M} + \text{Na}]^+$; HRMS (MALDI) $m/z = \text{C}_{35}\text{H}_{53}\text{N}_3\text{O}_6$ $[\text{M} + \text{Na}]^+$ calcd. 634.3832, found 634.3845.

General procedure VI for the Zemlén-deacetylation

20 To a solution of an *O*-acetyl-protected compound in dry MeOH 1–2 drops of a ~ 1 M methanolic NaOMe solution were added, and the reaction mixture was kept at rt until completion of the transformation (TLC, CHCl_3 –MeOH, 1:1). Amberlyst 15 (H^+ form) was then added to remove sodium ions, the resin was filtered off, and the solvent removed *in vacuo*. If the residue was chromatographically non-uniform it was purified by column chromatography or crystallisation.

General procedure VII for reduction

30 An *N*-(ω -azidoalkanoyl)- β -D-glucopyranosyl-amine was dissolved in dry MeOH (12 mL/mmol). To the solution RANEY[®]-Ni (~ 2 mmol) was added, and H_2 gas was bubbled through the mixture at 70 °C until the complete transformation of the starting azide TLC (CHCl_3 –MeOH, 1:1). The solution was filtered over a Celite pad and the solvent was removed *in vacuo*.

40 ***N*-Azidoacetyl- β -D-glucopyranosylamine (39).** Prepared from **12** (0.10 g, 0.23 mmol) according to General procedure VI. The residue was purified by column chromatography (CHCl_3 –MeOH, 7:3). Yield: 0.058 g, 95%, colourless oil, $R_f = 0.34$ (CHCl_3 –MeOH, 7:3); $[\alpha]_D = -12$ ($c = 0.22$, MeOH), (lit.⁴⁰ $[\alpha]_D = -61$ ($c = 1$, H_2O)); ^1H NMR (360 MHz, CD_3OD): δ (ppm) 3.29–3.37 (m, 3H, H-3, H-4, H-5), 3.44 (t, 1H, $J = 9.2$, 9.2 Hz, H-2), 3.68 (dd, 1H, $J = 4.0$, 11.9 Hz, H-6b), 3.83–3.99 (m, 3H, CH_2 , H-6a), 4.95 (d, 1H, $J = 9.2$ Hz, H-1). ^{13}C NMR (90 MHz, D_2O): δ (ppm) 52.8 (CH_2), 62.6 (C-6), 71.2, 73.8, 78.8, 79.7 (C-2, C-3, C-4, C-5), 81.0 (C-1), 171.3 (CONH). Analysis: Calcd for $\text{C}_8\text{H}_{14}\text{N}_4\text{O}_6$ (262.22): C, 36.64; H, 5.38; N, 21.37. Found: C, 36.73; H, 5.42; N, 21.25.

55 ***N*-Glycyl- β -D-glucopyranosylamine (40).** Prepared from **39** (0.097 g, 0.37 mmol) according to General procedure VII. Yield: 0.07 g, 79%, amorphous oil, $R_f = 0.16$ (MeOH); $[\alpha]_D = +34$ ($c = 0.08$, DMSO); ^1H NMR (360 MHz, CD_3OD): δ (ppm) 3.48–3.61 (m, 6H, H-2, H-3, H-4, H-5, CH_2), 3.76 (dd, 1H, $J = 5.3$, 11.9 Hz, H-6a), 3.90 (dd, 1H, $J = 1.2$, 11.9 Hz, H-6b), 5.04 (d, 1H, $J = 9.2$ Hz, H-1). ^{13}C NMR (90 MHz,

1 D₂O): δ(ppm) 44.1 (CH₂), 62.6 (C-6), 69.8, 72.4, 77.1, 78.2
(C-2, C-3, C-4, C-5), 79.9 (C-1), 176.5 (CONH). Analysis:
Calcd for C₈H₁₆N₂O₆ (236.22): C, 40.68; H, 6.83; N, 11.86.
Found: C, 40.75; H, 6.68; N, 11.79.

5 **N-(6-Azidohexanoyl)-β-D-glucopyranosylamine (41)**. Prepared
from **13** (0.50 g, 1.03 mmol) according to General procedure
VI. The residue was purified by column chromatography
(CHCl₃-MeOH, 7:3). Yield: 0.31 g (97%) colourless oil,
10 *R_f* = 0.66 (CHCl₃-MeOH, 7:3); [α]_D = +13 (*c* = 0.22,
MeOH); ¹H NMR (360 MHz, MeOD): δ(ppm) 1.42–1.49 (m,
2H, CH₂), 1.60–1.72 (m, 4H, CH₂), 2.25–2.32 (m, 2H, CH₂),
3.36–3.24 (m, 5H, H-3, H-4, H-5, CH₂), 3.44 (pseudo t, 1H,
15 *J* = 7.9, 9.2 Hz, H-2), 3.69 (dd, 1H, *J* = 5.3, 11.9 Hz, H-6b),
3.85 (dd, 1H, *J* = 1.2, 11.9 Hz, H-6a), 4.92 (d, 1H, *J* = 7.9 Hz,
H-1); ¹³C NMR (90 MHz, MeOD): δ (ppm) 26.0, 27.4, 29.6,
36.9, 52.3 (5 × CH₂), 62.6 (C-6), 71.4, 73.9, 79.0, 79.5 (C-2,
C-3, C-4, C-5), 80.9 (C-1), 177.0 (NHCO); Anal. calcd. for
20 C₁₂H₂₂N₄O₆ (318.33): C 45.28, H 6.97, N 17.60. Found:
C 45.36, H 6.84, N 17.49.

N-(6-Aminohexanoyl)-β-D-glucopyranosylamine (42). Prepared
from **41** (0.18 g 0.57 mmol) according to General procedure
VII. Yield: 0.09 g (57%) colourless oil, *R_f* = 0.05 (MeOH);
25 [α]_D = +11 (*c* = 0.15, MeOH); ¹H NMR (360 MHz, MeOD):
δ(ppm) 1.34–1.38 (m, 2H, CH₂), 1.47–1.51 (m, 2H, CH₂),
1.60–1.64 (m, 2H, CH₂), 2.21–2.27 (m, 2H, CH₂), 2.62–2.68
(m, 2H, CH₂), 3.23–3.35 (m, 3H, H-3, H-4, H-5), 3.38 (t, 1H,
30 *J* = 7.9, 7.9 Hz, H-2), 3.62 (dd, 1H, *J* = 5.3, 11.9 Hz, H-6a),
3.81 (dd, 1H, *J* = 1.2, 11.9 Hz, H-6a), 4.84 (d, 1H, *J* = 7.9 Hz,
H-1); ¹³C NMR (MeOD, 90 MHz): δ (ppm) 26.2, 27.4, 32.4,
36.9, 41.9 (5 × CH₂), 62.7 (C-6), 71.4, 73.9, 79.0, 79.7
(C-2, C-3, C-4, C-5), 81.0 (C-1), 177.2 (NHCO). Anal. calcd.
for C₁₂H₂₄N₄O₆ (292.33): C 49.30, H 8.28, N 9.58. Found:
35 C 49.36, H 8.18, N 9.45.

**1,4-Bis-[1-(2,6-tetra-*O*-acetyl-β-D-glucopyranosylaminocarbonyl-
methyl)-1*H*-1,3-triazol-4-yl]butane (44)**. Prepared from **12**
(0.30 g 0.41 mmol) according to General procedure IVb. The
residue was purified by column chromatography (EtOAc).
40 Yield: 0.10 g, 89%, white crystalline product, mp 197–199 °C,
[α]_D = +35 (*c* = 0.20, DMSO); ¹H NMR (360 MHz,
DMSO-*d*₆): δ(ppm) 1.64 (brs, 4H, 2 × CH₂), 1.93, 1.95,
1.99, 2.00 (4s, 24H, 8 × OCOCH₃), 2.65 (brs, 4H, 2 × CH₂),
3.96–4.16 (m, 6H, 2 × H-5-Glc, 2 × H-6a-Glc, 2 ×
45 H-6b-Glc), 5.07 (brs, 4H, 2 × CH₂), 4.86, 4.92, 5.34, 5.42
(4t, 8 H, *J* = 9.2, 9.2 Hz in each, 2 H-1-Glc, 2 × H-2-Glc, 2 ×
H-3-Glc, 2 × H-4-Glc), 7.78 (s, 2H, 2 triazole CH), 9.20
(d, 2H, *J* = 9.2 Hz, 2 × NH); ¹³C NMR (90 MHz, DMSO-*d*₆):
50 δ(ppm) 20.3, 20.5, (8 × OCOCH₃), 24.7, 28.4, 51.3 (6 × CH₂),
61.6 (2 × C-6-Glc), 67.7, 70.5, 72.1, 72.7 (2 C-2-Glc, 2 ×
C-3-Glc, 2 × C-4-Glc, 2 × C-5-Glc), 76.8 (2 × C-1-Glc), 123.4
(2 triazole C-5), 146.4 (2 triazole C-4), 166.4 (2 × CONH),
169.2, 169.3, 169.5, 170.0 (8 × OCOCH₃). Anal. calcd. for
55 C₄₀H₅₄N₈O₂₀ (966.92): C, 49.69; H, 5.63; N, 11.59; Found: C,
49.59; H, 5.71; N, 11.67.

**1,4-Bis-[1-(2,6-tetra-*O*-acetyl-β-D-glucopyranosylaminocarbonyl-
pentyl)-1*H*-1,3-triazol-4-yl]butane (45)**. Prepared from **13**
(0.20 g, 0.41 mmol) according to General procedure IVb.

The residue purified by column chromatography (EtOAc-
MeOH, 95:5). Yield: 0.21 g, 96%, colourless oil, *R_f* = 0.32
(EtOAc); [α]_D = +16 (*c* = 0.16, CHCl₃); ¹H NMR (360
MHz, CDCl₃): δ(ppm) 1.16–1.30 (m, 4H, 2 × CH₂), 1.62–1.74
5 (m, 4H, 2 × CH₂), 1.85–1.91 (m, 4H, 2 × CH₂), 2.01, 2.03,
2.04, 2.07 (4 s, 24H, 8 × OCOCH₃), 2.18–2.22 (m, 4H, 2 ×
CH₂), 2.52–2.56 (m, 2H, CH₂), 2.72–2.76 (m, 4H, 2 × CH₂),
3.84 (ddd, 2H, *J* = 1.1, 2.6, 10.6 Hz, 2 × H-5-Glc), 4.13–4.07
(m, 4H, 2 × CH₂), 4.27–4.31 (m, 6H, 2 × H-6a-Glc, 2 ×
10 H-6b-Glc, CH₂), 5.28, 5.24, 5.06, 4.93 (4 pseudo t, 8H, *J* =
9.2, 10.6 Hz in each, 2 × H-1-Glc, 2 × H-2-Glc, 2 × H-3-Glc,
2 × H-4-Glc), 6.57 (d, 2H, *J* = 7.9 Hz, 2 × NH), 7.34 (s, 2H,
2 triazole CH); ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.3,
20.4, (8 × OCOCH₃), 24.1, 25.1, 25.7, 28.6, 29.7, 35.7, 49.5
(14 × CH₂), 61.6 (2 × C-6-Glc), 68.0, 70.4, 72.7, 73.3 (2 ×
15 C-2-Glc, 2 × C-3-Glc, 2 × C-4-Glc, 2 × C-5-Glc), 77.8 (2 ×
C-1-Glc), 120.6 (2 triazole C-5), 147.7 (2 triazole C-4), 169.4,
169.6, 170.0, 170.4 (8 × OCOCH₃), 172.9 (2 × NHCO). Anal.
calcd. for C₄₈H₇₀N₈O₂₀ (1079.13): C 53.43, H 6.54, N 10.38.
Found: C 53.49, H 6.62, N 10.45.

**1,4-Bis-[1-(2,6-tetra-*O*-acetyl-β-D-glucopyranosylaminocarbonyl-
decyl)-1*H*-1,3-triazol-4-yl]butane (46)**. Prepared from **14**
(0.20 g 0.36 mmol) according to General procedure IVb. The
residue purified by column chromatography (EtOAc). Yield:
25 0.136 g, 62%, colourless oil; [α]_D = +7 (*c* = 0.62, CHCl₃); ¹H
NMR (CDCl₃, 90 MHz): δ (ppm) 1.26 (brs, 24H, 12 × CH₂),
1.56–1.59 (m, 4H, 2 × CH₂), 1.73–1.78 (m, 4H, 2 × CH₂),
1.85–1.89 (m, 4H, 2 × CH₂), 2.02, 2.03, 2.04, 2.08 (4s, 24H,
8 × OCOCH₃), 2.10–2.25 (m, 4H, 2 × CH₂), 2.73–2.76 (m, 4H,
30 2 × CH₂), 3.83 (m, 2H, *J* = 2.4, 4.3, 9.9 Hz, 2 × H-5-Glc), 4.08
(dd, 2H, *J* = 2.3, 12.3 Hz, 2 × H-6b-Glc), 4.27–4.34 (m, 6H, 2
× H-6a-Glc, 2 × CH₂), 4.93, 5.06, 5.27, 5.31 (4 pseudo t, 8 H,
J = 9.6, 9.9 Hz in each, 2 × H-1-Glc, 2 × H-2-Glc, 2 × H-3-
35 Glc, 2 × H-4-Glc), 6.42 (d, 2H, *J* = 9.2 Hz, 2 × NH), 7.28
(s, 2H, 2 triazole CH); ¹³C NMR (90 MHz, CDCl₃): δ (ppm)
20.4, 20.5, 20.6 (8 × OCOCH₃), 25.0, 25.3, 26.3, 28.7, 28.8,
28.9, 29.0, 29.1, 29.2, 29.6, 30.2, 50.0, (24 × CH₂), 61.6 (2 ×
C-6-Glc), 68.1, 70.5, 72.6, 73.4 (2 C-2-Glc, 2 × C-3-Glc, 2 ×
C-4-Glc, 2 × C-5-Glc), 78.0 (2 × C-1-Glc), 120.4 (2 triazole
40 C-5), 147.8 (2 triazole C-4), 169.5, 169.8, 170.5, 170.8 (8 ×
OCOCH₃), 173.4 (2 × CONH). Anal. calcd. for C₅₈H₉₀N₈O₂₀
(1219.38): C, 57.13; H, 7.44; N, 9.19; Found: C, 57.22; H, 7.32;
N, 9.30.

**1,4-Bis-[1-(2,6-tetra-*O*-acetyl-β-D-glucopyranosylaminocarbonyl-
pentadecyl)-1*H*-1,3-triazol-4-yl]butane (47)**. Prepared from **15**
(0.30 g 0.41 mmol) according to General procedure IV. The
residue purified by column chromatography (EtOAc). Yield:
50 0.28 g, 50%, white crystalline product, mp 144–146 °C; [α]_D =
+14 (*c* = 0.24, CHCl₃); ¹H NMR (360 MHz, CDCl₃): δ(ppm)
1.20–1.35 (m, 26H, 13CH₂), 1.40–1.50 (m, 8H, 2 × CH₂),
1.60–1.77 (m, 8H, 2 × CH₂), 1.82–1.90 (m, 8H, 2 × CH₂), 2.02,
2.03, 2.05, 2.06 (4 s, 24H, 8 × OCOCH₃), 2.32–2.27 (m, 2H,
CH₂), 2.60–2.52 (m, 4H, 2 × CH₂), 2.73–2.80 (m, 2H, CH₂),
55 3.70 (ddd, 2H, *J* = 1.1, 2.6, 10.6 Hz, 2 × H-5-Glc), 4.11–4.20
(m, 8H, 4 CH₂), 4.37–4.44 (m, 4H, 2 × H-6a-Glc, 2 × H-6b-Glc),
4.94, 5.00, 5.11, 5.19 (4 pseudo t, 8H, *J* = 9.2, 10.6 Hz in each,
2 × H-1-Glc, 2 × H-2-Glc, 2 × H-3-Glc, 2 × H-4-Glc),

1 6.73 (d, 2H, $J = 7.9$ Hz, 2 NH), 7.34 (s, 2H, 2 triazole CH); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 20.4, 20.5 (8 \times OCOCH_3), 22.1, 22.5, 23.0, 23.6, 24.8, 25.1, 25.8, 26.1, 26.6, 27.0, 27.5, 29.8, 30.7, 31.7, 35.7, 36.7, 50.1 (32 \times CH_2), 60.6 (2 \times C-6-Glc), 69.3, 71.0, 72.0, 72.8 (2 \times C-2-Glc, 2 \times C-3-Glc, 2 \times C-4-Glc, 2 \times C-5-Glc), 77.6 (2 \times C-1-Glc), 121.0 (2 triazole C-5), 146.6 (2 triazole C-4), 169.4, 169.6, 169.9, 170.1 (8 \times OCOCH_3), 171.8 (2 \times HHCO). Anal. calcd. for $\text{C}_{68}\text{H}_{110}\text{N}_8\text{O}_{20}$ (1359.68): C, 60.07; H, 8.15; N, 8.24; Found: C, 60.16; H, 8.22; N, 8.33.

Q10 **1,4-Bis-[1-(β -D-glucopyranosylaminocarbonylmethyl)-1H-1,3-triazol-4-yl],2]butane (48).** Prepared from **44** (0.10 g, 0.10 mmol) according to General procedure VI. Yield: 0.042 g, 65%, white crystalline product, mp 235–236 °C, $[\alpha]_{\text{D}} = 42$ ($c = 0.22$, DMSO); ^1H NMR (360 MHz, D_2O): δ (ppm) 1.64 (brs, 4H, 2 \times CH_2), 2.70 (brs, 4H, 2 \times CH_2), 3.39–3.57 (8H, m, 2 \times H-2-Glc, 2 \times H-3-Glc, 2 \times H-4-Glc, 2 \times H-5-Glc), 3.71 (dd, 2H, $J = 5.3$ 11.9 Hz, 2 \times H-6b-Glc), 5.01 (d, 2H, $J = 9.2$ Hz, 2 \times H-1-Glc), 3.85 (dd, 2H, $J = <1$, 11.9 Hz, 2 \times H-6a-Glc), 5.24–5.27 (s, 4H, 2 \times CH_2), 7.73 (s, 2H, triazole CH); ^{13}C NMR (90 MHz, DMSO- d_6): δ (ppm) 24.7, 28.4, 51.5 (6 \times CH_2), 60.8 (2 \times C-6-Glc), 69.9, 72.6, 77.3, 78.7 (2 \times C-2-Glc, 2 \times C-3-Glc, 2 \times C-4-Glc, 2 \times C-5-Glc), 79.7 (2 \times C-1-Glc), 123.5 (2 triazole C-5), 146.5 (2 triazole C-4), 166.1 (2 \times CONH). Anal. calcd. for $\text{C}_{24}\text{H}_{38}\text{N}_8\text{O}_{12}$ (630.62): C, 45.71; H, 6.07; N, 17.77; Found: C, 45.80; H, 5.97; N, 17.54.

1,4-Bis-[1-(β -D-glucopyranosylaminocarbonylpentyl)-1H-1,3-triazol-4-yl],2]butane (49). Prepared from **45** (0.21 g 0.19 mmol) according to General procedure VI. Yield: 0.09 g, 66%, white crystalline product, mp: 150–152 °C; $[\alpha]_{\text{D}} = +14$ ($c = 0.12$, MeOH); ^1H NMR (360 MHz, D_2O): δ (ppm) 1.20–1.25 (m, 4H, 2 \times CH_2), 1.52–1.64 (m, 8H, 4 \times CH_2), 1.80–1.88 (m, 4H, 2 \times CH_2), 2.22–2.28 (m, 4H, 2 \times CH_2), 2.66–2.70 (m, 4H, 2 \times CH_2), 3.33–3.42 (m, 6H, 2 \times H-3-Glc, 2 \times H-4-Glc, CH_2), 3.50 (ddd, 2H, $J = 1.2$, 5.3, 9.2 Hz, 2 \times H-5-Glc), 3.52 (t, 2H, $J = 9.2$, 9.2 Hz, 2 \times H-2-Glc), 3.70 (dd, 2H, $J = 5.3$, 11.9 Hz, 2 \times H-6b-Glc), 3.85 (dd, 2H, $J = 1.2$, 11.9 Hz, 2 \times H-6a-Glc), 4.31–4.37 (m, 4H, 2 \times CH_2), 4.91 (d, 2H, $J = 9.2$ Hz, 2 \times H-1-Glc), 7.82 (s, 2H, 2 triazole CH); ^{13}C NMR (90 MHz, D_2O): δ (ppm) 24.0, 24.1, 25.2, 27.8, 29.1, 35.6, 50.4 (14 \times CH_2), 60.7 (2 \times C-6-Glc), 69.4, 71.9, 76.7, 77.7 (2 \times C-2-Glc, 2 \times C-3-Glc, 2 \times C-4-Glc, 2 \times C-5-Glc), 79.4 (2 \times C-1-Glc), 123.8 (2 triazole C-5), 146.2 (2 triazole C-4), 178.1 (2 \times NHCO). Anal. calcd. for $\text{C}_{32}\text{H}_{54}\text{N}_8\text{O}_{12}$ (742.83): C 51.74, H 7.33, N 15.08. Found: C 51.69, H 7.25, N 15.02.

1,4-Bis-[1-(β -D-glucopyranosylaminocarbonyldecyl)-1H-1,3-triazol-4-yl],2]butane (50). Prepared from **46** (0.07 g, 0.06 mmol) according to General procedure VI. Precipitated from the reaction mixture. Yield: 0.048 g, 95%, white amorphous product; $[\alpha]_{\text{D}} = +16$ ($c = 0.37$, DMSO); ^1H NMR (360 MHz, DMSO- $d_6 + \text{D}_2\text{O}$): δ (ppm) 1.20 (brs, 24H, 12 \times CH_2), 1.45, 1.60, 1.75, 2.06, 2.60 (5 brs, 20H, 10 \times CH_2), 3.00–3.19 (m, 8H, 2 \times H-2-Glc, 2 \times H-3-Glc, 2 \times H-4-Glc, 2 \times H-5-Glc), 3.38 (dd, 2H, $J = 4.6$, 11.6 Hz, 2 \times H-6b-Glc), 3.61 (dd, 2H, $J < 1.0$, 11.2 Hz, 2 \times H-6a-Glc), 4.23–4.27 (m, 4H, 2 \times CH_2), 4.68 (d, 2H, $J = 8.9$ Hz, 2 \times H-1-Glc), 7.80 (s, 2H, 2 triazole CH); ^{13}C NMR (90 MHz, DMSO- $d_6 + \text{D}_2\text{O}$):

δ (ppm) 24.8, 25.0, 25.9, 28.4, 28.5, 28.9, 29.7, 35.5, 49.2 (24 \times CH_2), 60.9 (2 \times C-6-Glc), 69.9, 72.3, 77.4, 78.4 (2 \times C-2-Glc, 2 \times C-3-Glc, 2 \times C-4-Glc, 2 \times C-5-Glc), 79.4 (2 \times C-1-Glc), 121.7 (2 triazole C-5), 146.8 (2 triazole C-4), 173.0 (2 \times CONH). Anal. calcd. for $\text{C}_{42}\text{H}_{74}\text{N}_8\text{O}_{12}$ (883.08): C, 57.12; H, 8.45; N, 12.69; Found: C, 57.23; H, 8.56; N, 12.58.

1,4-Bis-[1-(β -D-glucopyranosylaminocarbonylpentadecyl)-1H-1,2,3-triazol-4-yl],2]butane (51). Prepared from **47** (0.2 g, 0.147 mmol) according to General procedure VI. Yield: 0.12 g, 80%, white crystalline product, mp 158–160 °C; $[\alpha]_{\text{D}} = +25$ ($c = 0.20$, MeOH); ^1H NMR (360 MHz, D_2O): δ (ppm) 1.19–1.32 (m, 30H, 15 \times CH_2), 1.49–1.70 (m, 4H, 2 \times CH_2), 1.91–1.98 (m, 4H, 2 \times CH_2), 2.18–2.38 (m, 8H, 4 \times CH_2), 2.51–2.90 (m, 20H, 10 \times CH_2), 3.32–3.39 (m, 4H, 2 \times H-3-Glc, 2 \times H-4-Glc), 3.45 (ddd, 2H, $J = 1.2$, 5.3, 9.2 Hz, 2 \times H-5-Glc), 3.49 (t, 2H, $J = 9.2$, 9.2 Hz, 2 \times H-2-Glc), 3.79 (dd, 2H, $J = 5.3$, 11.9 Hz, 2 \times H-6b-Glc), 3.87 (2H, dd, $J = 1.2$, 11.9 Hz, 2 \times H-6a-Glc), 4.41–4.47 (m, 4H, 2 \times CH_2), 5.01 (d, 2H, $J = 9.2$ Hz, 2 \times H-1-Glc), 7.67 (s, 2H, 2 triazole CH); ^{13}C NMR (90 MHz, D_2O): δ (ppm) 21.7, 22.9, 23.3, 24.1, 24.8, 25.3, 25.8, 27.1, 27.6, 28.1, 29.5, 31.2, 32.0, 33.1, 36.7, 37.4, 51.2 (34 \times CH_2), 61.5 (2 \times C-6-Glc), 69.0, 72.1, 76.3, 77.3 (2 \times C-2-Glc, 2 \times C-3-Glc, 2 \times C-4-Glc, 2 \times C-5-Glc), 78.1 (2 \times C-1-Glc), 121.5 (2 triazole C-5), 146.5 (2 triazole C-4), 179.3 (2 \times NHCO). Anal. calcd. for $\text{C}_{52}\text{H}_{94}\text{N}_8\text{O}_{12}$ (1023.38): C, 61.06; H, 9.26; N, 10.95; Found: C, 61.13; H, 9.36; N, 10.88.

Enzyme assays

(a) Against RMGP_a. The inhibitory activity of the prepared compounds against rabbit muscle glycogen phosphorylase a (RMGP_a) was monitored using microplate reader (BIO-RAD) based on the published method.³⁵ In brief, GP_a activity was measured in the direction of glycogen synthesis by the release of phosphate from glucose-1-phosphate. Each prepared compound was dissolved in DMSO and diluted to different concentrations for IC_{50} determination. The enzyme was added into 100 μL of buffer containing 50 mM Hepes (pH = 7.2), 100 mM KCl, 2.5 mM MgCl_2 , 0.5 mM glucose-1-phosphate, 1 mg/ml glycogen and the test compound in 96-well microplates (Costar). After the addition of 150 μL of 1 M HCl containing 10 mg/ml ammonium molybdate and 0.38 mg/ml malachite green, reactions were run at 22 °C for 25 min, and then the phosphate absorbance was measured at 655 nm. The IC_{50} values were estimated by fitting the inhibition data to a dose-dependent curve using a logistic derivative equation.

(b) Against RMGP_b. Glycogen phosphorylase b (RMGP_b) was prepared from rabbit skeletal muscle according to the method of Fischer and Krebs,⁴¹ using dithiothreitol instead of L-cysteine, and recrystallized at least three times before use. Kinetic experiments were performed in the direction of glycogen synthesis using RMGP_b as described.^{36,42} IC_{50} values were determined in the presence of 4 mM α -D-glucose-1-phosphate, 1 mM AMP, 1% glycogen and varying concentrations of the inhibitor.⁴³ Inhibitors were dissolved in dimethyl sulfoxide (DMSO) and diluted in the assay buffer (50 mM triethanolamine, 1 mM EDTA and 1 mM dithiothreitol) so that the DMSO concentration in the assay should

1 be lower than 1%. The means of standard errors for all
calculated kinetic parameters averaged to less than 10%.

Acknowledgements

5 HBS thanks financial support by National Natural Science
Foundation of China (Grants 30672523 and 90713037),
research grants from Chinese Ministry of Education (Grants
706030 and 20050316008) and the program for New Century
10 Excellent Talents in University (NCET-05-0495). In Hungary
this research was supported by the National Office for
Research and Technology as well as the Hungarian Scientific
Research Fund (Grants NI-61336, NK-68578, and
CK-77712). The authors also thank NORT (Hungary, CHN-
15 25/05) and the Program for Chinese–Hungarian Scientific and
Technological Cooperation (China, No. CHN-7/2006) for
supporting this bilateral R&D project.

References

- 1 D. E. Moller, *Nature*, 2001, **414**, 821–827.
- 2 P. Zimmet, K. G. M. M. Alberti and J. Shaw, *Nature*, 2001, **414**, 782–861.
- 3 J. Diamond, *Nature*, 2003, **423**, 599–602.
- 4 P. Staehr, O. Hother-Nielsen and H. Beck-Nielsen, *Diabetes, Obes. Metab.*, 2002, **4**, 215–223.
- 5 N. G. Oikonomakos and L. Somsák, *Curr. Opin. Invest. Drugs*, 2008, **9**, 379–395.
- 6 L. Somsák, K. Czifrák, M. Tóth, É. Bokor, E. D. Chrysina, K. M. Alexacou, J. M. Hayes, C. Tiraidis, E. Lazoura, D. D. Leonidas, S. E. Zographos and N. G. Oikonomakos, *Curr. Med. Chem.*, 2008, **15**, 2933–2983.
- 7 D. J. Baker, P. L. Greenhaff and J. A. Timmons, *Expert Opin. Ther. Pat.*, 2006, **16**, 459–466.
- 8 L. Agius, *Best Pract. Res. Clin. Endocrinol. Metab.*, 2007, **21**, 587–605.
- 9 N. G. Oikonomakos, *Curr. Protein Pept. Sci.*, 2002, **3**, 561–586.
- 10 E. D. Chrysina, M. N. Kosmopolou, C. Tiraidis, R. Kardarakis, N. Bischler, D. D. Leonidas, Z. Hadady, L. Somsák, T. Docsa, P. Gergely and N. G. Oikonomakos, *Protein Sci.*, 2005, **14**, 873–888.
- 11 L. Somsák, V. Nagy, Z. Hadady, T. Docsa and P. Gergely, *Curr. Pharm. Des.*, 2003, **9**, 1177–1189.
- 12 L. Somsák, V. Nagy, Z. Hadady, N. Felföldi, T. Docsa and P. Gergely, in *Frontiers in Medicinal Chemistry*, ed. A. B. Reitz, C. P. Kordik, M. I. Choudhary and A. U. Rahman, Bentham, 2005, pp. 253–272.
- 13 J. Chen, J. Liu, Y. C. Gong, L. Y. Zhang, W. Y. Hua and H. B. Sun, *J. China Pharm. Univ.*, 2006, **37**, 397–402.
- 14 J. Chen, J. Liu, L. Y. Zhang, G. Z. Wu, W. Y. Hua, X. M. Wu and H. B. Sun, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 2915–2919.
- 15 K. G. Cheng, P. Zhang, J. Liu, J. Xie and H. B. Sun, *J. Nat. Prod.*, 2008, **71**, 1877–1880.
- 16 X. Wen, H. Sun, J. Liu, K. Cheng, P. Zhang, L. Zhang, J. Hao, L. Zhang, P. Ni, S. E. Zographos, D. D. Leonidas, K. M. Alexacou, T. Gimisis, J. M. Hayes and N. G. Oikonomakos, *J. Med. Chem.*, 2008, **51**, 3540–3554.
- 17 P. Dzubak, M. Hajdúch, D. Vydra, A. Hustova, M. Kvasnica, D. Biedermann, L. Markova, M. Urban and J. Sarek, *Nat. Prod. Rep.*, 2006, **23**, 394–411.

- 18 C. D. Shomin, S. C. Meyer and I. Ghosh, *Bioorg. Med. Chem.*, 2009, **17**, 6196–6202.
- 19 H. Li, Y. J. Guan, A. Szczepanska, A. J. Moreno-Vargas, A. T. Carmona, I. Robina, G. K. Lewis and L. X. Wang, *Bioorg. Med. Chem.*, 2007, **15**, 4220–4228.
- 20 J. L. Arolas, G. M. Popowicz, J. Lorenzo, C. P. Sommerhoff, R. Huber, F. X. Aviles and T. A. Holak, *J. Mol. Biol.*, 2005, **350**, 489–498.
- 21 T. Selwood, K. C. Elrod and N. M. Schechter, *Biol. Chem.*, 2003, **384**, 1605–1611.
- 22 X. Wu and B. S. Cooperman, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 2387–2389.
- 23 T. Steinmetzer, M. Renatus, S. Kunzel, A. Eichinger, W. Bode, P. Wikstrom, J. Hauptmann and J. Sturzebecher, *Eur. J. Biochem.*, 1999, **265**, 598–605.
- 24 J. J. Lundquist and E. J. Toone, *Chem. Rev.*, 2002, **102**, 555–578.
- 25 S. Cecioni, O.-A. Argintaru, T. Docsa, P. Gergely, J.-P. Praly and S. Vidal, *New J. Chem.*, 2009, **33**, 148–156.
- 26 V. L. Rath, M. Ammirati, D. E. Danley, J. L. Ekstrom, E. M. Gibbs, T. R. Hynes, A. M. Mathiowetz, R. K. McPherson, T. V. Olson, J. L. Treadway and D. J. Hoover, *Chem. Biol.*, 2000, **7**, 677–682.
- 27 S. Furukawa, Y. Tsurumi, K. Murakami, T. Nakanishi, K. Ohsumi, M. Hashimoto, M. Nishikawa, S. Takase, O. Nakayama and M. Hino, *J. Antibiot.*, 2005, **58**, 497–502.
- 28 C. Tiraidis, K. M. Alexacou, S. E. Zographos, D. D. Leonidas, T. Gimisis and N. G. Oikonomakos, *Protein Sci.*, 2007, **16**, 1773–1782.
- 29 K. Cheng, J. Liu, X. Liu, H. Li, H. Sun and J. Xie, *Carbohydr. Res.*, 2009, **344**, 841–850.
- 30 M. Meldal and C. W. Tornøe, *Chem. Rev.*, 2008, **108**, 2952–3015.
- 31 X. Wen, P. Zhang, J. Liu, L. Zhang, X. Wu, P. Ni and H. Sun, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 722–726.
- 32 Z. Györgydeák and J. Thiem, *Adv. Carbohydr. Chem. Biochem.*, 2006, **60**, 103–182.
- 33 L. Kovács, E. Ósz, V. Domokos, W. Holzer and Z. Györgydeák, *Tetrahedron*, 2001, **57**, 4609–4621.
- 34 E. I. Petsalakis, E. D. Chrysina, C. Tiraidis, T. Hadjiloi, D. D. Leonidas, N. G. Oikonomakos, U. Aich, B. Varghese and D. Loganathan, *Bioorg. Med. Chem.*, 2006, **14**, 5316–5324.
- 35 W. H. Martin, D. J. Hoover, S. J. Armento, I. A. Stock, R. K. McPherson, D. E. Danley, R. W. Stevenson, E. J. Barrett and J. L. Treadway, *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 1776–1781.
- 36 E. Ósz, L. Somsák, L. Szilágyi, L. Kovács, T. Docsa, B. Tóth and P. Gergely, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 1385–1390.
- 37 E. D. Chrysina, É. Bokor, K.-M. Alexacou, M.-D. Charavgi, G. N. Oikonomakos, S. E. Zographos, D. D. Leonidas, N. G. Oikonomakos and L. Somsák, *Tetrahedron: Asymmetry*, 2009, **20**, 733–740.
- 38 K. M. Alexacou, J. M. Hayes, C. Tiraidis, S. E. Zographos, D. D. Leonidas, E. D. Chrysina, G. Archontis, N. G. Oikonomakos, J. V. Paul, B. Varghese and D. Loganathan, *Proteins: Struct., Funct., Bioinf.*, 2008, **71**, 1307–1323.
- 39 É. Bokor, T. Docsa, P. Gergely and L. Somsák, *Bioorg. Med. Chem.*, 2009, submitted.
- 40 U. Aich and D. Loganathan, *J. Carbohydr. Chem.*, 2005, **24**, 1–12.
- 41 E. H. Fischer and E. G. Krebs, *Methods Enzymol.*, 1962, **5**, 369–372.
- 42 L. Somsák, L. Kovács, M. Tóth, E. Ósz, L. Szilágyi, Z. Györgydeák, Z. Dinya, T. Docsa, B. Tóth and P. Gergely, *J. Med. Chem.*, 2001, **44**, 2843–2848.
- 43 M. Benlifa, S. Vidal, B. Fenet, M. Msaddek, P. G. Goekjian, J.-P. Praly, A. Brunyánszki, T. Docsa and P. Gergely, *Eur. J. Org. Chem.*, 2006, 4242–4256.