



Published in final edited form as:

*Sci Signal*. ; 7(328): ra53. doi:10.1126/scisignal.2005051.

## The ABA Receptor PYL8 Promotes Lateral Root Growth by Enhancing MYB77-Dependent Transcription of Auxin-Responsive Genes

Yang Zhao<sup>#1,2</sup>, Lu Xing<sup>#2,3</sup>, Xingang Wang<sup>2</sup>, Yueh-Ju Hou<sup>2</sup>, Jinghui Gao<sup>2,4</sup>, Pengcheng Wang<sup>2</sup>, Cheng-Guo Duan<sup>2</sup>, Xiaohong Zhu<sup>2</sup>, and Jian-Kang Zhu<sup>1,2,†</sup>

<sup>1</sup>Shanghai Center for Plant Stress Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200032, China.

<sup>2</sup>Department of Horticulture and Landscape Architecture, Purdue University, West Lafayette, IN 47907, USA.

<sup>3</sup>School of Life Sciences, University of Science and Technology of China, Hefei 230026, China.

<sup>4</sup>College of Animal Science and Technology, Northwest A&F University, Yangling, Shaan'xi 712100, China.

# These authors contributed equally to this work.

### Abstract

The phytohormone abscisic acid (ABA) regulates plant growth, development, and abiotic stress responses. ABA signaling is mediated by a group of receptors known as the PYR1/PYL/RCAR family, which includes the pyrabactin resistance 1-like protein PYL8. Under stress conditions, ABA signaling activates SnRK2 protein kinases to inhibit lateral root growth after emergence from the primary root. However, even in the case of persistent stress, lateral root growth eventually recovers from inhibition. We showed that PYL8 is required for the recovery of lateral root growth, following inhibition by ABA. PYL8 directly interacted with the transcription factors MYB77, MYB44, and MYB73. The interaction of PYL8 and MYB77 increased the binding of MYB77 to its target MBSI motif in the promoters of multiple auxin-responsive genes. Compared to wild-type seedlings, the lateral root growth of *pyl8* mutant seedlings and *myb77* mutant seedlings was more sensitive to inhibition by ABA. The recovery of lateral root growth was delayed in *pyl8* mutant seedlings in the presence of ABA, and the defect was rescued by exposing *pyl8* mutant seedlings to the auxin IAA (3-indoleacetic acid). Thus, PYL8 promotes lateral root growth independently of the core ABA-SnRK2 signaling pathway by enhancing the activities of MYB77 and its paralogs, MYB44 and MYB73, to augment auxin signaling.

<sup>†</sup>Corresponding author. jkzhu@purdue.edu.

**Author contributions:** Y.Z., P.W., and J.-K.Z. designed the research; Y.Z., L.X., X.W., Y.-J.H., J.G., and C.-G.D. performed the experiments and analyzed the data; and Y.Z., X.Z., and J.-K.Z. wrote the manuscript.

**Competing interests:** The authors declare that they have no competing interests.

**Citation:** Y. Zhao, L. Xing, X. Wang, Y.-J. Hou, J. Gao, P. Wang, C.-G. Duan, X. Zhu, J.-K. Zhu, The ABA receptor PYL8 promotes lateral root growth by enhancing MYB77-dependent transcription of auxin-responsive genes. *Sci. Signal*. 7, ra53 (2014).

## INTRODUCTION

Abscisic acid (ABA) is an important phytohormone that regulates many growth and developmental processes, including seed longevity, dormancy and germination, vegetative growth, root architecture, and abiotic stress responses (1–4). ABA, together with auxin and other hormones, regulates root growth and architecture to acclimate to the variable growth environment, such as drought and salt stress, nutrient availability, gravity, and light (5–12). The root system of higher plants consists of primary, lateral, and adventitious roots. High concentrations of ABA inhibit both primary and lateral root growth (5, 9, 13, 14). Osmotic stress and salt stress inhibit primary and lateral root growth in a partially ABA-dependent manner (1, 6, 9, 15). During salt stress, both primary roots and lateral roots that have emerged from the primary root undergo a quiescent phase followed by a recovery phase (9, 15). ABA signaling functions in growth suppression during the quiescent phase of primary roots and lateral roots that have emerged from the primary root (post-emergence), although the quiescent phases for primary and lateral roots last for different periods and show different sensitivities to salt (9, 15). A low concentration of ABA is required for primary root elongation in response to water deficit and salt stress (15, 16). ABA is required for the growth recovery of primary root from inhibition and for long-term growth of primary roots during salt stress (15). In addition, the phytohormone gibberellic acid promotes growth recovery of both primary and lateral roots (9, 15).

In contrast to the inhibitory role of ABA on lateral root initiation and growth, auxin stimulates lateral root initiation and promotes lateral root growth (8, 12). The repression of lateral root formation under osmotic stress can be overcome by exogenous application of the synthetic auxin 1-naphthaleneacetic acid (6), implying that the balance between the repression by ABA and promotion by auxin may determine the fate of the lateral root primordium. As reported previously, ABA antagonizes auxin by promoting production of reactive oxygen species (ROS), leading to the inhibition of primary root growth (17–20). ABA also reduces auxin transport in roots by suppressing the expression of *PIN-FORMED 1* through *ABA-insensitive (ABI) 4*, resulting in suppression of lateral root formation and elongation (21). Furthermore, ABA greatly reduces the auxin reporter *ProDR5:GUS* expression in roots, indicating a reduction in auxin concentration or response (20, 22). In contrast, ABA can enhance auxin signaling without increasing the amount of auxin by activating auxin-responsive promoters to repress embryonic axis elongation (23). Similarly, auxin has both antagonistic and synergistic effects on responses to ABA. Auxin response factor (ARF) 2 directly suppresses the expression of *HOMEODOMAIN PROTEIN 33*, which is a positive regulator in ABA repression of primary root growth (22). However, auxin enhances ABA signaling through ARF10- and ARF16-mediated activation of *ABI3* expression in the regulation of lateral root formation and seed dormancy (24, 25). Plants with mutations in the *auxin resistant (AXR)* genes, *axr2-1* and *axr3-1*, and auxin transport mutants *aux1* and *pin2* are insensitive to the effects of ABA and auxin on embryonic axis elongation and root growth (23, 25, 26).

The core ABA signaling pathways consist of ABA-specific receptors and their downstream protein phosphatases and kinases. The pyrabactin resistance 1 (PYR1) and PYR1-like proteins (PYLs), also known as the regulatory component of ABA receptor (RCAR) family

proteins, are ABA receptors (27, 28). PYR1/PYL/RCARs interact with and inhibit clade-A protein phosphatase type 2Cs (PP2Cs) in the presence of ABA and release the SnRK2 protein kinases from inhibition by the PP2Cs (27, 28). Activated SnRK2s phosphorylate transcription factors such as ABA-responsive element binding factors (29), ABI5 (30), enhanced late embryogenesis abundant level (30), and flowering basic helix-loop-helix-type transcription factor 3 (30) to alter gene expression. Activated SnRK2s also phosphorylate ion channels such as slow anion channel 1 (31–36), K<sup>+</sup> channel in *Arabidopsis* 1 (37), and mechanosensitive channel of small conductance-like 9 (38) to stimulate stomatal closure. SnRK2s also phosphorylate the nicotinamide adenine dinucleotide phosphate oxidase RBOHF, leading to the production of ROS (17–19). In addition, several other substrates of SnRK2s include proteins related to RNA binding, microRNA and epi-genetic regulation, chloroplast function, and many other processes based on phosphoproteomics analyses (38, 39). Thus, PYR1/PYL/RCARs are thought to primarily regulate cellular functions using the PP2C-SnRK2 pathway.

Here, we found a previously uncharacterized function of PYL8 (also known as RCAR3) in the regulation of lateral root growth of *Arabidopsis thaliana* seedlings exposed to ABA treatment. PYL8 has a high affinity for ABA (40) and was recently shown to mediate the ability of ABA to inhibit primary root growth through the PP2C-SnRK2 pathway (7). We found that PYL8 mediates a synergistic action of ABA and auxin in promoting the growth recovery of post-emergence lateral roots. In addition, we found evidence for the function of the core ABA signaling components, PYR1/PYL/RCARs, PP2Cs, and SnRK2s, in the inhibition of lateral root growth. Our results suggest a signaling pathway in which the ABA receptor PYL8 directly interacts with a group MYB transcription factors to mediate crosstalk between ABA and auxin signaling to promote the growth recovery of lateral root from inhibition.

## RESULTS

### PYL8 promotes recovery from lateral root growth inhibition by ABA

ABA regulates the architecture of plant root systems during abiotic stress. To investigate how ABA controls root architecture, we analyzed the lateral root growth of *A. thaliana* plants with mutations in components of the ABA signaling pathway. To assess the role of PYL8 in ABA signaling, we analyzed two plants with transferred DNA (T-DNA) insertion mutations in *PYL8* (*pyl8-1* and *pyl8-2*) (Fig. 1A) that result in loss of function of PYL8 (7). We found that *pyl8* mutant seedlings showed more suppression of lateral root growth than did wild-type seedlings when exposed to ABA for 5 days (Fig. 1B). Seedlings of *pyl8* mutant seedlings grown on ABA-containing media for 30 days exhibited altered root architecture compared to wild-type seedlings (Fig. 1B, right panel). Lateral root growth was severely suppressed, whereas primary root growth was less inhibited, in *pyl8* mutant seedlings exposed to ABA (Fig. 1B, right panel). Both the total and average lateral root lengths of *pyl8* mutant seedlings were reduced compared to the wild-type seedlings when grown on ABA-containing media for 5 days (Fig. 1, C and D), suggesting that the lateral root growth of *pyl8* mutant seedlings is more sensitive to ABA. To confirm this observation, we examined lateral root growth with different concentrations of ABA. The lateral root

growth of *pyl8* mutant seedlings was more sensitive to ABA inhibition at all concentrations tested (Fig. 1, E and F). Thus, PYL8 promotes lateral root growth in response to ABA.

A previous study suggests that after being shifted to salt stress, post-emergence lateral roots undergo a quiescent phase followed by a recovery phase, and their growth is spatially and temporally regulated (9). To test whether the retarded lateral root growth of *pyl8* mutants exposed to ABA resulted from an extended quiescent phase, we monitored lateral roots for 5 days after emergence. Lateral roots that were less than 0.5 mm in length were considered quiescent, which is defined as stage C in the development of lateral root (5, 9). We found that the quiescent phase was extended in *pyl8* mutant compared to wild-type seedlings when grown on ABA-containing media, whereas the quiescent phase of *pyl8* mutant seedlings was comparable to that of wild-type seedlings when grown on control media (Fig. 1G). As a result, growth recovery was delayed in *pyl8* mutant compared to wild-type seedlings when exposed to ABA (Fig. 1H). Thus, inhibition of lateral root growth by ABA in *pyl8* mutant seedlings may be caused by prolonged quiescence, implying that PYL8 may function in promoting lateral root growth recovery in the presence of ABA.

Because salt or osmotic stress is known to induce ABA accumulation (41, 42), we analyzed the lateral and primary root growth of *pyl8* seedlings in media supplemented with various concentrations of NaCl or mannitol. However, we did not observe significant differences in lateral root growth between *pyl8* mutant and wild-type seedlings (fig. S1). Plant responses to these stressors are more complicated than responses to ABA because NaCl causes both osmotic stress and ion toxicity (43) and mannitol causes osmotic stress and other effects due to mannitol entry into cells (44). These differences may explain the differences in the response of lateral root growth in *pyl8* mutant plants to ABA and these other stressors.

### **PYLs, PP2Cs, and SnRK2s are required for lateral root growth suppression by ABA**

To test whether the lateral root phenotype of *pyl8* mutant seedlings is due to reduced response to ABA, we analyzed lateral root growth of seedlings with mutations in the core ABA signaling pathway, including *pyr1*, *pyl1*, *pyl2*, and *pyl4* (*pyr1pyl1/2/4*); *snrk2.2*, *snrk3*, and *snrk6* (*snrk2.2/3/6*); or *abi1-1*, each of which is resistant to ABA-induced inhibition of seed germination and seedling growth (13, 27, 45). Unlike *pyl8* mutant seedlings, the lateral root growth of *abi1-1*, *pyr1pyl1/2/4*, or *snrk2.2/3/6* mutant seedlings was less sensitive than that of wild-type seedlings to inhibition by ABA (Fig. 2, A to C). The lateral roots of wild-type and *abi1-1*, *pyr1pyl1/2/4*, or *snrk2.2/3/6* mutant seedlings had similar length of quiescent phases when grown in control media. However, the quiescent phase was shorter in *abi1-1*, *pyr1pyl1/2/4*, or *snrk2.2/3/6* mutant compared to wild-type seedlings when grown on ABA-containing media (Fig. 2D). Thus, the core ABA signaling pathway promotes lateral root quiescence in plants exposed to ABA.

### **Suppression of primary root growth and lateral root formation by ABA is reduced in *pyl8* and ABA-resistant mutants**

Primary and lateral roots use common and distinct molecular mechanisms to initiate and grow (5, 6, 9, 46, 47). We compared primary root elongation and the number of lateral roots in *pyl8* or ABA-resistant mutant seedlings. As reported previously (7, 13, 14, 48), the

primary root growth of *pyl8-1*, *abi1-1*, or *snrk2.2/3/6* mutant seedlings was less sensitive to ABA exposure compared to that of wild-type seedlings (Fig. 3, A and B). In wild-type seedlings, exposure to ABA reduced the number of lateral roots. In contrast, in *pyl8-1*, *abi1-1*, or *snrk2.2/3/6* mutant seedlings, the number of lateral roots was less affected by exposure to ABA (Fig. 3, C and D). We found that the expression of ABA-responsive genes was similar in the roots of *pyl8* mutant and wild-type plants exposed to ABA (fig. S2), suggesting that PYL8 has functional redundancy with other PYLs in mediating ABA-responsive gene expression. Thus, PYL8 likely plays a similar role to other PYLs in primary root growth and lateral root initiation.

We also assessed whether exposing wild-type seedlings to ABA could affect the expression of *PYL8*. We found that growing seedlings for 12 hours on ABA-containing media reduced the abundance of *PYL8* transcripts to about 33% of that in seedlings grown on control media (fig. S3). However, because ABA stabilizes PYL8 protein by reducing ubiquitination (49), the abundance of PYL8 protein may not substantially decrease in response to ABA.

### **Application of exogenous auxin promotes lateral root growth in *pyl8* mutant seedlings exposed to ABA**

Suppression of lateral root development by osmotic stress can be overcome by exogenous application of auxin (6). To test whether the inhibition of lateral root growth recovery in *pyl8* mutants could be rescued by auxin, we analyzed the lateral root growth of *pyl8* seedlings grown on ABA-containing media supplemented with different concentrations of the auxin IAA (3-indoleacetic acid). We found that 10 nM IAA was sufficient to rescue the inhibition of lateral root growth in *pyl8* mutant seedlings exposed to ABA for 5 days (Fig. 4, A to C). We also analyzed lateral root growth over time and found that wild-type and *pyl8* mutant seedlings showed similar lateral root growth on control media, and that lateral root elongation of *pyl8* mutant seedlings was suppressed by exposure to ABA (Fig. 4, D and E). However, in *pyl8* mutant seedlings exposed to both IAA and ABA, lateral root growth was enhanced compared to that of wild-type seedlings (Fig. 4, D and E).

To test whether IAA treatment alters the quiescent or recovery phases of lateral roots, we monitored the lateral roots of wild-type and *pyl8* mutant seedlings for 5 days. The *pyl8* mutant seedlings had a similar length of quiescent phase as wild-type seedlings in the presence of both IAA and ABA (Fig. 4F). The recovery of post-emergence lateral root growth in *pyl8* mutant seedlings exposed to ABA treatment was accelerated by exposure to IAA compared to exposure to ABA alone, resulting in comparable lengths of lateral roots to wild-type seedlings (Figs. 1H and 4G). Thus, the prolonged quiescent phase in lateral roots of *pyl8* mutant seedlings exposed to ABA was shortened by exposure to IAA, suggesting that the delayed recovery of lateral root growth may be due to auxin deficiency or reduced response to auxin.

### **PYL8 enhances MYB77 activity to promote lateral root growth**

Auxin signaling can activate the transcription factor MYB77 to promote lateral root growth (50). Recently, systematic yeast two-hybrid screens found MYB77 as an interacting protein of PYL8 (51). To test whether the interaction between PYL8 and MYB77 was specific, we

fused several PYLs to the GAL4 DNA binding domain (BD) and MYB77 or MYB61 to the GAL4-activating domain (AD) and performed yeast two-hybrid assays. We found that PYL8, but not other PYLs, interacted with MYB77, but not MYB61, in an ABA-independent manner (Fig. 5A). To further characterize the interaction between PYL8 and MYB77 in plant cells, we fused PYL8 to the N-terminal domain of firefly luciferase (LUC) (PYL8-nLUC) and MYB77 to the C-terminal domain of LUC (MYB77-cLUC) and co-expressed the two constructs in *A. thaliana* protoplasts. Coexpression of PYL8-nLUC, but not nLUC, with MYB77-cLUC produced measurable luciferase activity, comparable to coexpression of MYB77-cLUC with the positive control ABI1-nLUC (7) (Fig. 5B). To test whether the interaction between PYL8 and MYB77 was altered in the presence of ABA and auxin in vivo, we treated wild-type protoplasts coexpressing PYL8-nLUC and MYB77-cLUC with different combinations of ABA and IAA.

We found that LUC activity was enhanced in cells exposed to both ABA and IAA, but not in cells exposed to either ABA or IAA alone (Fig. 5C, left panel). To investigate the effect of endogenous ABA, we performed LUC complementation assays in protoplasts derived from the ABA-deficient mutant *aba2-1* (52) and found that ABA exposure promoted the interaction between PYL8 and MYB77 in the absence or presence of IAA (Fig. 5C, right panel). The interaction between PYL8 and MYB77 was not reduced by overexpression of ABI1 or ABI2 in *aba2-1* mutant protoplasts (fig. S4), suggesting that MYB77 and the PP2Cs may interact with different regions of PYL8. Thus, PYL8 specifically interacts with MYB77, and this interaction may be enhanced by ABA and IAA.

We asked whether the interaction between PYL8 and MYB77 affected the ability of MYB77 to bind to DNA and activate transcription. A previous study found that MYB77 $\Delta$ C1 (amino acids 1 to 200) recognizes several DNA motifs, including MBSI (CNGTTR) and MBSII (GTTAGTTA), and preferentially binds to MBSI motif-containing promoters (53). Using an electrophoretic mobility shift assay (EMSA), we found that full-length MYB77 bound to MBSI motif-containing DNA oligonucleotides (CRGTTA), but not to oligonucleotides containing M1 (CTGTTC), M2 (CAGTGA), or M3 (CCATTA) motifs (fig. S5, A and B). To test whether PYL8 affects the ability of MYB77 to bind to MBSI motifs, we preincubated recombinant MYB77 and PYL8 with or without ABA before performing the EMSA and found that PYL8 enhanced the binding of MYB77 to MBSI motifs in an ABA-independent manner (Fig. 5, D and E). PYL8 alone did not bind to MBSI motifs (fig. S5C). We identified putative MBSI and MBSII motifs in the 500-base pair (bp) promoter regions of multiple auxin-responsive genes, including *IAA1*, *IAA7*, *IAA17*, *IAA19*, *GH3.2*, *PIN1*, and *HAT2* (fig. S5, D and E). To test whether PYL8 can enhance MYB77 transcriptional activity, we fused a 439-bp fragment of *IAA19* promoter to the gene encoding LUC (*IAA19pro-LUC*, fig. S5D) and used it as an auxin-responsive reporter. We expressed *PYL8*, *MYB77*, *IAA19pro-LUC*, or combinations thereof in wild-type protoplasts. As expected, IAA, but not ABA, enhanced LUC activity in control protoplasts expressing only the reporter (Fig. 5F, left panel). However, co-overexpression of *PYL8* and *MYB77* increased LUC activity in the presence of IAA (Fig. 5F, left panel). To investigate the effects of endogenous ABA, we repeated the experiments in protoplasts from *aba2-1* mutant plants and found that ABA increased the ability of *PYL8* and *MYB77* overexpression to enhance

LUC activity in the presence of IAA (Fig. 5F, middle panel). Because ABA can inhibit auxin-responsive gene expression (20, 22), we performed reporter assays in protoplasts of *abil-1* mutant plants (54), which show inhibition of the core ABA signaling pathway (45), and found that cooverexpression of *PYL8* and *MYB77* increased LUC activity with or without ABA in the presence of IAA in these cells (Fig. 5F, right panel). Thus, *PYL8* enhances the binding of *MYB77* to its target motif MBSI, leading to increased response to auxin in an ABA-enhanced manner.

To further understand whether *MYB77* can activate auxin responses to control lateral root growth in response to ABA, we analyzed the lateral root growth of two plants with T-DNA insertion mutations in *MYB77* (50). We found that lateral root growth in *myb77* mutant seedlings was more sensitive to inhibition by ABA than that in wild-type seedlings and that exposure to IAA could reverse ABA-induced inhibition of lateral root growth in *myb77* mutant seedlings (Fig. 5, H and I). Thus, *MYB77*, like *PYL8*, promotes lateral root growth in seedlings exposed to ABA.

### **PYL8 interacts with paralogs of MYB77 and enhances their transcriptional activity**

We asked whether *PYL8* could bind paralogs of *MYB77* and modify their transcriptional activity. *MYB77*, *MYB73*, *MYB70*, and *MYB44* belong to the R2R3 MYB family subgroup 22, which is defined by two conserved motifs, TGLYMSPxSP and GxFMxVVQEMIxxEVRSYM (fig. S6) (55). *PYL8* is a putative interacting protein with *MYB44* (56). To test whether *MYB73* and *MYB44* could specifically bind *PYL8*, we performed yeast two-hybrid assays and found that *PYL8*, but not other *PYLs*, bound to both *MYB44* and *MYB73* in an ABA-independent manner (Fig. 6, A and B). *PYL8* showed a much stronger interaction with *MYB77* and *MYB44* than with *MYB73* (Figs. 5A and 6B). To test whether *MYB44* and *MYB73* could enhance the transcriptional response to auxin, we performed LUC assays in *IAA19pro-LUC*-expressing protoplasts from *aba2-1* and *abil-1* mutant plants. We found that co-overexpression with *PYL8* enhanced the ability of *MYB44* or *MYB73* to increase LUC activity in the presence of IAA (Fig. 6C and fig. S7) and that exposure to ABA enhanced the ability of *PYL8* to activate *MYB73* (Fig. 6C). Thus, *PYL8* likely binds to and enhances the activities of paralogs of *MYB77*, which may function redundantly with *MYB77* to enhance auxin signaling.

## **DISCUSSION**

The root system of plants is one of the most sensitive organs in sensing the availability of water and nutrients or other adverse soil conditions. As a critical part of plant responses to environmental cues, phytohormone signaling finely controls the growth and development of roots for acclimating to environmental changes. Here, we found a synergistic action of ABA and auxin in controlling the growth of plant lateral roots. Our study reveals that ABA promotes lateral root growth recovery through a pathway mediated by a previously uncharacterized action of ABA receptor *PYL8*. *PYL8* directly interacts with a group of transcription factors, *MYB77*, *MYB44*, and *MYB73*, leading to the enhancement of auxin-dependent transcription. Together with evidence from previous studies (5, 6, 9), our results suggest that ABA has dual roles in controlling lateral root growth via two different pathways

(Fig. 7). In this model, ABA signaling inhibits lateral root growth during the quiescent phase and promotes lateral root growth during the recovery phase of post-emergence lateral root growth during abiotic stress.

The growth of post-emergence lateral roots under salt stress undergoes a quiescent phase followed by a recovery phase. Endogenous ABA stimulated by salt stress promotes lateral root quiescence, and application of exogenous ABA treatment mimics this effect (9). Therefore, together with our results, it is likely that lateral root growth undergoes quiescent and recovery phases when exposed to exogenous ABA. We found that lateral root growth was less sensitive to ABA in ABA-resistant mutant seedlings including *pyr1pyl1/2/4*, *abi1-1*, and *snrk2.2/3/6* (Fig. 2, A to C). Consistent with this result, lateral root growth is less sensitive to salt stress in transgenic seedlings expressing *ABI1-1* in the root endodermis (9). The *ABI4* gene encodes an ABA-regulated AP2 domain transcription factor, and lateral root growth of *abi4* mutant seedlings is insensitive to ABA (21). Similarly, the double mutant seedlings *abi4-1* and *fus3-3* are less sensitive to salt stress (9, 21), suggesting that *ABI4* may be the downstream regulator in controlling lateral root quiescence during exposure to ABA or salt stress. Thus, lateral root quiescence is likely controlled by core ABA signaling through the PYL receptors, PP2C co-receptors, SnRK2 protein kinases, ABI4 transcription factor, and downstream ABA-responsive genes, and our results suggest that PYL8-mediated lateral root growth recovery is independent of the ABA core signaling pathway. Instead, PYL8 interacts directly with the MYB77 paralogs to enhance auxin-dependent gene expression.

In support of the proposed model, our data showed that the recovery of post-emergence lateral root growth was delayed in *pyl8* mutant seedlings exposed to ABA and that the application of exogenous IAA accelerated the lateral root growth of *pyl8* mutant seedlings exposed to ABA (Figs. 1 and 4). The direct interaction of PYL8 with the MYB transcription factors bridges ABA and auxin signaling in controlling lateral root growth. We showed that ABA enhances the effects of PYL8 on the activities of MYB73 and MYB77 in the presence of IAA (Figs. 5F and 6C), suggesting a synergistic action of ABA and IAA on MYB44, MYB73, and MYB77 activities through their interactions with PYL8. It is known that MYB77 positively regulates auxin signaling to promote lateral root formation (50). MYB77 forms a heterodimer with ARF7 and increases the transcription of ARF7 target genes (50), as indicated in our model (Fig. 7). In addition, both ARF7 and ARF19 increase the expression of early auxin response genes such as *LBD16* (also known as *ASL18*) and *LBD29* (also known as *ASL16*) in lateral roots, promoting lateral root formation and elongation (57–59).

ABA signaling can enhance auxin-dependent responses by activating an auxin-responsive promoter (23). Truncated MYB77 and MYB44 proteins bind to the motifs of MBSI and MBSII (53, 60) that were identified in the 500-bp fragments of the promoters of multiple auxin-responsive genes (fig. S5, D and E). Auxin-responsive genes were highly induced by salt stress in plants overexpressing *MYB44* (61), suggesting that salt stress or ABA can enhance auxin signaling when MYB44 abundance is high. *MYB73* and *MYB77* are expressed in lateral roots (50, 62), whereas *PYL8* is expressed extensively in roots (7). The spatial expression patterns of MYBs and PYL8 indicate that the interaction of PYL8 and MYBs



may occur specifically in roots. In addition, we observed that the inhibition of lateral root growth in *myb77* mutant seedlings was not as severe as that in *pyl8* mutant seedlings when exposed to ABA, suggesting that MYB44 and MYB73 may be functionally redundant with MYB77. In protoplasts, ABA enhanced the ability of IAA and PYL8 overexpression to increase the ability of MYB73 and MYB77 to activate an auxin reporter gene (Figs. 5F and 6C). This result implies that the synergistic action of ABA and IAA can be mediated by PYL8-MYB73 and PYL8-MYB77. Furthermore, the expression of *MYB44*, *MYB73*, and *MYB77* is strongly induced by dehydration, salt, cold, ABA, or auxin (50, 61, 63).

PYL8 has been shown to mediate ABA inhibition of primary root growth (7). This role is likely mediated through the PP2C-SnRK2 pathway because primary root elongation in *abi1-1* mutant seedlings and *snrk2.2/3/6* mutant seedlings was not sensitive to ABA exposure (Fig. 3, A and B). Our results show that PYL8 also has a role in promoting lateral root elongation exposed to ABA, and suggest that this function is mediated through the previously uncharacterized PYL8-MYB77 pathway. Therefore, PYL8 has different roles mediated through distinct pathways in the ABA-dependent regulation of different aspects of root growth and development.

## MATERIALS AND METHODS

### Plant materials

The *pyl8-1* (SAIL\_1269\_A02) (7, 14), *pyl8-2* (SALK\_033867) (7), *pyr1pyl1/2/4* quadruple mutant (27), *snrk2.2/3/6* triple mutant (13), *myb77-1* (50), and *myb77-2* (50) were in Col-0 background, except *pyl2* (27) and *abi1-1* (45), which were in the Landsberg erecta (Ler) background. The *pyl8* and *myb77* mutants were obtained from the Arabidopsis Biological Resource Center (ABRC).

### Plant growth conditions

Seeds were surface-sterilized for 10 min in 20% bleach and then rinsed in sterile deionized water four times. Sterilized seeds were grown on 0.6% Phytigel (Sigma) media containing 1/2 MS nutrients or 1/4 MS nutrients as indicated (catalog no. M524, PhytoTechnology Laboratories), 1% sucrose adjusted to pH 5.7 (control media), and kept at 4° to 8°C for 3 days. Seedlings were grown vertically for 3 days before transfer to control media or media with the indicated concentrations of ABA (Sigma, A1049) or IAA (Sigma, I2886). Seedlings were grown in a Percival CU36L5 incubator at 23°C under a 16-hour light, 8-hour dark photoperiod. Plates were sealed with micropore tape (3M).

For protoplast analysis, seedlings were grown on Jiffy 7 peat soil (42mm Pellets) in a Percival chamber with a relatively short photoperiod (12 hours of light at 23°C, 12 hours of dark at 20°C) under low light (about 100  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) and 50 to 70% relative humidity under well-watered conditions (no drought and no flooding). Protoplasts were isolated as described (64, 65).

## Phenotype analysis

The length of all lateral roots was quantified using a Leica EZ4HD stereo microscope. The total and average lateral root lengths of individual plants were quantified by summing or averaging the lengths of all the lateral roots on each plant. Lateral roots shorter than 0.5 mm were categorized as quiescent (9).

## Plasmid constructs

All primer sequences are shown in table S1. To generate *IAA19pro-LUC* construct, the 439-bp *IAA19* promoter fragment amplified from Col-0 genomic DNA with primers IAA19proF and IAA19proR was cloned into the Bam HI and Nco I of the *RD29B-LUC* vector (65). To generate pET28a-PYL8 construct, the coding sequence of *PYL8* was amplified from *His-PYL8* (65) using *PYL8* primers and cloned into pET28a vectors. To generate pENTR-MYB77 construct, *MYB77* CDS was amplified using pENTR-MYB77 primers and cloned into pENTR vector using the pENTR/D-TOPO Cloning Kit (Invitrogen). *MYB77* was recombined into the pEarley-nLUC and pEarley-cLUC binary vectors using an LR reaction (Invitrogen). *MYB44*, *MYB73*, and *MYB77* were cloned into pHBT95 using transfer polymerase chain reaction with pHBT-MYBs primers, cloned into pGADT7 with AD-MYBs primers, and cloned into pGEX-6P1 with 6P1-MYB77 primers. *ABI1*, *ABI2*, BD-PYLs, and *PYL8-nLUC* were the same as reported (64). *ABI1* was cloned into pHBT95-cLUC using cLUC-ABI1 primers. *His-PYL8* was the same as reported (65). *ZmUBQ::GUS* was provided by J. Sheen.

## Yeast two-hybrid assay

Yeast two-hybrid assays were performed as described (66). pGBKT7-PYLs were the same as reported (64). PYLs fused to the GAL4 DNA binding domain were used as baits. *MYB44*, *MYB73*, *MYB77*, and *MYB61* fused to the GAL4-activating domain were used as preys. Interaction was determined by growth assay on media lacking His or His and Ade in the presence and absence of 10  $\mu$ M ABA. Dilutions ( $10^{-1}$ ,  $10^{-2}$ , and  $10^{-3}$ ) of saturated cultures were spotted onto the plates and photographed after 5 days.

## Transient expression assay in *Arabidopsis*

Assays for transient expression in protoplasts were performed as described (64). All steps were kept at room temperature. *IAA19pro-LUC* was used as the auxin-responsive reporter. *ZmUBQ-GUS* was used as the internal control. After transfection, protoplasts were incubated in washing and incubation solution (0.5 M mannitol, 20 mM KCl, 4 mM MES, pH 5.7) with or without ABA and IAA at the indicated concentrations with 12 hours of light. LUC complementation assays were performed as described (64).

## Electrophoretic mobility shift assay

GST-MYB77 and His-PYL8 were purified as described (67). EMSA assay was performed as described (64, 68). The radiolabeled DNA probes were labeled with [ $\gamma$ - $^{32}$ P]adenosine 5'-triphosphate using T<sub>4</sub> polynucleotide kinase and purified using Micro Bio-Spin 6 columns (Bio-Rad, 732-6221). EMSAs were performed with radiolabeled oligonucleotides (0.6 nM) and recombinant GST-MYB77 protein (17 nM) with or without the indicated unlabeled

oligonucleotides (40 nM). Recombinant His-PYL8 protein (17 nM) was preincubated with GST-MYB77 (17 nM) for 10 min with or without ABA (0.2 mM) on ice before the radiolabeled oligo-nucleotide was added. The DNA binding reactions were performed in binding buffer containing 10 mM Hepes-KOH (pH 7.5), 0.1 mM EDTA, 75 mM KCl, 1.25 mM MgCl<sub>2</sub>, 0.2 mM dithiothreitol, bovine serum albumin (2 µg/ml), and 5% glycerol. The reaction was incubated with the radio-labeled oligonucleotide for 30 min on ice and then loaded onto 5% polyacrylamide native gels, which had been pre-run at 180 V in 0.5× tris-borate EDTA buffer for 30 min at 4°C, and run at 180 V for 35 min at 4°C. After electrophoresis, the gel was dried under vacuum at 80°C for 15 min on a filter paper and exposed to a phosphorimager for 2 to 10 hours.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank J. Sheen for the *ZmUBQ::GUS* construct and S. R. Cutler for the *pyr1pyl1/2/4* quadruple mutant.

**Funding:** This work was supported by NIH grant R01GM059138 and by the Chinese Academy of Sciences.

## REFERENCES AND NOTES


- Zhu JK. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* 2002; 53:247–273. [PubMed: 12221975]
- Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. Abscisic acid: Emergence of a core signaling network. *Annu. Rev. Plant Biol.* 2010; 61:651–679. [PubMed: 20192755]
- Raghavendra AS, Gonugunta VK, Christmann A, Grill E. ABA perception and signalling. *Trends Plant Sci.* 2010; 15:395–401. [PubMed: 20493758]
- Ooms J, Leon-Kloosterziel KM, Bartels D, Koornneef M, Karssen CM. Acquisition of desiccation tolerance and longevity in seeds of *Arabidopsis thaliana* (a comparative study using abscisic acid-insensitive *abi3* mutants). *Plant Physiol.* 1993; 102:1185–1191. [PubMed: 12231895]
- De Smet I, Signora L, Beeckman T, Inzé D, Foyer CH, Zhang H. An abscisic acid-sensitive checkpoint in lateral root development of *Arabidopsis*. *Plant J.* 2003; 33:543–555. [PubMed: 12581312]
- Deak KI, Malamy J. Osmotic regulation of root system architecture. *Plant J.* 2005; 43:17–28. [PubMed: 15960613]
- Antoni R, Gonzalez-Guzman M, Rodriguez L, Peirats-Llobet M, Pizzio GA, Fernandez MA, De Winne N, De Jaeger G, Dietrich D, Bennett MJ, Rodriguez PL. PYRABACTIN RESISTANCE1-LIKE8 plays an important role for the regulation of abscisic acid signaling in root. *Plant Physiol.* 2013; 161:931–941. [PubMed: 23370718]
- Lavenus J, Goh T, Roberts I, Guyomarc'h S, Lucas M, De Smet I, Fukaki H, Beeckman T, Bennett M, Laplaze L. Lateral root development in *Arabidopsis*: Fifty shades of auxin. *Trends Plant Sci.* 2013; 18:450–458. [PubMed: 23701908]
- Duan L, Dietrich D, Ng CH, Chan PM, Bhalerao R, Bennett MJ, Dinneny JR. Endodermal ABA signaling promotes lateral root quiescence during salt stress in *Arabidopsis* seedlings. *Plant Cell.* 2013; 25:324–341. [PubMed: 23341337]
- Overvoorde P, Fukaki H, Beeckman T. Auxin control of root development. *Cold Spring Harb. Perspect. Biol.* 2010; 2:a001537.
- Uga Y, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Kitomi Y, Inukai Y, Ono K, Kanno N, Inoue H, Takehisa H, Motoyama R, Nagamura Y, Wu J, Matsumoto T, Takai T, Okuno K, Yano M. Control of root system architecture by *DEEPER ROOTING 1* increases rice yield under drought conditions. *Nat. Genet.* 2013; 45:1097–1102. [PubMed: 23913002]

12. Smith S, De Smet I. Root system architecture: Insights from *Arabidopsis* and cereal crops. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 2012; 367:1441–1452. [PubMed: 22527386]
13. Fujii H, Zhu JK. *Arabidopsis* mutant deficient in 3 abscisic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:8380–8385. [PubMed: 19420218]
14. Gonzalez-Guzman M, Pizzio GA, Antoni R, Vera-Sirera F, Merilo E, Bassel GW, Fernández MA, Holdsworth MJ, Perez-Amador MA, Kollist H, Rodriguez PL. *Arabidopsis* PYR/PYL/RCAR receptors play a major role in quantitative regulation of stomatal aperture and transcriptional response to abscisic acid. *Plant Cell.* 2012; 24:2483–2496. [PubMed: 22739828]
15. Geng Y, Wu R, Wee CW, Xie F, Wei X, Chan PM, Tham C, Duan L, Dinneny JR. A spatio-temporal understanding of growth regulation during the salt stress response in *Arabidopsis*. *Plant Cell.* 2013; 25:2132–2154. [PubMed: 23898029]
16. Sharp RE, Poroyko V, Hejlek LG, Spollen WG, Springer GK, Bohnert HJ, Nguyen HT. Root growth maintenance during water deficits: Physiology to functional genomics. *J. Exp. Bot.* 2004; 55:2343–2351. [PubMed: 15448181]
17. Mustilli AC, Merlot S, Vavasseur A, Fenzi F, Giraudat J. *Arabidopsis* OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production. *Plant Cell.* 2002; 14:3089–3099. [PubMed: 12468729]
18. Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JD, Schroeder JI. NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J.* 2003; 22:2623–2633. [PubMed: 12773379]
19. Sirichandra C, Gu D, Hu HC, Davanture M, Lee S, Djaoui M, Valot B, Zivy M, Leung J, Merlot S, Kwak JM. Phosphorylation of the *Arabidopsis* AtrbohF NADPH oxidase by OST1 protein kinase. *FEBS Lett.* 2009; 583:2982–2986. [PubMed: 19716822]
20. He J, Duan Y, Hua D, Fan G, Wang L, Liu Y, Chen Z, Han L, Qu LJ, Gong Z. DEXH box RNA helicase-mediated mitochondrial reactive oxygen species production in *Arabidopsis* mediates crosstalk between abscisic acid and auxin signaling. *Plant Cell.* 2012; 24:1815–1833. [PubMed: 22652060]
21. Shkolnik-Inbar D, Bar-Zvi D. ABI4 mediates abscisic acid and cytokinin inhibition of lateral root formation by reducing polar auxin transport in *Arabidopsis*. *Plant Cell.* 2010; 22:3560–3573. [PubMed: 21097710]
22. Wang L, Hua D, He J, Duan Y, Chen Z, Hong X, Gong Z. *Auxin Response Factor2 (ARF2)* and its regulated homeodomain gene *HB33* mediate abscisic acid response in *Arabidopsis*. *PLOS Genet.* 2011; 7:e1002172. [PubMed: 21779177]
23. Belin C, Megies C, Hauserová E, Lopez-Molina L. Abscisic acid represses growth of the *Arabidopsis* embryonic axis after germination by enhancing auxin signaling. *Plant Cell.* 2009; 21:2253–2268. [PubMed: 19666738]
24. Brady SM, Sarkar SF, Bonetta D, McCourt P. The *ABSCISIC ACID INSENSITIVE 3 (ABI3)* gene is modulated by farnesylation and is involved in auxin signaling and lateral root development in *Arabidopsis*. *Plant J.* 2003; 34:67–75. [PubMed: 12662310]
25. Liu X, Zhang H, Zhao Y, Feng Z, Li Q, Yang HQ, Luan S, Li J, He ZH. Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 2013; 110:15485–15490. [PubMed: 23986496]
26. Wilson AK, Pickett FB, Turner JC, Estelle M. A dominant mutation in *Arabidopsis* confers resistance to auxin, ethylene and abscisic acid. *Mol. Gen. Genet.* 1990; 222:377–383. [PubMed: 2148800]
27. Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez PL, McCourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR. Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science.* 2009; 324:1068–1071. [PubMed: 19407142]

28. Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E. Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science*. 2009; 324:1064–1068. [PubMed: 19407143]
29. Fujii H, Verslues PE, Zhu JK. Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in *Arabidopsis*. *Plant Cell*. 2007; 19:485–494. [PubMed: 17307925]
30. Nakashima K, Fujita Y, Kanamori N, Katagiri T, Umezawa T, Kidokoro S, Maruyama K, Yoshida T, Ishiyama K, Kobayashi M, Shinozaki K, Yamaguchi-Shinozaki K. Three *Arabidopsis* SnRK2 protein kinases, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, involved in ABA signaling are essential for the control of seed development and dormancy. *Plant Cell Physiol*. 2009; 50:1345–1363. [PubMed: 19541597]
31. Brandt B, Brodsky DE, Xue S, Negi J, Iba K, Kangasjärvi J, Ghassemian M, Stephan AB, Hu H, Schroeder JI. Reconstitution of abscisic acid activation of SLAC1 anion channel by CPK6 and OST1 kinases and branched ABI1 PP2C phosphatase action. *Proc. Natl. Acad. Sci. U.S.A.* 2012; 109:10593–10598. [PubMed: 22689970]
32. Geiger D, Maierhofer T, Al-Rasheid KA, Scherzer S, Mumm P, Liese A, Ache P, Wellmann C, Marten I, Grill E, Romeis T, Hedrich R. Stomatal closure by fast abscisic acid signaling is mediated by the guard cell anion channel SLAH3 and the receptor RCAR1. *Sci. Signal*. 2011; 4:ra32. [PubMed: 21586729]
33. Vahisalu T, Puzõrjova I, Brosché M, Valk E, Lepiku M, Moldau H, Pechter P, Wang YS, Lindgren O, Salojärvi J, Loog M, Kangasjärvi J, Kollist H. Ozone-triggered rapid stomatal response involves the production of reactive oxygen species, and is controlled by SLAC1 and OST1. *Plant J*. 2010; 62:442–453. [PubMed: 20128877]
34. Geiger D, Scherzer S, Mumm P, Marten I, Ache P, Matschi S, Liese A, Wellmann C, Al-Rasheid KA, Grill E, Romeis T, Hedrich R. Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca<sup>2+</sup> affinities. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:8023–8028. [PubMed: 20385816]
35. Lee SC, Lan W, Buchanan BB, Luan S. A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:21419–21424. [PubMed: 19955427]
36. Geiger D, Scherzer S, Mumm P, Stange A, Marten I, Bauer H, Ache P, Matschi S, Liese A, Al-Rasheid KA, Romeis T, Hedrich R. Activity of guard cell anion channel SLAC1 is controlled by drought-stress signaling kinase-phosphatase pair. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:21425–21430. [PubMed: 19955405]
37. Sato A, Sato Y, Fukao Y, Fujiwara M, Umezawa T, Shinozaki K, Hibi T, Taniguchi M, Miyake H, Goto DB, Uozumi N. Threonine at position 306 of the KAT1 potassium channel is essential for channel activity and is a target site for ABA-activated SnRK2/OST1/SnRK2.6 protein kinase. *Biochem. J*. 2009; 424:439–448. [PubMed: 19785574]
38. Wang P, Xue L, Batelli G, Lee S, Hou YJ, Van Oosten MJ, Zhang H, Tao WA, Zhu JK. Quantitative phosphoproteomics identifies SnRK2 protein kinase substrates and reveals the effectors of abscisic acid action. *Proc. Natl. Acad. Sci. U.S.A.* 2013; 110:11205–11210. [PubMed: 23776212]
39. Umezawa T, Sugiyama N, Takahashi F, Anderson JC, Ishihama Y, Peck SC, Shinozaki K. Genetics and phosphoproteomics reveal a protein phosphorylation network in the abscisic acid signaling pathway in *Arabidopsis thaliana*. *Sci. Signal*. 2013; 6:rs8. [PubMed: 23572148]
40. Szostkiewicz I, Richter K, Kepka M, Demmel S, Ma Y, Korte A, Assaad FF, Christmann A, Grill E. Closely related receptor complexes differ in their ABA selectivity and sensitivity. *Plant J*. 2010; 61:25–35. [PubMed: 19769575]
41. Pierce M, Raschke K. Correlation between loss of turgor and accumulation of abscisic acid in detached leaves. *Planta*. 1980; 148:174–182. [PubMed: 24309706]
42. Jia W, Wang Y, Zhang S, Zhang J. Salt-stress-induced ABA accumulation is more sensitively triggered in roots than in shoots. *J. Exp. Bot*. 2002; 53:2201–2206. [PubMed: 12379787]
43. Niu X, Bressan RA, Hasegawa PM, Pardo JM. Ion homeostasis in NaCl stress environments. *Plant Physiol*. 1995; 109:735–742. [PubMed: 12228628]

44. Tholalakabavi A, Zwiazek J, Thorpe T. Effect of mannitol and glucose-induced osmotic stress on growth, water relations, and solute composition of cell suspension cultures of poplar (*Populus deltoides* var. *Occidentalis*) in relation to anthocyanin accumulation. *In Vitro Cell Dev. Biol.* 1994; 30:164–170.
45. Gosti F, Beaudoin N, Serizet C, Webb AA, Vartanian N, Giraudat J. ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. *Plant Cell.* 1999; 11:1897–1910. [PubMed: 10521520]
46. Roychoudhry S, Del Bianco M, Kieffer M, Kepinski S. Auxin controls gravitropic setpoint angle in higher plant lateral branches. *Curr. Biol.* 2013; 23:1497–1504. [PubMed: 23891109]
47. Rosquete MR, von Wangenheim D, Marhavý P, Barbez E, Stelzer EH, Benková E, Maizel A, Kleine-Vehn J. An auxin transport mechanism restricts positive orthogravitropism in lateral roots. *Curr. Biol.* 2013; 23:817–822. [PubMed: 23583551]
48. Leung J, Merlot S, Giraudat J. The *Arabidopsis* *ABSCISIC ACID-INSENSITIVE2 (ABI2)* and *ABI1* genes encode homologous protein phosphatases 2C involved in abscisic acid signal transduction. *Plant Cell.* 1997; 9:759–771. [PubMed: 9165752]
49. Irigoyen ML, Iniesto E, Rodriguez L, Puga MI, Yanagawa Y, Pick E, Strickland E, Paz-Ares J, Wei N, De Jaeger G, Rodriguez PL, Deng XW, Rubio V. Targeted degradation of abscisic acid receptors is mediated by the ubiquitin ligase substrate adaptor DDA1 in *Arabidopsis*. *Plant Cell.* 2014; 26:712–728. [PubMed: 24563205]
50. Shin R, Burch AY, Huppert KA, Tiwari SB, Murphy AS, Guilfoyle TJ, Schachtman DP. The *Arabidopsis* transcription factor MYB77 modulates auxin signal transduction. *Plant Cell.* 2007; 19:2440–2453. [PubMed: 17675404]
51. Arabidopsis Interactome Mapping Consortium. Evidence for network evolution in an *Arabidopsis* interactome map. *Science.* 2011; 333:601–607. [PubMed: 21798944]
52. González-Guzmán M, Apostolova N, Bellés JM, Barrero JM, Piqueras P, Ponce MR, Micol JL, Serrano R, Rodríguez PL. The short-chain alcohol dehydrogenase ABA2 catalyzes the conversion of xanthoxin to abscisic aldehyde. *Plant Cell.* 2002; 14:1833–1846. [PubMed: 12172025]
53. Romero I, Fuertes A, Benito MJ, Malpica JM, Leyva A, Paz-Ares J. More than 80 *R2R3-MYB* regulatory genes in the genome of *Arabidopsis thaliana*. *Plant J.* 1998; 14:273–284. [PubMed: 9628022]
54. Koornneef M, Reuling G, Karssen CM. The isolation and characterization of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiol. Plant.* 1984; 61:377–383.
55. Stracke R, Werber M, Weisshaar B. The R2R3-MYB gene family in *Arabidopsis thaliana*. *Curr. Opin. Plant Biol.* 2001; 4:447–456. [PubMed: 11597504]
56. Jaradat MR, Feurtado JA, Huang D, Lu Y, Cutler AJ. Multiple roles of the transcription factor AtMYBR1/AtMYB44 in ABA signaling, stress responses, and leaf senescence. *BMC Plant Biol.* 2013; 13:192. [PubMed: 24286353]
57. Wilmoth JC, Wang S, Tiwari SB, Joshi AD, Hagen G, Guilfoyle TJ, Alonso JM, Ecker JR, Reed JW. NPH4/ARF7 and ARF19 promote leaf expansion and auxin-induced lateral root formation. *Plant J.* 2005; 43:118–130. [PubMed: 15960621]
58. Okushima Y, Overvoorde PJ, Arima K, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Lui A, Nguyen D, Onodera C, Quach H, Smith A, Yu G, Theologis A. Functional genomic analysis of the *AUXIN RESPONSE FACTOR* gene family members in *Arabidopsis thaliana*: Unique and overlapping functions of ARF7 and ARF19. *Plant Cell.* 2005; 17:444–463. [PubMed: 15659631]
59. Okushima Y, Fukaki H, Onoda M, Theologis A, Tasaka M. ARF7 and ARF19 regulate lateral root formation via direct activation of *LBD/ASL* genes in *Arabidopsis*. *Plant Cell.* 2007; 19:118–130. [PubMed: 17259263]
60. Jung C, Kim YK, Oh NI, Shim JS, Seo JS, Choi YD, Nahm BH, Cheong JJ. Quadruple 9-mer-based protein binding microarray analysis confirms AACnG as the consensus nucleotide sequence sufficient for the specific binding of AtMYB44. *Mol. Cells.* 2012; 34:531–537. [PubMed: 23161171]
61. Jung C, Seo JS, Han SW, Koo YJ, Kim CH, Song SI, Nahm BH, Choi YD, Cheong JJ. Overexpression of *AtMYB44* enhances stomatal closure to confer abiotic stress tolerance in transgenic *Arabidopsis*. *Plant Physiol.* 2008; 146:623–635. [PubMed: 18162593]

62. Kim JH, Nguyen NH, Jeong CY, Nguyen NT, Hong SW, Lee H. Loss of the R2R3 MYB, AtMyb73, causes hyper-induction of the *SOS1* and *SOS3* genes in response to high salinity in *Arabidopsis*. *J. Plant Physiol.* 2013; 170:1461–1465. [PubMed: 23809151]
63. Fowler S, Thomashow MF. *Arabidopsis* transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell.* 2002; 14:1675–1690. [PubMed: 12172015]
64. Zhao Y, Chan Z, Xing L, Liu X, Hou YJ, Chinnusamy V, Wang P, Duan C, Zhu JK. The unique mode of action of a divergent member of the ABA-receptor protein family in ABA and stress signaling. *Cell Res.* 2013; 23:1380–1395. [PubMed: 24189045]
65. Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, Cutler SR, Sheen J, Rodriguez PL, Zhu JK. In vitro reconstitution of an abscisic acid signalling pathway. *Nature.* 2009; 462:660–664. [PubMed: 19924127]
66. Bai G, Yang DH, Zhao Y, Ha S, Yang F, Ma J, Gao XS, Wang ZM, Zhu JK. Interactions between soybean ABA receptors and type 2C protein phosphatases. *Plant Mol. Biol.* 2013; 83:651–664. [PubMed: 23934343]
67. Zhao Y, Zhao S, Mao T, Qu X, Cao W, Zhang L, Zhang W, He L, Li S, Ren S, Zhao J, Zhu G, Huang S, Ye K, Yuan M, Guo Y. The plant-specific actin binding protein SCAB1 stabilizes actin filaments and regulates stomatal movement in *Arabidopsis*. *Plant Cell.* 2011; 23:2314–2330. [PubMed: 21719691]
68. Zhao J, Zhang W, Zhao Y, Gong X, Guo L, Zhu G, Wang X, Gong Z, Schumaker KS, Guo Y. SAD2, an importin  $\beta$ -like protein, is required for UV-B response in *Arabidopsis* by mediating MYB4 nuclear trafficking. *Plant Cell.* 2007; 19:3805–3818. [PubMed: 17993626]




**Fig. 1. *pyl8* mutant seedlings have enhanced sensitivity to ABA-induced inhibition of lateral root growth**

(A) Diagram of T-DNA insertions in the *PYL8* gene. The T-DNA insertions in the two SALK lines are inserted in exons, which are presented as closed boxes. (B) The root architecture of wild-type (Col-0) or *pyl8* mutant seedlings was documented at 5 dpt (days post-transfer) (left panel) or 30 dpt (right panel) to media (1/2 MS, 1% sucrose) with or without ABA. The primary root tips are marked by white asterisks. (C and D) Lateral root (LR) length of Col-0 and *pyl8* mutant seedlings at 5 dpt to media with or without ABA.  $n \geq 12$  seedlings. Data are means  $\pm$  SEM.  $***P < 0.001$ , Student's *t* test. (E and F) Lateral root length of Col-0 and *pyl8-1* mutant seedlings at 5 dpt to media with varying concentrations of ABA.  $n = 9$  seedlings. Data are means  $\pm$  SEM.  $**P < 0.01$ ,  $***P < 0.001$ , Student's *t* test. (G) Pie charts showing the percentage of quiescent lateral roots for the indicated number of days for seedlings grown in media with or without ABA media.  $n = 10$  to 24 seedlings per condition. (H) Percentage of quiescent lateral roots (<0.5 mm at 0 dpt) that exited quiescence within 5 dpt to the media with or without ABA. Percentage was calculated as




total lateral roots exiting quiescence (>0.5 mm) at the indicated days divided by total number of quiescent lateral roots at 0 dpt.  $n = 10$  to 24 seedlings per condition. Data are means  $\pm$  SEM.  $**P < 0.01$ ,  $***P < 0.001$ , analysis of variance (ANOVA).




**Fig. 2. Lateral root growth is not suppressed by exposure to ABA in ABA-resistant mutant seedlings**

(A) The root architecture of wild-type (Col-0 or Ler) or *abi1-1*, *pyr1pyl1/2/4*, or *snrk2.2/3/6* mutant seedlings was documented at 3 dpt to media (1/2 MS, 1% sucrose) with or without ABA. (B and C) Lateral root (LR) length of seedlings at 3 dpt to media with varying concentrations of ABA.  $n \geq 11$  seedlings. Data are means  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , Student's  $t$  test. (D) Pie charts for the percentage of quiescent lateral roots for the indicated number of days on seedlings grown in media with or without ABA.  $n = 10$  to 12 seedlings per condition.




**Fig. 3. Suppression of primary root growth and lateral root formation by exposure to ABA is reduced in *pyl8* and ABA-resistant mutant seedlings**

(A and C) The primary root length and lateral root number of wild-type (Col-0 or Ler) or *abi1-1* or *snrk2.2/3/6* mutant seedlings were documented at 6 dpt to media (1/2 MS, 1% sucrose) with varying concentrations of ABA.  $n \geq 10$  seedlings. Data are means  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , Student's  $t$  test. (B and D) The primary root length and lateral root number of Col-0 and *pyl8-1* mutant were documented at 7 dpt to media with varying concentrations of ABA.  $n = 9$  seedlings. Data are means  $\pm$  SEM. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , Student's  $t$  test.



**Fig. 4. Exogenous IAA rescues the lateral root growth defects of *pyl8* mutant seedlings**  
 (A) The root architecture of wild-type (*Col-0*) or *pyl8* mutant seedlings was documented at 5 dpt to media (1/2 MS, 1% sucrose) with or without ABA or ABA plus IAA. (B and C) Lateral root (LR) length of seedlings at 5 dpt to media with or without ABA or ABA plus IAA.  $n \geq 11$  seedlings for *Col-0* and *pyl8-2*, and  $n = 6$  seedlings for *pyl8-1*. Data are means  $\pm$  SEM.  $**P < 0.01$ ,  $***P < 0.001$ , Student's *t* test. (D and E) The lateral root length of seedlings was measured at the indicated days after transfer to media with or without ABA or ABA plus IAA.  $n \geq 11$  seedlings for *Col-0* and *pyl8-2*, and  $n = 6$  seedlings for *pyl8-1*. Data are means  $\pm$  SEM.  $*P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ , Student's *t* test. (F) Pie charts for


the percentage of quiescent lateral roots for the indicated number of days for seedling grown in media with or without ABA or ABA plus IAA.  $n = 6$  to 18 seedlings per treatment. (G) Percentage of quiescent lateral roots ( $<0.5$  mm at 0 dpt) that exited within 5 dpt to the media with ABA plus IAA.  $n = 6$  to 18 seedlings per treatment. Data are means  $\pm$  SEM.  $*P < 0.05$ , ANOVA.



**Fig. 5. PYL8 enhances auxin signaling through MYB77**


(A) PYL-MYB77 interactions in the yeast two-hybrid assay. Interaction was determined by yeast growth in media lacking His or His and Ade in the presence and absence of ABA. Dilutions ( $10^{-1}$ ,  $10^{-2}$ , and  $10^{-3}$ ) of saturated cultures were spotted onto the plates, which were photographed after 5 days. (B) PYL8-MYB77 interaction in LUC complementation assays in wild-type Col-0 protoplasts.  $n = 4$  experiments. Data are means  $\pm$  SEM.  $***P < 0.001$ , Student's  $t$  test. (C) PYL8-MYB77 interaction in Col-0 wild-type (left panel) and *aba2-1* mutant protoplasts (right panel) with or without ABA in LUC complementation

assays.  $n \geq 7$  experiments. Data are means  $\pm$  SEM.  $*P < 0.05$ ,  $***P < 0.001$ , Student's  $t$  test. (D and E) Binding between MYB77 and the MBSI motif-containing DNA oligonucleotides with or without PYL8 in EMSA assay.  $n = 4$  experiments. Data are means  $\pm$  SEM.  $**P < 0.01$ ,  $***P < 0.001$ , Student's  $t$  test. (F) *IAA19pro-LUC* expression in Col-0 wild-type (left panel), *aba2-1* mutant (middle panel), and *abi1-1* mutant (right panel) protoplasts cotransformed with the indicated combinations of *PYL8*, *MYB77*, and *IAA19pro-LUC* with varying concentrations of ABA and IAA.  $n \geq 3$  experiments. Data are means  $\pm$  SEM.  $*P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ , Student's  $t$  test. (G) The root architecture of Col-0 or *myb77* mutant seedlings was documented at 5 dpt to media (1/4 MS, 1% sucrose) with or without ABA. (H and I) Lateral root (LR) length of Col-0 and *myb77* mutant seedlings at 5 dpt to media with or without ABA.  $n \geq 11$  seedlings. Data are means  $\pm$  SEM.  $*P < 0.05$ ,  $**P < 0.01$ , Student's  $t$  test.



**Fig. 6. PYL8 interacts with and activates MYB44 and MYB73**  
 (A and B) PYL-MYB44 and PYL-MYB73 interactions in the yeast two-hybrid assay. Interaction was determined by yeast growth in media lacking His or His and Ade in the presence and absence of ABA. Dilutions ( $10^{-1}$ ,  $10^{-2}$ , and  $10^{-3}$ ) of saturated cultures were spotted onto the plates, which were photographed after 5 days. (C) *IAA19pro-LUC* expression in *aba2-1* mutant protoplasts cotransformed with the indicated combinations of *PYL8*, *MYB44*, *MYB73*, and *IAA19pro-LUC* with varying concentrations of ABA and IAA.  $n = 3$  experiments. Data are means  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , Student's *t* test.





**Fig. 7. Proposed model of ABA response pathways in regulating lateral root growth**  
 ABA has dual roles in controlling lateral root growth in *Arabidopsis*. ABA exposure promotes periods of quiescence and growth recovery in post-emergence lateral roots. The inhibitory effect of ABA on lateral root growth is mediated by the core ABA signaling pathway through the PYL receptors, PP2C co-receptors, and SnRK2 protein kinases. PYL8 has a previously undescribed function in accelerating lateral root growth during the recovery phase through activating MYB77. PYL8 directly interacts with MYB77 and enhances the activity of MYB77 to up-regulate the expression of multiple auxin-responsive genes at the time when lateral roots enter the recovery phase. MYB77 forms heterodimers with ARF7 and promotes the transcription of ARF7 target genes. At high concentrations of auxin, Aux/IAA is degraded by transport inhibitor response 1 (TIR1)- and auxin signaling F-box (AFB)-mediated ubiquitination to release the inhibition on the ARFs and induce auxin-responsive gene expression, bypassing PYL8.