



The Alleviation of Heat Damage to Photosystem II and Enzymatic Antioxidants by Exogenous Spermidine in Tall Fescue

Liang Zhang^{1,2}, Tao Hu¹, Erick Amombo^{1,2}, Guangyang Wang^{1,2}, Yan Xie^{1*} and Jinmin Fu^{1,3*}

¹ Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan, China, ² Graduate University of Chinese Academy of Sciences, Beijing, China, ³ School of Resources and Environmental Engineering, Ludong University, Yantai, China

OPEN ACCESS

Edited by:

Rosa M. Rivero,
Centro de Edafología y Biología
Aplicada del Segura (CSIC), Spain

Reviewed by:

Marian Brestic,
Slovak University of Agriculture,
Slovakia
Jianming Li,
Northwest A&F University, China

*Correspondence:

Yan Xie
xieyan60b@126.com
Jinmin Fu
jfu@wbpcas.cn

Specialty section:

This article was submitted to
Plant Abiotic Stress,
a section of the journal
Frontiers in Plant Science

Received: 15 June 2017

Accepted: 25 September 2017

Published: 12 October 2017

Citation:

Zhang L, Hu T, Amombo E, Wang G,
Xie Y and Fu J (2017) The Alleviation
of Heat Damage to Photosystem II
and Enzymatic Antioxidants by
Exogenous Spermidine in Tall Fescue.
Front. Plant Sci. 8:1747.
doi: 10.3389/fpls.2017.01747

Tall fescue (*Festuca arundinacea* Schreb) is a typical cool-season grass that is widely used in turf and pasture. However, high temperature as an abiotic stress seriously affects its utilization. The objective of this study was to explore the effect of spermidine (Spd) on heat stress response of tall fescue. The samples were exposed to 22°C (normal condition) or 44°C (heat stress) for 4 h. The results showed that exogenous Spd partially improved the quality of tall fescue leaves under normal temperature conditions. Nevertheless, after heat stress treatment, exogenous Spd significantly decreased the electrolyte leakage of tall fescue leaves. Spd also profoundly reduced the H₂O₂ and O₂^{·-} content and increased antioxidant enzymes activities. In addition, PAs can also regulate antioxidant enzymes activities including SOD, POD, and APX which could help to scavenge ROS. Moreover, application of Spd could also remarkably increase the chlorophyll content and had a positive effect on the chlorophyll α fluorescence transients under high temperature. The Spd reagent enhanced the performance of photosystem II (PSII) as observed by the JIP-test. Under heat stress, the Spd profoundly improved the partial potentials at the steps of energy bifurcations (PI_{ABS} and PI_{total}) and the quantum yields and efficiencies (ϕ P₀, δ R₀, ϕ R₀, and γ RC). Exogenous Spd could also reduce the specific energy fluxes per Q_A⁻ reducing PSII reaction center (RC) (TP₀/RC and ET₀/RC). Additionally, exogenous Spd improved the expression level of *psbA* and *psbB*, which encoded the proteins of PSII core reaction center complex. We infer that PAs can stabilize the structure of nucleic acids and protect RNA from the degradation of ribonuclease. In brief, our study indicates that exogenous Spd enhances the heat tolerance of tall fescue by maintaining cell membrane stability, increasing antioxidant enzymes activities, improving PSII, and relevant gene expression.

Keywords: spermidine, tall fescue, heat stress, antioxidant enzymes, photosystem II, gene expression

INTRODUCTION

Tall fescue (*Festuca arundinacea* Schreb) is a major cool-season grass that is widely used for turf, on the sports field, and as a forage grass with an optimal growth temperature of 16–24°C (Emmons, 2007). However, it is sensitive to heat stress which affects tall fescue turf quality and utilization. When the temperature exceeds the optimal range, heat stress could lead to the photosynthesis

inhibition, cell membrane damage, senescence, severe obstruction in growth, development, and even death (Xu et al., 2006; Mostofa et al., 2014). Therefore, the high temperature is the key limiting factor for tall fescue utilization. It is crucial to explore a convenient method to reduce the damage of tall fescue by heat stress.

Heat stress brings great challenge to the utilization of cool-season turfgrass worldwide. It results in the loss of balance between the scavenging and producing of reactive oxygen species (ROS) (Smirnoff, 1998). ROS can be produced constantly in the process of plant growth and development which includes hydrogen peroxide (H_2O_2), singlet oxygen (1O_2), hydroxyl radical (OH^\cdot), and superoxide anion ($O_2^{\cdot-}$) (Mostofa et al., 2014). Under normal circumstances, the balance of generation and ROS scavenging is harmless to plants. When the botanical species are under environmental stress including heat, on one hand, the accumulation of ROS can lead to injury to the cell membrane via increasing electrolyte leakage (EL) (Liu and Huang, 2000). On the other hand, heat stress could also decrease the activities of antioxidants causing injury to plants (Monk et al., 1989). Overproduction of ROS also has negative influences on biomacromolecule containing membrane lipids, proteins, nucleic acids, and chlorophyll which are necessary for growth and development to plants (Blokhina et al., 2003). The evolution of higher plants has developed the ROS-scavenging system to defend against oxidative stress (Mittler et al., 2004). Enzymatic antioxidants include superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), and ascorbate peroxidase (APX) which can scavenge H_2O_2 , $O_2^{\cdot-}$, and oxy-intermediates (Apel and Hirt, 2004; Lee et al., 2007). The SOD is regarded as the first line of defense against ROS and catalyzes $O_2^{\cdot-}$ to H_2O_2 and oxygen (Sigaud-Kutner et al., 2002), while H_2O_2 can be further removed by POD and APX through dismutation or ascorbateglutathione cycle (Mostofa et al., 2014; Liu et al., 2016).

Photosynthesis involves a series of complex metabolic reactions which are not only vital for biological survival but also forms a critical carbon-oxygen balance on earth (Berry and Downton, 1982). The structural and functional photosynthetic machinery is sensitive and vulnerable to severe or mild heat stress (Essemine et al., 2011). Photosystem II (PSII) as the core portion of the photosynthetic process whose components are susceptible to high-temperature stress and are seriously damaged (Hideg et al., 2002). PSII located in the thylakoid membranes of oxygenic photosynthetic organisms is a membrane protein complex with multi-subunit that catalyzes a series of electron transfer reactions (Umena et al., 2011). Basically, the PSII catalyzes the unique reactions resulting in the splitting of water and the production of dioxygen and reducing equivalents (Barber, 1998). The PS II is composed of two different parts of structure and function. One as the reaction center (RC) of PS II is constituted by a D1-D2 heterodimer binding capture complex named CP43 and CP47. The other part is the pigment protein complex which binds plenty of Chl a, Chl b, and lutein (Barber, 1998; Pfannschmidt, 2003; Pospíšil, 2012). Thereinto, D1 protein is the most important subunit which can provide a position for cofactors to bind, maintain the structure of PS II reaction center, and have a close

connection with the separation and transmission of the original charge (Kruse et al., 1997).

It has been previously reported that many low molecular compounds have an essential role in plants to respond to abiotic stress. Bartwal et al. (2013) reported that brassinosteroids could enhance the tolerance of chilling, heat, salt, and drought in rice, tomato, beet, and wheat by maintaining membrane stability and modulating the expression of relevant genes. Tan et al. (2011) study showed that $CaCl_2$ treatment has a positive effect on improving heat tolerance of tobacco by elevating net photosynthetic rate, thermostability of reaction center of PSII, antioxidative enzymes activity, and HSP70 level. Salicylic Acid has been found to exert some positive effects on the improvement of malting barley resistances to heavy metals by increasing the activities of SOD and CAT (Song et al., 2014). Except above, polyamines (PAs), including putrescine (Put), spermidine (Spd), and spermine (Spm), is one of the vital compounds. The PAs play crucial roles in various abiotic stresses, including salt, drought, high temperature, wounding, ozone, flooding, heavy metals, acid, and oxidative stresses (Shi and Chan, 2014). PAs are a type of ubiquitous nitrogenous compounds containing two or more amines and exist in almost all organisms (Alcázar et al., 2010). They are widely involved in the regulation of growth and development in plants, such as morphogenesis, root elongation, pollen viability, leaf senescence, fruit ripening, programmed cell death, DNA synthesis, gene transcription, protein translation, and chromatin organization (Shi and Chan, 2014). Additionally, PAs are also considered as vital secondary messengers in the signaling pathway (Kusano et al., 2008). They could maintain membrane stability by their interaction with phospholipids as well as scavenge ROS (Roberts et al., 1986; Besford et al., 1993). PAs will largely accumulate under abiotic stresses including high temperature (Todorova et al., 2007; Goyal and Asthir, 2010).

Exogenous application low molecular compound is regarded as one of the efficient methods to alleviate environmental stresses of plants. There is tremendous progress concerning exogenous PAs response to abiotic stresses in rice, tomato, and cucumber (Tian et al., 2012; Mostofa et al., 2014; Hu L. et al., 2016). Most of them had a close association with enhanced levels of antioxidant capacity. However, studies on the effect of Spd on PSII and the expression of relevant genes in tall fescue under heat stress have still been obscure. Therefore, the objectives of this study were to explore the influence of Spd on chlorophyll α fluorescence, antioxidant enzyme activity, and photosynthetic gene transcription level in tall fescue under high temperature to enrich the information in our attempts to comprehend turf breeding and management.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

This research was conducted at Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan, China in 2016. The plant used in this study was PI234881 seeded in plastic pots (13 cm in diameter and 15 cm deep) with matrix (brown coal soil and sand = 1:1). Plants were maintained in the greenhouse

after germination with day/night temperature 22/18°C ($\pm 2^\circ\text{C}$), humidity 80% and illumination 14 h (with light intensity of $300 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) for 50 days. The seedlings were watered daily and fertilized once a week with 100 mL of half-strength Hoagland's solution (Hoagland and Arnon, 1950).

Reagent Treatments

The reagent treatments were performed using a vacuum infiltration to allow efficient transportation of reagents into the leaves (Filippou et al., 2012) as described by Carolina et al. and Luisa et al. with little modification (Ederli et al., 2006; Attallah et al., 2007). The collection was done by snipping third fully expanded leaves from petiole which were subsequently immersed in half-strength Hoagland's solution with various pharmacological reagents in 15 cm length containers. Later, the leaves were infiltrated with a desiccator under dark vacuum condition for 15 min where they are maintained for 4 h to enable fully recovering before heat stress treatment.

Heat Stress Treatment

For the heat stress treatment, the leaf petioles were immersed in a 1-cm deep half-strength Hoagland's solution kept in falcon tubes. Subsequently, the tubes were transferred into two growth chambers with temperature regimes of 22°C (CK and S) or 44°C (H and HS). The light intensity was set at $450 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ at 80% humidity. Each treatment was repeated at least three times.

Evaluation of the Optimum Spd Concentration

To determine the adequate effective Spd concentration for alleviating heat stress, we performed a preliminary experiment by applying different concentration Spd. The concentration of Spd (0, 0.5, 1, and 2 mM) were chosen preliminarily according to the Mostofa experiment on rice (Mostofa et al., 2014). Subsequently, we selected the optimum concentration (0.5 mM) by comparing the fluorescence transients after heat stress 4 h (Figure 1). Figure 1 shows the differential changes in chlorophyll fluorescence transients after treatment with different concentration of Spd under heat stress. A 0.5 mM of Spd had the positive impact on photosynthesis by improving F_j , F_i , and F_p .

Chlorophyll Content

The leaf chlorophyll content was measured based on the method described by Hiscox and Israelstam (1979). Briefly, fresh leaves (0.1 g) were immersed in a 10-mL dimethylsulfoxide tube, then the absorbance of the samples at 645 and 663 nm was measured by spectrophotometer (UV-2600, UNICO, Shanghai) after 72 h dark treatment. The chlorophyll content was calculated by the following formula:

$$\text{Chl a (mg/g FW)} = (12.72 * \text{OD}_{663} - 2.59 * \text{OD}_{645}) * 0.3$$

$$\text{Chl b (mg/g FW)} = (22.88 * \text{OD}_{645} - 4.67 * \text{OD}_{663}) * 0.3$$

$$\text{Chl total (mg/g FW)} = (20.2 * \text{OD}_{645} + 8.02 * \text{OD}_{663}) * 0.3.$$

Where OD_{645} and OD_{663} are the absorbance of the extract solution at 645 and 663 nm, respectively, and FW is the fresh weight of the leaf.




FIGURE 1 | Evaluation of the optimum Spd concentration by OJIP curves. The OJIP fluorescence transients in tall fescue leaves with 0, 0.5, 1, and 2 mM Spd at high temperature (44°C) for 4 h.

Electrolyte Leakage (EL)

To quantify the EL, 0.1 g of treated leaves were washed three times with deionized water. The leaves were cut into 0.5 cm long debris and put into test tubes filled with 15 mL deionized water. The tubes were shaken for 24 h at 25°C and the initial conductivity (C_i) was measured by a conductivity meter (JENCO-3173, Jenco Instruments, Inc., San Diego, CA, United States). Subsequently, the leaves were autoclaved at 121°C for 30 min to release the electrolytes of the tissue completely. The final conductivity (C_{max}) was measured after the solution had been cooled to room temperature. The relative EL was calculated with the formula:

$$\text{EL (\%)} = (C_i/C_{max}) * 100\%.$$

Crude Enzyme Extraction

For enzyme extracts, a 0.2 g of leaves powder with liquid nitrogen was immersed in 4 mL phosphate buffer (150 mM, pH 7.0) precooled at 4°C homogenized with 0.2 M Na_2HPO_4 and 0.2 M NaH_2PO_4 . Then, the homogenate was centrifuged at $15,000 \times g$ at 4°C for 30 min. Finally, the supernatant was collected and stored at 4°C to determine enzyme activities.

Antioxidant Enzyme Activity

For the SOD activity assay, a 0.1 mL enzyme extract was added into 2.9 mL solution plus 50 mM phosphate buffer (pH 7.8), 1.125 mM nitro blue tetrazolium (NBT), 60 μM riboflavin, 195 mM methionine and 3 μM ethylene diamine tetraacetic acid (EDTA). Then, the solution was incubated under 4000 lx irradiance for 30 min. The change of absorbance at 560 nm was recorded with 3 mL of solution without enzyme extract as the control. One unit of SOD activity was defined as the inhibition of NBT reduction by 50%.

The POD activity was measured based on the method described by Fan et al. (2014). In brief, a 50 μL enzyme extract

TABLE 1 | Primer sequences and information used for reserved transcription real-time PCR (RT-PCR) analyses.

Gene	Encoded polypeptide		Primers sequences (5'–3')	Size (bp)	Gene ID
psb A	D1 protein	F	GTATTTATTATCGCCTTCATCG	284	7095419
		R	AGGACGCATACCCAAACG		
psb B	CP47	F	TAGGCGTAACGGTGGGA	254	7095420
		R	AACATCTCGGAACAAGG		
psb C	CP43	F	TAATACGGCTTATCCGAGTGAGTTT	288	7095484
		R	TCTTGCCAAGGTTGTATGTCTTT		

F and R represent forward and reverse, respectively.

was added into 2.95 mL solution containing 0.075% H₂O₂, 0.1 M sodium acetate-acetic buffer (pH 5.0), 0.25 mL guaiacol (dissolved in 50% ethanol solution). Then we recorded the absorbance changes at 460 nm per minute for 3 min. One unit POD activity is defined as the increase in absorbance per minute.

The APX activity was measured using Plant APX Elisa Kit (Huijia Biotechnology Institute, Xiamen, China).

H₂O₂ and O₂^{•-} Content

The H₂O₂ content was determined using a H₂O₂ Assay Kit (Nanjing Jiancheng Bioengineering Institute, China).

The O₂^{•-} content was measured using Plant SOA Elisa Kit (Huijia Biotechnology Institute, Xiamen, China).

Chlorophyll (Chl) α Fluorescence Transient

Chl α fluorescence transients were recorded by pulse-amplitude modulation (PAM) fluorometer (PAM 2500, Heinz Walz GmbH). After 30 min of adaption in the darkness, leaves were triggered with the red light of 3000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to attain OJIP transients which were measured and digitized between 10 and 320 ms. The data of OJIP transients analyzed method was initially reported by Strasser et al. (2004). In the present assay, the data of OJIP transients were analyzed by using JIP-test as reported by Chen et al. (2014). The JIP-test is used for analysis parameters of OJIP transient, which is based on the energy fluxes in the biofilm. These parameters digitize photosynthesis to further study of the photosystem.

Quantitative RT-PCR Analysis

The levels of gene expression were analyzed by approximately 0.1 g crushed leaves. Total RNA was extracted and purified by Trizol-reagent (Invitrogen, Carlsbad, CA, United States) according to the instruction. About 0.2 μg RNA was used for synthesizing the first-strand cDNA using M-MLV reverse transcriptase (Promega, Madison, WI, United States) with an oligo (dT) primer. Then we examined the quality of cDNA by gel electrophoresis and maintained it at -80°C for qRT-PCR analysis. Specific primers (Table 1) were designed for analyzing gene expression, fluorescent dye SYBR Green (Toyobo, Osaka, Japan) was applied in the detection system. Real-time PCR reaction was performed by the real-time PCR Master Mix (Toyobo) according to the manual. The TUB gene was used as an internal control. The method used to determine the relative

quantity of the target gene expression was proposed by Chen et al. (2009).

Statistical Analysis

In the experiment, all values were shown as mean \pm (Standard Error) SE with at least triplication. One-way analysis of variance (ANOVA) and Duncan's multiple range test were conducted using SPSS (version 20.0, SPSS Institute, Chicago, IL, United States) statistical software and 5% level of probability was used to test the significant effects of treatments. The graphs were produced by Origin 8.0 (Origin Lab, Inc., Hampton, MA, United States) and Excel 2010 for Windows.

RESULTS

Electrolyte Leakage

The EL alteration was measured to investigate the role of exogenous Spd in maintaining cell membrane stability of tall fescue under heat stress. The result showed that EL values in the





FIGURE 2 | Effects of EL in tall fescue leaves after 0.5 mM Spd treatment under heat stress. CK was normal temperature of 22°C. H was high temperature where tall fescue leaves were treated at 44°C. S was treated with Spd at normal temperature. HS was treated with Spd at high temperature. Values are given as means \pm SE of three independent experiments. Different letters indicate statistical difference significance at $P < 0.05$ among the treatments by Duncan's multiple range test.



leaves of control and Spd treatment regime had no significant difference. However, under heat stress, the EL value increased five folds, compared to the control. The application of Spd to stressed leaves significantly reduced EL by 28.64% compared to heat stress treatment (Figure 2).

Chlorophyll Content



When plants were exposed to various abiotic stresses, the leaves would exhibit chlorosis and a significant change in the chlorophyll content. Therefore, chlorophyll content is usually taken as an indicator that reflected plants resistance to stress. Under the control condition, there was no significant difference in the Chl a, Chl b, total chlorophyll content, and the ratio of Chl a to Chl b (Figure 3). Conversely, after heat stress treatment, Chl a, Chl b, and the total chlorophyll content notably decreased by 16.47, 14.95, and 14.96%, respectively. Interestingly, Spd application under heat stress could notably enhance the Chl a, Chl b, total chlorophyll content, and the ratio of Chl a to Chl b by 22.04, 17.32, 19.40, and 4.06%, respectively.

Antioxidant Enzyme Activities

To investigate the effect of Spd on the antioxidant enzymes, several antioxidant enzymes activities were assayed (Figure 4), which included SOD, POD, and APX activities. Under normal temperature, SOD activity had no change after applied Spd. However, after heat stress, the activity of SOD decreased by 9.59%. Under the heat stress, the activity of SOD after Spd treatment increased significantly by 20.67%. Heat stress damaged the activity of POD which decreasing by 15.74% compared to control. After treatment with Spd, POD activity increased significantly by 21.51%, returned to normal levels. The activity of APX was also remarkably elevated 13.52% that treated by Spd under high temperature.

H₂O₂ and O₂^{•-} Contents

There was not a notable difference of H₂O₂ content in leaves under normal temperature whether application Spd or not (Figure 5). However, after heat treatment, the H₂O₂ content significantly increased about 10% compared to the control. Interestingly, the Spd treatment decreased the H₂O₂ content



remarkably to normal condition. Similarly, under normal temperature, the content of O₂^{•-} decreased sharply by 43.3% after applied Spd treatment, then it rose again in the leaves after heat stress treatment. However, the Spd treatment reduced the O₂^{•-} contents to a normal level.

The OJIP Fluorescence Transient and JIP-Test

On one hand, under normal temperature, Spd application increased the OJIP curve compared to non-Spd treated regime (**Figure 6**). On the other hand, heat stress treatment made the OJIP curve decline dramatically after 4 h, which was partially ameliorated by Spd treatment.

To further study the effect of Spd on the photosynthetic system under heat stress, we employed the JIP-test to analyze the parameters of OJIP transient curves. We extracted F₀, F_K, F_J, F_I, F_P, and M₀ as basic parameters as displayed in **Table 2**. As shown, there was no difference in the F_K among all treatments. However, heat stress treatment profoundly reduced the F_J, F_I, F_M, while increased the F₀, M₀. Meanwhile, the parameters had

the tendency return normal values in Spd treatment under high temperature. The leaves treated with high temperature without Spd had the highest F₀, M₀, and the lowest other parameters.

All of the above basic parameters were analyzed by the JIP-test to deduce further the structural and functional parameters to quantify the photosynthesis of tall fescue leaves. There was almost no significant difference except for the value of δR_0 in control and Spd treatment on the condition of normal temperature. In terms of quantum yields and efficiencies or probabilities, heat stress decreased the values of ϕP_0 (maximum quantum yield), δR_0 (efficiency with which an electron from Q_B is transferred until PSI acceptors), γRC (probability that PSII Chl molecule functions as RC), and ϕR_0 (quantum yield for reduction of end electron acceptors at the PSI acceptor side) compared to normal condition. On the other hand, exogenous Spd notably enhanced the values of the ϕP_0 , δR_0 , γRC , and ϕR_0 (**Figures 7A–D**).

Meanwhile, several parameters which were also analyzed to specify energy fluxes were displayed in **Figures 7E,F**. In the absence of heat stress, there were no obvious effects on the




FIGURE 6 | The effect of chlorophyll fluorescence transients (OJIP curve) in tall fescue leaves after 0.5 mM Spd treatment under heat stress. CK was normal temperature of 22°C. H was high temperature where tall fescue leaves were treated at 44°C. S was treated with Spd at normal temperature. HS was treated with Spd at high temperature. Values are given as means \pm SE of three independent experiments. Different letters indicate statistical difference significance at $P < 0.05$ among the treatments by Duncan's multiple range test.

TP_0/RC (trapped excitation flux per RC), and ET_0/RC (electron transport flux per RC) for CK and Spd treatments. However, the values of TP_0/RC and ET_0/RC were higher under high temperature, while these values reduced after Spd application.

Performance index (PI) including PI_{total} and PI_{ABS} are important indices to describe the overall activity of PSII. It was shown in the **Figure 8** that PI was much higher in those under heat stress. After exogenous Spd treatment, PI_{ABS} became conspicuously higher than the leaves with heat treatment. In addition, the PI_{total} increased by approximately two folds compared with the high-temperature treatment.

Relevant Gene Expression to Photosynthetic System

Three genes, *psbA*, *psbB*, and *psbC*, encoding D1 protein, CP47, and CP43 involved in the photosynthetic system were measured with expression levels to further explore the protective role of Spd in tall fescue leaves against heat stress. We observed

that exogenous Spd made tremendous contributions toward enhancing the expression levels of *psbA* and *psbB*, but it did not affect on *psbC* under the normal conditions (**Figure 9**). Heat stress suppressed the expression of the three genes compared to normal temperature. However, Spd application profoundly promoted the *psbA* and *psbB* expressions while did not affect on *psbC*.

DISCUSSION


Tall fescue, as a typical cool-season turf and forage grass, is limited by heat stress. Therefore, it is vital to improving the thermotolerance for its efficient use in turf industry. It has been reported that Spd plays a crucial role of against abiotic stresses, such as salt, drought, heat, and salinity-alkalinity stresses in tomato, cucumber, rice (Farooq et al., 2009; Tian et al., 2012; Hu L. et al., 2016). In the present study, we would explore the alleviative effect of Spd to heat stress in tall fescue.

Cell membrane breakage is one of the major damages caused by heat stress which leads to cell permeability and EL (Liu and Huang, 2000). Therefore, the EL could usually be used as an indicator to reflect the damage of membrane in heat stress (Blum and Ebercon, 1981; Marcum, 1998). It was reported that exogenous application of Spd could induce endogenous PAs generating (Li et al., 2016). And PAs, in the state of polycation, could attach to the phosphate groups with anions which results in stability of the membrane and intracellular material outflow (Galston and Sawhney, 1990). Additionally, PAs are ideally suitable for physiological channel modulator to block fast vacuolar channel activity and K^+/Na^+ homeostasis partially at a physiological pH (Alcázar et al., 2010). Moreover, some studies previously indicated that PAs can also non-covalently bind to some low molecules including proteins in the plasma membrane, antibiotics, phenolic acids, and hydroxycinnamic acid (Feuerstein and Marton, 1989; Martin-Tanguy, 2001). In the present study (**Figure 2**), the value of EL soared after high-temperature treatment which suggested that tall fescue leaf cells experienced serious EL. However, the EL was then reduced after Spd treatment. Those results indicated that exogenous applied Spd could induce the PAs generating in tall fescue leaf cells. Besides, PAs may interact with negatively charged phosphate groups and low molecules mentioned above. As a result, the treatment of Spd had a positive effect on balancing the pH and ionic exchange through plasma membrane. Therefore, exogenous

TABLE 2 | Basic photosynthetic parameters extracted from the OJIP transient curves.

Treatment	F_0	F_K	F_J	F_I	F_M	M_0
CK	0.25 \pm 0.01b	0.57 \pm 0.01a	0.67 \pm 0.02b	0.97 \pm 0.01a	1.05 \pm 0.01a	1.59 \pm 0.08b
S	0.26 \pm 0.02b	0.59 \pm 0.01a	0.72 \pm 0.02a	1.00 \pm 0.02a	1.08 \pm 0.02a	1.61 \pm 0.03b
H	0.31 \pm 0.01a	0.57 \pm 0.01a	0.61 \pm 0.01c	0.82 \pm 0.01c	0.87 \pm 0.01c	1.83 \pm 0.05a
HS	0.28 \pm 0.01ab	0.58 \pm 0.01a	0.71 \pm 0.01ab	0.88 \pm 0.01b	0.98 \pm 0.01b	1.75 \pm 0.02ab



CK was normal temperature of 22°C. H was high temperature where tall fescue leaves were treated at 44°C. S was treated with Spd at normal temperature. HS was treated with Spd at high temperature. Values are given as means \pm SE of three independent experiments. Different letters indicate statistical difference significance at $P < 0.05$ among the treatments by Duncan's multiple range test.



Spd could decrease the EL of tall fescue leaf cells which under heat stress.

Under optimum conditions, the plants could balance the generating and scavenging ROS (Bowler et al., 1992) through a

well-organized scavenging system (Tian et al., 2012). However, the ROS, such as H_2O_2 and $O_2^{\cdot-}$, will largely accumulate when the plants suffer from various abiotic stress. These abiotic stresses could lead to plant metabolic inactivation, cell death, net



photosynthesis rate and photosynthate reduction or even cause the loss of plant quality and serious yield reduction (Mallick and Mohn, 2000). Under heat stress, the ROS production always exceeds the defense capability of antioxidant resulting in macromolecules being damaged in living cells (Liu and Huang, 2000; Tian et al., 2012). The SOD-POD and APX line plays crucial role in scavenging ROS. The SOD could catalyze $O_2^{\cdot-}$ to H_2O_2 and oxygen (Sigaud-Kutner et al., 2002). The H_2O_2 can be further removed by POD and APX through dismutation or ascorbateglutathione cycle afterward (Mostofa et al., 2014; Liu et al., 2016). In the present study, the H_2O_2 , $O_2^{\cdot-}$, SOD, POD, and APX activities were measured to explore the effect of Spd in scavenging ROS and promoting antioxidant enzymes activities (Figures 4, 5). The result showed that exogenous Spd could significantly decrease the H_2O_2 and $O_2^{\cdot-}$ contents which indicated the application of Spd alleviating the heat stress suffered by tall fescue leaves. On the other hand, antioxidant enzymes

activities increased obviously regardless of whether the activities of these enzymes were inhibited or raised by high-temperature with the treatment of Spd. Shi et al. (2013) has reported that Spd could regulate nucleoside diphosphate kinase (NDPK) and three antioxidant enzymes (2-Cys POD, APX, Cu/Zn SOD) in bermudagrass. PAs increased level of NDPK₂ protein which is directly related with activated activities of antioxidant enzymes (Shi and Chan, 2014). Namely, exogenous PAs treatment regulated the level of NDPK₂ protein related with activating the activities of antioxidant enzymes to inhibit accumulations of $O_2^{\cdot-}$ and H_2O_2 under stress conditions (Shi et al., 2013). In addition, exogenous Spd may alter the antioxidant enzymes isozymes zymogram expression, which could also enhance plant tolerance at high temperature (Tian et al., 2012). Some studies reported that PAs could also be as efficient antioxidants by their character of polycation under various environmental stresses (Groppa et al., 2001). From the results, we observed that

exogenous Spd could have the positive effects on inhibiting ROS generation and promoting antioxidative enzymes activities. Therefore, we also deduce that Spd could enhance the tall fescue thermotolerance resulting from removing ROS by regulating the expression of relevant gene.

Photosynthesis, inhibited by cold, heat, drought, saline-alkaline, and nutritional deficiency, is one of the most processes sensitive to diverse abiotic stresses (Jiang et al., 2001; Bi et al., 2016; Hu Z. et al., 2016; Xiang et al., 2016). PSII, as an important membrane structure of the photosynthesis processes, is also vulnerable to high temperature (Chen et al., 2014; Bi et al., 2016). Chlorophyll fluorescence kinetics transients could provide abundant information of the original photochemical reaction of the mechanism of photosynthesis, mainly in PSII donor side, receptor side, and reaction centers (Murkowski, 2001; Chen et al., 2014). To figure out the adaption mechanism of high temperature in tall fescue leaves treated by Spd, chlorophyll fluorescence transients and JIP-test were investigated. As shown in **Table 2** and **Figure 6**, there was subtle difference between the treatments regardless of application of Spd under normal condition. However, heat stress aggravated the function of PSII as observed after altering the F_0 , F_j , F_i , F_M , and M_0 . Exogenous Spd obviously alleviated the inhibition of heat stress through observing the values which mentioned above. The difference indicates that exogenous Spd is vital to tall fescue heat stress-resistance by protecting the PSII. This result accorded with Murkowski (2001) study that adequate concentration Spd could alleviate heat stress damage of tomato PS II. Then we evaluated the quantum yields and efficiencies by maximum quantum yield for primary photochemistry (ϕP_0 , TR_0/ABS), efficiency with which an electron from QB is transferred to PSI acceptors (δR_0 , RE_0/ET_0), the quantum yield for reduction of end electron acceptors at the PSI acceptor side (ϕR_0 , RE_0/ABS), and the probability that a PSII Chl molecule functions as RC (γRC) in PSII. The result indicated that heat stress notably decreased the efficiency of electron transportation and Chl molecule functions of PSII. Exogenous Spd remarkably improved the ϕP_0 , δR_0 , ϕR_0 , and γRC . Videlicet, Spd has a positive effect on the side of donor and acceptor of PSII under high temperature. Specific energy fluxes including TP_0/RC and ET_0/RC were analyzed to detect the functional properties of PSII (**Figures 7E,F**). The result determined that the plant could not balance between light absorption and utilization under heat treatment, and it had a negative effect on the RC. After application of Spd, the trapped excitation flux (leading to Q_A reduction) and electron transport flux (further than Q_A^-) per RC resumed to normal level. These indicate that Spd increased the active RC and alleviated the damage on RC under the heat stress. Performance index (PI) including PI_{ABS} and PI_{total} , is the most sensitive parameters of the JIP-test which is used for assessing the photochemical activities of stressed plants (Fan et al., 2015). The PI integrates several parameters containing light energy absorption, excitation energy trapping, and conversion of excitation energy to electron transport (Fan et al., 2015). In the present study, we ascertained that Spd treatment has a positive effect on PI_{total} and PI_{ABS} which were much higher than heat stress alone (**Figure 8**). As analyzed above, we can conclude that exogenous Spd has protective effects

on PS II which is a very heat-sensitive membrane structure. To a certain degree, the results in PSII are consistent with Chen et al. (2013) studies on NO alleviation of heat damage in tall fescue.

The chlorophyll content of leaves displays crucial information concerning the physiological condition of the plants (Gitelson et al., 2003). The chlorophylls, Chl a and Chl b, are really important pigments in the process of photosynthesis which related to the transforming light energy to chemical energy (Gitelson et al., 2003). It is known that both Chl a and Chl b can absorb light energy, but only a handful of the excited state Chl a can transform light energy into electrical energy. Chl b is a pigment of antenna complexes in green algae (Green and Durnford, 1996). It also plays an important role in regulating the size of the photosynthetic antenna and maintaining the stability of light-harvesting complex associated with PSII (LHCII) in plants (Yamasato et al., 2005). The ratio of chlorophyll a to b (Chl a/b) is a vital value of LHCII to other components that contain chlorophyll (Leong and Anderson, 1984). Tanaka et al. (2001) reported that the overexpression of CAO (chlorophyllide an oxygenase) in *Arabidopsis thaliana*, a key enzyme in the process of synthesis of Chl b, could increase the expression of antenna by 20%, and LHCII and CP47 content also increased by 40%. However, heat stress may induce a decline in chlorophyll level and the ratio of Chl a/Chl b in tall fescue leaves (Bi et al., 2016). In the present study, the content of Chl a, Chl b, and total chlorophyll content decreased under the high-temperature treatment. The impact on degrading chlorophyll by heat stress was improved after application Spd. As well, the ratio of Chl a to Chl b was higher after Spd treatment under heat stress. The result implies that exogenous Spd may act as a regulator to prevent chlorophyll from disintegrating and thus further protecting the photosynthetic antenna and structure of PSII to improve the heat tolerance and photosynthetic efficiency of tall fescue. Heat stress may induce that chlorophyll decreasing by generating ROS. We deduced that the Spd could improve the chlorophyll content via scavenging ROS, directly or indirectly. Previous study also indicated that low chlorophyll content also has disadvantages to photosynthesis and signaling (Brestic et al., 2016). That is according with our results of JIP-test. High temperature has a negative effect on chlorophyll and further influence on the electron transforming resulting to the function of PS II been damaged.

In higher plants, CP43 (*psbC* encoded protein) and CP47 (*psbB* encoded protein) are the core antenna protein complexes with the composition of chlorophyll a located in the RC of PSII (Bricker, 1990; Barber, 2003). They can transfer the excitation energy (captured by the peripheral antenna chlorophyll a/b protein complex, LHCII) to RC and also be involved in water splitting and maintain PSII core complex structure (Bassi et al., 1987; Bricker, 1990). D1 protein is the most essential subunit which can bind to cofactors, protect the structure of PS II RC, and have a close connection with the separation and transmission of the original chemical reaction (Kruse et al., 1997). Under high temperature, CP43 will uncouple the light harvesting antenna from the RC (Yoshioka et al., 2006). The damaged D1 protein

will also splits after CP43 is released from RC, then the active RCs decrease leads to inefficient energy utilization as a result of a reduction in CP43 and CP47 (Vani et al., 2001). The data in **Figures 7–9** agree with these reports. However, the application of Spd promoted the transcription of CP47 and improves the behavior of RC to some extent. Thus, Spd could partially alleviate and heat damage for RC. D1 protein encoded by *psbA* is the most fundamental structure of PSII and a variety of cofactors related to the original charge separation and electron transfer are combined in an orderly in this structure (Bredenkamp and Baker, 1994). It was reported that a D1 protein was dramatically damaged when exposed to singlet oxygen produced from the water-oxidation reaction (Takahashi et al., 2004). Therefore, the protection and recovery of D1 proteins damaged by heat stress which are crucial for the RC of PSII. As observed in this study, the expression of *psbA* increased significantly in the presence of Spd, which could be a contributory toward the stability of PSII under heat stress. Polyamines can stabilize the structure of nucleic acids by its cation combined with the negative charge on phosphate groups of nucleic acid. It has been reported that exogenous PAs can protect RNA from the degradation of ribonuclease (Goyns, 1982). PAs may also as a mediator or secondary messenger to activate gene network with a potential to protect plants from biotic and abiotic stresses (Paschalidis and Roubelakis-Angelakis, 2005; Cona et al., 2006). It has also been reported that a number of genes were activated by PAs. These genes mainly about the transcription, translation, signal transduction, stress protein biosynthesis (Cheng et al., 2012). Exogenous Spd may regulate

gene expression as discussed above. However, the detail of the mechanism is still unclear.

CONCLUSION

Exogenous PAs has been regarded as an efficient method to alleviate plants heat stress. In this research, we found that exogenous Spd could improve the heat tolerance of tall fescue by protecting all kinds of membranes structures from ROS. As well, it has positive effects on increasing antioxidant enzymes activities and stabilizing the structure of nucleic acids. Besides, the heat damage to PS II of tall fescue is also been alleviated.

AUTHOR CONTRIBUTIONS

JF and YX designed the experiments. LZ performed the experiments, and wrote the manuscript. GW assisted with doing the experiments. TH analyzed the data. JF and EA helped to draft the manuscript and revised manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGMENT

This research was funded by the National Natural Science Foundation of China (No. 31470363, 31772349).

REFERENCES

- Alcázar, R., Altabella, T., Marco, F., Bortolotti, C., Reymond, M., Koncz, C., et al. (2010). Polyamines: molecules with regulatory functions in plant abiotic stress tolerance. *Planta* 231, 1237–1249. doi: 10.1007/s00425-010-1130-0
- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373. doi: 10.1146/annurev.arplant.55.031903.141701
- Attallah, C. V., Welchen, E., and Gonzalez, D. H. (2007). The promoters of arabidopsis thaliana genes atcox17-1 and -2, encoding a copper chaperone involved in cytochrome c oxidase biogenesis, are preferentially active in roots and anthers and induced by biotic and abiotic stress. *Physiol. Plant.* 129, 123–134. doi: 10.1111/j.1399-3054.2006.00776.x
- Barber, J. (1998). Photosystem two. *Biochim. Biophys. Acta Bioenerget.* 1365, 269–277.
- Barber, J. (2003). Photosystem II: the engine of life. *Q. Rev. Biophys.* 36, 71–89. doi: 10.1017/S0033583502003839
- Bartwal, A., Mall, R., Lohani, P., Guru, S. K., and Arora, S. (2013). Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *J. Plant Growth Regul.* 32, 216–232. doi: 10.1007/s00344-012-9272-x
- Bassi, R., Hoyer-Hansen, G., Barbato, R., Giacometti, G. M., and Simpson, D. J. (1987). Chlorophyll-proteins of the photosystem II antenna system. *J. Biol. Chem.* 262, 13333–13341.
- Berry, J. A., and Downton, W. J. S. (1982). Environmental regulation of photosynthesis. *Photosynthesis* 2, 263–343. doi: 10.1016/B978-0-12-294302-7.50017-3
- Besford, R. T., Richardson, C. M., Campos, J. L., and Tiburcio, A. F. (1993). Effect of polyamines on stabilization of molecular complexes in thylakoid membranes of osmotically stressed oat leaves. *Planta* 189, 201–206. doi: 10.1007/BF00195077
- Bi, A., Fan, J., Hu, Z., Wang, G., Amombo, E., Fu, J., et al. (2016). Differential acclimation of enzymatic antioxidant metabolism and photosystem II photochemistry in tall fescue under drought and heat and the combined stresses. *Front. Plant Sci.* 7:453. doi: 10.3389/fpls.2016.00453
- Blokhina, O., Virolainen, E., and Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann. Bot.* 91, 179–194. doi: 10.1093/aob/mcf118
- Blum, A., and Ebercon, A. (1981). Cell membrane stability as a measure of drought and heat tolerance in wheat. *Crop Sci.* 21, 43–47. doi: 10.2135/cropsci1981.0011183X002100010013x
- Bowler, C., Van Montagu, M., and Inz'e, D. (1992). Superoxide dismutase and stress tolerance. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43, 83–116. doi: 10.1146/annurev.pp.43.060192.000503
- Bredenkamp, G. J., and Baker, N. R. (1994). Temperature sensitivity of D1 protein metabolism in isolated *Zea mays*, chloroplasts. *Plant Cell Environ.* 17, 205–210. doi: 10.1111/j.1365-3040.1994.tb00284.x
- Brestic, M., Zivcak, M., Kunderlikova, K., and Allakhverdiev, S. I. (2016). High temperature specifically affects the photoprotective responses of chlorophyll b-deficient wheat mutant lines. *Photosynth. Res.* 130, 251–266. doi: 10.1007/s11120-016-0249-7
- Bricker, T. M. (1990). The structure and function of CPa-1 and CPa-2 in photosystem II. *Photosyn. Res.* 24, 1–13. doi: 10.1007/BF00032639
- Chen, K., Fan, J., and Fu, J. (2013). Alleviation of heat damage to photosystem II by nitric oxide in tall fescue. *Photosyn. Res.* 116, 21–31. doi: 10.1007/s11120-013-9883-5
- Chen, K., Sun, X., Amombo, E., Zhu, Q., Zhao, Z., Chen, L., et al. (2014). High correlation between thermotolerance and photosystem II activity in tall fescue. *Photosyn. Res.* 122, 305–314. doi: 10.1007/s11120-014-0035-3
- Chen, L., Ren, F., Zhong, H., Jiang, W., and Li, X. (2009). Identification and expression analysis of genes in response to high-salinity and drought stresses in *Brassica napus*. *Acta Biochim. Biophys. Sin.* 42, 154–164. doi: 10.1093/abbs/gmp113
- Cheng, L., Sun, R. R., Wang, F. Y., Peng, Z., Kong, F. L., Wu, J., et al. (2012). Spermidine affects the transcriptome responses to high temperature stress in

- ripening tomato fruit. *J. Zhejiang Univ. Sci. B.* 13, 283–297. doi: 10.1631/jzus. B1100060
- Cona, A., Rea, G., Angelini, R., Federico, R., and Tavladoraki, P. (2006). Functions of amine oxidases in plant development and defence. *Trends Plant Sci.* 11, 80–88. doi: 10.1016/j.tplants.2005.12.009
- Ederli, L., Moretini, R., Borgogni, A., Wasternack, C., Miersch, O., Reale, L., et al. (2006). Interaction between nitric oxide and ethylene in the induction of alternative oxidase in ozone-treated tobacco plants. *Plant Physiol.* 142, 595–608. doi: 10.1104/pp.106.085472
- Emmons, R. D. (2007). *Turfgrass Science and Management*. Albany, NY: Delmar.
- Essemine, J., Govindachary, S., Ammar, S., Bouzid, S., and Carpentier, R. (2011). Abolition of photosystem I cyclic electron flow in *Arabidopsis thaliana* following thermal stress. *Plant Physiol. Biochem.* 49, 235–243. doi: 10.1016/j.plaphy.2010.11.002
- Fan, J., Hu, Z., Xie, Y., Chan, Z., Chen, K., Amombo, E., et al. (2015). Alleviation of cold damage to photosystem II and metabolisms by melatonin in bermudagrass. *Front. Plant Sci.* 6:925. doi: 10.3389/fpls.2015.00925
- Fan, J., Ren, J., Zhu, W., Amombo, E., Fu, J., and Chen, L. (2014). Antioxidant responses and gene expression in bermudagrass under cold stress. *J. Am. Soc. Hortic. Sci.* 139, 699–705. doi: 10.1093/jxb/eru373
- Farooq, M., Wahid, A., and Lee, D. J. (2009). Exogenously applied polyamines increase drought tolerance of rice by improving leaf water status, photosynthesis and membrane properties. *Acta Physiol. Plant.* 31, 937–945. doi: 10.1007/s11738-009-0307-2
- Feuerstein, B. G., and Marton, L. J. (1989). Specificity and binding in polyamine/nucleic acid interactions. *The physiology of polyamines.* 1, 109–124.
- Filippou, P., Antoniou, C., Yelamanchili, S., and Fotopoulos, V. (2012). No loading: efficiency assessment of five commonly used application methods of sodium nitroprusside in medicago truncatula plants. *Plant Physiol. Biochem.* 60, 115–118. doi: 10.1016/j.plaphy.2012.07.026
- Galston, A. W., and Sawhney, R. K. (1990). Polyamines in plant physiology. *Plant Physiol.* 94, 406–410. doi: 10.1104/pp.94.2.406
- Gitelson, A. A., Gritz, Y., and Merzlyak, M. N. (2003). Relationships between leaf chlorophyll content and spectral reflectance and algorithms for non-destructive chlorophyll assessment in higher plant leaves. *J. Plant Physiol.* 160, 271–282. doi: 10.1078/0176-1617-00887
- Goyal, M., and Asthir, B. (2010). Polyamine catabolism influences antioxidative defense mechanism in shoots and roots of five wheat genotypes under high temperature stress. *Plant Growth Regul.* 60, 13–25. doi: 10.1007/s10725-009-9414-8
- Goyns, M. H. (1982). The role of polyamines in animal cell physiology. *J. Theor. Biol.* 97, 577–589. doi: 10.1016/0022-5193(82)90359-9
- Green, B. R., and Durnford, D. G. (1996). The chlorophyll-carotenoid proteins of oxygenic photosynthesis. *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 47, 685. doi: 10.1146/annurev.arplant.47.1.685
- Groppa, M. D., Tomaro, M. L., and Benavides, M. P. (2001). Polyamines as protectors against cadmium or copper-induced oxidative damage in sunflower leaf discs. *Plant Sci.* 161, 481–488. doi: 10.1016/S0168-9452(01)00432-0
- Hideg, É., Barta, C., Kálai, T., Vass, I., Hideg, K., and Asada, K. (2002). Detection of singlet oxygen and superoxide with fluorescent sensors in leaves under stress by photoinhibition or UV radiation. *Plant Cell Physiol.* 43, 1154–1164. doi: 10.1093/pcp/pcf145
- Hiscox, J. T., and Israelstam, G. (1979). A method for the extraction of chlorophyll from leaf tissue without maceration. *Can. J. Bot.* 57, 1332–1334. doi: 10.1139/b79-163
- Hoagland, D., and Arnon, D. (1950). *The Water-Culture Method for Growing Plants without Soil*. Berkeley, CA: University of California.
- Hu, L., Xiang, L., Li, S., Zou, Z., and Hu, X. H. (2016). The beneficial role of spermidine chlorophyll metabolism and D1 protein content in tomato seedlings under salinity-alkalinity stress. *Physiol. Plant.* 156, 468. doi: 10.1111/pp.12398
- Hu, Z., Fan, J., Chen, K., Amombo, E., Chen, L., and Fu, J. (2016). Effects of ethylene on photosystem II and antioxidant enzyme activity in bermudagrass under low temperature. *Photosyn. Res.* 128, 1–14. doi: 10.1007/s11120-015-0199-5
- Jiang, C. D., Gao, H. Y., and Zou, Q. (2001). Enhanced thermal energy dissipation depending on xanthophyll cycle and D1 protein turnover in iron-deficient maize leaves under high irradiance. *Photosynthetic.* 39, 269–274. doi: 10.1023/A:1013701224847
- Kruse, O., Zheleva, D., and Barber, J. (1997). Stabilization of photosystem two dimers by phosphorylation: implication for the regulation of the turnover of D1 protein. *FEBS Lett.* 408, 276–280. doi: 10.1016/S0014-5793(97)00439-0
- Kusano, T., Berberich, T. C., and Takahashi, Y. (2008). Polyamines: essential factors for growth and survival. *Planta* 228, 367–381. doi: 10.1007/s00425-008-0772-7
- Lee, S. H., Ahsan, N., Lee, K. W., Kim, D. H., Lee, D. G., Kwak, S. S., et al. (2007). Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses. *J. Plant Physiol.* 164, 1626–1638. doi: 10.1016/j.jplph.2007.01.003
- Leong, T. Y., and Anderson, J. M. (1984). Adaptation of the thylakoid membranes of pea chloroplasts to light intensities. I. Study on the distribution of chlorophyll-protein complexes. *Photosyn. Res.* 5, 105–115. doi: 10.1007/BF00028524
- Li, Z., Zhang, Y., Zhang, X., Peng, Y., Merewitz, E., Ma, X., et al. (2016). The alterations of endogenous polyamines and phytohormones induced by exogenous application of spermidine regulate antioxidant metabolism, metallothionein and relevant genes conferring drought tolerance in white clover. *Environ. Exp. Bot.* 124, 22–38. doi: 10.1016/j.envexpbot.2015.12.004
- Liu, A., Hu, Z., Bi, A., Fan, J., Gitau, M. M., Amombo, E., et al. (2016). Photosynthesis, antioxidant system and gene expression of bermudagrass in response to low temperature and salt stress. *Ecotoxicolog* 25, 1445–1457. doi: 10.1007/s10646-016-1696-9
- Liu, X., and Huang, B. (2000). Heat stress injury in relation to membrane lipid peroxidation in creeping bentgrass. *Crop Sci.* 40, 503–510. doi: 10.2135/cropsci2000.402503x
- Mallick, N., and Mohn, F. H. (2000). Reactive oxygen species: response of algal cells. *J. Plant Physiol.* 157, 183–193. doi: 10.1016/S0176-1617(00)80189-3
- Marcum, K. B. (1998). Cell membrane thermostability and whole-plant heat tolerance of Kentucky bluegrass. *Crop Sci.* 38, 1214–1218. doi: 10.2135/cropsci1998.0011183X003800050017x
- Martin-Tanguy, J. (2001). Metabolism and function of polyamines in plants: recent development (new approaches). *Plant Growth Regulat.* 34, 135–148. doi: 10.1023/A:1013343106574
- Mittler, R., Vanderauwera, S., Gollery, M., and Van, B. F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–. doi: 10.1016/j.tplants.2004.08.009
- Monk, L. S., Fagerstedt, K. V., and Crawford, R. M. M. (1989). Oxygen toxicity and superoxide dismutase as an antioxidant in physiological stress. *Physiol. Plant.* 76, 456–459. doi: 10.1111/j.1399-3054.1989.tb06219.x
- Mostofa, M. G., Yoshida, N., and Fujita, M. (2014). Spermidine pretreatment enhances heat tolerance in rice seedlings through modulating antioxidative and glyoxalase systems. *Plant Growth Regul.* 73, 31–44. doi: 10.1007/s10725-013-9865-9
- Murkowski, A. (2001). Heat stress and spermidine: effect on chlorophyll fluorescence in tomato plants. *Biol. Plant.* 44, 53–57. doi: 10.1023/A:1017966203859
- Paschalidis, K. A., and Roubelakis-Angelakis, K. A. (2005). Sites and regulation of polyamine catabolism in the tobacco plant. Correlations with cell division/expansion, cell cycle progression, and vascular development. *Plant Physiol.* 138, 2174–2184. doi: 10.1104/pp.105.063941
- Pfannschmidt, T. (2003). Chloroplast redox signals: how photosynthesis controls its own genes. *Trends Plant Sci.* 8, 33–41. doi: 10.1016/S1360-1385(02)00005-5
- Pospíšil, P. (2012). Molecular mechanisms of production and scavenging of reactive oxygen species by photosystem II. *Biochim. Biophysica Acta* 1817, 218–231. doi: 10.1016/j.bbabi.2011.05.017
- Roberts, D. R., Dumbroff, E. B., and Thompson, J. E. (1986). Exogenous polyamines alter membrane fluidity in bean leaves a basis for potential misinterpretation of their true physiological role. *Planta* 167, 395–401.
- Shi, H., and Chan, Z. (2014). Improvement of plant abiotic stress tolerance through modulation of the polyamine pathway. *J. Integr. Plant Biol.* 56, 114–121. doi: 10.1111/jipb.12128
- Shi, H., Ye, T., and Chan, Z. (2013). Comparative proteomic and physiological analyses reveal the protective effect of exogenous polyamines in the bermudagrass (*Cynodon dactylon*) response to salt and drought stresses. *J. Proteome Res.* 12:4951. doi: 10.1021/pr400479k
- Sigaud-Kutner, T. C., Pinto, E., Okamoto, O. K., Latorre, L. R., and Colepicolo, P. (2002). Changes in superoxide dismutase activity and photosynthetic pigment

- content during growth of marine phytoplankters in batch cultures. *Physiol. Plant.* 114, 566–571. doi: 10.1034/j.1399-3054.2002.1140409.x
- Smirnoff, N. (1998). Plant resistance to environmental stress. *Curr. Opin. Biotechnol.* 9, 214–219. doi: 10.1016/S0958-1669(98)80118-3
- Song, W. Y., Yang, H. C., Shao, H. B., Zheng, A. Z., and Brestic, M. (2014). The alleviative effects of salicylic acid on the activities of catalase and superoxide dismutase in malting barley (*Hordeum vulgare* L.) seedling leaves stressed by heavy metals. *Clean Soil Air Water* 42, 88–97. doi: 10.1002/clen.201200310
- Strasser, R. J., Tsimilli-Michael, M., and Srivastava, A. (2004). “Analysis of the chlorophyll a fluorescence transient,” in *Chlorophyll A Fluorescence. Advances in Photosynthesis and Respiration*, Vol. 19, eds G. C. Papageorgiou and Govindjee (Dordrecht: Springer), 321–362. doi: 10.1007/978-1-4020-3218-9_12
- Takahashi, S., Nakamura, T., Sakamizu, M., Van, W. R., and Yamasaki, H. (2004). Repair machinery of symbiotic photosynthesis as the primary target of heat stress for reef-building corals. *Plant Cell Physiol.* 45, 251–255. doi: 10.1093/pcp/pch028
- Tan, W., Meng, Q. W., Brestic, M., Olsovska, K., and Yang, X. (2011). Photosynthesis is improved by exogenous calcium in heat-stressed tobacco plants. *J. Plant Physiol.* 168, 2063–2071. doi: 10.1016/j.jplph.2011.06.009
- Tanaka, R., Koshino, Y., Sawa, S., Ishiguro, S., Okada, K., and Tanaka, A. (2001). Overexpression of chlorophyllide a oxygenase (CAO) enlarges the antenna size of photosystem-II in *Arabidopsis thaliana*. *Plant J.* 26, 365–373. doi: 10.1046/j.1365-313X.2001.2641034.x
- Tian, J., Wang, L. P., Yang, Y. J., Sun, J., and Guo, S. R. (2012). Exogenous spermidine alleviates the oxidative damage in cucumber seedlings subjected to high temperatures. *J. Am. Soc. Hortic. Sci.* 137, 11–19.
- Todorova, D., Sergiev, I., Alexieva, V., Karanov, E., Smith, A., and Hall, M. (2007). Polyamine content in *Arabidopsis thaliana*, (L.) heynh during recovery after low and high-temperature treatments. *Plant Growth Regul.* 51, 185–191. doi: 10.1007/s10725-006-9143-1
- Umena, Y., Kawakami, K., Shen, J.-R., and Kamiya, N. (2011). Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 angstrom. *Nature* 473, 55–65. doi: 10.1038/nature09913
- Vani, B., Saradhi, P. P., and Mohanty, P. (2001). Alteration in chloroplast structure and thylakoid membrane composition due to heat treatment of rice seedlings: correlation with the functional changes. *J. Plant Physiol.* 158, 583–592. doi: 10.1078/0176-1617-00260
- Xiang, L., Hu, L., Xu, W., Ai, Z., Liang, Z., and Hu, X. (2016). Exogenous γ -aminobutyric acid improves the structure and function of photosystem II in muskmelon seedlings exposed to salinity-alkalinity stress. *PLOS ONE* 11:e0164847. doi: 10.1371/journal.pone.0164847
- Xu, S., Li, J., Zhang, X., Wei, H., and Cui, L. (2006). Effects of heat acclimation pretreatment on changes of membrane lipid peroxidation, antioxidant metabolites, and ultrastructure of chloroplasts in two cool-season turfgrass species under heat stress. *Environ. Exp. Bot.* 56, 274–285. doi: 10.1016/j.envexpbot.2005.03.002
- Yamasato, A., Nagata, N., Tanaka, R., and Tanaka, A. (2005). The N-terminal domain of chlorophyllide a oxygenase confers protein instability in response to chlorophyll b accumulation in *Arabidopsis*. *Plant Cell* 17, 1585–1597. doi: 10.1105/tpc.105.031518
- Yoshioka, M., Uchida, S., Mori, H., Komayama, K., Ohira, S., Morita, N., et al. (2006). Quality control of photosystem II. Cleavage of reaction center D1 protein in spinach thylakoids by FtsH protease under moderate heat stress. *J. Biol. Chem.* 281, 21660–21669. doi: 10.1074/jbc.M602896200

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Zhang, Hu, Amombo, Wang, Xie and Fu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.