

The Association Between Physical Activity, Sitting Time, Sleep Duration, and Sleep Quality as Correlates of Presenteeism

Diana Guertler, Dipl Psych, Corneel Vandelanotte, PhD, Camille Short, PhD, Stephanie Alley, B BSc, Stephanie Schoeppe, M SSc, and Mitch J. Duncan, PhD

Objective: This study aims to examine the relationship of lifestyle behaviors (physical activity, work and non-work sitting time, sleep quality, and sleep duration) with presenteeism while controlling for sociodemographics, work- and health-related variables. **Methods:** Data were collected from 710 workers (aged 20 to 76 years; 47.9% women) from randomly selected Australian adults who completed an online survey. Linear regression was used to examine the relationship between lifestyle behaviors and presenteeism. **Results:** Poorer sleep quality (standardized regression coefficients [B] = 0.112; $P < 0.05$), suboptimal duration (B = 0.081; $P < 0.05$), and lower work sitting time (B = -0.086; $P < 0.05$) were significantly associated with higher presenteeism when controlling for all lifestyle behaviors. Engaging in three risky lifestyle behaviors was associated with higher presenteeism (B = 0.150; $P < 0.01$) compared with engaging in none or one. **Conclusions:** The results of this study highlight the importance of sleep behaviors for presenteeism and call for behavioral interventions that simultaneously address sleep in conjunction with other activity-related behaviors.

Presenteeism is an individual's loss of productivity at work because of physical and psychosocial conditions and illness.^{1,2} The economic cost of lost productivity because of presenteeism is higher than the cost of absenteeism, that is, being not at work because of illness.^{3,4} Furthermore, longitudinal research suggests that presenteeism may increase the likelihood of future absenteeism.⁵ To reduce presenteeism in the workplace, the associated economic and social burden effective interventions are required. To inform this process, a greater understanding of the factors that affect presenteeism is needed.

There is some evidence that poor lifestyle behaviors may adversely affect presenteeism.^{1,6,7} For example, studies have shown that low levels of physical activity,^{6,8-10} higher sitting time before and after work,⁶ sleep disorders¹¹⁻¹³ and poor sleep quality⁷ are associated with higher presenteeism. Nevertheless, the extent to which these behaviors affect presenteeism is still relatively unclear. This is because of the paucity of studies conducted, inconsistent findings

reported (in the case of physical activity^{14,15}), important aspects of behavior not examined (eg, examining sleep disorders¹¹⁻¹³ and sleep quality⁷ but not sleep duration), the tendency to examine these behaviors in isolation, as well as lack of control for other important factors, such as health issues.^{1,6-13}

Strong evidence shows that sleep duration and sleep quality are important for maintaining good physical and mental health.¹⁶⁻¹⁸ Compared with the prevalence of sleep disorders (insomnia, obstructive sleep apnea, and restless leg syndrome), which affects 8.9% of the population, suboptimal sleep duration (sleeping less than 7 hours and more than 8 hours) affects 35% to 60% of the population and poor sleep quality (difficulty falling asleep or remaining asleep¹⁹) affects 23% of the population.²⁰⁻²⁴ Yet, previous studies^{7,11-13} examining sleep behavior and presenteeism have predominantly focused on sleep disorders and rarely considered less severe sleep issues (such as too little or not enough sleep and difficulty falling asleep) that also affect on health and well-being and are more prevalent in the population.

The lack of studies examining the effect of these behaviors in conjunction with each other is particularly concerning, given that 58.5% of the population report having at least two unhealthy lifestyle behaviors simultaneously.²⁵ It may be that lifestyle behaviors have a greater effect on presenteeism than what is shown in the current literature focusing on single behaviors as having several unhealthy lifestyle behaviors increases the risk of poor health, chronic disease, and mortality.^{16,26-28} The association of physical activity, sitting behavior, and sleep with presenteeism is of particular importance. Nonoccupational sitting time is increasing,^{29,30} and changes to work environments and job requirements are influencing workers' sleep behaviors³¹ and have resulted in reduced levels of occupational physical activity while increasing the proportion of occupational sitting.³² This is illustrated by studies reporting that office workers spend approximately 66% of their workday sitting.³³

Furthermore, when examining the effect of lifestyle behaviors on presenteeism, it is important to take into account other factors that may influence this association. Particularly, health-related factors such as self-rated health should be considered given the strong interrelationships between lifestyle behaviors, self-rated health, and health outcomes.^{20,34,35}

Therefore, this study aims to examine associations of physical activity, sitting time, sleep duration, and sleep quality with presenteeism when adjusting for self-rated health, as well as examine the association between having multiple unhealthy lifestyle behaviors simultaneously and presenteeism.

METHODS

The Human Ethics Research Review Panel at the Central Queensland University provided ethical approval for the project (Project H12/06-126).

Participant Recruitment

Participants were members of the Australian Health and Social Science Panel study funded by the Institute for Health and Social Science Research at the Central Queensland University, Australia. Panel members were recruited between 2009 and 2012

From the Institute for Health and Social Science Research (Ms Guertler, Dr Vandelanotte, Dr Short, Ms Alley, Ms Schoeppe, and Dr Duncan), Centre for Physical Activity Studies, School of Health and Human Services, Central Queensland University, Bruce Highway, Rockhampton, QLD, Australia; Institute of Social Medicine and Prevention (Ms Guertler), University Medicine, Walther-Rathenau-Strasse; DZHK (German Centre for Cardiovascular Research) (Ms Guertler), Greifswald Partner Site, Greifswald, Germany; and School of Medicine & Public Health (Dr Duncan), Priority Research Centre for Physical Activity and Nutrition, Faculty of Health and Medicine, The University of Newcastle, University Drive, Callaghan, NSW, Australia.

Disclosure of funding: Dr Duncan is supported by a Future Leader Fellowship (ID 100029) from the National Heart Foundation of Australia.

There are no further funding sources to disclose.

Conflict of interest: The authors report no conflicts of interest.

Supplemental digital content is available for this article. Direct URL citation appears in the printed text and is provided in the HTML and PDF versions of this article on the journal's Web site (www.joem.org).

Address correspondence to: Diana Guertler, Dipl Psych, Institute of Social Medicine and Prevention, University Medicine, Walther-Rathenau-Strasse 48, 17475 Greifswald, Germany (diana.guertler@uni-greifswald.de).

Copyright © 2015 by American College of Occupational and Environmental Medicine

DOI: 10.1097/JOM.0000000000000355

using computer-assisted telephone interviewing. Panel members were randomly selected adults (aged 18 years and older) contactable via telephone across all states and territories of Australia. Further details on the recruitment methods for the Australian Health and Social Science Panel are available elsewhere.³⁶

In 2012, all panel members ($n = 3932$) were invited to a Web-based survey via e-mail with up to four reminders. A total of 1843 (46.9%) respondents completed the survey. Only respondents who were employed in any type of paid work ($n = 1073$) were included in analyses. Respondents were excluded if their body mass index (BMI) exceeded 50 ($n = 10$), if they reported any health condition that prevented them from increasing physical activity or decreasing sitting time ($n = 41$), or if they had missing data for any variables included in the analyses conducted for this study ($n = 312$).

MEASURES

Sociodemographic, Work-Related, and Health-Related Variables

Participants provided information on their sex, age, marital status, employment, educational level, and household income. Highest level of education was categorized into lower education (high school or less, TAFE—a provider of vocational nonbachelor education up to level of advanced diploma) or higher education (technical college and higher education including university bachelor degree or higher). Household income was dichotomized into less than \$1500 per week and \$1500 or more per week. Respondents were asked to report employment status (full-time, part-time, or casual worker) and primary working time (during the day, during the night, or during the day and night). Occupation was classified using standardized measures³⁷ into the following three categories similar to other research³⁸: professionals (managers and administrators, professionals, and associate professionals); white-collar workers (elementary, intermediate, and advanced clerical, sales, and service workers); and blue-collar workers (tradespersons, intermediate production, and transport workers, laborers, and related workers). Self-rated health was assessed with one question “Would you say that in general your health is excellent, very good, good, fair, poor?” from an existing measure of health-related quality of life—HRQOL-4.³⁹ Participants BMI was calculated from their self-reported height and weight (weight in kg/[height in m]²). The presence of a chronic health condition was assessed by asking respondents if they have at least one of the following diseases: coronary heart disease, hypertension, hypercholesterolemia, diabetes, chronic bronchitis, cancer, osteoporosis, osteoarthritis, rheumatoid arthritis, irritable bowel syndrome, celiac disease, food allergy/intolerance, Alzheimer disease, and dementia. Mental health was assessed with one question “Have you ever been diagnosed with or treated for any kind of mental health condition including depression or anxiety?” (response options “yes” or “no”).

Presenteeism

Presenteeism was quantified as reduced performance while at work using a question of the World Health Organization’s validated Health and Work Performance Questionnaire (HPQ).^{40,41} Participants were asked to rate their overall job performance on working days during the past 4 weeks from 0 (worst job performance) to 10 (performance of a top worker). Response options were inversely coded and expressed as percentage to ease interpretation resulting in a score ranging from 0 (no presenteeism) to 100 (maximal presenteeism).

Physical Activity

The Active Australia Survey (AAQ)⁴² was used to measure frequency and time spent performing walking, moderate and vigorous physical activity in the last week. In this study, two measures of

physical activity were examined. The first was a continuous measure of total minutes of physical activity, calculated as the sum of time participating in walking, moderate and vigorous physical activity (multiplied by two); this is consistent with guidelines for analysis and reporting of AAQ items.⁴² A second measure of activity defined as the accumulation of at least 150 minutes of activity in five or more sessions was also used to classify participants as insufficiently or sufficiently active. Agreement of the AAQ with other questionnaires regarding the proportion of participants categorized as sufficiently active (150 minutes per week including at least five sessions) was between 59.2% and 74.3%,⁴³ and reliability for total minutes of physical activity was also high (intraclass correlation coefficient = 0.59).⁴⁴

Sitting Behavior

Daily sitting time in the last week was assessed using the Workforce Sitting Questionnaire (WSQ).⁴⁵ The WSQ assesses sitting time (on weekdays and weekend days) separately for traveling to and from places, being at work, watching TV, using a computer at home, and doing other leisure activities. The WSQ was found to have a good test-retest reliability (intraclass correlation coefficient = 0.46 to 0.90) and criterion validity against accelerometry ($r = 0.18$ to 0.46).⁴⁵ Average daily sitting time at work was calculated by $([\text{time spent sitting while at work on workdays} * \text{the number of workdays}] + [\text{time spent sitting while at work on non-workdays} * \text{the number of non-workdays}]/7)$. Average daily non-work-related sitting time was calculated by $([\text{time spent sitting while traveling, watching TV, using a computer, and other leisure activities on workdays} * \text{the number of workdays}] + [\text{time spent sitting while traveling, watching TV, using a computer, and other leisure activities on non-workdays} * \text{the number of non-workdays}]/7)$.

Sleep Duration and Quality

Sleep duration was assessed using one question “During the past month, how many hours of sleep did you usually get each night? This may be quite different to the number of hours you spent in bed. (Enter total number of hours sleep per night).” Sleep quality during the past month was assessed with a four-point rating scale (1 = very good, 2 = fairly good, 3 = fairly bad, and 4 = very bad). These items were adopted from the Pittsburgh Sleep Quality Index, which has demonstrated good psychometric properties (Cronbach $\alpha = 0.83$; test-retest reliability = 0.85).⁴⁶

Statistical Analysis

Descriptive statistics (means, standard deviation, and/or proportions) were calculated for all variables. Univariate linear regression analyses including sociodemographic, work- and health-related variables from Table 2 as predictors and presenteeism as the outcome variable were performed to identify possible confounders for the relationship between lifestyle behaviors and presenteeism. The associations of physical activity, work and non-work sitting time, sleep quality, and sleep duration with presenteeism were examined using linear regression analyses for each separate lifestyle behavior adjusted for significant sociodemographic, work- and health-related confounders identified in step one (models 1a to 1e). The lifestyle behaviors were included as continuous predictor variables in the described models. Furthermore, because lifestyle behaviors are likely to influence each other, a linear regression analysis including all lifestyle behaviors in a single model and adjusted for all significant sociodemographic, work- and health-related confounders was performed (model 2).

For lifestyle behaviors that are significantly associated with presenteeism graphs were produced in Excel showing the predicted presenteeism scores for the different levels of the lifestyle behavior. Predicted presenteeism scores were derived from the regression equation using unstandardized regression coefficients and sample

TABLE 1. Description of Sociodemographics, Work- and Health-Related Variables, Lifestyle Behaviors, and Presenteeism (*n* = 710)

Variables	<i>n</i> (%), If Not Otherwise Stated
Sociodemographics	
Sex	
Men	340 (47.9)
Women	370 (52.1)
Age, yrs	
Mean (SD), range	51.0 (10.8), 20–76
Educational level	
Low	232 (32.7)
High	478 (67.3)
Marital status	
Single	64 (9.0)
Divorced, separated, widowed	75 (10.6)
Married, de facto	571 (80.4)
Income	
< \$1,500 per week	222 (31.3)
≥ \$1,500 per week	488 (68.7)
Work-related factors	
Employment status	
Full-time worker	445 (62.7)
Part-time worker	208 (29.3)
Casual worker	57 (8.0)
Occupation	
Professional	502 (70.7)
White collar	131 (18.5)
Blue collar	77 (10.9)
Working time	
During the day	606 (85.4)
During the night, or day and night	104 (14.7)
Health-related variables	
Body mass index	
Mean (SD), range	27.2 (5.0), 17–48
Normal (<25 kg/m ²)	257 (36.2)
Overweight/obese (≥25 kg/m ²)	453 (63.8)
General health	
Poor	3 (0.4)
Fair	68 (9.6)
Good	254 (35.8)
Very good	301 (42.4)
Excellent	84 (11.8)
Chronic diseases	
Yes	447 (63.0)
No	263 (37.0)
Mental health issue	
Yes	161 (22.7)
No	549 (77.3)
Lifestyle behaviors	
Physical activity	
Minutes mean (SD), median	364.0 (375.2), 240.0
Sufficient*	415 (58.5)
Insufficient	254 (35.8)
No activity	41 (5.8)

(continues)

TABLE 1. (Continued)

Variables	<i>n</i> (%), If Not Otherwise Stated
Sitting time	
Work-related mean (SD)	232.3 (141.1)
Non-work-related mean (SD)	388.1 (197.0)
<8 h	224 (31.5)
≥8 h	486 (68.5)
Sleep quality	
Very good	125 (17.6)
Fairly good	430 (60.6)
Fairly bad	137 (19.3)
Very bad	18 (2.5)
Sleep duration	
Mean (SD), range	6.9 (1.0), 3–12
<7 h	240 (33.8)
≥7 to <8 h	266 (37.5)
≥8 h	204 (28.7)
Number of risky lifestyle behaviors respondents engage in	
0 or 1 behaviors	97 (13.7)
2 behaviors	230 (32.4)
3 behaviors	265 (37.3)
4 behaviors	118 (16.6)
Presenteeism	
Loss of productivity in %	Mean (SD) 19.8 (14.9)

*The accumulation of at least 150 minutes of activity per week including at least five sessions.

means for covariates to control for sociodemographic, work- and health-related variables, as well as other lifestyle behaviors.

Since research suggested that both shorter and longer sleep duration may have adverse effects on health,⁴⁷ it was examined whether the relationship between sleep duration and presenteeism was non-linear. First, an augmented partial residuals plot⁴⁸ was produced to identify nonlinearity in the data. Second, to test for a nonlinear relationship, a squared term of sleep duration (after centering by the mean) was added to the regression model 1e, and change in model fit was explored using likelihood ratio test. Because the squared term was significant (Standardized regression coefficients [B] = 0.105; *P* < 0.005) and model fit was improved (likelihood ratio $\chi^2(1) = 8.54$; *P* < .005) compared with the model including only a linear association, we only report on results for models 1e and 2 while including the squared term in addition to the linear term of sleep duration. Variance inflation factors for model 2 were less than 1.6 for all lifestyle behaviors indicating no multicollinearity.

To examine the effect of multiple lifestyle behaviors on presenteeism, an index was created where each participant was allocated a single point for each of the following risky lifestyle behaviors they engaged in—insufficient physical activity (not accumulating at least 150 minutes of physical activity with at least five sessions of activity over 1 week⁴²), reporting sitting time of 8 or more hours a day,⁴⁹ reporting not very good sleep quality,¹⁸ and a sleep duration less than 7 hours or 8 or more hours.²⁵ Overall sitting time was included in the index because there are no separate recommendations for work and non-work sitting time. Behaviors were categorized according to established guidelines for the behavior (physical activity) or evidence that risk of poor health outcomes (eg, overall mortality) was increased based on that pattern of behavior. Because of the low number of participants engaging in none risky lifestyle behavior (2.0%), we collapsed participants reporting to engage in none

or one risky lifestyle behavior (13.7%) into one category. Linear regression analysis with the number of risky lifestyle behaviors as the dummy coded predictor and significant sociodemographics, work- and health-related variables as covariates (from univariate analysis) was performed to examine the association of multiple lifestyle behaviors and presenteeism. Standardized regression coefficients and standard errors were calculated. Statistical analysis was performed with Stata version 12 (StataCorp LP, College Station, TX). All models were evaluated using a significance level of $P < 0.05$.

RESULTS

Respondents

A description of the sample ($n = 710$) regarding sociodemographics, work- and health-related variables, as well lifestyle behaviors, and presenteeism is presented in Table 1.

Associations of Lifestyle Behaviors With Presenteeism

Results of univariate regression analyses on the association of sociodemographic, work- and health-related factors with presenteeism are presented in Table 2. There was a significant association between the following sociodemographic, work- and health-related factors and presenteeism: sex, marital status, employment status, occupation class, general health status, and mental health. Therefore, all the following analyses were adjusted for these factors.

Individual regression analyses with adjustment for the significant sociodemographic, work- and health-related variables (Table 3, models 1a to 1e) revealed a significant association of presenteeism with poor sleep quality ($B = 0.132$; $P < 0.001$). For sleep duration, both the linear ($B = -0.076$; $P < 0.05$) and the squared term

($B = 0.105$; $P < 0.01$) were significantly associated with presenteeism. Physical activity ($B = -0.054$; $P = 0.156$), work-related ($B = -0.057$; $P = 0.149$), and non-work-related sitting time ($B = 0.060$; $P = 0.106$) were not associated with presenteeism.

After controlling for all lifestyle behaviors and socio-demographic, work- and health-related variables (Table 3, model 2), poor sleep quality ($B = 0.112$; $P < 0.05$) and the squared term of sleep duration ($B = 0.081$; $P < 0.05$) remained significantly associated with presenteeism, whereas the linear term of sleep duration was no longer significant ($B = -0.023$; $P = 0.582$). Furthermore, work-related sitting was significantly associated with presenteeism in this model ($B = -0.086$; $P < 0.05$), whereas physical activity ($B = -0.057$; $P = 0.132$) and non-work-related sitting time ($B = 0.060$; $P = .116$) remained not associated with presenteeism. Figures 1 and 2 display the association of sleep quality and sleep duration with

TABLE 3. Association of Lifestyle Behaviors With Presenteeism

Variables	Presenteeism	
	Models 1a–1e ^a B (SE)	Model 2 ^b B (SE)
Physical activity	-0.054 (0.002)	-0.057 (0.001)
Work sitting time	-0.057 (0.004)	-0.086 (0.004)*
Non-work sitting time	0.060 (0.003)	0.060 (0.003)
Sleep quality ^c	0.132 (0.833)***	0.112 (0.065)*
Sleep duration ^d		
Sleep duration	-0.076 (0.525)*	-0.023 (0.601)
Sleep duration ²	0.105 (0.296)**	0.081 (0.301)*

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

^aModel 1, adjusted for sex, marital status, employment status, occupation, general health, and mental health.

^bModel 2, same as model 1 but additionally adjusted for all other lifestyle behaviors.

^cCoded from 1 "very good" to 4 "very bad."

^dThe effect of sleep duration on presenteeism is represented by a linear and a squared term as adding the squared term was shown to improve model fit.

B, standardized regression coefficients; SE, standard error.

TABLE 2. Univariate Associations of Sociodemographics, Work- and Health-Related Variables With Presenteeism

Variables	Presenteeism B (SE)
Sociodemographics	
Sex (0 male; 1 female)	-0.180 (1.100)***
Age, yrs (continuous)	-0.060 (0.052)
Educational level (0 no tertiary; 1 tertiary)	0.024 (1.191)
Marital status (0 single, divorced, widowed; 1 married, de facto)	-0.103 (1.401)**
Income (0 <1500\$; 1 ≥ 1500\$ per week)	0.068 (1.203)
Work-related factors	
Employment status	
Full-time	Reference
Part-time	-0.080 (1.248)*
Casual	-0.008 (2.090)
Occupation	
Professional	Reference
White collar	-0.090 (1.455)*
Blue collar	-0.060 (1.815)
Working time (0 night, night and day; 1 day)	-0.018 (1.580)
Health-related variables	
BMI (continuous)	0.044 (0.112)
General health (continuous; 1 excellent to 5 poor)	0.187 (0.656)***
Chronic diseases (0 none; 1 yes)	0.007 (1.157)
Mental health issue (0 none; 1 yes)	-0.099 (1.328)**

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

B, standardized regression coefficients; BMI, body mass index; SE, standard error.




FIGURE 1. Association between reported sleep quality and presenteeism. Note: Predicted values based on model 2 adjusted for sex, marital status, employment status, occupation, general health and mental health, physical activity, work-related sitting time, non-work-related sitting time, and sleep duration.

presenteeism on the basis of model 2; the pattern of association in these models is similar to that of models 1d and 1e. Figure 3 illustrates the association of work-related sitting time with presenteeism on the basis of model 2.

Table 4 shows the association between presenteeism and the number of risky lifestyle behaviors participants engage in. Engaging in three risky lifestyle behaviors was associated with a significantly higher presenteeism score ($B = 0.150$; $P < 0.01$) compared with engaging in none or one risky lifestyle behavior. There were no significant associations with presenteeism scores when engaging in two ($B = 0.050$; $P = 0.368$) or four ($B = 0.067$; $P = 0.188$) risky lifestyle behaviors compared with engaging in none or one risky behavior.




FIGURE 2. Association between reported hours of sleep and presenteeism. Note: Predicted values based on model 2 including both sleep duration and sleep duration², and adjusted for sex, marital status, employment status, occupation, general health and mental health, physical activity, work-related sitting time, non-work-related sitting time, and sleep quality.




FIGURE 3. Association between reported work-related sitting time and presenteeism. Note: Predicted values based on model 2, adjusted for sex, marital status, employment status, occupation, general health, mental health, physical activity, non-work-related sitting time, sleep quality, and sleep duration.

TABLE 4. Association Between Number of Risky Lifestyle Behaviors Respondents Engage in and Presenteeism^a

Number of Risky Lifestyle Behaviors ^b Respondents Engage in	Presenteeism B (SE)
0 or 1	Reference
2	0.050 (1.743)
3	0.150 (1.725)*
4	0.067 (2.036)

* $P < 0.01$.

^aAdjusted for sex, marital status, employment status, occupation, general health, and mental health.

^bIncluding physical activity, sitting time, sleep quality, and a sleep duration. B, standardized regression coefficients; SE, standard error.

DISCUSSION

Results of this study demonstrate that poor sleep quality and suboptimal sleep duration are associated with higher presenteeism in workers when accounting for sociodemographics, work- and health-related variables, as well as for other lifestyle behaviors. Furthermore, presenteeism was heightened when respondents engaged in three risky lifestyle behaviors compared with engaging in none or only one risky behavior.

This study extends previous studies on the association of presenteeism and sleep disorders^{16–18} by concurrently examining the role of sleep quality and sleep duration and by controlling for a broad range of factors. As depression and chronic disease presence are related to both presenteeism and suboptimal sleep durations,^{50,51} depression, chronic diseases, and sleep quality were adjusted for in our analysis. This allowed us to more clearly examine the relationship between sleep duration, sleep quality, and presenteeism. Analyses also adjusted for physical activity, which is important as physical activity and sleep behaviors are interrelated.⁵² In line with previous studies,^{7,53} this study showed that poorer sleep quality was associated with higher presenteeism, and that this relationship remains when accounting for sociodemographics, work- and health-related variables, and other lifestyle behaviors. Furthermore, the squared term of sleep duration was significantly associated with presenteeism. This indicates that medium sleep duration is associated with lower presenteeism compared with shorter and longer sleep duration. In this study, this medium sleep duration was broadly comparable with the sleep duration suggested by several health agencies.⁵⁴ This is in line with previous studies showing adverse health effects of both shorter and longer sleep durations.^{16,23} These findings are important, given that 22% of the employees in this study reported fairly bad or very bad sleep quality, and further 62.5% reported sleep durations less than 7 hours or 8 or more hours. Other potential reasons for these associations may be due to impaired cognitive performance associated with shorter or longer sleep durations; however, this was not assessed in this study.^{55,56} Further research examining this issue is needed including the use of study designs that can limit the effect of any potential bidirectional relationships.

To date, workplace health promotion programs have typically focused on improving physical activity, smoking, and nutrition. Although these have had some success reducing presenteeism,⁵³ results from this study suggest that the inclusion of sleep behaviors may further enhance intervention outcomes. Notwithstanding that brief educational interventions in sleep hygiene have been found to be effective in enhancing sleep quality,^{57,58} sleep hygiene has rarely been included in health promotion programs. This may be due to the fact that sleep disturbances were traditionally treated pharmacological, and only recently studies started to focus on the benefits of nonpharmacological treatment.⁵⁹

When examining the combined effect of lifestyle behaviors on presenteeism, this study showed that engaging in three risky lifestyle behaviors is associated with higher presenteeism compared with engaging in none or one risky behavior. Thus, interventions targeting multiple lifestyle behaviors simultaneously are likely to be more useful for reducing presenteeism compared with interventions that focus on single health behaviors, as many previous interventions have done.⁶⁰ Therefore, employers should encourage their workers to a healthier lifestyle, including being physically active, reducing sitting time, and enhancing sleep behaviors. This is of particular importance as a supportive work environment was found to be associated with lower presenteeism.¹⁰

When adjusting for sociodemographic, work- and health-related factors, there was no association between physical activity and sitting time with presenteeism in this study. Previous studies^{6,8-10} have reported a relationship of physical activity and sitting time with presenteeism. Nevertheless, these studies have not controlled for health-related variables, which are known to influence presenteeism. This may suggest that the association between these lifestyle behaviors and presenteeism is partially mediated through health variables. This is supported by subsequent analysis in this study (see Supplemental Digital Content Table 1, <http://links.lww.com/JOM/A182>), which demonstrated that low physical activity and higher non-work sitting time were indeed associated with presenteeism when only controlling for sociodemographic variables. This analysis was not reported because we deemed it important to account for health-related variables in the association between lifestyle behaviors and presenteeism. Nevertheless, this is in line with studies showing that both high physical activity and low sitting time are associated with better health.^{27,28,61}

In relation to sitting time, two aspects should be considered when evaluating the results of this study. First, as there is a relatively high proportion of casual and part-time workers in our sample, this may have affected the results because casual and part-time workers are likely to differ from the full-time workers in their sitting time.⁶² Second, after controlling for other lifestyle behaviors (model 2), higher work-related sitting was significantly associated with lower presenteeism. Given previous studies demonstrating higher sedentary time was related to heightened presenteeism,⁶ this is unexpected and may be due to several factors. Differences in the behaviors being measured, sedentary behavior versus sitting time, and the methods used to quantify them may contribute to differences between studies. The high number of professional and white-collar employees in the sample (89%), many of whom are required to be seated to conduct their work^{63,64} regardless of their presenteeism, may have confounded this relationship. Study design prohibited exploring this association in depth; however, when examining this association separately by occupation (data not shown), high work-related sitting time was associated with lower presenteeism only in professionals. Thus, it may be useful to examine this in future studies to better understand this association.

Some methodological limitations have been focused in this study. First, as we used cross-sectional data, we cannot provide information on the causality of the observed associations. Even though it seems reasonable to assume that lifestyle behaviors lead to differences in presenteeism, there is evidence from prospective studies suggesting that presenteeism predicts future health as well.⁶⁵ Second, even though we found that engaging in three risky lifestyle behaviors is associated with higher presenteeism, there was no heightened presenteeism for engaging in four risky lifestyle behaviors compared with engaging in none or one. This may be due to a lack of power to detect differences between groups because only 118 respondents (16.6%) reported engaging in four risky health behaviors. Third, we assessed presenteeism using a self-report measure (the HPQ), which may have led to bias from memory effects or social desirability. Nevertheless, respondents' reported score on the HPQ is in line with

previous studies, showing a similar score for general workers⁶⁶ and a lower score for workers with medical conditions⁶⁷⁻⁶⁹ indicating this sample is comparable with other published data. Finally, the HPQ did not assess productivity loss because of health problems, so we may have assessed productivity loss because of other reasons as well. Although computer-based tracking systems, for work productivity, exist, the objective measurement of work productivity remains challenging when looking at measurement of work quality and in occupations where discrete endpoints (eg, produced pieces, finished calls) are lacking.⁷⁰ Strengths of this study include examining the association of a range of lifestyle behaviors and presenteeism while taking into account health-related variables and the effect of multiple lifestyle behaviors.

CONCLUSIONS

This study demonstrated that higher presenteeism is associated with poor sleep quality and suboptimal sleep duration even after controlling for health-related variables and other lifestyle behaviors. Presenteeism was heightened for employees engaging in three risky lifestyle behaviors compared with engaging in none or one. Hence, the outcomes of this study suggest that encouraging employees to be more physically active, reducing sitting time, and enhancing sleep behavior can reduce effects of presenteeism. To reduce presenteeism associated costs, employers should consider implementing workplace programs to improve multiple health behaviors in employees.

ACKNOWLEDGMENTS

The authors thank the Australian Health and Social Science (AHSS) Panel study founded by the Institute for Health and Social Science Research (IHSSR) at the Central Queensland University, Australia, for providing data used in this study.

REFERENCES

1. Brown HE, Gilson ND, Burton NW, Brown WJ. Does physical activity impact on presenteeism and other indicators of workplace well-being? *Sports Med.* 2011;41:249–262.
2. Mattke S, Balakrishnan A, Bergamo G, Newberry SJ. A review of methods to measure health-related productivity loss. *Am J Manag C.* 2007;13:211–217.
3. Collins JJ, Baase CM, Sharda CE, et al. The assessment of chronic health conditions on work performance, absence, and total economic impact for employers. *J Occup Environ Med.* 2005;47:547–557.
4. The Sainsbury Centre for Mental Health. Mental health at work: developing the business case. Policy paper 8. Available at: http://www.centreformentalhealth.org.uk/pdfs/mental_health_at_work.pdf. Published 2007. Accessed February 27, 2014.
5. Janssens H, Clays E, De Clercq B, De Bacquer D, Braeckman L. The relation between presenteeism and different types of future sickness absence. *J Occup Health.* 2013;55:132–141.
6. Brown HE, Ryde GC, Gilson ND, Burton NW, Brown WJ. Objectively measured sedentary behavior and physical activity in office employees. *J Occup Environ Med.* 2013;55:945–953.
7. Zammit GK, Joish VN, Kong MC, Balkrishnan R, Lerner D, Rosekind M. Impact of nighttime awakenings on worker productivity and performance. *J Occup Environ Med.* 2010;52:513–518.
8. Arvidson E, Börjesson M, Ahlborg G Jr, Lindegård A, Jonsdottir IH. The level of leisure time physical activity is associated with work ability—a cross sectional and prospective study of health care workers. *BMC Public Health.* 2013;13:1–6.
9. Williden M, Schofield G, Duncan S. Establishing links between health and productivity in the New Zealand workforce. *J Occup Environ Med.* 2012;54:545–550.
10. Merrill RM, Aldana SG, Pope JE, Anderson DR, Coberley CR, Subcommittee RW. Presenteeism according to healthy behaviors, physical health, and work environment. *Popul Health Manag.* 2012;15:293–301.
11. Bolge S, Doan J, Kannan H, Baran R. Association of insomnia with quality of life, work productivity, and activity impairment. *Qual Life Res.* 2009;18:415–422.
12. Daley M, Morin CM, LeBlanc M, Grégoire JP, Savard J, Baillargeon L. Insomnia and its relationship to health-care utilization, work absenteeism, productivity and accidents. *Sleep Med.* 2009;10:427–438.

13. Swanson LM, Arnedt JT, Rosekind MR, Belenky G, Balkin TJ, Drake C. Sleep disorders and work performance: Findings from the 2008 National Sleep Foundation Sleep in America Poll. *J Sleep Res.* 2011;20:487–494.
14. Cash SW, Beresford SAA, Henderson JA, et al. Dietary and physical activity behaviours related to obesity-specific quality of life and work productivity: Baseline results from a worksite trial. *Brit J Nutr.* 2012;108:1134–1142.
15. Block G, Sternfeld B, Block CH, et al. Development of Alive! (A Lifestyle Intervention Via Email), and its effect on health-related quality of life, presenteeism, and other behavioral outcomes: randomized controlled trial. *J Med Internet Res.* 2008;10:e43.
16. Alvarez GG, Ayas NT. The impact of daily sleep duration on health: a review of the literature. *Prog Cardiovasc Nurs.* 2004;19:56–59.
17. Liu Y, Croft J, Wheaton A, et al. Association between perceived insufficient sleep, frequent mental distress, obesity and chronic diseases among US adults, 2009 Behavioral Risk Factor Surveillance System. *BMC Public Health.* 2013;13:84.
18. Doi Y, Minowa M, Tango T. Impact and correlates of poor sleep quality in Japanese white-collar employees. *Sleep.* 2003;26:467–471.
19. Buysse DJ, Ancoli-Israel S, Edinger JD, Lichstein KL, Morin CM. Recommendations for a standard research assessment of insomnia. *Sleep.* 2006;29:1155–1173.
20. Duncan MJ, Kline CE, Vandelanotte C, Sargent C, Rogers NL, Di Milia L. Cross-sectional associations between multiple lifestyle behaviors and health-related quality of life in the 10,000 Steps Cohort. *PLoS One.* Available at: <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0094184>. Published 2014. Accessed August 2, 2014.
21. Krueger PM, Friedman EM. Sleep duration in the United States: a cross-sectional population-based study. *Am J Epidemiol.* 2009;169:1052–1063.
22. Shankar A, Charumathi S, Kalidindi S. Sleep duration and self-rated health: the National Health Interview Survey 2008. *Sleep.* 2011;34:1173–1177.
23. Patel SR, Malhotra A, Gottlieb DJ, White DP, Hu FB. Correlates of long sleep duration. *Sleep.* 2006;29:881–889.
24. Sleep Health Foundation. Re-awakening Australia. The economic cost of sleep disorders in Australia. Available at: <http://www.sleephealthfoundation.org.au/pdfs/news/Reawakening%20Australia.pdf>. Published 2011. Accessed February 2, 2014.
25. Martínez-Gómez D, Guallar-Castillón P, León-Muñoz LM, López-García E, Rodríguez-Artalejo F. Combined impact of traditional and non-traditional health behaviors on mortality: a national prospective cohort study in Spanish older adults. *BMC Med.* 2013;11:1–10.
26. Bauman AE. Updating the evidence that physical activity is good for health: an epidemiological review 2000–2003. *J Sci Med Sport.* 2004;7:6–19.
27. Matthews CE, George SM, Moore SC, et al. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr.* 2012;95:437–445.
28. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sport Exerc.* 2009;41:998–1005.
29. Duncan MJ, Vandelanotte C, Caperchione C, Hanley C, Mummery WK. Temporal trends and in relationships between screen time, physical activity, overweight and obesity. *BMC Public Health.* 2012;12:1060.
30. Chau JY, Merom D, Grunseit A, Rissel C, Bauman AE, van der Ploeg HP. Temporal trends in non-occupational sedentary behaviours from Australian time use surveys 1992, 1997 and 2006. *Int J Behav Nutr Phy.* 2012;9:76–83.
31. Akerstedt T, Fredlund P, Gillberg M, Jansson B. Work load and work hours in relation to disturbed sleep and fatigue in a large representative sample. *J Psychosom Res.* 2002;53:585–588.
32. Church TS, Thomas DM, Tudor-Locke C, et al. Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PLoS One.* 2011;6:1–7.
33. Ryan CG, Dall PM, Granat MH, Grant PM. Sitting patterns at work: objective measurement of adherence to current recommendations. *Ergonomics.* 2011;54:531–538.
34. Li C, Ford ES, Mokdad AH, Balluz LS, Brown DW, Giles WH. Clustering of cardiovascular disease risk factors and health-related quality of life among US adults. *Value Health.* 2008;11:689–699.
35. Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: a systematic review and meta-analysis. *Prev Med.* 2012;55:163–170.
36. Population Research Laboratory Institute for Health and Social Science Research. The correlates of sedentary behavior: an examination of health and sitting time. *An Online Survey of Members of the CQ University Australian Health and Social Science Panel. Final Sampling and Technical Report.* Rockhampton, Australia: Central Queensland University; 2012.
37. Australian Bureau of Statistics. *Australian Standard Classification of Occupations (ASCO).* 2nd ed. Canberra, Australia: Australian Government Publishing Service; 1997.
38. Duncan MJ, Badland HM, Mummery WK. Physical activity levels by occupational category in non-metropolitan Australian adults. *J Phys Act Health.* 2010;7:718–723.
39. Centers for Disease Control and Prevention. Measuring healthy days. Available at: <http://www.cdc.gov/hrqol/pdfs/mhd.pdf>. Published 2000. Accessed March 15, 2014.
40. Kessler RC, Ames M, Hymel PA, et al. Using the World Health Organization Health and Work Performance Questionnaire (HPQ) to evaluate the indirect workplace costs of illness. *J Occup Environ Med.* 2004;46:S23–S37.
41. Kessler RC, Barber C, Beck A, et al. The World Health Organization Health and Work Performance Questionnaire (HPQ). *J Occup Environ Med.* 2003;45:156–174.
42. Australian Institute of Health and Welfare. The Active Australia Survey: a guide and manual for implementation, analysis and reporting. Available at: <http://www.aihw.gov.au/publication-detail/?id=6442467449>. Published 2003. Accessed March 10, 2014.
43. Brown W, Bauman A, Chey T, Trost S, Mummery K. Comparison of surveys used to measure physical activity. *Aus NZ J Public Health.* 2004;28:128–134.
44. Brown WJ, Trost SG, Bauman A, Mummery K, Owen N. Test-retest reliability of four physical activity measures used in population surveys. *J Sci Med Sport.* 2004;7:205–215.
45. Chau JY, van der Ploeg H, Dunn S, Kurko J, Bauman AE. A tool for measuring workers' sitting time by domain: the Workforce Sitting Questionnaire. *Brit J Sport Med.* 2011;45:1216–1222.
46. Buysse DJ, Reynolds CF III, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psych Res.* 1989;28:193–213.
47. Gallicchio L, Kalesan B. Sleep duration and mortality: a systematic review and meta-analysis. *J Sleep Res.* 2009;18:148–158.
48. Mallows C. Augmented partial Residuals. *Technometrics.* 1986;28:313–319.
49. van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med.* 2012;172:494–500.
50. Lerner D, Henke RM. What does research tell us about depression, job performance, and work productivity? *J Occup Environ Med.* 2008;50:401–410.
51. Kaneita Y, Ohida T, Uchiyama M, et al. The relationship between depression and sleep disturbances: a Japanese nationwide general population survey. *J Clin Psychiatr.* 2006;67:196–203.
52. Loprinzi P, Cardinal BJ. Association between objectively-measured physical activity and sleep, NHANES 2005–2006. *Ment Health Phys.* 2011;4: 65–69.
53. Barber L, Grawitch MJ, Munz DC. Are better sleepers more engaged workers? A self-regulatory approach to sleep hygiene and work engagement. *Stress Health.* 2013;29:307–316.
54. US Department of Health & Human Services. National Heart, Lung, and Blood Institute. How much sleep is enough? Available at: <http://www.nhlbi.nih.gov/health/health-topics/topics/sdd/howmuch.html>. Published 2012. Accessed August 8, 2014.
55. Ferrie JE, Shipley MJ, Akbaraly TN, Marmot MG, Kivimaki M, Singh-Manoux A. Change in sleep duration and cognitive function: findings from the Whitehall II Study. *Sleep.* 2011;34:565–573.
56. Xu L, Jiang CQ, Lam TH, et al. Short or long sleep duration is associated with memory impairment in older Chinese: the Guangzhou Biobank Cohort Study. *Sleep.* 2011;34:575–580.
57. Kaku A, Nishinoue N, Takano T, et al. Randomized controlled trial on the effects of a combined sleep hygiene education and behavioral approach program on sleep quality in workers with insomnia. *Ind Health.* 2012;50:52–59.
58. Chen P-H, Kuo H-Y, Chueh K-H. Sleep hygiene education: efficacy on sleep quality in working women. *J Nurs Res.* 2010;18:283–289.
59. Montgomery P, Dennis J. A systematic review of non-pharmacological therapies for sleep problems in later life. *Sleep Med Rev.* 2004;8:47–62.
60. Rongen A, Robroek SJW, Lenthe FJV, Burdorf A. Workplace health promotion: a meta-analysis of effectiveness. *Am J Prev Med.* 2013;44:406–415.
61. Danaei G, Ding EL, Mozaffarian D, et al. The Preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med.* 2009;6:1–23.
62. Vandelanotte C, Duncan MJ, Short C, et al. Associations between occupational indicators and total, work-based and leisure-time sitting: a cross-sectional study. *BMC Public Health.* 2013;13:1110.

63. Mummery W, Schofield G, Steele R, Eakin E, Brown W. Occupational sitting time and overweight and obesity in Australian workers. *Am J Prev Med.* 2005;29:91–97.
64. Miller R, Brown W. Steps and sitting in a working population. *Int J Behav Med.* 2004;11:219–224.
65. Taloyan M, Aronsson G, Leineweber C, Hanson LM, Alexanderson K, Westerlund H. Sickness presenteeism predicts suboptimal self-rated health and sickness absence: a nationally representative study of the Swedish working population. *PLoS One.* Available at: <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0044721>. Published 2012. Accessed August 8, 2014.
66. Shi Y, Sears LE, Coberley CR, Pope JE. Classification of individual well-being scores for the determination of adverse health and productivity outcomes in employee populations. *Popul Health Manag.* 2013;16:90–98.
67. Vries H, Reneman M, Groothoff J, Geertzen J, Brouwer S. Self-reported work ability and work performance in workers with chronic nonspecific musculoskeletal pain. *J Occup Rehab.* 2013;23:1–10.
68. Osilla KC, dela Cruz E, Miles JN, et al. Exploring productivity outcomes from a brief intervention for at-risk drinking in an employee assistance program. *Addict Behav.* 2010;35:194–200.
69. Furukawa TA, Horikoshi M, Kawakami N, et al. Telephone cognitive-behavioral therapy for subthreshold depression and presenteeism in workplace: a randomized controlled trial. *PLoS One.* 2012;7:1–9.
70. Prasad M, Wahlqvist P, Shikar R, Shih Y-CT. A review of self-report instruments measuring health-related work productivity: a patient-reported outcomes perspective. *Pharmacoeconomics.* 2004;22:225–244.