



The Auxin Signaling Repressor IAA8 Promotes Seed Germination Through Down-Regulation of *ABI3* Transcription in *Arabidopsis*

Shah Hussain¹, Sun Ho Kim¹, Sunghwa Bahk¹, Akhtar Ali², Xuan Canh Nguyen³, Dae-Jin Yun² and Woo Sik Chung^{1*}

¹ Division of Applied Life Science (BK21 Plus program), Plant Molecular Biology and Biotechnology Research Center, Gyeongsang National University, Jinju, South Korea, ² Department of Biomedical Science and Engineering, Konkuk University, Seoul, South Korea, ³ Faculty of Biotechnology, Vietnam National University of Agriculture, Hanoi, Vietnam

OPEN ACCESS

Edited by:

Stephan Pollmann,
National Institute of Agricultural and
Food Research and Technology,
Spain

Reviewed by:

Shucui Wang,
Linyi University, China
He Junna,
China Agricultural University
(CAU), China

*Correspondence:

Woo Sik Chung
chungws@gnu.ac.kr

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 04 September 2019

Accepted: 24 January 2020

Published: 20 February 2020

Citation:

Hussain S, Kim SH, Bahk S, Ali A,
Nguyen XC, Yun D-J and Chung WS
(2020) The Auxin Signaling Repressor
IAA8 Promotes Seed Germination
Through Down-Regulation of *ABI3*
Transcription in *Arabidopsis*.
Front. Plant Sci. 11:111.
doi: 10.3389/fpls.2020.00111

Seed germination is a complex biological process controlled by various regulators, including phytohormones. Among these, abscisic acid and gibberellic acid inhibit and promote seed germination, respectively. Many studies have addressed the biological roles of auxin in plant growth and development, but very few have considered its role in seed germination. Here, we identified a novel function of the auxin signaling repressor Aux/IAA8 during seed germination. The *IAA8* loss-of-function mutant *iaa8-1* exhibited delayed seed germination. The phenotype of *iaa8-1* was restored by ectopic expression of *IAA8*. Interestingly, *IAA8* accumulated to high levels during seed germination, which was achieved not only by increased protein synthesis but also by the stabilization of *IAA8* protein. We also showed that *IAA8* down-regulates the transcription of *ABSCISIC ACID INSENSITIVE3 (ABI3)*, a negative regulator of seed germination. Our study, thus strongly suggest that the auxin signaling repressor *IAA8* acts as a positive regulator of seed germination in *Arabidopsis thaliana*.

Keywords: *ABI3*, *Arabidopsis*, auxin, *IAA8*, protein stability, seed germination

INTRODUCTION

Seed maturation is the final stage of embryogenesis. The embryo becomes protected by a hard outer cover of dead tissue, the testa, underneath which the endosperm is deposited (Debeaujon et al., 2000). Upon germination, the testa ruptures, allowing the embryonic axis to protrude at the proper time to ensure propagation to the next generation (Piskurewicz et al., 2008). Germination occupies a critical position in the life cycle of seed plants, converting dormant seeds into active seedlings. This process is strictly regulated by genetic and environmental factors (Chen et al., 2008; Bassel et al., 2011; Liu et al., 2013). From an agronomic point of view, uniform germination is required for high crop yield. However, lack of dormancy may lead to pre-harvest sprouting (Gubler et al., 2005), resulting in decreased seed longevity (Rajjou and Debeaujon, 2008). Therefore, it is important to

Abbreviations: ABA, Abscisic Acid; *ABI3*, *ABSCISIC ACID INSENSITIVE3*; ARF, Auxin Response Factor; Aux/IAA, Auxin/Indole-3-Acetic Acid; GA, Gibberellic Acid; ROS, Reactive Oxygen Species.

maintain seeds in a dormant condition until suitable timing for germination is established (Shu et al., 2016).

Releasing dormancy is a prerequisite to germination that can be induced by various external and internal stimuli (Arc et al., 2012). Germination is induced most commonly through the imbibition of water at a species-specific temperature. Imbibition of dry seeds activates a series of events (Bewley, 1997) including oxidation, degradation, and mobilization of accumulated reserve components (Penfield et al., 2005a). Reactive oxygen species (ROS) also accumulate in seeds to a level that positively regulates seed germination (Leymarie et al., 2012). ROS are proposed to up-regulate abscisic acid (ABA) catabolism and promote gibberellic acid (GA) biosynthesis, thereby maintaining a dynamic balance between ABA and GA during seed germination (Liu et al., 2010).

Various endogenous regulators including phytohormones and associated proteins control the transition from dormant to active, germinating seeds (Shu et al., 2016; Nee et al., 2017). Among phytohormones, ABA is considered to maintain seed dormancy and inhibit seed germination (Hubbard et al., 2010). Meanwhile, GA, brassinosteroids, ethylene, and cytokinin promote seed germination (Steber and McCourt, 2001; Wang et al., 2011). Several ABA signaling genes such as *ABSCISIC ACID INSENSITIVE3* (*ABI3*), *ABI4*, and *ABI5* was identified as negative regulators of seed germination. Among these, *ABI3* acts as a major downstream components of ABA signaling (Giraudat et al., 1992; Bentsink and Koornneef, 2008). The transcript levels of *ABI3* are high in dormant seeds but low after germination (Liu et al., 2013). The antagonistic roles of ABA and GA in dormancy and germination have been studied extensively (Shu et al., 2016). However, the relationship between ABA and auxin in seed germination remains unresolved.

Auxin is involved in almost every aspect of plant growth and development (Zhao, 2010); however, its role in seed germination is still unknown. Rapid turnover of auxin/indole-3-acetic acid (Aux/IAA) repressor proteins is required (Overvoorde et al., 2005) to trigger auxin-mediated transcriptional activation (Tiwari et al., 2003). These short-lived transcriptional repressors are mainly targeted for degradation by polyubiquitination (Kepinski and Leyser, 2005; Overvoorde et al., 2005; Gilkerson et al., 2015). Recent reports have suggested that auxin inhibits seed germination in an ABA dependent manner. For instance, seeds of auxin over-producing transgenic plants (*iaaM-OX*) display arrested germination (Liu et al., 2013). By contrast, mutants with reduced levels of auxin (*yuc1/yuc6*) showed enhanced seed germination rate (Liu et al., 2013). Similarly, auxin signaling mutants with impaired ability to degrade Aux/IAA repressor proteins show enhanced germination (Kepinski and Leyser, 2005; Liu et al., 2013). Gain-of-function mutants of Aux/IAA repressors, such as *IAA7/AXR2* and *IAA17/AXR3*, also display enhanced germination rate (Liu et al., 2013). The underlying genetic and biochemical evidence suggests that *ABI3* is the downstream regulatory component of auxin-mediated seed dormancy (Belin et al., 2009; Liu et al., 2013). These molecular observations imply that inhibition of auxin signaling *via* Aux/IAA might be responsible for promoting seed germination.

Although the gain-of-function mutation of *IAA8* negatively regulates flower development (Wang et al., 2013), the loss-of-function mutant show no visible developmental phenotype (Overvoorde et al., 2005). To decipher the molecular mechanism explaining how auxin signaling regulates seed germination, we characterized the biological role of *IAA8* during seed germination. We provide evidence that *IAA8* protein accumulates during seed germination, promoting germination through the inhibition of *ABI3* transcription.

MATERIALS AND METHODS

Plant Material and Growth Conditions

Arabidopsis thaliana ecotype Columbia (Col-0) was used in all experiments. T-DNA insertion mutants *iaa8-1* (CS25210) and *iaa8-2* (SALK_202296) were obtained from SALK. T-DNA insertion was confirmed by genotyping PCR using *IAA8* gene-specific and T-DNA border primers (listed in **Supplementary Table 1**). The transcript was confirmed by semi-quantitative RT-PCR using *IAA8* gene specific forward and reverse primers (**Supplementary Table 1**).

Seeds were surface sterilized and then stratified at 4°C for 4 days in the dark. All seeds were germinated on plates containing half-strength Murashige and Skoog (½ MS) medium supplemented with 2% sucrose and 0.25% Phytigel. Plates were then transferred to a growth chamber at 22 ± 2°C under long day conditions (16-h-light/8-h-dark photoperiod) with 100 E m⁻² s⁻¹ light intensity.

Generation of Transgenic Plants Overexpressing *IAA8*

The *Cauliflower mosaic virus* (*CaMV*) 35S::3xflag-*IAA8* construct in binary vector pCambia 1300 was introduced into *Agrobacterium tumefaciens* strain GV3101 and used for transformation of *iaa8-1* mutant plants by floral dipping. Transformed lines were selected on ½ MS medium containing hygromycin (40 g/mL). Three independent homozygous lines overexpressing *IAA8* were selected from the T₃ generation and used for all experiments.

Seed Germination Assay

Seeds were carefully harvested after siliques were fully mature. The germination assay was performed according to the method of Nguyen et al. (2012). After surface sterilization, seeds of all genotypes were stratified at 4°C for 4 days in the dark and allowed to germinate on ½ MS medium or ½ MS supplemented with 5 μM NAA or 1 μM ABA alone or together at 22 ± 2°C in a growth chamber under a 16-h-light/8-h-dark cycle. Seed germination based on radicle protrusion was quantified from day 0 until day 5. Seeds were considered germinated after radicle protrusion at the indicated time. Statistical analysis was performed, and data are presented as percentage germination rate from three independent experiments with three biological replicates.

Protein Extraction and Immunoblot Analysis

Immunoblot analysis was performed according to the method of Kim et al. (2017). Seedlings were treated with or without MG132, cycloheximide (CHX), or H₂O₂. Tissues were ground in liquid nitrogen to fine powder, and total proteins were extracted using extraction buffer containing 50 mM HEPES, pH 7.5, 5 mM EDTA, 5 mM EGTA, 2 mM DTT, 25 mM NaF, 1 mM Na₃VO₄, 50 mM β-glycerophosphate, 20% glycerol (v/v), 2 mM PMSF, 1% Triton X-100 (v/v), and protease inhibitor cocktail (Roche diagnostics, Germany). Following two rounds of centrifugation at 12,000 × g for 15 min, supernatants were transferred to pre-chilled micro centrifuge tubes. Protein concentration was determined using a protein assay kit (Bio-Rad Laboratories, USA) with bovine serum albumin (BSA) as a standard. For immunoblot analysis, 80 μg of total protein from each sample was separated by 10% Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE) and transferred to polyvinylidene fluoride (PVDF) membranes (Bio-Rad Laboratories, USA). Proteins were probed using mouse anti-flag (1:5000; Sigma, USA) as primary antibody and horseradish peroxidase (HRP) conjugated anti-mouse as secondary antibody (1:5000) and visualized using an ECL kit (Bio-Rad Laboratories, USA).

RNA Extraction and Gene Expression Analysis by Reverse-Transcription Quantitative PCR (RT-qPCR) and Semi-Quantitative RT-PCR

Total RNA was extracted from seeds by the LiCl/phenol method according to the protocol of Nguyen et al. (2012). RNA (2 μg) was reverse transcribed using SuperScript II RNase-Reverse Transcriptase (Invitrogen, USA). RT-qPCR was performed according to the method of Kim et al. (2017) with some modification. RT product (1 μL) was mixed with gene specific primers (10 pmol) in 10 μL reaction volume. SYBR Green PCR Master Mix kit (Bio-Rad SYBR Green Supermix) was added to the mixture and incubated in a CFX384 real-time PCR detection system (Bio-Rad Laboratories, USA). Gene expression was quantified during the logarithmic phase using expression of the housekeeping gene *Tubulin2* as an internal control. Semi-quantitative RT-PCR was carried out as described by Nguyen et al. (2012). Primers used for PCR are listed in **Supplementary Table 2**.

Chromatin Immunoprecipitation Assay

Ten-day-old *iaa8-1* and *iaa8-1/IAA8 OX* seedlings were treated with cold (4°C) and H₂O₂ for 12 h. Chromatin immunoprecipitation (ChIP) was carried out as described (Gendrel et al., 2002) using mouse polyclonal anti-flag antibody (1:3000; Sigma, USA). PCR amplification was performed quantitatively using the CFX384 Real-Time System (Bio-Rad, USA). The immunoprecipitation was replicated three times. The ChIP primers are listed in **Supplementary Table S2**.

RESULTS

IAA8 Is Involved in Seed Germination

IAA8 negatively regulates flower development (Wang et al., 2013). To investigate the biological function of IAA8, we obtained the T-DNA insertion mutants *iaa8-1* (CS25210) and *iaa8-2* (SALK_202296) from SALK (**Supplementary Figure 1A**) and confirmed the T-DNA insertion by diagnostic PCR and semi-quantitative RT-PCR assays (**Supplementary Figure 1B**). The full length transcript of *IAA8* was disrupted in the *iaa8-1* line, whereas the transcript in *iaa8-2* was similar to the Col-0 plants, suggesting that only *iaa8-1* is a loss-of-function mutant (**Supplementary Figure 1C**).

To identify the physiological function of *IAA8*, we intensively screened the *iaa8-1* phenotype. In a time course experiment, seeds of the selected genotypes were germinated on ½ MS medium for 5 days under long day conditions. Germination rate based on radicle protrusion was lower in *iaa8-1* mutant seeds than in Col-0 and *iaa8-2* seeds (**Figure 1A**), indicating that *IAA8* positively regulates seed germination.

To further confirm the *iaa8-1* mutant phenotype, we generated complementation plants expressing *p35S::3xflag-IAA8* in the *iaa8-1* background (*iaa8-1/IAA8 OX*). Three independent homozygous lines were selected from the T₃ generation. To verify ectopic expression, semi-quantitative RT-PCR and immunoblot analyses were performed (**Figures 1B, D**). Aux/IAA repressor proteins are rapidly degraded by the auxin-mediated 26S proteasomal pathway, thereby regulating auxin-dependent transcriptional regulation (Tiwari et al., 2003; Park et al., 2011). We assumed that *IAA8* might also be degraded by the 26S proteasomal pathway, similar to other Aux/IAA proteins (Gilkerson et al., 2015). To test this hypothesis, 10-day-old seedlings of *iaa8-1/IAA8 OX* plants from three independent lines were treated with or without 10 μM MG132 (a proteasome inhibitor) for 3 h. Proteins were extracted and immunoblot analysis was performed. As expected, *IAA8* protein was almost undetectable under mock conditions (control); however, high levels of *IAA8* protein accumulated in the presence of MG132 (**Figure 1D**), indicating that *IAA8* is degraded by the 26S proteasomal pathway.

We next tested seed germination phenotypes using Col-0, *iaa8-1*, and three independent *iaa8-1/IAA8 OX* complemented lines. Seeds of the selected genotypes were allowed to germinate on ½ MS medium under long day conditions. The germination rate of *iaa8-1/IAA8 OX* complemented seeds was almost the same as that of Col-0 (**Figure 1C** and **Supplementary Figure 2**), indicating that the lower germination in the *iaa8-1* mutant line resulted from loss-of-function of *IAA8*. Three independent complemented lines showed similar expression patterns and seed germination phenotype. Therefore, we used the *iaa8-1/IAA8 OX* (#6-3) for further experiments, and hereafter referred to as *iaa8-1/IAA8 OX*.

Previous study suggests that auxin inhibits seed germination through ABA-dependent manner (Liu et al., 2013). Therefore, we tested the effect of auxin on germination by using Col-0, *iaa8-1*, and *iaa8-1/IAA8 OX* seeds in the presence and absence of ABA (**Supplementary Figure 3**). Seeds of the selected genotypes were allowed to germinate on ½ MS or ½ MS supplemented with 5 μM NAA and 1 μM ABA alone or together. In our experimental




FIGURE 1 | IAA8 positively regulates seed germination. **(A)** Seeds of Col-0, *iaa8-1*, and *iaa8-2* mutants were germinated on ½ MS medium after stratification for 4 days in the dark at 4°C in a long-day chamber. Radicle protrusion was quantified at the indicated times (days). Data are presented as mean values of three independent experiments of three biological replicates. Error bars represent SE. Significant difference was determined by Student's t-test (* $P < 0.05$ and ** $P < 0.01$). **(B)** Construction of transgenic lines overexpressing *IAA8* in the *iaa8-1* background. Abundance of *IAA8* transcript in Col-0 and transgenic lines expressing *p35S::3xflag-IAA8* was determined by semi-quantitative RT-PCR. Total RNA was extracted from 10-day-old seedlings, and semi-quantitative RT-PCR analysis was performed using gene-specific primers; *Tubulin2* was used as an internal control. **(C)** *IAA8* rescues the germination phenotype of *iaa8-1*. Seeds of Col-0, *iaa8-1*, and an *iaa8-1/IAA8 OX* (#6-3) complemented line were germinated on ½ MS medium in a long-day chamber. Radicle protrusion was quantified at the indicated times (days). Data are presented as mean values. Error bars represent SE. Significant difference was determined by Student's t-test (* $P < 0.05$ and ** $P < 0.01$). **(D)** Immunoblot analysis of protein extracts prepared from samples used in **(B)** to confirm the expression of flag-tagged *IAA8* protein in the absence (mock) or presence of 10 μ M MG132. Proteins were extracted and probed with anti-flag antibody. Rubisco was stained with Coomassie brilliant blue (CBB) and used as a loading control.

condition, exogenous NAA alone did not inhibit seed germination in Col-0 and *iaa8-1/IAA8 OX* compared to *iaa8-1* mutant (**Supplementary Figures 3A, B**). These results suggest that even upon NAA treatment, small amount of IAA8 protein can be existed in Col-0 and *iaa8-1/IAA8 OX*, which inhibits *ABI3* expression to promote seed germination compared to *iaa8-1* mutant seeds. However, co-treatment of NAA and ABA synergistically inhibited seed germination in Col-0 and *iaa8-1/IAA8 OX* seeds similar to those of *iaa8-1* (**Supplementary Figures 3C, D**), indicating that *ABI3*, *ABI4*, and *ABI5* transcription by ABA may cause inhibitory effect on germination, which cannot be rescued by small amount of IAA8 proteins.

IAA8 Protein Accumulates During Seed Germination

To further investigate the specific role of *IAA8* in seed germination, we searched the public *Arabidopsis* microarray database (<http://bar.utoronto.ca/eplant/>) and found that the *IAA8* transcript is induced by seed imbibition. We therefore proposed that *IAA8* transcription might be induced during seed

germination. To determine the expression pattern of *IAA8* during seed germination, we performed RT-qPCR using freshly harvested Col-0 seeds. The *IAA8* transcript level peaked in germinating seeds at day 2 (**Figure 2A**), indicating that *IAA8* transcription is induced during seed germination.

Because *IAA8* transcription was induced during seed germination, we assumed that the *IAA8* protein might also accumulate following initiation of germination signals. We therefore germinated seeds of *iaa8-1/IAA8 OX* plants and performed immunoblot analysis. As expected, *IAA8* protein accumulated to high levels during seed germination. Interestingly, *IAA8* protein was detectable at day 2 and accumulated rapidly until day 5 (**Figure 2B**). By day 10, however, *IAA8* protein was almost undetectable (**Figure 1D**), suggesting that *IAA8* proteins accumulate only during germination.

IAA8 Protein Stability Is Maintained by Cold and ROS Signals

Cold treatment is considered one of the most efficient stimuli of seed dormancy release (Penfield et al., 2005b). Germinating seeds




FIGURE 2 | IAA8 transcript levels and protein abundance during seed germination. **(A)** IAA8 transcript level in germinating seeds of Col-0 plants. Seeds of Col-0 plants were germinated on $\frac{1}{2}$ MS medium for the indicated time periods (days). Total RNA was extracted, and RT-qPCR analysis was performed using gene-specific primers; *Tubulin2* was used as an internal control. Data are presented as mean values. Error bars represent SE of three biological replicates. Different letters indicate statistically significant differences ($P < 0.05$). **(B)** IAA8 protein accumulation during seed germination. Seeds of *iaa8-1/IAA8 OX* (#6-3) plants were stratified for 4 days in the dark at 4°C and then allowed to germinate on $\frac{1}{2}$ MS medium. Proteins were extracted at the indicated time points (days), and immunoblot analysis was performed using 80 μ g of total protein in each well. IAA8 protein was detected using anti-flag antibody. Membranes were stained using Coomassie brilliant blue (CBB). A representative image of three independent experiments is shown.

produce ROS upon imbibition, which promotes radicle protrusion by rupturing the testa (Leymarie et al., 2012). Cold and imbibition therefore work in parallel during seed germination. As IAA8 protein accumulated to high levels during seed germination after cold imbibition (Figure 2B), we investigated whether cold and ROS affect IAA8 protein level. Seedlings of *iaa8-1/IAA8 OX* plants were exposed to cold (4°C) and 5 mM H₂O₂ alone or together for 12 h. Total proteins were then extracted and immunoblot analysis was performed

(Figure 3). Under mock conditions, IAA8 protein was detected at low levels, indicating that *iaa8-1/IAA8 OX* plants were under mild stress during experimental handling. However, treatment with cold or H₂O₂ induced a rapid accumulation of IAA8 (Figure 3A). Interestingly, more IAA8 protein accumulated when *iaa8-1/IAA8 OX* seedlings were incubated under cold and H₂O₂ together (Figure 3A). These results demonstrate that cold and ROS act together to promote IAA8 protein accumulation during germination.




FIGURE 3 | IAA8 protein is stabilized by cold and H₂O₂. **(A)** Stabilization of IAA8 in the presence of MG132, cold, and H₂O₂. Ten-day-old *iaa8-1/IAA8 OX* (#6-3) seedlings were treated with 10 μ M MG132, cold (4°C), 5 mM H₂O₂, or cold and H₂O₂ together for 12 h. **(B–D)** *De novo* protein synthesis and auxin-mediated turnover of stabilized IAA8 in presence of MG132, cold, or H₂O₂. Ten-day-old *iaa8-1/IAA8 OX* (#6-3) seedlings were treated with 10 μ M MG132. **(B)** cold (4°C). **(C)** or 5 mM H₂O₂ **(D)** for 12 h and then incubated with 500 μ M CHX with or without 20 μ M NAA for the indicated time periods (h). Total proteins were extracted, and immunoblot analysis was performed using anti-flag antibody. Col-0 seedlings were used as a negative control. Rubisco was stained using Coomassie brilliant blue (CBB) and used as a loading control.

We next asked whether accumulation of IAA8 protein under cold and H₂O₂ conditions was caused by *de novo* protein synthesis or increased protein stability. We exposed *iaa8-1/IAA8 OX* seedlings to MG132, cold or H₂O₂ for 12 h, and then incubated them in medium containing 500 μM cycloheximide (CHX; an inhibitor of translation) with or without 20 μM 1-naphthaleneacetic acid (NAA) for 1, 3, and 6 h. As expected, IAA8 protein accumulated rapidly in the presence of MG132; however, preventing *de novo* protein synthesis by treatment with CHX caused significant depletion of IAA8 (Figure 3B). Next, we treated *iaa8-1/IAA8 OX* seedlings with cold or H₂O₂, followed by incubation with CHX. Surprisingly, preventing *de novo* protein synthesis by treatment with CHX did not result in complete loss of IAA8 protein in cold and H₂O₂ treated seedlings (Figures 3C, D), indicating that the increased abundance of IAA8 protein caused by cold and H₂O₂ is independent of *de novo* protein synthesis. Furthermore, treatment with exogenous NAA potentiated the effect of CHX, leading to faster turnover of IAA8 in MG132-treated seedlings (Figure 3B). Surprisingly, however, IAA8 protein was still detectible after 6 h of NAA treatment in both cold- and H₂O₂-treated seedlings (Figures 3C, D). Together, these results suggest that cold and H₂O₂ delay auxin-mediated degradation of IAA8 protein.

IAA8 Down-Regulates *ABI3* Expression During Seed Germination

ABI3 functions in ABA-mediated seed dormancy and inhibition of seed germination. *ABI3* transcript levels are high in dormant seeds and decrease rapidly after germination (Liu et al., 2013). Likewise, *ABI3* protein levels decrease in germinating seeds in a light dependent manner (Lopez-Molina et al., 2002). We therefore proposed that IAA8 accumulation might regulate *ABI3* expression during or after seed germination. To test this hypothesis, we germinated seeds of Col-0, *iaa8-1*, and *iaa8-1/IAA8 OX* under long day conditions and collected samples from day 0 to day 5. Total RNA was then extracted, and RT-qPCR analysis was performed. Interestingly, the transcript levels of *ABI3* were comparatively high in non-germinating Col-0 seeds but declined during germination. The *ABI3* transcript levels were very high in *iaa8-1* seeds compared to those in Col-0 seeds (Figure 4A). Furthermore, the transcript levels of other ABA-responsive genes such as *ABI4*, *ABI5*, *Em1*, *Em6*, and *RAV1* were also analyzed. The results showed that transcript levels of *ABI4*, *ABI5*, *Em1*, and *Em6* were significantly elevated in *iaa8-1* (Figures 4B–E). In contrast, the transcript level of *RAV1* was very low in *iaa8-1* compared to Col-0 plants (Figure 4F). Consistent with the germination results (Figures 1A, C, Supplementary Figure 2), *IAA8* complementation down-regulated *ABI3*, *ABI4*, *ABI5*, *Em1*, and *Em6* expression to levels almost comparable to those in Col-0 (Figure 4), indicating that *IAA8* negatively regulates *ABI3*, *ABI4*, *ABI5*, *Em1*, and *Em6* expression during seed germination.

Aux/IAA proteins are transcriptional repressors that regulate the expression of auxin responsive genes by inactivating ARFs activity (Tiware et al., 2003). Recent studies suggest that *ABI3* functions as a major downstream regulatory component of auxin-mediated seed dormancy (Belin et al., 2009; Liu et al., 2013). We

therefore asked how IAA8 might affect the expression of *ABI3*. For this purpose, chromatin immunoprecipitation (ChIP) assay was performed to confirm the binding ability of IAA8 with *ABI3* promoter using anti-flag antibody. We treated the *iaa8-1* and *iaa8-1/IAA8 OX* plants with mock, cold and H₂O₂ for 12 h, and then ChIP-qPCR analysis was performed. The results showed that IAA8 associates to AuxRE motif on *ABI3* promoter under mock condition in *iaa8-1/IAA8 OX* plants (Figure 5). Interestingly, the *ABI3* promoter fragment were highly enriched in *iaa8-1/IAA8 OX* plants in response to cold and H₂O₂ signals compared to ChIP samples in mock and without anti-flag treated plants (Figure 5B), suggesting that binding of IAA8 to *ABI3* promoter presumably through ARFs could down-regulate *ABI3* transcription during seed germination.

DISCUSSION

Seed germination is mainly controlled by two hormones (Leymarie et al., 2012): ABA inhibits germination, while GA promotes it. Catabolism of ABA and biosynthesis of GA are up-regulated by imbibition to release seeds from dormancy (Liu et al., 2010; Leymarie et al., 2012; Wang et al., 2018). Other hormones, including brassinosteroids, ethylene, and cytokinin, are also known to contribute to seed germination (Steber and McCourt, 2001; Wang et al., 2011; Corbineau et al., 2014). Auxin is generally considered to negatively regulate seed germination in an ABA-dependent manner (Liu et al., 2013). However, the biological role of auxin in seed germination has not been fully elucidated. In this study, we uncovered a novel physiological function of IAA8 in seed germination.

IAA8 Is a Positive Regulator of Seed Germination

The biological functions of most *Aux/IAA* genes have been identified using gain-of-function mutants, with *axr5-1* (*iaa1*), *shy2* (*iaa3*), *iaa6* (*shy1*), *axr2-1* (*iaa7*), *iaa8*, *iaa12* (*bd1*), *iaa14* (*slr*), *iaa16*, *axr3-1* (*iaa17*), *iaa18*, *iaa19* (*msg2*), and *iaa28* reported to display auxin related phenotypes (Timpte et al., 1994; Kim et al., 1996; Leyser et al., 1996; Rouse et al., 1998; Hamann et al., 1999; Nagpal et al., 2000; Rogg et al., 2001; Fukaki et al., 2002; Tian et al., 2003; Tatematsu et al., 2004; Yang et al., 2004; Uehara et al., 2008; Rinaldi et al., 2012; Wang et al., 2013). Among these, the *IAA8* gain-of-function mutant negatively regulates flower development (Wang et al., 2013); however, the *IAA8* loss-of-function mutant shows no developmental phenotype in aerial plants parts (Overvoorde et al., 2005). In this study, we found that the *IAA8* loss-of-function mutant (*iaa8-1*) showed delayed seed germination compared to the wild type (Figures 1A, C). This result correlates well with previous observations. First, external auxin application negatively regulates seed germination (Liu et al., 2013). Second, the gain-of-function mutants *axr2-1* (*IAA7*) and *axr3-1* (*IAA17*) show enhanced seed germination and exhibit stronger ABA insensitivity than the wild type (Belin et al., 2009; Liu et al., 2013). Third, *tir1*, *tir1afb2*, *tir1afb3*, and *tir1afb1afb2afb3*




FIGURE 4 | IAA8 down-regulates ABA responsive genes transcription. Transcript abundance of *ABI3* (A), *ABI4* (B), *ABI5* (C), *Em1* (D), *Em6* (E), and *RAV1* (F) in germinating seeds of Col-0, *iaa8-1* mutant, and *iaa8-1/IAA8 OX* (#6-3) plants. Seeds of the selected genotypes were germinated on 1/2 MS medium. Total RNA was extracted and RT-qPCR analysis was performed using gene-specific primers; *Tubulin2* was used as an internal control. Error bars represent SE in three biological replicates. Significant difference was determined by Student's t-test (*P < 0.05, **P < 0.01 and ***P < 0.001).

mutants with impaired ability to degrade Aux/IAA proteins show enhanced seed germination (Liu et al., 2013). These observations suggest that Aux/IAA proteins act as positive regulators of seed germination. The biological roles of other Aux/IAA proteins during seed germination should be investigated to better understand which and how many Aux/IAA proteins are involved.

IAA8 Accumulation Is Caused by De Novo Synthesis and Increased Stability During Germination

The amount of different proteins in cells is controlled by translation of mRNA, which is dependent on gene expression. In addition, protein accumulation can also be achieved by increasing protein stability in response to external or internal

signals (Finch-Savage et al., 2007; Carrera et al., 2008; Holdsworth et al., 2008). We found that IAA8 protein rapidly accumulated to high levels during seed germination in response to germination signals (Figure 2B). The increase in IAA8 protein was achieved not only by *de novo* protein synthesis but also by protein stabilization (Figure 3). Similarly, the expression of many germination related genes is highly up-regulated in response to germination-inducing signals (Wang et al., 2018). Several Aux/IAA genes, such as *IAA1*, *IAA2*, *IAA3*, *IAA16*, *IAA20*, *IAA26*, *IAA28*, and *IAA29*, are highly expressed during germination (Fujii et al., 2000; Winter et al., 2007; Carranco et al., 2010). Furthermore, transcription of *IAA30*, *IAA11*, and *IAA19* is enhanced under stress conditions in germinating seeds (Park et al., 2011), suggesting that many Aux/IAA genes may play important roles during seed germination in a stress-dependent or -independent manner.





FIGURE 5 | IAA8 associate to *ABI3* gene promoter. **(A)** Structure of the *ABI3* promoter and amplicon regions (P1 and P2). The arrow indicates the transcription start site. The black line and black rectangle indicate 1.5-kb promoter and CDS region, respectively. TGTCTC sequence is auxin responsive element (AuxRE). **(B–C)** Germination signals induce binding of IAA8 to *ABI3* promoter at AuxRE region. Chromatins from *iaa8-1* and *iaa8-1/IAA8 OX* (#6-3) plants were treated with cold (4°C) and H₂O₂ for 12 h. Flag-tagged IAA8 chromatin complex were immunoprecipitated with anti-Flag antibody. A control reaction was processed in parallel without antibody. ChIP-DNA was applied to RT-qPCR using primers specifically targeting to *ABI3* promoter regions, **(B)** (P1) and **(C)** (P2). The ChIP results are presented as fold-enrichment of nontarget DNA. Error bars indicate SE ($n = 3$). Significant difference was determined by Student's t-test (* $P < 0.05$ and ** $P < 0.01$). The experiments were repeated three times with similar results.

This study showed that IAA8 protein is stabilized by cold and ROS signals during seed germination (Figure 3). Similarly, IAA7 and IAA17 are highly stabilized by salt and salicylic acid (Wang et al., 2007; Liu et al., 2015). Thus, we suspect that the stabilization of IAA8 is achieved by unidentified post-translational modification (PTM). Various PTMs, such as phosphorylation, ubiquitination, and sumoylation, can alter protein stability, subcellular localization, and protein interactions in response to external signals (Yang et al., 2003). Interestingly, various Aux/IAA proteins, such as IAA1, IAA5, IAA8, IAA11, IAA13, IAA15, and IAA31, are putative substrates of kinases (Popescu et al., 2009). Therefore, there is a strong possibility that IAA8 is stabilized by cold and ROS through phosphorylation during seed germination.

IAA8 Promotes Seed Germination Through Suppression of *ABI3* Transcription

ABI3 is a well-known key negative regulator in seed germination (Liu et al., 2013). *ABI3* is transcriptionally induced by ABA and involved in ABA-mediated inhibition of seed germination (Bentsink and Koornneef, 2008; Liu et al., 2013). Auxin also negatively regulates seed germination through auxin response factor 10 (ARF10)- and ARF16-mediated increase of *ABI3* transcription (Nemhauser et al., 2006; Santner and Estelle,

2009; Liu et al., 2013). We showed that the *iaa8-1* mutant has delayed seed germination and higher levels of *ABI3*, *ABI4*, *ABI5*, *Em1*, and *Em6* transcript (Figures 4A–E), while the *RAV1* transcript level was markedly repressed in *iaa8-1* mutant compared to wild type (Figure 4F). Similar observations in seeds of various auxin biosynthesis and signaling mutants (Liu et al., 2013) demonstrate that auxin signaling can negatively control seed germination through ARF-mediated regulation of *ABI3* transcription. In addition, exogenous auxin application up-regulates the transcription of *ABI3*, resulting in inhibition of seed germination (Liu et al., 2007; Liu et al., 2013). By contrast, the *arf10arf16* double mutant shows enhanced germination with reduced transcription levels of *ABI3* (Liu et al., 2013), suggesting that ARF10 and ARF16 are required for maintenance of *ABI3* transcription. Moreover, *ABI3* transcription factor positively regulates *ABI4* and *ABI5* expression, which in turn promotes *Em1* and *Em6* transcription, thereby, negatively regulating seed germination (Soderman et al., 2000; Skubacz et al., 2016). In contrast, the higher transcript level of *RAV1* positively regulates seed germination (Feng et al., 2014). Taken together, these results suggest that IAA8 acts as a key upstream regulator during seed germination. Additionally, ChIP assay confirmed that IAA8 associates to *ABI3* promoter, specifically to AuxRE motif via unidentified interacting ARFs (Figure 5). These observations



indicate that *ABI3* transcription can be down-regulated through IAA8-mediated transcriptional inactivation of ARFs proteins. Interestingly, various ARFs, such as ARF4, ARF5, ARF6, ARF7, ARF8, ARF11, ARF14, ARF15, ARF16, ARF19, ARF20, and ARF22 have been previously investigated as IAA8 interacting partners (Vernoux et al., 2011; Arase et al., 2012; Wang et al., 2013; Piya et al., 2014). Therefore, we suspect that stabilized IAA8 down-regulates transcription of *ABI3* by inactivating ARF proteins, which directly bind to the promoter of *ABI3*. A better understanding of auxin signaling in seed germination will be achieved by investigating which specific ARF proteins bind to IAA8 and regulate the transcription of *ABI3*.

We propose a model (Figure 6) explaining how Aux/IAA proteins regulate seed germination. Germination-inducing signals promote seed germination by increasing ROS, which triggers ABA catabolism and GA biosynthesis (Liu et al., 2010).

REFERENCES

- Arase, F., Nishitani, H., Egusa, M., Nishimoto, N., Sakurai, S., Sakamoto, N., et al. (2012). IAA8 involved in lateral root formation interacts with the TIR1 auxin receptor and ARF transcription factors in Arabidopsis. *PLoS One* 7, e43414. doi: 10.1371/journal.pone.0043414
- Arc, E., Chibani, K., Grappin, P., Jullien, M., Godin, B., Cuff, G., et al. (2012). Cold stratification and exogenous nitrates entail similar functional proteome adjustments during Arabidopsis seed dormancy release. *J. Proteome Res.* 11, 5418–5432. doi: 10.1021/pr3006815
- Bassel, G. W., Lan, H., Glaab, E., Gibbs, D. J., Gerjets, T., Krasnogor, N., et al. (2011). Genome-wide network model capturing seed germination reveals

In response to these signals, IAA8 accumulates not only by induction of protein synthesis, but also by protein stabilization. The accumulated IAA8 increases seed germination through transcriptional inhibition of *ABI3* by inactivating ARFs activity.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

AUTHOR CONTRIBUTIONS

SH, SK, and WC conceived the study. XN and WC supervised this study. SH and SB performed the experiments. SH, SK, and AA analyzed the data. SH, AA, DJ-Y, and WC wrote the manuscript with feedback from all authors.

FUNDING

This work was supported by the Next-Generation BioGreen 21 Program (#PJ01325401) funded by RDA and the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIP) (no. 2019R1A2C1009932), partly by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (no. 2018R1A6A3A11042628), and by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 106.02-2017.09. SH and SB were supported by the Brain Korea 21 Plus (BK21+) fellowship program.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2020.00111/full#supplementary-material>

- coordinated regulation of plant cellular phase transitions. *Proc. Natl. Acad. Sci. U. S. A.* 108, 9709–9714. doi: 10.1073/pnas.1100958108
- Belin, C., Megies, C., Hauserova, E., and Lopez-Molina, L. (2009). Abscisic acid represses growth of the Arabidopsis embryonic axis after germination by enhancing auxin signaling. *Plant Cell* 21, 2253–2268. doi: 10.1105/tpc.109.067702
- Bentsink, L., and Koornneef, M. (2008). Seed dormancy and germination. *Arabidopsis Book* 6, e0119. doi: 10.1199/tab.0119
- Bewley, J. D. (1997). Seed germination and dormancy. *Plant Cell* 9, 1055–1066. doi: 10.1105/tpc.9.7.1055
- Carranco, R., Espinosa, J. M., Prieto-Dapena, P., Almoguera, C., and Jordano, J. (2010). Repression by an auxin/indole acetic acid protein connects auxin

- signaling with heat shock factor-mediated seed longevity. *Proc. Natl. Acad. Sci. U. S. A.* 107, 21908–21913. doi: 10.1073/pnas.1014856107
- Carrera, E., Holman, T., Medhurst, A., Dietrich, D., Footitt, S., Theodoulou, F. L., et al. (2008). Seed after-ripening is a discrete developmental pathway associated with specific gene networks in Arabidopsis. *Plant J.* 53, 214–224. doi: 10.1111/j.1365-313X.2007.03331.x
- Chen, H., Zhang, J., Neff, M. M., Hong, S. W., Zhang, H., Deng, X. W., et al. (2008). Integration of light and abscisic acid signaling during seed germination and early seedling development. *Proc. Natl. Acad. Sci. U. S. A.* 105, 4495–4500. doi: 10.1073/pnas.0710778105
- Corbineau, F., Xia, Q., Bailly, C., and El-Maarouf-Bouteau, H. (2014). Ethylene, a key factor in the regulation of seed dormancy. *Front. Plant Sci.* 5, 539. doi: 10.3389/fpls.2014.00539
- Debeaujon, I., Leon-Kloosterziel, K. M., and Koornneef, M. (2000). Influence of the testa on seed dormancy, germination, and longevity in Arabidopsis. *Plant Physiol.* 122, 403–414. doi: 10.1104/pp.122.2.403
- Feng, C. Z., Chen, Y., Wang, C., Kong, Y. H., Wu, W. H., and Chen, Y. F. (2014). Arabidopsis RAV1 transcription factor, phosphorylated by SnRK2 kinases, regulates the expression of ABI3, ABI4, and ABI5 during seed germination and early seedling development. *Plant J.* 80, 654–668. doi: 10.1111/tpj.12670
- Finch-Savage, W. E., Cadman, C. S., Toorop, P. E., Lynn, J. R., and Hilhorst, H. W. (2007). Seed dormancy release in Arabidopsis Cvi by dry after-ripening, low temperature, nitrate and light shows common quantitative patterns of gene expression directed by environmentally specific sensing. *Plant J.* 51, 60–78. doi: 10.1111/j.1365-313X.2007.03118.x
- Fujii, N., Kamada, M., Yamasaki, S., and Takahashi, H. (2000). Differential accumulation of Aux/IAA mRNA during seedling development and gravity response in cucumber (*Cucumis sativus* L.). *Plant Mol. Biol.* 42, 731–740. doi: 10.1023/a:1006379804678
- Fukaki, H., Tameda, S., Masuda, H., and Tasaka, M. (2002). Lateral root formation is blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. *Plant J.* 29, 153–168. doi: 10.1046/j.0960-7412.2001.01201.x
- Gendrel, A. V., Lippman, Z., Yordan, C., Colot, V., and Martienssen, R. A. (2002). Dependence of heterochromatic histone H3 methylation patterns on the Arabidopsis gene DDM1. *Science* 297, 1871–1873. doi: 10.1126/science.1074950
- Gilkerson, J., Kelley, D. R., Tam, R., Estelle, M., and Callis, J. (2015). Lysine residues are not required for proteasome-mediated proteolysis of the Auxin/Indole Acetic Acid Protein IAA1. *Plant Physiol.* 168, 708–720. doi: 10.1104/pp.15.00402
- Giraudat, J., Hauge, B. M., Valon, C., Smalle, J., Parcy, F., and Goodman, H. M. (1992). Isolation of the Arabidopsis ABI3 gene by positional cloning. *Plant Cell* 4, 1251–1261. doi: 10.1105/tpc.4.10.1251
- Gubler, F., Millar, A. A., and Jacobsen, J. V. (2005). Dormancy release, ABA and pre-harvest sprouting. *Curr. Opin. Plant Biol.* 8, 183–187. doi: 10.1016/j.pbi.2005.01.011
- Hamann, T., Mayer, U., and Jurgens, G. (1999). The auxin-insensitive bodenlos mutation affects primary root formation and apical-basal patterning in the Arabidopsis embryo. *Development* 126, 1387–1395.
- Holdsworth, M. J., Finch-Savage, W. E., Grappin, P., and Job, D. (2008). Post-genomics dissection of seed dormancy and germination. *Trends Plant Sci.* 13, 7–13. doi: 10.1016/j.tplants.2007.11.002
- Hubbard, K. E., Nishimura, N., Hitomi, K., Getzoff, E. D., and Schroeder, J. I. (2010). Early abscisic acid signal transduction mechanisms: newly discovered components and newly emerging questions. *Genes Dev.* 24, 1695–1708. doi: 10.1101/gad.1953910
- Kepinski, S., and Leyser, O. (2005). The Arabidopsis F-box protein TIR1 is an auxin receptor. *Nature* 435, 446–451. doi: 10.1038/nature03542
- Kim, B. C., Soh, M. C., Kang, B. J., Furuya, M., and Nam, H. G. (1996). Two dominant photomorphogenic mutations of Arabidopsis thaliana identified as suppressor mutations of hy2. *Plant J.* 9, 441–456. doi: 10.1046/j.1365-313X.1996.09040441.x
- Kim, S. H., Kim, H. S., Bahk, S., An, J., Yoo, Y., Kim, J. Y., et al. (2017). Phosphorylation of the transcriptional repressor MYB15 by mitogen-activated protein kinase 6 is required for freezing tolerance in Arabidopsis. *Nucleic Acids Res.* 45, 6613–6627. doi: 10.1093/nar/gkx417
- Leymarie, J., Vitkauskaitė, G., Hoang, H. H., Gendreau, E., Chazoule, V., Meimoun, P., et al. (2012). Role of reactive oxygen species in the regulation of Arabidopsis seed dormancy. *Plant Cell Physiol.* 53, 96–106. doi: 10.1093/pcp/pcr129
- Leyser, H. M., Pickett, F. B., Dharmasiri, S., and Estelle, M. (1996). Mutations in the AXR3 gene of Arabidopsis result in altered auxin response including ectopic expression from the SAUR-AC1 promoter. *Plant J.* 10, 403–413. doi: 10.1046/j.1365-313x.1996.10030403.x
- Liu, P. P., Montgomery, T. A., Fahlgren, N., Kasschau, K. D., Nonogaki, H., and Carrington, J. C. (2007). Repression of AUXIN RESPONSE FACTOR10 by microRNA160 is critical for seed germination and post-germination stages. *Plant J.* 52, 133–146. doi: 10.1111/j.1365-313X.2007.03218.x
- Liu, Y., Ye, N., Liu, R., Chen, M., and Zhang, J. (2010). H₂O₂ mediates the regulation of ABA catabolism and GA biosynthesis in Arabidopsis seed dormancy and germination. *J. Exp. Bot.* 61, 2979–2990. doi: 10.1093/jxb/erq125
- Liu, X., Zhang, H., Zhao, Y., Feng, Z., Li, Q., Yang, H. Q., et al. (2013). Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* 110, 15485–15490. doi: 10.1073/pnas.1304651110
- Liu, W., Li, R. J., Han, T. T., Cai, W., Fu, Z. W., and Lu, Y. T. (2015). Salt stress reduces root meristem size by nitric oxide-mediated modulation of auxin accumulation and signaling in Arabidopsis. *Plant Physiol.* 168, 343–356. doi: 10.1104/pp.15.00030
- Lopez-Molina, L., Mongrand, S., McLachlin, D. T., Chait, B. T., and Chua, N. H. (2002). ABI5 acts downstream of ABI3 to execute an ABA-dependent growth arrest during germination. *Plant J.* 32, 317–328. doi: 10.1046/j.1365-313X.2002.01430.x
- Nagpal, P., Walker, L. M., Young, J. C., Sonawala, A., Timpte, C., Estelle, M., et al. (2000). AXR2 encodes a member of the Aux/IAA protein family. *Plant Physiol.* 123, 563–574. doi: 10.1104/pp.123.2.563
- Nee, G., Xiang, Y., and Soppe, W. J. (2017). The release of dormancy, a wake-up call for seeds to germinate. *Curr. Opin. Plant Biol.* 35, 8–14. doi: 10.1016/j.pbi.2016.09.002
- Nemhauser, J. L., Hong, F., and Chory, J. (2006). Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* 126, 467–475. doi: 10.1016/j.cell.2006.05.050
- Nguyen, X. C., Hoang, M. H., Kim, H. S., Lee, K., Liu, X. M., Kim, S. H., et al. (2012). Phosphorylation of the transcriptional regulator MYB44 by mitogen-activated protein kinase regulates Arabidopsis seed germination. *Biochem. Biophys. Res. Commun.* 423, 703–708. doi: 10.1016/j.bbrc.2012.06.019
- Overvoorde, P. J., Okushima, Y., Alonso, J. M., Chan, A., Chang, C., Ecker, J. R., et al. (2005). Functional genomic analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in Arabidopsis thaliana. *Plant Cell* 17, 3282–3300. doi: 10.1105/tpc.105.036723
- Park, J., Kim, Y. S., Kim, S. G., Jung, J. H., Woo, J. C., and Park, C. M. (2011). Integration of auxin and salt signals by the NAC transcription factor NTM2 during seed germination in Arabidopsis. *Plant Physiol.* 156, 537–549. doi: 10.1104/pp.111.177071
- Penfield, S., Graham, S., and Graham, I. A. (2005a). Storage reserve mobilization in germinating oilseeds: Arabidopsis as a model system. *Biochem. Soc Trans.* 33, 380–383. doi: 10.1042/BST0330380
- Penfield, S., Josse, E.-M., Kannangara, R., Gilday, A. D., Halliday, K. J., and Graham, I. A. (2005b). Cold and light control seed germination through the bHLH transcription factor SPATULA. *Curr. Biol.* 15, 1998–2006. doi: 10.1016/j.cub.2005.11.010
- Piskurewicz, U., Jikumaru, Y., Kinoshita, N., Nambara, E., Kamiya, Y., and Lopez-Molina, L. (2008). The gibberellic acid signaling repressor RGL2 inhibits Arabidopsis seed germination by stimulating abscisic acid synthesis and ABI5 activity. *Plant Cell* 20, 2729–2745. doi: 10.1105/tpc.108.061515
- Piya, S., Shrestha, S. K., Binder, B., Stewart, C. N., and Hewezi, T. (2014). Protein-protein interaction and gene co-expression maps of ARFs and Aux/IAAs in Arabidopsis. *Front. Plant Sci.* 5, 744. doi: 10.3389/fpls.2014.00744
- Popescu, S. C., Popescu, G. V., Bachan, S., Zhang, Z., Gerstein, M., Snyder, M., et al. (2009). MAPK target networks in Arabidopsis thaliana revealed using functional protein microarrays. *Genes Dev.* 23, 80–92. doi: 10.1101/gad.1740009

- Rajjou, L., and Debeaujon, I. (2008). Seed longevity: survival and maintenance of high germination ability of dry seeds. *C. R. Biol.* 331, 796–805. doi: 10.1016/j.crvi.2008.07.021
- Rinaldi, M. A., Liu, J., Enders, T. A., Bartel, B., and Strader, L. C. (2012). A gain-of-function mutation in IAA16 confers reduced responses to auxin and abscisic acid and impedes plant growth and fertility. *Plant Mol. Biol.* 79, 359–373. doi: 10.1007/s11103-012-9917-y
- Rogg, L. E., Lasswell, J., and Bartel, B. (2001). A gain-of-function mutation in IAA28 suppresses lateral root development. *Plant Cell* 13, 465–480. doi: 10.1105/tpc.13.3.465
- Rouse, D., Mackay, P., Stirnberg, P., Estelle, M., and Leyser, O. (1998). Changes in auxin response from mutations in an AUX/IAA gene. *Science* 279, 1371–1373. doi: 10.1126/science.279.5355.1371
- Santner, A., and Estelle, M. (2009). Recent advances and emerging trends in plant hormone signalling. *Nature* 459, 1071. doi: 10.1038/nature08122
- Shu, K., Liu, X. D., Xie, Q., and He, Z. H. (2016). Two faces of one seed: hormonal regulation of dormancy and germination. *Mol. Plant* 9, 34–45. doi: 10.1016/j.molp.2015.08.010
- Skubacz, A., Daszkowska-Golec, A., and Szarejko, I. (2016). The role and regulation of ABI5 (ABA-Insensitive 5) in plant development, abiotic stress responses and phytohormone crosstalk. *Front. In Plant Sci.* 7, 1884. doi: 10.3389/fpls.2016.01884
- Soderman, E. M., Brocard, I. M., Lynch, T. J., and Finkelstein, R. R. (2000). Regulation and function of the Arabidopsis ABA-insensitive 4 gene in seed and abscisic acid response signaling networks. *Plant Physiol.* 124, 1752–1765. doi: 10.1104/pp.124.4.1752
- Steber, C. M., and McCourt, P. (2001). A role for brassinosteroids in germination in Arabidopsis. *Plant Physiol.* 125, 763–769. doi: 10.1104/pp.125.2.763
- Tatematsu, K., Kumagai, S., Muto, H., Sato, A., Watahiki, M. K., Harper, R. M., et al. (2004). MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. *Plant Cell* 16, 379–393. doi: 10.1105/tpc.018630
- Tian, Q., Nagpal, P., and Reed, J. W. (2003). Regulation of Arabidopsis SHY2/IAA3 protein turnover. *Plant J.* 36, 643–651. doi: 10.1046/j.1365-313X.2003.01909.x
- Timppte, C., Wilson, A. K., and Estelle, M. (1994). The axr2-1 mutation of Arabidopsis thaliana is a gain-of-function mutation that disrupts an early step in auxin response. *Genetics* 138, 1239–1249.
- Tiwari, S. B., Hagen, G., and Guilfoyle, T. (2003). The roles of auxin response factor domains in auxin-responsive transcription. *Plant Cell* 15, 533–543. doi: 10.1105/tpc.008417
- Uehara, T., Okushima, Y., Mimura, T., Tasaka, M., and Fukaki, H. (2008). Domain II mutations in CRANE/IAA18 suppress lateral root formation and affect shoot development in Arabidopsis thaliana. *Plant Cell Physiol.* 49, 1025–1038. doi: 10.1093/pcp/pcn079
- Vernoux, T., Brunoud, G., Farcot, E., Morin, V., Van den Daele, H., Legrand, J., et al. (2011). The auxin signalling network translates dynamic input into robust patterning at the shoot apex. *Mol. Syst. Biol.* 7, 508. doi: 10.1038/msb.2011.39
- Wang, D., Pajerowska-Mukhtar, K., Culler, A. H., and Dong, X. (2007). Salicylic acid inhibits pathogen growth in plants through repression of the auxin signaling pathway. *Curr. Biol.* 17, 1784–1790. doi: 10.1016/j.cub.2007.09.025
- Wang, Y., Li, L., Ye, T., Zhao, S., Liu, Z., Feng, Y. Q., et al. (2011). Cytokinin antagonizes ABA suppression to seed germination of Arabidopsis by downregulating ABI5 expression. *Plant J.* 68, 249–261. doi: 10.1111/j.1365-313X.2011.04683.x
- Wang, J., Yan, D. W., Yuan, T. T., Gao, X., and Lu, Y. T. (2013). A gain-of-function mutation in IAA8 alters Arabidopsis floral organ development by change of jasmonic acid level. *Plant Mol. Biol.* 82, 71–83. doi: 10.1007/s11103-013-0039-y
- Wang, X., Yesbergenova-Cuny, Z., Biniek, C., Bailly, C., El-Maarouf-Bouteau, H., and Corbineau, F. (2018). Revisiting the role of ethylene and N-end rule pathway on chilling-induced dormancy release in Arabidopsis seeds. *Int. J. Mol. Sci.* 19, 3577. doi: 10.3390/ijms19113577
- Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G. V., and Provart, N. J. (2007). An “Electronic Fluorescent Pictograph” browser for exploring and analyzing large-scale biological data sets. *PLoS One* 2, e718. doi: 10.1371/journal.pone.0000718
- Yang, S. H., Sharrocks, A. D., and Whitmarsh, A. J. (2003). Transcriptional regulation by the MAP kinase signaling cascades. *Gene* 320, 3–21. doi: 10.1016/s0378-1119(03)00816-3
- Yang, X., Lee, S., So, J. H., Dharmasiri, S., Dharmasiri, N., Ge, L., et al. (2004). The IAA1 protein is encoded by AXR5 and is a substrate of SCF(TIR1). *Plant J.* 40, 772–782. doi: 10.1111/j.1365-313X.2004.02254.x
- Zhao, Y. (2010). Auxin biosynthesis and its role in plant development. *Annu. Rev. Plant Biol.* 61, 49–64. doi: 10.1146/annurev-arplant-042809-112308

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Hussain, Kim, Bahk, Ali, Nguyen, Yun and Chung. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.