

T. Kimura
Nagoya Math. J.
Vol. 85 (1982), 1-80

THE b -FUNCTIONS AND HOLONOMY DIAGRAMS OF IRREDUCIBLE REGULAR PREHOMOGENEOUS VECTOR SPACES

TATSUO KIMURA

Introduction

The purpose of this paper is to investigate the micro-local structure and to calculate, by constructing the holonomy diagrams, the b -functions (See [2]) of irreducible regular prehomogeneous vector spaces (See [1]).

Since we know the relation of b -functions with respect to casting transformations (See § 12), it is enough to calculate them only when they are reduced. In this paper, we shall deal with twenty of all twenty nine reduced regular P.V.'s in the Table in [1]. Together with other articles, this completes the list of b -functions of irreducible reduced regular prehomogeneous vector spaces (See § 12) except $(SL(5) \times GL(4), A_2 \otimes A_1, V(10) \otimes V(4))$ which is the most complicated case (See I. Ozeki [11]). This paper consists of the following twelve sections and one Appendix with I. Ozeki.

- § 1. Preliminaries
- § 2. Regular P.V.'s related with $GL(n)$
- § 3. $(Sp(n) \times GL(2m), A_1 \otimes A_1, V(2n) \otimes V(2m))$
- § 4. $(Spin(10) \times GL(2), \text{half-spin rep.} \otimes A_1, V(16) \otimes V(2))$
- § 5. $(GL(1) \times Spin(12), \square \otimes \text{half-spin rep.}, V(1) \otimes V(32))$
- § 6. $(GL(1) \times E_6, \square \otimes A_1, V(1) \otimes V(27))$
- § 7. $(GL(1) \times E_7, \square \otimes A_6, V(1) \otimes V(56))$
- § 8. $(GL(6), A_3, V(20))$
- § 9. $(GL(1) \times Sp(3), \square \otimes A_1, V(1) \otimes V(14))$
- § 10. $(GL(7), A_3, V(35))$
- § 11. $(SL(5) \times GL(3), A_2 \otimes A_1, V(10) \otimes V(3))$
- § 12. Table of the b -functions of irreducible reduced regular P.V.'s

Received January 11, 1979
Revised September 7, 1979

Appendix with I. Ozeki. ($GL(1) \times Spin(14)$, $\square \otimes$ half-spin rep., $V(1) \otimes V(64)$)

In § 1, we shall review the main results of [2] which will be used later. From § 2 to § 11, we do the classification of the orbits, construction of the holonomy diagrams and calculation of the b -functions. In § 12, we shall give the list of b -functions for irreducible reduced regular P.V.'s. Some of them have been already calculated by M. Sato and the author using the different method (See [7]). The holonomy diagrams in § 2, § 8 and § 10 are first obtained by M. Sato. The author would like to express his hearty thanks to Professors Mikio Sato and Masaki Kashiwara for their invaluable advice and encouragement.

§ 1. Preliminaries

Let (G, ρ, V) be an irreducible regular prehomogeneous vector space (abbrev. P.V.) with the singular set S . Then S is the zeros of the relative invariant $f(x) : S = \{x \in V; f(x) = 0\}$, $f(\rho(g)x) = \chi(g)f(x)$ for all $g \in G$ and $x \in V$. We shall consider the micro-differential equations $\mathfrak{M} = \mathcal{E}f^s$ where \mathcal{E} is the sheaf of micro-differential operators of finite order on the cotangent bundle $T^*V = V \times V^*$ (See [2]). Note that the group G acts on T^*V by $(x, y) \mapsto (\rho(g)x, \rho^*(g)y)$ for $x \in V$, $y \in V^*$ and $g \in G$ where ρ^* denotes the contragredient representation of ρ . Let A be the Zariski-closure of a conormal bundle of some G -orbit $\rho(G) \cdot x_0 (x_0 \in V)$. Since we consider only the Zariski-closure of a conormal bundle, we shall omit the word "the Zariski-closure" for simplicity. Assume that A is G -prehomogeneous and is contained in $W = \overline{\{(x, \text{grad } \log f(x)^s); x \in V - S, s \in \mathbb{C}\}}$. In this case, A is called a good holonomic variety. It is an irreducible component of the characteristic variety of \mathfrak{M} . We can show that there exists a local b -function $b_A(s)$ which is unique up to a constant multiple (See [2]). We have $b_{V \times \{0\}}(s) = 1$ and $b_{\{0\} \times V^*}(s) = b(s)$ where $b(s)$ denotes the b -function of this P.V. When two good holonomic varieties A_0 and A_1 intersect with codimension one, we have the relation between $b_{A_0}(s)$ and $b_{A_1}(s)$ as follows (See [2]).

THEOREM 1-1 ([2] Theorem 7-5). *Let A_0 and A_1 be good holonomic varieties whose intersection is of codimension one with the intersection exponent $(\mu : \nu)$. Assume that $\mathfrak{M} = \mathcal{E}f^s$ is a simple holonomic system with support $A_0 \cup A_1$ and $A_0 \cap A_1 \not\subset \overline{\text{supp } \mathfrak{M}} - (A_0 \cup A_1)$. Assume that $m_0 > m_1$, where*

$\text{ord}_{A_i} f^s = -m_i s - \mu_i/2$ ($i = 0, 1$). Then we have, up to a constant multiple,

$$(1.1) \quad b_{A_0}(s)/b_{A_1}(s) = \prod_{k=0}^{\nu} \left[\frac{1}{\nu+1} (\text{ord}_{A_1} f^s - \text{ord}_{A_0} f^s) + \frac{\mu+2k}{2(\nu+\mu)} \right]^{(m_0-m_1)/(\nu+1)}$$

where $[\alpha]^k = \alpha(\alpha+1)\cdots(\alpha+k-1)$.

Here we denote by $\text{ord}_A f^s$ the order of f^s at A (See [2]). Note that m_0 and m_1 are non-negative integers, and $(\mu:\nu) = (1:0)$ or $(\mu:\nu)$ is a pair of positive integers satisfying $\mu \geq 2$, $\nu \geq 1$, and $(m_0 - m_1)$ is a multiple of $(\nu + 1)$.

COROLLARY 1-2 ([2] Corollary 7-6). *If A_0 and A_1 intersect regularly, i.e., $\mu = 1$ and $\nu = 0$, we have*

$$(1.2) \quad b_{A_0}(s)/b_{A_1}(s) = \prod_{k=1}^{m_0-m_1} \left((m_0 - m_1)s + \frac{\mu_0 - \mu_1 - 1}{2} + k \right)$$

where $\text{ord}_{A_i} f^s = -m_i s - \frac{\mu_i}{2}$ ($i = 0, 1$).

Let A be a good holonomic variety. Then $A = \overline{G(x_0, y_0)}$ for some $x_0 \in V$, $y_0 \in V^*$ where $G(x_0, y_0) = \{(\rho(g)x_0, \rho^*(g)y_0); g \in G\}$. In this case, we can calculate the order $\text{ord}_A f^s$ by the following proposition.

PROPOSITION 1-3 ([2] Proposition 4-14). *Let A_0 be an element of the Lie algebra \mathfrak{g} of G satisfying $d\rho(A_0)x_0 = 0$ and $d\rho^*(A_0)y_0 = y_0$. Then we have*

$$(1.3) \quad \text{ord}_A f^s = s\delta\chi(A_0) - \text{tr}_{V_{x_0}^*} d\rho_{x_0}(A_0) + \frac{1}{2} \dim V_{x_0}^*$$

where $V_{x_0}^*$ denotes the conormal vector space $(d\rho(\mathfrak{g}) \cdot x_0)^\perp$, and $d\rho_{x_0}$ denotes the representation of $\mathfrak{g}_{x_0} = \{A \in \mathfrak{g}; d\rho(A)x_0 = 0\}$ induced by $d\rho^*$.

Now let $A_0 = \overline{G(x_0, y_0)}$ and $A_1 = \overline{G(x_1, y_1)}$ be good holonomic varieties such that $(x_0, y_1) \in A_0 \cap A_1$ and $\dim G(x_0, y_1) = \dim V - 1$. In this case, the intersection exponent $(\mu:\nu)$ is given by the following proposition.

PROPOSITION 1-4 ([2] Proposition 6-5). *Let A_1 be an element of \mathfrak{g} satisfying $d\rho(A_1)x_0 = 0$ and $d\rho^*(A_1)y_1 = y_1$. Then A_1 acts on the one-dimensional vector space $\tilde{V} = V_{x_0}^*$ modulo $d\rho_{x_0}(\mathfrak{g}_{x_0})y_1$. Let β be its eigenvalue, i.e., $\beta = \text{tr}_V A_1$. Then μ and ν are given by $\beta = \mu/(\mu + \nu)$, $(\mu, \nu) = 1$. If β is not determined uniquely, i.e., β depends on A_1 , then we have $\mu = 1$, $\nu = 0$, and A_0, A_1 intersect regularly.*

Let $A = \overline{T(\rho(G)x_0)}^\perp$ be a conormal bundle of a G -orbit $\rho(G)x_0$. Then G acts on A prehomogeneously if and only if the colocalization $(G_{x_0}, \rho_{x_0}, V_{x_0}^*)$ at x_0 is a P.V. We shall consider some sufficient conditions that $A \subset W$, i.e., A is a good holonomic variety.

PROPOSITION 1-5 ([2] Proposition 6-6). *Let A_0 and A_1 be two conormal bundles of some G -orbits. Assume that $\dim \mathfrak{g}_0 \cdot p = \dim V - 1$ for some $p \in A_0 \cap A_1$ where $\mathfrak{g}_0 = \{A \in \mathfrak{g}; \delta\chi(A) = 0\}$. Assume that A_0 (or A_1) $\subset W$. Then we have $A_0 \cup A_1 \subset W$. Moreover W is non-singular and $W = \{(x, y) \in V \times V^*; \langle d\rho(A)x, y \rangle = 0 \text{ for all } A \in \mathfrak{g}_0\}$ near p .*

Let $V_{x_0} = V \bmod d\rho(\mathfrak{g})x_0$ be the normal vector space. Then the isotropy subgroup G_{x_0} acts on V_{x_0} . We denote this action by $\tilde{\rho}_{x_0}$. Let f_{x_0} be the localization of $f(x)$ at x_0 (See [2]). This is a relative invariant of $(G_{x_0}, \tilde{\rho}_{x_0}, V_{x_0})$ corresponding to $\chi|_{G_{x_0}}$. Let S_{x_0} be the singular set of $(G_{x_0}, \tilde{\rho}_{x_0}, V_{x_0})$.

PROPOSITION 1-6 ([2] Proposition 6-9). *If $\text{grad log } f_{x_0}: V_{x_0} - S_{x_0} \rightarrow V_{x_0}^*$ is generically surjective, then $A_0 = \overline{T(\rho(G)x_0)}^\perp \subset W$, i.e., A_0 is a good holonomic variety.*

COROLLARY 1-7 ([2] Corollary 6-10). *Assume that the colocalization $(G_{x_0}, \rho_{x_0}, V_{x_0}^*)$ of (G, ρ, V) at x_0 ($\in V$) is a regular P.V. If $\delta\chi|_{\mathfrak{g}_{x_0}}$ is a non-degenerate element, then the conormal bundle $A_0 = \overline{T(\rho(G)x_0)}^\perp$ of the G -orbit $\rho(G)x_0$ is a good holonomic variety.*

COROLLARY 1-8 ([2] Corollary 6-11). *Assume that the colocalization $(G_{x_0}, \rho_{x_0}, V_{x_0}^*)$ of (G, ρ, V) at x_0 ($\in V$) is an irreducible regular P.V. Then the conormal bundle $A_0 = \overline{T(\rho(G)x_0)}^\perp$ of the orbit $\rho(G)x_0$ is a good holonomic variety.*

PROPOSITION 1-9 ([1] Proposition 14 in § 4).

(1) *For $d = \deg f$ and $n = \dim V$, we have $d|2n$ and $\chi(g)^{2n/d} = \det_V \rho(g)^2$ for $g \in G$.*

(2) *$\delta\chi(A) = (d/n) \text{tr } d\rho(A)$ for $A \in \mathfrak{g}$.*

Remark 1-10. Let (G, ρ, V) be an irreducible regular P.V. with finitely many orbits. Let $\mathcal{L} = \{A, A', \dots, A''\}$ be the set of all conormal bundles in W , of some G -orbits in V . The holonomy diagram is, by definition, given as follows.

If $\dim A \cap A' = \dim V - 1$, and $A \cap A' \not\subset A''$ for any other A'' in \mathcal{L} , then we write the diagram as in Figure 1-1. Moreover, if A and A' are good



Figure 1-1.




Figure 1-2.




Figure 1-3. ($m_{A'} > m_A$)

holonomic varieties, we write the orders $\text{ord}_A f^s = -m_A s - \mu_A/2$ for A and A' , and the ratio of the b -functions as in Figure 1-3. If $\dim A \cap A' = \dim V - 1$ and $A \cap A' \subset A''$ for some A'' , then we write the diagram as in Figure 1-2 (e.g. Figure 11-1). Although some general theory for such cases has been established, it is not published yet and hence in this paper we avoid to argue this case. Actually, only in § 11, such case will appear and to calculate the b -function in § 11, we can use another part of the holonomy diagram. Although usually we do not write the conormal bundles outside W (e.g. Figure 3-2), sometimes we write them (e.g. Figure 4-1, Figure 11-1). Since G is reductive, we have $(G, \rho, V) \cong (G, \rho^*, V^*)$ and we identify them.

We sometimes write as $\textcircled{A} \text{---} \textcircled{A'}$ when T and T' are the dual orbits of each other (See § 11) where A and A' are the conormal bundles of T and T' respectively. If $T = T'$, we write as $\textcircled{A} \text{---}$ (e.g. Figure 4-1 and Figure 11-1).

§ 2. Regular P.V.'s related with $GL(n)$

We shall use the same notations as in [1].

2-1. $(\tilde{G} \times GL(m), \tilde{\rho} \otimes A_1, V(m) \otimes V(m))$ where $\tilde{\rho}: \tilde{G} \rightarrow GL(V(m))$ is an m -dimensional irreducible representation of a connected semi-simple algebraic group \tilde{G} (or $\tilde{G} = \{1\}$ and $m = 1$)

The representation space $V = V(m) \otimes V(m)$ can be identified with the totality of $m \times m$ matrices $M(m, C)$. Then the action $\rho = \tilde{\rho} \otimes A_1$ is given by $\rho(g)X = \tilde{\rho}(g_1)X^t g_2$ for $g = (g_1, g_2) \in G = \tilde{G} \times GL(m)$, $X \in M(m, C)$. The relative invariant $f(X)$ is given by the determinant: $f(X) = \det X$. Since we are concerned with relative invariants, we may assume that $\tilde{G} = SL(m)$ and $\tilde{\rho} = A_1$. It is well-known that there exist $(m + 1)$ -orbits

$$\rho(G)X_\mu = \{X \in M(m, \mathbf{C}); \text{rank } X = \mu\}$$

$$\text{where } X_\mu = \left(\begin{array}{ccc|ccc} 1 & & & & & \\ & \ddots & & & & \\ & & \ddots & & & \\ & & & 1 & & \\ & & & & 0 & \dots \\ & & & & & \ddots \\ 0 & & & & & & 0 \end{array} \right) \text{ for } \mu = 0, 1, \dots, m.$$

We identify the dual V^* of V with $V = M(m, \mathbf{C})$ by $\langle X, Y \rangle = \text{tr } {}^tXY$ for $X, Y \in M(m, \mathbf{C})$. Then the dual ρ^* of ρ is given by $\rho^*(g)Y = {}^t g_1^{-1} Y g_2^{-1}$ for $g = (g_1, g_2) \in G = SL(m) \times GL(m)$, $Y \in M(m, \mathbf{C})$.

Since $d\rho(\tilde{A})X_\mu = AX_\mu + X_\mu {}^t B = \left(\begin{array}{ccc|ccc} A_1 + {}^t B_1 & {}^t B_3 & & & & \\ & A_2 & & & & \\ & & A_3 & & & \\ \hline & & & 0 & & \\ & & & & 0 & \\ & & & & & 0 \end{array} \right)$ for $\tilde{A} = (A, B) \in \mathfrak{g}$ with $A = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix}$ and $B = \begin{pmatrix} B_1 & B_2 \\ B_3 & B_4 \end{pmatrix}$, the conormal vector space $V_{X_\mu}^* = (d\rho(\mathfrak{g})X_\mu)^\perp$ is given by $V_{X_\mu}^* = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & Y_\mu \end{pmatrix}; Y_\mu \in M(m - \mu, \mathbf{C}) \right\}$. The isotropy subalgebra $\mathfrak{g}_{X_\mu} = \{\tilde{A} \in \mathfrak{g}; d\rho(\tilde{A})X_\mu = 0\}$ is given by

$$(2.1) \quad \mathfrak{g}_{X_\mu} = \left\{ \left(\begin{pmatrix} A_1 & A_2 \\ 0 & A_4 \end{pmatrix}, \begin{pmatrix} -{}^t A_1 & B_2 \\ 0 & B_4 \end{pmatrix} \right) \in \mathfrak{g}; A_1 \in M(\mu, \mathbf{C}), A_2, B_2 \in M(\mu, m - \mu, \mathbf{C}), A_4, B_4 \in M(m - \mu, \mathbf{C}) \right\}.$$

This \mathfrak{g}_{X_μ} acts on $V_{X_\mu}^*$ as $d\rho_{X_\mu}(\tilde{A})Y_\mu = -{}^t A_1 Y_\mu - Y_\mu B_4$ for $\tilde{A} \in \mathfrak{g}_{X_\mu}$. Therefore we have $(G_{X_\mu}, \rho_{X_\mu}, V_{X_\mu}^*) \cong (SL(m - \mu) \times GL(m - \mu), A_1 \otimes A_4, V(m - \mu) \otimes V(m - \mu))$. Put $Y_\mu = \begin{pmatrix} 0 & 0 \\ 0 & I_{m - \mu} \end{pmatrix}$ where $I_{m - \mu}$ denotes the unit matrix of size $m - \mu$ ($\mu = 0, 1, \dots, m$). Then Y_μ is a generic point of the colocalization $(G_{X_\mu}, \rho_{X_\mu}, V_{X_\mu}^*)$, and $Y_{\mu+1}$ is a point of the one-codimensional orbit ($\mu \leq m - 1$). We denote by A_μ the conormal bundle of $\rho(G)X_\mu$ ($0 \leq \mu \leq m$). Then, we have $\dim A_\mu \cap A_{\mu+1} = \dim V - 1$. Note that the colocalization $(G_{X_\mu}, \rho_{X_\mu}, V_{X_\mu}^*)$ ($\mu = 0, 1, \dots, m$) has finitely many orbits with the unique one-codimensional orbit, it is clear that we have obtained all one-codimensional intersections among A_μ ($\mu = 0, 1, \dots, m$). Since $\mathfrak{g}_0 = \mathfrak{sl}(m) \oplus \mathfrak{sl}(m)$, we have $\dim \mathfrak{g}_0(X_\mu, Y_{\mu+1}) = m^2 - 1$ for $\mu = 0, 1, \dots, m - 1$, and hence by Proposition 1-5, we have $A_\mu \subset W$, i.e., A_μ is a good holonomic variety ($0 \leq \mu \leq m$). Note that $A_m = V \times \{0\}$ is always a good holonomic variety. We shall calculate the intersection exponent $(\tilde{\nu} : \tilde{\nu})$ of A_μ and $A_{\mu+1}$ by using Proposition 1-4. For any $\beta \in \mathbf{C}$, put $A_\mu^\beta = \left((0), \begin{pmatrix} 0 & 0 \\ 0 & -\beta \\ & & -I_{m - \mu - 1} \end{pmatrix} \right) \in \mathfrak{g}$. Then we have $d\rho(A_\mu^\beta)X_\mu = 0$, $d\rho^*(A_\mu^\beta)Y_{\mu+1} = Y_{\mu+1}$, and $\beta = \text{tr } A_\mu^\beta$ where tr denotes

the trace of A_μ^s on $V_{X_\mu}^*$ modulo $d\rho_{X_\mu}(\mathfrak{g}_{X_\mu})Y_{\mu+1}$ since $V_{X_\mu}^*$ modulo $d\rho_{X_\mu}(\mathfrak{g}_{X_\mu})Y_{\mu+1} \cong \{yE_{\mu+1, \mu+1} \in M(m, \mathbb{C}); y \in \mathbb{C}\}$ where E_{ij} denotes the matrix unit. Therefore we have $\tilde{\nu} = 1$ and $\tilde{\nu} = 0$, i.e., they intersect regularly. Now by Proposition 1-3, we shall calculate the order $\text{ord}_{A_\mu} f^s$ of $\mathfrak{m} = \mathcal{E}f^s$ at A_μ where $f(X) = \det X$.

Put $A_\mu = \left((0, \begin{pmatrix} 0 & 0 \\ 0 & -I_{m-\mu} \end{pmatrix}) \right) \in \mathfrak{g}$. Then $d\rho(A_\mu)X_\mu = 0$ and $d\rho^*(A_\mu)Y_\mu = Y_\mu$ ($0 \leq \mu \leq m$). The character $\delta\chi$ corresponding to $f(X) = \det X$ is given by $\delta\chi(\tilde{A}) = \text{tr } B$ for $\tilde{A} = (A, B) \in \mathfrak{g} = \mathfrak{sl}(m) \oplus \mathfrak{gl}(m)$. Since $\dim V_{X_\mu}^* = (m - \mu)^2$ and $\text{tr}_{V_{X_\mu}^*} d\rho_{X_\mu}(A_\mu) = (m - \mu)^2$, we have $\text{ord}_{A_\mu} f^s = s\delta\chi(A_\mu) - \text{tr}_{V_{X_\mu}^*} d\rho_{X_\mu}(A_\mu) + (1/2)\dim V_{X_\mu}^* = -(m - \mu)s - ((m - \mu)^2/2)$. Thus we obtain the holonomy diagram (Figure 2-1).

By Corollary 1-2, we have $b_{A_\mu}(s)/b_{A_{\mu+1}}(s) = s + (m - \mu)$ ($0 \leq \mu \leq m - 1$). Hence

$$\begin{aligned} b(s) &= b_{A_0}(s) = b_{A_m}(s) \cdot \prod_{\mu=0}^{m-1} b_{A_\mu}(s)/b_{A_{\mu+1}}(s) \\ &= \prod_{\mu=0}^{m-1} (s + m - \mu) = (s + 1)(s + 2) \cdots (s + m). \end{aligned}$$

Note that $b_{A_m}(s) = 1$.




Figure 2-1. Holonomy diagram of $(SL(m) \times GL(m), A_1 \otimes A_1, V(m) \otimes V(m))$

Remark 2-1. The b -function of 2-1 is classically known by using Capelli's identity (See H. Weyl: Classical Groups).

2-2. $(GL(n), 2A_1, V(\frac{1}{2}n(n+1)))$ ($n \geq 2$)

The representation space can be identified with the totality of $n \times n$ symmetric matrices $V = \{X \in M(n, \mathbb{C}); {}^t X = X\}$. Then the action $\rho = 2A_1$ of $GL(n)$ on V is given by $\rho(g)X = gXg^t$ for $g \in GL(n)$, $X \in V$. It is well-known that there exist $(n+1)$ -orbits $\rho(G)X_\nu = \{X \in V; \text{rank } X = \nu\}$ where

$$X_\nu = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & \ddots & & & \\ & & & 1 & & \\ & & & & 0 & \\ & 0 & & & & \ddots \\ & & & & & & 0 \end{pmatrix} \text{ for } \nu = 0, 1, \dots, n.$$

The relative invariant is given by the determinant $f(X) = \det X$. If we identify the dual V^* of V with V by $\langle X, Y \rangle = \text{tr } XY$, we have $\rho^*(g)Y = {}^t g^{-1} Y g^{-1}$ for $g \in GL(n)$. We have

$$(2.2) \quad d\rho(A)X_\nu = AX_\nu + X_\nu {}^t A = \left(\begin{array}{c|c} A_1 + {}^t A_1 & {}^t A_3 \\ \hline A_3 & 0 \end{array} \right)$$

$$\text{for } A = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix} \in \mathfrak{gl}(n).$$

Therefore we have

$$(2.3) \quad \mathfrak{g}_{X_\nu} = \left\{ \begin{pmatrix} A_1 & A_2 \\ 0 & A_4 \end{pmatrix}; {}^t A_1 = -A_1, A_1 \in M(\nu), A_2 \in M(\nu, n-\nu), \right.$$

$$\left. A_4 \in M(n-\nu) \right\}.$$

Since $\dim \mathfrak{g}_{X_\nu} = n(n-\nu) + (\nu(\nu-1)/2)$, we have $\dim \rho(G)X_\nu = n\nu - (\nu(\nu-1)/2)$. The conormal vector space $V_{X_\nu}^*$ is given by

$$(2.4) \quad V_{X_\nu}^* = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & W_\nu \end{pmatrix}; {}^t W_\nu = W_\nu, W_\nu \in M(n-\nu) \right\}.$$

Since $d\rho^*\left(\begin{pmatrix} A_1 & A_2 \\ 0 & A_4 \end{pmatrix}\right)\begin{pmatrix} 0 & 0 \\ 0 & W_\nu \end{pmatrix} = \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & -{}^t A_4 W_\nu - W_\nu A_4 \end{array}\right)$, the colocalization $(G_{X_\nu}, \rho_{X_\nu}, V_{X_\nu}^*)$ at X_ν is isomorphic to $(GL(n-\nu), 2A_1, V(\frac{1}{2}(n-\nu)(n-\nu+1)))$. Put $Y_\nu = \begin{pmatrix} 0 & 0 \\ 0 & I_{n-\nu} \end{pmatrix}$ ($\nu = 0, 1, \dots, n$). Then Y_ν is a generic point of the

colocalization at X_ν and $Y_{\nu+1}$ is a point of the unique one-codimensional orbit. Thus we have $\dim A_\nu \cap A_{\nu+1} = \dim V - 1$, where A_ν denotes the conormal bundle of $\rho(G)X_\nu$. Since $\dim d\rho^*(\mathfrak{g}_{X_\nu} \cap \mathfrak{g}_0)Y_{\nu+1} = \dim d\rho^*(\mathfrak{g}_{X_\nu})Y_{\nu+1}$, we have $\dim \mathfrak{g}_0(X_\nu, Y_{\nu+1}) = (n(n+1)/2) - 1$, and hence A_ν is a good holonomic variety by Proposition 1-5 ($\nu = 0, 1, \dots, n$).

Put $A_\nu^\beta = \left(\begin{array}{c|c|c} 0 & & \\ \hline & -\beta & \\ \hline & & -\frac{1}{2}I_{n-\nu-1} \\ \hline \end{array} \right) \in \mathfrak{gl}(n)$. Then we have $d\rho(A_\nu^\beta)X_\nu = 0$,

$d\rho^*(A_\nu^\beta)Y_{\nu+1} = Y_{\nu+1}$ and $2\beta = \text{tr } A_\nu^\beta$ where tr denotes the trace of A_ν^β on $V_{X_\nu}^*$ modulo $d\rho_{X_\nu}(\mathfrak{g}_{X_\nu})Y_{\nu+1}$. Hence, A_ν and $A_{\nu+1}$ intersect regularly, i.e., the intersection exponent of A_ν and $A_{\nu+1}$ equals $(1:0)$. We shall calculate the order $\text{ord}_{A_\nu} f^s$ by Proposition 1-3. Put $A_\nu = \begin{pmatrix} 0 & 0 \\ 0 & -\frac{1}{2}I_{n-\nu} \end{pmatrix}$ ($0 \leq \nu \leq n$). Then $d\rho(A_\nu)X_\nu = 0$ and $d\rho^*(A_\nu)Y_\nu = Y_\nu$. Since $\delta\chi(A_\nu) = 2 \text{tr } A_\nu = -(n-\nu)$, and $\text{tr}_{V_{X_\nu}^*} d\rho_{X_\nu}(A_\nu) = \dim V_{X_\nu}^* = \frac{1}{2}(n-\nu)(n-\nu+1)$, we have $\text{ord}_{A_\nu} f^s = s\delta\chi(A_\nu) - \text{tr}_{V_{X_\nu}^*} d\rho_{X_\nu}(A_\nu) + \frac{1}{2}\dim V_{X_\nu}^* = -(n-\nu)s - \frac{1}{4}(n-\nu)(n-\nu+1)$.

Thus we obtain the holonomy diagram (Figure 2-2). By Corollary




Figure 2-2. Holonomy diagram of $(GL(n), 2A_1, V(\frac{1}{2}n(n+1)))$ ($n \geq 2$)

1-2, we have $b_{A_\nu}(s)/b_{A_{\nu+1}}(s) = s + ((n - \nu + 1)/2)$ ($0 \leq \nu \leq n - 1$), and hence $b(s) = b_{A_0}(s) = \prod_{\nu=1}^n (s + (\nu + 1)/2)$.

Remark. The b -function of 2-2 is also already known. It can be obtained by using Capelli's identity or by a direct calculation of the Fourier transform of $f(x)^*$.

2-3. ($GL(2m), A_2, V(m(2m - 1))$) ($m \geq 3$)

The representation space can be identified with $V_m = \{X \in M(2m, \mathbb{C}) \mid {}^tX = -X\}$. Then the action $\rho = A_2$ is given by $\rho(g)X = gX{}^tg$ for $g \in GL(2m)$, $X \in V_m$. The relative invariant $f(X)$ is the Pfaffian of X . It is well-known that there exists $(m + 1)$ -orbits $\rho(G)X_\mu = \{X \in V_m; \text{rank } X = 2\mu\}$

$$\text{where } X_\mu = \begin{pmatrix} 0 & 0 & I_\mu & 0 \\ 0 & 0 & 0 & 0 \\ -I_\mu & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (0 \leq \mu \leq m).$$

By simple calculation, we have

$$(2.5) \quad d\rho(\tilde{A})X_\mu = \begin{pmatrix} \overbrace{{}^tB_1 - B_1}^\mu & \overbrace{{}^tB_3}^{m-\mu} & \overbrace{A_1 + {}^tD_1}^\mu & \overbrace{{}^tD_3}^{m-\mu} \\ -B_3 & & A_3 & \\ -D_1 - {}^tA_1 & -{}^tA_3 & C_1 - {}^tC_1 & -{}^tC_3 \\ -D_3 & & C_3 & \end{pmatrix}$$

$$\text{where } \tilde{A} = \begin{pmatrix} A_1 & A_2 & B_1 & B_2 \\ A_3 & A_4 & B_3 & B_4 \\ C_1 & C_2 & D_1 & D_2 \\ C_3 & C_4 & D_3 & D_4 \end{pmatrix}$$

and hence,

$$(2.6) \quad \mathfrak{g}_{X_\mu} = \left\{ \begin{pmatrix} A_1 & A_2 & B_1 & B_2 \\ 0 & A_4 & 0 & B_4 \\ C_1 & C_2 & -{}^tA_1 & D_2 \\ 0 & C_4 & 0 & D_4 \end{pmatrix}; {}^tB_1 = B_1, {}^tC_1 = C_1 \right\}$$

$$V_{X_\mu}^* = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & X & 0 & Y \\ 0 & 0 & 0 & 0 \\ 0 & -{}^tY & 0 & Z \end{pmatrix}; {}^tX = -X, {}^tZ = -Z \right\} \cong \left\{ \begin{pmatrix} X & Y \\ {}^tY & Z \end{pmatrix}; {}^tX = -X, {}^tZ = -Z \right\} = V_{m-\mu}.$$

Since \mathfrak{g}_{X_μ} acts on $V_{X_\mu}^*$ as $\tilde{X} \mapsto -{}^t\tilde{A}_4\tilde{X} - \tilde{X}\tilde{A}_4$ where $\tilde{A}_4 = \begin{pmatrix} A_4 & B_4 \\ C_4 & D_4 \end{pmatrix}$ and

$\tilde{X} = \left(\begin{array}{c|c} X & Y \\ \hline -{}^t Y & Z \end{array} \right)$ with ${}^t X = -X$, ${}^t Z = -Z$, the colocalization $(G_{X_\mu}, \rho_{X_\mu}, V_{X_\mu}^*)$ at X_μ is isomorphic to $(GL(2m - 2\mu), A_2, V((m - \mu)(2m - 2\mu - 1)))$. Here we identified the dual V_m^* of V_m with V_m by $\langle X, Y \rangle = \text{tr } XY$.

$$\text{Put } Y_\mu = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{m-\mu} \\ 0 & 0 & 0 & 0 \\ 0 & -I_{m-\mu} & 0 & 0 \end{pmatrix} \quad (0 \leq \mu \leq m).$$

Then Y_μ is a generic point of the colocalization $(G_{X_\mu}, \rho_{X_\mu}, V_{X_\mu}^*)$ at X_μ , and $Y_{\mu+1}$ is a point of the one-dimensional orbit and hence we have $\dim A_\mu \cap A_{\mu+1} = \dim V - 1$ ($0 \leq \mu \leq m - 1$) where A_μ denotes the conormal bundle of $\rho(G)X_\mu$.

By (2.6), we have $\mathfrak{g}_{X_\mu} \not\subset \mathfrak{g}_0$ for $\mu \neq m$, and hence $\dim d\rho(\mathfrak{g})X_\mu = \dim d\rho(\mathfrak{g}_0)X_\mu$ for $\mu \neq m$. Applying this fact to the colocalization at X_μ , we have $\dim \mathfrak{g}_0(X_\mu, Y_{\mu+1}) = \dim \mathfrak{g}(X_\mu, Y_{\mu+1}) = m(2m - 1) - 1$. This implies that A_μ is a good holonomic variety by Proposition 1-5.


$$\text{Put } A_\mu^\beta = \left(\begin{array}{c|cc|c} 0 & & 0 & 0 \\ \hline 0 & -\beta & 0 & 0 \\ & -\frac{1}{2}I_{m-\mu-1} & & \\ \hline 0 & & 0 & 0 \\ \hline 0 & & 0 & -\beta \\ & & & -\frac{1}{2}I_{m-\mu-1} \end{array} \right) \quad \text{for } \beta \in \mathbb{C}.$$

Then we have $d\rho(A_\mu^\beta)X_\mu = 0$, $d\rho^*(A_\mu^\beta)Y_{\mu+1} = Y_{\mu+1}$ and $\text{tr } A_\mu^\beta = 2\beta$ where tr denotes the trace of A_μ^β on $V_{X_\mu}^*$ modulo $d\rho(\mathfrak{g}_{X_\mu})Y_{\mu+1}$, and hence by Proposition 1-4, A_μ and $A_{\mu+1}$ intersect regularly, i.e., the intersection exponent of A_μ and $A_{\mu+1}$ equals $(1:0)$. We shall calculate the order $\text{ord}_{A_\mu} f^s$.

$$\text{Put } A_0 = \left(\begin{array}{c|cc|c} & & & \\ \hline & -\frac{1}{2}I_{m-\mu} & & \\ \hline & & & \\ \hline & & & -\frac{1}{2}I_{m-\mu} \end{array} \right). \quad \text{Then we have } d\rho(A_0)X_\mu = 0 \text{ and}$$

$d\rho^*(A_0)Y_\mu = Y_\mu$. Since $\delta\chi(A_0) = -(m - \mu)$, $\text{tr}_{V_{X_\mu}^*} A_0 = \dim V_{X_\mu}^* = (m - \mu)(2m - 2\mu - 1)$, we have $\text{ord}_{A_\mu} f^s = s\delta\chi(A_0) - \text{tr}_{V_{X_\mu}^*} A_0 + \frac{1}{2} \dim V_{X_\mu}^* = -(m - \mu)s - \frac{1}{2}(m - \mu)(2m - 2\mu - 1)$.

By Corollary 1-2, we have $b_{A_\mu}(s)/b_{A_{\mu+1}}(s) = s + 2(m - \mu) - 1$ ($0 \leq \mu \leq m - 1$). Hence we obtain the holonomy diagram (Figure 2-3) and b -function $b(s) = \prod_{\mu=0}^{m-1} (s + 2(m - \mu) - 1) = \prod_{k=1}^m (s + 2k - 1)$.

Figure 2-3. Holonomy diagram of $(GL(2m), A_2, V(m(2m-1)))$ ($m \geq 3$).

Remark. These three P.V.'s have many common properties: (1) $(GL(m), 2A_1, V\left(\binom{m}{2}\ell + m\right))$ with $\ell = 1$ (2) $(SL(m) \times GL(m), A_1 \otimes A_1, V\left(\binom{m}{2}\ell + m\right))$ with $\ell = 2$ (3) $(GL(2m), A_2, V\left(\binom{m}{2}\ell + m\right))$ with $\ell = 4$. They have $(m+1)$ -orbits and their relative invariants are of degree m of $\binom{m}{2}\ell + m$ variables. We denote (A) by (μ) if A is the conormal bundle of a μ -codimensional orbit. Then their holonomy diagrams are as in Figure 2-4.

§ 3. $(Sp(n) \times GL(2m), A_1 \otimes A_1, V(2n) \otimes V(2m))$ with $n \geq 2m$

The representation space V can be identified with the totality of $2n \times 2m$ matrices. Then the action $\rho = A_1 \otimes A_1$ is given by $\rho(g)X = g_1 X' g_2$ for $g = (g_1, g_2) \in G = Sp(n) \times GL(2m)$, $X \in V$. Let X be an element of V such that $\text{rank } X = \nu$ and $\text{rank } {}^t X J X = 2\mu (2m \geq \nu \geq 2\mu \geq 0)$ where $J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$. Then by the action of $GL(2m)$, we may assume that $X = (X', 0)$ with $X' \in M(2n, \nu)$ satisfying ${}^t X' J X' = \begin{pmatrix} 0 & 0 & I_\mu \\ 0 & 0 & 0 \\ -I_\mu & 0 & 0 \end{pmatrix}$. Put $X_{\nu, 2\mu}$ as follows.





Figure 2-4

$$(3.1) \quad X_{\nu, 2\mu} = \left(\begin{array}{ccc|ccc} I_{\mu} & & & & & \\ & I_{\nu-2\mu} & & & & \\ & & & & & \\ & & & I_{\mu} & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \end{array} \right) \left. \begin{array}{l} \\ \\ \\ \\ \\ \\ \\ \end{array} \right\} \begin{array}{l} n \\ \\ \\ \\ \\ \\ \\ \end{array}$$

$\nu-2\mu \left\{ \begin{array}{l} \\ \\ \\ \\ \\ \\ \end{array} \right\} \begin{array}{l} n \\ \\ \\ \\ \\ \\ \\ \end{array}$

$\underbrace{\hspace{10em}}_{2m-\nu}$

Then $X_{\nu, 2\mu}$ satisfies the same condition as X and hence there exists an element g_1 of $Sp(n)$ satisfying $g_1 X = X_{\nu, 2\mu}$. This implies that $S_{\nu, 2\mu} = \{X \in V; \text{rank } X = \nu, \text{rank } {}^t X J X = 2\mu\}$ ($2m \geq \nu \geq 2\mu \geq 0$) consists of a single G -orbit, and we complete the orbital decomposition of this space. Put $A \in \mathfrak{sp}(n)$ and $D \in \mathfrak{gl}(2m)$ as follows:

$$(3.2) \quad A = \left(\begin{array}{ccc|ccc} A_1 & A_{12} & A_{13} & B_1 & B_{12} & B_{13} \\ A_{21} & A_2 & A_{23} & {}^t B_{12} & B_2 & B_{23} \\ A_{31} & A_{32} & A_3 & {}^t B_{13} & {}^t B_{23} & B_3 \\ \hline C_1 & C_{12} & C_{13} & -{}^t A_1 & -{}^t A_{21} & -{}^t A_{31} \\ {}^t C_{12} & C_2 & C_{23} & -{}^t A_{12} & -{}^t A_2 & -{}^t A_{32} \\ {}^t C_{13} & {}^t C_{23} & C_3 & -{}^t A_{13} & -{}^t A_{23} & -{}^t A_3 \end{array} \right),$$

$\underbrace{\hspace{2em}}_{\mu} \quad \underbrace{\hspace{2em}}_{\nu-2\mu} \quad \underbrace{\hspace{2em}}_{n-\nu+\mu} \quad \underbrace{\hspace{2em}}_{\mu} \quad \underbrace{\hspace{2em}}_{\nu-2\mu} \quad \underbrace{\hspace{2em}}_{n-\nu+\mu}$

$$D = \left(\begin{array}{cccc} D_1 & D_{12} & D_{13} & D_{14} \\ D_{21} & D_2 & D_{23} & D_{24} \\ D_{31} & D_{32} & D_3 & D_{34} \\ \hline D_{41} & D_{42} & D_{43} & D_4 \end{array} \right)$$

$\underbrace{\hspace{2em}}_{\mu} \quad \underbrace{\hspace{2em}}_{\nu-2\mu} \quad \underbrace{\hspace{2em}}_{\mu} \quad \underbrace{\hspace{2em}}_{2m-\nu}$

where ${}^t B_i = B_i, {}^t C_i = C_i$ ($i = 1, 2, 3$).

Then, for $\tilde{A} = A \oplus D \in \mathfrak{g}$, we have

$$(3.3) \quad \begin{aligned} d\rho(\tilde{A})X_{\nu, 2\mu} &= AX_{\nu, 2\mu} + X_{\nu, 2\mu} {}^t D \\ &= \left(\begin{array}{cc|cc} A_1 + {}^t D_1 & A_{12} + {}^t D_{21} & B_1 + {}^t D_{31} & {}^t D_{41} \\ A_{21} + {}^t D_{12} & A_2 + {}^t D_2 & {}^t B_{12} + {}^t D_{32} & {}^t D_{42} \\ \hline A_{31} & A_{32} & {}^t B_{13} & 0 \\ C_1 + {}^t D_{13} & C_{12} + {}^t D_{23} & -{}^t A_1 + {}^t D_3 & {}^t D_{43} \\ {}^t C_{12} & C_2 & -{}^t A_{12} & 0 \\ {}^t C_{13} & {}^t C_{23} & -{}^t A_{13} & 0 \end{array} \right) \end{aligned}$$

and hence the isotropy subalgebra $\mathfrak{g}_{X_{\nu, 2\mu}}$ is given as follows:

$$\begin{aligned}
 \mathfrak{g}_{X_{\nu, 2\mu}} = & \left\{ \tilde{A} = \left[\begin{array}{ccc|ccc} A_1 & 0 & 0 & B_1 & B_{12} & 0 \\ A_{21} & A_2 & A_{23} & {}^t B_{12} & B_2 & B_{23} \\ 0 & 0 & A_3 & 0 & {}^t B_{23} & B_3 \\ \hline C_1 & 0 & 0 & -{}^t A_1 & -{}^t A_{21} & 0 \\ 0 & 0 & 0 & 0 & -{}^t A_2 & 0 \\ 0 & 0 & C_3 & 0 & -{}^t A_{23} & -{}^t A_3 \end{array} \right] \right. \\
 & \oplus \left. \left[\begin{array}{cc|cc} -{}^t A_1 & -{}^t A_{21} & -C_1 & D_{14} \\ 0 & -{}^t A_2 & 0 & D_{24} \\ \hline -B_1 & -B_{12} & A_1 & D_{34} \\ 0 & 0 & 0 & D_4 \end{array} \right] \right\} \\
 (3.4) \quad & \cong \left\{ \left[\begin{array}{cc|cc|c} A_2 & A_{21} & {}^t B_{12} & A_{23} & B_{23} & B_2 \\ \hline 0 & A_1 & B_1 & & 0 & B_{12} \\ C_1 & -{}^t A_1 & & & & -{}^t A_{21} \\ \hline 0 & & 0 & A_3 & B_3 & {}^t B_{23} \\ & & & C_3 & -{}^t A_3 & -{}^t A_{23} \\ \hline 0 & & 0 & & 0 & -{}^t A_2 \end{array} \right] \right. \\
 & \oplus \left. \left[\begin{array}{cc|cc} A_1 & B_1 & -B_{12} & D_{34} \\ C_1 & -{}^t A_1 & {}^t A_{21} & -D_{14} \\ \hline 0 & & -{}^t A_2 & D_{24} \\ & & 0 & D_4 \end{array} \right] \right\} \\
 & \cong (\mathfrak{gl}(\nu - 2\mu) \oplus \mathfrak{gl}(2m - \nu) \oplus \mathfrak{sp}(\mu) \oplus \mathfrak{sp}(n - \nu + \mu)) \oplus \mathfrak{u}(k)
 \end{aligned}$$

where $\mathfrak{u}(k)$ denotes the Lie algebra of a k -dimensional unipotent group with $k = \frac{1}{2}(4n + 1)(\nu - 2\mu) - \frac{3}{2}(\nu - 2\mu)^2 + \nu(2m - \nu)$. In this paper, we make a convention that the first (resp. second) \oplus implies the direct sum as Lie algebras (resp. vector spaces) for $(\mathfrak{g}_1 \oplus \mathfrak{g}_2) \oplus \mathfrak{g}_3$.

We identify the dual space V^* of V with V by $\langle X, Y \rangle = \text{tr } X^t Y$ for $X, Y \in V = M(2n, 2m)$, and hence we have $\rho^*(g)Y = {}^t g_1^{-1} Y g_2^{-1}$ for $g = (g_1, g_2) \in G$, $Y \in V$ and $d\rho^*(\tilde{A})Y = -{}^t A Y - Y D$ for $\tilde{A} = (A, D) \in \mathfrak{g}$. From (3.3), the conormal vector space $V_{X_{\nu, 2\mu}}^*$ is given by

$$(3.5) \quad V_{X_{\nu, 2\mu}}^* = \left\{ \tilde{Y} = \left[\begin{array}{ccc|c} & & & 0 \\ & 0 & 0 & 0 \\ & & & Y \\ & & 0 & 0 \\ \hline & 0 & X & Z \\ & & & W \end{array} \right] ; {}^t X = -X \right\}$$

$\underbrace{\quad}_{\mu} \quad \underbrace{\quad}_{\nu-2\mu} \quad \underbrace{\quad}_{\mu} \quad \underbrace{\quad}_{2m-\nu}$

$$\cong \left\{ \tilde{Y}' = \left(\underbrace{\begin{array}{c|c} Y_1 & Y_2 \\ \hline 0 & Y_3 \end{array}}_{\substack{\nu-2\mu \\ 2m-\nu}} \right)^{\nu-2\mu} ; {}^t Y_1 = -Y_1 \right\}.$$

Here the isomorphism is obtained by putting $Y_1 = X$, $Y_2 = Z$ and $Y_3 = \begin{bmatrix} Y \\ W \end{bmatrix}$. Then the action $d\rho_{X\nu, 2\mu}$ of $\mathfrak{g}_{X\nu, 2\mu}$ on $V_{X\nu, 2\mu}^*$ is given as follows.

$$d\rho_{X\nu, 2\mu}(\tilde{A})\tilde{Y}' = \left(\begin{array}{c|cc} A_2 & -B_{23} & A_{23} \\ \hline 0 & -{}^t A_3 & -C_3 \\ & -B_3 & A_3 \end{array} \right) \left(\begin{array}{c|c} Y_1 & Y_2 \\ \hline 0 & Y_3 \end{array} \right) + \left(\begin{array}{c|c} Y_1 & Y_2 \\ \hline 0 & Y_3 \end{array} \right) \left(\begin{array}{c|c} {}^t A_2 & -D_{24} \\ \hline 0 & -D_4 \end{array} \right).$$

Thus the action on Y_1 -space is isomorphic to $(GL(\nu - 2\mu), A_2, V(\frac{1}{2}(\nu - 2\mu) \times (\nu - 2\mu - 1)))$ and the action on Y_3 -space is isomorphic to $(Sp(n - \nu + \mu) \times GL(2m - \nu), A_1 \otimes A_1, V(2n - 2\nu + 2\mu) \otimes V(2m - \nu))$. First we shall consider the case when ν is even, i.e., $\nu = 2\nu'$. Let \tilde{Y}_0 be an element of $V_{X\nu, 2\mu}^*$ with $X = \begin{pmatrix} 0 & I_{\nu'-\mu} \\ -I_{\nu'-\mu} & 0 \end{pmatrix}$, $Y = \begin{pmatrix} I_{m-\nu'} & 0 \\ 0 & 0 \end{pmatrix}$, $W = \begin{pmatrix} 0 & I_{m-\nu} \\ 0 & 0 \end{pmatrix}$ and $Z = 0$ in (3.5). Then \tilde{Y}_0 is a generic point and $\tilde{Y}_0 \in S_{2m-2\mu, 2m-2\nu'}^*$, i.e., $A_{2\nu', 2\mu} = A_{2m-2\mu, 2m-2\nu'}^*$ where $A_{\nu, 2\mu}$ (resp. $A_{\nu, 2\mu}^*$) denotes the conormal bundle of $S_{\nu, 2\mu}$ (resp. $S_{\nu, 2\mu}^*$). We shall calculate the order $\text{ord}_{A_{2\nu', 2\mu}} f^s$ where $f(X) = Pf^t X J X$. Let \tilde{A}_0 be the element of $\mathfrak{g}_{X\nu, 2\mu}$ with $A_2 = \frac{1}{2}I_{2(\nu'-\mu)}$, $D_4 = -I_{2(m-\nu')}$, all remaining parts zero in (3.4). Then we have $d\rho(\tilde{A}_0)X_{\nu, 2\mu} = 0$ and $d\rho^*(\tilde{A}_0)\tilde{Y}_0 = \tilde{Y}_0$. Since $\delta\chi(\tilde{A}_0) = -(2m - \nu' - \mu)$, $\text{tr}_{V_{X\nu, 2\mu}^*} \tilde{A}_0 = (\nu' - \mu)(2\nu' - 2\mu - 1) + 4(m - \nu')(n - 2\nu' + \mu) + 6(m - \nu')(\nu' - \mu)$ and $\dim V_{X\nu, 2\mu}^* = (\nu' - \mu)(2\nu' - 2\mu - 1) + 4(m - \nu')(n - 2\nu' + \mu) + 4(m - \nu')(\nu' - \mu)$, we have

$$(3.6) \quad \text{ord}_{A_{2\nu', 2\mu}} f^s = -(2m - \nu' - \mu)s - \frac{1}{2}(\nu' - \mu)(2\nu' - 2\mu - 1) - 2(m - \nu')(n - 2\nu' + \mu) - 4(m - \nu')(\nu' - \mu).$$

Let \tilde{Y}_1 be the element of $V_{X\nu, 2\mu}^*$ with $X = \begin{pmatrix} 0 & I_{\nu'-\mu} \\ -I_{\nu'-\mu} & 0 \end{pmatrix}$, $Y = \begin{pmatrix} I_{m-\nu'+1} & 0 \\ 0 & 0 \end{pmatrix}$, $W = \begin{pmatrix} 0 & I_{m-\nu'-1} \\ 0 & 0 \end{pmatrix}$ and $Z = 0$. Since \tilde{Y}_1 is a point of a one-codimensional orbit and $\tilde{Y}_1 \in S_{2m-2\mu, 2(m-\nu'-1)}^*$, we have $A_{2\nu', 2\mu} \cap A_{2(\nu'+1), 2\mu} = \dim V - 1$. They intersect regularly. By Corollary 1-2, we have

$$(3.7) \quad b_{A_{2\nu', 2\mu}}(s)/b_{A_{2(\nu'+1), 2\mu}}(s) = s + 2n - 2\nu' \quad (m - 1 \geq \nu' \geq 0).$$

Now let \tilde{Y}_2 be the element of $V_{X\nu, 2\mu}^*$ with $X = \left(\begin{array}{c|c} 0 & I_{\nu'-\mu-1} \\ \hline -I_{\nu'-\mu-1} & 0 \end{array} \middle| \begin{array}{c} 0 \\ 0 \end{array} \right)$, $Y = \begin{pmatrix} I_{m-\nu'} & 0 \\ 0 & 0 \end{pmatrix}$, $W = \begin{pmatrix} 0 & I_{m-\nu'} \\ 0 & 0 \end{pmatrix}$ and $Z = 0$. Since \tilde{Y}_2 is a point of the

other one-codimensional orbit and $\tilde{Y}_2 \in S_{2(m-\mu-1), 2(m-\nu')}$, we have $\dim A_{2\nu', 2\mu} \cap A_{2\nu', 2(\mu+1)} = \dim V - 1$. They intersect regularly. By Corollary 1-2, we have

$$(3.8) \quad b_{A_{2\nu', 2\mu}}(s)/b_{A_{2\nu', 2(\mu+1)}} = s + 2m - 2\mu - 1 \quad (m - 1 \geq \mu \geq 0).$$

Now we shall show that $A_{\nu, 2m}$ is not a good holonomic variety when ν is odd, i.e., $\nu = 2\nu' + 1$. Let \tilde{Y}_0 be the element of $V_{X_{\nu, 2\mu}}^*$ with $X = \left(\begin{array}{c|c} 0 & I_{\nu'-\mu} \\ \hline -I_{\nu'-\mu} & 0 \end{array} \right)$, $Y = \begin{pmatrix} I_{m-\nu'-1} & 0 \\ 0 & 0 \end{pmatrix}$, $W = \begin{pmatrix} 0 & I_{m-\nu'} \\ 0 & 0 \end{pmatrix}$ and $Z = 0$ in (3.5).

Then it is a generic point of the conormal vector space. Let \tilde{A}_0 be the element of $\mathfrak{g}_{V_{\nu, 2\mu}}$ with $A_2 = \begin{pmatrix} \frac{1}{2}I_{2(\nu'-\mu)} & 0 \\ 0 & \beta \end{pmatrix}$, $D_4 = -I_{2m-\nu}$, all remaining parts zero. Then we have $d\rho(\tilde{A}_0)X_{\nu, 2\mu} = 0$ and $d\rho^*(\tilde{A}_0)\tilde{Y}_0 = \tilde{Y}_0$. Therefore, if $A_{\nu, 2\mu}$ is a good holonomic variety, $m_{A_{\nu, 2\mu}} = -\delta\chi(\tilde{A}_0) = 2m - \nu' - \mu - 1 + \beta$ is a non-negative integer which is a contradiction. Thus we obtain the following proposition.

PROPOSITION 3-1. *The irreducible regular P.V. $(Sp(n) \times GL(2m), A_1 \otimes A_1, V(2n) \otimes V(2m))$ ($n \geq 2m$) has finitely many orbits $S_{\nu, 2\mu} = \{X \in M(2n, 2m); \text{rank } X = \nu, \text{rank } {}^tXJX = 2\mu\}$ ($2m \geq \nu \geq 2\mu \geq 0$). When ν is odd, the conormal bundle $A_{\nu, 2\mu}$ of $S_{\nu, 2\mu}$ is outside W , i.e., $A_{\nu, 2\mu}$ is not a good holonomic variety. When ν is even ($\nu = 2\nu'$), $A_{\nu, 2\mu}$ is a good holonomic variety and $\text{ord}_{A_{\nu, 2\mu}} f^s = -(2m - \nu' - \mu)s - \frac{1}{2}(\nu' - \mu)(2\nu' - 2\mu - 1) - 2(m - \nu')(n - 2\nu' + \mu) - 4(m - \nu')(\nu' - \mu)$. We have $\dim A_{\nu, 2\mu} \cap A_{\nu, 2(\mu+1)} = \dim A_{\nu, 2\mu} \cap A_{\nu+2, 2\mu} = \dim V - 1$. The b-function $b(s)$ is given by $b(s) = \prod_{k=1}^m (s + 2k - 1) \cdot \prod_{\ell=0}^{m-1} (s + 2n - 2\ell)$.*




Figure 3-1. (ν : even)




Figure 3-2. Holonomy diagram of $(Sp(n) \times GL(2m), A_1 \otimes A_1, V(2n) \otimes V(2m))$ with $n \geq 2m$.

§ 4. ($\text{Spin}(10) \times \text{GL}(2)$, half-spin rep. $\otimes A_1$, $V(16) \otimes V(2)$)

The representation space $V(16) \otimes V(2)$ is identified with $V = V(16) \oplus V(16)$ where $V(16)$ is spanned by $1, e_i e_j, e_k e_\ell e_m e_n$ ($1 \leq i < j \leq 5, 1 \leq k < \ell < m < n \leq 5$) (See p. 110–112 in [1]). The action $\rho = \rho_1 \otimes A_1$ is given by $\rho(g)x = (\rho_1(g_1)X, \rho_1(g_1)Y)'g_2$ for $g = (g_1, g_2) \in \text{Spin}(10) \times \text{GL}(2)$, $x = (X, Y) \in V = V(16) \oplus V(16)$ where ρ_1 denotes the even half-spin representation of $\text{Spin}(10)$ on $V(16)$. First of all, we shall complete the orbital decomposition of this space. J-I. Igusa completed the orbital decomposition of $(\text{Spin}(10), \rho_1, V(16))$ (See [3]). There exist three orbits $S'_m = \rho_1(\text{Spin}(10)) \cdot x'_m$ ($m = 0, 5, 16$) where S'_m denotes the m -codimensional $\text{Spin}(10)$ -orbit and $x'_0 = 1 + e_1 e_2 e_3 e_4$, $x'_5 = 1$, $x'_{16} = 0$. If $\lambda \in \mathbb{C}^\times$, for any index i satisfying $1 \leq i \leq 5$, we put $S_i(\lambda) = \lambda^{-1} + (\lambda - \lambda^{-1})e_i f_i$. Then $S_i(\lambda)$ is an element of $\text{Spin}(10)$. For any two distinct indices i, j satisfying $1 \leq i, j \leq 10, j \neq i + 5, i \neq j + 5$, we put $S_{ij}(\lambda) = 1 + \lambda e_i e_j = \exp(\lambda e_i e_j)$ where $e_k = f_{k-5}$ for $6 \leq k \leq 10$ (See [1], [3]). Then $S_{ij}(\lambda)$ is an element of $\text{Spin}(10)$ satisfying $S_{ij}(\lambda)S_{ji}(\lambda) = 1$.

PROPOSITION 4-1. *The triplet $(\text{Spin}(10) \times \text{GL}(2)$, half-spin rep. $\otimes A_1$, $V(16) \otimes V(2)$) has nine orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 4, 8, 9, 13, 15, 20, 32$) where S_m denotes the m -codimensional orbit.*

- (1) $x_0 = (1 + e_1 e_2 e_3 e_4, e_1 e_5 + e_2 e_3 e_4 e_5)$
- (2) $x_1 = (1 + e_1 e_2 e_3 e_4, e_1 e_2 + e_2 e_3 e_4 e_5)$
- (3) $x_4 = (1, e_1 e_5 + e_2 e_3 e_4 e_5)$
- (4) $x_8 = (1, e_1 e_2 e_3 e_4)$
- (5) $x_9 = (1, e_1 e_2 + e_3 e_4)$
- (6) $x_{13} = (1, e_1 e_2)$
- (7) $x_{15} = (1 + e_1 e_2 e_3 e_4, 0)$
- (8) $x_{20} = (1, 0)$
- (9) $x_{32} = (0, 0)$

Proof. Let $\tilde{x} = (x, y)$ be a representative of one of the orbits of $V = V(16) \oplus V(16)$. Then we may assume that $x = 0, 1$, or $1 + e_1 e_2 e_3 e_4$ by the action of $\text{Spin}(10)$. If $x = 0$, then we have also $y = 1 + e_1 e_2 e_3 e_4, 1, 0$, i.e., (7), (8), (9) respectively. Note that we can exchange x and y in $\tilde{x} = (x, y)$ by the action of $\text{GL}(2)$. Assume that $x = 1$. We may put $y = y_0 + y_2 + y_4 \neq 0$ where $y_0 = y_0 \cdot 1$, $y_2 = \sum y_{ij} e_i e_j$ and $y_4 = \sum y_{rstu} e_r e_s e_t e_u$. We may assume that $y_0 = 0$ by the action of $\begin{pmatrix} 1 & 0 \\ -y_0 & 1 \end{pmatrix}$. If $y = y_2 \neq 0$, we may assume that $y_{12} = 1$ by the action of some $S_{ij}(\lambda)$ ($i = 1, 2; j \geq 6$) and $\begin{pmatrix} 1 & 0 \\ 0 & \lambda \end{pmatrix}$ if necessary.

In this case, we have $y = e_1e_2 + y_{34}e_3e_4 + y_{35}e_3e_5 + y_{45}e_4e_5$ by $S_{j7}(-y_{1j})$ and $S_{j6}(y_{2j})$ for $j = 3, 4, 5$. If $y_{34} = y_{35} = y_{45} = 0$, we have (6), and otherwise we may assume that $y_{34} = 1, y_{35} = y_{45} = 0$ by the action of suitable elements of $\{S_{3,10}(\lambda), S_{4,10}(\lambda), S_{58}(\lambda), S_{59}(\lambda); \lambda \in \mathcal{C}\}$, i.e., (5). If $y_4 \neq 0$, we may assume that $y_4 = e_1e_2e_3e_4$. By the action of $S_{89}(y_{12})$ and $\begin{pmatrix} 1 & 0 \\ y_{12}y_{34} & 1 \end{pmatrix}$, we have $y_{12} = 0$. Similarly $y_{ij} = 0$ for $1 \leq i < j \leq 4$, and hence $y = \sum_{j=1}^4 y_{j5}e_je_5 + e_1e_2e_3e_4$. If $y_{j5} = 0$ for all $j = 1, \dots, 4$, we have (4). In the other case, we may assume that $y_{15} = 1$ and $y_{j5} = 0$ ($2 \leq j \leq 4$). By the action of $S_{56}(-1)$ and $S_{1,10}(1)$, we have (3).

Finally assume that $x = 1 + e_1e_2e_3e_4$. We may put $y = y_2 + y_4$. If $y_4 \neq 0$, we may assume that $y_4 = e_2e_3e_4e_5$ or $y_4 = e_1e_2e_3e_4$. In the former case, if $y_{15} \neq 0$, we may assume that $y_{15} = 1$ by the action of $S_1(\lambda)S_5(\lambda)S_2(\lambda^{-1})$ and λI_2 where $\lambda^4 \cdot y_{15} = 1$. Then by the action of $S_{j10}(-y_{1j}), S_{j6}(-y_{j5})$ ($j = 2, 3, 4$), $S_{9,10}(y_{23}), S_{8,10}(-y_{24})$ and $S_{7,10}(y_{34})$, we have (1). If $y_{15} = 0$, we may assume that $y_{35} = y_{45} = 0$ by $\{S_{28}(\lambda), S_{29}(\lambda), S_{37}(\lambda), S_{47}(\lambda); \lambda \in \mathcal{C}\}$. Then by $S_{8,10}(-y_{24})$ and $S_{9,10}(y_{23})$, we may assume that $y_{24} = y_{23} = 0$. By some $S_{39}(\lambda)$ and $S_{28}(\lambda)$, we may also assume that $y_{14} = 0$, i.e., $y = y_{12}e_1e_2 + y_{13}e_1e_3 + y_{34}e_3e_4 + y_{25}e_2e_5 + e_2e_3e_4e_5$. By the action of $S_{7,10}(y_{34}), \begin{pmatrix} 1 & 0 \\ y_{25}y_{34} & 1 \end{pmatrix}$ and $S_{1,10}(y_{25}y_{34})$, we have $y_{34} = 0$. By $S_{89}(y_{25})$ and $S_{12}(y_{25})$, we also have $y_{25} = 0$, i.e., $y = y_{12}e_1e_2 + y_{13}e_1e_3 + e_2e_3e_4e_5$, where we may assume that $y_{13} = 0$. If $y_{12} \neq 0$, we have (2). If $y_{12} = 0$, it is transferred to x_4 by $S_{12}(-1), S_{89}(-1), S_{34}(-1), S_{87}(-1), S_{17}(1), S_{26}(-1), S_{56}(-1), S_{1,10}(1), \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, S_{6,10}(1), S_{15}(1)$ and $\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$.

Now consider the latter case, i.e., $y_4 = e_1e_2e_3e_4$. If some of y_{j5} ($1 \leq j \leq 4$) is not zero, we may assume that $y = e_1e_5 + y_{23}e_2e_3 + e_1e_2e_3e_4$. If $y_{23} = 0$, it is transferred to x_4 by $S_{15}(1), \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, S_{56}(-1)$ and $S_{1,10}(1)$. If $y_{23} \neq 0$, we may assume that $y_{23} = 1$. In this case, it is transferred to x_1 by $S_{89}(1), S_{23}(1), S_{4,10}(-1), S_{78}(1), S_{14}(1), S_{46}(1), S_{19}(-1), S_{29}(-1), S_{47}(1)$. When all $y_{j5} = 0$ for $1 \leq j \leq 4$, $y = \sum_{1 \leq i < j \leq 4} y_{ij}e_i e_j + e_1e_2e_3e_4$. If all $y_{ij} = 0$ for $1 \leq i < j \leq 4$, it is transferred to x_8 by $\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$. In the other case, we may assume that $y = e_1e_2 + y_{34}e_3e_4 + e_1e_2e_3e_4$. By the action of $S_{67}(\lambda), S_{34}(\lambda)$ and $\begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}$ with $\lambda^2 - \lambda - y_{34} = 0$, we have $y = e_1e_2 + (1 + 2\lambda)e_1e_2e_3e_4$. If $(1 + 2\lambda) \neq 0$, it is transferred to x_8 by $S_{89}(\mu), S_{12}(\mu)$ and $\begin{pmatrix} 1 & -\mu \\ 0 & \mu \end{pmatrix}$ with $\mu = \frac{1}{1 + 2\lambda}$. If $(1 + 2\lambda) = 0$, it is equivalent to x_8 by $S_{67}(-1), S_{12}(-1), \begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix}, S_i(\sqrt{-1})$, and $\sqrt{-1} I_2$.

Finally consider the case $y = y_2$, i.e., $y_4 = 0$. Since $y \neq 0$, we may assume that $y = e_1 e_2 + y_{34} e_3 e_4 + \sum_{j=1}^4 y_{j5} e_j e_5$. If $y_{34} = y_{j5} = 0$ for $1 \leq j \leq 4$, it is equivalent to x_6 as we have already seen. If $y_{34} \neq 0$ and $y_{j5} = 0$ for $1 \leq j \leq 4$, it is transferred to x_8 by $S_{34}(\lambda)$, $\begin{pmatrix} 1 & -1/\lambda \\ 1 & 0 \end{pmatrix}$, $S_{12}(1/\lambda)$, $\begin{pmatrix} 1 & 0 \\ 0 & 1/2\lambda \end{pmatrix}$, $S_{67}(\lambda/2)$, $S_{89}(1/2\lambda)$, $\begin{pmatrix} 1 & 0 \\ 1/4 & 1 \end{pmatrix}$ with $\lambda^2 = y_{34}$. If some of y_{j5} ($1 \leq j \leq 4$) is not zero, y is equivalent to an element of the form $e_1 e_5 + y_{23} e_2 e_3 + y_{24} e_2 e_4 + y_{34} e_3 e_4$. If $y_{ij} = 0$ ($2 \leq i < j \leq 4$), it is equivalent to x_4 as we have already seen. In the other case, we have $y = e_1 e_5 + e_3 e_4$. By the action of $S_{26}(-1)$, $S_{17}(1)$, $S_{34}(-1)$, $S_{67}(-1)$, $S_{7,10}(1)$, $\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$, $S_{66}(-1)$, $S_{12}(-1)$, $S_{1,10}(-1)$, it is equivalent to x_1 . About the codimension of these orbits, we will see later. Q.E.D.

By the degree formula (See Proposition 15, § 4 in [1]), we know that there exists a relatively invariant irreducible polynomial $f(x, y)$ of degree four which is unique up to a constant multiple. We shall give an explicit form of $f(x, y)$ after H. Kawahara's work (Master Thesis in Japanese, University of Tokyo, 1974).

For an element $x = x_0 + \sum_{i < j} x_{ij} e_i e_j + \sum_k x_k^* e_k^*$ of $V(16)$ where $e_k e_k^* = e_1 e_2 e_3 e_4 e_5$ for $1 \leq k \leq 5$, let $X = (x_{ij})$ be the skew-symmetric matrix of degree five determined by x_{ij} , and X_i the skew-symmetric matrix of degree four obtained from $(-1)^i X$ by crossing out its i -th line and column ($1 \leq i \leq 5$). We denote by $\text{Pf}(Y)$ the Pfaffian of the skew-symmetric matrix $Y = (y_{ij})$ of degree four, i.e., $\text{Pf}(Y) = y_{12} y_{34} - y_{13} y_{24} + y_{14} y_{23}$. We define ten quadratic forms $Q_i(x)$ on $V(16)$ by $Q_i(x) = \sum_{j=1}^5 x_{ij} x_j^*$ and $Q_{i+5}(x) = x_0 x_i^* + \text{Pf}(X_i)$ for $1 \leq i \leq 5$.

PROPOSITION 4-2 (H. Kawahara).

(1) $\rho_1(\text{Spin}(10)) \cdot 1 = \{x \in V(16); Q_i(x) = 0 \ (1 \leq i \leq 10)\} - \{0\}$, where ρ_1 denotes the even half-spin representation. Moreover, this is the totality of pure spinors.

(2) The relative invariant $f(x, y)$ of $(\text{Spin}(10) \times GL(2), \rho_1 \otimes A_1, V(16) \oplus V(16))$ is given by $f(x, y) = \sum_{i=1}^5 B_i(x, y) B_{i+5}(x, y)$ for $(x, y) \in V(16) \oplus V(16)$ where $B_i(x, y) = Q_i(x + y) - Q_i(x) - Q_i(y)$ is the associated bilinear form of $Q_i(x)$ for $1 \leq i \leq 10$.

Proof. We shall use the same notation as in [4]. By simple calculation, we have $\beta_i(x, x) = (1/8) \sum_{i=1}^{10} Q_i(x) e_i$. Since $\beta_1(\rho_1(s)x, \rho_1(s)x) = \lambda(s) \cdot \zeta_1(\chi(s)) \cdot \beta_1(x, x)$ for $s \in \text{Spin}(10)$ where ζ_1 is the representation A_1 of $SO(10) = \chi(\text{Spin}(10))$ (See p. 90 in [4]), we have

$$(4.1) \quad \sum_{i=1}^{10} Q_i(\rho_1(s)x)e_i = \lambda(s) \cdot \zeta_1(\chi(s)) \cdot \sum_{i=1}^{10} Q_i(x)e_i.$$

This implies that $W = \{x \in V(16); Q_i(x) = 0, 1 \leq i \leq 10\}$ is a Spin(10)-invariant subspace. From the orbital decomposition, it is clear that $W = S'_5 \cup S'_{16}$, i.e., $S'_5 = W - \{0\}$. Since the totality of pure spinors in $V(16)$ is a single Γ^+ -orbit where Γ^+ denotes the even Clifford group, and $\beta_1(x, x) = 0$ for a pure spinor x (See [4]), we have (1). From (4.1), $F(x) = \sum_{i=1}^5 Q_i(x) Q_{i+5}(x)$ is invariant under the action ρ_1 of Spin(10) since $\tilde{f}(y) = \sum_{i=1}^5 y_i y_{i+5}$ for $y = \sum_{i=1}^{10} y_i e_i$ is invariant under the action ζ_1 of $SO(10) = \chi(\text{Spin}(10))$. The triplet $(\text{Spin}(10), \rho_1, V(10))$ has no relative invariant (See [1]) and hence we have $F(x) \equiv 0$. By using (4.1), it is clear that $f(x, y)$ is invariant under the action of Spin(10). We shall show that $f(x, y)$ is relatively invariant under $GL(2)$. Assume that $Q_i(x)$ (resp. $Q_{i+5}(x)$) has a term $x_i x_{i_2}$ (resp. $x_{i_3} x_{i_4}$) ($1 \leq i \leq 5$). Since $F(x) \equiv 0$, we may assume that $Q_j(x)$ (resp. $Q_{j+5}(x)$) has a term $x_{i_1} x_{i_3}$ (resp. $x_{i_2} x_{i_4}$) for some j satisfying $1 \leq j \leq 5$. This implies that $f(x, y) = \sum_{i=1}^5 B_i(x, y) B_{i+5}(x, y)$ is a linear combination of terms of the following form:

$$(4.2) \quad \begin{aligned} & (x_{i_1} y_{i_2} + y_{i_1} x_{i_2})(x_{i_3} y_{i_4} + y_{i_3} x_{i_4}) - (x_{i_1} y_{i_3} + y_{i_1} x_{i_3})(x_{i_2} y_{i_4} + y_{i_2} x_{i_4}) \\ & = \det \begin{pmatrix} x_{i_2} & y_{i_2} \\ x_{i_3} & y_{i_3} \end{pmatrix} \cdot \det \begin{pmatrix} x_{i_4} & y_{i_4} \\ x_{i_1} & y_{i_1} \end{pmatrix}. \end{aligned}$$

Hence it is clear that $f(x, y)$ is relatively invariant under $GL(2)$. Since $f(1 + e_1 e_2 e_3 e_4, e_1 e_5 + e_2 e_3 e_4 e_5) = 1$, it is not identically zero. Q.E.D.

Now we shall consider the micro-differential equation $\mathfrak{M} = \mathcal{E}f(x, y)^s$ and by constructing its holonomy diagram, we shall calculate the b -function of this space.

Since $G = \text{Spin}(10) \times GL(2)$ is reductive, we have $(G, \rho^*, V^*) \cong (G, \rho, V)$ and hence the dual space V^* has also nine G -orbits S_m^* ($m = 0, 1, 4, 8, 9, 13, 15, 20, 32$). We identify V and V^* by taking $(e_{i_1} \cdots e_{i_k}, e_{j_1} \cdots e_{j_\ell})$ ($k, \ell = 0, 2, 4$) as a dual basis, where $e_{i_1} \cdots e_{i_k} = 1$ for $k = 0$. We denote by A_m (resp. A_m^*) the conormal bundle of S_m (resp. S_m^*).

(1) The isotropy subalgebra \mathfrak{g}_{x_0} at $x_0 = (1 + e_1 e_2 e_3 e_4, e_1 e_5 + e_2 e_3 e_4 e_5)$ is isomorphic to $(\mathfrak{g}_2) \oplus \mathfrak{sl}(2)$ (See (5.40) and (5.42) in [1]). Since $A_0 = V \times \{0\} = A_{32}^*$, A_0 is a good holonomic variety and we have $\text{ord}_{A_0} f^s = 0$.

(2) The isotropy subalgebra \mathfrak{g}_{x_1} at $x_1 = (1 + e_1 e_2 e_3 e_4, e_1 e_2 + e_2 e_3 e_4 e_5)$ is isomorphic to $(\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(11)$ (See (5.43) in [1]). The conormal vector space $V_{x_1}^*$ is spanned by $(e_1 e_3 e_4 e_5, -e_1 e_5) = y_1 \in S_{13}^*$. Hence $A_1 =$

$\overline{G(x_1, y_1)} = A_{13}^*$ and $A_{13} = A_1^*$. Let A_0 be an element of \mathfrak{g}_{x_1} with $d_{11} = d_{22} = -1/4$, all remaining parts zero in (5.43) of [1]. Then we have $d\rho(A_0)x_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$. Since $\delta\chi(A_0) = 2(d_{11} + d_{22}) = -1$, $\text{tr}_{V_{x_1}^*} A_0 = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_1} f^s = -s - 1/2$. It is clear that A_0 and A_1 intersect regularly and G_0 -prehomogeneously with codimension one. Hence we have $b_{A_1}(s)/b_{A_0}(s) = (s + 1)$. Note that $G_0 = \{g \in G; \chi(g) = 1\}$.

(3) The isotropy subalgebra \mathfrak{g}_{x_4} at $x_4 = (1, e_1e_5 + e_2e_3e_4e_5)$ is, by simple calculation using (5.38) in [1], given as follows:

$$(4.3) \quad \mathfrak{g}_{x_4} = \left\{ \tilde{A} = \left(\begin{array}{c|c} A & 0 \\ \hline C & -{}^tA \end{array} \right) \oplus \left(\begin{array}{cc} 3\varepsilon - \eta & 0 \\ c & \eta \end{array} \right); A = \left(\begin{array}{c|c|c} 3\varepsilon & 0 & 0 \\ * & \varepsilon I_3 + X & 0 \\ * & * & -2n \end{array} \right), \right. \\ \left. X \in \mathfrak{sl}(3), {}^tC = -C \quad \text{with} \quad c_{i5} = 0, i = 1, \dots, 4 \right\}.$$

Put $\omega_1 = (e_1e_3e_4e_5, 0)$, $\omega_2 = (-e_1e_2e_4e_5, 0)$, $\omega_3 = (e_1e_2e_3e_5, 0)$, and $\omega_4 = (e_1e_2e_3e_4, 0)$. Then the conormal vector space $V_{x_4}^*$ is spanned by $\omega_1, \dots, \omega_4$. The action $d\rho_{x_4}$ of \mathfrak{g}_{x_4} on $V_{x_4}^*$ is given as follows:

$$(4.4) \quad d\rho_{x_4}(\tilde{A})(\omega_1, \dots, \omega_4) = (\omega_1, \dots, \omega_4) \left(\begin{array}{c|c|c} (2\eta - 5\varepsilon)I_3 + X & 0 \\ * & * & * \\ * & * & -6\varepsilon \end{array} \right).$$

Since ω_1 is a generic point, we have $A_4 = A_{20}^*$ and $A_{20} = A_4^*$. Let A_0 be an element of \mathfrak{g}_{x_4} with $2\eta - 5\varepsilon = 1$, all remaining parts zero except ε and η in (4.3). Then $d\rho(A_0)x_4 = 0$ and $d\rho^*(A_0)\omega_1 = \omega_1$. However we have $\delta\chi(A_0) = 6\varepsilon$ which is not definite. If A_4 is a good holonomic variety, this must be definite by Proposition 1-3, and hence A_4 is not a good holonomic variety, i.e., $A_4 \not\subset W$. Note that the P.V. $(G_{x_4}, \rho_{x_4}, V_{x_4}^*)$ has no relative invariant.

(4) The isotropy subalgebra \mathfrak{g}_{x_8} at $x_8 = (1, e_1e_2e_3e_4)$ is given as follows:

$$(4.5) \quad \mathfrak{g}_{x_8} = \left\{ \tilde{X} = \left[\begin{array}{c|c|c|c} \varepsilon I_4 + X & \gamma & 0 & 0 \\ \hline 0 & 2\eta & 0 & 0 \\ \hline 0 & \delta & -\varepsilon I_4 - {}^tX & 0 \\ \hline -{}^t\delta & 0 & -{}^t\gamma & -2\eta \end{array} \right] \oplus \left(\begin{array}{cc} \eta + 2\varepsilon & 0 \\ 0 & \eta - 2\varepsilon \end{array} \right); \right. \\ \left. X \in \mathfrak{sl}(4), \gamma, \delta \in \mathbf{C}^4 \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(4) \oplus \mathfrak{u}(8)).$$

Put $\omega_1 = (e_2e_3e_4e_5, 0)$, $\omega_2 = -(e_1e_3e_4e_5, 0)$, $\omega_3 = (e_1e_2e_4e_5, 0)$, $\omega_4 = -(e_1e_2e_3e_5, 0)$, $\omega_5 = (0, e_1e_5)$, $\omega_6 = (0, e_2e_5)$, $\omega_7 = (0, e_3e_5)$, $\omega_8 = (0, e_4e_5)$. Then the conormal vector space $V_{x_8}^*$ is spanned by $\omega_1, \dots, \omega_8$, and the action $d\rho_{x_8}$ of \mathfrak{g}_{x_8} on $V_{x_8}^*$ is given as follows.

$$(4.6) \quad d\rho_{x_8}(\tilde{X})(\omega_1, \dots, \omega_8) = (\omega_1, \dots, \omega_8) \left(\frac{(3\varepsilon - 2\eta)I_4 + X}{0} \middle| \frac{0}{-(3\varepsilon + 2\eta)I_4 - {}^tX} \right)$$

where $\tilde{X} \in \mathfrak{g}_{x_8}$ in (4.5).

Any relative invariant of $(G_{x_8}, \rho_{x_8}, V_{x_8}^*)$ is of the form $c \cdot g(x)^m$ ($c \in \mathbf{C}$, $m \in \mathbf{Z}$) where $g(x) = \sum_{i=1}^4 x_i x_{i+4}$ for $x = \sum_{i=1}^8 x_i \omega_i$. Clearly $y_8 = \omega_1 + \omega_5 = (e_2e_3e_4e_5, e_1e_5)$ is a generic point, and $y'_8 = \omega_1 + \omega_6 = (e_2e_3e_4e_5, e_2e_5)$ is a point of the one-codimensional orbit. Hence we have $A_3 = A_8^*$ and $\dim A_1 \cap A_8 = \dim V - 1$. Since $A_{13} = A_1^*$, we have also $\dim A_8 \cap A_{13} = \dim V - 1$. Note that $(G_{x_8}, \rho_{x_8}, V_{x_8}^*)$ is a regular P.V. since $\rho_{x_8}(G_{x_8})$ and its generic isotropy subgroup are reductive (See [1]). By Corollary 1-7, A_8 is a good holonomic variety. Let \tilde{X}_0 be an element of \mathfrak{g}_{x_8} with $\eta = -\frac{1}{2}$, all remaining parts zero in (4.5). Then $d\rho(\tilde{X}_0)x_8 = 0$ and $d\rho^*(\tilde{X}_0)y_8 = y_8$. Since $\delta\chi(\tilde{X}_0) = 4\eta = -2$, $\text{tr}_{V_{x_8}^*} \tilde{X}_0 = -16\eta = 8$ and $\dim V_{x_8}^* = 8$, we have $\text{ord}_{A_8} f^s = -2s - \frac{8}{2}$. Since $m_{A_8} - m_{A_1} = 1$, they intersect regularly. By Corollary 1-2, we have $b_{A_8}(s)/b_{A_1}(s) = (s + 4)$.

(5) We shall calculate the isotropy subalgebra at $x'_9 = (1, e_1e_3 + e_2e_4)$ instead of $x_9 = (1, e_1e_2 + e_3e_4)$. It is given as follows.

$$(4.7) \quad \mathfrak{g}_{x'_9} = \left\{ \tilde{A} = \left[\begin{array}{c|c|c} \varepsilon I_4 + A & B & 0 \\ \hline 0 & 2\eta & \\ \hline C & & \frac{-\varepsilon I_4 - {}^tA}{-{}^tB} \quad \frac{0}{-2\eta} \end{array} \right] + \left(\begin{array}{c|c} 2\varepsilon + \eta & 0 \\ \hline c_{13} + c_{24} & \eta \end{array} \right); \right. \\ \left. A \in \mathfrak{sp}(2), B \in \mathbf{C}^4, C = -{}^tC = (c_{ij}) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sp}(2)) \oplus \mathfrak{u}(4).$$

Put $\omega_1 = (e_2e_3e_4e_5, 0)$, $\omega_2 = -(e_1e_3e_4e_5, 0)$, $\omega_3 = (e_1e_2e_4e_5, 0)$, $\omega_4 = -(e_1e_2e_3e_5, 0)$, $\omega_5 = (e_1e_2e_3e_4, 0)$, $\omega_6 = -(e_3e_5, e_2e_3e_4e_5)$, $\omega_7 = (-e_4e_5, e_1e_3e_4e_5)$, $\omega_8 = (e_1e_5, -e_1e_2e_4e_5)$, $\omega_9 = (e_2e_5, e_1e_2e_3e_5)$. Then the conormal vector space $V_{x'_9}^*$ is spanned by these $\omega_1, \dots, \omega_9$ and the action $d\rho_{x'_9}$ of $\mathfrak{g}_{x'_9}$ on $V_{x'_9}^*$ is given as follows:

$$(4.8) \quad d\rho_{x_9}(\tilde{A})(\omega_1, \dots, \omega_9) = (\omega_1, \dots, \omega_9) \left(\begin{array}{c|c|c} A - (3\varepsilon + 2\eta)I_4 & B & C' \\ \hline 0 & -4\varepsilon & 0 \\ \hline 0 & 0 & A - (\varepsilon + 2\eta)I_4 \end{array} \right)$$

with $C' = \left(\begin{array}{c|c|c|c} c_{13} + 2c_{24} & -c_{23} & 0 & c_{34} \\ \hline -c_{14} & 2c_{13} + c_{24} & -c_{34} & 0 \\ \hline 0 & -c_{12} & c_{13} + 2c_{24} & -c_{14} \\ \hline c_{12} & 0 & -c_{23} & 2c_{13} + c_{24} \end{array} \right).$

Clearly, $y_9 = \omega_5 + \omega_8$ is its generic point and $y'_9 = \omega_1 + \omega_8$ is a point of the one-codimensional orbit. Note that $(G_{x_9'}, \rho_{x_9'}, V_{x_9}^*)$ has only one orbit of codimension one. Since $y_9, y'_9 \in S_9^*$, we have $A_9 = A_9^*$, and A_9 has no one-codimensional intersection with other conormal bundles. Let \tilde{A}_0 be an element of $\mathfrak{g}_{x_9'}$ with $\varepsilon = -\frac{1}{4}$, $\eta = -\frac{3}{8}$, all remaining parts zero in (4.7). Then $d\rho(\tilde{A}_0)x'_9 = 0$ and $d\rho^*(\tilde{A}_0)y_9 = y_9$. We have $\delta\chi(\tilde{A}_0) = 2\{(2\varepsilon + \eta) + \eta\} = -\frac{5}{2}$, we have $m_{A_9} = \frac{5}{2}$. This implies that the conormal bundle A_9 is not a good holonomic variety, i.e., $A_9 \not\subset W$ since otherwise m_{A_9} must be a non-negative integer (See §1 or [1]).

(6) The isotropy subalgebra $\mathfrak{g}_{x_{13}}$ at $x_{13} = (1, e_1e_2)$ is given as follows.

$$(4.9) \quad \mathfrak{g}_{x_{13}} = \left\{ \tilde{X} = \left(\begin{array}{c|c|c|c} \varepsilon_1 I_2 + X & Z & 0 & -b \\ \hline 0 & 2\varepsilon I_3 + Y & 0 & 0 \\ \hline C & -\varepsilon_1 I_2 - {}^t X & 0 & 0 \\ \hline & -{}^t Z & -2\varepsilon I_3 - {}^t Y & \end{array} \right) \right.$$

$$\left. \oplus \left(\begin{array}{c|c} 3\varepsilon + \varepsilon_1 & b \\ \hline c_{12} & 3\varepsilon - \varepsilon_1 \end{array} \right); X \in \mathfrak{sl}(2), Y \in \mathfrak{sl}(3), {}^t C = -C \right\}$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(3) \oplus \mathfrak{u}(15)).$$

Since $A_{13} = A_1^*$, $A_3 = A_3^*$, and $\dim A_1^* \cap A_3^* = \dim V - 1$, the conormal bundle A_{13} is a good holonomic variety and $\dim A_3 \cap A_{13} = \dim V - 1$. They intersect regularly. Put $\omega_1 = (0, e_2e_3e_4e_5)$, $\omega_2 = (e_2e_3e_4e_5, 0)$, $\omega_3 = (0, e_1e_3e_4e_5)$, $\omega_4 = (e_1e_3e_4e_5, 0)$, $\omega_5 = (e_1e_5, -e_1e_2e_4e_5)$, $\omega_6 = (e_3e_5, -e_1e_2e_3e_5)$, $\omega_7 = (e_3e_4, -e_1e_2e_3e_4)$, $\omega_8 = (0, e_4e_5)$, $\omega_9 = (0, e_3e_5)$, $\omega_{10} = (0, e_3e_4)$, $\omega_{11} = (e_1e_2e_4e_5, 0)$, $\omega_{12} = (e_1e_2e_3e_5, 0)$, $\omega_{13} = (e_1e_2e_3e_4, 0)$. Then the conormal vector space $V_{x_{13}}^*$ is spanned by these $\omega_1, \dots, \omega_{13}$ and the action $d\rho_{x_{13}}$ of $\mathfrak{g}_{x_{13}}$ on $V_{x_{13}}^*$ is given as follows:

$$(4.10) \quad \begin{aligned} & d\rho_{x_{13}}(\tilde{X})(\omega_1, \dots, \omega_{13}) \\ &= (\omega_1, \dots, \omega_{13}) \left(\frac{-6\varepsilon I_4 + d\rho_1(X \oplus W)}{0} \middle| \frac{*}{-4\varepsilon I_9 + d\rho_2(W \oplus Y)} \right) \end{aligned}$$

where $\rho_1 = A_1 \otimes A_1$ for $SL(2) \times SL(2)$, $\rho_2 = (2A_1) \otimes A_1$ for $SL(2) \times SL(3)$ and $W = \begin{pmatrix} \varepsilon_1 & b \\ c_{12} & -\varepsilon_1 \end{pmatrix} \in \mathfrak{sl}(2)$.

As a generic point, we may take $y_{13} = \omega_5 + \omega_9 + \omega_{13} = (e_4e_5 + e_1e_2e_3e_4, e_3e_5 - e_1e_2e_4e_5)$. Let \tilde{X}_0 be an element of $\mathfrak{g}_{x_{13}}$ with $\varepsilon = -\frac{1}{4}$, all remaining parts zero. Then $d\rho(\tilde{X}_0)x_{13} = 0$ and $d\rho^*(\tilde{X}_0)y_{13} = y_{13}$. Since $\delta\chi(\tilde{X}_0) = 12\varepsilon = -3$, $\text{tr}_{V_{x_{13}}^*} \tilde{X}_0 = -60\varepsilon = 15$ and $\dim V_{x_{13}}^* = 13$, we have $\text{ord}_{A_{13}} f^s = -3s - \frac{17}{2}$. By Corollary 1-2, we have $b_{A_{13}}(s)/b_{A_6}(s) = (s+5)$. By (4.10), we can see that $(G_{x_{13}}, \rho_{x_{13}}, V_{x_{13}}^*)$ has the unique relative invariant (See Lemma 4 and Proposition 5 in § 4 in [1]), i.e., it has the unique one-codimensional orbit.

(7) The isotropy subalgebra $\mathfrak{g}_{x_{15}}$ at $x_{15} = (1 + e_1e_2e_3e_4, 0)$ is given as follows.

$$(4.11) \quad \mathfrak{g}_{x_{15}} = \left\{ \tilde{X} = \left(\begin{array}{c|c|c|c} X & Y & C'' & 0 \\ \hline 0 & 2\varepsilon & 0 & 0 \\ \hline C & C' & -{}^tX & 0 \\ \hline -{}^tC' & 0 & -{}^tY & -2\varepsilon \end{array} \right) \oplus \begin{pmatrix} \varepsilon & b \\ 0 & \eta \end{pmatrix}; \right. \\ \left. \begin{array}{l} X \in \mathfrak{sl}(4), CC'' = -\text{Pf } C \cdot I_4, {}^tC = -C \in M(4) \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{o}(7) \oplus \mathfrak{u}(9)). \end{array} \right\}$$

Note that, in (4.11), $\tilde{X}_0 = \left(\begin{array}{c|c} X & C'' \\ \hline C & -{}^tX \end{array} \right)$ is the spin representation of X_0 in $\mathfrak{o}(7)$. Put $\omega_1 = (0, e_1e_5)$, $\omega_2 = (0, e_2e_5)$, $\omega_3 = (0, e_3e_5)$, $\omega_4 = (0, e_4e_5)$, $\omega_5 = (0, e_2e_3e_4e_5)$, $\omega_6 = (0, -e_1e_3e_4e_5)$, $\omega_7 = (0, e_1e_2e_4e_5)$, $\omega_8 = (0, -e_1e_2e_3e_5)$, $\omega_9 = (0, \frac{1}{2}(1 - e_1e_2e_3e_4))$, $\omega_{10} = (0, e_2e_3)$, $\omega_{11} = (0, -e_2e_4)$, $\omega_{12} = (0, e_3e_4)$, $\omega_{13} = (0, e_1e_4)$, $\omega_{14} = (0, e_1e_3)$, $\omega_{15} = (0, e_1e_2)$. The conormal vector space $V_{x_{15}}^*$ is spanned by $\omega_1, \dots, \omega_{15}$. Then $y_{15} = \omega_9$ is its generic point and $y'_{15} = \omega_{10} + \omega_{14}$ is a point of the unique one-codimensional orbit. Since $y_{15}, y'_{15} \in S_{15}^*$, we have $A_{15} = A_{15}^*$, and A_{15} has no one-codimensional intersection with any other conormal bundle. Let \tilde{X}_1 be an element of $\mathfrak{g}_{x_{15}}$ with $\varepsilon = \beta + 1$, $\eta = \beta$, all remaining parts zero in (4.11). Then $d\rho(\tilde{X}_1)x_{15} = 0$ and $d\rho^*(\tilde{X}_1)y_{15} = y_{15}$. Since $\delta\chi(\tilde{X}_1) = 2(\varepsilon + \eta) = 2(2\beta + 1)$ is not definite, the conormal bundle A_{15} is not a good holonomic variety, i.e., $A_{15} \not\subset W$.

(8) Since $A_{20} = A_4^*$ and $A_4 \not\subset W$, the conormal bundle A_{20} is not a good holonomic variety. Note that $W \subset V \times V^*$ is symmetric with respect to V and V^* .

(9) Since $A_{32} = \{0\} \times V^*$, the conormal bundle A_{32} is a good holonomic variety. Put $A_0 = (0) \oplus (-I_2)$. Then $d\rho(A_0)x_{32} = 0$ and $d\rho^*(A_0)y_{32} = y_{32}$ where y_{32} is a generic point of (G, ρ^*, V^*) . Since $\delta\chi(A_0) = -4$, $\text{tr}_{V_{x_{32}}}^* A_0 = 32$ and $\dim V_{x_{32}}^* = 32$, we have $\text{ord}_{A_{32}} f^s = -4s - \frac{32}{2}$ and hence by Corollary 1-2, we have $b_{A_{32}}(s)/b_{A_{13}}(s) = s + 8$. Note that $A_{32} = A_0^*$ and $A_{13} = A_1^*$. Since $b_{A_0}(s) = 1$ and $b_{A_{13}}(s) = b(s)$, we have the b -function $b(s) = (s + 1)(s + 4)(s + 5)(s + 8)$, and the holonomy diagram (Figure 4-1). We denote A_m by \textcircled{m} .




Figure 4-1. Holonomy diagram of $(\text{Spin}(10) \times GL(2), \text{half-spin rep.} \otimes A_1, V(16) \otimes V(2))$

§ 5. $(GL(1) \times \text{Spin}(12), \square \otimes \text{half-spin rep.}, V(1) \otimes V(32))$

The representation space $V = V(1) \otimes V(32)$ is spanned by $1, e_i e_j, e_r e_s e_t e_u, e_i e_2 e_3 e_4 e_5 e_6$ ($1 \leq i < j \leq 6, 1 \leq r < s < t < u \leq 6$) (See [1], [4]). J-I. Igusa has completed the orbital decomposition of this space (See [3]). There exist five G -orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 7, 16, 32$) where S_m denotes the m -codimensional orbit and $x_0 = 1 + e_1 e_2 e_3 e_4 e_5 e_6, x_1 = 1 + e_2 e_3 e_4 e_5 e_6 + e_1 e_3 e_4 e_5 e_6, x_7 = 1 + e_2 e_3 e_4 e_5, x_{16} = 1, x_{32} = 0$. We identify V^* with V by taking $\{1, e_i e_j, e_r e_s e_t e_u, e_1 e_2 e_3 e_4 e_5 e_6\}$ as a dual basis. Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, there exist also five orbits $S_m^* (m = 0, 1, 7, 16, 32)$ in V^* . We denote by A_m (resp. A_m^*) the conormal bundle of S_m (resp. S_m^*). Clearly, we have $A_0 = V \times \{0\} = A_{32}^*$ and $A_{32} = \{0\} \times V^* = A_0^*$. The Lie algebra \mathfrak{g} of $GL(1) \times \text{Spin}(12)$ is given as follows:

$$(5.1) \quad \mathfrak{g} = \left\{ (d) \oplus \left(\begin{array}{c|c} A & B \\ \hline C & -{}^t A \end{array} \right); A, B, C \in M(6), {}^t B = -B, {}^t C = -C \right\}.$$

(1) The isotropy subalgebra \mathfrak{g}_{x_0} at x_0 is given as follows (See [1]).

$$(5.2) \quad \mathfrak{g}_{x_0} = \left\{ (0) \oplus \left(\begin{array}{c|c} A & 0 \\ \hline 0 & -{}^t A \end{array} \right); A \in \mathfrak{sl}(6) \right\} \cong \mathfrak{sl}(6).$$

Since $A_0 = V \times \{0\}$, we have $\text{ord}_{A_0} f^s = 0$, where f denotes the relative invariant of degree four (See [1], [3]).

(2) By using (5.29) in [1], we can calculate the isotropy subalgebra \mathfrak{g}_{x_1} .

$$(5.3) \quad \begin{aligned} \mathfrak{g}_{x_1} &= \left\{ \tilde{A} = (d) \oplus \left(\begin{array}{c|c} A & B \\ \hline C & -{}^t A \end{array} \right); a_1 + a_4 = a_2 + a_5 = -a_3 - a_6 = 2d, c_{36} = 0 \right\} \\ &\cong \left\{ (d) \oplus \left(\begin{array}{c|c} -dI_6 + A_0 & B_0 \\ \hline 0 & -dI_6 - {}^t A_0 \end{array} \right); A_0 \in \mathfrak{sp}(3), {}^t B_0 = -B_0, \text{tr } B_0 J = 0 \right\} \\ &\quad \text{with } J = \left(\begin{array}{c|c} 0 & I_3 \\ \hline -I_3 & 0 \end{array} \right), \\ &\cong (\mathfrak{gl}(1) \oplus \mathfrak{sp}(3)) \oplus V(14) \quad \text{where} \end{aligned}$$

$$A = \begin{pmatrix} a_1 & a_{12} & 0 & a_{14} & a_{15} & 0 \\ a_{21} & a_2 & 0 & a_{15} & a_{25} & 0 \\ a_{31} & a_{32} & a_3 & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & 0 & a_4 & -a_{21} & 0 \\ a_{42} & a_{52} & 0 & -a_{12} & a_5 & 0 \\ a_{61} & a_{62} & a_{63} & a_{64} & a_{65} & a_6 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & c_{46} & 0 & 0 & c_{34} \\ 0 & 0 & c_{56} & 0 & 0 & c_{35} \\ -c_{46} & -c_{56} & 0 & c_{16} & c_{26} & b_{36} \\ 0 & 0 & -c_{16} & 0 & 0 & c_{13} \\ 0 & 0 & -c_{26} & 0 & 0 & c_{23} \\ -c_{34} & -c_{35} & -b_{36} & -c_{13} & -c_{23} & 0 \end{pmatrix}$$

with $b_{36} + c_{14} + c_{25} = 0$.

The conormal vector space $V_{x_1}^*$ is spanned by $e_1 e_2 e_4 e_5$ on which \mathfrak{g}_{x_1} acts as $d\rho_{x_1}(\tilde{A})e_1 e_2 e_4 e_5 = -4de_1 e_2 e_4 e_5$ for $\tilde{A} \in \mathfrak{g}_{x_1}$. This implies that $A_1 = \overline{G(x_1, e_1 e_2 e_4 e_5)} = A_{16}^*$. Since 0 is the point of the one-codimensional orbit, we have $\dim A_0 \cap A_1 = \dim V - 1$ and $A_0 \cap A_1$ is G_0 -prehomogeneous, i.e., A_1 is a good holonomic variety by Proposition 1-5. Let A_0 be an element of \mathfrak{g}_{x_1} with $-4d = 1$. Then $d\rho(A_0)x_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$ where $y_1 = e_1 e_2 e_4 e_5$. Since $\delta\chi(A_0) = 4d = -1$, $\text{tr}_{V_{x_1}^*} A_0 = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_1} f^s = -s - \frac{1}{2}$. By Proposition 1-4, A_0 and A_1 intersect regularly and hence $b_{A_1}(s)/b_{A_0}(s) = (s+1)$ by Corollary 1-2.

(3) By using (5.29) in [1], we can calculate the isotropy subalgebra \mathfrak{g}_{x_7} .

$$\mathfrak{g}_{x_7} = \left\{ \tilde{A} = (d) \oplus \left(\begin{array}{c|c} A & C \\ \hline C & -{}^t A \end{array} \right); d = \frac{a_1 + a_4}{2}, a_2 + a_3 + a_5 + a_6 = 0, \right.$$

$$\begin{aligned}
 (5.4) \quad & \left. {}^t C = C, A \text{ and } C' \text{ are given as follows} \right\} \\
 & \cong \left\{ (d) \oplus \left(\begin{array}{c|c|c} dI_2 + A_1 & 0 & 0 \\ \hline U & d\rho_1(V) & 0 \\ \hline W & -{}^t US & -dI_2 - {}^t A_1 \end{array} \right); A_1 \in \mathfrak{sl}(2), W \in \mathfrak{o}(2), \right. \\
 & \left. V \in \mathfrak{o}(7), U \in M(8,2) \right\} \\
 & \cong (\mathfrak{gl}(1) \oplus \mathfrak{o}(7) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(17)
 \end{aligned}$$

where ρ_1 is the spin-representation of $\text{Spin}(7)$, $S = \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}$, and

$$A = \begin{pmatrix} a_1 & 0 & 0 & a_{14} & 0 & 0 \\ a_{21} & a_2 & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_3 & a_{34} & a_{35} & a_{36} \\ a_{41} & 0 & 0 & a_4 & 0 & 0 \\ a_{51} & a_{52} & a_{53} & a_{54} & a_5 & a_{56} \\ a_{61} & a_{62} & a_{63} & a_{64} & a_{65} & a_6 \end{pmatrix}, \quad C' = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & c_{56} & 0 & -c_{36} & c_{35} \\ 0 & -c_{56} & 0 & 0 & c_{26} & -c_{25} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_{36} & -c_{26} & 0 & 0 & c_{23} \\ 0 & -c_{35} & c_{25} & 0 & -c_{23} & 0 \end{pmatrix}.$$

Put $\omega_1 = e_1 e_4 - e_1 e_2 e_3 e_4 e_5 e_6$, $\omega_2 = e_1 e_2 e_4 e_5$, $\omega_3 = e_1 e_2 e_3 e_4$, $\omega_4 = e_1 e_2 e_4 e_6$, $\omega_5 = e_1 e_3 e_4 e_5$, $\omega_6 = e_1 e_3 e_4 e_6$, $\omega_7 = e_1 e_4 e_5 e_6$. The conormal vector space $V_{x_7}^*$ is spanned by these $\omega_1, \dots, \omega_7$, and $(G_{x_7}, \rho_{x_7}, V_{x_7}^*) \cong (GL(1) \times SO(7), \square \otimes A_1, V(1) \otimes V(7))$. Then ω_1 is its generic point and $\omega_2 = e_1 e_2 e_4 e_5$ is a point of the one-codimensional orbit. Since $A_1 = \overline{G(x_1, \omega_2)}$, we have $\text{codim } A_1 \cap A_7 = 1$. Since $A_1 \cap A_7$ is G_0 -prehomogeneous, A_7 is a good holonomic variety. We have $A_7 = A_7^*$.

Let A'_β be an element of \mathfrak{g}_{x_7} with $4d = 2(a_1 + a_3) = -\beta - 1$, $2(a_3 + a_6) = -2(a_2 + a_5) = 1 - \beta$, all remaining parts zero in (5.4). Then we have $d\rho(A'_\beta)x_7 = 0$, $d\rho^*(A'_\beta)\omega_2 = \omega_2$ and $\text{tr}_V A'_\beta = \beta$ where $\tilde{V} = V_{x_7}^* \text{ mod } d\rho_{x_7}(\mathfrak{g}_{x_7})\omega_2$. This implies that A_1 and A_7 intersect regularly by Proposition 1-4. Let A_0 be an element of \mathfrak{g}_{x_7} with $d = -\frac{1}{2}$, all remaining parts zero in (5.4). Then $d\rho(A_0)x_7 = 0$, $d\rho^*(A_0)\omega_1 = \omega_1$. Since $\delta\chi(A_0) = 4d = -2$, $\text{tr}_{V_{x_7}^*} A_0 = -14d = 7$, $\dim V_{x_7}^* = 7$, we have $\text{ord}_{A_7} f^s = -2s - \frac{7}{2}$. By Corollary 1-2, we have $b_{A_7}(s)/b_{A_1}(s) = (s + \frac{7}{2})$.

(4) Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, $A_7^* = A_7$ and $A_1^* = A_{16}$, we have $\dim A_7 \cap A_{16} = \dim V - 1$ and they intersect regularly. Since $A_7 \cap A_{16} = A_7^* \cap A_1^*$ is G_0 -prehomogeneous, A_{16} is a good holonomic variety. Since $d\rho(\tilde{A}) \cdot 1 = d - (\text{tr } A/2) + \sum_{i < j} b_{ij} e_i e_j$, the isotropy subalgebra $\mathfrak{g}_{x_{16}}$ at $x_{16} = 1$ is

$$\begin{aligned}
 (5.5) \quad \mathfrak{g}_{x_{16}} &= \left\{ \tilde{A} = \begin{pmatrix} \text{tr } A \\ 2 \end{pmatrix} \oplus \begin{pmatrix} A & 0 \\ C & -{}^t A \end{pmatrix}; {}^t C = -C, A, C \in M(3) \right\} \\
 &\cong \mathfrak{gl}(6) \oplus V(15).
 \end{aligned}$$

The conormal vector space $V_{x_{16}}^*$ is spanned by $e_1 e_2 e_3 e_4 e_5 e_6$ and $e_i e_j e_k e_\ell$ ($1 \leq i < j < k < \ell \leq 6$). Then the action $d\rho_{x_{16}}$ of $\mathfrak{g}_{x_{16}}$ on $V_{x_{16}}^*$ is given by

$$(5.6) \quad d\rho_{x_{16}}(\tilde{A})(\omega_1, \dots, \omega_{16}) = (\omega_1, \dots, \omega_{16}) \begin{pmatrix} -\text{tr } A & c' \\ 0 & d\rho_1^*(A) \end{pmatrix}$$

where $\omega_1 = e_1 e_2 e_3 e_4 e_5 e_6$, $\{\omega_2, \dots, \omega_{16}\} = \{e_i e_j e_k e_\ell, 1 \leq i < j < k < \ell \leq 6\}$, $c' \in \mathbf{C}^{15}$, $\rho_1 = A_2$ for $GL(6)$.

Then $y_{16} = e_1 e_2 e_4 e_5 + e_1 e_3 e_4 e_6 + e_2 e_3 e_5 e_6$ is its generic point. Let A_0 be an element of $\mathfrak{g}_{x_{16}}$ with $A = -\frac{1}{4}I_6$, $C = 0$ in (5.5). Then $d\rho(A_0)x_{16} = 0$ and $d\rho^*(A_0)y_{16} = y_{16}$. Since $\delta\chi(A_0) = 4d = 2\text{tr}(-\frac{1}{4}I_6) = -3$, $\text{tr}_{V_{x_{16}}^*} A_0 = -11\text{tr}(-\frac{1}{4}I_6) = \frac{33}{2}$ and $\dim V_{x_{16}}^* = 16$, we have $\text{ord}_{A_{16}} f^s = s\delta\chi(A_0) - \text{tr}_{V_{x_{16}}^*} A_0 + \frac{1}{2}\dim V_{x_{16}}^* = -3s - \frac{17}{2}$. By Corollary 1-2, we have $b_{A_{16}}(s)/b_{A_7}(s) = (s + \frac{11}{2})$. By (5.6), the character group of $\rho_{x_{16}}(G_{x_{16}})$ is one-dimensional and hence $(G_{x_{16}}, \rho_{x_{16}}, V_{x_{16}}^*)$ has (at most) the unique one-codimensional orbit.

(5) Since $A_{32} = A_0^*$ and $A_{16} = A_1^*$, they intersect regularly with codimension one. We shall calculate the order $\text{ord}_{A_{32}} f^s$. Since $(G_{x_{32}}, \rho_{x_{32}}, V_{x_{32}}) \cong (G, \rho^*, V^*)$, $y_{32} = 1 + e_1 e_2 e_3 e_4 e_5 e_6$ is its generic point. Let A_0 be an element of \mathfrak{g} with $d = -1$, all remaining parts zero in (5.1). Then $d\rho(A_0)x_{32} = 0$, $d\rho^*(A_0)y_{32} = y_{32}$. Since $\delta\chi(A_0) = -4$, $\text{tr}_{V_{x_{32}}^*} A_0 = -32d = 32$, $\dim V_{x_{32}}^* = 32$, we have $\text{ord}_{A_{32}} f^s = -4s - \frac{32}{2}$. By Corollary 1-2, we have $b_{A_{32}}(s)/b_{A_{16}}(s) = s + 8$. Since $b_{A_0}(s) = 1$ and $b_{A_{32}}(s) = b(s)$, we obtain the b -function $b(s) = (s + 1)(s + \frac{7}{2})(s + \frac{11}{2})(s + 8)$ and the holonomy diagram (Figure 5-1).

We denote $\textcircled{A_m}$ by \textcircled{m} .




Figure 5-1. Holonomy diagram of $(GL(1) \times \text{Spin}(12))$,
 $\square \otimes$ half-spin rep., $V(1) \otimes V(32)$.

- Remark.* (1) $(GL(1) \times Spin(7), \square \otimes \text{spin rep.}, V(1) \otimes V(8))$
 (2) $(Spin(7) \times GL(2), \text{spin rep.} \otimes A_1, V(8) \otimes V(2))$
 (3) $(Spin(7) \times GL(3), \text{spin rep.} \otimes A_1, V(8) \otimes V(3))$
 (4) $(GL(1) \times Spin(9), \square \otimes \text{spin rep.}, V(1) \otimes V(16))$
 (5) $(GL(1) \times (G_2), \square \otimes A_2, V(1) \otimes V(7))$
 (6) $((G_2) \times GL(2), A_2 \otimes A_1, V(7) \otimes V(2))$
 (7) $(GL(1) \times Spin(11), \square \otimes \text{spin rep.}, V(1) \otimes V(32))$

Since $Spin(7) \longrightarrow SO(8)$ by the spin representation, the first three triplets (1), (2), (3) are reduced to the triplet $(SO(8) \times GL(n), A_1 \otimes A_1, V(8) \otimes V(n))$ ($n = 1, 2, 3$) (See [1]). Since $Spin(9) \longrightarrow SO(16)$ by the spin representation, (4) is reduced to $(SO(16) \times GL(1), A_1 \otimes A_1, V(16) \otimes V(1))$ (See [1]). Since $(G_2) \longrightarrow SO(7)$ by A_2 , (5) and (6) are reduced to $(SO(7) \times GL(n), A_1 \otimes A_1, V(7) \otimes V(n))$ ($n = 1, 2$) (See [1]).

Since the spin representation of $Spin(11)$ is obtained by the restriction of the half-spin representation of $Spin(12)$ to $Spin(11)$, (7) is reduced to $Spin(12)$ in § 5. Note that the b -function depends essentially on the relative invariant itself, not on the group.

§ 6. $(GL(1) \times E_6, \square \otimes A_1, V(1) \otimes V(27))$

The Lie algebra \mathfrak{g} of $G = GL(1) \times E_6$ can be written as $\mathfrak{g} = \mathcal{D}_0 \oplus \mathcal{T}_1 \oplus \mathcal{T}_2 \oplus \mathcal{T}_3 \oplus \{R_\nu\}_{\nu \in \mathcal{J}}$ (See Proposition 37 and Example 39 of § 1 in [1]). The representation space is identified with the simple Jordan algebra \mathcal{J} .

$$(6.1) \quad \mathcal{J} = \left\{ X = \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}; \xi_1, \xi_2, \xi_3 \in \mathbf{C}; x_1, x_2, x_3 \in \mathcal{L} \right\}$$

where \mathcal{L} denotes the complex Cayley algebra.

DEFINITION 6-1. For $a \in \mathcal{L}$, we define elements $T_i(a)$ and $T'_i(a)$ ($i = 1, 2, 3$) of \mathfrak{g} as follows:

$$\begin{aligned} T_1(a) \cdot X &= [R_{A_1(a)} + \mathcal{T}_1(2a)]X = \begin{pmatrix} 0 & 0 & x_3 a \\ 0 & 0 & a \xi_2 \\ \bar{x}_3 a & \bar{a} \xi_2 & \text{tr}(\bar{x}_1 a) \end{pmatrix} \\ T'_1(a) \cdot X &= [R_{A_1(a)} - \mathcal{T}_1(2a)]X = \begin{pmatrix} 0 & \bar{a} \bar{x}_2 & 0 \\ a x_2 & \text{tr}(a \bar{x}_1) & a \xi_3 \\ 0 & \bar{a} \xi_3 & 0 \end{pmatrix} \\ T_2(a) \cdot X &= [R_{A_2(\bar{a})} + \mathcal{T}_2(2\bar{a})]X = \begin{pmatrix} 0 & 0 & a \xi_1 \\ 0 & 0 & \bar{x}_3 a \\ \bar{a} \xi_1 & \bar{a} x_3 & \text{tr}(x_2 a) \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
T'_2(a) \cdot X &= [R_{A_2(\bar{a})} - \mathcal{F}_2(2\bar{a})]X = \begin{pmatrix} \text{tr}(ax_2) & a\bar{x}_1 & a\xi_3 \\ x_1\bar{a} & 0 & 0 \\ \overline{a\xi_3} & 0 & 0 \end{pmatrix} \\
T_3(a) \cdot X &= [R_{A_3(a)} + \mathcal{F}_3(2a)]X = \begin{pmatrix} 0 & a\xi_1 & 0 \\ \overline{a\xi_1} & \text{tr}(\bar{a}x_3) & \overline{x_2a} \\ 0 & x_2a & 0 \end{pmatrix} \\
T'_3(a) \cdot X &= [R_{A_3(a)} - \mathcal{F}_3(2a)]X = \begin{pmatrix} \text{tr}(a\bar{x}_3) & a\xi_2 & ax_1 \\ \overline{a\xi_2} & 0 & 0 \\ \overline{ax_1} & 0 & 0 \end{pmatrix}
\end{aligned}$$

where $A_i(a)$ denotes the element of \mathcal{J} with $x_i = a$, all remaining terms zero in (6.1) for $i = 1, 2, 3$, and $\text{tr } b = b + \bar{b}$ for $b \in \mathcal{L}$. Thus we have $\mathfrak{g} = \mathcal{D}_0 \oplus T_1 \oplus T_2 \oplus T_3 \oplus T'_1 \oplus T'_2 \oplus T'_3 \oplus \{R_{\begin{pmatrix} \bar{\rho}_1 & 0 & 0 \\ 0 & \bar{\rho}_2 & 0 \\ 0 & 0 & \bar{\rho}_3 \end{pmatrix}}\}$. For $a \in \mathcal{L}$, we put $t_i(a) = \exp T_i(a)$ and $t'_i(a) = \exp T'_i(a)$ for $i = 1, 2, 3$. They are elements of G . For $\xi \in \mathcal{C}$, let $B_i(\xi)$ be the element of \mathcal{J} with $\xi_i = \xi$, all remaining terms zero in (6.1) for $i = 1, 2, 3$ and put $c = \exp \xi$. We define an element $S_i(c)$ of G by $S_i(c) = \exp R_{B_i(\xi)}$ for $i = 1, 2, 3$. The following proposition is well-known.

PROPOSITION 6-2. *There exist four orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 10, 27$) where S_m denotes the m -codimensional orbit, and x_m is given as follows:*

$$x_0 = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}, \quad x_1 = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 0 \end{pmatrix}, \quad x_{10} = \begin{pmatrix} 1 & & \\ & 0 & \\ & & 0 \end{pmatrix}, \quad x_{27} = (0).$$

Proof. Let X be a non-zero element of \mathcal{J} . Then we may assume that $\xi_1 = 1$ by t_i , t'_i and S_1 . By $t_2(-\bar{x}_2)$ and $t_3(-x_3)$, we have $x_2 = x_3 = 0$. Unless $\xi_2 = \xi_3 = x_1 = 0$, we have $\xi_2 = 1$ by t'_1 and S_2 . Then by $t_1(-x_1)$, we have $x_1 = 0$. If $\xi_3 \neq 0$, we have $\xi_3 = 1$ by S_3 . Thus we obtain four orbits. We shall calculate their codimension later. Q.E.D.

DEFINITION 6-3. We identify the dual vector space V^* of $V = \mathcal{J}$ with V by $\langle X, Y \rangle = \text{tr } X \circ Y$. Then the dual actions are given as follows: (i) $D^*Y = DY$ for $D \in \mathcal{D}_0$. (ii) $T_i^*(a)Y = -T'_i(a)Y$ for $a \in \mathcal{L}$ and $i = 1, 2, 3$. (iii) $T'_i{}^*(a)Y = -T_i(a)Y$ for $i = 1, 2, 3$ and $a \in \mathcal{L}$. (iv) $R_z^*Y = -R_zY$ for $z \in \mathcal{J}$.

DEFINITION 6-4. Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, the dual space has also four orbits S_m^* ($m = 0, 1, 10, 27$). We denote by A_m (resp. A_m^*) the conormal

bundle of S_m (resp. S_m^*). Clearly we have $A_0 = V \times \{0\} = A_{27}^*$ and $A_{27} = \{0\} \times V^* = A_0^*$.

(1) The isotropy subalgebra \mathfrak{g}_{x_0} at x_0 is \mathcal{D}_0 which is the Lie algebra of F_4 (See [1]). Since $A_0 = V \times \{0\}$, we have $\text{ord}_{A_0} f^s = 0$.

(2) For $\tilde{A} = D \oplus \sum_{i=1}^3 (T_i(a_i) \oplus T'_i(a'_i)) \oplus R \begin{pmatrix} a_1 & & 0 \\ & a_2 & \\ 0 & & a_3 \end{pmatrix}$, we have $d\rho(\tilde{A})x_1 = \begin{pmatrix} \alpha_1 & a_3 + a'_3 & a_2 \\ \bar{a}_3 + \bar{a}'_3 & \alpha_2 & \alpha_1 \\ \bar{a}_2 & \bar{a}_1 & 0 \end{pmatrix}$, and hence \tilde{A} is an element of the isotropy subalgebra \mathfrak{g}_{x_1} at x_1 if and only if $\tilde{A} = D \oplus T'_1(a'_1) \oplus T'_2(a'_2) \oplus [T_3(a_3) \oplus T'_3(-a_3)] \oplus R \begin{pmatrix} 0 & & \\ & 0 & \\ & & a_3 \end{pmatrix}$.

The conormal vector space $V_{x_1}^*$ is given by $V_{x_1}^* = \left\{ \begin{pmatrix} 0 & & 0 \\ 0 & 0 & \\ 0 & & \eta \end{pmatrix}; \eta \in \mathbb{C} \right\} \cong \{\eta\}$ and $d\rho_{x_{10}}(\tilde{A})\eta = -\alpha_3\eta$. For $\tilde{A}_0 = R \begin{pmatrix} 0 & & \\ & 0 & \\ & & -1 \end{pmatrix}$, we have $d\rho(\tilde{A}_0)x_1 = 0$ and $d\rho^*(\tilde{A}_0)y_1 = y_1$ where $y_1 = \begin{pmatrix} 0 & & \\ & 0 & \\ & & 1 \end{pmatrix}$. Since $\delta\chi(\tilde{A}_0) = -1$, $\text{tr}_{V_{x_1}^*} \tilde{A}_0 = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_1} f^s = -s - \frac{1}{2}$. By Corollary 1-2, we have $b_{A_1}(s)/b_{A_0}(s) = (s+1)$.

(3) For $\tilde{A} = D \oplus \sum_{i=1}^3 (T_i(a_i) \oplus T'_i(a'_i)) \oplus R \begin{pmatrix} a_1 & & \\ & a_2 & \\ & & a_3 \end{pmatrix}$, we have $d\rho(\tilde{A})x_{10} = \begin{pmatrix} \alpha_1 & a_3 & a_2 \\ \bar{a}_3 & 0 & 0 \\ \bar{a}_2 & 0 & 0 \end{pmatrix}$ and hence $\tilde{A} \in \mathfrak{g}_{x_{10}}$ if and only if $\tilde{A} = D \oplus T_1(a_1) \oplus T'_1(a'_1) \oplus T'_2(a'_2) \oplus T'_3(a'_3) \oplus R \begin{pmatrix} 0 & & \\ & a_2 & \\ & & a_3 \end{pmatrix}$. In this case, \tilde{A} acts on the conormal vector space as follows:

$$d\rho_{x_{10}}(\tilde{A}) \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & \eta_2 & y_1 \\ 0 & \bar{y}_1 & \eta_3 \end{pmatrix} = \left[\begin{array}{c|cc} 0 & & \\ \hline & -\alpha_2\eta_2 & Uy_1 - a_1\eta_3 - a'_1\eta_2 \\ & -\text{tr}(y_1\bar{a}_1) & -\frac{1}{2}(\alpha_2 + \alpha_3)y_1 \\ \hline & Uy_1 - a_1\eta_3 - a'_1\eta_2 & \\ & -\frac{1}{2}(\alpha_2 + \alpha_3)\bar{y}_1 & -\alpha_3\eta_3 - \text{tr}(\bar{y}_1 a'_1) \end{array} \right].$$

Let y_{10} (resp. y'_{10}) be the element of \mathcal{J} with $x_1 = 1$ (resp. $\xi_3 = 1$), all remaining parts zero in (6.1). Then y_{10} is a generic point of $(G_{x_{10}}, \rho_{x_{10}}, V_{x_{10}}^*)$ and y'_{10} is a point of the one-codimensional orbit. Thus we have $A_{10} = A_1^*$, $A_1 = A_{10}^*$ and $\dim A_1 \cap A_{10} = \dim V - 1$. Put $\tilde{A}_0 = R \begin{pmatrix} 0 & & \\ & -1 & \\ & & -1 \end{pmatrix}$. Then we have $d\rho(\tilde{A}_0)x_{10} = 0$ and $d\rho^*(\tilde{A}_0)y_{10} = y_{10}$. Since $\delta\chi(\tilde{A}_0) = -2$, $\text{tr}_{V_{x_{10}}^*} \tilde{A}_0 = 10 = \dim V_{x_{10}}^*$, we have $\text{ord}_{A_{10}} f^s = -2s - \frac{1}{2}$. By Corollary 1-2, we have $b_{A_{10}}(s)/b_{A_1}(s) = (s+5)$.

(4) The isotropy subalgebra $\mathfrak{g}_{x_{27}}$ is \mathfrak{g} . Put $y_{27} = x_0$ and $y'_{27} = x'_1$. Then $d\rho(\tilde{A}_0)x_{27} = 0$ and $d\rho^*(\tilde{A}_0)y_{27} = y_{27}$ for $\tilde{A}_0 = R_{-I_3}$. Since $\delta\chi(\tilde{A}_0) = -3$, $\text{tr}_{V_{x_{27}}^*} \tilde{A}_0$

$= \dim V_{x_{27}}^* = 27$, we have $\text{ord}_{A_{27}} f^s = -3s - \frac{27}{2}$. Since $A_{10} = A_1^*$, $A_{27} = A_0^*$, we have $\text{codim } A_{10} \cap A_{27} = 1$ and $b_{A_{27}}(s)/b_{A_{10}}(s) = (s+9)$. Thus we obtain the b -function $b(s) = (s+1)(s+5)(s+9)$ and the holonomy diagram (Figure 6-1). Note that the relative invariant $f(X)$ is given by the determinant $\det X$ of X in \mathcal{I} .




Figure 6-1. Holonomy diagram of $(GL(1) \times E_6, \square \otimes A_1, V(1) \otimes V(27))$

§ 7. $(GL(1) \times E_7, \square \otimes A_6, V(1) \otimes V(56))$

The representation space $V(1) \otimes V(56)$ is identified with

$$(7.1) \quad V = \{X = (x, x'); x, x' \in M(8), {}^t x = -x, {}^t x' = -x'\}.$$

Then the infinitesimal action $d\rho$ of $\mathfrak{g} = \mathfrak{gl}(1) \oplus E_7$ is given by

$$(7.2) \quad \begin{aligned} (1) \quad & (x, x') \xrightarrow{p} (px + x{}^t p, -{}^t p y - y p) \quad \text{for } p \in SL(8, \mathbf{C}) \\ (2) \quad & (x, x') \xrightarrow{c} (cx, cx') \quad \text{for } c \in \mathfrak{gl}(1) \\ (3) \quad & ((x_{ij}), (x'_{ij})) \xrightarrow{\theta} \left(\left(\sum_{m,n=1}^8 \mathcal{G}^{ijmn} y_{mn} \right), \left(- \sum_{m,n=1}^8 \mathcal{G}_{ijmn} x_{mn} \right) \right) \end{aligned}$$

where \mathcal{G} denotes a tensor, antisymmetric in its indices, and upper, lower indices satisfy the relation

$$\mathcal{G}_{i_1, \dots, i_4} = \frac{1}{4!} \sum_{j_1, \dots, j_4} I_{i_1, \dots, i_4, j_1, \dots, j_4}^{1, \dots, 8} \mathcal{G}^{j_1, \dots, j_4}.$$

Here $I_{i_1, \dots, i_4, j_1, \dots, j_4}^{1, \dots, 8}$ denotes the signature of the permutation $(1, \dots, i_1, \dots, i_4, j_1, \dots, j_4, 8)$ when $\{i_1, \dots, i_4, j_1, \dots, j_4\} = \{1, \dots, 8\}$, and 0 otherwise. The product as a Lie algebra is given as follows:

$$(7.3) \quad \begin{aligned} (1) \quad & [p, p'] = pp' - p'p \text{ where } pp' \text{ denotes the matrix multiplication} \\ (2) \quad & [p, \mathcal{P}] = \mathcal{P}' \text{ where } (\mathcal{P}')^{ijkl} = \sum_m (\mathcal{P}^{mjkl} p_{im} + \mathcal{P}^{imkl} p_{jm} + \mathcal{P}^{ijml} p_{km} + \\ & \mathcal{P}^{ijkm} p_{lm}) \\ (3) \quad & [\mathcal{P}, \mathcal{P}'] = p \text{ where } p_{ij} = \frac{2}{3} \left(\sum_{\ell, m, n} (\mathcal{P}^{\ell m ni} (\mathcal{P}')_{\ell m n j} - \frac{1}{8} \left(\sum_r \mathcal{P}^{\ell m nr} (\mathcal{P}')_{\ell m nr} \right) \delta_{ij} \right) \end{aligned}$$

PROPOSITION 7-1 (Stephen J. Haris). *There exist five orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 11, 28, 56$) where S_m denotes the m -codimensional G -orbit and x_m is given as follows.*

$$\begin{aligned} x_0 &= \left\{ \left(\begin{array}{c|c} \begin{matrix} 1 \\ -1 \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ -1 \end{matrix} & \begin{matrix} 1 \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ -1 \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ \\ 1 \\ -1 \end{matrix} \end{array} \right), 0 \right\}, & x_1 &= \left\{ \left(\begin{array}{c|c} \begin{matrix} 1 \\ -1 \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ -1 \end{matrix} & \begin{matrix} 1 \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ -1 \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ \\ 0 \\ \end{matrix} \end{array} \right), 0 \right\} \\ x_{11} &= \left\{ \left(\begin{array}{c|c} \begin{matrix} 1 \\ -1 \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ -1 \end{matrix} & \begin{matrix} 1 \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ \\ 0 \\ \end{matrix} \end{array} \right), 0 \right\}, & x_{28} &= \left\{ \left(\begin{array}{c|c} \begin{matrix} 1 \\ -1 \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ 0 \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \\ \hline \begin{matrix} \\ \\ \\ \end{matrix} & \begin{matrix} \\ \\ \\ \end{matrix} \end{array} \right), 0 \right\} \end{aligned}$$

and $x_{56} = (0, 0)$.

Proof. See [5].

Q.E.D.

We identify the dual vector space V^* with V by $\langle X, Y \rangle = \text{tr } xx' + \text{tr } yy'$ for $X = (x, x')$, $Y = (y, y') \in V$.

Then the dual action $d\rho^*$ is given as follows:

$$(7.4) \quad \begin{aligned} (1) \quad & (y, y') \xrightarrow{p^*} (-{}^t py - yp, py' + y'{}^t p) \\ (2) \quad & (y, y') \xrightarrow{c^*} (-cy, -cy') \\ (3) \quad & ((y_{ij}), (y'_{ij})) \xrightarrow{\theta^*} \left(\left(\sum_{m,n=1}^8 \mathcal{P}_{ijmn} y'_{mn} \right), \left(- \sum_{m,n=1}^8 \mathcal{P}^{ijmn} y_{mn} \right) \right) \end{aligned}$$

Since $G = GL(1) \times E_7$ is reductive, the dual triplet (G, ρ^*, V^*) has also five orbits S_m^* ($m = 0, 1, 11, 28, 56$). We denote by A_m (resp. A_m^*) the conormal bundle of S_m (resp. S_m^*).

(1) The isotropy subalgebra \mathfrak{g}_{x_0} at x_0 is the Lie algebra of E_6 (See [5], [1]). Since $A_0 = V \times \{0\} = A_{56}^*$, we have $\text{ord}_{A_0} f^s = 0$ where $f(X) = \text{Pf}(x) + \text{Pf}(x') - \frac{1}{4} \text{tr}(xx'xx') + \frac{1}{16} \text{tr}(xx')^2$ for $X = (x, x') \in V$.

(2) The isotropy subalgebra \mathfrak{g}_{x_1} at x_1 is the set $\{c \oplus p \oplus \theta\}$ satisfying the following conditions:

$$(7.4) \quad p = \begin{bmatrix} -\frac{c}{2} I_6 & \\ & \frac{3}{2} c I_2 \end{bmatrix} + \begin{bmatrix} p_1 & p_{12} & p_{13} & p_{14} \\ p'_{12} & p_2 & p_{23} & p_{24} \\ p'_{13} & p'_{23} & p_3 & p_{34} \\ 0 & 0 & 0 & p_4 \end{bmatrix} \quad \text{where } \text{tr } p_i = 0$$

for $i = 1, \dots, 4$, and $p_{ij}p'_{ij} = -\det p_{ij} \cdot I_2$ for $1 \leq i < j \leq 3$. $\mathcal{D} = (\mathcal{D}_{ijkl})$ satisfies $\mathcal{D}_{12ij} + \mathcal{D}_{34ij} + \mathcal{D}_{56ij} = 0$.

In fact, the isotropy subgroup is connected, and is isomorphic to $(GL(1) \times F_4) \cdot U$ where U is unipotent of dimension 26 (See [5]). The conormal vector space $V_{x_1}^*$ is given by

$$(7.5) \quad V_{x_1}^* = \left\{ x = \left(\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & 0 \ x \\ -x & 0 \end{array} \right), 0 \right); x \in \mathcal{C} \right\}.$$

Let y_1 be the element of $V_{x_1}^*$ with $x = 1$ in (7.5). Then it is a generic point, and $y'_1 = 0$ is the point of the one-codimensional orbit. Let A_0 be an element of \mathfrak{g}_{x_1} with $c = -\frac{1}{4}$, all remaining parts zero in (7.4). Then $d\rho(A_0)x_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$. Since $\delta_X(A_0) = -1$, $\text{tr}_{V_{x_1}^*} A_0 = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_0} f^s = -s - \frac{1}{2}$ and $b_{A_0}(s)/b_{A_0}(s) = (s + 1)$.

(3) The isotropy subalgebra $\mathfrak{g}_{x_{11}}$ at x_{11} is the set $\{c \oplus p \oplus \theta\}$ satisfying the following conditions

$$(7.6) \quad p = \left[\begin{array}{c|c} -\frac{c}{2} I_4 + p_1 & p_2 \\ \hline 0 & \frac{c}{2} I_4 + p_4 \end{array} \right] \quad \text{where } K^{-1}p_1K \in \mathfrak{sp}(2) \text{ with}$$

$$K = \begin{bmatrix} 1 & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & 1 \end{bmatrix}, \quad \text{tr } p_4 = 0, \quad (\mathcal{D}) \text{ with } \mathcal{D}_{12ij} + \mathcal{D}_{34ij} = 0 \text{ for all } i, j.$$

The conormal vector space $V_{x_{11}}^*$ is given by

$$(7.7) \quad V_{x_{11}}^* = \left\{ \tilde{Y} = \left(\begin{pmatrix} 0 & 0 \\ 0 & Y_4 \end{pmatrix}, \begin{pmatrix} Y'_1 & 0 \\ 0 & 0 \end{pmatrix} \right); y'_{12} + y'_{34} = 0 \right\}.$$

Then, for $A = c \oplus p \oplus \theta$ in $\mathfrak{g}_{x_{11}}$, we have $d\rho^*(A)\tilde{Y} = \left(\begin{pmatrix} 0 & 0 \\ 0 & Z_4 \end{pmatrix}, \begin{pmatrix} Z'_1 & 0 \\ 0 & 0 \end{pmatrix} \right)$ where

$$Z_4 = -2cY_4 - {}^tP_4Y_4 - Y_4P_4 + \left(\sum_{m,n=1}^4 \mathfrak{g}^{ijmn} y'_{mn} \right)$$

and

$$Z'_1 = -2cY'_1 + P_1Y'_1 + Y'_1{}^tP_1 - \left(\sum_{m,n=5}^8 \mathfrak{g}^{ijmn} y_{mn} \right).$$

$$\text{Put } y_{11} = \left\{ \left(\begin{array}{c|ccc} 0 & & & 0 \\ \hline & & 1 & \\ 0 & -1 & & \\ & & & 1 \\ & & & & -1 \end{array} \right), 0 \right\} \quad \text{and} \quad y'_{11} = \left\{ \left(\begin{array}{c|ccc} 0 & & & 0 \\ \hline & & & \\ 0 & & & \\ & & & 1 \\ & & & & -1 \end{array} \right), 0 \right\}.$$

Then y_{11} is a generic point and y'_{11} is a point of the unique one-codimensional orbit. Thus we have $A_{11} = A_{11}^*$ and $\dim A_1 \cap A_{11} = \dim V - 1$. Let A_c be an element of $\mathfrak{g}_{x_{11}}$ with $c = -\frac{1}{2}$, all remaining parts zero in (7.7). Then $d\rho(A_0)x_{11} = 0$ and $d\rho^*(A_0)y_{11} = y_{11}$. Since $\delta\chi(A_0) = 4c = -2$, $\text{tr}_{V_{x_{11}}}^* A_0 = -22c = 11$ and $\dim V_{x_{11}}^* = 11$, we have $\text{ord}_{A_{11}} f^s = -2s - \frac{1}{2}$ and hence $b_{A_{11}}(s)/b_{A_1}(s) = (s + \frac{1}{2})$.

(4) The isotropy subalgebra $\mathfrak{g}_{x_{28}}$ at x_{28} is the set $\{c \oplus p \oplus \theta\}$ satisfying the following conditions:

$$(7.8) \quad p = \left[\begin{array}{c|ccc} -\frac{c}{2}I_2 + p_1 & & & p_2 \\ \hline & & & \\ 0 & & & \frac{c}{6}I_2 + p_4 \\ & & & \end{array} \right] \quad \text{with} \quad \text{tr } p_1 = \text{tr } p_4 = 0$$

$$\mathfrak{g} = (\mathfrak{g}^{ijkl}) \quad \text{with} \quad \mathfrak{g}_{12ij} = 0 \quad \text{for all } i, j.$$

The conormal vector space $V_{x_{28}}^*$ is given by

$$(7.9) \quad V_{x_{28}}^* = \left\{ \tilde{Y} = \left(\begin{pmatrix} 0 & 0 \\ 0 & \underbrace{Y_4}_6 \end{pmatrix}, \begin{pmatrix} \underbrace{Y'_1}_2 & \underbrace{Y'_3}_6 \\ -{}^t\underbrace{Y'_3}_2 & \underbrace{0}_6 \end{pmatrix} \right) \in V^* \right\}.$$

Then for $A = c \oplus p \oplus \theta$ in (7.8), we have

$$d\rho^*(A)\tilde{Y} = \left(\begin{pmatrix} 0 & 0 \\ 0 & Z_4 \end{pmatrix}, \begin{pmatrix} Z'_1 & Z'_3 \\ -{}^tZ'_3 & 0 \end{pmatrix} \right)$$

where

$$\begin{aligned}
 Z_4 &= -\frac{4}{3}cY_4 - {}^t p_4 Y_4 - Y_4 p_4 + \left(\sum_{m,n} \mathcal{G}^{ijmn} y'_{mn} \right) \\
 Z'_1 &= -2cY'_1 + p_1 Y'_1 + Y'_1 p_1 - p_2 {}^t Y'_3 + Y'_3 p_2 - \left(\sum_{m,n=3}^8 \mathcal{G}^{ijmn} y_{mn} \right) \\
 Z'_3 &= -\frac{4}{3}cY'_3 + p_1 Y'_3 + Y'_3 p_1 - \left(\sum_{m,n=3}^8 \mathcal{G}^{ijmn} y_{mn} \right).
 \end{aligned}$$

Therefore, one can see that the colocalization at x_{28} has at most unique one-codimensional orbit.

Since $A_{28} = A_1^*$ and $A_{11} = A_{11}^*$, A_{28} is a good holonomic variety and $\dim A_{28} \cap A_{11} = \dim V - 1$.




Figure 7-1. Holonomy diagram of $(GL(1) \times E_7, \square \otimes A_1, V(1) \otimes V(56))$

$$\text{Put } y_{28} = \left\{ \left(\begin{array}{c|cccc} 0 & & & & \\ \hline & 1 & & & \\ & -1 & & & \\ & & 1 & & \\ & & & -1 & \\ & & & & 1 \\ & & & & & -1 & \\ & & & & & & 1 \end{array} \right), 0 \right\}. \quad \text{Then } y_{28} \text{ is a generic point.}$$

Let A_0 be an element of $\mathfrak{g}_{x_{28}}$ with $c = -\frac{3}{4}$, all remaining parts zero in (7.8). Then we have $d\rho(A_0)x_{28} = 0$ and $d\rho^*(A_0)y_{28} = y_{28}$. Since $\delta\chi(A_0) = 4c = -3$, $\text{tr}_{V_{x_{28}}} A_0 = -38c = \frac{57}{2}$ and $\dim V_{x_{28}}^* = 28$, we have $\text{ord}_{A_{28}} f^s = -3s - \frac{29}{2}$ and hence $b_{A_{28}}(s)/b_{A_{11}}(s) = (s + \frac{1}{2})$ by Corollary 1-2.

(5) Since $x_{56} = 0$, we have $(G_{x_{56}}, \rho_{x_{56}}, V_{x_{56}}^*) = (G, \rho^*, V^*)$. Since $A_{56} = \{0\} \times V^* = A_0^*$ and $A_{28} = A_1^*$, we have $\dim A_{56} \cap A_{28} = \dim V - 1$ and they intersect regularly. Let A_0 be an element of $\mathfrak{g}_{x_{28}}$ with $c = -1$, all remaining parts zero in (7.2). Then $d\rho(A_0)x_{56} = 0$ and $d\rho^*(A_0)y_{56} = y_{56}$ where $y_{56} = x_0$. Since $\delta\chi(A_0) = -4$, $\text{tr}_{V_{x_{56}}} A_0 = \dim V_{x_{56}}^* = 56$, we have $\text{ord}_{A_{56}} f^s = -4s - \frac{56}{2}$, and hence $b_{A_{56}}(s)/b_{A_{28}}(s) = s + 14$. Thus we obtain the b -function $b(s) = (s + 1)(s + \frac{1}{2})(s + \frac{1}{2})(s + 14)$ and the holonomy diagram (Figure 7-1).

§ 8. $(GL(6), A_3, V(20))$

Let V_1 be a 6-dimensional vector space spanned by u_1, \dots, u_6 . Then $G = GL(6)$ acts on V_1 by $\rho_1(g)(u_1, \dots, u_6) = (u_1, \dots, u_6)g$ for $g \in G$. The representation space $V = V(20)$ is spanned by skew-tensors $u_i \wedge u_j \wedge u_k$ ($1 \leq i < j < k \leq 6$), and $\rho = A_3$ is given by $\rho(g)(u_i \wedge u_j \wedge u_k) = \rho_1(g)u_i \wedge \rho_1(g)u_j \wedge \rho_1(g)u_k$ for $1 \leq i < j < k \leq 6$, and $g \in G$. Then it is well-known (and also one can easily check) that there exist five G -orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 5, 10, 20$) where S_m denotes the m -codimensional orbit, and $x_0 = u_1 \wedge u_2 \wedge u_3 + u_4 \wedge u_5 \wedge u_6$, $x_1 = u_1 \wedge u_2 \wedge u_3 + u_1 \wedge u_4 \wedge u_5 + u_2 \wedge u_4 \wedge u_6$, $x_5 = u_1 \wedge u_2 \wedge u_3 + u_1 \wedge u_4 \wedge u_5$, $x_{10} = u_1 \wedge u_2 \wedge u_3$, and $x_{20} = 0$. We identify the dual space V^* with V by $(\sum a_{ijk}u_i \wedge u_j \wedge u_k, \sum b_{rst}u_r \wedge u_s \wedge u_t) = \sum_{1 \leq i < j < k \leq 6} a_{ijk}b_{ijk}$. Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, there exist also five orbits S_m^* ($m = 0, 1, 5, 10, 20$) in V^* . We denote by A_m the conormal bundle of S_m . The isotropy subalgebra \mathfrak{g}_x at $x \in V(20)$ is, by definition, $\mathfrak{g}_x = \{A \in \mathfrak{gl}(6); d\rho(A)x = 0\}$ where $d\rho(A)(u_i \wedge u_j \wedge u_k) = d\rho_1(A)u_i \wedge u_j \wedge u_k + u_i \wedge d\rho_1(A)u_j \wedge u_k + u_i \wedge u_j \wedge d\rho_1(A)u_k$.

(1) The isotropy subalgebra \mathfrak{g}_{x_0} is, by simple calculation, given as follows:

$$(8.1) \quad \mathfrak{g}_{x_0} = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \in \mathfrak{gl}(6); A, B \in \mathfrak{sl}(3) \right\}.$$

We have $\mathcal{A}_0 = V \times \{0\}$, and hence $\text{ord}_{\mathcal{A}_0} f^s = 0$ where f denotes the relatively invariant irreducible polynomial of degree four (See [1], [14]).

(2) Put $x'_1 = u_1 \wedge u_2 \wedge u_6 + u_2 \wedge u_3 \wedge u_4 - u_1 \wedge u_3 \wedge u_5$. Then $x'_1 \in S_1$. By simple calculation, the isotropy subalgebra $\mathfrak{g}_{x'_1}$ at x'_1 is given by

$$(8.2) \quad \mathfrak{g}_{x'_1} = \left\{ \tilde{A} = \left(\begin{array}{c|c} A & B \\ \hline 0 & A - (\text{tr } A) \cdot I_3 \end{array} \right) \in \mathfrak{gl}(6); A, B \in M(3), \text{tr } B = 0 \right\}.$$

Therefore we have $G_{x'_1} \sim GL(3) \cdot (G_a)^3$ where \cdot denotes a semi-direct product and $G_a \cong C$. The conormal vector space $V_{x'_1}^*$ is of one-dimension with a basis $u_4 \wedge u_5 \wedge u_6$. The action $d\rho_{x'_1}$ of $\mathfrak{g}_{x'_1}$ on $V_{x'_1}^*$ is $d\rho_{x'_1}(\tilde{A})u_4 \wedge u_5 \wedge u_6 = -2\text{tr } \tilde{A} \cdot u_4 \wedge u_5 \wedge u_6$. Take $A_0 \in \mathfrak{g}_{x'_1}$ with $\text{tr } A_0 = -\frac{1}{2}$. Then we have $d\rho(A_0)x'_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$ where $y_1 = u_4 \wedge u_5 \wedge u_6$. Since $\delta\chi(A_0) = (\text{deg } f / \dim V) \text{tr } d\rho(A_0) = \frac{4}{2 \cdot 6} \times (10 \text{tr } A_0) = -1$ and $\text{tr}_{V_{x'_1}^*} A_0 = \dim V_{x'_1}^* = 1$, we have $\text{ord}_{\mathcal{A}_1} f^s = -s - \frac{1}{2}$ by Proposition 1-3. Since 0 is the point of the one-codimensional orbit, we have $\dim \mathcal{A}_0 \cap \mathcal{A}_1 = \dim V - 1$ and $\mathcal{A}_0 \cap \mathcal{A}_1$ is G_0 -prehomogeneous, i.e., \mathcal{A}_1 is a good holonomic variety by Proposition 1-5. Also we have $\mu = 1$ and $\nu = 0$ by Proposition 1-4, i.e., \mathcal{A}_0 and \mathcal{A}_1 intersect regularly. By Corollary 1-2, we have $b_{\mathcal{A}_1}(s)/b_{\mathcal{A}_0}(s) = (s + 1)$.

(3) Put $x'_5 = u_1 \wedge (u_2 \wedge u_4 + u_3 \wedge u_5) \in S_5$. Then the isotropy subalgebra $\mathfrak{g}_{x'_5}$ is given as follows:

$$(8.3) \quad \mathfrak{g}_{x'_5} = \left\{ \left(\begin{array}{c|c|c} -2\varepsilon & B & C \\ \hline 0 & A + \varepsilon I_4 & D \\ \hline 0 & 0 & \eta \end{array} \right) \in \mathfrak{gl}(6); A \in \mathfrak{sp}(2), {}^t B, D \in C^4, C \in C \right\} \\ = (\mathfrak{sp}(2) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(9))$$

where $\mathfrak{u}(9)$ denotes the Lie algebra of 9-dimensional unipotent group. Put $\omega_1 = (u_2 \wedge u_4 - u_3 \wedge u_5) \wedge u_6$, $\omega_2 = u_4 \wedge u_5 \wedge u_6$, $\omega_3 = u_3 \wedge u_4 \wedge u_6$, $\omega_4 = u_2 \wedge u_5 \wedge u_6$ and $\omega_5 = u_2 \wedge u_3 \wedge u_6$. Then the conormal vector space $V_{x'_5}^*$ is spanned by $\omega_1, \dots, \omega_5$ and $(G_{x'_5}, \rho_{x'_5}, V_{x'_5}^*) \cong (GL(1) \times Sp(2), \mathcal{A}_1 \otimes \mathcal{A}_2, V(1) \otimes V(5)) \cong (GL(1) \times SO(5), \mathcal{A}_1 \otimes \mathcal{A}_1, V(1) \otimes V(5))$, where ω_1 is a generic point and $\omega_2 = u_4 \wedge u_5 \wedge u_6$ is a point of the one-codimensional orbit. Therefore we have $\dim \mathcal{A}_1 \cap \mathcal{A}_5 = \dim V - 1$. Since the $(G_{x'_5} \cap G_0)$ -orbit of ω_2 is one-codimensional in $V_{x'_5}^*$, i.e., $\mathcal{A}_1 \cap \mathcal{A}_5$ is G_0 -prehomogeneous, \mathcal{A}_5 is a good holonomic variety by (2) and Proposition 1-5. Let A_0 be an element of $\mathfrak{g}_{x'_5}$ with $\eta = -1$ and all remaining parts zero in (8.3). Then we have

$d\rho(A_0)x'_5 = 0$ and $d\rho^*(A_0)\omega_1 = \omega_1$. Since $\delta\chi(A_0) = 2\text{tr } A_0 = -2$, $\text{tr}_{V_{x'_5}} A_0 = -5(2\varepsilon + \eta) = 5$, and $\dim V_{x'_5}^* = 5$, we have $\text{ord}_{A_5} f^s = -2s - \frac{5}{2}$. Put $A_\beta = \beta(E_{22} - E_{44}) + (\beta + 1)E_{66}$ for $\beta \in \mathbb{C}$ where E_{ij} denotes the matrix unit. Then $d\rho(A_\beta)x'_5 = 0$ and $d\rho^*(A_\beta)\omega_2 = \omega_2$. Since $\tilde{V} = V_{x'_5}^* \bmod d\rho_{x'_5}(\mathfrak{g}_{x'_5})\omega_2$ is spanned by $u_2 \wedge u_3 \wedge u_6$, we have $\text{tr}_{\mathbb{F}} A_\beta = 2\beta + 1$. Hence we have $\mu = 1$ and $\nu = 0$ by Proposition 1-4, i.e., A_1 and A_5 intersect regularly. One can also get this from the fact $m_{A_5} - m_{A_1} = 1$. By Corollary 1-2, we have $b_{A_5}(s)/b_{A_1}(s) = s + \frac{5}{2}$.

(4) Put $x_{10} = u_1 \wedge u_2 \wedge u_3 \in S_{10}$. Then the isotropy subalgebra $\mathfrak{g}_{x_{10}}$ is given as follows:

$$(8.4) \quad \mathfrak{g}_{x_{10}} = \left\{ \tilde{A} = \left(\begin{array}{c|c} A & B \\ \hline 0 & C \end{array} \right) \in \mathfrak{gl}(6); A, B, C \in M(3), \text{tr } A = 0 \right\} \\ \cong (\mathfrak{sl}(3) \oplus \mathfrak{gl}(3)) \oplus V(9), \text{ i.e., } G_{x_{10}} \sim (SL(3) \times GL(3)) \cdot (G_a)^3.$$

In general, we write $G_1 \sim G_2$ when two groups G_1 and G_2 are locally isomorphic to each other. Put $\omega_1 = u_1 \wedge u_4 \wedge u_5$, $\omega_2 = u_1 \wedge u_4 \wedge u_6$, $\omega_3 = u_1 \wedge u_5 \wedge u_6$, $\omega_4 = u_2 \wedge u_4 \wedge u_5$, $\omega_5 = u_2 \wedge u_4 \wedge u_6$, $\omega_6 = u_2 \wedge u_5 \wedge u_6$, $\omega_7 = u_3 \wedge u_4 \wedge u_5$, $\omega_8 = u_3 \wedge u_4 \wedge u_6$, $\omega_9 = u_3 \wedge u_5 \wedge u_6$ and $\omega_{10} = u_4 \wedge u_5 \wedge u_6$. Then the conormal vector space $V_{x_{10}}^*$ is spanned by $\omega_1, \dots, \omega_{10}$. The action $d\rho_{x_{10}}$ of $\mathfrak{g}_{x_{10}}$ on $V_{x_{10}}^*$ is given by

$$(8.5) \quad d\rho_{x_{10}}(\tilde{A})(\omega_1, \dots, \omega_{10}) = (\omega_1, \dots, \omega_{10}) \left(\begin{array}{c|c} d\tilde{\rho}(A \oplus C) & 0 \\ \hline B' & -\text{tr } A \end{array} \right) ('B' \in \mathbb{C}^9)$$

where $\tilde{\rho} = A_2 \otimes A_2^*$, i.e., the action of G_{x_0} induced on the subspace spanned by $\omega_1, \dots, \omega_9$ is isomorphic to a triplet $(SL(3) \times GL(3), A_1 \otimes A_1, V(3) \otimes V(3))$ as a triplet (See [1]). Then $\omega_1 + \omega_5 + \omega_9 \in S_1^*$ is a generic point and $\omega_1 + \omega_5 \in S_5^*$ is a point of the one-codimensional orbit. This implies that $\dim A_5 \cap A_{10} = \dim V - 1$. Since $A_5 \cap A_{10}$ is G_0 -prehomogeneous, A_{10} is a good holonomic variety by Proposition 1-5. Let \tilde{A} be an element of $\mathfrak{g}_{x_{10}}$ with $A = B = 0$ and $C = -\frac{1}{2}I_3$ in (8.4). Then $d\rho(\tilde{A})x_{10} = 0$ and $d\rho^*(\tilde{A})(\omega_1 + \omega_5 + \omega_9) = (\omega_1 + \omega_5 + \omega_9)$. Since $\delta\chi(\tilde{A}) = 2 \cdot \text{tr } \tilde{A} = -3$, $\text{tr}_{V_{x_{10}}^*} \tilde{A} = -7\text{tr } C = \frac{21}{2}$ and $\dim V_{x_{10}}^* = 10$, we have $\text{ord}_{A_{10}} f^s = -3s - \frac{11}{2}$ by Proposition 1-3. Let A_β be an element of $\mathfrak{g}_{x_{10}}$ with $A = B = 0$ and $C = \begin{pmatrix} 1 - (\beta/2) & 0 \\ 0 & (\beta/2)I_2 \end{pmatrix}$ in (8.4). Then $d\rho(A_\beta)x_{10} = 0$ and $d\rho(A_\beta)(\omega_1 + \omega_5) = (\omega_1 + \omega_5)$. Since $\tilde{V} = V_{x_{10}}^* \bmod d\rho_{x_{10}}(\mathfrak{g}_{x_{10}})(\omega_1 + \omega_5)$ is spanned by $u_3 \wedge u_5 \wedge u_6$, we have $\text{tr}_{\mathbb{F}} A_\beta = \beta$. This implies that $\mu = 1$ and $\nu = 0$, i.e., A_5 and A_{10} intersect regularly by Proposition 1-4. One can also get this from the fact $m_{A_{10}} - m_{A_5} = 1$. By

Corollary 1-2, we have $b_{A_{10}}(s)/b_{A_5}(s) = s + \frac{7}{2}$.

(5) Put $x_{20} = 0 \in S_{20}$. In this case, $(G_{x_{20}}, \rho_{x_{20}}, V_{x_{20}}^*) \cong (GL(6), A_3, V(20))$. $A_{x_{20}} = \{0\} \times V^*$ is a good holonomic variety. Put $\tilde{A} = -\frac{1}{3}I_6$. Then $d\rho(\tilde{A})x_{20} = 0$ and $d\rho^*(\tilde{A})x_0^* = x_0^*$ where $x_0^* = u_1 \wedge u_2 \wedge u_3 + u_4 \wedge u_5 \wedge u_6 \in S_0^*$. Since $\delta\chi(\tilde{A}) = 2\text{tr}\tilde{A} = -4$, $\text{tr}_{V_{x_{20}}^*}\tilde{A} = 20$ and $\dim V_{x_{20}}^* = 20$, we have $\text{ord}_{A_{20}}f^s = -4s - \frac{2 \cdot 0}{2}$ by Proposition 1-3. Put $A_\beta = \begin{pmatrix} a_1 & & & & & 0 \\ & \ddots & & & & \\ & & \ddots & & & \\ 0 & & & & & a_6 \end{pmatrix}$ with $a_1 = a_2 = a_4 = 1/2 - \beta/6$, $a_3 = a_5 = a_6 = \beta/3$. Then $d\rho(A_\beta)x_{20} = 0$ and $d\rho^*(A_\beta)x_1^* = x_1^*$ where $x_1^* = u_1 \wedge u_2 \wedge u_3 + u_1 \wedge u_4 \wedge u_5 + u_2 \wedge u_4 \wedge u_6$. Since $\tilde{V} = V_{x_{20}}^* \text{ mod } d\rho_{x_{20}}(\mathfrak{g}_{x_{20}})x_1^*$ is spanned by $u_3 \wedge u_5 \wedge u_6$, we have $\text{tr}_{\tilde{V}}A_\beta = \beta$. This implies that $\mu = 1$ and $\nu = 0$, i.e., A_{20} and A_{10} intersect regularly. One can also get this from $m_{A_{20}} - m_{A_{10}} = 1$. By Corollary 1-2, we have $b_{A_{20}}(s)/b_{A_{10}}(s) = s + 5$. Thus we obtain the b -function $b(s) = (s + 1)(s + \frac{5}{2})(s + \frac{7}{2})(s + 5)$ and the holonomy diagram (Figure 8-1). We denote $\textcircled{A_m}$ by \textcircled{m} .




Figure 8-1. Holonomy diagram of $(GL(6), A_3, V(20))$.

§ 9. $(GL(1) \times Sp(3), \square \otimes A_3, V(1) \otimes V(14))$

Put $\omega_1 = u_1 \wedge u_2 \wedge u_3$, $\omega_2 = u_4 \wedge u_5 \wedge u_6$, $\omega_3 = u_2 \wedge u_3 \wedge u_4$, $\omega_4 = u_1 \wedge u_5 \wedge u_6$, $\omega_5 = u_1 \wedge u_3 \wedge u_5$, $\omega_6 = u_2 \wedge u_4 \wedge u_6$, $\omega_7 = u_1 \wedge u_2 \wedge u_6$, $\omega_8 = u_3 \wedge u_4 \wedge u_5$, $\omega_9 = u_1 \wedge u_2 \wedge u_5 - u_1 \wedge u_3 \wedge u_6$, $\omega_{10} = u_2 \wedge u_4 \wedge u_5 - u_3 \wedge u_4 \wedge u_6$,

$\omega_{11} = u_1 \wedge u_2 \wedge u_4 + u_2 \wedge u_3 \wedge u_6$, $\omega_{12} = u_1 \wedge u_4 \wedge u_5 + u_3 \wedge u_5 \wedge u_6$, $\omega_{13} = u_1 \wedge u_3 \wedge u_4 - u_2 \wedge u_3 \wedge u_5$, $\omega_{14} = u_1 \wedge u_4 \wedge u_6 - u_2 \wedge u_5 \wedge u_6$. Then the representation space V is identified with the subspace of $V(20)$ in § 8 generated by $\omega_1, \dots, \omega_{14}$. Then the representation $\rho = \square \otimes A_3$ is the restriction of A_3 for $GL(6)$ to $G = GL(1) \times Sp(3)$. The orbital decomposition of this space has been completed by J-I. Igusa (See [3]). There exist five G -orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 4, 7, 14$) where S_m denotes the m -codimensional orbit, and $x_0 = \omega_1 + \omega_2$, $x_1 = \omega_7 + \omega_{13}$, $x_4 = \omega_{13}$, $x_7 = \omega_1$, $x_{14} = 0$. We identify the dual space V^* with V by $(\sum_{i=1}^{14} a_i \omega_i, \sum_{j=1}^{14} b_j \omega_j) = \sum_{k=1}^{14} a_k b_k$. Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, there exist also five G -orbits S_m^* ($m = 0, 1, 4, 7, 14$) in V^* . We denote by A_m the conormal bundle of S_m . The Lie algebra \mathfrak{g} of $G = GL(1) \times Sp(3)$ is given as follows:

$$(9.1) \quad \mathfrak{g} = \left\{ \tilde{A} = (d) \oplus \begin{pmatrix} A & B \\ C & -{}^t A \end{pmatrix}; A, B, C \in M(3), {}^t B = B, {}^t C = C \right\}.$$

(1) Since $d\rho(\tilde{A})x_0 = (d + \text{tr } A)\omega_1 + (d - \text{tr } A)\omega_2 + c_1\omega_3 + b_1\omega_4 - c_2\omega_5 - b_2\omega_6 + c_3\omega_7 + b_3\omega_8 + c_{23}\omega_9 + b_{23}\omega_{10} + c_{13}\omega_{11} + b_{13}\omega_{12} - c_{12}\omega_{13} - b_{12}\omega_{14}$ where $c_i = c_{ii}$ and $b_i = b_{ii}$ for $i = 1, 2, 3$, we have

$$(9.2) \quad \mathfrak{g}_{x_0} = \left\{ \tilde{A} = (0) \oplus \begin{pmatrix} A & 0 \\ 0 & -{}^t A \end{pmatrix}; A \in \mathfrak{sl}(3) \right\} \cong \mathfrak{sl}(3).$$

We have $A_0 = V \times \{0\}$, and hence $\text{ord}_{A_0} f^s = 0$ where f denotes the relatively invariant irreducible polynomial of degree four (See [1], [3]).

(2) Since $d\rho(\tilde{A})x_1 = (b_3 - 2b_{12})\omega_1 + 2a_{21}\omega_3 + c_2\omega_4 - 2a_{12}\omega_5 - c_1\omega_6 + (d + a_1 + a_2 - a_3)\omega_7 + 2c_{12}\omega_8 + (a_{13} - a_{32})\omega_9 - c_{13}\omega_{10} + (a_{23} - a_{31})\omega_{11} - c_{23}\omega_{12} + (d + a_3)\omega_{13} + (c_{12} - c_3)\omega_{14}$ where $a_i = a_{ii}$ for $i = 1, 2, 3$, we have

$$(9.3) \quad \mathfrak{g}_{x_1} = \left((d) \oplus \left[\begin{array}{ccc|ccc} -d+\alpha & 0 & \beta & b_1 & b_{12} & b_{13} \\ 0 & -d-\alpha & \gamma & b_{12} & b_2 & b_{23} \\ \gamma & \beta & -d & b_{13} & b_{23} & 2b_{12} \\ \hline & 0 & & d-\alpha & 0 & -\gamma \\ & & & 0 & d+\alpha & -\beta \\ & & & -\beta & -\gamma & d \end{array} \right] \right) \cong (\mathfrak{gl}(1) \oplus \mathfrak{o}(3)) \oplus V(5)$$

where $V(5)$ denotes the Lie algebra of $(G_a)^5$.

The conormal vector space $V_{x_1}^*$ is one-dimensional with a basis ω_2 . The action $d\rho_{x_1}$ of \mathfrak{g}_{x_1} on $V_{x_1}^*$ is given by $d\rho_{x_1}(\tilde{A})\omega_2 = (-d + a_1 + a_2 + a_3)\omega_2 = -4d\omega_2$. Therefore we have $A_1 = \overline{G(x_1, y_1)}$ where $y_1 = \omega_2$. Let A_0 be an

element of \mathfrak{g}_{x_1} with $d = -\frac{1}{4}$, all remaining parts zero in (9.3). Then we have $d\rho(A_0)x_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$. Since $\delta\chi(A_0) = 4d = -1$, $\text{tr}_{V_{x_1}^*}\tilde{A} = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_1}f^s = -s - \frac{1}{2}$ by Proposition 1-3. Since 0 is the point of the one-codimensional orbit, we have $\dim A_0 \cap A_1 = \dim V - 1$ and $A_0 \cap A_1$ is G_0 -prehomogeneous, i.e., A_1 is a good holonomic variety by Proposition 1-5. Also we have $\mu = 1$ and $\nu = 0$ by Proposition 1-4, i.e., A_0 and A_1 intersect regularly. By Corollary 1-2, we have $b_{A_1}(s)/b_{A_0}(s) = (s + 1)$.

(3) Since $d\rho(\tilde{A})x_i = -2b_{12}\omega_1 + 2a_{21}\omega_3 - 2a_{12}\omega_5 + 2c_{12}\omega_8 + a_{13}\omega_9 - c_{13}\omega_{10} + a_{23}\omega_{11} - c_{23}\omega_{12} + (d + a_3)\omega_{13} - c_3\omega_{14}$, we have

$$(9.4) \quad \mathfrak{g}_{x_4} = \left\{ (d) \oplus \left(\begin{array}{ccc|ccc} a_1 & 0 & 0 & b_1 & 0 & \beta \\ 0 & a_2 & 0 & 0 & b_2 & \delta \\ \alpha & \gamma & -d & \beta & \delta & \varepsilon \\ \hline c_1 & 0 & 0 & -a_1 & 0 & -\alpha \\ 0 & c_2 & 0 & 0 & -a_2 & -\gamma \\ 0 & 0 & 0 & 0 & 0 & d \end{array} \right) \right\} \\ \cong \left\{ (d) \oplus \left(\begin{array}{c|ccc|cc} -d & \alpha & \beta & \gamma & \delta & \varepsilon \\ \hline 0 & a_1 & b_1 & & 0 & \beta \\ & c_1 & -a_1 & & & -\alpha \\ \hline 0 & & 0 & a_2 & b_2 & \delta \\ & & & c_2 & -a_2 & -\gamma \\ \hline 0 & 0 & & 0 & 0 & d \end{array} \right) \right\} \\ \simeq (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)) \oplus V(5).$$

The conormal vector space $V_{x_4}^*$ is spanned by $\omega_2, \omega_4, \omega_6, \omega_7$ on which \mathfrak{g}_{x_4} acts as follows:

$$(\omega_2, \omega_4, \omega_6, \omega_7) \mapsto (\omega_2, \omega_4, \omega_6, \omega_7) \begin{pmatrix} A_1 & -b_1 & b_2 & 0 \\ -c_1 & A_2 & 0 & -b_2 \\ c_2 & 0 & A_3 & b_1 \\ 0 & -c_2 & c_1 & A_4 \end{pmatrix}$$

where $A_1 = a_1 + a_2 - 2d$, $A_2 = -a_1 + a_2 - 2d$, $A_3 = a_1 - a_2 - 2d$, $A_4 = -a_1 - a_2 - 2d$.

Hence we have $(G_{x_4}, \rho_{x_4}, V_{x_4}^*) \cong (SL(2) \times GL(2), A_1 \otimes A_1, V(2) \otimes V(2)) \cong (GL(1) \times SO(4), \square \otimes A_1, V(1) \otimes V(4))$.

Clearly, $y_4 = \omega_4 + \omega_6$ is its generic point, and ω_2 is a point of the one-codimensional orbit. Since $A_1 = \overline{G(x_1, \omega_2)}$, we have $\dim A_1 \cap A_4 = \dim V - 1$. Since $A_1 \cap A_4$ is G_0 -prehomogeneous, A_4 is a good holonomic variety by (2) and Proposition 1-5. Let A_0 be an element of \mathfrak{g}_{x_4} with d

$= -\frac{1}{2}$ and all remaining parts zero in (9.4). Then $d\rho(A_0)x_4 = 0$ and $d\rho^*(A_0)y_4 = y_4$. Since $\delta\chi(A_0) = 4d = -2$, $\text{tr}_{V_{x_4}^*}A_0 = -8d = 4$ and $\dim V_{x_4}^* = 4$, we have $\text{ord}_{A_1}f^s = -2s - \frac{4}{2}$. Let A_β be an element of \mathfrak{g}_{x_4} with $a_1 = \frac{1}{2}(1 - \beta)$, $d = -\frac{1}{4}(\beta + 1)$, all remaining parts zero in (9.4). Then we have $d\rho(A_\beta)x_4 = 0$, $d\rho^*(A_\beta)\omega_2 = \omega_2$ and $\text{tr}_{V_{x_4}}A_\beta = \beta$ where $\tilde{V} = V_{x_4}^* \bmod d\rho_{x_4}(\mathfrak{g}_{x_4})\omega_2$. This implies that A_4 and A_1 intersect regularly by Proposition 1-4. By Corollary 1-2, we have $b_{A_4}(s)/b_{A_1}(s) = (s + 2)$.

(4) Since $d\rho(\tilde{A})x_7 = (d + a_1 + a_2 + a_3)\omega_1 + c_1\omega_3 - c_2\omega_5 + c_3\omega_7 + c_{23}\omega_9 + c_{13}\omega_{11} - c_{12}\omega_{13}$, we have

$$(9.5) \quad \mathfrak{g}_{x_7} = \left\{ \tilde{A} = (-\text{tr } A) \oplus \left(\frac{A}{0} \middle| \frac{B}{-{}^t A} \right); {}^t B = B \right\} \simeq \mathfrak{gl}(3) \oplus V(6).$$

The conormal vector space $V_{x_7}^*$ is spanned by $\omega_2, \omega_4, \omega_6, \omega_8, \omega_{10}, \omega_{12}, \omega_{14}$, and \mathfrak{g}_{x_7} acts on $V_{x_7}^*$ as follows:

$$(9.6) \quad d\rho_{x_7}(\tilde{A})(\omega_2, \omega_4, \dots, \omega_{14}) = (\omega_2, \omega_4, \dots, \omega_{14}) \left(\frac{2 \text{tr } A}{0} \middle| \frac{B}{2 \text{tr } A \cdot I_6 \oplus d\rho_1^*(A)} \right)$$

where ${}^t B \in C^6$ and $\rho_1 = 2A_1$.

Then $y_7 = \omega_4 + \omega_{10}$ is its generic point, and $\omega_4 + \omega_6$ is a point of the one-codimensional orbit. Since $A_4 = \overline{G(x_4, \omega_4 + \omega_6)}$, we have $\dim A_4 \cap A_7 = \dim V - 1$. Since $A_4 \cap A_7$ is G_0 -prehomogeneous, A_7 is a good holonomic variety by (3) and Proposition 1-5. Let A_0 be an element of \mathfrak{g}_{x_7} with $A = \frac{1}{4}I_3$ and $B = 0$ in (9.5). Then $d\rho(A_0)x_7 = 0$ and $d\rho^*(A_0)y_7 = y_7$. Since $\delta\chi(A_0) = -4 \text{tr } A = -3$, $\text{tr}_{V_{x_7}^*}A_0 = 10 \text{tr } A = \frac{15}{2}$ and $\dim V_{x_7}^* = 7$, we have $\text{ord}_{A_7}f^s = -3s - \frac{6}{2}$. Let A_β be an element of \mathfrak{g}_{x_7} with $a_1 = a_2 = \frac{\beta}{4}$, $a_3 = \frac{1}{2} - \frac{\beta}{4}$, all remaining parts zero in (9.5). Then $d\rho(A_\beta)x_7 = 0$, $d\rho(A_\beta)(\omega_4 + \omega_6) = (\omega_4 + \omega_6)$ and $\text{tr}_{V_{x_7}}A_\beta = \beta$ where $\tilde{V} = V_{x_7}^* \bmod d\rho_{x_7}(\mathfrak{g}_{x_7})(\omega_4 + \omega_6) = C\omega_8$. This implies that A_4 and A_7 intersect regularly by Proposition 1-4. By Corollary 1-2, we have $b_{A_7}(s)/b_{A_4}(s) = (s + \frac{5}{2})$.

(5) Since $x_{14} = 0$, we have $(G_{x_{14}}, \rho_{x_{14}}, V_{x_{14}}^*) \cong (GL(1) \times Sp(3), \square \otimes A_3, V(1) \otimes V(14))$ and $A_{14} = \{0\} \times V^*$ is a good holonomic variety. Take $\tilde{A} = (-1) \oplus (0) \in \mathfrak{g} = \mathfrak{gl}(1) \oplus \mathfrak{sp}(3)$. Then $d\rho(\tilde{A})x_{14} = 0$, $d\rho^*(\tilde{A})(\omega_1 + \omega_2) = (\omega_1 + \omega_2)$. Since $\delta\chi(\tilde{A}) = -4$, $\text{tr}_{V_{x_{14}}^*}\tilde{A} = 14$ and $\dim V_{x_{14}}^* = 14$, we have $\text{ord}_{A_{14}}f^s = -4s - \frac{14}{2}$. Since $A_{14} = A_0^*$, $A_7 = A_1^*$ where A_m^* denotes the conormal bundle of $S_m^*(\subset V^*)$, they intersect regularly by (2). Note that $(G, \rho, V) \cong (G, \rho^*, V^*)$ since $G = GL(1) \times Sp(3)$ is reductive. By Corollary 1-2, we have $b_{A_{14}}(s)/b_{A_7}(s) = s + \frac{7}{2}$. Since $b_{A_{14}}(s) = b(s)$ and $b_{A_0}(s) = 1$, we obtain the b -function




Figure 9-1. Holonomy diagram of $(GL(1) \times Sp(3), \square \otimes A_3, V(1) \otimes V(14))$.

$b(s) = (s + 1)(s + 2)(s + \frac{5}{2})(s + \frac{7}{2})$ and the holonomy diagram (Figure 9-1). We denote A_m by \textcircled{m} .

§ 10. $(GL(7), A_3, V(35))$

The representation space $V = V(35)$ is spanned by the skew-tensors $u_i \wedge u_j \wedge u_k$ ($1 \leq i < j < k \leq 7$) of degree three, on which $G = GL(7)$ acts as in § 8. Then it is known (See [6], [7]) that there exist ten orbits $S_m = \rho(G)x_m$ ($m = 0, 1, 4, 7, 9, 10, 14, 15, 22, 35$), where S_m denotes the m -codimensional orbit, and $x_0 = u_2 \wedge u_3 \wedge u_4 + u_5 \wedge u_6 \wedge u_7 + u_1 \wedge (u_2 \wedge u_5 + u_3 \wedge u_6 + u_4 \wedge u_7)$, $x_1 = u_2 \wedge u_3 \wedge u_5 + u_3 \wedge u_4 \wedge u_6 + u_1 \wedge (u_2 \wedge u_7 - u_4 \wedge u_5)$, $x_4 = u_1 \wedge u_3 \wedge u_4 + u_2 \wedge u_5 \wedge u_6 + u_1 \wedge u_2 \wedge u_7$, $x_7 = u_2 \wedge u_3 \wedge u_4 + u_1 \wedge (u_2 \wedge u_5 + u_3 \wedge u_6 + u_4 \wedge u_7)$, $x_9 = u_1 \wedge u_2 \wedge u_3 + u_4 \wedge u_5 \wedge u_6$, $x_{10} = u_1 \wedge u_2 \wedge u_6 - u_1 \wedge u_3 \wedge u_5 + u_2 \wedge u_3 \wedge u_4$, $x_{14} = u_1 \wedge (u_2 \wedge u_5 + u_3 \wedge u_6 + u_4 \wedge u_7)$, $x_{15} = u_1 \wedge (u_2 \wedge u_4 + u_3 \wedge u_5)$, $x_{22} = u_1 \wedge u_2 \wedge u_3$ and $x_{35} = 0$. Note that we chose these representative points x_m so that the isotropy subalgebra \mathfrak{g}_{x_m} at x_m will be the standard form. The relative invariant $f(x)$ of this space is of degree seven (See [1], [14]). Since $(G, \rho, V) \cong (G, \rho^*, V^*)$, there exist also ten G -orbits S_m^* ($m = 0, 1, 4, 7, 9, 10, 14, 15, 22, 35$) in V^* . We denote by A_m (resp. A_m^*) the conormal bundle of S_m (resp. S_m^*). Clearly we have $A_0 = V \times \{0\} = A_{35}^*$ and $A_{35} = \{0\} \times V^* = A_0^*$.

(1) The isotropy subalgebra \mathfrak{g}_{x_0} at x_0 is given as follows (See [1]).

$$(10.1) \quad \mathfrak{g}_{x_0} = \left\{ \left(\begin{array}{ccc|ccc} 2d & 2e & 2f & 2a & 2b & 2c \\ \hline a & & & 0 & f & -e \\ b & & X & -f & 0 & d \\ c & & & e & -d & 0 \\ \hline d & 0 & -c & b & & \\ e & c & 0 & -a & & -{}^tX \\ f & -b & a & 0 & & \end{array} \right) ; X \in \mathfrak{sl}(3) \right\} \cong \mathfrak{g}_2 .$$

Since $A_0 = V \times \{0\}$, we have $\text{ord}_{A_0} f^s = 0$.

(2) The isotropy subalgebra \mathfrak{g}_{x_1} at x_1 is given as follows.

$$(10.2) \quad \mathfrak{g}_{x_1} = \left\{ \left(\begin{array}{cc|cc|ccc} \frac{d}{2} + \alpha + \beta & a_{12} & & b_{12}I_2 & \gamma_1 & \gamma_2 & \gamma_3 \\ a_{21} & \frac{d}{2} - \alpha + \beta & & & \gamma_4 & \gamma_5 & \gamma_6 \\ \hline & b_{21}I_2 & \frac{d}{2} + \alpha - \beta & a_{12} & -\gamma_2 & \gamma_7 & -\gamma_1 \\ & & a_{21} & \frac{d}{2} - \alpha - \beta & -\gamma_5 & \gamma_8 & -\gamma_4 \\ \hline & & & & -d & 2b_{21} & 2b_{12} \\ & & 0 & & b_{12} & -d + 2\beta & 0 \\ & & & & b_{21} & 0 & -d - 2\beta \end{array} \right) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)) \oplus V(8).$$

The conormal vector space $V_{x_1}^*$ is spanned by $u_5 \wedge u_6 \wedge u_7$. Then $d\rho_{x_1}(A)u_5 \wedge u_6 \wedge u_7 = 3d u_5 \wedge u_6 \wedge u_7$ for $A \in \mathfrak{g}_{x_1}$. Since 0 is the point of the one-codimensional G -orbit, A_0 and A_1 intersect regularly with codimension one. Let A_0 be an element of \mathfrak{g}_{x_1} with $d = \frac{1}{3}$, all remaining parts zero in (10.2). Then $d\rho(A_0)x_1 = 0$ and $d\rho^*(A_0)y_1 = y_1$ where $y_1 = u_5 \wedge u_6 \wedge u_7$. Since $\delta_\chi(A_0) = (\deg f / \dim V) \cdot \text{tr}_V A_0 = 3 \text{tr} A_0 = -3d = -1$ (See Proposition 1-9), $\text{tr}_{V_{x_1}^*} A_0 = \dim V_{x_1}^* = 1$, we have $\text{ord}_{A_1} f^s = -s - \frac{1}{2}$ and hence $b_{A_1}(s)/b_{A_0}(s) = (s + 1)$. We have also $A_1 = A_{22}^*$, and hence $A_{22} = A_1^*$.

(3) The isotropy subalgebra \mathfrak{g}_{x_4} at x_4 is given as follows.

$$(10.3) \quad \mathfrak{g}_{x_4} = \left\{ \left(\begin{array}{cc|cc|c} -\text{tr} X & 0 & \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ 0 & -\text{tr} Y & \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \hline & 0 & X & 0 & & & \beta_2 \\ & & & & & & -\beta_1 \\ \hline & 0 & 0 & Y & & & -\alpha_4 \\ & & & & & & \alpha_3 \\ \hline & 0 & 0 & 0 & & & \text{tr}(X+Y) \end{array} \right) \right\} \\ \cong (\mathfrak{gl}(2) \oplus \mathfrak{gl}(2)) \cdot \mathfrak{u}(10).$$

The conormal vector space $V_{x_4}^*$ is spanned by $\omega_1 = u_3 \wedge u_5 \wedge u_7$, $\omega_2 = u_3$

$\wedge u_6 \wedge u_7$, $\omega_3 = u_4 \wedge u_5 \wedge u_7$, $\omega_4 = u_4 \wedge u_6 \wedge u_7$. Then we have $(G_{x_4}, \rho_{x_4}, V_{x_4}^*) \cong (GL(2) \times GL(2), A_1 \otimes A_1, V(2) \otimes V(2))$, and $y_4 = \omega_1 + \omega_4$ is its generic point, $y'_4 = \omega_1$ is a point of the one-codimensional orbit. Since the colocalization $(G_{x_4}, \rho_{x_4}, V_{x_4}^*)$ is an irreducible regular P.V., A_4 is a good holonomic variety by Corollary 1-8. Let A_0 be an element of \mathfrak{g}_{x_4} with $\mathfrak{g}_X = -\frac{1}{3}I_2$, all remaining parts zero in (10.3). Then $d\rho(A_0)x_4 = 0$ and $d\rho^*(A_0)y_4 = y_4$. Since $\delta\chi(A_0) = 3 \operatorname{tr} A_0 = -2$, $\operatorname{tr}_{V_{x_4}^*} A_0 = 4 \dim V_{x_4}^*$, we have $\operatorname{ord}_{A_0} f^s = -2s - \frac{4}{2}$ by Proposition 1-3. We have also $d\rho^*(A_0)y'_4 = y'_4$ and $\operatorname{tr}_{\mathfrak{V}} A_0 = 1$ where $\tilde{V} = V_{x_4}^* \bmod d\rho_{x_4}(\mathfrak{g}_{x_4})y'_4 = C\omega_4$. This implies that A_1 and A_4 intersect regularly with codimension one by Proposition 1-4. By Corollary 1-2, we have $b_{A_4}(s)/b_{A_1}(s) = (s+2)$. We have also $A_4 = A_{15}^*$ and hence $A_{15} = A_4^*$.

(4) The isotropy subalgebra \mathfrak{g}_{x_7} at x_7 is given as follows.

$$(10.4) \quad \mathfrak{g}_{x_7} = \left\{ \left(\begin{array}{c|ccc|ccc} \varepsilon & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 \\ \hline & & & & \delta_4 & (\gamma_3 + \delta_3) & \delta_2 \\ 0 & & X & & \delta_3 & \delta_5 & (\gamma_1 + \delta_1) \\ \hline & & & & (\gamma_2 + \delta_2) & \delta_1 & \delta_6 \\ 0 & & 0 & & -\varepsilon I_3 - X & & \end{array} \right) ; X \in \mathfrak{sl}(3) \right\}$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(3)) \oplus \mathfrak{u}(12).$$

Put $\omega_1 = u_5 \wedge u_6 \wedge u_7$, $\omega_2 = u_2 \wedge u_6 \wedge u_7$, $\omega_3 = u_4 \wedge u_5 \wedge u_6$, $\omega_4 = u_3 \wedge u_5 \wedge u_7$, $\omega_5 = u_2 \wedge u_5 \wedge u_7 - u_3 \wedge u_6 \wedge u_7$, $\omega_6 = u_4 \wedge u_5 \wedge u_7 - u_3 \wedge u_5 \wedge u_6$, $\omega_7 = u_2 \wedge u_5 \wedge u_6 + u_4 \wedge u_6 \wedge u_7$. Then the conormal vector space $V_{x_7}^*$ is spanned by these $\omega_1, \dots, \omega_7$. The action $d\rho_{x_7}$ of \mathfrak{g}_{x_7} on $V_{x_7}^*$ is given by

$$(10.5) \quad d\rho_{x_7}(A)(\omega_1, \dots, \omega_7) = (\omega_1, \dots, \omega_7) \left(\begin{array}{c|cccccc} 3\varepsilon & * & * & * & * & * \\ \hline 0 & 2\varepsilon I_6 & \oplus & d\rho_1^*(X) & & \end{array} \right)$$

where $\rho_1 = 2A_1$ for $SL(3)$.

Here $y_7 = \omega_2 + \omega_3 + \omega_4$ is its generic point, and $y'_7 = \omega_2 + \omega_3$ is a point of the one-codimensional orbit. This implies that $A_7 = A_{10}^*$, $A_{10} = A_7^*$ and $\dim A_4 \cap A_7 = \dim V - 1$. Since $A_4 \cap A_7$ is G_0 -prehomogeneous, A_7 is a good holonomic variety. Let A_0 be an element of \mathfrak{g}_{x_7} with $\varepsilon = \frac{1}{2}$, all remaining parts zero in (10.4). Then $d\rho(A_0)x_7 = 0$ and $d\rho^*(A_0)y_7 = y_7$. Since $\delta\chi(A_0) = 3 \operatorname{tr} A_0 = -6\varepsilon = -3$, $\operatorname{tr}_{V_{x_7}^*} A_0 = \frac{1}{2} \cdot 7$ and $\dim V_{x_7}^* = 7$, we have $\operatorname{ord}_{A_0} f^s = -3s - \frac{7}{2}$ by Proposition 1-3. Let A_β be an element of \mathfrak{g}_{x_7} with $\varepsilon = \frac{\beta}{6} + \frac{1}{3}$, $X = \begin{pmatrix} \eta & & & & & & \\ & \eta & & & & & \\ & & -2\eta & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \end{pmatrix}$ with $\eta = \frac{\beta}{6} - \frac{1}{6}$, all remaining parts zero in (10.4). Then we have $d\rho(A_\beta)x_7 = 0$, $d\rho^*(A_\beta)y'_7 = y'_7$ and $\operatorname{tr}_{\mathfrak{V}} A_\beta = \beta$ where $\tilde{V} = V_{x_7}^* \bmod d\rho_{x_7}(\mathfrak{g}_{x_7})y'_7 = C\omega_4$. This implies that A_4 and A_7 intersect regularly

by Proposition 1-4. By Corollary 1-2, we have $b_{A_7}(s)/b_{A_4}(s) = (s + \frac{5}{2})$.

(5) The isotropy subalgebra \mathfrak{g}_{x_9} at x_9 is given as follows.

$$(10.6) \quad \mathfrak{g}_{x_9} = \left\{ \begin{pmatrix} X & 0 & Z \\ 0 & Y & W \\ 0 & 0 & \varepsilon \end{pmatrix}; X, Y \in \mathfrak{sl}(3), Z, W \in \mathbf{C}^3 \right\}$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(3) \oplus \mathfrak{sl}(3) \oplus V(6)).$$

The conormal vector space $V_{x_9}^*$ is spanned by $u_i \wedge u_j \wedge u_7 (1 \leq i \leq 3; 4 \leq j \leq 6)$. By seeing the weights, we have $(G_{x_9}, \rho_{x_9}, V_{x_9}^*) \cong (SL(3) \times GL(3), A_1 \otimes A_1, V(3) \otimes V(3))$. Since this is an irreducible regular P.V., A_9 is a good holonomic variety by Corollary 1-8. As a generic point, we may take $y_9 = (u_1 \wedge u_4 + u_2 \wedge u_5 + u_3 \wedge u_6) \wedge u_7$, and $y'_9 = (u_1 \wedge u_4 + u_2 \wedge u_5) \wedge u_7$ is a point of the one-codimensional orbit. This implies that $A_9 = A_{14}^*$, $A_{14} = A_9^*$ and $\dim A_4 \cap A_9 = \dim V - 1$. Let A_0 be an element of \mathfrak{g}_{x_9} with $\varepsilon = -1$, all remaining parts zero in (10.6). Then $d\rho(A_0)x_9 = 0$, $d\rho^*(A_0)y_9 = y_9$. Since $\delta\chi(A_0) = 3 \operatorname{tr} A_0 = -3$, $\operatorname{tr}_{V_{x_9}^*} A_0 = 9\varepsilon = -9$, $\dim V_{x_9}^* = 9$, we have $\operatorname{ord}_{A_0} f^s = -3s - \frac{9}{2}$. Let A_β be an element of \mathfrak{g}_{x_9} with $\varepsilon = ((\beta + 2)/3)$, $X = \begin{pmatrix} \eta & & \\ & \eta & \\ & & -2\eta \end{pmatrix}$ with $\eta = ((1 - \beta)/3)$, all remaining parts zero in (10.6). Then we have $d\rho(A_0)x_9 = 0$, $d\rho^*(A_0)y'_9 = y'_9$ and $\operatorname{tr}_{\tilde{V}} A_0 = \beta$ where $\tilde{V} = V_{x_9}^* \bmod d\rho_{x_9}(\mathfrak{g}_{x_9})y'_9 = \mathbf{C}u_3 \wedge u_6 \wedge u_7$. This implies that A_4 and A_9 intersect regularly. By Corollary 1-2, we have $b_{A_9}(s)/b_{A_4}(s) = (s + 3)$.

(6) The isotropy subalgebra $\mathfrak{g}_{x_{10}}$ at x_{10} is given as follows.

$$(10.7) \quad \mathfrak{g}_{x_{10}} = \left\{ A = \left(\begin{array}{c|c|c} \varepsilon I_3 + X & B & C \\ \hline 0 & -2\varepsilon I_3 + X & F \\ \hline 0 & 0 & \eta \end{array} \right); X \in \mathfrak{sl}(3), \operatorname{tr} B = 0, C, D \in \mathbf{C}^3 \right\}$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(3)) \oplus \mathfrak{u}(14).$$

Put $\omega_1 = u_5 \wedge u_6 \wedge u_7$, $\omega_2 = u_4 \wedge u_6 \wedge u_7$, $\omega_3 = u_4 \wedge u_5 \wedge u_7$, $\omega_4 = u_1 \wedge u_4 \wedge u_7$, $\omega_5 = u_2 \wedge u_5 \wedge u_7$, $\omega_6 = u_3 \wedge u_6 \wedge u_7$, $\omega_7 = (u_1 \wedge u_5 + u_2 \wedge u_4) \wedge u_7$, $\omega_8 = (u_2 \wedge u_6 + u_3 \wedge u_5) \wedge u_7$, $\omega_9 = (u_1 \wedge u_6 + u_3 \wedge u_4) \wedge u_7$, $\omega_{10} = u_4 \wedge u_5 \wedge u_6$. Then the conormal vector space $V_{x_{10}}^*$ is spanned by $\omega_1, \dots, \omega_{10}$, and the action $d\rho_{x_{10}}$ of $\mathfrak{g}_{x_{10}}$ on $V_{x_{10}}^*$ is given as follows.

$$(10.8) \quad d\rho_{x_{10}}(A)(\omega_1, \dots, \omega_{10}) = (\omega_1, \dots, \omega_{10}) \left(\begin{array}{c|c|c} (4\varepsilon - \eta)I_3 + X & B' & F \\ \hline 0 & (\varepsilon - \eta)I_6 + d\rho_1^*(X) & 0 \\ \hline 0 & 0 & 6\varepsilon \end{array} \right)$$

where $\rho_1 = 2A_1$ for $SL(3)$.

Then $y_{10} = \omega_4 + \omega_8 + \omega_{10}$ is a generic point. There exist two one-codimensional orbits. As a representative point, we may take $y'_{10} = \omega_8 + \omega_{10}$ and $y''_{10} = \omega_4 + \omega_5 + \omega_6$ respectively. This implies that $\dim A_7 \cap A_{10} = \dim A_9 \cap A_{10} = \dim V - 1$. Since $A_{10} = A_7^*$, A_{10} is a good holonomic variety. Let A_0 be an element of $\mathfrak{g}_{x_{10}}$ with $\varepsilon = \frac{1}{6}$, $\eta = -\frac{5}{6}$, all remaining parts zero in (10.7). Then $d\rho(A_0)x_{10} = 0$ and $d\rho^*(A_0)y_{10} = y_{10}$. Since $\delta\chi(A_0) = -9\varepsilon + 3\eta = -4$, $\text{tr}_{V_{x_{10}}} A_0 = 24\varepsilon - 9\eta = \frac{23}{2}$, and $\dim V_{x_{10}}^* = 10$, we have $\text{ord}_{A_{10}} f^s = -4s - \frac{1}{2}$ by Proposition 1-3.

Since $d\rho^*(A_0)y'_{10} = y'_{10}$ and $\text{tr}_V A_0 = 1$ where $\tilde{V} = V_{x_{10}}^* \bmod d\rho_{x_{10}}(\mathfrak{g}_{x_{10}})y'_{10} = C\omega_4$, A_7 and A_{10} intersect regularly by Proposition 1-4. Let A_β be an element of $\mathfrak{g}_{x_{10}}$ with $\varepsilon = \frac{\beta}{6}$, $\eta = \frac{\beta}{6} - 1$, all remaining parts zero in (10.7). Then we have $d\rho(A_\beta)x_{10} = 0$, $d\rho^*(A_\beta)y'_{10} = y''_{10}$, and $\text{tr}_V A_\beta = \beta$ where $\tilde{V} = V_{x_{10}}^* \bmod d\rho_{x_{10}}(\mathfrak{g}_{x_{10}})y''_{10} = C\omega_{10}$. This implies that A_9 and A_{10} intersect regularly by Proposition 1-4. By Corollary 1-2, we have $b_{A_{10}}(s)/b_{A_7}(s) = (s + 3)$ and $b_{A_{10}}(s)/b_{A_9}(s) = (s + \frac{5}{2})$.

(7) The isotropy subalgebra $\mathfrak{g}_{x_{14}}$ at x_{14} is given as follows.

$$(10.9) \quad \mathfrak{g}_{x_{14}} = \left\{ \left(\begin{array}{c|c} -2\varepsilon & Y \\ \hline 0 & \varepsilon I_6 + X \end{array} \right); X \in \mathfrak{sp}(3), {}^t Y \in \mathbf{C}^6 \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{sp}(3)) \oplus V(6).$$

Put $\omega_1 = u_2 \wedge u_3 \wedge u_4$, $\omega_2 = u_5 \wedge u_6 \wedge u_7$, $\omega_3 = u_3 \wedge u_4 \wedge u_5$, $\omega_4 = u_2 \wedge u_6 \wedge u_7$, $\omega_5 = u_2 \wedge u_4 \wedge u_6$, $\omega_6 = u_3 \wedge u_5 \wedge u_7$, $\omega_7 = u_2 \wedge u_3 \wedge u_7$, $\omega_8 = u_4 \wedge u_5 \wedge u_6$, $\omega_9 = u_2 \wedge u_3 \wedge u_6 - u_2 \wedge u_4 \wedge u_7$, $\omega_{10} = u_3 \wedge u_5 \wedge u_6 - u_4 \wedge u_5 \wedge u_7$, $\omega_{11} = u_2 \wedge u_3 \wedge u_5 + u_3 \wedge u_4 \wedge u_7$, $\omega_{12} = u_2 \wedge u_5 \wedge u_6 + u_4 \wedge u_6 \wedge u_7$, $\omega_{13} = u_2 \wedge u_4 \wedge u_5 - u_3 \wedge u_4 \wedge u_6$, $\omega_{14} = u_2 \wedge u_5 \wedge u_7 - u_3 \wedge u_6 \wedge u_7$. The conormal vector space $V_{x_{14}}^*$ is spanned by these $\omega_1, \dots, \omega_{14}$. By seeing the weights, we have $(G_{x_{14}}, \rho_{x_{14}}, V_{x_{14}}^*) \cong (GL(1) \times Sp(3), \square \otimes A_3, V(1) \otimes V(14))$. Since this is an irreducible regular P.V., A_{14} is a good holonomic variety by Corollary 1-8. As we have seen in § 9, $y_{14} = \omega_1 + \omega_2$ is a generic point. Let A_0 be an element of $\mathfrak{g}_{x_{14}}$ with $\varepsilon = -\frac{1}{3}$, $X = Y = 0$ in (10.9). Then $d\rho(A_0)x_{14} = 0$ and $d\rho^*(A_0)y_{14} = y_{14}$. Since $\delta\chi(A_0) = 3 \text{tr} A_0 = -4$, $\text{tr}_{V_{x_{14}}} A_0 = 14 \times 3\varepsilon = -14$ and $\dim V_{x_{14}}^* = 14$, we have $\text{ord}_{A_{14}} f^s = -4s - \frac{1}{2}$. Since $A_{14} = A_9^*$, $A_7 = A_{10}^*$, and A_9 and A_{10} intersect regularly with codimension one, so do A_{14} and A_7 . By Corollary 1-2, we have $b_{A_{14}}(s)/b_{A_7}(s) = (s + \frac{7}{2})$.

(8) The isotropy subalgebra $\mathfrak{g}_{x_{15}}$ at x_{15} is given as follows.

$$(10.10) \quad \mathfrak{g}_{x_{15}} = \left\{ \left(\begin{array}{c|c|c} -2\varepsilon & W & U \\ \hline 0 & \varepsilon I_4 + X & Z \\ \hline 0 & 0 & \eta I_2 + Y \end{array} \right); X \in \mathfrak{sp}(2), Y \in \mathfrak{sl}(2), Z \in M(4, 2) \right\} \\ {}^t W \in \mathbf{C}^2, {}^t U \in \mathbf{C}^4, \varepsilon, \eta \in \mathbf{C}$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{sp}(2) \oplus \mathfrak{gl}(2) \oplus \mathfrak{u}(14)).$$

Put $\omega_i = u_{i+1} \wedge u_6 \wedge u_7$ ($1 \leq i \leq 4$), $\omega_5 = u_1 \wedge u_6 \wedge u_7$, $\omega_{6+j} = (u_2 \wedge u_4 - u_3 \wedge u_5) \wedge u_{6+j}$, $\omega_{8+j} = u_2 \wedge u_3 \wedge u_{6+j}$, $\omega_{10+j} = u_2 \wedge u_5 \wedge u_{6+j}$, $\omega_{12+j} = u_3 \wedge u_4 \wedge u_{6+j}$, $\omega_{14+j} = u_4 \wedge u_5 \wedge u_{6+j}$ ($j = 0, 1$). Then the conormal vector space $V_{x_{15}}^*$ is spanned by $\omega_1, \dots, \omega_{15}$. The action $d\rho_{x_{15}}$ of $\mathfrak{g}_{x_{15}}$ on $V_{x_{15}}^*$ is as follows.

$$(10.11) \quad d\rho_{x_{15}}(A)(\omega_1, \dots, \omega_{15}) = (\omega_1, \dots, \omega_{15}) \times \left(\begin{array}{c|c|c} -(\varepsilon + 2\eta)I_4 + d\rho_1(X) & * & * \\ \hline 0 & 2\varepsilon - 2\eta & * \\ \hline 0 & 0 & -(2\varepsilon + \eta)I_{10} + d\rho_2^*(X \oplus Y) \end{array} \right)$$

where $\rho_1 = A_1$ for $Sp(2)$ and $\rho_2 = A_2 \otimes A_1$ for $Sp(2) \times SL(2)$. Since $A_{15} = A_4^*$ and A_4 is a good holonomic variety, A_{15} is also a good holonomic variety. $y_{15} = \omega_5 + \omega_{11} + \omega_{12}$ is a generic point. Let A_0 be an element of $\mathfrak{g}_{x_{15}}$ with $\varepsilon = -\frac{1}{6}$, $\eta = -\frac{2}{3}$, all remaining parts zero in (10.10). Then $d\rho(A_0)x_{15} = 0$ and $d\rho^*(A_0)y_{15} = y_{15}$. Since $\delta\chi(A_0) = 3 \operatorname{tr} A_0 = 6(\varepsilon + \eta) = -5$, $\operatorname{tr}_{V_{x_{15}}^*} A_0 = -22\varepsilon - 20\eta = \frac{19}{6}$, and $\dim V_{x_{15}}^* = 15$, we have $\operatorname{ord}_{A_{15}} f^s = -5s - \frac{19}{2}$. Since $A_{15} = A_4^*$, $A_{14} = A_9^*$, $A_{10} = A_7^*$, we have $\dim A_{15} \cap A_{14} = \dim A_{15} \cap A_{10} = \dim V - 1$ and they intersect regularly. By Corollary 1-2, we have $b_{A_{15}}(s)/b_{A_{14}}(s) = (s + 3)$ and $b_{A_{15}}(s)/b_{A_{10}}(s) = (s + \frac{7}{2})$.

(9) The isotropy subalgebra $\mathfrak{g}_{x_{22}}$ at x_{22} is given as follows.

$$(10.12) \quad \mathfrak{g}_{x_{22}} = \left\{ \tilde{X} = \left(\begin{array}{c|c} X & Z \\ \hline 0 & \varepsilon I_4 + Y \end{array} \right); X \in \mathfrak{sl}(3), Y \in \mathfrak{sl}(4), Z \in M(3, 4) \right\} \\ \cong (\mathfrak{sl}(3) \oplus \mathfrak{gl}(4) \oplus V(12)).$$

The conormal vector space $V_{x_{22}}^*$ is spanned by $u_i \wedge u_j \wedge u_k$ ($4 \leq i < j < k \leq 7$) and $u_i \wedge u_j \wedge u_k$ ($1 \leq i \leq 3, 4 \leq j < k \leq 7$). The action $d\rho_{x_{22}}$ of $\mathfrak{g}_{x_{22}}$ is given by

$$(10.13) \quad d\rho(\tilde{X})(u_5 \wedge u_6 \wedge u_7, \dots) \\ = (u_5 \wedge u_6 \wedge u_7, \dots) \left(\begin{array}{c|c} Y - 3\varepsilon I_4 & * \\ \hline 0 & -2\varepsilon I_{18} + d\rho_1^*(X \oplus Y) \end{array} \right)$$

where $\rho_1 = A_1 \otimes A_2$ for $SL(3) \times SL(4)$. For example, $y_{22} = u_1 \wedge (u_4 \wedge u_5 + u_6 \wedge u_7) + u_2 \wedge u_4 \wedge u_6 + u_3 \wedge u_5 \wedge u_7$ is a generic point. Since $A_{22} = A_1^*$, A_{22} is a good holonomic variety. Let A_0 be an element of $\mathfrak{g}_{x_{22}}$ with $\varepsilon = -\frac{1}{2}$, $X = Y = Z = 0$ in (10.12). Then $d\rho(A_0)x_{22} = 0$ and $d\rho^*(A_0)y_{22} = y_{22}$. Since $\delta\chi(A_0) = 12\varepsilon = -6$, $\operatorname{tr}_{V_{x_{22}}^*} A_0 = -48\varepsilon = 24$ and $\dim V_{x_{22}}^* = 22$, we have $\operatorname{ord}_{A_{22}} f^s = -6s - \frac{26}{2} (= -6s - 13)$. Since $A_{22} = A_1^*$ and $A_{15} = A_4^*$, we have

$\dim A_{22} \cap A_{15} = \dim V - 1$ and they intersect regularly.

By Corollary 1-2, we have $b_{A_{22}}(s)/b_{A_{15}}(s) = (s + 4)$.

(10) The isotropy subalgebra $\mathfrak{g}_{x_{35}}$ at $x_{35} = 0$ is \mathfrak{g} itself and we have $(G_{x_{35}}, \rho_{x_{35}}, V_{x_{35}}^*) = (G, \rho^*, V^*) \cong (GL(7), A_3, V(35))$. Then $y_{35} = x_0 = u_2 \wedge u_3 \wedge u_4 \wedge u_5 \wedge u_6 \wedge u_7 + u_1 \wedge (u_2 \wedge u_5 + u_3 \wedge u_6 + u_4 \wedge u_7)$ is its generic point. Put $A_0 = -\frac{1}{3}I_7$. Then $d\rho(A_0)x_{35} = 0$ and $d\rho^*(A_0)y_{35} = y_{35}$. Since $\delta\chi(A_0) = 3 \operatorname{tr} A_0 = -7$, $\operatorname{tr}_{V_{x_{35}}^*} A_0 = -35$ and $\dim V_{x_{35}}^* = 35$, we have $\operatorname{ord}_{A_{35}} f^s = -7s - \frac{3 \cdot 5}{2}$. Since $A_{22} = A_1^*$ and $A_{35} = A_0^*$, they intersect regularly with codimension one. By Corollary 1-2, we have $b_{A_{35}}(s)/b_{A_{22}}(s) = (s + 5)$. Since $b_{A_0}(s) = 1$ and $b_{A_{35}}(s) = b(s)$, we obtain the b -function $b(s) = (s + 1)(s + 2)(s + \frac{5}{2})(s + \frac{7}{2})(s + 3)(s + 4)(s + 5)$, and the holonomy diagram (Figure 10-1). We denote A_m by \textcircled{m} .

Note that the colocalization at $x_1, x_4, x_7, x_9, x_{14}, x_{22}$ and x_{35} has the unique one-codimensional orbit respectively, and the colocalization at x_{10} and x_{15} has the two one-codimensional orbits respectively. Therefore we have obtained all one-codimensional intersections among the conormal bundles.




Figure 10-1. Holonomy diagram of $(GL(7), A_3, V(35))$.

§ 11. $(SL(5) \times GL(3), A_2 \otimes A_1, V(10) \otimes V(3))$

Let $V(10)$ be a vector space spanned by 2-forms $u_i \wedge u_j$ ($1 \leq i < j \leq 5$). Then the representation space is identified with $V = V(10) \oplus V(10) \oplus V(10)$ (See [1]). Let A be the conormal bundle of an orbit S in V and A^* that of an orbit S^* in V^* . When $A = A^*$, we say that S and S^* are the dual orbits of each other. We denote by $S_{i,j}^{(k)}$ the i -codimensional orbit whose dual orbit is j -codimensional, where k denotes the dimension of the central torus of the isotropy subgroup of this orbit. When there is no confusion, denote this by S_i or $S_{i,j}$. We denote by $A_{i,j}^{(k)}$ (resp. $A_{i,j}, A_i$) the conormal bundle of $S_{i,j}^{(k)}$ (resp. $S_{i,j}, S_i$). We identify V and its dual V^* by taking $(u_i \wedge u_{i'}, u_j \wedge u_{j'}, u_k \wedge u_{k'})$ ($i < i', j < j', k < k'$) as a dual basis.

PROPOSITION 11-1. *This space has following twenty five orbits $S_{i,j}^{(k)}$.*

- (1) $S_{0,30}^{(0)}: (u_1 \wedge u_2 + u_3 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5, u_1 \wedge u_3 + u_2 \wedge u_5) (= x_0)$
- (2) $S_{1,21}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_5 + u_3 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5) (= x_1)$
- (3) $S_{2,16}^{(2)}: (u_1 \wedge u_2, u_2 \wedge u_3 + u_3 \wedge u_4, u_1 \wedge u_3 + u_4 \wedge u_5) (= x_2)$
- (4) $S_{3,15}^{(3)}: (u_1 \wedge u_2, u_3 \wedge u_4, u_1 \wedge u_3 + u_4 \wedge u_5) (= x_3)$
- (5) $S_{3,13}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5) (= x'_3)$
- (6) $S_{4,11}^{(3)}: (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_4 \wedge u_5) (= x_4)$
- (7) $S_{5,8}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_1 \wedge u_5 + u_3 \wedge u_4) (= x_5)$
- (8) $S_{6,12}^{(1)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_2 \wedge u_3 + u_4 \wedge u_5) (= x_6)$
- (9) $S_{7,9}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_4 \wedge u_5) (= x_7)$
- (10) $S_{7,7}^{(1)}: (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_2 \wedge u_3 + u_1 \wedge u_5) (= x'_7)$
- (11) $S_{7,7}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_2 \wedge u_4 + u_3 \wedge u_5) (= x''_7)$
- (12) $S_{8,18}^{(2)}: (u_1 \wedge u_2 + u_3 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5, 0) (= x_8)$
- (13) $S_{8,14}^{(1)}: (u_1 \wedge u_2, u_3 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_4) (= x'_8)$
- (14) $S_{8,5}^{(3)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_1 \wedge u_5 + u_2 \wedge u_4) (= x''_8)$
- (15) $S_{9,7}^{(4)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_2 \wedge u_4) (= x_9)$
- (16) $S_{10,10}^{(3)}: (u_1 \wedge u_2, u_3 \wedge u_4 + u_1 \wedge u_5, 0) (= x_{10})$
- (17) $S_{11,4}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_1 \wedge u_4 + u_2 \wedge u_3) (= x_{11})$
- (18) $S_{12,6}^{(3)}: (u_1 \wedge u_2, u_3 \wedge u_4, 0) (= x_{12})$
- (19) $S_{13,3}^{(3)}: (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, 0) (= x_{13})$
- (20) $S_{14,8}^{(2)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_1 \wedge u_4) (= x_{14})$
- (21) $S_{15,3}^{(1)}: (u_1 \wedge u_2, u_1 \wedge u_3, u_2 \wedge u_3) (= x_{15})$
- (22) $S_{16,2}^{(3)}: (u_1 \wedge u_2, u_1 \wedge u_3, 0) (= x_{16})$
- (23) $S_{18,8}^{(2)}: (u_1 \wedge u_2 + u_3 \wedge u_4, 0, 0) (= x_{18})$
- (24) $S_{21,1}^{(2)}: (u_1 \wedge u_2, 0, 0) (= x_{21})$

$$(25) \quad S_{30,0}^{(1)}: (0, 0, 0) (= x_{30}).$$

Proof. It is easy to check that the non-regular P.V. $(SL(5) \times GL(2), A_2 \otimes A_1, V(10) \otimes V(2))$ has eight orbits which are represented by the following points; [1] $(0, 0)$, [2] $(u_1 \wedge u_2, 0)$, [3] $(u_1 \wedge u_2 + u_3 \wedge u_4, 0)$, [4] $(u_1 \wedge u_2, u_1 \wedge u_3)$, [5] $(u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4)$, [6] $(u_1 \wedge u_2, u_3 \wedge u_4)$, [7] $(u_1 \wedge u_2, u_3 \wedge u_4 + u_1 \wedge u_5)$, [8] $(u_1 \wedge u_2 + u_3 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5)$. Therefore, for a point $x = (x_1, x_2, x_3)$ of V , we may assume that (x_1, x_2) is one of these points. In the first three cases, repeating the same argument, we obtain (12), (16), (18), (19), (22), (23), (24) and (25). For $\lambda \in \mathcal{C}$, we define $S_{ij}(\lambda)$ by $S_{ij}(\lambda)u_k = u_k$ for $k \neq i$ and $S_{ij}(\lambda)u_i = u_i + \lambda u_j$. Then $S_{ij}(\lambda)$ is an element of $\rho(G)$. Put $x_3 = \sum_{i < j} a_{ij}u_i \wedge u_j$. First we consider the case [4], i.e., $(x_1, x_2) = (u_1 \wedge u_2, u_1 \wedge u_3)$. Assume that $a_{45} \neq 0$. Then we may assume that $x_3 = a_{23}u_2 \wedge u_3 + u_4 \wedge u_5$. In fact, we have $a_{35} = 0$ by $S_{43}(-a_{35}/a_{45})$ and so on. If $a_{23} = 0$, then we have (9). If $a_{23} \neq 0$, then we have (8). Next assume that $a_{45} = 0$. If one of a_{ij} ($i = 2, 3; j = 4, 5$) is not zero, we may assume that $x_3 = u_2 \wedge u_4 + a_{15}u_1 \wedge u_5 + a_{35}u_3 \wedge u_5$. If $a_{35} \neq 0$ (resp. $a_{35} = 0$ and $a_{15} \neq 0$, $a_{35} = a_{15} = 0$), then we have (11) (resp. (14), (15)). If any $a_{ij} = 0$ ($i = 2, 3; j = 4, 5$), then we may assume that $x_3 = a_{14}u_1 \wedge u_4 + a_{23}u_2 \wedge u_3$. If $a_{14} \neq 0$ and $a_{23} \neq 0$ (resp. $a_{14} \neq 0$ and $a_{23} = 0$, $a_{14} = 0$ and $a_{23} \neq 0$, $a_{14} = a_{23} = 0$), then we have (17) (resp. (20), (21), (22)). Next we consider the case [5], i.e., $(x_1, x_2) = (u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4)$. If $a_{35} \neq 0$ or $a_{45} \neq 0$, we may assume that $x_3 = a_{23}u_2 \wedge u_3 + u_4 \wedge u_5$ and hence we have (5) (resp. (6)) for $a_{23} \neq 0$ (resp. $a_{23} = 0$). If $a_{35} = a_{45} = 0$ and one of a_{k5} ($k = 1, 2$) is not zero, then we may assume that $x_3 = a_{23}u_2 \wedge u_3 + a_{34}u_3 \wedge u_4 + u_1 \wedge u_5$ and hence we have (7) (resp. (10), (14)) for $a_{34} \neq 0$ (resp. $a_{34} = 0$ and $a_{23} \neq 0$, $a_{34} = a_{23} = 0$). If $a_{k5} = 0$ for $1 \leq k \leq 4$, we may assume that $x_3 = a_{13}u_1 \wedge u_3 + a_{14}u_1 \wedge u_4 + a_{23}u_2 \wedge u_3 + a_{34}u_3 \wedge u_4$. Then we have (13) (resp. we have (19); it is reduced to the case [4]) for $a_{34} \neq 0$ (resp. $x_3 = 0; a_{34} = 0$ and $x_3 \neq 0$). Now we consider the case [6], i.e., $(x_1, x_2) = (u_1 \wedge u_2, u_3 \wedge u_4)$. (i) If $a_{35} \neq 0$ or $a_{45} \neq 0$, we may assume that $x_3 = a_{12}u_1 \wedge u_3 + a_{15}u_1 \wedge u_5 + a_{25}u_2 \wedge u_5 + u_4u_5$. Moreover if $a_{25} \neq 0$, then we have (3) (resp. (4)) for $a_{13} \neq 0$ (resp. $a_{13} = 0$). If $a_{25} = 0$, then we have (4) (resp. it is reduced to the case [4] or [5]) for $a_{15} \neq 0$ (resp. $a_{15} = 0$). (ii) If $a_{35} = a_{45} = 0$, it is reduced to the previous cases. Next we shall consider the case [7], i.e., $(x_1, x_2) = (u_1 \wedge u_2, u_3 \wedge u_4 + u_1 \wedge u_5)$. (i) If $a_{35} \neq 0$ or $a_{45} \neq 0$, then we may assume that $a_{45} = 1$ and $a_{35} = a_{24} = a_{14} = a_{12} = 0$. By $S_{53}(\lambda)$, $S_{41}(\mu)$, $S_{21}(\nu)$ and $GL(3)$, we have $x_3 = (a_{13} + (a_{15} - a_{34})\lambda + (a_{23} + a_{25})\nu + \lambda^2)u_1 \wedge u_3 + (a_{23} + \lambda a_{25})u_2 \wedge u_3$

+ $a_{25}u_2 \wedge u_5 + (a_{15} + \nu a_{25} + \mu + \lambda)u_1 \wedge u_5 + (a_{34} + \mu - \lambda)u_3 \wedge u_4 + u_4 \wedge u_5$.
 If $a_{25} \neq 0$, we may take λ, μ, ν so that $a_{13} + (a_{15} - a_{34})\lambda + \nu(a_{23} + \lambda a_{25}) + \lambda^2 = a_{15} + \nu a_{25} + \mu + \lambda = a_{34} + \mu - \lambda = 0$ and hence we have $x_3 = \alpha u_2 \wedge u_3 + u_2 \wedge u_5 + u_4 \wedge u_5$. If $\alpha \neq 0$ (resp. $\alpha = 0$), then we have (2) (resp. (3)) by $S_{35}(-1/2\alpha), S_{42}(-1/2), S_{12}(1/4\alpha), \{u_3 \mapsto (1/\sqrt{\alpha})u_3, u_4 \mapsto \sqrt{\alpha}u_4, u_j \mapsto u_j (j \neq 3, 4)\}$ and $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1/2\alpha \\ 0 & 0 & 1/\sqrt{\alpha} \end{pmatrix}$ (resp. by $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$) and $\{u_5 \mapsto u_3, u_3 \mapsto u_5, u_4 \mapsto -u_4, u_j \mapsto u_j (j = 1, 2)\}$. If $a_{25} = 0$, taking λ and μ satisfying $a_{15} + \mu + \lambda = a_{34} + \mu - \lambda = 0$, we have $x_3 = a'_{13}u_1 \wedge u_3 + a'_{23}u_2 \wedge u_3 + u_4 \wedge u_5$. If $a'_{23} \neq 0$ (resp. $a'_{23} = a'_{13} = 0$), then we have (2) (resp. (6)). If $a'_{23} = 0$ and $a'_{13} \neq 0$, then we have (4) by $S_{41}(\lambda), \begin{pmatrix} 1 & & & \\ & 1 & -1/\gamma & \\ & 0 & 1 & \end{pmatrix}, S_{35}\left(-\frac{1}{\gamma}\right), S_{53}\left(\frac{\gamma}{2}\right), \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \gamma/2 & 1 \end{pmatrix}$ and $\{u_1 \mapsto (1/2\gamma)u_1, u_2 \mapsto 2\gamma u_2, u_j \mapsto u_j (j \neq 1, 2)\}$ where $\gamma = \sqrt{-a'_{13}}$. (ii) If $a_{35} = a_{45} = 0$, it is reduced to the previous cases. Finally we shall consider the case [8], i.e., $(x_1, x_2) = (u_1 \wedge u_2 + u_3 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5)$. The isotropy subalgebra \mathfrak{h} of $\mathfrak{sl}(5) \oplus \mathfrak{gl}(2)$ at this point (x_1, x_2) is given by

$$\mathfrak{h} = \left\{ \begin{pmatrix} a_1 & & \gamma_1 & & \\ a_1 & a_2 & \beta_1 & \gamma_2 & \beta_2 \\ -2\gamma_2 & & a_3 & & 2\gamma_1 \\ \beta_1 & -\gamma_2 & \beta_2 & a_4 & a_2 \\ & & -\gamma_2 & & a_5 \end{pmatrix} \oplus \begin{pmatrix} -(a_1 + a_2) & -\gamma_2 \\ \gamma_1 & -(a_4 + a_5) \end{pmatrix} \right\}$$

with $a_1 + a_2 = a_3 + a_4, a_2 + a_3 = a_4 + a_5$ and $\sum_{i=1}^5 a_i = 0$.

Taking one-parameter subgroups from \mathfrak{h} , we obtain the following actions which fix (x_1, x_2) . (i) $\alpha_1(\lambda): u_1 \mapsto u_1 + \lambda u_2, u_j \mapsto u_j (j \neq 1)$, (ii) $\alpha_2(\lambda): u_5 \mapsto u_5 + \lambda u_4, u_j \mapsto u_j (j \neq 5)$, (iii) $\beta_1(\lambda): u_1 \mapsto u_1 + \lambda u_4, u_3 \mapsto u_3 + \lambda u_2, u_j \mapsto u_j (j \neq 1, 3)$ (iv) $\beta_2(\lambda): u_3 \mapsto u_3 + \lambda u_4, u_5 \mapsto u_5 + \lambda u_2, u_j \mapsto u_j (j \neq 3, 5)$ (v) $\gamma_1(\lambda): u_2 \mapsto u_2 - \lambda u_4, u_3 \mapsto u_3 + \lambda u_1, u_5 \mapsto u_5 + 2\lambda u_3 + \lambda^2 u_1, u_j \mapsto u_j (j \neq 2, 3, 5)$ and $(x_1, x_2, x_3) \mapsto (x_1, \lambda x_1 + x_2, x_3)$, (vi) $\gamma_2(\lambda): u_1 \mapsto u_1 - 2\lambda u_3 + \lambda^2 u_5, u_3 \mapsto u_3 - \lambda u_5, u_4 \mapsto u_4 + \lambda u_2, u_j \mapsto u_j (j \neq 1, 3, 4)$ and $(x_1, x_2, x_3) \mapsto (x_1 - \lambda x_2, x_2, x_3)$. We have also $\xi_1(\lambda)$ (resp. $\xi_2(\lambda)$): $(x_1, x_2, x_3) \mapsto (x_1, x_2, \lambda x_1 + x_3)$ (resp. $(x_1, x_2, \lambda x_2 + x_3)$) and $\eta(\mu): (x_1, x_2, x_3) \mapsto (x_1, x_2, \mu x_3)$ with $\mu \neq 0$. By using these actions, we shall do the orbital decomposition leaving (x_1, x_2) fixed. If at least one of a_{13}, a_{15} and a_{35} is not zero, then by $\gamma_1, \gamma_2, \xi_1, \xi_2$ and η , we may assume that $a_{13} = 1, a_{34} = a_{35} = a_{45} = 0$. (i) If $a_{15} \neq 0$, by $\alpha_1, \beta_1, \xi_1, \xi_2, \beta_2, \alpha_2, \gamma_1$ and η , we may assume that $x_3 = a_{24}u_2 \wedge u_4 + u_1 \wedge u_5$. If $a_{24} \neq 0$ (resp. $a_{24} = 0$), we have (1) (resp. (2)). (ii) If $a_{15} = 0$, by $\xi_1, \xi_2, \beta_1, \beta_2$ and α_1 , we may assume that $x_3 = u_1 \wedge u_3 + a_{24}u_2 \wedge u_4 + a_{25}u_2 \wedge u_5$. If $a_{25} \neq 0$, then we have (1).

If $a_{25} = 0$, then it is reduced to previous cases. Finally, if $a_{13} = a_{15} = a_{35} = 0$, we may assume that $a_{15} = a_{25} = a_{35} = a_{45} = 0$ by the action of ξ_1, ξ_2 and γ_2 . By considering (x_1, x_3) instead of (x_1, x_2) , it is reduced to the previous cases. We shall see later, by calculating the isotropy subalgebras, that these orbits are different from each other. Q.E.D.

(1) Put $x'_0 = (3u_3 \wedge u_4 - u_2 \wedge u_5, u_1 \wedge u_5 - 2u_2 \wedge u_4, 3u_2 \wedge u_3 - u_1 \wedge u_4)$. Then the isotropy subalgebra $\mathfrak{g}_{x'_0}$ is the following standard form.

$$(11.1) \quad \mathfrak{g}_{x'_0} = \left\{ \begin{pmatrix} 4\alpha & \beta & & & \\ 4\gamma & 2\alpha & 2\beta & & \\ & 3\gamma & 0 & 3\beta & \\ & & 2\gamma & -2\alpha & 4\beta \\ & & & \gamma & -4\alpha \end{pmatrix} \oplus \begin{pmatrix} 2\alpha & \beta & & & \\ 2\gamma & & 2\beta & & \\ & \gamma & & -2\alpha & \end{pmatrix} \right\} \\ \cong \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha \end{pmatrix} \right\} = \mathfrak{sl}(2).$$

Since $A_0 = V \times \{0\}$, we have $\text{ord}_{r_0^{0,30}} f^s = 0$ where f denotes the relative invariant of degree 15 (See [1]).

(2) The isotropy subgroup at x_1 is locally isomorphic to $(GL(1) \times GL(1)) \cdot U(2)$ where $U(2)$ denotes a 2-dimensional unipotent group (See [1]). The conormal vector space $V_{x_1}^*$ is spanned by $(u_3 \wedge u_5, 0, 0) \in S_{21,1}^*$. We have $\dim A_0 \cap A_1 = \dim V - 1$; $\text{ord}_{A_1} f^s = -s - \frac{1}{2}$ and $b_{A_1}(s)/b_{A_0}(s) = (s + 1)$.

(3) The isotropy subalgebra \mathfrak{g}_{x_2} at x_2 is given as follows.

$$(11.2) \quad \mathfrak{g}_{x_2} = \left\{ A = \begin{pmatrix} \varepsilon & 0 & 0 & 0 & -\beta \\ \alpha & -2(\varepsilon + \eta) & \beta & 0 & 0 \\ 0 & 0 & \eta & 0 & -\alpha \\ 0 & 0 & -\beta & -2(\varepsilon + \eta) & \gamma \\ 0 & 0 & 0 & 0 & 3(\varepsilon + \eta) \end{pmatrix} \oplus \begin{pmatrix} \varepsilon + 2\eta & & & & \\ & 2\varepsilon + \eta & & & \\ -\beta & -\alpha & -(\varepsilon + \eta) & & \end{pmatrix} \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(3)).$$

The conormal vector space $V_{x_2}^*$ is spanned by $v_1 = (u_3 \wedge u_5, 0, 0)$, and $v_2 = (0, u_1 \wedge u_5, 0)$, and the action $d\rho_{x_2}$ is given by

$$d\rho_{x_2}(A)(v_1, v_2) = (v_1, v_2) \begin{pmatrix} -(4\varepsilon + 6\eta) & 0 \\ 0 & -(6\varepsilon + 4\eta) \end{pmatrix}$$

- i) $V_{x_2}^* - S_{x_2}^* \leftrightarrow v_1 + v_2 = (u_3 \wedge u_5, u_1 \wedge u_5, 0)$, i.e., $v_1 + v_2$ is a generic point of $V_{x_2}^*$, where $S_{x_2}^*$ is the singular set of the P.V. $(G_{x_2}, \rho_{x_2}, V_{x_2}^*)$. We use this notation from now in § 11. Put $y = y_1 v_1 + y_2 v_2$.
- ii) $(S_{x_2}^*)_1 \leftrightarrow d\rho_1(A) = -(4\varepsilon + 6\eta) \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 = (0, u_1 \wedge u_5, 0) \in S_{21,1}^*$, i.e., $(S_{x_2}^*)_1 = \{y \in V_{x_2}^*; f_1^*(y) = 0\} = \overline{\rho_{x_2}(G_{x_2}) \cdot v_2}$ and $f_1^*(\rho_{x_2}(g)y) = \rho_1(g)f_1^*(y)$ for $y \in V_{x_2}^*$, $g \in G_{x_2}$. From now on, we use this notation in § 11.
- iii) $(S_{x_2}^*)_2 \leftrightarrow d\rho_2(A) = -(6\varepsilon + 4\eta) \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 = (u_3 \wedge u_5, 0, 0) \in S_{21,1}^*$
- iv) $-\delta\chi = d\rho_1 + d\rho_2, \text{tr}_{V_{x_2}^*} = d\rho_1 + d\rho_2$
- v) $\text{ord}_{A_2} f^s = -2s - 2/2$.

Since the Hessian of the localization $f_{x_2}(z) = z_1 z_2$ ($z = z_1 v_1 + z_2 v_2 \in V_{x_2}$) of $f(x)$ is not identically zero, $A_2 = A_{2,16}^{(2)}$ is a good holonomic variety. We have $\dim A_1 \cap A_2 = \dim V - 1$ and $b_{A_2}(s)/b_{A_1}(s) = (s + 1)$.

(4) The isotropy subalgebra \mathfrak{g}_{x_3} is given as follows.

$$(11.3) \quad \mathfrak{g}_{x_3} = \left\{ A = \begin{pmatrix} \varepsilon & \alpha & 0 & 0 & \beta \\ 0 & \eta & 0 & 0 & 0 \\ 0 & 0 & \xi & 0 & 0 \\ 0 & 0 & \gamma & \varepsilon & \beta \\ 0 & 0 & 0 & 0 & -(2\varepsilon + \eta + \xi) \end{pmatrix} \right. \\ \left. \oplus \begin{pmatrix} -(\varepsilon + \eta) & & & & \\ & -(\varepsilon + \xi) & & & \\ & & & & (\varepsilon + \eta + \xi) \end{pmatrix} \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(3)).$$

The conormal vector space $V_{x_3}^*$ is spanned by $v_1 = (u_3 \wedge u_5, 0, 0)$, $v_2 = (0, u_2 \wedge u_5, 0)$, $v_3 = (0, 0, u_2 \wedge u_3)$, and

$$d\rho_{x_3}(A)(v_1, v_2, v_3) = (v_1, v_2, v_3) \begin{pmatrix} 3\varepsilon + 2\eta \\ 3\varepsilon + 2\xi \\ -(\varepsilon + 2\eta + 2\xi) \end{pmatrix}$$

- i) $V_{x_3}^* - S_{x_3}^* \leftrightarrow v_1 + v_2 + v_3 = (u_3 \wedge u_5, u_2 \wedge u_5, u_2 \wedge u_3) \in S_{15,3}^*$
- ii) $(S_{x_3}^*)_1 \leftrightarrow d\rho_1(A) = 3\varepsilon + 2\eta \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 + v_3 = (0, u_2 \wedge u_5, u_2 \wedge u_3) \in S_{16,2}^*$
- iii) $(S_{x_3}^*)_2 \leftrightarrow d\rho_2(A) = 3\varepsilon + 2\xi \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 + v_3 = (u_3 \wedge u_5, 0, u_2 \wedge u_3) \in S_{16,2}^*$
- iv) $(S_{x_3}^*)_3 \leftrightarrow d\rho_3(A) = -(\varepsilon + 2\eta + 2\xi) \leftrightarrow f_3^*(y) = y_3 \leftrightarrow v_1 + v_2 = (u_3 \wedge u_5, u_2 \wedge u_5, 0) \in S_{16,2}^*$
- v) $-\delta\chi = d\rho_1 + d\rho_2 + d\rho_3, \text{tr}_{V_{x_3}^*} = d\rho_1 + d\rho_2 + d\rho_3$.

Since the localization $f_{x_3}(z) = z_1 z_2 z_3$ ($z = \sum z_i v_i \in V_{x_3}$) is non-degenerate, $A_{3,15}$ is a good holonomic variety and $\text{ord}_{A_{3,15}} f^s = -3s - \frac{3}{2}$. We have $\dim A_2 \cap A_{3,15} = \dim V - 1$ and $b_{A_{3,15}}(s)/b_{A_2}(s) = (s+1)$.

(5) The isotropy subalgebra \mathfrak{g}_{x_3} at x_3 is given as follows.

$$(11.4) \quad \mathfrak{g}_{x_3} = \left\{ A = \begin{pmatrix} -2(\varepsilon+\eta) & \alpha & \beta & 0 & 0 \\ 0 & \varepsilon & \gamma & 0 & -2\alpha \\ 0 & 0 & \eta & 0 & 0 \\ 0 & 0 & -\alpha & -(3\varepsilon+\eta) & \delta \\ 0 & 0 & 0 & 0 & 4\varepsilon+2\eta \end{pmatrix} \right. \\ \left. \oplus \begin{pmatrix} \varepsilon+2\eta & 0 & 0 \\ -\gamma & 2\varepsilon+\eta & 0 \\ \beta & -\alpha & -\varepsilon-\eta \end{pmatrix} \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(4)).$$

The conormal vector space $V_{x_3}^*$ is spanned by $v_1 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_2 = (u_3 \wedge u_4, 0, 0)$, $v_3 = (u_3 \wedge u_5, 0, 0)$, and

$$d\rho_{x_3}(A)(v_1, v_2, v_3) = (v_1, v_2, v_3) \begin{pmatrix} -(6\varepsilon+4\eta) & 0 & 0 \\ 0 & 2(\varepsilon-\eta) & 0 \\ -2\gamma & -\delta & -5(\varepsilon+\eta) \end{pmatrix}$$

- i) $V_{x_3}^* - S_{x_3}^* \leftrightarrow v_1 + v_2 = (u_2 \wedge u_5 + u_3 \wedge u_4, -u_3 \wedge u_5, 0) \in S_{13,3}^*$
- ii) $(S_{x_3}^*)_1 \leftrightarrow d\rho_1(A) = -(6\varepsilon+4\eta) \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 \in S_{21,1}^*$
- iii) $(S_{x_3}^*)_2 \leftrightarrow d\rho_2(A) = 2(\varepsilon-\eta) \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 \in S_{16,2}^*$
- iv) $-\delta\chi = 2d\rho_1 + d\rho_2 = -10(\varepsilon+\eta)$, $\text{tr}_{V_{x_3}^*} = 2d\rho_1 + \frac{3}{2}d\rho_2 = -9\varepsilon - 11\eta$.

Since $\dim A_{3,13} \cap A_2 = \dim A_{3,13} \cap A_1 = \dim V - 1$ and they intersect G_0 -prehomogeneously, $A_{3,13}$ is a good holonomic variety and $\text{ord}_{A_{3,13}} f^s = -3s - \frac{4}{2}$. The intersection exponent of $A_{3,13}$ and A_1 is $(1:0)$. We have $b_{A_{3,13}}(s)/b_{A_1}(s) = (s+1)(s+\frac{3}{2})$ and $b_{A_{3,13}}(s)/b_{A_2}(s) = (s+\frac{3}{2})$.

(6) The isotropy subalgebra \mathfrak{g}_{x_4} at x_4 is given as follows.

$$(11.5) \quad \mathfrak{g}_{x_4} = \left\{ A = \begin{pmatrix} \varepsilon & \alpha & \gamma & 0 & 0 \\ 0 & \eta & \beta & 0 & 0 \\ 0 & 0 & \xi & 0 & 0 \\ 0 & 0 & -\alpha & \varepsilon-\eta+\xi & \delta \\ 0 & 0 & 0 & 0 & -2(\varepsilon+\xi) \end{pmatrix} \right. \\ \left. \oplus \begin{pmatrix} -(\varepsilon+\eta) & 0 & 0 \\ -\beta & -(\varepsilon+\xi) & 0 \\ 0 & 0 & (\varepsilon+\eta+\xi) \end{pmatrix} \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(4)).$$

The conormal vector space $V_{x_4}^*$ is spanned by $v_1 = (u_3 \wedge u_4, 0, 0)$, $v_2 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_3 = (u_3 \wedge u_5, 0, u_2 \wedge u_3)$, $v_4 = (0, 0, u_2 \wedge u_3)$, and

$$d\rho_{x_4}(A)(v_1, \dots, v_4) = (v_1, \dots, v_4) \begin{bmatrix} 2(\eta - \varepsilon) & 0 & 0 & 0 \\ 0 & 3\varepsilon + 2\eta & 0 & 0 \\ \alpha & -2\beta & 3\varepsilon + \eta + \xi & 0 \\ 0 & 0 & 0 & -(\varepsilon + 2\eta + 2\xi) \end{bmatrix}$$

- i) $V_{x_4}^* - S_{x_4}^* \leftrightarrow v_1 + v_2 + v_4 = (u_3 \wedge u_4 + u_2 \wedge u_5, -u_3 \wedge u_5, u_2 \wedge u_3) \in S_{11,4}^*$
- ii) $(S_{x_4}^*)_1 \leftrightarrow d\rho_1(A) = 2(\eta - \xi) \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 + v_4 = (u_2 \wedge u_5, -u_3 \wedge u_5, u_2 \wedge u_3) \in S_{15,3}^*$
- iii) $(S_{x_4}^*)_2 \leftrightarrow d\rho_2(A) = 3\varepsilon + 2\xi \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 + v_4 = (u_3 \wedge u_4, 0, u_2 \wedge u_3) \in S_{16,2}^*$
- iv) $(S_{x_4}^*)_3 \leftrightarrow d\rho_3(A) = -(\varepsilon + 2\eta + 2\xi) \leftrightarrow f_3^*(y) = y_4 \leftrightarrow v_1 + v_2 = (u_2 \wedge u_5 + u_3 \wedge u_4, -u_3 \wedge u_5, 0) \in S_{13,3}^*$
- v) $-\delta\chi = d\rho_1 + 2d\rho_2 + d\rho_3, \text{tr}_{V_{x_4}^*} = \frac{3}{2}d\rho_1 + 2d\rho_2 + d\rho_3.$

The conormal bundle A_4 is a good holonomic variety with $\text{ord}_{A_4} f^s = -4s - \frac{5}{2}$. We have $b_{A_4}(s)/b_{A_2}(s) = (s+1)(s+\frac{3}{2})$, $b_{A_4}(s)/b_{A_3,13}(s) = (s+1)$ and $b_{A_4}(s)/b_{A_3,15}(s) = (s+\frac{3}{2})$. Note that these intersections are regular and G_0 -prehomogeneous.

(7) The isotropy subalgebra \mathfrak{g}_{x_5} at x_5 is given as follows.

$$(11.6) \quad \mathfrak{g}_{x_5} = \left\{ A = \begin{bmatrix} \varepsilon & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ 0 & 2\varepsilon + 4\eta & \gamma_5 & 0 & \gamma_6 \\ 0 & 0 & \eta & 0 & \gamma_3 + \gamma_5 \\ 0 & 0 & -\gamma_1 & -(\varepsilon + 3\eta) & -\gamma_2 \\ 0 & 0 & 0 & 0 & -2(\varepsilon + \eta) \end{bmatrix} \right\} \\ \oplus \left(\begin{bmatrix} -(3\varepsilon + 4\eta) & 0 & 0 \\ \gamma_3 - \gamma_5 & -(\varepsilon + \eta) & 0 \\ -\gamma_6 & -\gamma_5 & \varepsilon + 2\eta \end{bmatrix} \right) \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(6)).$$

Then $V_{x_5}^*$ is spanned by $v_1 = (u_1 \wedge u_5 - u_3 \wedge u_4, u_4 \wedge u_5, 0)$, $v_2 = (u_2 \wedge u_3, -u_2 \wedge u_5, u_3 \wedge u_5)$, $v_3 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_4 = (u_4 \wedge u_5, 0, 0)$, $v_5 = (u_3 \wedge u_5, 0, 0)$, and

$$(11.7) \quad d\rho_{x_5}(A)(v_1, \dots, v_5) = (v_1, \dots, v_5) \begin{bmatrix} 4\varepsilon + 6\eta & 0 & 0 & 0 & 0 \\ 0 & \varepsilon - \eta & 0 & 0 & 0 \\ -\gamma_1 & -\gamma_5 & 3\varepsilon + 2\eta & 0 & 0 \\ -3\gamma_3 & 0 & 0 & 6\varepsilon + 9\eta & 0 \\ -2\gamma_5 & 2\gamma_6 & \gamma_3 - 2\gamma_5 & \gamma_1 & 5(\varepsilon + \eta) \end{bmatrix}$$

- i) $V_{x_5}^* - S_{x_5}^* \leftrightarrow v_1 + v_2 = (u_1 \wedge u_5 - u_3 \wedge u_4 + u_2 \wedge u_3, u_4 \wedge u_5 - u_2 \wedge u_5, u_3 \wedge u_5) \in S_{8,5}^*$
- ii) $(S_{x_5}^*)_1 \leftrightarrow d\rho_1(A) = 4\varepsilon + 6\eta \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 + v_4 \in S_{11,4}^*$
- iii) $(S_{x_5}^*)_2 \leftrightarrow d\rho_2(A) = \varepsilon - \eta \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 \in S_{13,3}^*$
- iv) $-\delta\chi = 3d\rho_1 + 3d\rho_2 (= 15\varepsilon + 15\eta), \text{tr}_{V_{x_5}^*} = 4d\rho_1 + 3d\rho_2 (= 19\varepsilon + 21\eta)$.

The conormal bundle A_5 is a good holonomic variety with $\text{ord}_{A_5} f^s = -6s - \frac{9}{2}$. We have $b_{A_5}(s)/b_{A_4}(s) = (s + \frac{4}{3})(s + \frac{5}{3})$ and $b_{A_5}(s)/b_{A_{3,13}}(s) = (s + 1)(s + \frac{4}{3})(s + \frac{5}{3})$. Note that the intersection exponent of A_5 and A_4 is $(2:1)$. The intersection of A_5 and $A_{3,13}$ is regular and G_0 -prehomogeneous.

(8) The isotropy subalgebra \mathfrak{g}_{x_6} at x_6 is given as follows.

$$(11.8) \quad \mathfrak{g}_{x_6} = \left\{ \tilde{A} = \left(\begin{array}{c|cc} -4\varepsilon & \gamma_1 & \gamma_2 \\ \hline 0 & \varepsilon I_2 + A & 0 \\ \hline 0 & 0 & \varepsilon I_2 + B \end{array} \right) \oplus \left(\begin{array}{c|c} 3\varepsilon I_2 - {}^t A & 0 \\ \hline \gamma_2 - \gamma_1 & -2\varepsilon \end{array} \right); A, B \in \mathfrak{sl}(2) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus V(2)).$$

Then $V_{x_6}^*$ is spanned by $v_1 = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$, $v_2 = (0, u_2 \wedge u_4, 0)$, $v_3 = (u_3 \wedge u_4, 0, 0)$, $v_4 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_5 = (0, u_2 \wedge u_5, 0)$, $v_6 = (u_3 \wedge u_5, 0, 0)$, and $(G_{x_6}, \rho_{x_6}, V_{x_6}^*) \cong (GL(1) \times SL(2) \times SL(2), 5A_1 \otimes 2A_1 \otimes A_1, V(1) \otimes V(3) \otimes V(2)) \cong (SO(3) \times GL(2), A_1 \otimes A_1, V(3) \otimes V(2))$ and hence A_6 is a good holonomic variety.

- i) $V_{x_6}^* - S_{x_6}^* \leftrightarrow v_2 + v_6 = (u_3 \wedge u_5, u_2 \wedge u_4, 0) \in S_{12,6}^*$
- ii) $(S_{x_6}^*)_1 \leftrightarrow v_2 + v_4 = (u_2 \wedge u_5, u_2 \wedge u_4 - u_3 \wedge u_5, 0) \in S_{13,3}^*$
- iii) $-\delta\chi = d\rho_1, \text{tr}_{V_{x_6}^*} = \frac{3}{2}d\rho_1$

We have $\text{ord}_{A_6} f^s = -4s - \frac{6}{2}$ and $b_{A_6}(s)/b_{A_{3,13}}(s) = (s + \frac{3}{2})$.

(9) The isotropy subalgebra \mathfrak{g}_{x_7} at x_7 is given as follows.

$$(11.9) \quad \mathfrak{g}_{x_7} = \left\{ \tilde{A} = \left(\begin{array}{c|cc} -2(\varepsilon + \eta) & F & 0 \\ \hline 0 & \varepsilon I_2 + A & 0 \\ \hline 0 & 0 & \eta I_2 + B \end{array} \right) \oplus \left(\begin{array}{c|c} (\varepsilon + 2\eta)I_2 - {}^t A & 0 \\ \hline 0 & -2\eta \end{array} \right); \right. \\ \left. A, B \in \mathfrak{sl}(2), {}^t F \in \mathcal{C}^2 \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus V(2)).$$

Then $V_{x_7}^*$ is spanned by $v_1 = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$, $v_2 = (0, u_2 \wedge u_4, 0)$, $v_3 = (u_3 \wedge u_4, 0, 0)$, $v_4 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_5 = (0, u_2 \wedge u_5, 0)$, $v_6 = (u_3 \wedge u_5, 0, 0)$, $v_7 = (0, 0, u_2 \wedge u_3)$, and $(G_{x_7}, \rho_{x_7}, V_{x_7}^*) \cong (GL(1) \times GL(1) \times SL(2) \times SL(2), (2A_1^* \otimes 3A_1^* \otimes 2A_1 \otimes A_1) \oplus (2A_1^* \otimes 2A_1 \otimes 1 \otimes 1), V(6) \oplus V(1))$.

- i) $V_{x_7}^* - S_{x_7}^* \leftrightarrow v_2 + v_6 + v_7 = (u_3 \wedge u_5, u_2 \wedge u_4, u_2 \wedge u_3) \in S_{9,7}^*$
- ii) $(S_{x_7}^*)_1 \leftrightarrow d\rho_1(\tilde{A}) = -8\varepsilon - 12\eta \leftrightarrow f_1^*(y) (\deg f_1^* = 4) \leftrightarrow v_2 + v_4 + v_7 \in S_{11,4}^*$

iii) $(S_{x_7}^*)_2 \leftrightarrow d\rho_2(\tilde{A}) = -2\varepsilon + 2\eta \leftrightarrow f_2^*(y) = y_7 \leftrightarrow v_2 + v_6 \in S_{12,6}^*$

iv) $-\delta\chi = d\rho_1 + d\rho_2, \text{tr}_{V_{x_7}^*} = \frac{3}{2}d\rho_1 + d\rho_2$.

Then by Corollary 1-7 conormal bundle $A_{7,9}$ is a good holonomic variety with $\text{ord}_{A_{7,9}} f^s = -5s - \frac{7}{2}$. We have $\dim A_{7,9} \cap A_4 = \dim A_{7,9} \cap A_6 = \dim V - 1$, $b_{A_{7,9}}(s)/b_{A_4}(s) = (s + \frac{3}{2})$, and $b_{A_{7,9}}(s)/b_{A_6}(s) = (s + 1)$.

(10) The isotropy subalgebra $\mathfrak{g}_{x_7'}$ at x_7' is given as follows.

$$(11.10) \quad \mathfrak{g}_{x_7'} = \left\{ A = \left(\begin{array}{cc|ccc} 3\varepsilon + \alpha & \beta & \gamma_1 & \gamma_2 & \gamma_3 \\ \gamma & 3\varepsilon - \alpha & \gamma_4 & \gamma_5 & \gamma_6 \\ \hline & \mathbf{0} & -2\varepsilon & -2\gamma & -2\beta \\ & & -\beta & -2\varepsilon + 2\alpha & 0 \\ & & -\gamma & 0 & -2\varepsilon - 2\alpha \end{array} \right) \oplus \left(\begin{array}{c|cc} -6\varepsilon & & 0 \\ \hline \gamma_2 - \gamma_4 & -\varepsilon - \alpha & \gamma \\ \gamma_1 - \gamma_6 & \beta & -\varepsilon + \alpha \end{array} \right) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus V(6).$$

Then $V_{x_7'}^*$ is spanned by $v_1 = (u_1 \wedge u_4, 0, u_3 \wedge u_4)$, $v_2 = (u_1 \wedge u_3 - u_2 \wedge u_4, -u_3 \wedge u_4, -2u_4 \wedge u_5)$, $v_3 = (u_1 \wedge u_5 - u_2 \wedge u_3, 2u_4 \wedge u_5, -u_3 \wedge u_5)$, $v_4 = (-u_2 \wedge u_5, u_3 \wedge u_5, 0)$, $v_5 = (u_4 \wedge u_5, 0, 0)$, $v_6 = (u_3 \wedge u_4, 0, 0)$, $v_7 = (u_3 \wedge u_5, 0, 0)$ and the action $d\rho_{x_7'}$ of $\mathfrak{g}_{x_7'}$ on $V_{x_7'}^*$ is given by

$$d\rho_{x_7'}(A)(v_1, \dots, v_7) = (v_1, \dots, v_7) \left(\begin{array}{c|ccc} 5\varepsilon I_4 + A_1 & & & 0 \\ \hline C & & & 10\varepsilon I_3 + A_2 \end{array} \right)$$

where

$$(C, A_2) = \begin{pmatrix} \gamma_3 & 2\gamma_1 - \gamma_6 & 3\gamma_2 - 2\gamma_4 & -\gamma_5 & 0 & -2\beta & 2\gamma \\ \gamma_6 - 2\gamma_1 & 2\gamma_2 & \gamma_5 & 0 & -\gamma & -2\alpha & 0 \\ 0 & \gamma_3 & 2\gamma_6 & \gamma_2 - 2\gamma_4 & \beta & 0 & 2\alpha \end{pmatrix}$$

and

$$A_1 = \begin{pmatrix} -3\alpha & 3\gamma & 0 & 0 \\ \beta & -\alpha & -2\gamma & 0 \\ 0 & -2\beta & \alpha & \gamma \\ 0 & 0 & 3\beta & 3\alpha \end{pmatrix}$$

- i) $V_{x_7'}^* - S_{x_7'}^* \leftrightarrow v_1 + v_4 = (u_1 \wedge u_4 - u_2 \wedge u_5, u_3 \wedge u_5, u_3 \wedge u_4) \in S_{7,7}^{(2)*}$
- ii) $(S_{x_7'}^*)_1 \leftrightarrow d\rho_1(A) = 20\varepsilon \leftrightarrow f_1^*(y_1, \dots, y_4)$: the discriminant of binary cubic forms $\leftrightarrow v_2 \in S_{8,5}^*$
- iii) $-\delta\chi = 2d\rho_1, \text{tr}_{V_{x_7'}^*} = \frac{5}{2}d\rho_1$.

The conormal bundle $A_{7,7}^{(1)}$ is a good holonomic variety with $\text{ord}_{V_{7,7}^{(1)}} f^s = -8s - \frac{13}{2}$. We have $\dim A_5 \cap A_{7,7}^{(1)} = \dim V - 1$ and $b_{A_{7,7}^{(1)}}(s)/b_{A_5}(s) = (s + \frac{5}{4})(s + \frac{7}{4})$. The intersection is regular and G_0 -prehomogeneous.

(11) The isotropy subalgebra $\mathfrak{g}_{x_7'}$ at x_7' is given as follows.

$$(11.11) \quad \mathfrak{g}_{x_7'} = \left\{ \tilde{A} = \left(\begin{array}{c|c|c} -2(\varepsilon+\eta) & 0 & C \\ \hline 0 & \varepsilon I_2 + A & D \\ \hline 0 & 0 & \eta I_2 - {}^t A \end{array} \right) \oplus \left(\begin{array}{c|c} (\varepsilon+2\eta)I_2 - {}^t A & 0 \\ \hline C & -(\varepsilon+\eta) \end{array} \right); \right. \\ \left. A \in \mathfrak{sl}(2) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(5).$$

Then $V_{x_7'}^*$ is spanned by $v_1 = (u_3 \wedge u_4, 0, 0)$, $v_2 = (u_2 \wedge u_4 - u_3 \wedge u_5, -u_3 \wedge u_4, 0)$, $v_3 = (u_2 \wedge u_5, u_2 \wedge u_4 - u_3 \wedge u_5, 0)$, $v_4 = (0, u_2 \wedge u_5, 0)$, $v_5 = (u_1 \wedge u_5, -u_1 \wedge u_4, u_4 \wedge u_5)$, $v_6 = (u_4 \wedge u_5, 0, 0)$, $v_7 = (0, u_4 \wedge u_5, 0)$, and

$$d\rho_{x_7'}(\tilde{A})(v_1, \dots, v_7) = (v_1, \dots, v_7) \left(\begin{array}{c|c|c} -(2\varepsilon+3\eta)I_4 + 3A_1(A) & 0 & 0 \\ \hline 0 & \varepsilon - \eta & 0 \\ \hline * & * & -(\varepsilon+4\eta)I_2 + A \end{array} \right).$$

- i) $V_{x_7'}^* - S_{x_7'}^* \leftrightarrow v_1 + v_4 + v_5 \in S_{7,7}^{(1)*}$
- ii) $(S_{x_7'}^*)_1 \leftrightarrow d\rho_1(\tilde{A}) = -8\varepsilon - 12\eta \leftrightarrow f_1^*(y_1, \dots, y_4)$: the discriminant of binary cubic forms $\leftrightarrow v_3 + v_4 + v_5 \in S_{8,6}^*$
- iii) $(S_{x_7'}^*)_2 \leftrightarrow d\rho_2(\tilde{A}) = \varepsilon - \eta \leftrightarrow f_2^*(y) = y_5 \leftrightarrow v_1 + v_4 \in S_{12,6}^*$
- iv) $-\delta\chi = d\rho_1 + 3d\rho_2, \text{tr}_{V_{x_7'}} = \frac{3}{2}d\rho_1 + 3d\rho_2$.

The conormal bundle $A_{7,7}^{(2)}$ is a good holonomic variety with $\text{ord}_{A_{7,7}^{(2)}} f^s = -7s - \frac{11}{2}$. We have $\dim A_5 \cap A_{7,7}^{(2)} = \dim A_6 \cap A_{7,7}^{(2)} = \dim V - 1$, $b_{A_{7,7}^{(2)}}(s)/b_{A_5}(s) = (s + \frac{3}{2})$ and $b_{A_{7,7}^{(2)}}(s)/b_{A_6}(s) = (s+1)(s + \frac{4}{3})(s + \frac{5}{3})$. The intersections are regular and G_0 -prehomogeneous.

(12) We shall calculate the isotropy subalgebra at $\tilde{x}_8 = (u_2 \wedge u_3 + u_1 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_5, 0)$ instead of x_8 .

$$(11.12) \quad \mathfrak{g}_{\tilde{x}_8} = \left\{ \tilde{A} = \left(\begin{array}{c|cc} 3\varepsilon + \alpha & \alpha_{12} & \gamma_1 & \gamma_2 & \gamma_3 \\ \alpha_{21} & 3\varepsilon - \alpha & \gamma_3 & \gamma_1 & \gamma_4 \\ \hline 0 & -2\varepsilon & -2\alpha_{12} & -2\alpha_{21} \\ -\alpha_{21} & -2\varepsilon - 2\alpha & & & \\ -\alpha_{12} & & & & -2\varepsilon + 2\alpha \end{array} \right) \oplus \left(\begin{array}{c|c} -\varepsilon + \alpha & \alpha_{12} & \gamma_5 \\ \alpha_{21} & -\varepsilon - \alpha & \alpha_6 \\ \hline 0 & \eta \end{array} \right) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus V(6).$$

Then $V_{\tilde{x}_8}^*$ is spanned by $v_1 = (0, 0, u_1 \wedge u_2)$, $v_2 = (0, 0, u_1 \wedge u_5)$, $v_3 = (0, 0, u_1 \wedge u_3 - u_2 \wedge u_5)$, $v_4 = (0, 0, u_1 \wedge u_4 - u_2 \wedge u_3)$, $v_5 = (0, 0, u_2 \wedge u_4)$, $v_6 = (0, 0, u_4 \wedge u_5)$, $v_7 = (0, 0, u_3 \wedge u_5)$, $v_8 = (0, 0, u_3 \wedge u_4)$ and the action $d\rho_{\tilde{x}_8}$ of $\mathfrak{g}_{\tilde{x}_8}$ on $V_{\tilde{x}_8}^*$ is given by

$$d\rho_{\tilde{x}_8}(\tilde{A})(v_1, \dots, v_8) = (v_1, \dots, v_8) \left(\begin{array}{c|c|c} -6\varepsilon - \eta & 0 & 0 \\ \hline C_1 & -(\varepsilon + \eta)L_4 + A_1 & 0 \\ \hline 0 & C_2 & (4\varepsilon - \eta)L_3 + A_2 \end{array} \right) \begin{matrix} \uparrow 1 \\ \uparrow 4 \\ \uparrow 3 \end{matrix}$$

where

$$(C_2, A_2) = \begin{pmatrix} -\gamma_2 & \gamma_1 & \gamma_3 & \gamma_4 & 0 & 2\alpha_{12} & -2\alpha_{21} \\ -\gamma_1 & 2\gamma_3 & -\gamma_4 & 0 & \alpha_{21} & -2\alpha & 0 \\ 0 & \gamma_2 & -2\gamma_1 & -\gamma_3 & -\alpha_{12} & 0 & 2\alpha \end{pmatrix}$$

and

$$(C_1, A_1) = \begin{pmatrix} -\gamma_4 & -3\alpha & 3\alpha_{21} & 0 & 0 \\ -\gamma_3 & \alpha_{12} & -\alpha & 2\alpha_{21} & 0 \\ -\gamma_1 & 0 & 2\alpha_{12} & \alpha & -\alpha_{21} \\ \gamma_2 & 0 & 0 & -3\alpha_{12} & 3\alpha \end{pmatrix}.$$

- i) $V_{\tilde{x}_8}^* - S_{\tilde{x}_8}^* \leftrightarrow v_1 + v_6 = (0, 0, u_1 \wedge u_2 + u_4 \wedge u_5) \in S_{18,8}^*$
- ii) $(S_{\tilde{x}_8}^*)_1 \leftrightarrow d\rho_1(\tilde{A}) = -4\varepsilon - 4\eta \leftrightarrow f_1^*(y_2, \dots, y_5)$: the discriminant of binary cubic forms $\leftrightarrow v_1 + v_7 = (0, 0, u_1 \wedge u_2 + u_3 \wedge u_5) \in S_{18,8}^*$
- iii) $(S_{\tilde{x}_8}^*)_2 \leftrightarrow d\rho_2(\tilde{A}) = -6\varepsilon - \eta \leftrightarrow f_2^*(y) = y_1 \leftrightarrow v_2 + v_5 = (0, 0, u_1 \wedge u_3 + u_2 \wedge u_4) \in S_{18,8}^*$
- iv) $-\delta\chi = 10\varepsilon - 5\eta = -3d\rho_1 + 2d\rho_2$, $\text{tr}_{V_{\tilde{x}_8}^*} = 2\varepsilon - 8\eta = 2d\rho_1 + \frac{5}{2}d\rho_2$.

The conormal bundle $A_{8,18}$ is a good holonomic variety with $\text{ord}_{A_{8,18}} f^s = -5s - \frac{s}{2}$. The conormal vector space $(G_{x_8}, \rho_{x_8}, V_{x_8}^*)$ is a regular P.V. In fact, for $z = \sum_{i=1}^8 z_i v_i \in V_{\tilde{x}_8}$, the localization $f_{\tilde{x}_8}(z)$ of $f(x)$ is given by $f_{\tilde{x}_8}(z) = z_1 z_6^4 + z_1 z_7^2 z_8^2 + 2z_1 z_6^2 z_7 z_8 + z_2^3 z_8^3 + z_2^2 z_6^2 z_8 + 2z_2 z_3 z_6 z_8^2 + z_2 z_5 z_6^3 + z_2 z_4 z_6^2 z_8 + z_3 z_4 z_6^3 + 3z_2 z_5 z_6 z_7 z_8 + z_3 z_5 z_6^2 z_7 - z_2 z_4 z_7 z_8^2 - z_3 z_4 z_6 z_7 z_8 - z_3 z_5 z_7^2 z_8 - z_5^2 z_7^2 - z_4^2 z_6^2 z_7 - 2z_4 z_5 z_6 z_7^2$, and hence its Hessian is not identically zero. Now we shall show that $\dim A_{7,7}^{(1)} \cap A_{8,18} = \dim V - 1$. From iii) above, $A = \overline{G(\tilde{x}_8, v_2 + v_5)}$ is one-codimensional and $A \subset A_{8,18}$. It is enough to show $(\tilde{x}_8, v_2 + v_5) = \{(u_2 \wedge u_3 + u_1 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_5, 0), (0, 0, u_1 \wedge u_5 + u_2 \wedge u_4)\} \in A_{7,7}^{(1)}$. Put $z = \{(u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_2 \wedge u_3 + u_1 \wedge u_5), (u_1 \wedge u_4 - u_2 \wedge u_5, u_3 \wedge u_5, u_3 \wedge u_4)\}$. Then $z \in A_{7,7}^{(1)}$ (See (10)). Then for $\varepsilon > 0$, put

$$g_\varepsilon = \left(\begin{array}{c|c|c} -\varepsilon^3 & & \\ \hline & \varepsilon^3 & \\ \hline & & \varepsilon^{-2} \\ \hline & & & -\varepsilon^{-2} \\ \hline & & & & -\varepsilon^{-2} \end{array} \right) \times \left(\begin{array}{ccc} & & \varepsilon^{-1} \\ & -\varepsilon^{-1} & \\ \varepsilon^{-1} & & \end{array} \right) \in G = SL(5) \times GL(3).$$

Then $g_\varepsilon \cdot z = \{(u_2 \wedge u_3 + u_1 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_5, -\varepsilon^5 u_1 \wedge u_2), (-\varepsilon^5 u_3 \wedge u_5, \varepsilon^5 u_3 \wedge u_4, u_1 \wedge u_5 + u_2 \wedge u_4)\}$. Since $g_\varepsilon \cdot z \in A_{7,7}^{(1)}$ and $A_{7,7}^{(1)}$ is closed, we have $(\tilde{x}_8, v_2 + v_5) = \lim_{\varepsilon \rightarrow 0} g_\varepsilon \cdot z \in A_{7,7}^{(1)}$ and hence $\dim A_{7,7}^{(1)} \cap A_{8,18} = \dim V - 1$. One can see easily that their intersection is regular and G_0 -prehomogeneous. We have $b_{A_{7,7}^{(1)}}(s)/b_{A_{8,18}}(s) = (s+1)(s+\frac{4}{3})(s+\frac{5}{3})$. Next we shall show that $\dim A_{8,18} \cap A_{3,13} = \dim V - 1$. From ii) above, $A = \overline{G(\tilde{x}_8, v_1 + v_7)}$ is one-codimensional and $A \subset A_{8,18}$, where $(\tilde{x}_8, v_1 + v_7) = \{(u_2 \wedge u_3 + u_1 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_5, 0), (0, 0, u_1 \wedge u_2 + u_3 \wedge u_5)\}$. Put $w = \{(u_1 \wedge u_2, u_1 \wedge u_3 + u_2 \wedge u_4, u_2 \wedge u_3 + u_4 \wedge u_5), (u_2 \wedge u_5 + u_3 \wedge u_4, -u_3 \wedge u_5, 0)\}$. Then $w \in A_{3,13}$ (See (5)). For $\varepsilon > 0$, put

$$g_\varepsilon = \left(\begin{array}{ccc|ccc} & & & -\sqrt{-1}\varepsilon^{-4} & & \\ & & & & & \varepsilon \\ & & & & & \\ & & -\sqrt{-1}\varepsilon & & & \\ \varepsilon^6 & & & & & \\ & & & & & \varepsilon^{-4} \end{array} \right) \times \begin{pmatrix} 0 & -\sqrt{-1}\varepsilon^{-2} & 0 \\ 0 & 0 & \varepsilon^3 \\ \sqrt{-1}\varepsilon^3 & 0 & 0 \end{pmatrix}$$

$\in G = SL(5) \times GL(3)$.

Then $g_\varepsilon \cdot w = \{(u_2 \wedge u_3 + u_1 \wedge u_4, u_1 \wedge u_3 + u_2 \wedge u_5, -\varepsilon^{10} u_3 \wedge u_4), (\varepsilon^{10} u_3 \wedge u_5, 0, u_1 \wedge u_2 + u_3 \wedge u_5)\}$. Since $g_\varepsilon \cdot w \in A_{3,13}$ and $A_{3,13}$ is closed, we have $(\tilde{x}_8, v_1 + v_7) = \lim_{\varepsilon \rightarrow 0} g_\varepsilon \cdot w \in A_{3,13}$, i.e., $A \subset A_{3,13} \cap A_{8,18}$. Hence we have $\dim A_{3,13} \cap A_{8,18} = \dim V - 1$. The intersection is G_0 -prehomogeneous and regular, and hence we have $b_{A_{8,18}}(s)/b_{A_{3,13}}(s) = (s+\frac{5}{4})(s+\frac{7}{4})$.

(13) The isotropy subalgebra \mathfrak{g}_{x_8} at x_8 is given as follows.

$$(11.13) \quad \mathfrak{g}_{x_8} = \left\{ A = \left(\begin{array}{cc|cc|c} \varepsilon + \alpha + \beta & \alpha_{12} & & \beta_{12} & \gamma_1 \\ \alpha_{21} & \varepsilon - \alpha + \beta & & -\beta_{12} & \gamma_2 \\ \hline & \beta_{21} & \varepsilon - \alpha - \beta & & \gamma_3 \\ -\beta_{21} & & \varepsilon + \alpha - \beta & & \gamma_4 \\ \hline & & 0 & & -4\varepsilon \end{array} \right) \right. \\ \left. \oplus \left(\begin{array}{cc|c} -2\varepsilon - 2\beta & & -\beta_{21} \\ & -2\varepsilon + 2\beta & \beta_{12} \\ 2\beta_{12} & & -2\varepsilon \end{array} \right) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus V(4)).$$

Then $V_{x_8}^*$ is spanned by $v_1 = (u_3 \wedge u_5, 0, 0)$, $v_2 = (u_2 \wedge u_5, 0, -u_3 \wedge u_5)$, $v_3 = (0, u_3 \wedge u_5, -u_2 \wedge u_5)$, $v_4 = (0, u_2 \wedge u_5, 0)$, $v_5 = (u_4 \wedge u_5, 0, 0)$, $v_6 = (u_1 \wedge u_5, 0, u_4 \wedge u_5)$, $v_7 = (0, u_4 \wedge u_5, u_1 \wedge u_5)$, $v_8 = (0, u_1 \wedge u_5, 0)$. We have $(G_{x_8}, \rho_{x_8}, V_{x_8}^*) \cong (GL(1) \times SL(2) \times SL(2), 5A_1 \otimes 3A_1 \otimes A_1, V(1) \otimes V(4) \otimes V(2))$. Since $\dim \rho_{x_8}(G_{x_8}) = 7 < \dim V_{x_8}^* = 8$, this is not a P.V. The dual of the orbit

$S_{8,14}$ is $S_{14,8}^*$ in V^* , i.e., $A_{8,14} = A_{14,8}^*$.

(14) The isotropy subalgebra $\mathfrak{g}_{x_8''}$ at x_8'' is given as follows.

$$(11.14) \quad \mathfrak{g}_{x_8''} = \left\{ A = \begin{pmatrix} \varepsilon & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ 0 & \eta & \gamma_5 & \gamma_6 & \gamma_7 \\ 0 & 0 & -2(\eta + \xi) & 0 & \gamma_8 \\ 0 & 0 & 0 & \xi & -\gamma_1 \\ 0 & 0 & 0 & 0 & -\varepsilon + \eta + \xi \end{pmatrix} \right\}$$

$$\oplus \left(\begin{pmatrix} -(\varepsilon + \eta) & 0 & 0 \\ -\gamma_5 & -\varepsilon + 2\eta + 2\xi & 0 \\ \gamma_3 - \gamma_7 & -\gamma_8 & -\eta - \xi \end{pmatrix} \right)$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{u}(8)).$$

Then $V_{x_8''}^*$ is spanned by $v_1 = (u_1 \wedge u_5 - u_2 \wedge u_4, 2u_3 \wedge u_4, u_4 \wedge u_5)$, $v_2 = (0, u_2 \wedge u_5, 0)$, $v_3 = (u_2 \wedge u_3, 0, u_3 \wedge u_5)$, $v_4 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_5 = (0, u_4 \wedge u_5, 0)$, $v_6 = (u_3 \wedge u_4, 0, 0)$, $v_7 = (u_4 \wedge u_5, 0, 0)$, $v_8 = (u_3 \wedge u_5, 0, 0)$.

The action $d\rho_{x_8''}$ of $\mathfrak{g}_{x_8''}$ on $V_{x_8''}^*$ is given by

$$d\rho_{x_8''}(A)(v_1, \dots, v_8) = (v_1, \dots, v_8) \left(\begin{array}{cccc|cccc} A_1 & & & & & & & & 0 \\ & A_2 & & & & & & & \\ & & A_3 & & & & & & \\ -2\gamma_1 & \gamma_5 & -\gamma_8 & A_4 & & & & & \\ \hline & 3\gamma_8 & -\gamma_6 & & A_5 & & & & \\ & 3\gamma_5 & & \gamma_6 & & A_6 & & & \\ -2\gamma_3 & & & & -\gamma_6 & \gamma_5 & \gamma_8 & A_7 & \\ -\gamma_2 & 2\gamma_7 - \gamma_3 & -2\gamma_5 & & & \gamma_1 & & A_8 & \end{array} \right)$$

where $A_1 = \varepsilon - \xi$, $A_2 = 2\varepsilon - 4\eta - 3\xi$, $A_3 = \varepsilon + 2\eta + 2\xi$, $A_4 = 2\varepsilon - \eta - \xi$, $A_5 = 2\varepsilon - 3\eta - 4\xi$, $A_6 = \varepsilon + 3\eta + \xi$, $A_7 = 2\varepsilon - 2\xi$ and $A_8 = 2\varepsilon + 2\eta + \xi$.

- i) $V_{x_8''}^* - S_{x_8''}^* \leftrightarrow v_1 + v_2 + v_3 \in S_{5,8}^*$
- ii) $(S_{x_8''}^*)_1 \leftrightarrow d\rho_1(A) = \varepsilon - \xi \leftrightarrow f_1^*(y) = y_1(y = \sum y_i v_i) \leftrightarrow v_2 + v_3 + v_5 \in S_{9,7}^*$
- iii) $(S_{x_8''}^*)_2 \leftrightarrow d\rho_2(A) = 2\varepsilon - 4\eta - 3\xi \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 + v_3 \in S_{7,7}^{(2)*}$
- iv) $(S_{x_8''}^*)_3 \leftrightarrow d\rho_3(A) = \varepsilon + 2\eta + 2\xi \leftrightarrow f_3^*(y) = y_3 \leftrightarrow v_1 + v_2 \in S_{7,7}^{(1)*}$
- v) $-\delta\chi = 6d\rho_1 + d\rho_2 + 2d\rho_3, \text{tr}_{V_{x_8''}^*} = \frac{1}{2}5d\rho_1 + \frac{3}{2}d\rho_2 + \frac{5}{2}d\rho_3.$

Since the intersection of $A_{8,5}^{(3)}$ and $A_{7,7}^{(1)}$ is G_0 -prehomogeneous, the conormal bundle $A_{8,5}^{(3)}$ is a good holonomic variety by Proposition 1-5. The order is given by $\text{ord}_{A_{8,5}^{(3)}} f^s = -9s - \frac{1}{2}5$. We have $\dim A_{8,5}^{(3)} \cap A_{7,9} = \dim A_{8,5}^{(3)} \cap A_{7,7}^{(1)} = \dim V - 1$ ($i = 1, 2$), $b_{A_{8,5}^{(3)}}(s)/b_{A_{7,7}^{(1)}}(s) = s + \frac{3}{2}$ and $b_{A_{8,5}^{(3)}}(s)/b_{A_{7,7}^{(2)}}(s) =$

$(s + \frac{5}{4})(s + \frac{7}{4})$. In (15), we shall prove that $\dim A_{8,5}^{(3)} \cap A_{7,9}^{(2)} \cap A_{5,8}^{(2)} = \dim A_{8,5}^{(3)} \cap A_{9,7}^{(4)} \cap A_{6,8}^{(2)} = \dim V - 1$.

(15) The isotropy subalgebra \mathfrak{g}_{x_9} at x_9 is given as follows.

$$(11.15) \quad \mathfrak{g}_{x_9} = \left\{ A = \begin{pmatrix} \varepsilon_1 & 0 & \gamma_1 & \gamma_2 & & \gamma_3 \\ 0 & \varepsilon_2 & \gamma_4 & \gamma_5 & & \gamma_6 \\ 0 & 0 & \varepsilon_3 & 0 & & \gamma_7 \\ 0 & 0 & 0 & \varepsilon_4 & & \gamma_8 \\ 0 & 0 & 0 & 0 & -(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4) & \end{pmatrix} \right. \\ \left. \oplus \begin{pmatrix} -\varepsilon_1 - \varepsilon_2 & 0 & 0 \\ -\gamma_4 & -\varepsilon_1 - \varepsilon_3 & 0 \\ \gamma_2 & 0 & -\varepsilon_2 - \varepsilon_4 \end{pmatrix} \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1)) \oplus \mathfrak{u}(8).$$

Then $V_{x_9}^*$ is spanned by $v_1 = (0, u_2 \wedge u_5, 0)$, $v_2 = (0, 0, u_1 \wedge u_5)$, $v_3 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_4 = (u_1 \wedge u_5, 0, u_4 \wedge u_5)$, $v_5 = (0, u_4 \wedge u_5, 0)$, $v_6 = (0, 0, u_3 \wedge u_5)$, $v_7 = (u_3 \wedge u_4, 0, 0)$, $v_8 = (u_3 \wedge u_5, 0, 0)$, $v_9 = (u_4 \wedge u_5, 0, 0)$.

The action $d\rho_{x_9}$ of \mathfrak{g}_{x_9} on $V_{x_9}^*$ is given by

$$d\rho_{x_9}(A)(v_1, \dots, v_9) = (v_1, \dots, v_9) \begin{pmatrix} A_1 & & & & & & & & \\ & A_2 & & & & & & & \\ & \gamma_4 & A_3 & & & & & & \\ & & -\gamma_2 & A_4 & & & & & \\ & -\gamma_5 & & & A_5 & & & & \\ & & -\gamma_1 & & & A_6 & & & \\ & & & & & & A_7 & & \\ & & & & & & & A_8 & \\ & & & & -2\gamma_4 & -\gamma_1 & -\gamma_2 & -\gamma_8 & \\ & & & & -\gamma_5 & -2\gamma_2 & \gamma_4 & \gamma_7 & A_9 \end{pmatrix}$$

where $A_1 = 2\varepsilon_1 + 2\varepsilon_3 + \varepsilon_4$, $A_2 = 2\varepsilon_2 + \varepsilon_3 + 2\varepsilon_4$, $A_3 = 2\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4$, $A_4 = \varepsilon_1 + 2\varepsilon_2 + \varepsilon_3 + \varepsilon_4$, $A_5 = 2\varepsilon_1 + \varepsilon_2 + 2\varepsilon_3$, $A_6 = \varepsilon_1 + 2\varepsilon_2 + 2\varepsilon_4$, $A_7 = \varepsilon_1 + \varepsilon_2 - \varepsilon_3 - \varepsilon_4$, $A_8 = 2\varepsilon_1 + 2\varepsilon_2 + \varepsilon_4$ and $A_9 = 2\varepsilon_1 + 2\varepsilon_2 + \varepsilon_3$.

- i) $V_{x_9}^* - S_{x_9}^* \leftrightarrow v_1 + v_2 + v_7 = (u_3 \wedge u_4, u_2 \wedge u_5, u_1 \wedge u_5) \in S_{7,9}^*$
- ii) $(S_{x_9}^*)_1 \leftrightarrow d\rho_1(A) = 2\varepsilon_1 + 2\varepsilon_3 + \varepsilon_4 \leftrightarrow f_1^*(y) = y_1 \leftrightarrow v_2 + v_3 + v_5 + v_7 \in S_{8,5}^*$
- iii) $(S_{x_9}^*)_2 \leftrightarrow d\rho_2(A) = 2\varepsilon_2 + \varepsilon_3 + 2\varepsilon_4 \leftrightarrow f_2^*(y) = y_2 \leftrightarrow v_1 + v_4 + v_6 + v_7 \in S_{8,5}^*$
- iv) $(S_{x_9}^*)_3 \leftrightarrow d\rho_3(A) = \varepsilon_1 + \varepsilon_2 - \varepsilon_3 - \varepsilon_4 \leftrightarrow f_3^*(y) = y_7 \leftrightarrow v_1 + v_2 + v_8 + v_9 \in S_{14,8}^*$
- v) $-\delta\chi = 3d\rho_1 + 3d\rho_2 + 4d\rho_3, \text{tr}_{V_{x_9}^*} = 4d\rho_1 + 4d\rho_2 + 5d\rho_3.$

The conormal bundle $A_{9,7}$ is a good holonomic variety with $\text{ord}_{A_{9,7}} f^s = -10s - \frac{17}{2}$.

Put $p = (u_1 \wedge u_2, u_1 \wedge u_3, u_2 \wedge u_4; u_1 \wedge u_5 + u_3 \wedge u_4, u_2 \wedge u_5, u_3 \wedge u_5 + u_4 \wedge u_5)$. Then by iii), we have $p \in A_{9,7}^{(4)} \cap A_{8,8}^{(2)}$ and $\dim G \cdot p = \dim V - 1$. We shall prove that $p \in A_{8,5}^{(3)}$, i.e., $\dim A_{6,8}^{(2)} \cap A_{8,5}^{(3)} \cap A_{9,7}^{(4)} = \dim V - 1$. By (14), for any $\varepsilon > 0$, we have $(u_1 \wedge u_2, u_1 \wedge u_3, u_1 \wedge u_5 + u_2 \wedge u_4; u_1 \wedge u_5 - u_2 \wedge u_4 + (1/\varepsilon)u_2 \wedge u_3 + (1/\varepsilon^2)u_3 \wedge u_4, 2u_3 \wedge u_4 + u_2 \wedge u_5, u_4 \wedge u_5 + (1/\varepsilon)u_3 \wedge u_5) \in (x'_8, V_{x_8}^*) \subset A_{8,5}^{(3)}$. Therefore, by the action of $g_\varepsilon = \begin{pmatrix} \varepsilon & & & & \\ & 1 & & & \\ & & \varepsilon^{-1} & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} \times \begin{pmatrix} \varepsilon^{-1} & & & \\ & 1 & & \\ & & & 1 \end{pmatrix} \in G = SL(5) \times GL(3)$, we have $p_\varepsilon = (u_1 \wedge u_2, u_1 \wedge u_3, \varepsilon u_1 \wedge u_5 + u_2 \wedge u_4; u_1 \wedge u_5 - \varepsilon u_2 \wedge u_4 + \varepsilon u_2 \wedge u_3 + u_3 \wedge u_4, 2\varepsilon u_3 \wedge u_4 + u_2 \wedge u_5, u_3 \wedge u_5 + u_4 \wedge u_5) \in A_{8,5}^{(3)}$. Hence, we have $p = \lim_{\varepsilon \rightarrow 0} p_\varepsilon \in A_{8,5}^{(3)}$. Since $A_{5,8}^{(2)} = A_{8,5}^{(3)*}$, $A_{8,5}^{(3)} = A_{5,8}^{(2)*}$, $A_{9,7}^{(4)} = A_{7,9}^{(2)*}$ and $(G, \rho, V) \cong (G, \rho^*, V^*)$, we have also $\dim A_{5,8}^{(2)} \cap A_{8,5}^{(3)} \cap A_{7,9}^{(2)} = \dim V - 1$.

(16) The isotropy subalgebra $\mathfrak{g}_{x_{10}}$ at x_{10} is given as follows.

$$(11.16) \quad \mathfrak{g}_{x_{10}} = \left\{ A = \left[\begin{array}{c|cc|cc|c} \eta & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ & -4\varepsilon & 0 & 0 & \gamma_5 \\ \hline 0 & \varepsilon + \alpha & \alpha_{12} & & \gamma_3 \\ & \alpha_{21} & \varepsilon - \alpha & & -\gamma_2 \\ \hline 0 & & 0 & & 2\varepsilon - \eta \end{array} \right] \oplus \left(\begin{array}{c|c|c} 4\varepsilon - \eta & 0 & \gamma_6 \\ \hline -\gamma_5 & -2\varepsilon & \gamma_7 \\ \hline 0 & & \xi \end{array} \right) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(7).$$

Then $V_{x_{10}}^*$ is spanned by $v_1 = (0, 0, u_1 \wedge u_3)$, $v_2 = (0, 0, u_1 \wedge u_4)$, $v_3 = (0, 0, u_1 \wedge u_5 - u_3 \wedge u_4)$, $v_4 = (0, 0, u_2 \wedge u_3)$, $v_5 = (0, 0, u_2 \wedge u_4)$, $v_6 = (u_3 \wedge u_5, 0, 0)$, $v_7 = (u_4 \wedge u_5, 0, 0)$, $v_8 = (0, 0, u_3 \wedge u_5)$, $v_9 = (0, 0, u_4 \wedge u_5)$, $v_{10} = (0, 0, u_2 \wedge u_5)$.

$$V_{x_{10}}^* - S_{x_{10}}^* \leftrightarrow v_1 + v_5 + v_7 = (u_4 \wedge u_5, 0, u_1 \wedge u_3 + u_2 \wedge u_4) = y_{10}.$$

Let A_0 be an element of $\mathfrak{g}_{x_{10}}$ with $\alpha = -\frac{1}{3} - 5\varepsilon$, $\eta = \frac{2}{3} + 6\varepsilon$, $\xi = -\frac{4}{3} - 2\varepsilon$, all remaining parts zero in (5.16). Then $d\rho(A_0)x_{10} = 0$ and $d\rho^*(A_0)y_{10} = y_{10}$. Since $-\delta\chi(A_0) = 10(1 + 3\varepsilon)$ is not definite, the conormal bundle $A_{10,10}$ is *not* a good holonomic variety.

(17) The isotropy subalgebra $\mathfrak{g}_{x_{11}}$ at x_{11} is given as follows.

$$(11.17) \quad \mathfrak{g}_{x_{11}} = \left\{ A = \left[\begin{array}{c|cc|cc|c} \varepsilon + \eta & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ \hline 0 & \varepsilon + \alpha & \alpha_{12} & \gamma_5 & \gamma_6 \\ & \alpha_{21} & \varepsilon - \alpha & \gamma_7 & \gamma_8 \\ \hline 0 & 0 & \varepsilon - \eta & \gamma_9 & \\ & & n_0 & -4\varepsilon & \end{array} \right] \oplus \left(\begin{array}{c|c|c} -2\varepsilon - \eta - \alpha & -\alpha_{21} & 0 \\ \hline -\alpha_{12} & -2\varepsilon - \eta + \alpha & \\ \hline \gamma_2 - \gamma_5 & -\gamma_1 - \gamma_7 & -2\varepsilon \end{array} \right) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(9).$$

The conormal vector space $V_{x_{11}}^*$ is spanned by $v_1 = (0, u_1 \wedge u_5, -u_2 \wedge u_5)$,
 $v_2 = (u_1 \wedge u_5, 0, u_3 \wedge u_5)$, $v_3 = (u_2 \wedge u_5, 0, -u_4 \wedge u_5)$, $v_4 = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$,
 $v_5 = (0, u_2 \wedge u_4, 0)$, $v_6 = (u_3 \wedge u_4, 0, 0)$, $v_7 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_8 = (0,$
 $u_2 \wedge u_5, 0)$, $v_9 = (u_3 \wedge u_5, 0, 0)$, $v_{10} = (0, u_4 \wedge u_5, 0)$, $v_{11} = (u_4 \wedge u_5, 0, 0)$.

The action $d\rho_{x_{11}}$ of $\mathfrak{g}_{x_{11}}$ on $V_{x_{11}}^*$ is given by

$$d\rho_{x_{11}}(A)(v_1, \dots, v_{11}) \left(\begin{array}{cccccc} \underbrace{A_1}_{2} & & & & & \\ \underbrace{B_1}_{2} & \underbrace{A_2}_{1} & & & & \\ & & \underbrace{A_3}_{3} & & & \\ \underbrace{B_2}_{2} & \underbrace{A_6}_{1} & \underbrace{B_3}_{3} & \underbrace{A_4}_{3} & & \\ \underbrace{B_4}_{2} & \underbrace{B_5}_{1} & \underbrace{B_6}_{3} & \underbrace{B_7}_{3} & \underbrace{A_5}_{2} & \end{array} \right) \left. \begin{array}{l} \end{array} \right\} \begin{array}{l} 2 \\ 1 \\ 3 \\ 3 \\ 2 \end{array}$$

where $(B_1, A_2) = (-\gamma_5, \gamma_7, 5\varepsilon + \eta)$, $B_3 = -\gamma_9 \cdot I_3$, $B_4 = -\gamma_3 \cdot I_2$, $A_1 = 5\varepsilon I_2 + A'$,
 $A_5 = (5\varepsilon + 2\eta)I_2 + A'$ with $A' = \begin{pmatrix} -\alpha & \alpha_{21} \\ \alpha_{12} & \alpha \end{pmatrix}$,

$$(B_5, B_6, B_7) = \left(\begin{array}{ccc|ccc} -\gamma_1 - \gamma_7 & -\gamma_8 & \gamma_6 & 0 & -\gamma_7 & -\gamma_5 & 0 \\ \gamma_2 - 2\gamma_5 & \gamma_6 & 0 & \gamma_8 & -\gamma_5 & 0 & -\gamma_7 \end{array} \right),$$

$A_3 = 2\eta \cdot I_3 + A''$, $A_4 = (5\varepsilon + \eta)I_3 + A''$ with $A'' = \begin{pmatrix} 0 & \alpha_{12} & -\alpha_{21} \\ 2\alpha_{21} & -2\alpha & 0 \\ -2\alpha_{12} & 0 & 2\alpha \end{pmatrix}$,

$$(B_2, A_6) = \begin{pmatrix} -\gamma_2 & -\gamma_1 - \gamma_7 & \\ -2\gamma_1 - \gamma_7 & & \alpha_{21} \\ 0 & \gamma_5 - 2\gamma_2 & -\alpha_{12} \end{pmatrix}.$$

Note that A_6 will disappear if we take $v_3 - \frac{1}{2}v_7$ instead of v_3 .

- i) $V_{x_{11}}^* - S_{x_{11}}^* \leftrightarrow v_1 + v_2 + v_4 = (u_1 \wedge u_5 + u_2 \wedge u_4, u_1 \wedge u_5 - u_3 \wedge u_4, u_3 \wedge u_5 - u_2 \wedge u_5) \in S_{4,11}^{(3)*}$
- ii) $(S_{x_{11}}^*)_1 \leftrightarrow v_1 + v_4 + v_9 = (u_2 \wedge u_4 + u_3 \wedge u_5, u_1 \wedge u_5 - u_3 \wedge u_4, -u_2 \wedge u_5) \in S_{5,8}^{(2)*} \leftrightarrow d\rho_1 = 10\varepsilon + 2\eta$
- iii) $(S_{x_{11}}^*)_2 \leftrightarrow v_2 + v_5 = (u_1 \wedge u_5, u_2 \wedge u_4, u_3 \wedge u_5) \in S_{7,9}^{(2)*} \leftrightarrow d\rho_2 = 4\eta \leftrightarrow f_2^*(y) = y_4^2 + y_5 y_6$
- iv) $-\delta\chi = 30\varepsilon + 10\eta = 3d\rho_1 + d\rho_2$, $\text{tr}_{V_{x_{11}}^*} = 40\varepsilon + 14\eta = 4d\rho_1 + \frac{3}{2}d\rho_2$.

Remark for calculation of $d\rho_1$. Let f_1^* be the relative invariant on $V_{x_{11}}^*$ corresponding to $d\rho_1$. Since $f_1^*(v_2 + v_5) \neq 0$, the restriction of $d\rho_1$ to the isotropy subalgebra of $\mathfrak{g}_{x_{11}}$ at $v_2 + v_5$ should be zero. Hence $d\rho_1$ must be of the form $d\rho_1 = 5\lambda\varepsilon + \lambda\eta$ for some λ . Take an element A_0 in $\mathfrak{g}_{x_{11}}$ satisfying $d\rho^*(A_0)x_{11}^* = x_{11}^*$ where $x_{11}^* = v_1 + v_2 + v_4 \in V_{x_{11}}^* - S_{x_{11}}^*$. Then, by the Euler's identity, we have $(\text{deg } f_1^*) \cdot f_1^*(x_{11}^*) = \langle d\rho^*(A_0)y, D_y \rangle f_1^*(y)|_{y=x_{11}^*} =$

$d\rho_1(A_0)f_1^*(x_{11})$ and hence $\deg f_1^* = d\rho_1(A_0) = \frac{3}{2}\lambda \in N$. Therefore $d\rho_1 = (10\varepsilon + 2\eta)\mu$ where μ is a natural number. Since $-\delta\chi$ is a linear combination of $d\rho_1$ and $d\rho_2$ with coefficients in Z , we have $\mu = 1$ or 3 . On the other hand, $2\text{tr}_{V_{x_{11}}}$ is also a linear combination of $d\rho_1$ and $d\rho_2$ with coefficients in Z , μ is a divisor of 8 , and hence $\mu = 1$, i.e., $d\rho_1 = 10\varepsilon + 2\eta$.

(18) The isotropy subalgebra $\mathfrak{g}_{x_{12}}$ at x_{12} is given as follows.

$$(11.18) \quad \mathfrak{g}_{x_{12}} = \left\{ A = \left[\begin{array}{cc|cc|c} \varepsilon + \alpha & \alpha_{12} & & & \gamma_1 \\ \alpha_{21} & \varepsilon - \alpha & & & \gamma_2 \\ \hline & & 0 & & \\ \hline & & \eta + \beta & \beta_{12} & \gamma_3 \\ & & \beta_{21} & \eta - \beta & \gamma_4 \\ \hline & & & & -2(\varepsilon + \eta) \end{array} \right] \oplus \left(\begin{array}{cc|c} -2\varepsilon & 0 & \gamma_5 \\ 0 & -2\eta & \gamma_6 \\ \hline & & \xi \end{array} \right) \right\} \\
 \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus V(6)).$$

The conormal vector space $V_{x_{12}}^*$ is spanned by $v_1 = (0, 0, u_1 \wedge u_5)$, $v_2 = (0, 0, u_2 \wedge u_5)$, $v_3 = (0, 0, u_3 \wedge u_5)$, $v_4 = (0, 0, u_4 \wedge u_5)$, $v_5 = (0, u_1 \wedge u_5, 0)$, $v_6 = (0, u_2 \wedge u_5, 0)$, $v_7 = (u_3 \wedge u_5, 0, 0)$, $v_8 = (u_4 \wedge u_5, 0, 0)$, $v_9 = (0, 0, u_1 \wedge u_3)$, $v_{10} = (0, 0, u_1 \wedge u_4)$, $v_{11} = (0, 0, u_2 \wedge u_3)$, $v_{12} = (0, 0, u_2 \wedge u_4)$.

The action $d\rho_{x_{12}}$ of $\mathfrak{g}_{x_{12}}$ on $V_{x_{12}}^*$ is given by

$$d\rho_{x_{12}}(A)(v_1, \dots, v_{12}) = (v_1, \dots, v_{12}) = \left[\begin{array}{ccc} \underbrace{A_1}_2 & \underbrace{B_1}_2 & \underbrace{B_2}_2 \\ & \underbrace{A_2}_2 & \underbrace{B_3}_2 \underbrace{B_4}_2 \\ & & \underbrace{A_3}_2 \\ & & & \underbrace{A_4}_2 \\ & & & & \underbrace{A_5}_4 \end{array} \right]$$

where $B_1 = -\gamma_5 I_2$, $B_3 = -\gamma_5 I_2$, $A_1 = (\varepsilon + 2\eta - \xi)I_2 + A'$, $A_2 = (2\varepsilon + \eta - \xi)I_2 + B'$, $A_3 = (\varepsilon + 4\eta)I_2 + A'$, $A_4 = (4\varepsilon + \eta)I_2 + B'$ with $A' = \begin{pmatrix} -\alpha & -\alpha_{21} \\ -\alpha_{12} & \alpha \end{pmatrix}$, $B' = \begin{pmatrix} -\beta & -\beta_{21} \\ -\beta_{12} & \beta \end{pmatrix}$, $A_5 = -(\varepsilon + \eta + \xi)I_4 + \begin{pmatrix} -\alpha I_2 + B' & -\alpha_{21} I_2 \\ -\alpha_{12} I_2 & \alpha I_2 + B' \end{pmatrix}$, $B_2 = \begin{pmatrix} -\gamma_3 & -\gamma_4 & 0 & 0 \\ 0 & 0 & -\gamma_3 & -\gamma_4 \end{pmatrix}$ and $B_4 = \begin{pmatrix} \gamma_1 & 0 & \gamma_2 & 0 \\ 0 & \gamma_1 & 0 & \gamma_2 \end{pmatrix}$.

i) $V_{x_{12}}^* - S_{x_{12}}^* \leftrightarrow v_6 + v_7 + v_{10} + v_{11} = (u_3 \wedge u_5, u_2 \wedge u_5, u_1 \wedge u_4 + u_2 \wedge u_3) \in S_{6,12}^{(1)*}$

ii) $(S_{x_{12}}^*)_1 \leftrightarrow v_6 + v_8 + v_9 = (u_4 \wedge u_5, u_2 \wedge u_5, u_1 \wedge u_3) \in S_{7,9}^{(2)*} \leftrightarrow d\rho_1 = -2(\varepsilon + \eta + \xi) \leftrightarrow f_1^*(y) = y_{10}y_{11} - y_9y_{12}$

iii) $(S_{x_{12}}^*)_2 \leftrightarrow v_5 + v_7 + v_{10} + v_{11} = (u_3 \wedge u_5, u_1 \wedge u_5, u_2 \wedge u_3 + u_2 \wedge u_4) \in S_{7,7}^{(2)*} \leftrightarrow d\rho_2 = 4(\varepsilon + \eta) - \xi \leftrightarrow \deg f_2^*(y) = 3$

iv) $-\delta\chi = 10(\varepsilon + \eta) - 5\xi = d\rho_1 + 3d\rho_2$, $\text{tr}_{V_{x_{12}}} = 12(\varepsilon + \eta) - 8\xi = 2d\rho_1 + 4d\rho_2$.

(19) The isotropy subalgebra $\mathfrak{g}_{x_{13}}$ at x_{13} is given as follows.

$$(11.19) \quad \mathfrak{g}_{x_{13}} = \left\{ A = \left(\begin{array}{cc|cc|c} \varepsilon + \alpha & \alpha_{12} & \gamma_1 & \gamma_2 & \gamma_5 \\ \alpha_{21} & \varepsilon - \alpha & \gamma_3 & \gamma_4 & \gamma_6 \\ \hline 0 & & -(\varepsilon + \eta) - \alpha & -\alpha_{21} & \gamma_7 \\ 0 & & -\alpha_{12} & -(\varepsilon + \eta) + \alpha & \gamma_8 \\ \hline 0 & & & 0 & 2\eta \end{array} \right) \right. \\ \left. \oplus \left(\begin{array}{ccc} -2\varepsilon & 0 & \delta_1 \\ \gamma_2 - \gamma_3 & \eta & \delta_2 \\ 0 & 0 & \xi \end{array} \right) \right\} \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2)) \oplus \mathfrak{u}(10).$$

The conormal vector space $V_{x_{13}}^*$ is spanned by $v_1 = (0, 0, -u_4 \wedge u_5)$, $v_2 = (0, 0, u_3 \wedge u_5)$, $v_3 = (0, 0, u_1 \wedge u_5)$, $v_4 = (0, 0, u_2 \wedge u_5)$, $v_5 = (0, 0, u_3 \wedge u_4)$, $v_6 = (0, 0, -u_1 \wedge u_4)$, $v_7 = (0, 0, u_1 \wedge u_3 - u_2 \wedge u_4)$, $v_8 = (0, 0, u_2 \wedge u_3)$, $v_9 = (-u_4 \wedge u_5, 0, 0)$, $v_{10} = (u_3 \wedge u_5, 0, 0)$, $v_{11} = (u_3 \wedge u_4, 0, 0)$, $v_{12} = (u_1 \wedge u_5, u_4 \wedge u_5, 0)$, $v_{13} = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$.

The action $d\rho_{x_{13}}$ of $\mathfrak{g}_{x_{13}}$ on $V_{x_{13}}^*$ is given by

$$d\rho_{x_{13}}(A)(v_1, \dots, v_{13}) = (v_1, \dots, v_{13}) \left(\begin{array}{cccccc} \underbrace{A_1}_2 & \underbrace{B_1}_2 & \underbrace{B_2}_1 & \underbrace{B_3}_3 & \underbrace{B_4}_2 & \underbrace{B_5}_1 \\ & \underbrace{A_2}_2 & & \underbrace{B_6}_3 & & \underbrace{B_7}_1 \\ & & \underbrace{A_3}_1 & \underbrace{B_8}_3 & & \underbrace{B_9}_1 \\ & & & \underbrace{A_4}_3 & & \underbrace{B_{10}}_2 \\ & & & & \underbrace{A_5}_2 & \underbrace{B_{11}}_1 \\ & & & & & \underbrace{A_6}_1 \\ & & & & & \underbrace{A_7}_2 \end{array} \right)^2$$

where $A_3 = 2\varepsilon + 2\eta - \xi$, $B_8 = (\gamma_1, \gamma_2 + \gamma_3, \gamma_4)$, $B_9 = -\delta_1$, $A_6 = 4\varepsilon + 2\eta$, $B_4 = B_7 = -\delta_1 I_2$, $B_5 = \delta_2 I_2$, $A_1 = (\varepsilon - \eta - \xi)I_2 + A'$ with $A' = \begin{pmatrix} -\alpha & -\alpha_{21} \\ -\alpha_{12} & \alpha \end{pmatrix}$, $(B_1, B_2, B_3) = \begin{pmatrix} \gamma_2 & \gamma_4 & -\gamma_7 & \gamma_5 & \gamma_6 & 0 \\ -\gamma_1 & -\gamma_3 & -\gamma_8 & 0 & \gamma_5 & \gamma_6 \end{pmatrix}$, $B_6 = \begin{pmatrix} \gamma_8 & -\gamma_7 \\ & \gamma_8 & -\gamma_7 \end{pmatrix}$, $(B_{10}, B_{11}) = \begin{pmatrix} -\gamma_7 & 2\gamma_2 - \gamma_3 & \gamma_4 \\ -\gamma_8 & -\gamma_1 & \gamma_2 - 2\gamma_3 \end{pmatrix}$, $A_2 = -(\varepsilon + 2\eta + \xi)I_2 + A'$, $A_5 = (3\varepsilon - \eta)I_2 + A'$, $A_7 = (\varepsilon - 2\eta)I_2 + A'$ and $A_4 = (\eta - \xi)I_3 + \begin{pmatrix} -2\alpha & -2\alpha_{21} & 0 \\ -\alpha_{12} & 0 & -\alpha_{21} \\ 0 & -2\alpha_{12} & 2\alpha \end{pmatrix}$.

- i) $V_{x_{13}}^* - S_{x_{13}}^* \leftrightarrow v_7 + v_{11} + v_{12} + v_{13} \in S_{3,13}^{(2)*}$
- ii) $(S_{x_{13}}^*)_1 \leftrightarrow v_7 + v_{12} + v_{13} \in S_{6,12}^{(1)*} \leftrightarrow d\rho_1 = 4\varepsilon + 2\eta \leftrightarrow f_1^*(y) = y_{11}$
- iii) $(S_{x_{13}}^*)_2 \leftrightarrow v_8 + v_{11} + v_{12} \in S_{4,11}^{(3)*} \leftrightarrow d\rho_2 = 2(\eta - \xi) \leftrightarrow f_2^*(y) = y_7^2 - y_6 y_8$
- iv) $(S_{x_{13}}^*)_3 \leftrightarrow v_7 + v_{11} + v_{12} \in S_{5,8}^{(2)*} \leftrightarrow d\rho_3 = 2\varepsilon - 3\eta - \xi \leftrightarrow \deg f_2^*(y) = 3$
- v) $-\delta\chi = d\rho_1 + d\rho_2 + 3d\rho_3$, $\text{tr}_{V_{x_{13}}^*} = \frac{3}{2}d\rho_1 + 2d\rho_2 + 4d\rho_3$.

(20) The isotropy subalgebra $\mathfrak{g}_{x_{14}}$ at x_{14} is given as follows.

$$(11.20) \quad \mathfrak{g}_{x_{14}} = \left\{ A = \left(\begin{array}{c|cc} \varepsilon & * & * \\ \hline 0 & \eta I_3 + X & * \\ \hline 0 & 0 & -\varepsilon - 3\eta \end{array} \right) \oplus (- (\varepsilon + \eta) I_3 - 'X); X \in \mathfrak{sl}(3) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(3)) \oplus \mathfrak{u}(7).$$

The conormal vector space $V_{x_{14}}^*$ is spanned by $v_1 = (u_3 \wedge u_4, 0, 0)$, $v_2 = (0, u_2 \wedge u_4, 0)$, $v_3 = (0, 0, u_2 \wedge u_3)$, $v_4 = (0, u_2 \wedge u_3, -u_2 \wedge u_4)$, $v_5 = (u_2 \wedge u_3, 0, u_3 \wedge u_4)$, $v_6 = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$, $v_7 = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_8 = (0, u_3 \wedge u_5, -u_4 \wedge u_5)$, $v_9 = (0, u_2 \wedge u_5, 0)$, $v_{10} = (0, 0, u_3 \wedge u_5)$, $v_{11} = (u_4 \wedge u_5, 0, 0)$, $v_{12} = (u_3 \wedge u_5, 0, 0)$, $v_{13} = (0, u_4 \wedge u_5, 0)$, $v_{14} = (0, 0, u_2 \wedge u_5)$.

Since $\dim \rho_{x_{14}}(G_{x_{14}}) = 13$ and $\dim V_{x_{14}}^* = 14$, the conormal vector space $(G_{x_{14}}, \rho_{x_{14}}, V_{x_{14}}^*)$ is not a P.V. Note that it is also obtained from the fact that $A_{8,14} = A_{14,8}^*$ is not G -prehomogeneous (See (13)).

(21) The isotropy subalgebra $\mathfrak{g}_{x_{15}}$ at x_{15} is given as follows.

$$(11.21) \quad \mathfrak{g}_{x_{15}} = \left\{ A = \left(\begin{array}{c|c} 2\varepsilon I_3 + X & Z \\ \hline 0 & -3\varepsilon I_2 + Y \end{array} \right) \oplus (-4\varepsilon I_3 + S^{-1}XS); X \in \mathfrak{sl}(3), \right. \\ \left. Y \in \mathfrak{sl}(2), Z \in V(6), S = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{sl}(3) \oplus \mathfrak{sl}(2)) \oplus V(6).$$

The conormal vector space $V_{x_{15}}^*$ is spanned by $v_1 = (0, 0, u_1 \wedge u_4)$, $v_2 = (0, u_2 \wedge u_4, 0)$, $v_3 = (u_3 \wedge u_4, 0, 0)$, $v_4 = (0, u_1 \wedge u_4, -u_2 \wedge u_4)$, $v_5 = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$, $v_6 = (u_1 \wedge u_4, 0, u_3 \wedge u_4)$, $v_7 = (0, 0, u_1 \wedge u_5)$, $v_8 = (0, u_2 \wedge u_5, 0)$, $v_9 = (u_3 \wedge u_5, 0, 0)$, $v_{10} = (0, u_1 \wedge u_5, -u_2 \wedge u_5)$, $v_{11} = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_{12} = (u_1 \wedge u_5, 0, u_3 \wedge u_5)$, $v_{13} = (0, 0, u_4 \wedge u_5)$, $v_{14} = (0, u_4 \wedge u_5, 0)$, $v_{15} = (u_4 \wedge u_5, 0, 0)$. Then the action $d\rho_{x_{15}}$ of $\mathfrak{g}_{x_{15}}$ on $V_{x_{15}}^*$ is given by

$$(v_1, \dots, v_{15}) \mapsto (v_1, \dots, v_{15}) \left(\begin{array}{c|c} GL(1) \times SL(3) \times SL(2) & 0 \\ \hline 5A_1 \otimes 2A_1^* \otimes A_1^* & \\ \hline * & GL(1) \times SL(3) \\ & 10A_1 \otimes A_1^* \end{array} \right)$$

- i) $V_{x_{15}}^* - S_{x_{15}}^* \leftrightarrow v_1 + v_2 + v_7 + v_9 = (u_3 \wedge u_5, u_2 \wedge u_4, u_1 \wedge u_4 + u_1 \wedge u_5) \in S_{3,15}^{(3)*}$
- ii) $(S_{x_{15}}^*)_1 \leftrightarrow v_4 + v_{11} + v_{12} \in S_{4,11}^{(3)*} \leftrightarrow d\rho_1 = 60\varepsilon$
- iii) $-\delta\chi = 60\varepsilon = d\rho_1, \text{tr}_{V_{x_{15}}^*} = 90\varepsilon = \frac{3}{2}d\rho_1.$

(22) The isotropy subalgebra $\mathfrak{g}_{x_{16}}$ at x_{16} is given as follows.

$$(11.22) \quad \mathfrak{g}_{x_{16}} = \left(A = \left[\begin{array}{c|cc|cc} -2(\varepsilon+\eta) & \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ \hline 0 & \varepsilon+\alpha_1 & \alpha_{12} & \gamma_5 & \gamma_6 \\ \hline & \alpha_{21} & \varepsilon-\alpha_1 & \gamma_7 & \gamma_8 \\ \hline 0 & & 0 & \eta+\beta_1 & \beta_{12} \\ & & & \beta_{21} & \eta-\beta_1 \end{array} \right] \oplus \left[\begin{array}{c|c} \varepsilon+2\eta-\alpha_1 & -\alpha_{21} \\ \hline -\alpha_{12} & \varepsilon+2\eta+\alpha_1 \\ \hline 0 & \xi \end{array} \right] \right)$$

$$\cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{u}(10)).$$

The conormal vector space $V_{x_{16}}^*$ is spanned by $v_1 = (0, 0, u_4 \wedge u_5)$, $v_2 = (0, -u_4 \wedge u_5, 0)$, $v_3 = (u_4 \wedge u_5, 0, 0)$, $v_4 = (0, 0, u_2 \wedge u_4)$, $v_5 = (0, 0, u_2 \wedge u_5)$, $v_6 = (0, 0, u_3 \wedge u_4)$, $v_7 = (0, 0, u_3 \wedge u_5)$, $v_8 = (0, 0, u_2 \wedge u_3)$, $v_9 = (0, 0, u_1 \wedge u_4)$, $v_{10} = (0, 0, u_1 \wedge u_5)$, $v_{11} = (0, -u_2 \wedge u_4, 0)$, $v_{12} = (u_2 \wedge u_4, -u_3 \wedge u_4, 0)$, $v_{13} = (u_3 \wedge u_4, 0, 0)$, $v_{14} = (0, -u_2 \wedge u_5, 0)$, $v_{15} = (u_2 \wedge u_5, -u_3 \wedge u_5, 0)$, $v_{16} = (u_3 \wedge u_5, 0, 0)$.

The action $d\rho_{x_{16}}$ of $\mathfrak{g}_{x_{16}}$ on $V_{x_{16}}^*$ is given by

$$d\rho(A)(v_1, \dots, v_{16}) = (v_1, \dots, v_{16}) \left[\begin{array}{cccc} \underbrace{A_1}_1 & \underbrace{B_1}_2 & \underbrace{B_2}_4 & \underbrace{B_3}_2 \\ & A_2 & & \underbrace{B_4}_6 \\ & & \underbrace{A_3}_1 & \underbrace{B_5}_2 \\ & & \underbrace{A_4}_1 & \underbrace{B_6}_2 \\ & & & \underbrace{A_5}_2 \\ & & & \underbrace{A_6}_6 \end{array} \right]$$

where $(A_1, B_1, B_2) = (-2\eta - \xi, \gamma_{10}, -\gamma_9, \gamma_6, -\gamma_5, \gamma_3, -\gamma_7)$, $B_3 = (\gamma_4, -\gamma_3)$, $A_4 = -2\varepsilon - \xi$, $A_2 = -(\varepsilon + 4\eta)I_2 + \begin{pmatrix} -\alpha_1 & -\alpha_{21} \\ -\alpha_{12} & \alpha_1 \end{pmatrix}$, $B_4 = \begin{pmatrix} \gamma_6 & \gamma_8 & 0 & -\gamma_5 & -\gamma_7 & 0 \\ 0 & \gamma_6 & \gamma_8 & 0 & -\gamma_5 & -\gamma_7 \end{pmatrix}$, $A_5 = (2\varepsilon + \eta - \xi)I_2 + B$ with $B = \begin{pmatrix} -\beta_1 & -\beta_{21} \\ -\beta_{12} & \beta_1 \end{pmatrix}$, $A_3 = -(\varepsilon + \eta + \xi)I_4 + \begin{pmatrix} -\alpha_1 I_2 + B & -\alpha_{21} I_2 \\ -\alpha_{12} I_2 & \alpha_1 I_2 + B \end{pmatrix}$, $A_6 = -(2\varepsilon + 3\eta)I_6 + \begin{pmatrix} -\beta_1 I_3 + A' & -\beta_{21} I_3 \\ -\beta_{12} I_3 & \beta_1 I_3 + A' \end{pmatrix}$ with $A' = \begin{pmatrix} -2\alpha_1 & -2\alpha_{21} & 0 \\ -\alpha_{12} & 0 & -\alpha_{21} \\ 0 & -2\alpha_{12} & 2\alpha_1 \end{pmatrix}$ and

$$(B_5, B_6, B_7) = \left[\begin{array}{c|cc|cc} -\gamma_7 & -\gamma_1 & \gamma_{10} & -\gamma_9 \\ \hline -\gamma_8 & & & \gamma_{10} & -\gamma_9 \\ \hline \gamma_5 & -\gamma_2 & \gamma_{10} & -\gamma_9 \\ \hline \gamma_6 & -\gamma_2 & & \gamma_{10} & -\gamma_9 \end{array} \right]$$

- i) $V_{x_{16}}^* - S_{x_{16}}^* \leftrightarrow v_8 + v_9 + v_{11} + v_{15} + v_{16} = (u_2 \wedge u_5 + u_3 \wedge u_5, -u_2 \wedge u_4 - u_3 \wedge u_5, u_2 \wedge u_3 + u_1 \wedge u_4) \in S_{2,16}^{(2)*}$
- ii) $(S_{x_{16}}^*)_1 \leftrightarrow v_9 + v_{10} + v_{11} + v_{16} = (u_3 \wedge u_5, -u_2 \wedge u_4, u_1 \wedge u_4 + u_1 \wedge u_5) \in S_{3,15}^{(3)*} \leftrightarrow d\rho_1 = -2\varepsilon - \xi \leftrightarrow f_1^*(y) = y_3$
- iii) $(S_{x_{16}}^*)_2 \leftrightarrow v_8 + v_9 + v_{10} + v_{11} + v_{15} = (u_2 \wedge u_5, -u_2 \wedge u_4 - u_3 \wedge u_5, u_2 \wedge u_3 + u_1 \wedge u_4 + u_1 \wedge u_5) \in S_{3,13}^{(2)*} \leftrightarrow d\rho_2 = -8\varepsilon - 12\eta \leftrightarrow f_2^*(y) = \det\begin{pmatrix} y_{11} & y_{14} \\ y_{13} & y_{16} \end{pmatrix}^2 - 4 \det\begin{pmatrix} y_{12} & y_{15} \\ y_{13} & y_{16} \end{pmatrix} \cdot \det\begin{pmatrix} y_{11} & y_{14} \\ y_{12} & y_{15} \end{pmatrix}$
- iv) $(S_{x_{16}}^*)_3 \leftrightarrow v_8 + v_{10} + v_{11} + v_{16} = (u_3 \wedge u_5, -u_2 \wedge u_4, u_1 \wedge u_5 + u_2 \wedge u_3) \in S_{4,11}^{(3)*} \leftrightarrow d\rho_3 = -4\eta - 2\xi \leftrightarrow f_3^*(y) = \det\begin{pmatrix} y_9 & y_{12} \\ y_{10} & y_{15} \end{pmatrix}^2 - \det\begin{pmatrix} y_9 & y_{11} \\ y_{10} & y_{14} \end{pmatrix} \cdot \det\begin{pmatrix} y_9 & y_{13} \\ y_{10} & y_{16} \end{pmatrix}$
- v) $-\delta\chi = d\rho_1 + d\rho_2 + 2d\rho_3, \text{tr}_{V_{x_{16}}^*} = 2d\rho_1 + \frac{3}{2}d\rho_2 + 3d\rho_3.$

(23) The isotropy subalgebra $\mathfrak{g}_{x_{18}}$ at x_{18} is given as follows.

$$(11.23) \quad \mathfrak{g}_{x_{18}} = \left\{ A = \left(\begin{array}{c|c} \varepsilon I_4 + X & * \\ \hline 0 & -4\varepsilon \end{array} \right) \oplus \left(\begin{array}{c|c} -2\varepsilon & * \\ \hline 0 & Y \end{array} \right); X \in \mathfrak{sp}(2), Y \in \mathfrak{gl}(2) \right\} \\ \cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sp}(2) \oplus \mathfrak{sl}(2) \oplus V(6)).$$

The conormal vector space $V_{x_{18}}^*$ is spanned by $v_1 = (0, u_1 \wedge u_3 - u_2 \wedge u_4, 0)$, $v_2 = (0, u_1 \wedge u_2, 0)$, $v_3 = (0, u_1 \wedge u_4, 0)$, $v_4 = (0, u_3 \wedge u_4, 0)$, $v_5 = (0, u_2 \wedge u_3, 0)$, $v_6 = (0, 0, u_1 \wedge u_3 - u_2 \wedge u_4)$, $v_7 = (0, 0, u_1 \wedge u_2)$, $v_8 = (0, 0, u_1 \wedge u_4)$, $v_9 = (0, 0, u_3 \wedge u_4)$, $v_{10} = (0, 0, u_2 \wedge u_3)$, $v_{11} = (0, u_1 \wedge u_5, 0)$, $v_{12} = (0, u_2 \wedge u_3, 0)$, $v_{13} = (0, u_3 \wedge u_5, 0)$, $v_{14} = (0, u_4 \wedge u_5, 0)$, $v_{15} = (0, 0, u_1 \wedge u_5)$, $v_{16} = (0, 0, u_2 \wedge u_5)$, $v_{17} = (0, 0, u_3 \wedge u_5)$, $v_{18} = (0, 0, u_4 \wedge u_5)$. The action $d\rho_{x_{18}}$ of $\mathfrak{g}_{x_{18}}$ on $V_{x_{18}}^*$ is given by

$$(v_1, \dots, v_{18})$$

$$\mapsto (v_1, \dots, v_{18}) \left(\begin{array}{c|c} \begin{array}{c} GL(1) \times GL(1) \times Sp(2) \times SL(2) \\ 2A_1^* \otimes A_1^* \otimes A_2 \otimes A_1^* \end{array} & 0 \\ \hline * & \begin{array}{c} GL(1) \times GL(1) \times Sp(2) \times SL(2) \\ 3A_1 \otimes A_1^* \otimes A_1 \otimes A_1^* \end{array} \end{array} \right)$$

- i) $V_{x_{18}}^* - S_{x_{18}}^* \leftrightarrow v_2 + v_4 + v_8 + v_{10} + v_{15} \in S_{8,18}^*$
- ii) $(S_{x_{18}}^*)_1 \leftrightarrow v_1 + v_8 + v_{10} + v_{11} \in S_{8,18}^* \leftrightarrow d\rho_1 = -2\varepsilon - 6\eta \leftrightarrow \text{deg } f_1^* = 6$
- iii) $(S_{x_{18}}^*)_2 \leftrightarrow v_1 + v_7 + v_{17} + v_{18} \in S_{8,18}^* \leftrightarrow d\rho_2 = -8\varepsilon - 4\eta \leftrightarrow \text{deg } f_2^* = 4$
- iv) $-\delta\chi = 3d\rho_1 - 2d\rho_2, \text{tr}_{V_{x_{18}}^*} = 4d\rho_1 - \frac{3}{2}d\rho_2.$

(24) The isotropy subalgebra $\mathfrak{g}_{x_{21}}$ at x_{21} is given as follows.

$$(11.24) \quad \mathfrak{g}_{x_{21}} = \left\{ A = \left(\begin{array}{c|c} 3\varepsilon I_3 + X & * \\ \hline 0 & -2\varepsilon I_3 + Y \end{array} \right) \oplus \left(\begin{array}{c|c} -6\varepsilon & * \\ \hline 0 & \eta I_2 + Z \end{array} \right); \right. \\
\left. X, Z \in \mathfrak{sl}(2), Y \in \mathfrak{sl}(3) \right\} \\
\cong (\mathfrak{gl}(1) \oplus \mathfrak{gl}(1) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(3) \oplus V(8)).$$

Then $V_{x_{21}}^*$ is spanned by $v_1 = (0, u_4 \wedge u_5, 0)$, $v_2 = (0, u_3 \wedge u_5, 0)$, $v_3 = (0, u_3 \wedge u_4, 0)$, $v_4 = (0, 0, u_4 \wedge u_5)$, $v_5 = (0, 0, u_3 \wedge u_5)$, $v_6 = (0, 0, u_3 \wedge u_4)$, $v_7 = (u_4 \wedge u_5, 0, 0)$, $v_8 = (u_3 \wedge u_5, 0, 0)$, $v_9 = (u_3 \wedge u_4, 0, 0)$, $v_{10} = (0, u_1 \wedge u_3, 0)$, $v_{11} = (0, u_1 \wedge u_4, 0)$, $v_{12} = (0, u_1 \wedge u_5, 0)$, $v_{13} = (0, u_2 \wedge u_3, 0)$, $v_{14} = (0, u_2 \wedge u_4, 0)$, $v_{15} = (0, u_2 \wedge u_5, 0)$, $v_{16} = (0, 0, u_1 \wedge u_3)$, $v_{17} = (0, 0, u_1 \wedge u_4)$, $v_{18} = (0, 0, u_1 \wedge u_5)$, $v_{19} = (0, 0, u_2 \wedge u_3)$, $v_{20} = (0, 0, u_2 \wedge u_4)$, $v_{21} = (0, 0, u_2 \wedge u_5)$. The action $d\rho_{x_{21}}$ of $\mathfrak{g}_{x_{21}}$ on $V_{x_{21}}^*$ is given by

(v_1, \dots, v_{21})

$$\mapsto (v_1, \dots, v_{21}) \left[\begin{array}{c|c|c} \begin{array}{c} GL(1) \times GL(1) \times SL(2) \\ \times SL(3) \\ 4A_1 \otimes A_1^* \otimes A_1^* \otimes A_1 \end{array} & * & * \\ \hline \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} GL(1) \times SL(3) \\ (10A_1) \otimes A_1 \end{array} & 0 \\ \hline \begin{array}{c} 0 \\ 0 \end{array} & 0 & \begin{array}{c} GL(1) \times GL(1) \times SL(2) \\ \times SL(2) \times SL(3) \\ A_1^* \otimes A_1^* \otimes A_1^* \otimes A_1^* \otimes A_1^* \end{array} \end{array} \right]$$

- i) $V_{x_{21}}^* - S_{x_{21}}^* \leftrightarrow v_7 + v_{10} + v_{14} + v_{18} + v_{19} = (u_4 \wedge u_5, u_1 \wedge u_3 + u_2 \wedge u_4, u_1 \wedge u_5 + u_2 \wedge u_3) \in S_{1,21}^{(2)*}$
- ii) $(S_{x_{21}}^*)_1 \leftrightarrow v_8 + v_{10} + v_{14} + v_{18} + v_{19} = (u_3 \wedge u_5, u_1 \wedge u_3 + u_2 \wedge u_4, u_1 \wedge u_5 + u_2 \wedge u_3) \in S_{3,13}^{(2)*} \leftrightarrow d\rho_1 = 18\varepsilon - 2\eta \leftrightarrow \deg f_1^* = 4$
- iii) $(S_{x_{21}}^*)_2 \leftrightarrow v_8 + v_9 + v_{10} + v_{14} + v_{18} = (u_3 \wedge u_4 + u_3 \wedge u_5, u_1 \wedge u_3 + u_2 \wedge u_4, u_1 \wedge u_5) \in S_{2,16}^{(2)*} \leftrightarrow d\rho_2 = -6\varepsilon - 6\eta \leftrightarrow \deg f_2^* = 6$
- iv) $-\delta\chi = 2d\rho_1 + d\rho_2, \text{tr}_{V_{x_{21}}^*} = 3d\rho_1 + 2d\rho_2.$

(25) The isotropy subalgebra $\mathfrak{g}_{x_{30}}$ at $x_{30} = 0$ is \mathfrak{g} itself.

This is a good holonomic variety and $\text{ord}_A f^s = -15s - \frac{3}{2}0$. Thus we obtain the holonomy diagram (Figure 11-1). From this diagram, we obtain the b -function $b(s) = ((s+1)(s+\frac{3}{2})(s+2))^3 \cdot ((s+\frac{4}{3})(s+\frac{5}{3}))^2 \cdot (s+\frac{5}{4})(s+\frac{7}{4})$.

Remark. Let $A_0 = \overline{G(x_0, y_0)}$ and $A_1 = \overline{G(x_1, y_1)}$ be good holonomic varieties satisfying $(x_0, y_1) \in A_0 \cap A_1$ and $\dim G(x_0, y_1) = \dim V - 1$. Then we can calculate β by Proposition 1-4. It is known that if β depends on the choice of A_1 , then (x_0, y_1) is not contained in other A_i ($i \neq 0, 1$), i.e., there are no three A_i 's which intersect at (x_0, y_1) with codimension one. (If more




Fig. 11-1. Holonomy diagram of $(SL(5) \times GL(3), A_2 \times A_1, V(10) \otimes V(3))$ where $\binom{(k)}{i j}$ denotes the conormal bundles of the orbit $S_{ij}^{(k)}$ in Proposition 6-1.

than two A_i 's intersect with codimension one, then $\beta = 1$ and it does not depend on A_1 .) All one-codimensional intersections obtained from (1)~(14) satisfy this condition except $A_{5,5}^{(2)} \cap A_{4,11}^{(3)}$ and $A_{8,5}^{(3)} \cap A_{7,9}^{(2)}$. In general, if $(x_0, y_1) \in \overline{G(x_2, y_2)}$, then we have $\text{codim}_V \rho(G)x_0 \geq \text{codim}_V \rho(G)x_1$ and $\text{codim}_{V^*} \rho^*(G)y_1 \geq \text{codim}_{V^*} \rho^*(G)y_2$. From this, there are no other A 's satisfying $\dim A \cap A_{i,5}^{(2)} \cap A_{4,11}^{(3)} = \dim V - 1$. For $A_{8,5}^{(3)} \cap A_{7,9}^{(2)}$, it is enough to check $A_{7,7}^{(1)}$, $A_{7,7}^{(2)}$ and $A_{6,8}^{(2)}$. By using the duality, i.e., $(G, \rho, V) \cong (G, \rho^*, V^*)$, we get all one-codimensional intersections of three good holonomic varieties.

§ 12. Table of the b -functions of irreducible reduced regular P.V.'s

- (1) $(G \times GL(m), \rho \otimes A_1, V(m) \otimes V(m))$ where $\rho : G \rightarrow GL(V(m))$ is an m -dimensional irreducible representation of a connected semi-simple algebraic group G (or $G = \{1\}$ and $m = 1$).

$$b(s) = (s+1)(s+2)\cdots(s+m) \text{ (See Figure 2-1 and 2-4).}$$

- (2) $(GL(n), 2A_1, V(\frac{1}{2}n(n+1)))$ ($n \geq 2$)

$$b(s) = \prod_{\nu=1}^n \left(s + \frac{\nu+1}{2} \right) = (s+1) \left(s + \frac{3}{2} \right) \cdots \left(s + \frac{n+1}{2} \right)$$

(See Figure 2-2 and 2-4).

- (3) $(GL(2m), A_2, V(m(2m-1)))$ ($m \geq 3$)

$$b(s) = \prod_{k=1}^m (s+2k-1) = (s+1)(s+3)\cdots(s+2m-1)$$

(See Figure 2-3 and 2-4).

- (4) $(GL(2), 3A_1, V(4))$

$$b(s) = (s+1)^2(s+\frac{5}{6})(s+\frac{7}{6}) \text{ (See [2]).}$$

- (5) $(GL(6), A_3, V(20))$

$$b(s) = (s+1)(s+\frac{5}{2})(s+\frac{7}{2})(s+5) \text{ (See Figure 8-1).}$$

- (6) $(GL(7), A_3, V(35))$

$$b(s) = (s+1)(s+2)(s+\frac{5}{2})(s+\frac{7}{2})(s+3)(s+4)(s+5)$$

(See Figure 10-1).

- (7) $(GL(8), A_3, V(56))$

$$b(s) = (s+1)(s+\frac{3}{2})^2(s+\frac{11}{6})(s+2)^3(s+\frac{13}{6})(s+\frac{7}{3})(s+\frac{5}{2})^3(s+\frac{8}{3})(s+3)^2(s+\frac{7}{2})$$

(See [10]).

- (8) $(SL(3) \times GL(2), 2A_1 \otimes A_1, V(6) \otimes V(2))$

$$b(s) = \{(s+1)^2(s+\frac{5}{6})(s+\frac{7}{6})(s+\frac{3}{4})(s+\frac{5}{4})\}^2 \text{ (See [12]).}$$

- (9) $(SL(6) \times GL(2), A_2 \otimes A_1, V(15) \otimes V(2))$

$$b(s) = (s+1)^2(s+\frac{5}{6})(s+\frac{7}{6})(s+\frac{3}{2})^2(s+2)^2(s+\frac{5}{2})(s+\frac{7}{2})(s+\frac{8}{3})$$

(See [12]).

- (10) $(SL(5) \times GL(3), A_2 \otimes A_1, V(10) \otimes V(3))$
 $b(s) = ((s+1)(s+\frac{3}{2})(s+2))^3 \cdot ((s+\frac{4}{3})(s+\frac{5}{3}))^2 \cdot (s+\frac{5}{4})(s+\frac{7}{4})$
 (See Figure 11-1).
- (11) $(SL(5) \times GL(4), A_2 \otimes A_1, V(10) \otimes V(4))$ (See [11]).
- (12) $(SL(3) \times SL(3) \times GL(2), A_1 \otimes A_1 \otimes A_1, V(3) \otimes V(3) \otimes V(2))$
 $b(s) = (s+1)^4(s+\frac{3}{2})^4(s+\frac{4}{3})(s+\frac{5}{3})(s+\frac{5}{6})(s+\frac{7}{6})$ (See [12]).
- (13) $(Sp(n) \times GL(2m), A_1 \otimes A_1, V(2n) \otimes V(2m))$ ($n \geq 2m \geq 2$)
 $b(s) = \prod_{k=1}^m (s+2k-1) \prod_{\ell=0}^{m-1} (s+2n-2\ell)$
 $= (s+1)(s+3) \cdots (s+2m-1)(s+2n)(s+2n-2) \cdots$
 $(s+2n-2m+2)$ (See Figure 3-1 and 3-2).
- (14) $(GL(1) \times Sp(3), \square \otimes A_3, V(1) \otimes V(14))$
 $b(s) = (s+1)(s+2)(s+\frac{5}{2})(s+\frac{7}{2})$ (See Figure 9-1).
- (15) $(SO(n) \times GL(m), A_1 \otimes A_1, V(n) \otimes V(m))$ ($n \geq 3, \frac{n}{2} \geq m \geq 1$)
 $b(s) = \prod_{k=1}^m \left(s + \frac{k+1}{2}\right) \prod_{\ell=1}^m \left(s + \frac{n-\ell+1}{2}\right)$
 $= (s+1) \left(s + \frac{3}{2}\right) \cdots \left(s + \frac{m+1}{2}\right) \left(s + \frac{n}{2}\right) \left(s + \frac{n-1}{2}\right) \cdots$
 $\left(s + \frac{n-m+1}{2}\right)$ (See [2]).
- (16) $(GL(1) \times Spin(7), \square \otimes \text{spin rep.}, V(1) \otimes V(8))$
 $b(s) = (s+1)(s+4)$ (See Remark in § 5).
- (17) $Spin(7) \times GL(2), \text{spin rep.} \otimes A_1, V(8) \otimes V(2)$
 $b(s) = (s+1)(s+\frac{3}{2})(s+4)(s+\frac{7}{2})$ (See Remark in § 5).
- (18) $(Spin(7) \times GL(3), \text{spin rep.} \otimes A_1, V(8) \otimes V(3))$
 $b(s) = (s+1)(s+\frac{3}{2})(s+2)(s+4)(s+\frac{7}{2})(s+3)$ (See Remark in § 5).
- (19) $(GL(1) \times Spin(9), \square \otimes \text{spin rep.}, V(1) \otimes V(16))$
 $b(s) = (s+1)(s+8)$ (See Remark in § 5).
- (20) $(Spin(10) \times GL(2), \text{half-spin rep.} \otimes A_1, V(16) \otimes V(2))$
 $b(s) = (s+1)(s+4)(s+5)(s+8)$ (See Figure 4-1).
- (21) $(Spin(10) \times GL(3), \text{half-spin rep.} \otimes A_1, V(16) \otimes V(3))$
 $b(s) = (s+1)(s+\frac{3}{2})(s+2)(s+3)(s+\frac{7}{2})(s+4)(s+\frac{5}{3})(s+\frac{6}{3})(s+\frac{7}{3}) \times$
 $\times (s+\frac{8}{3})(s+\frac{9}{3})(s+\frac{10}{3})$ (See [15]).
- (22) $(GL(1) \times Spin(11), \square \otimes \text{spin rep.}, V(1) \otimes V(32))$
 $b(s) = (s+1)(s+\frac{7}{2})(s+\frac{11}{2})(s+8)$ (See Remark in § 5).
- (23) $(GL(1) \times Spin(12), \square \otimes \text{half-spin rep.}, V(1) \otimes V(32))$
 $b(s) = (s+1)(s+\frac{7}{2})(s+\frac{11}{2})(s+8)$ (See Figure 5-1).

- (24) $(GL(1) \times Spin(14), \square \otimes \text{half-spin rep.}, V(1) \otimes V(64))$
 $b(s) = (s+1)(s+\frac{5}{2})(s+\frac{7}{2})(s+4)(s+5)(s+\frac{11}{2})(s+\frac{13}{2})(s+8)$
 (See Appendix).
- (25) $(GL(1) \times (G_2), \square \otimes A_2, V(1) \otimes V(7))$
 $b(s) = (s+1)(s+\frac{7}{2})$ (See Remark in § 5).
- (26) $((G_2) \times GL(2), A_2 \otimes A_1, V(7) \otimes V(2))$
 $b(s) = (s+1)(s+\frac{3}{2})(s+\frac{7}{2})(s+3)$ (See Remark in § 5).
- (27) $(GL(1) \times E_6, \square \otimes A_1, V(1) \otimes V(27))$
 $b(s) = (s+1)(s+5)(s+9)$ (See Figure 6-1).
- (28) $(E_6 \times GL(2), A_1 \otimes A_1, V(27) \otimes V(2))$
 $b(s) = (s+1)^2(s+\frac{5}{6})(s+\frac{7}{6})(s+\frac{5}{2})^2(s+3)^2(s+\frac{9}{2})^2(s+\frac{13}{3})(s+\frac{14}{3})$
 (See [12]).
- (29) $(GL(1) \times E_7, \square \otimes A_6, V(1) \otimes V(56))$
 $b(s) = (s+1)(s+\frac{11}{2})(s+\frac{13}{2})(s+14)$ (See Figure 7-1).

We can obtain the b -functions of all irreducible regular P.V.'s, except for those in the castling class of (11), from the Table above and the following theorem due to T. Shintani.

THEOREM (T. Shintani). *Let (G', ρ', V') be a castling transform of an irreducible regular P.V. (G, ρ, V) , i.e., there exists a triplet $(\tilde{G}, \tilde{\rho}, V(m))$ and a positive number n with $m > n \geq 1$ such that*

$$\begin{aligned} (G, \rho, V) &\cong (\tilde{G} \times GL(n), \tilde{\rho} \otimes A_1, V(m) \otimes V(n)) \\ (G', \rho', V') &\cong (\tilde{G} \times GL(m-n), \tilde{\rho}^* \otimes A_1, V(m)^* \otimes V(m-n)). \end{aligned}$$

Then the b -functions $b(s)$ and $b'(s)$ of them satisfy

$$\begin{aligned} b(s) &\prod_{i=1}^d (ds-i)(ds-i+1) \cdots (ds-i+m-n-1) \\ &= b'(s) \prod_{i=1}^d (ds-i)(ds-i+1) \cdots (ds-i+n-1) \end{aligned}$$

where $\deg f = dm$ and $\deg f' = d(m-n)$. Here f and f' are the basic relative invariants of (G, ρ, V) and (G', ρ', V') respectively.

Appendix with I. Ozeki

Here we consider the regular irreducible P.V. $(GL(1) \times Spin(14), \square \otimes \text{half-spin rep.}, V(1) \otimes V(64))$. The orbital decomposition of this space has been done by the author and I. Ozeki ([7]), by Popov ([9]), by V. Gatti and E. Viniberghi ([10]). There exist ten orbits, and the conormal bundle

of each orbit is a good Lagrangian variety. The relative invariant of this space is of degree eight ([1]), and its b -function is given by $b(s) = (s + 1)(s + \frac{5}{2})(s + \frac{7}{2})(s + 4)(s + 5)(s + \frac{11}{2})(s + \frac{13}{2})(s + 8)$. Its holonomy diagram is given by Figure A, where we denote by \textcircled{m} the conormal bundle A of the m -codimensional orbit.




Figure A. Holonomy diagram of $(GL(1) \times Spin(14))$,
 $\square \otimes$ half-spin rep., $V(1) \otimes V(64)$

REFERENCES

- [1] M. Sato and T. Kimura, A classification of irreducible prehomogeneous vector spaces and their relative invariants, Nagoya Math. J., **65** (1977), 1–155.
- [2] M. Sato, M. Kashiwara, T. Kimura and T. Oshima, Micro-local analysis of prehomogeneous vector spaces, Inv. math., **62** (1980), 117–179.
- [3] J-I. Igusa, A classification of spinors up to dimension twelve, Amer. J. of Math., **92**, No. 4 (1970), 997–1028.
- [4] C. Chevalley, The Algebraic Theory of Spinors, Columbia University Press, 1954.
- [5] S. J. Haris, Some irreducible representation of exceptional algebraic groups, Amer. J. of Math., **93** (1971).
- [6] G. B. Gurevich, Foundations of the Theory of Algebraic Invariants, P. Noordhoff LTD-Groningen, 1964.
- [7] T. Kimura, Study of irreducible prehomogeneous vector spaces, Master Thesis in Japanese, University of Tokyo, 1973, 373 pp.
- [8] H. Kawahara, Prehomogeneous vector spaces related with the spin group, Master Thesis in Japanese, University of Tokyo, 1974.
- [9] V. L. Popov, A classification of spinors of dimension fourteen, Uspechi Math. Nauk, **32**, (1977).
- [10] V. Gatti and E. Viniberghi, Spinors of 13-dimensional space, Adv. Math., **30**, No. 2 (1978).
- [11] I. Ozeki, On the micro-local structure of the regular prehomogeneous vector space associated with $SL(5) \times GL(4)$, I, Proc. Japan Acad., **55**, Ser. A, (1979).
- [12] T. Kimura and M. Muro, On some series of regular irreducible prehomogeneous vector spaces, Proc. Japan Acad., **55**, Ser A, No. 10 (1979), 384–389.
- [13] I. Ozeki, On the micro-local structure of a regular prehomogeneous vector space associated with $GL(8)$, Proc. Japan Acad., **56**, Ser. A, No. 1 (1980), 18–21.
- [14] T. Kimura, Remark on some combinatorial construction of relative invariants, Tsukuba J. of Math., **5**, No. 1 (1981).
- [15] T. Kimura and I. Ozeki, On the micro-local structure of regular prehomogenous vector space associated with $Spin(10) \times GL(3)$. (in preparation).

*Institute of Mathematics
University of Tsukuba*