

The Bacterial Microflora of Fish, Revised

B. Austin

School of Life Sciences, John Muir Building, Heriot-Watt University, Riccarton,
Edinburgh EH14 4AS, Scotland

Original review published March 4, 2002; Revised edition published August 11, 2006

This Revised Review is a revision of the review article published, [Vol. 2, 2002](#).

The results of numerous studies indicate that fish possess bacterial populations on or in their skin, gills, digestive tract, and light-emitting organs. In addition, the internal organs (kidney, liver, and spleen) of healthy fish may contain bacteria, but there is debate about whether or not muscle is actually sterile. Using traditional culture-dependent techniques, the numbers and taxonomic composition of the bacterial populations generally reflect those of the surrounding water. More modern culture-independent approaches have permitted the recognition of previously uncultured bacteria. The role of the organisms includes the ability to degrade complex molecules (therefore exercising a potential benefit in nutrition), to produce vitamins and polymers, and to be responsible for the emission of light by the light-emitting organs of deep-sea fish. Taxa, including *Pseudomonas*, may contribute to spoilage by the production of histamines in fish tissue.

KEYWORDS: bacteria, fish, microflora, methods, digestive tract, gills, skin, population size, taxonomy, biodiversity, luminescence, degradative ability, effect of antibiotics, polymers, enzymes, spoilage

INTRODUCTION

Traditionally, studies on fish-associated microorganisms involved culture-dependent techniques of dubious sensitivity, which highlighted only the bacteria (typically the aerobic heterotrophic bacterial component[1]) to the exclusion of eukaryotes. Anaerobic bacteria have been comparatively neglected[2,3,4,5] by culturists, possibly reflecting the need for more exacting techniques, although there is increasing evidence that such organisms occur in large numbers especially within the digestive tract of freshwater and marine fish[2]. More recently and in line with other studies of microbial biodiversity, emphasis has been placed on molecular-based culture-independent techniques, which have been generating some exciting data, and have revealed the presence of uncultured organisms including anaerobes[6].

This article, which is an updated version first published in 2002[143], will synthesise the available information on fish-associated bacteria, focusing on the numbers, nature, and role of bacteria on or in healthy finfish. Aspects of fish pathology will be ignored, as a wealth of information sufficient to fill several textbooks already exists[7]. However, it is apparent that some pathogens may be found on healthy fish in the absence of disease. It is questionable whether such associations represent the asymptomatic carrier state of the disease cycle, a preliminary colonisation step prior to pathogenesis, or commensalism-synergism. For example, *Flavobacterium psychrophilum*, the causal agent of coldwater disease (of

salmon) and rainbow trout fry syndrome, has been found in the kidney, spleen, brain, ovarian fluid, unfertilised eggs, and milt of healthy Baltic salmon (*Salmo salar*)[8].

It is apparent that fish are continuously exposed to the microorganisms present in water and in sediment including the contaminants in sewage/faeces[9]. These organisms will undoubtedly influence the microflora on external surfaces, including the gills, of fish. Similarly, the digestive tract will receive water and food that is populated with microorganisms. Certainly, colonisation may well start at the egg and/or larval stage, and continue with the development of the fish[10]. Thus, the numbers and range of microorganisms present in the eggs, on food, and in the water, will influence the microflora of the developing fish. Also, it is recognised that, to some extent, it is possible to manipulate the microflora of the developing fish by use of prebiotics, i.e., nondigestible food ingredients that beneficially affect the host by stimulating growth[11] and probiotics, i.e., live microbial food supplements, which may colonise the digestive tract for short or prolonged periods[12]. This action may have benefit for the host, such as the moderation of fish diseases[12].

From the published literature, it may be deduced that there are three likely scenarios for the fate of bacteria coming into contact with fish:

1. The organisms from the environment around the fish may become closely associated with and even colonise the external surfaces of the fish. There may be accumulation of the organisms at sites of damage, such as missing scales or abrasions[13].
2. The organisms may enter the mouth with water[10] or food and pass through and/or colonise the digestive tract[13].
3. The organisms coming into contact with fish surfaces may be inhibited by the resident microflora or by natural inhibitory compounds present on or in the fish[13].

The overriding problem concerns whether or not it is possible to differentiate members of the indigenous (fish) microflora from transients, which could be in the water film around fish or in water/food within the digestive tract. This is a problem particularly with the culture-dependent approaches. Unfortunately, most publications do not address this issue. Yet this is not so unusual insofar as similar arguments have centred on the nature of the true microflora of other habitats, e.g., the distinction between microbial populations of the rhizoplane (root surface) vs. the rhizosphere (habitat around roots), and of the phylloplane (leaf surface) vs. the phyllosphere (habitat around leaves).

It is recognised that extraneous bacteria are capable of surviving in fish. For example, the faecal indicator organism, *Escherichia coli*, has been found to survive and even multiply in the digestive tract of rainbow trout (*Oncorhynchus mykiss*) after ingestion via contaminated food[14].

RESEARCH ACTIVITIES

Research has focused on six principal aspects of the microbiology of fish:

- The microbiology of the surface (including gills)
- The populations in the digestive tract (an area of current interest particularly involving use of modern molecular-based culture-independent techniques)
- The possible presence of bacteria in muscle and the internal organs of healthy fish
- The microflora of eggs
- The presence and role of bacteria associated with the light-emitting organs, particularly of deep-sea fish
- The bacterial populations in food

As a simplification, publications have tended to emphasise either quantitative or qualitative aspects or the supposed role of the organisms on/in fish. It is unusual for research articles to address more than one of these aspects.

METHODS USED TO STUDY FISH-ASSOCIATED BACTERIA

Dilution and Spread-Plating Techniques

There has been a tendency for scientists to study fish-associated bacteria using only traditional, comparatively simple, culture-dependent techniques, such as dilution or spread plating, incorporating media and incubation conditions often of dubious relevance. As an example, to isolate bacteria from the skin, the surface of the fish may be swabbed over an indeterminate area, and the inoculum spread over the surface of nutrient-rich medium, such as tryptone soya agar (TSA[15]), with incubation at 15–25°C for 7–14 days. A disadvantage of using swabbing techniques is that it is difficult to equate the data with a defined unit of measurement. In addition, the relevance of the resulting data for ascertaining the indigenous bacterial populations on the surface of fish is questionable. Yet culturing methods are still used extensively and publications appear regularly[e.g., 16,17,18].

Criticism may also be levied at the time taken between sample collection and examination, which may often be measured in hours or days. It is not uncommon for whole or parts (e.g., the digestive tract) of fish to be transported on ice, cooled or at ambient temperature, to distant laboratories for examination. Sections or the entire digestive tract, together with the contents, have then been homogenised, in which case it is impossible to decide from the data if the resulting bacteria have originated from the food particles, lumen, and/or the wall. Some workers have studied the bacterial populations of the digestive tract by swabbing the anus and faeces[13]. Unfortunately, most scientists do not consider whether or not there may be significant changes in the microflora during the period from collection of the fish to microbiological examination.

As a final criticism, it is noted that the proportion of the bacterial microflora contributing to the colony count is rarely considered in quantitative-type studies. Circumstantial evidence suggests that populations deduced from colony counts on agar plates greatly underestimate the total microflora likely to be present. Nevertheless, a study considered that at much as 50% of the microflora from the intestine of rainbow trout produced colonies on TSA[15]; thus, a 50% error might be inferred!

Some comparisons between methods have been conducted. For example, gentle rinsing techniques have been evaluated and compared to excision and homogenisation with a stomacher to recover bacteria from the surface of fish[19]. In this case, stomaching was regarded as superior for the enumeration of total bacterial populations, but rinsing was better for rainbow trout than striped bass (*Morone saxatilis*)[19].

Microscopic Techniques

Microscopic techniques have found increasing use in the study of fish microflora, and include direct microscopic counts by light[15,20,21] and electron microscopy[20,22,23,24]. These have been used to visualise the organisms present, particularly in the digestive tract.

Automated Direct Epifluorescent Filter Technique

An automated direct epifluorescent filter technique instrument, COBRA, has been evaluated and offers promise for enumerating bacterial populations[25].

Molecular Techniques

Molecular methods are increasingly being used. For example, numerous publications have discussed the sequencing of 16S rRNA genes[15,26,27,28,29,30]. Also, microplate hybridisation has been successful at identifying aeromonads in the digestive tract of freshwater fish[31].

QUANTITATIVE ASPECTS OF THE BACTERIAL MICROFLORA

Surface Populations

Most workers have opted for the comparatively easy approach of studying the aerobic heterotrophic bacteria populations, with data suggesting that the numbers of bacteria on the surface of fish approximate those of the surrounding water. Yet, in retrospect, it is apparent that the units of measurement between water (populations ml^{-1}) and fish surfaces (populations cm^{-2}) are very different, and comparisons are not especially helpful.

Scrutiny of the literature suggests that fish have only low bacterial populations on the skin. For example, Atlantic salmon (*Salmo salar*) from the U.K. were reported to possess populations of 10^2 – 10^3 culturable bacteria cm^{-2} of skin[32], whereas rainbow trout from Turkey contained bacterial populations of 10^1 – 10^7 g^{-1} [33]. However, it should be emphasised that the relevance of using weight as a unit of measurement to estimate bacterial populations on skin is debatable. Interestingly, freshly caught mullet (*Mugil cephalus*), whiting (*Sillago ciliato*), and flathead (*Platycephalus fiscus*) from Australia were reported to have seemingly higher populations of 4×10^3 to 8×10^4 bacteria cm^{-2} [34]. Not surprisingly, there are data suggesting that the bacterial population size reflects the level of water pollution, i.e., higher counts are present on fish in polluted waters[35]. Also, there is some evidence that the population of aerobes exceeds that of the anaerobes[36].

Bacterial populations of skin

10^2 to $\sim 10^4$ cm^{-2}

Overall, these low counts, which to some extent have been supported by scanning electron microscopic evidence[13], indicate that only a minute area of the fish surface is populated with bacteria. However, it is conceded that the preparation for scanning electron microscopy may well have removed some organisms from the skin. Thus, it could be inferred that the surface microflora is only loosely associated with fish skin. Coincidentally, this low level of colonisation. on fish skin corresponds well to that of other habitats, such as the leaf surface of terrestrial plants[37].

Gills

Gill tissue has been found to harbour high bacterial populations, e.g., up to 10^6 bacteria g^{-1} of gill tissue[38].

Eyes

There is anecdotal evidence that the eyes of healthy fish are devoid of bacterial colonisation[13].

Muscle and the Internal Organs

Muscle has been considered by some to be sterile[39], whereas other investigators have reported the presence of bacteria[40]. Also, some workers have found bacteria in the kidney and liver of healthy fish, i.e., turbot (*Scophthalmus maximus*)[41].

Digestive Tract

A consensus view is that dense bacterial populations occur in the digestive tract (i.e., populations of up to $\sim 10^8$ heterotrophs g^{-1} [42,43,44,45,46]) and $\sim 10^5$ anaerobes g^{-1} [3,42,43] have been reported with numbers appearing to be much higher than those of the surrounding water. For example, by including contents with

the intestine, maximal bacterial populations of 2×10^7 colony-forming units (cfu) g^{-1} were recorded in the pinfish (*Lagodon rhomboides*)[47]. Moreover, counts of 1.1×10^6 to 3.6×10^8 bacteria were recorded for the intestinal contents of deep-sea fish[45], when it was noted that more cultures were obtained at *in situ* (barophilic), rather than atmospheric, pressure[45]. Also, some differences have been considered to reflect seasonality, i.e., with maximum and minimum counts occurring in summer and winter, respectively[48]. Indeed, an effect of water temperature on the size of the microflora of pike perch (*Stizostedion lucioperca*) has been reported[49]. In another study, seasonal variation was attributed to the monsoon season, with maximal and minimal populations in green chromides (*Eetroplus suratensis*) and orange chromides (*E. maculates*) corresponding with postmonsoon (September to December) and premonsoon (January to April), respectively[50]. Also, the population densities are likely to be influenced by the feeding regime, with fish receiving live feeds having higher populations than those with artificial diets[46].

Differences in population size have been detected in specific regions of the digestive tract. Thus, an estimation of aerobic heterotrophs in the digestive tract of yellowtail (*Seriola* sp.) revealed counts of 2×10^4 bacteria g^{-1} , 2.5×10^5 bacteria g^{-1} , and 6.5×10^4 to 5.9×10^6 bacteria g^{-1} in the pyloric caeca, stomach, and intestine, respectively[51]. Parallel data were published separately[49], when the presence of 5.5×10^3 to 5.0×10^4 cfu of aerobic heterotrophic bacteria g^{-1} and 1.0×10^4 to 1.0×10^7 cfu aerobic heterotrophic bacteria g^{-1} were found in the stomach and intestine of pike perch, respectively. However, higher populations were noted in the digestive tract of juvenile compared with adult farmed Dover sole (*Solea solea*), with 5.2×10^5 , 8.0×10^5 , and 9.8×10^6 aerobic heterotrophs g^{-1} recovered from the stomach-foregut, midgut, and hindgut-rectum (of juvenile fish), respectively[4,52]. It should be emphasised that anaerobes (7.1×10^5 anaerobic bacteria g^{-1}) have been found in addition within the intestines[51]. Following a familiar theme, it was observed that there was an increase in bacterial populations, especially of adherent organisms, along the digestive tract of herring (*Clupea harengus*) larvae[20].

Some differences have been noted according to the feeding pattern of fish. Thus, detritivorous fish possessed higher bacterial populations than filter feeders[53]. Of course, it is likely that many organisms in the digestive tract will have been derived from the food. In this connection, it was found that there were between 10^3 and 10^7 aerobic heterotrophs g^{-1} in commercial fish food in North America[54], whereas comparable data from Japan indicated counts of 1.8×10^3 to 8.0×10^5 bacteria g^{-1} [55].

Electron microscopy has substantiated the presence of high bacterial populations in the digestive tract. In particular, scanning and transmission electron microscopy demonstrated the presence of large populations of ovoid and rod-shaped bacterial-like objects in association with the microvillous brush borders of the enterocytes of Arctic charr (*Salvelinus alpinus*)[23]. Also, bacteria were observed at and between the tips of microvilli, and rod-shaped bacteria were seen between the microvilli of common wolffish (*Anarhichas lupus*)[22]. Evidence has pointed to endocytosis of bacteria by epithelial cells in the pyloric caeca and midgut[23].

Bacterial populations in the digestive tract

$\sim 10^8$ aerobic heterotrophs g^{-1}

$\sim 10^5$ anaerobic g^{-1}

Fish Eggs and Larvae

Fish eggs may be populated by high numbers of bacteria, with 10^3 – 10^6 bacteria g^{-1} reported for salmonid eggs[56]. There is evidence that adhesion and colonisation of the egg by bacteria occurs within a few hours of fertilization[57]. Undoubtedly, these organisms and those of the food and surrounding water are important for the establishment of a microflora in the digestive tract of fish larvae. Incidentally, the digestive tract of newly hatched larvae contains scant bacterial populations, but is quickly colonised[58].

TAXONOMY (BIODIVERSITY)

Approaches have gone from the traditional[59], through numerical taxonomy studies involving large numbers of isolates (e.g., 197 cultures investigated in one study[60]; 473 isolates studied in another[1]), to culture-independent molecular approaches (e.g., partial sequencing of the 16S rRNA gene[15,29]). The benefit of the latter is the recognition of organisms that may or may not be culturable by conventional techniques[e.g., 21]. Sometimes, the phenetic approach has centered on the use of rapid systems, such as BIOLOG or MIDI[36,61]. It is encouraging that some comparative studies have pointed to congruence between phenotypic and molecular analyses[15]. Overall, it would appear that narrow-based studies focusing on a limited number of bacterial groups are often more successful than those that attempt to be broad-based, trying to consider all of the bacteria from fish. From the published literature, it is apparent that there are many similarities between the bacterial populations in fish and water[33,39,40,59,62,63, 64,65,66,67,68].

Surface Microflora

The bacteria from the surface of freshwater fish have been reported to include *Acinetobacter johnsonii*[69], aeromonads (notably *Aeromonas hydrophila*, *A. bestiarum*, *A. caviae*, *A. jandaei*, *A. schubertii*, and *A. veronii* biovar *sobria*[70]), *Alcaligenes piechaudii*, *Enterobacter aerogenes*, *Escherichia coli*, *Flavobacterium*[35], *Flexibacter* spp., *Micrococcus luteus*, *Moraxella* spp., *Pseudomonas fluorescens*, psychrobacters[69], and *Vibrio fluvialis*[33,67,71]. To some extent, the presence of aeromonads reflected whether or not the water in which the fish occurred was polluted or clean[70]. Bacteria, typical of those in seawater, have been recovered from the surface of marine fish and include *Acinetobacter calcoaceticus*, *Alcaligenes faecalis*, *Bacillus cereus*, *B. firmus*, *Caulobacter*, coryneforms, *Cytophaga/Flexibacter*, *E. coli*, *Hyphomicrobium vulgare*, *Lucibacterium (Vibrio) harveyi*, *Photobacterium angustum*, *P. logei*, *Prosthecomicrobium*, *Pseudomonas fluorescens*, *P. marina*, and *Vibrio* spp. (including *V. albensis*, *V. anguillarum*, *V. splendidus* biotype I, *V. fischeri*, *V. ordalii*, and *V. scophthalmi* on the surface of turbot)[1,65,66].

As a result of a detailed numerical taxonomic study of Gram-negative, oxidase-positive bacteria recovered from sharks, the dominance of vibrios was noted, with representatives including *V. harveyi* (= *V. carchariae*), and *V. alginolyticus*. Other groups included *Aeromonas*, *Photobacterium* (including *P. damsela* and *P. damsela* subsp. *piscicida*), *Alteromonas*, *Plesiomonas shigelloides*, *Moraxella*, and *Neisseria*[60].

Gill Microflora

Yellow-pigmented, Gram-negative rods, especially *Cytophaga* spp., dominate on gills[38]. Aeromonads, coryneforms, enterobacteria, Gram-positive cocci, pseudomonads, and vibrios have also been recovered from the gills of healthy juvenile rainbow trout[68].

Gills of marine fish accommodate *Achromobacter*, *Alcaligenes*, *Bacillus*, *Flavobacterium*, and *Micrococcus*[72] and yellow-pigmented bacteria, loosely associated with *Chryseobacterium-Flavobacterium-Flexibacter-Cytophaga*[73].

Microflora in the Digestive Tract

Studies on the microflora of the digestive tract have led the way in the use of culture-independent approaches[e.g., 21]. However, the bulk of the historical data stems from culturing methods, which will be discussed first. Ringø *et al.*[74] have written an excellent review on the topic.

Initially in the sac fry, only a few taxa (coryneforms and *Pseudomonas*) occur within the digestive tract[56]. It is likely that some bacteria become ingested at the yolk-sac stage, leading to the establishment

of an initial intestinal microflora[57]. In addition, it has been reported that bacterial colonisation of the digestive tract of turbot larvae coincided with the start of feeding, when the microflora was dominated by *Aeromonas* and *Vibrio*[75]. In an investigation of the intestinal microflora of larval sea bream (*Dicentrarchus labrax*) and sea bass (*Sparus aurata*), it was observed that when the larvae were fed with rotifers, there was a high incidence of *V. anguillarum*, *V. tubiashii*, and nonvibrio groups[76]. However, feeding with *Artemia* led to the recovery of mostly *V. alginolyticus*, *V. proteolyticus*, *V. harveyi*, and *V. natriegens*[76]. It was concluded from these experiments that the fluctuations in the dominant components of the microflora reflected the bacteria in the live feed. Indeed, the dominance of vibrios was not recorded until the end of the larval stage[76]. The comparative lack of diversity in larvae continues into older fish, and it has been suggested that the flora may be subjected to as-yet undescribed selective effects leading to a restricted number of taxa being present[59,77,78,79,80].

A comparatively wide range of taxa have been associated with the digestive tract of adult freshwater fish and include *Acinetobacter*, *Enterobacter*, *Escherichia*, *Klebsiella*, *Proteus*, *Serratia*[42], *Aeromonas*[42,43,48,68,81,82], *Mycoplasma*[30], *Clostridium*[42] and *Fusobacterium*[42,74]. Isolates have been identified by microplate hybridization as *A. caviae*, *A. hydrophila*, *A. jandaei*, *A. sobria*, and *A. veronii*[31]. *Alcaligenes*, *Eikenella*[4], *Bacteroides*[3,83,84], *Citrobacter freundii*[39], *Hafnia alvei*[81], *Cytophaga/Flexibacter*[68], *Bacillus*, *Listeria*, *Propionibacterium*, *Staphylococcus*[39], *Moraxella*[49], and *Pseudomonas*[4,39,68,80]. In one study involving pike perch, it was concluded that *Moraxella* and *Staphylococcus* were unique to the habitat when compared with the digestive tract of other fish species[49].

Modern phenetic and molecular-based studies, including 16S rRNA sequencing have indicated variability in the intestinal microflora of salmonids, notably rainbow trout and Atlantic salmon reflecting the fish farm of origin[15,30], with analyses revealing the dominance of the gamma subclass[15,21] (i.e., enterics, *Aeromonas*, and *Pseudomonas*) and beta subclass of Proteobacteria, and Gram-positive bacteria with a low G + C-content of the DNA (*Carnobacterium*)[15]. The approaches have permitted the recognition of potentially new taxa. For example, a 16S rRNA gene sequence with similarity to *Anaerofilum pentosovorans* has been detected[30].

In one detailed study, 41 culturable microbial phylotypes, and 39 sequences from 16S rRNA and 2 from 18S rRNA genes were retrieved from the digestive and intestinal mucus of rainbow trout and equated largely with Aeromonadaceae, Enterobacteriaceae (i.e., *Buttiauxella*, *Enterobacter*, *Hafnia*, *Pantoea*, *Plesiomonas*, and *Proteus*) and Pseudomonadaceae representatives. Intestinal contents contained *Arthrobacter*, *Bacillus*, *Carnobacterium*, *Exiguobacterium*, *Flavobacterium*, *Kokuria*, *Microbacterium*, *Micrococcus*, *Rhodococcus*, *Sporocytophaga*, and *Ultramicrobacterium*. Genomic DNA isolated from intestinal contents and mucus was used to generate 104 random clones, which were mostly affiliated with Proteobacteria (>70% of the total). Twelve sequences were retrieved from denaturing gradient gel electrophoresis analysis of the digestive tract of rainbow trout, and dominant bands were mostly related to *Clostridium*[29]. One of the outcomes of the study was the realization that *Capnocytophaga*, *Cetobacterium*, *Erwinia*, *Porphyromonas*, *Prevotella*, *Rahnella*, *Ralstonia*, *Serratia*, and *Veillonella* were recognised as occurring for the first time as culturable components of the microflora in the digestive tract of freshwater fish[29]. Using a parallel approach, the digestive tract of wild and farmed salmon from Norway and Scotland were found to be populated with *Acinetobacter junii* and a novel *Mycoplasma* phylotype, the latter of which comprised almost all, i.e., ~96%, of the microflora of the distal intestine of wild salmon[30].

The digestive tract of adult marine fish has been reported to contain *Aeromonas*[81], *Alcaligenes*[52,85], *Alteromonas*[20], *Carnobacterium*[86], *Flavobacterium*[52,85], *Micrococcus*[52], *Photobacterium*[52,85], *Pseudomonas*[59,85], *Staphylococcus*[52], and *Vibrio*[20,52,59,62,64,80,82,85], including *V. iliopiscarius*[87]. Terminal restriction fragment length polymorphism data point to a greater diversity in the posterior compared to the anterior gut in large herbivorous fish, i.e., *Kyphosus sydneyanus*[6].

Special groups, such as large (gigantobacteria) symbiotic bacteria, have been observed in the digestive tract of surgeonfish from the Red Sea and Indo-Pacific Region[88]. Also, using a methanogen-

specific nested polymerase chain reaction, methanogens have been detected in the digestive tract and faeces of flounder (*Platichthys flesus*) from the North Sea[28]. Indeed, in this study, 16S rDNA sequences revealed 97.6–99.5% similarity to the archaea representative *Methanococcoides methylutens*[28].

Lactic acid bacteria, notably carnobacteria, are common on/in fish, particularly in the digestive tract[89,90,91] with investigations highlighting the presence of *Lactococcus* (notably *L. lactis* and *L. raffinolactis*[90], *Lactobacillus*, *Aerococcus*-like bacteria, *Leuconostoc*, *Pediococcus*, *Streptococcus*, *Vagococcus*, and *Weissella*) as part of the normal microflora[92]. To date, studies have emphasised the taxonomy of the organisms[89], highlighting the presence of *Carnobacterium*[21,86,91,92,93] particularly *C. piscicola*[89,91,94] and *C. piscicola*-like bacteria[95], and their role as putative probiotics for use in aquaculture. Other lactic-acid bacteria present in the epithelial mucosa have been equated with *Lactobacillus plantarum*, *Leuconostoc mesenteroides*, and *Streptococcus* spp.[89]. In a separate investigation, *Lactobacillus*, *Enterococcus durans*, *Lactococcus*, *Vagococcus*, *C. divergens*, and *C. piscicola* were recovered from freshwater fish, notably brown trout (*Salmo trutta*), and characterised phenotypically by numerical analyses[96]. A previously undescribed species, *C. inhibens*, was recovered from the intestine of Atlantic salmon, and demonstrated antibacterial activity against fish pathogens, notably *Aeromonas salmonicida* and *Vibrio anguillarum*[97].

Diets

Aeromonads, *Bacillus*, pseudomonads, and *Staphylococcus* dominate in diets[35,55].

Eggs

Healthy eggs are populated by *Cytophaga/Flavobacterium* and, to a lesser extent, *Pseudomonas*[56,98], reflecting the organisms present in water[57].

Internal Organs

The liver and kidney of healthy turbot have been found to be populated by mostly *Pseudomonas* and *Vibrio*, including *V. fischeri*, *V. harveyi*, *V. pelagius*, and *V. splendidus*[41]. Similarly, *Shewanella* spp. have been recovered from the internal organs[99]. The reasons for the presence of some of these bacteria are unclear. Moreover, it is speculative whether or not the fish are at the earliest stage of an infection cycle.

Human Pathogens Recovered from Fish Tissue

Attention has focused on the presence of potential human pathogens in and around fish, namely *Aeromonas* spp., *Campylobacter jejuni*, *Clostridium botulinum*, *C. perfringens*, *Erysipelothrix rhusiopathiae*, *Edwardsiella tarda*, *Legionella pneumophila*, *Mycobacterium* spp., *Photobacterium damsela*, *Plesiomonas shigelloides*, *Staphylococcus aureus*, *Streptococcus iniae*, *V. cholerae*, *V. parahaemolyticus*, and *V. vulnificus*[100]. For example, *Plesiomonas shigelloides* has been cultured from the digestive tract of pike[5]. Similarly, *Staphylococcus aureus* and *V. mimicus* have been isolated repeatedly from striped bass reared in flow-through and recirculating systems[101]. *V. cholerae* was recovered from presumably healthy sharks[60]. *V. vulnificus*, enumerated by the most probable number technique with serological identification, has been found in the contents of the digestive tracts of numerous fish from the U.S. Gulf Coast[102]. In this study, a seasonal fluctuation was recorded with minimum and maximum numbers occurring in winter and April to October, respectively. Indeed, the highest populations of *V. vulnificus* (10^8 bacterial 100 g^{-1}) were associated with the gut contents of bottom-feeding fish, especially those that consumed molluscs and crustacea[102]. In contrast, the

plankton-feeding fish contained 10^5 cells of *V. vulnificus* 100 g^{-1} . Overall, it was apparent from this study that the incidence of *V. vulnificus* was comparatively uncommon in offshore fish, instead being restricted to those specimens from estuaries, i.e., closer to shore[102]. In contrast, there has not been any evidence of *Listeria monocytogenes*[5], *Salmonella*, or *Yersinia enterocolitica*[36,61].

THE ROLE OF FISH BACTERIA

Although the relative numbers and types of bacteria associated with healthy fish are interesting, it is the role of these bacteria that is of importance. However, the information is generally patchy. For a start, it is relevant to inquire whether fish-associated bacteria are active metabolically or could some be inactive-dormant-nonculturable[103]. By piecing together various data, it becomes apparent that components of the bacterial microflora of fish have been associated with numerous functions, including: (1) the production of friction-preventing polymers (bacteria on fish skin, perhaps, important for the movement of fish through the water column[104]); (2) eicosapentaenoic acid (intestinal bacteria[105]); (3) the degradation of complex molecules, including starch (amylase production by intestinal bacteria[106,107,108]), cellulose[47,109], phospholipids (intestinal bacteria[110]), proteins[111], chitin, and collagen[52,107]; and (4) the production of neuraminidase (in *Photobacterium damsela*, from the intestines of coastal fish[112]) and vitamins (e.g., vitamin B₁₂, which may be of value to the host[83,113,114,115]). Moreover by use of DNA microarrays, gnotobiotic zebrafish (*Danio rerio*) revealed the presence of 212 host genes, which were regulated by the intestinal microflora[116].

Some taxa, such as *Pseudomonas*, have been implicated as causes of fish spoilage[117,118] by the production of histamines[119,120], principally during storage of fish[72].

Thus, it is likely that bacteria are often beneficial by contributing to the nutritional processes of fish, namely by degrading complex molecules in the digestive tract[52] and by producing vitamins[83].

Luminescent Bacteria

Luminous bacteria, principally *Photobacterium*[121,122], including *P. phosphoreum* and *P. leiognathi*[27], organisms related to *P. phosphoreum* as determined by 16S rRNA sequencing[26], and *Vibrio* spp.[122], including *V. fischeri*[103,123,124], are responsible for the light-emitting properties of fish from ten families and five orders[27,125,126]. In addition, obligately symbiotic luminous bacteria that have been equated by 16S rRNA analyses as new species of *Vibrio* have been found in members of the beryciform family Anomalopidae and nine families in the lophiiform suborder Ceratioidei[27]. Generally, luminous bacteria are extracellular, and appear to be tightly arranged in tubules with communication with the exterior of the light-emitting organ[122]. A second site for luminous bacteria has been found in apogonid fish, *Siphamia permutata* and *S. cephalotes*[127]. The tubules release bacteria into the digestive tract of the host and thus into the surrounding seawater, where the released organisms are viable and culturable, and may well contribute to the planktonic microbial populations[122]. Superficially, it would seem that this work has been largely substantiated by others who have also recognised the presence of luminous bacteria, namely *Photobacterium* (*P. phosphoreum*) and *V. harveyi*, in the digestive tracts of some marine fish[128]. However, it should be emphasised that many fish without light-emitting organs have also been found to possess luminescent bacteria in their intestines[128]. Therefore, light-emitting organs are clearly not always the source of luminous bacteria in the digestive tract or, for that matter, in seawater.

The production of light by the light-emitting organ is a direct function of synergism or symbiosis between luminous bacteria and fish. There is some evidence that luminous bacteria pass from the adults to offspring[129]. In particular, it was found that offspring from spotnape ponyfish (*Leiognathus nuchalis*) eggs, which were hatched in the absence of adults, did not develop luminescence activity[129]. Conversely, juvenile fish developed bioluminescence within 48 h of contact with adults or inoculation with a homogenate of the adult light-emitting organs[129]. From this work, the inference was that

juvenile fish became infected with symbiotic luminous bacteria from the light-emitting organ of adult fish, thereby gaining the ability to become bioluminescent[129].

Luminous bacteria in the intestine appear to be involved in chitin degradation, and may therefore have a role in the digestion of complex molecules[130]. Also, some luminous bacteria have been attributed with the ability to produce histamine, and could, therefore, be involved in fish spoilage[131].

Production of Inhibitory Compounds

Some bacteria produce inhibitory compounds, particularly in the digestive tract, and may be responsible for controlling the colonisation of potential pathogens in fish[95,132]. For example a *Vibrio* sp. recovered from the intestine of a spotnape ponyfish in Japanese coastal waters inhibited the causal agent of pasteurellosis/pseudotuberculosis, i.e., *P. damsela* subsp. *piscicida*[133]. Specifically, the inhibitory compound was heat-labile and proteinaceous, with a molecular mass of <5 kDa, and was considered to be possibly a bacteriocin or a bacteriocin-like substance[133].

Similarly, bacteria were isolated and found to be capable of inhibiting growth of pathogenic *Vibrio* sp. from the digestive tract of halibut (*Hippoglossus hippoglossus*) larvae[134]. Here, the fraction of pathogen inhibitors among the total number of isolates ranged between 0–100% (first feeding) and 0–66% (weaning). All antagonists were Gram-negative rods, most of which were fermentative, and produced catalase and oxidase, being equated with *Aeromonas* and *Vibrio*[134].

Using a double agar layer method, 940 aerobic and anaerobic isolates obtained from the digestive tract of river fish, water, and sediment in Japan were examined for antagonism[84]. Some of the isolates (i.e., *Bacteroides* type A and other Bacteroidaceae representatives) from the digestive tract inhibited the target organisms, which included *A. hydrophila*, *A. salmonicida*, *E. coli*, and *Staphylococcus aureus*. The implication of the data was that these antagonistic bacteria may well influence the composition of the microflora in the digestive tract by the production of inhibitory compounds[84]. In another study by the same group, it was reported that, of 1,055 intestinal bacteria derived from 7 coastal fish in Japan, 28 isolates (2.7% of the total) inhibited the human and eel pathogen *V. vulnificus*[135]. Thus, marked inhibition was displayed by 15 isolates, comprising 11 Vibrionaceae representatives, 3 coryneforms, and 1 *Bacillus* strain NM 12; the latter demonstrated the most profound antimicrobial activity, and was therefore chosen for detailed study[135]. This revealed that one of the inhibitory compounds, which was determined to be a heat labile siderophore of <5 kDa molecular weight, inhibited the growth of 227 out of 363 (62.5% of the total) intestinal bacterial cultures derived from 7 fish[135]. Others have achieved this level of success. For example, of >400 bacteria recovered from turbot, 89 inhibited the growth of the fish pathogen *V. anguillarum*[136]. Similarly, of >400 isolates from the intestine and the external surface of farmed turbot, 28% (mostly from the digestive tract) inhibited *A. salmonicida*, *A. hydrophila*, and *V. anguillarum*[137].

Effect of Antimicrobial Compounds on the Microflora

When fish become exposed to antimicrobial compounds, there will undoubtedly be an impact on the composition of the microflora and on antibiotic resistance patterns[20,138,139,140,141,142]. This, in turn, may impact upon the transmission of antibiotic resistance, such as via R-factors[139], to other bacteria, and perhaps of significance to humans.

CONCLUSIONS

Fish possess a diverse array of bacterial taxa, often reflecting the composition of the microflora of the surrounding water. It is argued that the role of many of these fish-associated bacteria is unclear, and future work should be directed at this aspect.

REFERENCES

1. Montes, M., Perez, M.J., and Nieto, T.P. (1999) Numerical taxonomy of gram-negative, facultative anaerobic bacteria isolated from skin of turbot (*Scophthalmus maximus*) and surrounding water. *Syst. Appl. Microbiol.* **22**, 604–618.
2. Sakata, T., Sugita, H., Mitsuoka, T., Kakimoto, D., and Kadota, H. (1981) Microflora in the gastrointestinal tracts of fresh-water fish. 2. Characteristics of obligate anaerobic-bacteria in the intestines of fresh-water fish. *Bull. Jpn. Soc. Sci. Fish.* **47**, 421–427.
3. Kamei, Y., Sakata, T., and Kakimoto, D. (1985) Microflora in the alimentary tract of the *Tilapia*: characteristics and distribution of anaerobic bacteria. *J. Gen. Appl. Microbiol.* **31**, 115–124.
4. Lee, S. and Lee, Y. (1995) Identification of intestinal microflora in rainbow trout. *J. Microbiol.* **33**, 273–277.
5. Gonzalez, C.J., Lopez-Diaz, T.M., Garcia-Lopez, M.L., Prieto, M., and Otero, A. (1999) Bacterial microflora of wild brown trout (*Salmo trutta*), wild pike (*Esox lucius*), and aquacultured rainbow trout (*Oncorhynchus mykiss*). *J. Food Protect.* **62**, 1270–1277.
6. Moran, D., Turner, S.J., and Clements, J.D. (2005) Ontogenetic development of the gastrointestinal microbiota in the marine herbivorous fish *Kyphosus sydneyanus*. *Microb. Ecol.* **49**, 590–597.
7. Austin, B. and Austin, D.A. (1999) *Bacterial Fish Pathogens, Disease of Farmed and Wild Fish*. 3rd ed. Springer-Praxis, Godalming.
8. Ekman, E., Borjeson, H., and Johansson, N. (1999) *Flavobacterium psychrophilum* in Baltic salmon *Salmo salar* brood fish and their offspring. *Dis. Aquat. Org.* **37**, 159–163.
9. El-Shafai, S.A., Gijzen, H.J., Nasr, F.A., and El-Gohary, F.A. (2004) Microbial quality of tilapia reared in fecal-contaminated ponds. *Environ. Res.* **95**, 231–238.
10. Olafsen, J.A. (2001) Interactions between fish larvae and bacteria in marine aquaculture. *Aquaculture* **200**, 223–247.
11. Burr, G., Gatlin, S., and Ricke, S. (2005) Microbial ecology of the gastrointestinal tract of fish and the potential application of prebiotics and probiotics in finfish aquaculture. *J. World Aquacult. Soc.* **36**, 425–436.
12. Robertson, P.A.W., O-Dowd, C., Burrells, C., Williams, P., and Austin, B. (2000) Use of *Carnobacterium* sp. as a probiotic for Atlantic salmon (*Salmo salar* L.) and rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Aquaculture* **185**, 235–243.
13. Austin, B. and Austin, D.A. (1987) *Bacterial Fish Pathogens, Disease of Farmed and Wild Fish*. Ellis Horwood, Chichester.
14. Del Rio Rodriguez, R.E., Inglis, V., and Millar, S.D. (1997) Survival of *Escherichia coli* in the intestine of fish. *Aquacult. Res.* **28**, 257–264.
15. Spanggaard, B., Huber, I., Nielsen, J., Nielsen, T., Appel, K.F., and Gram, L. (2000) The microflora of rainbow trout intestine: a comparison of traditional and molecular identification. *Aquaculture* **182**, 1–15.
16. Eddy, S.D. and Jones, S.H. (2002) Microbiology of the summer flounder *Paralichthys dentatus* fingerling production at a marine fish hatchery. *Aquaculture* **211**, 9–28.
17. Al-Harbi, A.H. and Uddin, M.N. (2004) Seasonal variation in the intestinal bacterial flora of hybrid tilapia (*Oreochromis niloticus* x *Oreochromis aureus*) cultured in earthen ponds in Saudi Arabia. *Aquaculture* **229**, 37–44.
18. Al-Harbi, A.H. and Uddin, N. (2005) Bacterial diversity of tilapia (*Oreochromis niloticus*) cultured in brackish water in Saudi Arabia. *Aquaculture* **250**, 566–572.
19. Nedoluha, P.C., Owens, S., Russek-Cohen, E., and Westhoff, D.C. (2001) Effect of sampling method on the representative recovery of microorganisms from the surfaces of aquacultured finfish. *J. Food Protect.* **64**, 1515–1520.
20. Hansen, G.H., Strom, E., and Olafsen, J.A. (1992) Effect of different holding regimens on the intestinal microflora of herring (*Clupea harengus*) larvae. *Appl. Environ. Microbiol.* **58**, 461–470.
21. Huber, I., Spanggaard, B., Appel, K.F., Rossen, L., Nielsen, T., and Gram, L. (2004) Phylogenetic analysis and in situ identification of the intestinal microbial community of rainbow trout (*Oncorhynchus mykiss*, Walbaum). *J. Appl. Microbiol.* **96**, 117–132.
22. Hellberg, H. and Bjerkas, I. (2000) The anatomy of the oesophagus, stomach and intestine in common wolffish (*Anarhichas lupus* L.): a basis for diagnostic work and research. *Acta Vet. Scand.* **41**, 283–297.
23. Ringø, E., Lodemel, J.B., Myklebust, R., Kaino, T., Mayhew, T.M., and Olsen, R.E. (2001) Epithelium-associated bacteria in the gastrointestinal tract of Arctic charr (*Salvelinus alpinus* L.). An electron microscopical study. *J. Appl. Microbiol.* **90**, 294–300.
24. Ringø, E., Lodemel, J.B., Myklebust, R., Kaino, T., Mayhew, T.M., and Olsen, R.E. (2001) Epithelium-associated bacteria in the gastrointestinal tract of Arctic charr (*Salvelinus alpinus* L.). An electron microscopical study. *J. Appl. Microbiol.* **90**, 294–300.
25. Pettipher, G.L., Watts, Y.B., Langford, S.A., and Kroll, R.G. (1992) Preliminary evaluation of COBRA, an automated Deft instrument, for the rapid enumeration of microorganisms in cultures, raw-milk, meat and fish. *Let. Appl. Microbiol.* **14**, 206–209.
26. Haygood, M.G., Distel, D.L., and Herring, P.J. (1992) Polymerase chain-reaction and 16S-ribosomal-RNA gene-sequences from the luminous bacterial symbionts of 2 deep-sea anglerfishes. *J. Mar. Biol. Assoc. U.K.* **72**, 149–159.
27. Haygood, M.G. (1993) Light organ symbioses in fishes. *Crit. Rev. Microbiol.* **19**, 191–216.
28. van der Maarel, M.J.E.C., Sprenger, W., Haanstra, R., and Forney, L.J. (1999) Detection of methanogenic archaea in seawater particles and the digestive tract of a marine fish species. *FEMS Microbiol. Lett.* **173**, 189–194.

29. Kim, D.-H., Brunt, J.W., and Austin, B. (2006) Microbial diversity in intestinal contents and mucus in rainbow trout (*Oncorhynchus mykiss*). *J. Appl. Microbiol.*, in press.
30. Holben, W.E., Williams, P., Saarinen, M., Sarkilahi, L.K., and Apajalahti, J.H.A. (2002) Phylogenetic analysis of intestinal microflora indicates a novel *Mycoplasma* phylotype in farmed and wild salmon. *Microb. Ecol.* **44**, 175–185.
31. Sugita, H., Nakamura, T., Tanaka, K., and Deguchi, Y. (1994) Identification of *Aeromonas* species isolated from fresh-water fish with the microplate hybridization method. *Appl. Environ. Microbiol.* **60**, 3036–3038.
32. Horsley, R.W. (1973) The bacterial flora of the Atlantic salmon (*Salmo salar* L.) in relation to its environment. *J. Appl. Bacteriol.* **36**, 377–386.
33. Diler, O., Altun, S., Calikusu, F., and Diler, A. (2000) A study on qualitative and quantitative bacterial flora of the rainbow trout (*Oncorhynchus mykiss*) living in different fish farms. *Turk. J. Vet. Anim. Sci.* **24**, 251–259.
34. Gillespie, N.C. and Macrae, I.C. (1975) The bacterial flora of some Queensland fish and its ability to cause spoilage. *J. Appl. Bacteriol.* **39**, 91–100.
35. Zmyslowska, I., Lewandowska, D., Nowakowski, T., and Kozlowski, J. (2001) Occurrence of bacteria in water and in vendace (*Coregonus albula*) during rearing in tanks. *Pol. J. Environ. Stud.* **10**, 51–56.
36. Nedoluha, P.C. and Westhoff, D. (1997) Microbiological analysis of striped bass (*Morone saxatilis*) grown in a recirculating system. *J. Food Protect.* **60**, 948–953.
37. Dickinson, C.H., Austin, B., and Goodfellow, M. (1975) Quantitative and qualitative studies on the phylloplane bacteria of *Lolium perenne*. *J. Gen. Microbiol.* **91**, 157–166.
38. Trust, T.J. (1975) Bacteria associated with the gills of salmonid fishes in freshwater. *J. Appl. Bacteriol.* **38**, 225–233.
39. Apun, K., Yusof, A.M., and Jugang, K. (1999) Distribution of bacteria in tropical freshwater fish and ponds. *Int. J. Environ. Health Res.* **9**, 285–292.
40. Evelyn, T.P.T. and McDermott, L.A. (1961) Bacteriological studies of freshwater fish. 1. Isolation of aerobic bacteria from several species of Ontario fish. *Can. J. Microbiol.* **7**, 357–382.
41. Toranzo, A.E., Novoa, B., Romalde, J.L., Nunez, S., Devesa, S., Marino, E., Silva, R., Martinez, E., Figueras, A., and Barja, J.L. (1993) Microflora associated with healthy and diseased turbot (*Scophthalmus maximus*) from 3 farms in Northwest Spain. *Aquaculture* **114**, 189–202.
42. Trust, T.J. and Sparrow, R.A.H. (1974). The bacterial flora in the alimentary tract of freshwater salmonid fish. *Can. J. Microbiol.* **20**, 1219–1228.
43. Yoshimizu, M., Kimura, T., and Sakai, M. (1976) Studies on the intestinal microflora of salmonids. 1. The intestinal microflora of fish reared in freshwater and seawater. *Bull. Jpn. Soc. Sci. Fish.* **42**, 91–99.
44. Campbell, A.C. and Buswell, J.A. (1983) The intestinal microflora of farmed Dover sole (*Solea solea*) at different stages of fish development. *J. Appl. Bacteriol.* **55**, 215–223.
45. Yano, Y., Nakayama, A., and Yoshida, K. (1995) Population sizes and growth pressure responses of intestinal microfloras of deep-sea fish retrieved from the abyssal zone. *Appl. Environ. Microbiol.* **61**, 4480–4483.
46. Savas, S., Kubilay, A., and Basmaz, N. (2005) Effect of bacterial load in feeds on intestinal microflora of seabream (*Sparus aurata*) larvae and juveniles. *Israeli J. Aquacult. – Bamidgeh* **57**, 3–9.
47. Luczkovich, J.J. and Stellwag, E.J. (1993) Isolation of cellulolytic microbes from the intestinal-tract of the pinfish, *Lagodon rhomboides*. size-related changes in diet and microbial abundance. *Mar. Biol.* **116**, 381–388.
48. Yoshimizu, M., Kamiyama, K., Kimura, T., and Sakai, M. (1976) Studies on the intestinal microflora of salmonids. IV. The intestinal microflora of freshwater salmon. *Bull. Jpn. Soc. Sci. Fish.* **42**, 1281–1290.
49. Diler O. and Diler A. (1998) Quantitative and qualitative changes of the gastrointestinal microflora of pike-perch (*Stizostedion lucioperca* L., 1758) in Egirdir Lake. *Turk. J. Vet. Anim. Sci.* **22**, 325–328.
50. Maya, R., Dhevendaran, K., Mathew, A., Georgekutty, M.I., and Natarajan, P. (1995) Seasonal variations of bacteria in fish *Eetroplus suratensis* and *Eetroplus maculatus* (Pisces: Cichlidae). *Indian J. Mar. Sci.* **24**, 225–228.
51. Sakata, T., Nakaji, M., and Kakimoto, D. (1978) Microflora in the digestive tract of marine fish. 1. General characterization of the isolates from yellowtail. *Mem. Fac. Fish. Kagoshima Univ.* **27**, 65–71.
52. MacDonald, N.L., Stark, J.R., and Austin, B. (1986) Bacterial microflora in the gastro-intestinal tract of Dover sole (*Solea solea* L.), with emphasis on the possible role of bacteria in the nutrition of the host. *FEMS Microbiol. Lett.* **35**, 107–111.
53. Balasubramanian, S., Rajan, M.R., and Raj, S.P. (1992) Microbiology of fish grown in a sewage-fed pond. *Biores. Technol.* **40**, 63–66.
54. Trust, T.J. (1971) Bacterial counts of commercial fish diets. *J. Fish. Res. Bd. Can.* **28**, 1185–1189.
55. Kitao, T. and Aoki, T. (1976) Microbial flora of artificial fish diets. *Fish Pathol.* **10**, 181–185.
56. Yoshimizu, M., Kimura, T., and Sakai, M. (1980) Microflora of the embryo and the fry of salmonids. *Bull. Jpn. Soc. Sci. Fish.* **46**, 967–975.
57. Hansen, G.H. and Olafsen, J.A. (1999) Bacterial interactions in early life stages of marine cold water fish. *Microb. Ecol.* **38**, 1–26.
58. Ringø, E. and Birkbeck, T.H. (1999) Intestinal microflora of fish larvae and fry. *Aquacult. Res.* **30**, 73–93.
59. Liston, J. (1957) The occurrence and distribution of bacterial types on flatfish. *J. Gen. Microbiol.* **16**, 205–216.
60. Grimes, D.J., Jacobs, D., Swartz, D.G., Brayton, P.R., and Colwell, R.R. (1993) Numerical taxonomy of Gram-negative, oxidase-positive rods from carcharhinid sharks. *Int. J. Syst. Bacteriol.* **43**, 88–98.
61. Nedoluha, P.C. and Westhoff, D. (1997) Microbiology of striped bass grown in three aquaculture systems. *Food Microbiol.* **14**, 255–264.




62. Colwell, R.R. (1962) The bacterial flora of Puget Sound fish. *J. Appl. Bacteriol.* **25**, 147–158.
63. Pacha, R.E. and Porter, A. (1968) Characteristics of myxobacteria isolated from the surface of freshwater fish. *Appl. Microbiol.* **16**, 1901–1906.
64. Simidu, U. and Kaneko, E. (1969) Microflora of fresh and stored flatfish *Kareius bicoloratus*. *Bull. Jpn. Soc. Sci. Fish.* **35**, 77–82.
65. Austin, B. (1982) Taxonomy of bacteria isolated from a coastal, marine fish-rearing unit. *J. Appl. Bacteriol.* **53**, 253–268.
66. Austin, B. (1983) Bacterial microflora associated with a coastal, marine fish-rearing unit. *J. Mar. Biol. Assoc. U. K.* **63**, 583–592.
67. Allen, D.A., Austin, B., and Colwell, R.R. (1983) Numerical taxonomy of bacterial isolates associated with a freshwater fishery. *J. Gen. Microbiol.* **129**, 2043–2062.
68. Nieto, T.P., Toranzo, A.E., and Barja, J.L. (1984) Comparison between the bacterial flora associated with fingerling rainbow trout cultured in two different hatcheries in the north-west of Spain. *Aquaculture* **42**, 193–206.
69. Gonzalez, C.J., Santos, J.A., Garcia-Lopez, M.L., and Otero A. (2000) Psychrobacters and related bacteria in freshwater fish. *J. Food Protect.* **63**, 315–321.
70. Gonzalez, C.J., Santos, J.A., Garcia-Lopez, M.L., Gonzalez, N., and Otero, A. (2001) Mesophilic aeromonads in wild and aquacultured freshwater fish. *J. Food Protect.* **64**, 687–691.
71. Christensen, P.J. (1977) The history, biology and taxonomy of the *Cytophaga* group. *Can. J. Microbiol.* **23**, 1599–1653.
72. Shewan, J.M. (1961) The microbiology of sea-water fish. In *Fish as Food*. Vol. 1. Borgstrom, G., Ed. Academic Press, New York. pp. 487–560.
73. Mudarris, M. and Austin, B. (1988) Quantitative and qualitative studies of the bacterial microflora of turbot, *Scophthalmus maximus* L., gills. *J. Fish Biol.* **32**, 223–229.
74. Ringø, E., Strøm, E., and Tabachek, J.A. (1995) Intestinal microflora: a review. *Aquacult. Res.* **26**, 773–789.
75. Munro, P.D., Barbour, A., and Birkbeck, T.H. (1994) Comparison of the gut bacterial-flora of start-feeding larval turbot reared under different conditions. *J. Appl. Bacteriol.* **77**, 560–566.
76. Grisez, L., Reyniers, J., Verdonck, L., Swings, J., and Ollevier, F. (1997) Dominant intestinal microflora of sea bream and sea bass larvae, from two hatcheries, during larval development. *Aquaculture* **155**, 387–399.
77. Shrivastava, K.P. and Floodgate, G.D. (1966) Studies on the intestinal microflora of the dab. *J. Mar. Biol. Assoc. India* **8**, 1–7.
78. Newman, J.T., Cosenza, B.J., and Buck, J.D. (1972) Aerobic microflora of the bluefish intestine. *J. Fish. Res. Bd. Can.* **29**, 33–336.
79. Sera, H., Ishida, Y., and Kadota, H. (1972) Bacterial microflora in the digestive tract of marine fish. IV. Effect of H⁺ concentration and gastric juices on the indigenous bacteria. *Bull. Jpn. Soc. Sci. Fish.* **38**, 859–863.
80. Yoshimizu, M. and Kimura, T. (1976) Study on the intestinal microflora of salmonids. *Fish Pathol.* **10**, 243–259.
81. Ugajin, M. (1979) Studies on the taxonomy of major microflora on the intestinal contents of salmonids. *Bull. Jpn. Soc. Sci. Fish.* **45**, 721–731.
82. Sakata, T., Okabayashi, J., and Kakimoto, D. (1980) Variations in the intestinal microflora of *Tilapia* reared in fresh and seawater. *Bull. Jpn. Soc. Sci. Fish.* **46**, 313–317.
83. Sugita, H., Miyajima, C., and Deguchi, Y. (1991) The vitamin-B₁₂-producing ability of the intestinal microflora of fresh-water fish. *Aquaculture* **92**, 267–276.
84. Sugita H., Shibuya K., Hanada H., and Deguchi Y. (1997) Antibacterial abilities of intestinal microflora of the river fish. *Fish. Sci.* **63**, 378–383.
85. LeaMaster, B.R., Walsh, W.A., Brock, J.A., and Fujioka, R.S. (1997) Cold stress-induced changes in the aerobic heterotrophic gastrointestinal tract bacterial flora of red hybrid tilapia. *J. Fish Biol.* **50**, 770–778.
86. Ringø, E., Wesmajervi, M.S., Bendiksen, H.R., Berg, A., Olsen, R.E., Johnsen, T., Mikkelsen, H., Seppola, M., Strom, E., and Holzapfel, W. (2001) Identification and characterization of carnobacteria isolated from fish intestine. *Syst. Appl. Microbiol.* **24**, 183–191.
87. Onarheim, A.M., Wiik, R., Burghardt, J., and Stackerbrandt, E. (1994) Characterization and identification of 2 *Vibrio* species indigenous to the intestine of fish in cold sea-water - description of *Vibrio iliopiscarius* sp nov. *Syst. Appl. Microbiol.* **17**, 370–379.
88. Fishelson, L. (1999) Polymorphism in giantobacterial symbionts in the guts of surgeonfish (Acanthuridae: Teleostei). *Mar. Biol.* **133**, 345–351.
89. Ringø, E., Bendiksen, H.R., Gausen, S.J., Sundsfjord, A., and Olsen, R.E. (1998) The effect of dietary fatty acids on lactic acid bacteria associated with the epithelial mucosa and from faecalia of Arctic charr, *Salvelinus alpinus* (L.). *J. Appl. Microbiol.* **85**, 855–864.
90. Hagi, T., Tanaka, D., Iwamura, Y., and Hoshino, T. (2004) Diversity and seasonal changes in lactic acid bacteria in the intestinal tract of cultured freshwater fish. *Aquaculture* **234**, 335–346.
91. Seppola, M., Olsen, R.E., Sandaker, E., Kanapathipillai, P., Holzapfel, W., and Ringø, E. (2006) Random amplification of polymorphic DNA (RAPD) typing of carnobacteria isolated from hindgut chamber and large intestine of Atlantic cod (*Gadus morhua* L.). *Syst. Appl. Microbiol.* **29**, 131–137.
92. Ringø, E. (2005) Lactic acid bacteria in fish and fish farming. In *Lactic Acid-Bacteria Microbiological and Functional Aspects*. Salminen, S., Von Wright, A., and Ouwehand, A., Eds. Marcel Dekker, Basel, Switzerland. pp.

- 581–610.
93. Ringø, E., Seppola, M., Berg, A., Olsen, R.E., Schillinger, U., and Holzapfel, W. (2002) Characterization of *Carnobacterium divergens* strain 6251 isolated from intestine of Arctic charr (*Salvelinus alpinus* L.). *Syst. Appl. Microbiol.* **25**, 120–129.
 94. Baya, A.M., Toranzo, A.E., Lupiani, B., Li, T., Roberson, B.S., and Hetrick, F.M. (1991) Biochemical and serological characterization of *Carnobacterium* spp isolated from farmed and natural-populations of striped bass and catfish. *Appl. Environ. Microbiol.* **57**, 3114–3120.
 95. Ringø, E., Bendiksen, H.R., Wesmajervi, M.S., Olsen, R.E., Jansen, P.A., and Mikkelsen, H. (2000) Lactic acid bacteria associated with the digestive tract of Atlantic salmon (*Salmo salar* L.). *J. Appl. Microbiol.* **89**, 317–322.
 96. Gonzalez, C.J., Encinas, J.P., Garcia-Lopez, M.L., and Otero, A. (2000) Characterization and identification of lactic acid bacteria from freshwater fishes. *Food Microbiol.* **17**, 383–391.
 97. Joborn, A., Dorsch, M., Olsson, J.C., Westerdahl, A., and Kjelleberg, S. (1999) *Carnobacterium inhibens* sp nov., isolated from the intestine of Atlantic salmon (*Salmo salar*). *Int. J. Syst. Bacteriol.* **49**, 1891–1898.
 98. Bell, G.R., Hoskins, G.E., and Hodgkiss, W. (1971) Aspects of the characterization, identification and ecology of the bacterial flora associated with the surface of stream-incubating Pacific salmon (*Oncorhynchus*) eggs. *J. Fish. Res. Bd. Can.* **28**, 1511–1525.
 99. Decostere, A., Haesebrouck, F., Devriese, L., and Ducatelle, R. (1996) Identification and pathogenic significance of *Shewanella* sp. from pond fish. *Vlaams Diergeneesk. Tijdschr.* **65**, 82–85.
 100. Novotny, L., Dvorska, L., Lorencova, A., Beran, V., and Pavlik, I. (2004) Fish: a potential source of bacterial pathogens for human beings. *Vet. Med.* **49**, 343–358.
 101. Nedoluha, P.C. and Westhoff, D. (1995) Microbiological analysis of striped bass (*Morone saxatilis*) grown in flow-through tanks. *J. Food Protect.* **58**, 1363–1368.
 102. DePaola, A., Capers, G.M., and Alexander, D. (1994) Densities of *Vibrio vulnificus* in the intestines of fish from the U.S. Gulf-Coast. *Appl. Environ. Microbiol.* **60**, 984–988.
 103. Ruby, E.G. and Lee, K.H. (1998) The *Vibrio fischeri*-*Euprymna scolopes* light organ association: current ecological paradigms. *Appl. Environ. Microbiol.* **64**, 805–812.
 104. Sar, N. and Rosenberg, E. (1989) Fish skin bacteria - production of friction-reducing polymers. *Mar. Ecol.* **17**, 27–38.
 105. Yazawa, K., Araki, K., Watanabe, K., Ishikawa, C., Inoue, A., Kondo, K., Watabe, S., and Hashimoto, K. (1988) Eicosapentaenoic acid productivity of the bacteria isolated from fish intestines. *Nippon Suisan Gakkaishi* **54**, 1835–1838.
 106. Sugita, H., Kawasaki, J., Kumazawa, J., and Deguchi, Y. (1996) Production of amylase by the intestinal bacteria of Japanese coastal animals. *Lett. Appl. Microbiol.* **23**, 174–178.
 107. Syvokiene, J. and Mickeniene, L. (1999) Microorganisms in the digestive tract of fish as indicators of feeding condition and pollution. *ICES J. Mar. Sci.* **56**, 147–149.
 108. Izvekova, G.I. (2005) Activity of carbohydrases of symbiotic microflora and their role in processes of digestion of fish and their parasitizing cestodes (on the example of pike and *Triaenophorus nodulosus*). *J. Evol. Biochem. Physiol.* **41**, 406–411.
 109. Saha, S., Roy, R.N., Sen, S.K., and Ray, A.K. (2006) Characterization of cellulase-producing bacteria from the digestive tract of tilapia, *Oreochromis mossambica* (Peters) and grass carp, *Ctenopharyngodon idella* (Valenciennes). *Aquacult. Res.* **37**, 380–388.
 110. Henderson, R.J. and Millar, R.M. (1998) Characterization of lipolytic activity associated with a *Vibrio* species of bacterium isolated from fish intestines. *J. Mar. Biotechnol.* **6**, 168–173.
 111. Izvekova, G.I. (2006) Hydrolytic activity of enzymes produced by symbiotic microflora and its role in digestion processes of bream and its intestinal parasite *Caryophyllaeus laticeps* (Cestoda, Caryophyllidae). *Biol. Bull.* **33**, 287–292.
 112. Sugita, H., Shinagawa, Y., and Okano, R. (2000) Neuraminidase-producing ability of intestinal bacteria isolated from coastal fish. *Lett. Appl. Microbiol.* **31**, 10–13.
 113. Kashiwada, K. and Teshima, S. (1966) Studies on the production of B vitamins by intestinal bacteria of fish. I. Nicotinic acid, pantothenic acid and vitamin B₁₂ in carp. *Bull. Jpn. Soc. Sci. Fish.* **32**, 961.
 114. Kashiwada, K., Teshima, S., and Kanazawa, A. (1970). Studies on the production of B vitamins by intestinal bacteria of fish. V. Evidence of the production of vitamin B₁₂ by microorganisms in the intestinal canal of carp. *Bull. Jpn. Soc. Sci. Fish.* **36**, 421–424.
 115. Kashiwada, K., Kanazawa, A., and Teshima, S. (1971) Studies on the production of B vitamins by intestinal bacteria. VI. Production of folic acid by intestinal bacteria of carp. *Mem. Fac. Fish. Kagoshima Univ.* **20**, 185–189.
 116. Rawls, J.F., Samuel, B.S., and Gordon, J.I. (2004) Gnotobiotic zebrafish reveal evolutionarily conserved responses to the gut microbiota. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 4596–4601.
 117. Gillespie, N.C. (1981) A numerical taxonomic study of *Pseudomonas*-like bacteria isolated from fish in southeastern Queensland and their association with spoilage. *J. Appl. Bacteriol.* **50**, 29–44.
 118. Malle, P. (1994) Bacterial microflora in marine fish and evaluation of spoilage. *Recl. Med. Vet.* **170**, 147–157.
 119. Yoguchi, R., Okuzumi, M., and Fujii, T. (1990) Seasonal-variation in number of halophilic histamine-forming bacteria on marine fish. *Nippon Suisan Gakkaishi* **56**, 1473–1479.
 120. Kim, S.H., Field, K.G., Chang, D.S., Wei, C.I., and An, H.J. (2001) Identification of bacteria crucial to histamine accumulation in Pacific mackerel during storage. *J. Food Protect.* **64**, 1556–1564.

121. Reichelt, J.L. and Baumann, P. (1977) Taxonomy of the marine luminous bacteria. *Arch. Mikrobiol.* **94**, 282–330.
122. Haygood, M.G. and Distel, D.L. (1993) Bioluminescent symbionts of flashlight fishes and deep-sea anglerfishes form unique lineages related to the genus *Vibrio*. *Nature (London)* **363**, 154–156.
123. Fitzgerald, D.M. (1977) Classification of luminous bacteria from the light organ of the Australian pinecone fish *Cleidopus gloriamans*. *Arch. Mikrobiol.* **112**, 153–156.
124. Dunlap, P.V. and Callahan, S.M. (1993) Characterization of a periplasmic 3'/5'-cyclic nucleotide phosphodiesterase gene, Cpdp, from the marine symbiotic bacterium *Vibrio-fischeri*. *J. Bacteriol.* **175**, 4615–4624.
125. Haygood, M.G., Tebo, B.M., and Nealson, K.H. (1984) Luminous bacteria of a monocentrid fish (*Monocentris japonicus*) and 2 anomalopid fishes (*Photoblepharon palpebratus* and *Kryptophanaron alfredi*) - population sizes and growth within the light organs, and rates of release into the seawater. *Mar. Biol.* **78**, 249–254.
126. Fukasawa, S., Suda, T., and Kubota, S. (1988) Identification of luminous bacteria isolated from the light organ of the fish, *Acropoma japonicum*. *Agric. Biol. Chem.* **52**, 285–286.
127. Fishelson, L., Gon, O., Goren, M., and Ben-David-Zaslow, R. (2005) The oral cavity and bioluminescent organs of the cardinal fish species *Siphamia permutata* and *S. cephalotes* (Perciformes, Apogonidae). *Mar. Biol.* **147**, 603–609.
128. Makemson, J.C. and Hermosa, G.V. (1999) Luminous bacteria cultured from fish guts in the Gulf of Oman. *Luminescence* **14**, 161–168.
129. Wada, M., Azuma, N., Mizuno, N., and Kurokura, H. (1999). Transfer of symbiotic luminous bacteria from parental *Leiognathus nuchalis* to their offspring. *Mar. Biol.* **135**, 683–687.
130. Ramesh, A. and Venugopalan, V.K. (1989) Role of luminous bacteria in chitin degradation in the intestine of fish. *MIRCEN-J. Appl. Microbiol. Biotechnol.* **5**, 55–59.
131. Morii, H., Cann, D.C., Taylor, L.Y., and Murray, C.K. (1986) Formation of histamine by luminous bacteria isolated from scombroid fish. *Bull. Jpn. Soc. Sci. Fish.* **52**, 2135–2141.
132. Makridis, P., Martins, S., Tsalavouta, M., Dionisio, L.C., Kotoulas, G., Magoulas, A., and Dinis, M.T. (2005) Antimicrobial activity in bacteria isolated from Senegalese sole, *Solea senegalensis*, fed with natural prey. *Aquacult. Res.* **36**, 1619–1627.
133. Sugita, H., Matsuo, N., Hirose, Y., Iwato, M., and Deguchi, Y. (1997) *Vibrio* sp. Strain NM 10, isolated from the intestine of a Japanese coastal fish, has an inhibitory effect against *Pasteurella piscicida*. *Appl. Environ. Microbiol.* **63**, 4986–4989.
134. Bergh, Ø. (1995) Bacteria associated with early-life stages of halibut, *Hippoglossus hippoglossus* L, inhibit growth of a pathogenic *Vibrio* sp. *J. Fish Dis.* **18**, 31–40.
135. Sugita, H., Hirose, Y., Matsuo, N., and Deguchi, Y. (1998) Production of the antibacterial substance by *Bacillus* sp. strain NM 12, an intestinal bacterium of Japanese coastal fish. *Aquaculture* **165**, 269–280.
136. Olsson, J.C., Westerdahl, A., Conway, P.L., and Kjelleberg, S. (1992) Intestinal colonization potential of turbot (*Scophthalmus maximus*)- and dab (*Limanda limanda*)-associated bacteria with inhibitory effects against *Vibrio anguillarum*. *Appl. Environ. Microbiol.* **58**, 551–556.
137. Westerdahl, A., Olsson, J.C., Kjelleberg, S., and Conway, P.L. (1991) Isolation and characterization of turbot (*Scophthalmus maximus*)-associated bacteria with inhibitory effects against *Vibrio anguillarum*. *Appl. Environ. Microbiol.* **57**, 2223–2228.
138. Austin, B. and Al-Zahrani, A.M.J. (1988) The effect of antimicrobial compounds on the gastrointestinal microflora of rainbow-trout, *Salmo-gairdneri* Richardson. *J. Fish Biol.* **33**, 1–14.
139. Alvarez, J.D., Austin, B., Alvarez, A.M., and Arguro, C.P. (2001) Antimicrobial resistance of vibrios isolated from fish and marine shrimp in Venezuela. *Rev. Cient.-Facul. Cien. Vet.* **64**, 139–148.
140. Cabello, F.C. (2004) Antibiotics and aquaculture in Chile: implications for humans and animal health. *Rev. Med. Chile* **132**, 1001–1006.
141. Moffitt, C.M. and Mobin, S.M.A. (2006) Profile of microflora of the posterior intestine of Chinook salmon before, during, and after administration of rations with and without erythromycin. *North Am. J. Aquacult.* **68**, 176–185.
142. Pedersen, A. and Dalsgaard, A. (2003) Antimicrobial resistance of intestinal *Aeromonas* spp. and *Enterococcus* spp. in fish cultured in integrated broiler-fish farms in Thailand. *Aquaculture* **219**, 71–82.
143. Austin, B. (2002) The bacterial microflora of fish. *TheScientificWorldJOURNAL* **2**, 558-572. DOI 10.1100/tsw.2002.137

This article should be cited as follows:

Austin, B. (2006) The bacterial microflora of fish, revised. *TheScientificWorldJOURNAL* **6**, 931–934. DOI 10.1100/tsw.2006.181.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

