

The Basic (G'/G) -Expansion Method for the Fourth Order Boussinesq Equation

Hasibun Naher, Farah Aini Abdullah

School of Mathematical Sciences, Universiti Sains Malaysia, Penang, Malaysia

Email: hasibun06tasauf@gmail.com, farahaini@usm.my

Received June 26, 2012; revised July 23, 2012; accepted August 1, 2012

ABSTRACT

The (G'/G) -expansion method is simple and powerful mathematical tool for constructing traveling wave solutions of nonlinear evolution equations which arise in engineering sciences, mathematical physics and real time application fields. In this article, we have obtained exact traveling wave solutions of the nonlinear partial differential equation, namely, the fourth order Boussinesq equation involving parameters via the (G'/G) -expansion method. In this method, the general solution of the second order linear ordinary differential equation with constant coefficients is implemented. Further, the solitons and periodic solutions are described through three different families. In addition, some of obtained solutions are described in the figures with the aid of commercial software Maple.

Keywords: The (G'/G) -Expansion Method; The Fourth Order Boussinesq Equation; Traveling Wave Solutions; Nonlinear Partial Differential Equations

1. Introduction

The complex physical phenomena are depicted of nonlinear evolution equations (NLEEs) which are widely used as models. One of the fundamental problems is to establish their traveling wave solutions. As a result, the attention of searching traveling wave solutions of NLEEs is enhancing and has now become a hot topic to researchers. At present time, various methods have been established to obtain exact solutions. For example, the Hirota's bilinear transformation method [1,2], the Backlund transformation method [3,4], the truncated Painleve expansion method [5,6], the inverse scattering method [7], the weierstrass elliptic function method [8], the homogeneous balance method [9], the Jacobi elliptic function expansion method [10-13], the generalized Riccati equation method [14], the tanh-coth method [15-19], the F-expansion method [20,21], the direct algebraic method [22], the Cole-Hopf transformation method [23], the Exp-function method [24-30] and others [31-36].

Recently, Wang *et al.* [37] presented a widely used method to construct traveling wave solutions of different nonlinear partial differential equations (PDEs), called the basic (G'/G) -expansion method. In addition, in this method $u(\xi) = \sum_{i=0}^m a_i (G'/G)^i$ is executed as traveling wave solutions, where $a_m \neq 0$. Later on, many researchers

established exact traveling wave solutions of various nonlinear PDEs via this method. For example, Feng *et al.* [38] studied the well-known Kolmogorov-Petrovskii - Piskunov equation to obtain analytical solutions by using the same method. Naher *et al.* [39] constructed abundant traveling wave solutions of the higher-order Caudrey-Dodd-Gibson equation via this powerful method. In [40], Abazari and Abazari concerned about the same method for obtaining hyperbolic, trigonometric and rational function solutions of Hirota-Ramani equation. Zayed [41] investigated some nonlinear evolution equations in mathematical physics to establish exact solutions by applying this method. Jabbari *et al.* [42] obtained some solutions of the coupled Higgs equation and the Miccari system via the same method while Ozis and Aslan [43] implemented the (G'/G) -expansion method to construct traveling wave solutions of the Kawahara type equations using symbolic computation. In [44], Gepreel studied some nonlinear PDEs with variable coefficients in mathematical physics by applying the same method for constructing exact solutions. Elagan *et al.* [45] executed this method to obtain innovative solutions of the generalized FitzHugh-Nagumo equation. Borhanifar and Moghanlu [46] established some solutions for the Zhiber-Sabat equation and other related equations via the same method whilst Borhanifar and Abazari [47] investigated two generalized form of Burgers equation for constructing ge-

neral solutions by using the (G'/G) -expansion method. In [48], Wang *et al.* constructed explicit solutions of the generalized (2+1)-dimensional Zakharov-Kuznetsov equation via this method and so on.

Many researchers studied the fourth order Boussinesq equation to construct analytical solutions by using different methods. For instance, Zhang [49] implemented the F-expansion method to investigate this equation for obtaining traveling wave solutions. In [50], Wazwaz employed the extended tanh method to construct exact solutions of the same equation.

The importance of this work is, the fourth order Boussinesq equation is considered to construct new exact traveling wave solutions including solitons, periodic and rational solutions by applying the basic (G'/G) -expansion method.

2. The Basic (G'/G) -Expansion Method

Suppose the general nonlinear partial differential equation:

$$F(u, u_t, u_x, u_{xt}, u_{tt}, u_{xx}, \dots) = 0, \tag{1}$$

where $u = u(x, t)$ is an unknown function, F is a polynomial in $u(x, t)$ and the subscripts stand for the partial derivatives.

The main steps of the basic (G'/G) -expansion method [37] are as follows:

Step 1. Consider the traveling wave transformation:

$$u(x, t) = A(\psi), \quad \psi = x - Ht, \tag{2}$$

where H is the wave speed. Now using Equation (2), Equation (1) is converted into an ordinary differential equation for $A(\psi)$:

$$Q(A, A', A'', A''', \dots) = 0, \tag{3}$$

where the superscripts indicate the ordinary derivatives with respect to ψ .

Step 2. According to possibility, Equation (3) can be integrated term by term one or more times, yields constant(s) of integration. The integral constant may be zero, for simplicity.

Step 3. Suppose that the traveling wave solution of Equation (3) can be expressed in the form [37]:

$$A(\psi) = \sum_{i=0}^n b_i (G'/G)^i \tag{4}$$

with $G = G(\psi)$ satisfies the second order linear ODE:

$$G'' + \lambda G' + \mu G = 0, \tag{5}$$

where $b_i (i = 0, 1, 2, \dots, n)$, λ and μ are constants.

Step 4. To determine the integer n , substituting Equation (4) along with Equation (5) into Equation (3) and then taking the homogeneous balance between the highest

order nonlinear terms and the highest order derivatives appearing in Equation (3).

Step 5. Substituting Equations (4) and (5) into Equation (3) with the value of n obtained in Step 4. Equating the coefficients of $(G'/G)^r, (r = 0, 1, 2, \dots)$, then setting each coefficient to zero, we obtain a set of algebraic equations for $b_i (i = 0, 1, 2, \dots, n), H, \lambda$ and μ .

Step 6. Solve the system of algebraic equations which are obtained in Step 5 with the aid of algebraic software Maple and we obtain values for $b_i (i = 0, 1, 2, \dots, n), H, \lambda$ and μ . Then, substitute obtained values in Equation (4) along with Equation (5) with the value of n , we can obtain the traveling wave solutions of Equation (1).

3. Applications of the Method

In this section, the fourth order Boussinesq equation involving parameters is investigated to establish traveling wave solutions including three different families by applying the (G'/G) -expansion method.

3.1. The Fourth Order Boussinesq Equation

In this work, we consider the fourth order Boussinesq equation followed by Wazwaz [50]:

$$u_{tt} - \alpha^2 u_{xx} - \beta(u^2)_{xx} + u_{xxxx} = 0. \tag{6}$$

Boussinesq established Equation (6) to illustrate the propagation of long waves in shallow water. This equation also arises in other physical applications, for example, ion sound waves in plasma, nonlinear lattice waves and in vibrations in nonlinear string. Further, the details of this equation can be found in references [49-54].

Equation (6) is integrable, therefore, integrating with respect ψ twice and setting the constants of integration equal to zero, yields:

$$(H^2 - \alpha^2)A - \beta A^2 + A'' = 0. \tag{7}$$

Taking the homogeneous balance between A^2 and A'' in Equation (7), we obtain $n = 2$.

Therefore, the solution of Equation (7) is of the form:

$$A(\psi) = b_0 + b_1(G'/G) + b_2(G'/G)^2, \tag{8}$$

where b_0, b_1 and b_2 are constants to be determined.

Substituting Equation (8) together with Equation (5) into the Equation (7), the left-hand side of Equation (7) is converted into a polynomial of $(G'/G)^r, (r = 0, 1, 2, \dots)$. According to Step 5, collecting all terms with the same power of (G'/G) . After that, setting each coefficient of the resulted polynomial to zero, yields a set of algebraic equations (for simplicity, which are not presented) for $b_0, b_1, b_2, H, \lambda$ and μ .

Solving the system of obtained algebraic equations

with the help of algebraic software Maple, we obtain two different values.

Case 1.

$$b_0 = \frac{6\mu}{\beta}, b_1 = \frac{6\lambda}{\beta}, b_2 = \frac{6}{\beta}, \tag{9}$$

$$H = \pm\sqrt{\alpha^2 - (\lambda^2 - 4\mu)},$$

where α, β, λ and μ are free parameters.

Case 2.

$$b_0 = \frac{2\mu + \lambda^2}{\beta}, b_1 = \frac{6\lambda}{\beta}, b_2 = \frac{6}{\beta}, \tag{10}$$

$$H = \pm\sqrt{\alpha^2 + (\lambda^2 - 4\mu)},$$

where α, β, λ and μ are free parameters.

Substituting the general solution Equation (5) into Equation (8), we obtain three different families of traveling wave solutions of Equation (7):

Family 3.1.1. Hyperbolic function solutions:

when $\lambda^2 - 4\mu > 0$, we obtain

$$A(\psi) = b_2 \left(\frac{-\lambda}{2} + \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \frac{U \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi + V \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi}{U \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi + V \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi} \right)^2 + b_1 \left(\frac{-\lambda}{2} + \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \frac{U \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi + V \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi}{U \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi + V \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi} \right) + b_0. \tag{11}$$

If U and V are taken as specific values, various known solutions can be rediscovered.

Family 3.1.2. Trigonometric function solutions:

when $\lambda^2 - 4\mu < 0$, we obtain

$$A(\psi) = b_2 \left(\frac{-\lambda}{2} + \frac{1}{2}\sqrt{4\mu - \lambda^2} \frac{-U \sin \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi + V \cos \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi}{U \cos \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi + V \sin \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi} \right)^2 + b_1 \left(\frac{-\lambda}{2} + \frac{1}{2}\sqrt{4\mu - \lambda^2} \frac{-U \sin \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi + V \cos \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi}{U \cos \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi + V \sin \frac{1}{2}\sqrt{4\mu - \lambda^2} \psi} \right) + b_0. \tag{12}$$

If U and V are taken as specific values, various known solutions can be rediscovered.

Family 3.1.3. Rational function solution:

when $\lambda^2 - 4\mu = 0$, we obtain

$$A(\psi) = b_0 + b_1 \left(\frac{-\lambda}{2} + \frac{V}{U + V\psi} \right) + b_2 \left(\frac{-\lambda}{2} + \frac{V}{U + V\psi} \right)^2. \tag{13}$$

Family 3.1.1.1. Hyperbolic function solutions:

Substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), yields the hyperbolic function solution Equation (11), our traveling wave solutions become respectively (if $U = 0$ but $V \neq 0$):

$$A_1(\psi) = \frac{3(\lambda^2 - 4\mu)}{2\beta} \left(\left(\coth \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi \right)^2 - 1 \right),$$

where

$$\psi = x \mp \left(\sqrt{\alpha^2 - (\lambda^2 - 4\mu)} \right) t.$$

$$A_2(\psi) = \frac{(\lambda^2 - 4\mu)}{2\beta} \left(3 \left(\coth \frac{1}{2}\sqrt{\lambda^2 - 4\mu} \psi \right)^2 - 1 \right),$$

where

$$\psi = x \mp \left(\sqrt{\alpha^2 + (\lambda^2 - 4\mu)} \right) t.$$

Again, substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), yields the hyperbolic function solution Equation (11), we obtain following exact solutions respectively (if $V = 0$ but $U \neq 0$):

$$A_3(\psi) = \frac{3(\lambda^2 - 4\mu)}{2\beta} \left(\left(\tanh \frac{1}{2} \sqrt{\lambda^2 - 4\mu} \psi \right)^2 - 1 \right).$$

$$A_4(\psi) = \frac{(\lambda^2 - 4\mu)}{2\beta} \left(3 \left(\tanh \frac{1}{2} \sqrt{\lambda^2 - 4\mu} \psi \right)^2 - 1 \right).$$

Moreover, substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), we obtain the hyperbolic function solution Equation (11), our obtained wave solutions become respectively (if $U \neq 0, U > V$):

$$A_5(\psi) = \frac{3(\lambda^2 - 4\mu)}{2\beta} \left(\left(\tanh \left(\frac{1}{2} \sqrt{\lambda^2 - 4\mu} \psi + \psi_0 \right) \right)^2 - 1 \right)$$

where $\psi_0 = \tanh^{-1}(V/U)$.

$$A_6(\psi) = \frac{(\lambda^2 - 4\mu)}{2\beta} \left(3 \left(\tanh \frac{1}{2} \sqrt{\lambda^2 - 4\mu} \psi + \psi_0 \right)^2 - 1 \right)$$

where $\psi_0 = \tanh^{-1}(V/U)$.

Family 3.1.2.1. Trigonometric function solutions:

Substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), yields the trigonometric function solution Equation (12), we obtain following solutions respectively (if $U = 0$ but $V \neq 0$):

$$A_7(\psi) = \frac{3(4\mu - \lambda^2)}{2\beta} \left(\left(\cot \frac{1}{2} \sqrt{4\mu - \lambda^2} \psi \right)^2 + 1 \right),$$

where $\psi = x \mp \left(\sqrt{\alpha^2 + (4\mu - \lambda^2)} \right) t$.

$$A_8(\psi) = \frac{(4\mu - \lambda^2)}{2\beta} \left(3 \left(\cot \frac{1}{2} \sqrt{4\mu - \lambda^2} \psi \right)^2 + 1 \right),$$

where $\psi = x \mp \left(\sqrt{\alpha^2 - (4\mu - \lambda^2)} \right) t$.

Also, substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), yields the trigonometric function solution Equation (12), our solutions become respectively (if $V = 0$ but $U \neq 0$):

$$A_9(\psi) = \frac{3(4\mu - \lambda^2)}{2\beta} \left(\left(\tan \frac{1}{2} \sqrt{4\mu - \lambda^2} \psi \right)^2 + 1 \right).$$

$$A_{10}(\psi) = \frac{(4\mu - \lambda^2)}{2\beta} \left(3 \left(\tan \frac{1}{2} \sqrt{4\mu - \lambda^2} \psi \right)^2 + 1 \right).$$

Furthermore, substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), yields the trigonometric function solution

Equation (12), our obtained traveling wave solutions become respectively (if $U \neq 0, U > V$):

$$A_{11}(\psi) = \frac{3(4\mu - \lambda^2)}{2\beta} \left(\left(\tan \frac{1}{2} \sqrt{4\mu - \lambda^2} \psi - \psi_0 \right)^2 + 1 \right),$$

where $\psi_0 = \tan^{-1}(V/U)$.

$$A_{12}(\psi) = \frac{(4\mu - \lambda^2)}{2\beta} \left(3 \left(\tan \frac{1}{2} \sqrt{\lambda^2 - 4\mu} \psi - \psi_0 \right)^2 + 1 \right),$$

where $\psi_0 = \tan^{-1}(V/U)$.

Family 3.1.3.1. Rational function solutions:

Substituting Equations (9) and (10) together with the general solution Equation (5) into the Equation (8), we obtain the rational function solution Equation (13), and our wave solutions become respectively (if $\lambda^2 - 4\mu = 0$):

$$A_{13}(\psi) = \frac{3}{2\beta} \left(\left(\frac{2V}{U + V\psi} \right)^2 - (\lambda^2 - 4\mu) \right),$$

where $\psi = x \mp \left(\sqrt{\alpha^2 - (\lambda^2 - 4\mu)} \right) t$.

$$A_{14}(\psi) = \frac{1}{2\beta} \left(3 \left(\frac{2V}{U + V\psi} \right)^2 - (\lambda^2 - 4\mu) \right),$$

where $\psi = x \mp \left(\sqrt{\alpha^2 + (\lambda^2 - 4\mu)} \right) t$.

4. Results and Discussion

It is worth declaring that some of our obtained solutions are in good agreement with already published results which are presented in **Table 1**. Moreover, some of obtained exact solutions are described in **Figures 1-12**.

Beyond this table, we obtain new exact traveling wave solutions $A_1, A_3, A_5, A_6, A_7, A_9, A_{11}, A_{12}, A_{13},$ and A_{14} ,




Figure 1. Solutions solution for $\lambda = 3, \mu = 1, \alpha = 3, \beta = 1$.




Figure 2. Periodic solution for $\lambda = 6, \mu = 8, \alpha = 12, \beta = 9$.




Figure 6. Solitons solution for $\lambda = 4, \mu = 5, \alpha = 11, \beta = 9$.




Figure 3. Solitons solution for $\lambda = 3, \mu = 3, \alpha = 9, \beta = 5$.




Figure 7. Solitons solution for $\lambda = 4, \mu = 4, \alpha = 1.5, \beta = 5 \times 10^{-11}, U = 0.75, V = 24$.




Figure 4. Solitons solution for $\lambda = 4, \mu = 4, \alpha = 0.5, \beta = 1, U = 0.5, V = 1$.




Figure 8. Periodic solution for $\lambda = 3, \mu = 3, \alpha = 16, \beta = 15$.




Figure 5. Solitons solution for $\lambda = 3, \mu = 2, \alpha = 7, \beta = 2$.




Figure 9. Periodic solution for $\lambda = 8, \mu = 16, \alpha = 5 \times 10^{-6}, \beta = 0.125, U = 5, V = 9$.




Figure 10. Solitons solution for $\lambda = 5, \mu = 7, \alpha = 15, \beta = 0.5$.




Figure 11. Solitons solution for $\lambda = 5, \mu = 7, \alpha = 7, \beta = 1$.




Figure 12. Solitons solution for $\lambda = 6, \mu = 9, \alpha = 1, \beta = 5 \times 10^{-11}, U = 0.75, V = 24$.

which are not being established in the previous literature.

5. Conclusion

In this article, the basic (G'/G) -expansion method with second order linear ordinary differential equation is implemented to investigate the fourth order Boussinesq equation involving parameters. The obtained solutions are presented through the hyperbolic functions, the trigonometric functions and the rational functions. Further, it is important mentioning that some of our solutions are coincided with published results and others have not been stated in earlier literature. The solutions show that this method is reliable and straightforward solution method to obtain exact solutions of nonlinear evolution equations. Therefore, this powerful method can be effectively used to construct new exact traveling wave solutions of diffe-

Table 1. Comparison between Wazwaz [50] solutions and newly obtained solutions.

Wazwaz [50] solutions	New solutions
1) If $c = 2, a = 1$ and $b = 3$ solution Equation (23) (from section 3) becomes: $u_2(x, t) = \frac{1}{2} \left(3 \coth^2 \frac{\sqrt{3}}{2} (x - 2t) - 1 \right).$	1) If $\mu = \frac{1}{4}, \lambda = 2, \alpha = 1, \beta = 3, A_2(\psi) = u_2(x, t)$, solution becomes: $u_2(x, t) = \frac{1}{2} \left(3 \coth^2 \frac{\sqrt{3}}{2} (x - 2t) - 1 \right).$
2) If $c = 3, a = 1$ and $b = 4$ solution Equation (22) (from section 3) becomes: $u_1(x, t) = 3 \tanh^2 \sqrt{2} (x - 3t) - 1.$	2) If $\mu = 2, \lambda = 4, \alpha = 1, \beta = 4, A_4(\psi) = u_1(x, t)$, solution becomes: $u_1(x, t) = 3 \tanh^2 \sqrt{2} (x - 3t) - 1.$
3) If $c = 4, a = 5$ and $b = \frac{9}{2}$ solution Equation (26) (from section 3) becomes: $u_5(x, t) = 3 \cot^2 \frac{3}{2} (x - 4t) + 1.$	3) If $\mu = \frac{10}{4}, \lambda = 1, \alpha = 5, \beta = \frac{9}{2}, A_8(\psi) = u_5(x, t)$, solution becomes: $u_5(x, t) = 3 \cot^2 \frac{3}{2} (x - 4t) + 1.$
4) If $c = 3, a = 5$ and $b = 8$ solution Equation (25) (from section 3) becomes: $u_4(x, t) = 3 \tan^2 2(x - 3t) + 1.$	4) If $\mu = 5, \lambda = 2, \alpha = 5, \beta = 8, A_{10}(\psi) = u_4(x, t)$, solution becomes: $u_4(x, t) = 3 \tan^2 2(x - 3t) + 1.$

rent nonlinear partial differential equations.

6. Acknowledgements

The authors would like to express their thanks to the referee(s) for their valuable comments and suggestions.

REFERENCES

- [1] R. Hirota, "Exact Solution of the KdV Equation for Multiple Collisions of Solutions," *Physics Review Letters*, Vol. 27, 1971, pp. 1192-1194.
[doi:10.1103/PhysRevLett.27.1192](https://doi.org/10.1103/PhysRevLett.27.1192)
- [2] J. Yan, "Soliton Resonances of the Nonisospectral Modified Kadomtsev-Petviashvili Equation," *Applied Mathematics*, Vol. 2, No. 6, 2011, pp. 685-693.
[doi:10.4236/am.2011.26090](https://doi.org/10.4236/am.2011.26090)
- [3] M. R. Mimura, "Backlund Transformation," Springer, Berlin, 1978.
- [4] C. Rogers and W. F. Shadwick, "Backlund Transformations," Academic Press, New York, 1982.
- [5] J. Weiss, M. Tabor and G. Carnevale, "The Painleve Property for Partial Differential Equations," *Journal of Mathematical Physics*, Vol. 24, No. 3, 1982, pp. 522-526.
[doi:10.1063/1.525721](https://doi.org/10.1063/1.525721)
- [6] S. L. Zhang, B. Wu and S. Y. Lou, "Painleve Analysis and Special Solutions of Generalized Broer-Kaup Equations," *Physics Letters A*, Vol. 300, No. 1, 2002, pp. 40-48.
[doi:10.1016/S0375-9601\(02\)00688-6](https://doi.org/10.1016/S0375-9601(02)00688-6)
- [7] M. J. Ablowitz and P. A. Clarkson, "Solitons, Nonlinear Evolution Equations and Inverse Scattering Transform," Cambridge University Press, Cambridge, 1991.
[doi:10.1017/CBO9780511623998](https://doi.org/10.1017/CBO9780511623998)
- [8] N. A. Kudryashov, "Exact Solutions of the Generalized Kuramoto-Sivashinsky Equation," *Physics Letters A*, Vol. 147, No. 5-6, 1990, pp. 287-291.
[doi:10.1016/0375-9601\(90\)90449-X](https://doi.org/10.1016/0375-9601(90)90449-X)
- [9] M. L. Wang, Y. B. Zhou and Z. B. Li, "Application of a Homogeneous Balance Method to Exact Solutions of Nonlinear Equations in Mathematical Physics," *Physics Letters A*, Vol. 216, 1996, pp. 67-75.
[doi:10.1016/0375-9601\(96\)00283-6](https://doi.org/10.1016/0375-9601(96)00283-6)
- [10] S. Liu, Z. Fu, S. Liu and Q. Zhao, "Jacobi Elliptic Function Expansion Method and Periodic Wave Solutions of Nonlinear Wave Equations," *Physics Letters A*, Vol. 289, 2001, pp. 69-74.
[doi:10.1016/S0375-9601\(01\)00580-1](https://doi.org/10.1016/S0375-9601(01)00580-1)
- [11] Z. Yan, "Abundant Families of Jacobi Elliptic Function Solutions of the (2+1)-Dimensional Integrable Davey-Stewartson-Type Equation via a New Method," *Chaos, Solitons and Fractals*, Vol. 18, No. 2, 2003, pp. 299-309.
[doi:10.1016/S0960-0779\(02\)00653-7](https://doi.org/10.1016/S0960-0779(02)00653-7)
- [12] Y. Chen and Q. Wang, "Extended Jacobi Elliptic Function Rational Expansion Method and Abundant Families of Jacobi Elliptic Function Solutions to (1+1)-Dimensional Dispersive Long Wave Equation," *Chaos, Solitons and Fractals*, Vol. 24, No. 3, 2005, pp. 745-757.
[doi:10.1016/j.chaos.2004.09.014](https://doi.org/10.1016/j.chaos.2004.09.014)
- [13] C. Xiang, "Jacobi Elliptic Function Solutions for (2+1) Dimensional Boussinesq and Kadomtsev-Petviashvili Equation," *Applied Mathematics*, Vol. 2, No. 11, 2011, pp. 1313-1316.
[doi:10.4236/am.2011.211183](https://doi.org/10.4236/am.2011.211183)
- [14] Z. Yan and H. Q. Zhang, "New Explicit Solitary Wave Solutions and Periodic Wave Solutions for Whitham-Broer-Kaup Equation in Shallow Water," *Physics Letters A*, Vol. 285, No. 5-6, 2001, pp. 355-362.
[doi:10.1016/S0375-9601\(01\)00376-0](https://doi.org/10.1016/S0375-9601(01)00376-0)
- [15] W. Malfliet, "Solitary Wave Solutions of Nonlinear Wave Equations," *American Journal of Physics*, Vol. 60, No. 7, 1992, pp. 650-654.
[doi:10.1119/1.17120](https://doi.org/10.1119/1.17120)
- [16] A. M. Wazwaz, "The tanh-coth Method for Solitons and Kink Solutions for Nonlinear Parabolic Equations," *Applied Mathematics and Computation*, Vol. 188, 2007, pp. 1467-1475.
[doi:10.1016/j.amc.2006.11.013](https://doi.org/10.1016/j.amc.2006.11.013)
- [17] A. Bekir and A. C. Cevikel, "Solitary Wave Solutions of Two Nonlinear Physical Models by tanh-coth Method," *Communications in Nonlinear Science and Numerical Simulation*, Vol. 14, No. 5, 2009, pp. 1804-1809.
[doi:10.1016/j.cnsns.2008.07.004](https://doi.org/10.1016/j.cnsns.2008.07.004)
- [18] F. Fan, "Extended tanh-Function Method and Its Applications to Nonlinear Equations," *Physics Letters A*, Vol. 277, No. 4-5, 2000, pp. 212-218.
[doi:10.1016/S0375-9601\(00\)00725-8](https://doi.org/10.1016/S0375-9601(00)00725-8)
- [19] N. H. Abdel-All, M. A. Abdel Razeq and A. K. Seddeek, "Expanding the tanh-Function Method for Solving Nonlinear Equations," *Applied Mathematics*, Vol. 2, No. 9, 2011, pp. 1096-1104.
[doi:10.4236/am.2011.29151](https://doi.org/10.4236/am.2011.29151)
- [20] M. L. Wang and X. Z. Li, "Applications of F-Expansion to Periodic Wave Solutions for a New Hamiltonian Amplitude Equation," *Chaos, Solitons and Fractals*, Vol. 24, No. 5, 2005, pp. 1257-1268.
[doi:10.1016/j.chaos.2004.09.044](https://doi.org/10.1016/j.chaos.2004.09.044)
- [21] M. A. Abou, "The Extended F-Expansion Method and Its Applications for a Class of Nonlinear Evolution Equation," *Chaos, Solitons and Fractals*, Vol. 31, 2007, pp. 95-104.
[doi:10.1016/j.chaos.2005.09.030](https://doi.org/10.1016/j.chaos.2005.09.030)
- [22] A. A. Soliman and H. A. Abdo, "New Exact Solutions of Nonlinear Variants of the RLW, the PHI-Four and Boussinesq Equations Based on Modified Extended Direct Algebraic Method," *International Journal of Nonlinear Science*, Vol. 7, No. 3, 2009, pp. 274-282.
- [23] A. H. Salas, and C. A. Gomez, "Application of the Cole-Hopf Transformation for Finding Exact Solutions to Several Forms of the Seventh-Order KdV Equation," *Mathematical Problems in Engineering*, 2010, Article ID 194329, 14 p.
- [24] J. H. He and X. H. Wu, "Exp-Function Method for Nonlinear Wave Equations," *Chaos Solitons and Fractals*, Vol. 30, No. 3, 2006, pp. 700-708.
[doi:10.1016/j.chaos.2006.03.020](https://doi.org/10.1016/j.chaos.2006.03.020)
- [25] H. Naher, F. A. Abdullah and M. A. Akbar, "New Traveling Wave Solutions of the Higher Dimensional Nonlinear Partial Differential Equation by the Exp-Function Method," *Journal of Applied Mathematics*, 2012, Article ID: 575387, 14 p.
- [26] S. T. Mohyud-din, M. A. Noor and K. I. Noor, "Exp-Function Method for Traveling Wave Solutions of Modi-

- fied Zakharov-Kuznetsov Equation,” *Journal of King Saud University*, Vol. 22, No. 4, 2010, pp. 213-216.
[doi:10.1016/j.jksus.2010.04.015](https://doi.org/10.1016/j.jksus.2010.04.015)
- [27] H. Naher, F. Abdullah and M. A. Akbar, “The Exp-Function Method for New Exact Solutions of the Nonlinear Partial Differential Equations,” *International Journal of the Physical Sciences*, Vol. 6, No. 29, 2011, pp. 6706-6716.
- [28] A. Yildirim and Z. Pinar, “Application of the Exp-Function Method for Solving Nonlinear Reaction-Diffusion Equations Arising in Mathematical Biology,” *Computers & Mathematics with Applications*, Vol. 60, No. 7, 2010, pp. 1873-1880. [doi:10.1016/j.camwa.2010.07.020](https://doi.org/10.1016/j.camwa.2010.07.020)
- [29] I. Aslan, “Application of the Exp-Function Method to Nonlinear Lattice Differential Equations for Multi-Wave and Rational Solutions,” *Mathematical Methods in the Applied Sciences*, Vol. 60, No. 7, 2011, pp. 1707-1710.
[doi:10.1002/mma.1476](https://doi.org/10.1002/mma.1476)
- [30] A. Bekir and A. Boz, “Exact Solutions for Nonlinear Evolution Equations Using Exp-Function Method,” *Physics Letters A*, Vol. 372, No. 4, 2008, pp. 1619-1625.
[doi:10.1016/j.physleta.2007.10.018](https://doi.org/10.1016/j.physleta.2007.10.018)
- [31] Q. Liu and R. Xu, “Periodic Solutions of a Cohen-Grossberg-Type BAM Neural Networks with Distributed Delays and Impulses,” *Journal of Applied Mathematics*, 2012, Article ID 643418, 17 p.
[doi:10.1155/2012/643418](https://doi.org/10.1155/2012/643418)
- [32] D. Feng and K. Li, “On Exact Traveling Wave Solutions for (1+1) Dimensional Kaup-Kupershmidt Equation,” *Applied Mathematics*, Vol. 2, No. 6, 2011, pp. 752-756.
[doi:10.4236/am.2011.26100](https://doi.org/10.4236/am.2011.26100)
- [33] K. A. Gepreel, S. Omran and S. K. Elagan, “The Travelling Wave Solutions for Some Nonlinear PDEs in Mathematical Physics,” *Applied Mathematics*, Vol. 2, No. 3, 2011, pp. 343-347. [doi:10.4236/am.2011.23040](https://doi.org/10.4236/am.2011.23040)
- [34] M. Noor, K. Noor, A. Waheed and E. A. Al-Said, “An Efficient Method for Solving System of Third-Order Nonlinear Boundary Value Problems,” *Mathematical Problems in Engineering*, 2011, Article ID: 250184, 14 p.
[doi:10.1155/2011/250184](https://doi.org/10.1155/2011/250184)
- [35] J. Wang, L. Wang and K. Yang, “Some New Exact Travelling Wave Solutions for the Generalized Benny-Luke (GBL) Equation with Any Order,” *Applied Mathematics*, Vol. 3, 2012, pp. 309-314.
[doi:10.4236/am.2012.34046](https://doi.org/10.4236/am.2012.34046)
- [36] J. F. Alzaidy, “Exact Travelling Wave Solutions of Nonlinear PDEs in Mathematical Physics,” *Applied Mathematics*, Vol. 3, No. 7, 2012, pp. 738-745.
[doi:10.4236/am.2012.37109](https://doi.org/10.4236/am.2012.37109)
- [37] M. Wang, X. Li and J. Zhang, “The (G'/G) -Expansion Method and Travelling Wave Solutions of Nonlinear Evolution Equations in Mathematical Physics,” *Physics Letters A*, Vol. 372, No. 4, 2008, pp. 417-423.
[doi:10.1016/j.physleta.2007.07.051](https://doi.org/10.1016/j.physleta.2007.07.051)
- [38] J. Feng, W. Li and Q. Wan, “Using (G'/G) -Expansion Method to Seek Traveling Wave Solution of Kolmogorov-Petrovskii-Piskunov Equation,” *Applied Mathematics and Computation*, Vol. 217, 2011, pp. 5860-5865.
[doi:10.1016/j.amc.2010.12.071](https://doi.org/10.1016/j.amc.2010.12.071)
- [39] H. Naher, F. A. Abdullah and M. A. Akbar, “The (G'/G) -Expansion Method for Abundant Traveling Wave Solutions of Caudrey-Dodd-Gibbon Equation,” *Mathematical Problems in Engineering*, 2011, Article ID: 218216, 11 p.
- [40] R. Abazari and R. Abazari, “Hyperbolic, Trigonometric and Rational Function Solutions of Hirota-Ramani Equation via (G'/G) -Expansion Method,” *Mathematical Problems in Engineering*, 2011, Article ID: 424801, 11 p.
- [41] E. M. E. Zayed, “Traveling Wave Solutions for Higher Dimensional Nonlinear Evolution Equations Using the (G'/G) -Expansion Method,” *Journal of Applied Mathematics & Informatics*, Vol. 28, No. 1-2, 2010, pp. 383-395.
- [42] A. Jabbari, H. Kheiri and A. Bekir, “Exact Solutions of the Coupled Higgs Equation and the Miccari System Using He’s Semi-Inverse Method and (G'/G) -Expansion Method,” *Computers and Mathematics with Applications*, Vol. 62, 2011, pp. 2177-2186.
[doi:10.1016/j.camwa.2011.07.003](https://doi.org/10.1016/j.camwa.2011.07.003)
- [43] T. Ozis and I. Aslan, “Application of the (G'/G) -Expansion Method to Kawahara Type Equations Using Symbolic Computation,” *Applied Mathematics and Computation*, Vol. 216, 2010, pp. 2360-2365.
[doi:10.1016/j.amc.2010.03.081](https://doi.org/10.1016/j.amc.2010.03.081)
- [44] K. A. Gepreel, “Exact Solutions for Nonlinear PDEs with the Variable Coefficients in Mathematical Physics,” *Journal of Information and Computing Science*, Vol. 6, No. 1, 2011, pp. 003-014.
- [45] S. K. Elagan, M. Sayed and Y. S. Hamed, “An Innovative Solutions for the Generalized FitzHugh-Nagumo Equation by Using the Generalized (G'/G) -Expansion Method,” *Applied Mathematics*, Vol. 2, 2011, pp. 470-474.
[doi:10.4236/am.2011.24060](https://doi.org/10.4236/am.2011.24060)
- [46] A. Borhanifar and A. Z. Moghanlu, “Application of the (G'/G) -Expansion Method for the Zhiber-Sabat Equation and Other Related Equations,” *Mathematical and Computer Modelling*, Vol. 54, 2011, pp. 2109-2116.
[doi:10.1016/j.mcm.2011.05.020](https://doi.org/10.1016/j.mcm.2011.05.020)
- [47] A. Borhanifar and R. Abazari, “General Solution of Two Generalized Form of Burgers Equation by Using the (G'/G) -Expansion Method,” *Applied Mathematics*, Vol. 3, 2012, pp. 158-168. [doi:10.4236/am.2012.32025](https://doi.org/10.4236/am.2012.32025)
- [48] G. Wang, X. Liu and Y. Zhang, “New Explicit Solutions of the Generalized (2+1)-Dimensional Zakharov-Kuznetsov Equation,” *Applied Mathematics*, Vol. 3, No. 6, 2012, pp. 523-527. [doi:10.4236/am.2012.36079](https://doi.org/10.4236/am.2012.36079)
- [49] H. Zhang, “New Exact Travelling Wave Solutions for Some Nonlinear Evolution Equations,” *Chaos, Solitons and Fractals*, Vol. 26, No. 3, 2005, pp. 921-925.
[doi:10.1016/j.chaos.2005.01.035](https://doi.org/10.1016/j.chaos.2005.01.035)
- [50] A. M. Wazwaz, “New Travelling Wave Solutions to the

- Boussinesq and Klein-Gordon Equations,” *Communications in Nonlinear Science and Numerical Simulation*, Vol. 13, No. 5, 2008, pp. 889-901.
[doi:10.1016/j.cnsns.2006.08.005](https://doi.org/10.1016/j.cnsns.2006.08.005)
- [51] Y. Guo and S. Lai, “New Exact Solutions for an (n+1)-Dimensional Generalized Boussinesq Equation,” *Nonlinear Analysis*, Vol. 72, 2010, pp. 2863-2873.
[doi:10.1016/j.na.2009.11.030](https://doi.org/10.1016/j.na.2009.11.030)
- [52] Sirendaoreji, “A New Auxiliary Equation and Exact Travelling Wave Solutions of Nonlinear Equations,” *Physics Letters A*, Vol. 356, No. 2, 2006, pp. 124-130.
[doi:10.1016/j.physleta.2006.03.034](https://doi.org/10.1016/j.physleta.2006.03.034)
- [53] A. Parker and J. M. Dye, “Boussinesq-Type Equations and Switching Solitons,” *Proceedings of Institute of Mathematics of NAS of Ukraine*, Vol. 43, No. 1, 2002, pp. 344-351.
- [54] Z. Yan, “Constructing Exact Solutions for Two-Dimensional Nonlinear Dispersion Boussinesq Equation. II: Solitary Patterns Solutions,” *Chaos, Solitons & Fractals*, Vol. 18, No. 4, 2003, pp. 869-880.
[doi:10.1016/S0960-0779\(03\)00059-6](https://doi.org/10.1016/S0960-0779(03)00059-6)