

THE BERRY-ESSEEN BOUND FOR CHARACTER RATIOS

QI-MAN SHAO AND ZHONG-GEN SU

(Communicated by Richard C. Bradley)

ABSTRACT. Let λ be a partition of n chosen from the Plancherel measure of the symmetric group S_n , let $\chi^\lambda(12)$ be the irreducible character of the symmetric group parameterized by λ evaluated on the transposition (12) , and let $\dim(\lambda)$ be the dimension of the irreducible representation parameterized by λ . Fulman recently obtained the convergence rate of $O(n^{-s})$ for any $0 < s < \frac{1}{2}$ in the central limit theorem for character ratios $\frac{(n-1)}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)}$ by developing a connection between martingale and character ratios, and he conjectures that the correct speed is $O(n^{-1/2})$. In this paper we confirm the conjecture via a refinement of Stein's method for exchangeable pairs.

1. INTRODUCTION AND MAIN RESULT

Let $n \geq 1$, let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_p)$ be a partition of n , i.e., $\lambda_1 + \lambda_2 + \dots + \lambda_p = n$, and write simply $\lambda \vdash n$. Denote by $\dim(\lambda)$ the number of standard Young tableaux associated with the shape λ . By the Robinson-Schensted-Knuth correspondence [18], we have

$$\sum_{\lambda \vdash n} \dim(\lambda)^2 = n!.$$

Thus we produce the so-called Plancherel measure

$$P(\{\lambda\}) = \frac{\dim(\lambda)^2}{n!}.$$

Recently there has been intensive interest in the statistical properties of partitions chosen from the Plancherel measure. We refer the reader to the surveys by Aldous and Diaconis [1], Defit [4] and the seminal papers of Borodin, Okounkov and Olshanski [2], Johansson [14], and Okounkov and Pandharipande [16] for details.

It turns out that the Plancherel measure can also be regarded as a probability measure on the irreducible representation of the symmetric group S_n . Observe that the irreducible representation of the symmetric group S_n is parameterized by partitions λ of n and $\dim(\lambda)$ is just the corresponding dimension of the irreducible representation.

Received by the editors September 28, 2004 and, in revised form, February 4, 2005.

2000 *Mathematics Subject Classification*. Primary 60F05; Secondary 05E10, 60C05.

Key words and phrases. Berry-Esseen bound, character ratio, Plancherel measure, Stein's method.

The first author was supported in part by Grant R-1555-000-035-112 at the National University of Singapore.

The second author was supported in part by NFS of China (No. 10371109).

Let $\chi^\lambda(12)$ be the irreducible character parameterized by λ evaluated on the transposition (12). The quantity $\frac{\chi^\lambda(12)}{\dim(\lambda)}$ is called a character ratio and is crucial for analyzing the convergence rate of the random walk on the symmetric group generated by transpositions in Diaconis and Shahshahani [5]. In fact, Diaconis and Shahshahani prove that the eigenvalues for this random walk are the character ratios $\frac{\chi^\lambda(12)}{\dim(\lambda)}$, each occurring with multiplicity $\dim(\lambda)^2$. Character ratios also play an essential role in work on the moduli spaces of curves; see Eskin and Okounkov [6], Okounkov and Pandharipande [16].

Kerov [15] first studies the asymptotic behavior for character ratios and outlines the proof of the following central limit theorem:

$$\frac{(n-1)}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)} \xrightarrow{d} N(0, 1).$$

A full proof of the result appears in Ivanov and Olshanski [13]; see also Hora [12] for another proof. A more probabilistic approach to Kerov's central limit theorem has recently been given by Fulman [7], in which a Stein's method for exchangeable pairs is used to obtain for all $n \geq 2, z \in \mathbb{R}$,

$$\left| P\left(\frac{(n-1)}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)} \leq z\right) - \Phi(z) \right| \leq 40.1n^{-1/4}$$

where $\Phi(z)$ is the standard normal distribution function.

More recently Fulman [8] developed a connection between martingales and character ratios of the symmetric group, and thereby improved the above speed of convergence to $O(n^{-s})$ for any $s < \frac{1}{2}$. He also conjectured that the correct speed is $O(n^{-1/2})$.

The main aim of this note is to confirm the following conjecture.

Theorem 1.1. *We have*

$$(1.1) \quad \sup_z \left| P\left(\frac{(n-1)}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)} \leq z\right) - \Phi(z) \right| \leq An^{-1/2}$$

where A is an absolute constant.

The proof of Theorem 1.1 will be given in Section 2. The main technique is a refinement of Stein's method for exchangeable pairs (see Theorem 2.1 below). Recall that two random variables W, W^* are called exchangeable if (W, W^*) and (W^*, W) have the same joint distribution function. In order to apply Stein's approach for exchangeable pairs, one needs to construct a W^* such that (W, W^*) is exchangeable and the difference $W - W^*$ is small. Fulman [7] uses the theory of harmonic functions on Bratelli diagrams and shows how it can be applied to generate a natural exchangeable pair (W, W^*) . The basic idea is to use a reversible Markov chain on the set of partitions of size n whose stationary distribution is the Plancherel measure. Let λ^* be obtained from λ by one step in the chain, and then set $(W, W^*) = (W(\lambda), W^*(\lambda))$. This construction also has the merit of being applicable to more general groups [9] and to measures arising from symmetric functions [10].

In the setting of Theorem 1.1, we let $W = \frac{(n-1)}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)}$. Let $\text{parents}(\lambda, \mu)$ denote the set of partitions above both λ, μ in the Young lattice (this set has size 0 or 1

unless $\lambda = \mu$), i.e.,

$$\text{parents}(\lambda, \mu) = \#\{\tau : \lambda \nearrow \tau, \mu \nearrow \tau\}.$$

Define

$$W^*(\lambda) = W(\lambda^*)$$

where, given λ , the partition λ^* is μ with probability

$$J(\lambda, \mu) = \frac{\dim(\mu)|\text{parents}(\lambda, \mu)|}{(n + 1) \dim(\lambda)}.$$

Then it follows from Proposition 2.1 of Fulman [7] that (W, W^*) is an exchangeable pair.

2. PROOF

The proof is based on the following refinement of Stein’s result [20] for exchangeable pairs.

Theorem 2.1. *Let (W, W^*) be an exchangeable pair of real-valued random variables such that*

$$(2.2) \quad E^W(W^*) = (1 - \tau)W$$

with $0 < \tau < 1$, where $E^W(W^*)$ denotes the conditional expected value of W^* given W . Assume $E(W^2) \leq 1$. Then for any $a > 0$,

$$(2.3) \quad \begin{aligned} & \sup_z |P(W \leq z) - \Phi(z)| \\ & \leq \sqrt{E\left(1 - \frac{1}{2\tau}E^W(\Delta^2)\right)^2} + \frac{0.41a^3}{\tau} + 1.5a + \frac{1}{2\tau}E\Delta^2 I_{\{|\Delta| \geq a\}}, \end{aligned}$$

where $\Delta = W - W^*$.

If Δ is bounded, say $|\Delta| \leq a_0$ for a constant a_0 , then (2.3) reduces to

$$\sup_z |P(W \leq z) - \Phi(z)| \leq \sqrt{E\left(1 - \frac{1}{2\tau}E^W(\Delta^2)\right)^2} + \frac{0.41a_0^3}{\tau} + 1.5a_0.$$

Similar results for the bounded case were obtained by Rinott and Rotar [17] and Rinott and Goldstein [11].

Theorem 1.1 is an easy consequence of Theorem 2.1.

Proof of Theorem 1.1. By [7], we can choose

$$\tau = \frac{2}{n + 1}, \quad \sqrt{E\left(1 - \frac{1}{2\tau}E^W(\Delta^2)\right)^2} \leq \frac{\sqrt{3}}{2n^{1/2}}.$$

Let $a = 4e\sqrt{2}n^{-1/2}$. Then, by the proof of Proposition 4.6 in [7],

$$\begin{aligned} E\Delta^2 I_{\{|\Delta| > a\}} & \leq 8P(|\Delta| > a) \\ & \leq 8P(\max(\lambda_1, \lambda'_1) > 2e\sqrt{n}) \\ & \leq 16e^{-2e\sqrt{n}}, \end{aligned}$$

and hence

$$\begin{aligned} \frac{1}{2\tau}E\Delta^2 I_{\{|\Delta| > a\}} & \leq 4(n + 1)e^{-2e\sqrt{n}} \\ & \leq n^{-1/2}4(n + 1)^{3/2}e^{-2e\sqrt{n}} \\ & \leq 0.05n^{-1/2}. \end{aligned}$$

Therefore, by Theorem 2.1,

$$\begin{aligned} & \sup_z |P(W \leq z) - \Phi(z)| \\ & \leq \frac{\sqrt{3}}{2n^{1/2}} + 0.205(n+1)(4e\sqrt{2})^3 n^{-3/2} + 4e\sqrt{2}n^{-1/2} + 0.05n^{-1/2} \\ & \leq An^{-1/2}, \end{aligned}$$

where A is an absolute constant. \square

We remark that if one uses

$$P(\lambda_1 \geq k) \leq \binom{n}{k}/k!$$

for $1 \leq k \leq n$ (see Lemma 1.4.1 in [19]) and chooses $a = \delta n^{-1/2}$ with $\delta > 0$ properly, then the constant A can be reduced to 150.

Now we turn to the proof of Theorem 2.1.

Proof of Theorem 2.1. For any measurable function f with $E\{|W| + 1\}|f(W)| < \infty$, exchangeability and (2.2) imply

$$\begin{aligned} 0 &= E\{(W - W^*)(f(W) + f(W^*))\} \\ &= 2E\{f(W)(W - W^*)\} + E\{(W - W^*)(f(W^*) - f(W))\} \\ &= 2\tau E\{Wf(W)\} - E\{(W - W^*)(f(W) - f(W^*))\}, \end{aligned}$$

and hence

$$(2.4) \quad E\{Wf(W)\} = \frac{1}{2\tau} E\{(W - W^*)(f(W) - f(W^*))\}.$$

Now let $f = f_z$ be the solution of the following Stein equation:

$$(2.5) \quad f'_z(x) - xf_z(x) = I_{\{x \leq z\}} - \Phi(z).$$

It is known (see [20, p.22]) that f is given by

$$f_z(x) = \begin{cases} \sqrt{2\pi}e^{x^2/2}\Phi(x)[1 - \Phi(z)] & \text{if } x \leq z, \\ \sqrt{2\pi}e^{x^2/2}\Phi(z)[1 - \Phi(x)] & \text{if } x \geq z, \end{cases}$$

satisfying

$$(2.6) \quad |xf_z(x)| \leq 1, \quad 0 < f_z(x) \leq \sqrt{2\pi}/4,$$

$$(2.7) \quad |f'_z(x)| \leq 1, \quad |f'_z(x) - f'_z(y)| \leq 1,$$

$$(2.8) \quad |(x+u)f_z(x+u) - xf_z(x)| \leq (|x| + \sqrt{2\pi}/4)|u|$$

for all real x , y , and u . For the proofs of the above inequalities, we refer to [20, p.23] for (2.6) and the first inequality of (2.7), and to Chen and Shao [3] for the second inequality of (2.7). (2.8) is a consequence of (2.6), (2.7) and the mean value theorem.

By (2.5), we have

$$\begin{aligned}
 P(W \leq z) - \Phi(z) &= Ef'_z(W) - EWf_z(W) \\
 &= Ef'_z(W) - \frac{1}{2\tau}E\{(W - W^*)(f_z(W) - f_z(W^*))\} \\
 &= E\left\{f'_z(W)\left(1 - \frac{1}{2\tau}\Delta^2\right)\right\} \\
 &\quad - \frac{1}{2\tau}E\{\Delta(f_z(W) - f_z(W - \Delta) - \Delta f'_z(W))\} \\
 (2.9) \qquad \qquad &:= J_1 + J_2.
 \end{aligned}$$

It follows from (2.6) that

$$\begin{aligned}
 |J_1| &= \left|E\left\{f'_z(W)\left(1 - \frac{1}{2\tau}E^W(\Delta^2)\right)\right\}\right| \\
 &\leq E\left|1 - \frac{1}{2\tau}E^W(\Delta^2)\right| \\
 (2.10) \qquad &\leq \sqrt{E\left(1 - \frac{1}{2\tau}E^W(\Delta^2)\right)^2}.
 \end{aligned}$$

To bound J_2 , write

$$\begin{aligned}
 &E\{\Delta(f_z(W) - f_z(W - \Delta) - \Delta f'_z(W))\} \\
 &= E\left\{\Delta \int_{-\Delta}^0 (f'_z(W + t) - f'_z(W))dt\right\} \\
 &= E\left\{\Delta I_{\{|\Delta|>a\}} \int_{-\Delta}^0 (f'_z(W + t) - f'_z(W))dt\right\} \\
 &\quad + E\left\{\Delta I_{\{|\Delta|\leq a\}} \int_{-\Delta}^0 (f'_z(W + t) - f'_z(W))dt\right\} \\
 (2.11) \qquad &:= J_{2,1} + J_{2,2}.
 \end{aligned}$$

By (2.7),

$$(2.12) \qquad |J_{2,1}| \leq E\Delta^2 I_{\{|\Delta|>a\}}.$$

Using (2.5) again, we have

$$\begin{aligned}
 J_{2,2} &= E\left\{\Delta I_{\{|\Delta|\leq a\}} \int_{-\Delta}^0 ((W + t)f_z(W + t) - Wf_z(W))dt\right\} \\
 &\quad + E\left\{\Delta I_{\{|\Delta|\leq a\}} \int_{-\Delta}^0 (I_{\{W+t\leq z\}} - I_{\{W\leq z\}})dt\right\} \\
 (2.13) \qquad &:= J_{2,2,1} + J_{2,2,2}.
 \end{aligned}$$

By (2.8),

$$\begin{aligned}
 |J_{2,2,1}| &\leq E\left\{\Delta I_{\{|\Delta|\leq a\}} \int_{-\Delta}^0 (|W| + \sqrt{2\pi}/4)|t|dt\right\} \\
 &\leq E\left\{0.5|\Delta|^3 I_{\{|\Delta|\leq a\}} (|W| + \sqrt{2\pi}/4)\right\} \\
 &\leq 0.5a^3(\sqrt{2\pi}/4 + E|W|) \\
 (2.14) \qquad &\leq 0.5a^3(\sqrt{2\pi}/4 + 1) \leq 0.82a^3.
 \end{aligned}$$

As for $J_{2,2,2}$, observe that

$$\begin{aligned}
 J_{2,2,2} &\leq E\left\{\Delta I_{\{0\leq\Delta\leq a\}} \int_{-\Delta}^0 I_{\{z\leq W\leq z-t\}} dt\right\} \\
 &\leq E(\Delta^2 I_{\{0\leq\Delta\leq a\}} I_{\{z\leq W\leq z+a\}}) \\
 (2.15) \quad &\leq 3a\tau,
 \end{aligned}$$

where in the last inequality we used the concentration inequality in Lemma 2.1 below.

Similarly, we have

$$J_{2,2,2} \geq -3a\tau.$$

This proves Theorem 2.1. □

Lemma 2.1. *Under the assumption of Theorem 2.1, we have*

$$(2.16) \quad E(\Delta^2 I_{\{0\leq\Delta\leq a\}} I_{\{z\leq W\leq z+a\}}) \leq 3a\tau$$

for $a > 0$.

Proof. Let

$$f(x) = \begin{cases} -1.5a & \text{for } x \leq z - a, \\ x - z - a/2 & \text{for } z - a \leq x \leq z + 2a, \\ 1.5a & \text{for } x \geq z + 2a. \end{cases}$$

By (2.4),

$$\begin{aligned}
 3a\tau &\geq 2\tau E(Wf(W)) \\
 &= E\{(W - W^*)(f(W) - f(W^*))\} \\
 &= E\left\{\Delta \int_{-\Delta}^0 f'(W+t) dt\right\} \\
 &\geq E\left\{\Delta \int_{-\Delta}^0 I_{\{|t|\leq a\}} I_{\{z\leq W\leq z+a\}} f'(W+t) dt\right\} \\
 &= E\left(|\Delta| \min(a, |\Delta|) I_{\{z\leq W\leq z+a\}}\right) \\
 &\geq E\left(\Delta^2 I_{\{0\leq\Delta\leq a\}} I_{\{z\leq W\leq z+a\}}\right)
 \end{aligned}$$

as desired. □

REFERENCES

- [1] Aldous, D. and Diaconis, P. (1999) Longest increasing subsequences: from patient sorting to Baik-Defit-Johansson theorem. *Bull. AMS (N.S.)* **36**, 413-432. MR1694204 (2000g:60013)
- [2] Borodin, A., Okounkov, A. and Olshanski, G. (2000). Asymptotics of Plancherel measures for symmetric groups. *J. Amer. Math. Soc.* **13**, 481-515. MR1758751 (2001g:05103)
- [3] Chen, L.H.Y. and Shao, Q.M. (2001). A non-uniform Berry-Esseen bound via Stein's method. *Probab. Theory Related Fields* **120**, 236-254. MR1841329 (2002h:60037)
- [4] Defit, P. (2000). Integrable systems and combinatorial theory. *Notices Amer. Math. Soc.* **47**, 631-640. MR1764262 (2001g:05012)
- [5] Diaconis, P. and Shahshahani, M. (1981). Generating a random permutation with random transpositions. *Z. Wahr. Verw. Gebiete* **57**, 159-179. MR0626813 (82h:60024)
- [6] Eskin, A. and Okounkov, A. (2001). Asymptotics of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials. *Invent. Math.* **145**, 59-103. MR1839286 (2002g:32018)
- [7] Fulman, J. (2005). Stein's method and Plancherel measure of the symmetric group. *Trans. AMS* **357**, 555-570.

- [8] Fulman, J. (2004). Martingales and character ratios, preprint.
- [9] Fulman, J. (2004). Card shuffling and the decomposition of tensor products. *Pacific J. Math.* **217** (2004), no. 2, 247–262. MR2109933
- [10] Fulman, J. (2004). Stein’s method, Jack measure and the metropolis algorithm, *J. Combin. Theory, A*. **108** (2004), no. 2, 275–296. MR2098845
- [11] Goldstein, L. and Rinott, Y. (1996). Multivariate normal approximations by Stein’s method and size bias couplings. *J. Appl. Probab.* **33**, 1–17. MR1371949 (97d:60038)
- [12] Hora, A. (1998). Central limit theorem for the adjacency operators on the infinite symmetric group. *Comm. Math. Phys.* **195**, 405–416. MR1637801 (99i:46058)
- [13] Ivanov, V. and Olshanski, G. (2002). Kerov’s central limit theorem for the Plancherel measure on Young diagrams. *Symmetric Functions 2001: Surveys of developments and perspectives*, Kluwer Academic Publishers, Dordrecht. MR2059361 (2005d:05148)
- [14] Johansson, K. (2001). Discrete orthogonal polynomial ensembles and the Plancherel measure. *Ann. Math.* **153**, 259–296. MR1826414 (2002g:05188)
- [15] Kerov, S. V. (1993). Gaussian limit for the Plancherel measure of the symmetric group. *Compt. Rend. Acad. Sci. Paris I*, 303–308. MR1204294 (93k:20106)
- [16] Okounkov, A. and Pandharipande, R. Gromov-Witten theory, Hurwitz numbers, and matrix models I, preprint.
- [17] Rinott, Y. and Rotar, V. (1997). On coupling constructions and rates in the CLT for dependent summands with applications to the antivoter model and weighted U -statistics. *Ann. Appl. Probab.* **7**, 1080–1105. MR1484798 (99g:60050)
- [18] Sagan, B. (1991). The symmetric group. Representations, combinatorial algorithms, and symmetric functions. Springer-Verlag.
- [19] Steele, J.M. (1997). *Probability Theory and Combinatorial Optimization*. Society for Industrial and Applied Mathematics, Philadelphia. MR1422018 (99d:60002)
- [20] Stein, C. (1986). Approximate computation of expectations. *Institute of Mathematical Statistics, Lect. Notes* **7**. MR0882007 (88j:60055)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OREGON 97403 – AND –
DEPARTMENT OF MATHEMATICS, DEPARTMENT OF STATISTICS AND APPLIED PROBABILITY, NATIONAL UNIVERSITY OF SINGAPORE

E-mail address: `qmshao@darkwing.uoregon.edu`

DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU, ZHEJIANG 310027, PEOPLE’S REPUBLIC OF CHINA