

The Bile Acid Membrane Receptor TGR5: A Valuable Metabolic Target

Thijs W.H. Pols Lilia G. Noriega Mitsunori Nomura Johan Auwerx
Kristina Schoonjans

Laboratory of Integrative and Systems Physiology at the Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

Key Words

Metabolism · Bile acids · Nuclear receptors · TGR5 · G-protein-coupled receptor

Abstract

Bile acids (BAs) are amphipathic molecules that facilitate the uptake of lipids, and their levels fluctuate in the intestines as well as in the circulation depending on food intake. Besides their role in dietary lipid absorption, BAs function as signaling molecules that activate specific BA receptors and trigger downstream signaling cascades. The BA receptors and the signaling pathways they control are not only important in the regulation of BA synthesis and their metabolism, but they also regulate glucose homeostasis, lipid metabolism and energy expenditure – processes relevant in the context of the metabolic syndrome. In addition to the function of the nuclear receptor FXR α in regulating local effects of BAs in the organs of the enterohepatic axis, increasing evidence points to a crucial role of the G-protein-coupled receptor TGR5 in mediating systemic actions of BAs. Here we review the current knowledge on BA receptors, with a strong focus on the cell membrane receptor TGR5, which has emerged as a promising target for intervention in metabolic diseases.

Copyright © 2011 S. Karger AG, Basel


Introduction

Bile acids (BAs) are a component of bile, which also contains phosphatidylcholine, bilirubin and cholesterol. Through their detergent properties, which allow the emulsification of lipids, BAs facilitate the uptake of lipids together with the fat-soluble vitamins A, D, E and K from the intestine [1]. BAs also control the intestinal microbial flora and play a role in the elimination of cholesterol from the body [2, 3]. More recently, BAs are increasingly being appreciated as signaling molecules that inform cells and organs concerning the fasting/feeding state, thereby regulating processes ranging from BA and lipid metabolism to glucose and energy homeostasis [4, 5]. This metabolic function is in fact not too surprising given their central role in dietary lipid absorption. We review here the most recent developments in the field of BA signaling, and the potential pharmaceutical repercussions in using them to treat various facets of the metabolic syndrome.

Signaling Pathways Activated by BAs

Nuclear Receptor Signaling Pathways

The farnesol X receptor (FXR) is a nuclear receptor (NR) that controls BA homeostasis. The FXR cDNA was first cloned from both mouse and rat in 1995 [6, 7]. Rodents have two FXR family members, FXR α and FXR β .



In humans only FXR α is expressed, although FXR β is still present in the genome as a nonexpressed pseudogene. FXR α is most potently activated by conjugated or unconjugated forms of chenodeoxycholic acid (CDCA) with an EC₅₀ in the range of 4.5–10 μ M [8–10], while FXR β in mice is activated by lanosterol, an intermediate of BA synthesis [11].

In addition to FXR α , other NRs directly activated by lithocholic acid (LCA) are the pregnane X receptor (PXR) and the vitamin D receptor (VDR) [12–14] (fig. 1a). PXR is not only activated by LCA, but also by certain BA precursors, such as 7 α -OH-4-cholesten-3-one [15]. PXR is like FXR α expressed in the liver and intestine [12, 13]. One of the physiological functions of PXR is to induce phase I detoxification through induction of CYP3A, which explains the finding that PXR transgenic mice are

protected against LCA-induced liver toxicity in a model of cholestasis [12]. Like PXR, VDR plays a role in the detoxification of BAs through induction of CYP3A [14]. In addition to their roles in detoxifying BAs, both NRs inhibit BA synthesis. Activation of VDR by vitamin D inhibits BA synthesis via the FGF19-CYP7A1 pathway [16], and activation of PXR with rifampicin, an agonist of hPXR, inhibits BA synthesis through a pathway involving HNF4 α [17]. It should be noted that high concentrations of LCA (30–100 μ M) are required to activate VDR and PXR, bringing into question the physiological relevance of the direct regulation of their activity by BAs.

Kinase Signaling Pathways Regulated by BAs

BAs also modulate kinase signaling pathways, such as the JNK pathway, which also downregulate CYP7A1 [18].

Another study has coupled JNK activation indirectly to BAs as the mechanism of the FGF19-FGFR4 signaling pathway to suppress CYP7A1 [19]. In addition to JNK, the p38 mitogen-activated protein kinases, the extracellular signal-regulated kinase (ERK) pathway and Akt are activated by BAs [20, 21]. Although the precise mechanisms remain elusive, a study indicated that phosphorylation of ERK1, ERK2 and Akt by BAs is induced through both radical oxygen species-dependent and G-protein-coupled receptor (GPCR)-dependent, pathways [21]. The activation of these phosphorylation cascades by BAs is, besides the suppression of CYP7A1, also involved in the regulation of apoptosis and cytoprotective effects [20, 22]. It is also tempting to speculate that some of these signaling pathways may contribute to the enhanced lifespan observed in yeast exposed to BAs, as yeast lack both the nuclear and membrane BA receptors [23].

GPCR Signaling Pathways Modulated by BAs

Most relevant to this review is that BAs also bind and modulate the activity of specific GPCRs (fig. 1a). The GPCR family comprises over 800 receptors, divided into 3 subgroups [24]. There are currently 3 GPCRs known that are modulated by BAs. Based on their sequence, these BA-modulated GPCRs are classified as class A or in the rhodopsin-like receptor class, the largest subgroup of GPCRs. Although BAs modulate the activity of muscarinic receptors (also designated as acetylcholine receptors) [25, 26] and inhibit the activity of formyl-peptide receptors [27, 28], the GPCR that has been most studied in relation to BAs is TGR5, also known as M-BAR, GPBAR or GPR131 [29, 30]. TGR5 is encoded by a single-exon gene, and its conservation among vertebrates underlines its physiological importance [29]. TGR5 is activated by several BAs, with LCA being the most potent natural agonist with an EC_{50} of $0.53 \mu\text{M}$ [29, 30]. Other BAs that activate TGR5 include conjugated and unconjugated forms of deoxycholic acid, CDCA and cholic acid with an EC_{50} of 1.0, 4.4 and $7.7 \mu\text{M}$, respectively [30].

TGR5 is abundantly expressed in gallbladder epithelium [31, 32] and in the intestine, particularly in the ileum and colon [29, 31, 33]. Also human spleen as well as human mononuclear and CD14+ white blood cells express high amounts of TGR5 [30]. TGR5 was also present in rabbit spleen and rabbit alveolar macrophages [30]. TGR5 is detected in several liver cells, including rat liver sinusoidal endothelial cells, as well as in rat Kupffer cells, which are resident macrophages of the liver [34, 35]. Furthermore, TGR5 is expressed in BAT, skeletal muscle and selected areas of the central nervous system [36, 37].

The BA Receptor TGR5

TGR5 in BA Homeostasis and Metabolism

In resting conditions, GPCRs, including TGR5, are in a so-called low-affinity state. In response to binding of BAs to the ligand-binding pocket of the receptor, a complex is released from TGR5 consisting of G-protein- α_s , $-\beta$ and $-\gamma$ [24, 30]. GDP is subsequently released from the G-protein and replaced by GTP, leading to dissociation of the G-protein complexes into G-protein- α_s and $-\beta\gamma$ dimers. G-protein- α_s then activates adenylyl cyclase inducing cAMP production, and the subsequent activation of protein kinase A, and its downstream signaling pathways [30] (fig. 1b). Whether TGR5 may also bind to other G-proteins, which have distinct downstream effector molecules, is still unresolved [24].

To explore the biological role of TGR5, several groups have independently generated TGR5^{-/-} mice [31, 33, 38]. Interestingly, the total BA pool size in TGR5^{-/-} mice was decreased [33], which was also observed by our lab (J.A. and K.S., unpubl. data). Why the BA pool decreases in TGR5^{-/-} mice is currently unknown, but appears independent of changes in fecal BA excretion [33]. In addition to changes in the BA pool size, TGR5^{-/-} mice fed a lithogenic diet are protected against cholesterol gallstone formation [39]. Hydrophobic bile salts decrease gallbladder smooth muscle function potentially via stimulation of TGR5, which could be a contributing factor in the manifestation of gallstone disease [40]. Additionally, TGR5 is expressed in human cholangiocytes and plays a role in bile composition, via the induction of chloride secretion in gallbladder epithelial cells [32, 41]. Taken together, the physiological changes observed in mice lacking TGR5 with regard to BA homeostasis together with expression of TGR5 in human tissue relevant to BA homeostasis hint towards a role of TGR5 in bile formation and homeostasis in man [32, 41].

Effect of TGR5 on Body Weight

TGR5 activation reduces the body weight of mice fed a high-fat diet. We first demonstrated that dietary supplementation of BAs significantly reduces body weight gain in C57Bl/6J mice fed a high-fat diet [36]. Cholic acid administration completely prevented high-fat diet-induced changes in adipose mass and morphology, and reversed 120 days of diet-induced weight gain, without apparent toxicity [36]. The weight-reducing effects were not due to reduced caloric intake, but were the consequence of enhanced energy expenditure. Using deiodinase-2-deficient mice, the effect of cholic acid on energy expenditure

was shown to require the induction of deiodinase-2 through a TGR5-cAMP-mediated pathway that was active in murine BAT and in human skeletal muscle myoblasts. Importantly, this effect of BAs on energy expenditure was independent of FXR, as the FXR agonist GW4064, did not increase cAMP levels in BAT and increased diet-induced obesity in mice. Deiodinase-2 increases mitochondrial oxidative phosphorylation and energy expenditure in brown adipose tissue and muscle via the conversion of inactive thyroxine (T4) into active 3,5,3'-tri-iodothyronine (T3), which activates the thyroid receptor and thereby induces energy expenditure [36]. This effect was subsequently confirmed using the semi-synthetic BA, 6-ethyl-23(S)methylcholic acid (6EMCA or INT-777), which acts as a specific TGR5, but not FXR, agonist [38].

In agreement with our data, Maruyama et al. [33] observed that female TGR5^{-/-} mice weigh more. The latter effect of TGR5 was already observed in heterozygous female mice. In agreement with this observation, female TGR5^{-/-} mice had a higher fat content, while the lean body weight was unaffected. Also the body composition of the male TGR5^{-/-} mice showed a tendency towards increased fat content [33]. Interestingly, in the TGR5^{-/-} mice line that we generated in our laboratory, body weight was also significantly increased in males (K.S. and J.A., unpubl. data). These two studies contrast with another publication [39], which did not report an effect of the absence of TGR5 on body weight. This difference may very well be explained by different diets or by other differences in animal experimental conditions in this specific study. Furthermore, the mere absence of a receptor, as in the TGR5^{-/-} mice, should not necessarily translate in the opposite phenotype as observed after receptor activation by exogenous ligands.

Effects of TGR5 on Glucose Metabolism and Insulin Sensitivity

High circulating levels of BAs have been linked to beneficial effects on glucose metabolism, and improved insulin sensitivity and better postprandial glycemic control have been reported [42, 43]. We have demonstrated that BAs regulate glucose homeostasis through activation of TGR5 [44]. In agreement with the report that TGR5 induces GLP-1 secretion in cultured mouse enteroendocrine STC-1 cells [45], the semisynthetic BA, 6-ethyl-23(S)methylcholic acid (6EMCA or INT-777), which is a specific TGR5 agonist, induces GLP-1 secretion in both STC-1 cells as well as in human intestinal NCI-H716 cells, and contributes as such to the effects of TGR5 in glucose

homeostasis. Silencing of TGR5 in STC-1 cells using shRNA prevented the secretion of GLP-1, illustrating the involvement of TGR5 in this response. Although the mechanism underlying TGR5-induced GLP-1 secretion is not yet completely established, stimulation of oxidative phosphorylation may be involved. The resulting increase in the ATP/ADP ratio can then induce membrane depolarization and Ca²⁺ mobilization in a way reminiscent to the cascade of events leading to insulin release in pancreatic β -cells [38].

Using obese and insulin-resistant mouse models, we have shown that mice with a gain-of-function of TGR5 are more glucose-tolerant, whereas TGR5^{-/-} mice have impaired glucose clearance. This effect was correlated with a healthier pancreatic islet phenotype in the TGR5 transgenic mice, and is at least partly explained by the tonic increase of GLP-1 secretion by TGR5 [44]. It was recently reported by Poole et al. [37] that TGR5 is also expressed in inhibitory motor neurons and modulates intestinal motility. This effect of TGR5 could be related to GLP-1 induction, which is also known to inhibit intestinal motility [46]. In apparent contrast to these observations are the findings in a recent study with independently generated TGR5^{-/-} mice, demonstrating that female and male chow-fed TGR5^{-/-} mice show improved insulin sensitivity [39]. However, it was also shown in this study that male TGR5^{-/-} mice on a high-fat diet are insulin-resistant, which is in agreement with our findings. Other evidence that TGR5 activation is beneficial with regard to diabetes comes from the observation that the triterpenoid oleanolic acid, a natural TGR5 agonist, also improves glucose homeostasis [47].

TGR5 Modulates Immune Response

One of the initial studies on TGR5 examined its role in immune cells and linked TGR5 to the immunomodulatory properties of BAs [30]. This action of TGR5 is relevant, as low-grade inflammation contributes to the development of the metabolic syndrome [48]. TGR5 is highly expressed in monocytes and macrophages, an observation derived from the finding that TGR5 is expressed in human spleen and human CD14⁺ monocytes, as well as in rabbit alveolar macrophages [30, 35]. In accordance with a report that cAMP inhibits LPS-induced cytokine secretion [49], BAs capable of activating TGR5 were found to increase cAMP production in alveolar macrophages [30]. In addition, BAs reduce the phagocytic activity of these cells and inhibit LPS-induced production of proinflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin (IL)-1 α , IL-1 β , IL-6 and

IL-8 [30]. Human monocytic leukemia THP-1 cells transfected with TGR5 exhibit increased cAMP production and reduced LPS-induced TNF- α expression. These effects were not observed in untransfected THP-1 cells, which express low levels of TGR5, suggesting that these BAs effects are TGR5-dependent [30]. Furthermore, stimulation of isolated rat Kupffer cells with tauroolithocholic acid or other TGR5 agonists, such as oleanolic acid, as well as cAMP stimulation reduced the expression of IL-1 α , IL-1 β , IL-6 and TNF- α following LPS treatment [35].

TGR5 in Liver Function

Administration of the specific TGR5 agonist INT-777 to high-fat-diet-fed mice reduces liver steatosis and associated hepatocyte damage, as measured by plasma liver enzymes LDH, ASAT and ALAT [38]. This is correlated with decreased plasma triglyceride and nonesterified fatty acid levels. The latter is consistent with the pronounced hepatosteatosis observed in male TGR5^{-/-} mice fed a high-fat diet for 8 weeks [39]. These data suggest that activation of TGR5 may prevent nonalcoholic fatty liver disease. The precise mechanisms by which TGR5 induces this effect remain, however, to be dissected. Although TGR5 seems not expressed in hepatocytes, it is detected in many cell types of the liver where it could directly or indirectly modulate liver function and triglyceride metabolism. TGR5 is, for instance, enriched in Kupffer cells, which are resident liver macrophages [35] that can secrete proinflammatory cytokines and as such contribute to the progression of nonalcoholic fatty liver disease [50]. TGR5 has also been shown to modulate microcirculation and fluid secretion in the endothelial and biliary epithelial cells of the liver [34, 41]. Although the increased energy expenditure and GLP-1 secretion following TGR5 activation may very well explain the significant improvement in liver steatosis, it will be challenging to examine whether any of these other cell types in the liver contribute to the protective effects of TGR5 activation against steatosis. Cell-type-specific TGR5^{-/-} mouse models will be extremely valuable for addressing these questions.

BA Signaling as a Target for Intervention in the Metabolic Syndrome

BA Signaling as Target for Intervention

The significance of BAs in human triglyceride metabolism is underlined by findings that BA-binding resins increase VLDL levels and that treatment of cholesterol

gallstones in humans with CDCA reduces hypertriglyceridemia [51–53]. In addition to the effects of BAs on human triglyceride metabolism, BAs are correlated to increased insulin sensitivity in humans [42]. Furthermore, patients after bariatric surgery to correct for obesity have higher circulating BA levels, which are positively correlated to peak GLP-1 concentrations [43]. The latter observation was recently confirmed in obese patients who have a decreased postprandial BA response and suboptimal GLP-1 secretion in comparison to subjects of normal weight [54].

The biological properties of TGR5 described in this review, mostly observed in animal models, strongly indicate that TGR5 is linked to the beneficial properties of BAs in humans. This is underlined by the findings that the improvements of BAs on metabolic homeostasis are, amongst others, linked to GLP-1, which is regulated by TGR5 [38, 45]. Furthermore, a recent human genetic study identified an association between the single nucleotide polymorphism rs3731859 within the human TGR5 gene and BMI, waist circumference, intramyocellular lipids and fasting GLP-1 levels [55]. This interesting result warrants further studies to explore the effect of genetic polymorphisms in the TGR5 gene.

Pharmacological Targeting of TGR5

BAs bind and activate both TGR5 and other (nuclear) receptors, including FXR. The binding pocket for the membrane BA receptor TGR5 and the nuclear BA receptor FXR α is only partially conserved since minor structural modifications on the steroid side chain of BAs can dictate selectivity of the ligand towards TGR5 [56, 57]. Therefore, this unique ligand-binding pocket of TGR5 allows the design of receptor selective ligands, leading to drugs that are able to target TGR5 exclusively.

TGR5 may be targeted by natural compounds as well as with synthetic agonists. Such TGR5 agonists include natural BAs, semisynthetic BAs (such as 6-ethyl-23(S) methylcholic acid) [58], bile alcohols and triterpenoid compounds of plant origin, such as oleanolic acid and betulinic acid [47, 59, 60]. Certain steroid hormones also potentially activate TGR5 [59], an observation that has been recently confirmed [61]. Significant progress has also been made with the search for synthetic TGR5 agonists, as 3-aryl-4-isoxazolecarboxamides were recently identified to activate TGR5, and found to induce GLP-1 secretion in canines [62]. Furthermore, many drug companies have active TGR5 programs, which have already resulted in the publication of several patents that describe additional TGR5 ligands.

Not All That Shines Is Gold – Issues with TGR5

Recently, several properties of TGR5 have been described that require further investigation as they could potentially be at the basis of certain side effects. For example, TGR5 has also been linked to epidermal growth factor receptor (EGFR) and c-Jun N-terminal kinase (JNK) signaling pathways in cell culture models, which modulates cell proliferation and apoptosis [63, 64]. Unfortunately evidence of such an effect was only ascertained in cultured cells and further studies in vivo are definitely required [63, 64]. TGR5 agonists, including certain steroids [59], were also reported to stimulate radical oxygen species generation in cultured astrocytes [61], which also merits further investigation in vivo. TGR5 activation has been suggested to influence cardiomyocytes as the TGR5-active BAs tauro-CDCA and LCA activated AKT and inhibited glycogen synthase kinase-3 β in these cells [65]. In view of the pleiotropic effects of BAs, it will be of interest to assess these effects in TGR5^{-/-} mice.

In addition to these cellular observations, it has been reported that TGR5^{-/-} mice have reduced severity of pancreatitis, induced by direct exposure of the pancreas to high concentrations of tauroolithocholic acid 3-sulfate sodium salt [66]. The fact that the pancreas in this study is exposed to very high concentrations of tauroolithocholic acid 3-sulfate sodium salt, which are never reached even under pathological conditions, may question the physiological relevance of this study. Finally, the fact that TGR5^{-/-} mice were protected against cholelithiasis [31], could imply that TGR5 agonism predisposes to this condition; however, there are no data to support a negative effect of TGR5 agonists on cholelithiasis. It needs to be stressed that many of these potential liabilities were either observed in cultured cells or in animal models where extremely high nonphysiological concentrations of BAs were used. Further detailed studies using more physiological conditions are therefore required to evaluate the clinical relevance of these observations.

Conclusions and Future Perspectives

The metabolic studies described above suggest that targeting TGR5 could provide an exciting new therapeutic approach to improve several aspects of the metabolic syndrome. Multiple studies reveal that TGR5 has beneficial effects on body weight in high-fat-diet-fed mice. In addition, TGR5 activation improves glucose homeostasis and reduces hepatic steatosis. Beneficial effects of TGR5 on macrophage-driven inflammation, as evidenced by the

reduction of proinflammatory cytokines, may also contribute to a potential positive effect of TGR5 with regard to the metabolic syndrome. These properties of TGR5 clearly suggest that activation of this GPCR is valuable within the framework of the metabolic syndrome. In view of the established role of BA or BA-like molecules (e.g. da-fachronic acids) to promote longevity in yeast [67] and in the worm *Caenorhabditis elegans* [68], it is also plausible that the various strategies to modulate BA signaling could increase lifespan through their potent hormonal activities that improve metabolism and reduce inflammation – two important contributors that determine lifespan [69].

The development of several novel natural, semisynthetic and synthetic TGR5 agonists is likely to further advance this receptor as a target for the metabolic syndrome. In addition, localized or tissue-specific gene targeting will shed light on ways to increase the efficacy and specificity of drugs that target this BA receptor. Despite the fact that we are convinced targeting BA signaling pathways through TGR5 holds great promise for the intervention in metabolic diseases, much work still needs to be done, especially to make sure that such compounds are safe.

Acknowledgements

We thank the Swiss National Science Foundation, ERC, NIH and Ecole Polytechnique Fédérale de Lausanne for funding. T.W.H.P. is supported by a long-term fellowship from FEBS, L.G.N. is supported by a CONACYT fellowship, and M.N. is supported by a fellowship from AXA. We especially acknowledge Mitsuhiro Watanabe, Hiroyuki Sato, and Charles Thomas for their invaluable contributions to many of the studies discussed in this review and the other members of the Auwerx-Schoonjans laboratory over the years for instructive discussions.

Disclosure Statement

J.A. consults for Intercept, a company that develops bile acid therapeutics.

References

- 1 Russell DW: Fifty years of advances in bile acid synthesis and metabolism. *J Lipid Res* 2009;50(Suppl):S120–S125.
- 2 Russell DW, Setchell KD: Bile acid biosynthesis. *Biochemistry* 1992;31:4737–4749.
- 3 Inagaki T, Moschetta A, Lee YK, Peng L, Zhao G, Downes M, Yu RT, Shelton JM, Richardson JA, Repa JJ, Mangelsdorf DJ, Kliewer SA: Regulation of antibacterial defense in the small intestine by the nuclear bile acid receptor. *Proc Natl Acad Sci USA* 2006; 103:3920–3925.

- 4 Houten SM, Watanabe M, Auwerx J: Endocrine functions of bile acids. *EMBO J* 2006; 25:1419–1425.
- 5 Thomas C, Pellicciari R, Pruzanski M, Auwerx J, Schoonjans K: Targeting bile-acid signalling for metabolic diseases. *Nat Rev Drug Discov* 2008;7:678–693.
- 6 Seol W, Choi HS, Moore DD: Isolation of proteins that interact specifically with the retinoid X receptor: two novel orphan receptors. *Mol Endocrinol* 1995;9:72–85.
- 7 Forman BM, Goode E, Chen J, Oro AE, Bradley DJ, Perlmann T, Noonan DJ, Burka LT, McMorris T, Lamph WW, Evans RM, Weinberger C: Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* 1995;81:687–693.
- 8 Makishima M, Okamoto AY, Repa JJ, Tu H, Learned RM, Luk A, Hull MV, Lustig KD, Mangelsdorf DJ, Shan B: Identification of a nuclear receptor for bile acids. *Science* 1999; 284:1362–1365.
- 9 Parks DJ, Blanchard SG, Bledsoe RK, Chandra G, Consler TG, Kliewer SA, Stimmel JB, Willson TM, Zavacki AM, Moore DD, Lehmann JM: Bile acids: natural ligands for an orphan nuclear receptor. *Science* 1999; 284:1365–1368.
- 10 Wang H, Chen J, Hollister K, Sowers LC, Forman BM: Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. *Mol Cell* 1999;3:543–553.
- 11 Otte K, Kranz H, Kober I, Thompson P, Hoefler M, Haubold B, Rimmel B, Voss H, Kaiser C, Albers M, Cheruvallath Z, Jackson D, Casari G, Koegl M, Paabo S, Mous J, Kremoser C, Deuschle U: Identification of farnesoid X receptor beta as a novel mammalian nuclear receptor sensing lanosterol. *Mol Cell Biol* 2003;23:864–872.
- 12 Xie W, Radomska-Pandya A, Shi Y, Simon CM, Nelson MC, Ong ES, Waxman DJ, Evans RM: An essential role for nuclear receptors SXR/PXR in detoxification of cholestatic bile acids. *Proc Natl Acad Sci USA* 2001;98: 3375–3380.
- 13 Staudinger JL, Goodwin B, Jones SA, Hawkins-Brown D, MacKenzie KI, LaTour A, Liu Y, Klaassen CD, Brown KK, Reinhard J, Willson TM, Koller BH, Kliewer SA: The nuclear receptor PXR is a lithocholic acid sensor that protects against liver toxicity. *Proc Natl Acad Sci USA* 2001;98:3369–3374.
- 14 Makishima M, Lu TT, Xie W, Whitfield GK, Domoto H, Evans RM, Haussler MR, Mangelsdorf DJ: Vitamin D receptor as an intestinal bile acid sensor. *Science* 2002;296: 1313–1316.
- 15 Goodwin B, Gauthier KC, Umetani M, Watson MA, Lochansky MI, Collins JL, Leitersdorf E, Mangelsdorf DJ, Kliewer SA, Repa JJ: Identification of bile acid precursors as endogenous ligands for the nuclear xenobiotic pregnane X receptor. *Proc Natl Acad Sci USA* 2003;100:223–228.
- 16 Schmidt DR, Holmstrom SR, Fon Tacer K, Bookout AL, Kliewer SA, Mangelsdorf DJ: Regulation of bile acid synthesis by fat-soluble vitamins A and D. *J Biol Chem* 2010.
- 17 Li T, Chiang JY: Mechanism of rifampicin and pregnane X receptor inhibition of human cholesterol 7 alpha-hydroxylase gene transcription. *Am J Physiol Gastrointest Liver Physiol* 2005;288:G74–G84.
- 18 Gupta S, Stravitz RT, Dent P, Hylemon PB: Down-regulation of cholesterol 7alpha-hydroxylase (CYP7A1) gene expression by bile acids in primary rat hepatocytes is mediated by the c-Jun N-terminal kinase pathway. *J Biol Chem* 2001;276:15816–15822.
- 19 Holt JA, Luo G, Billin AN, Bisi J, McNeill YY, Kozarsky KF, Donahee M, Wang DY, Mansfield TA, Kliewer SA, Goodwin B, Jones SA: Definition of a novel growth factor-dependent signal cascade for the suppression of bile acid biosynthesis. *Genes Dev* 2003;17: 1581–1591.
- 20 Qiao D, Stratagouleas ED, Martinez JD: Activation and role of mitogen-activated protein kinases in deoxycholic acid-induced apoptosis. *Carcinogenesis* 2001;22:35–41.
- 21 Dent P, Fang Y, Gupta S, Studer E, Mitchell C, Spiegel S, Hylemon PB: Conjugated bile acids promote ERK1/2 and AKT activation via a pertussis toxin-sensitive mechanism in murine and human hepatocytes. *Hepatology* 2005;42:1291–1299.
- 22 Qiao L, Han SI, Fang Y, Park JS, Gupta S, Gilford D, Amorino G, Valerie K, Sealy L, Engelhardt JF, Grant S, Hylemon PB, Dent P: Bile acid regulation of C/EBPbeta, CREB, and c-Jun function, via the extracellular signal-regulated kinase and c-Jun NH2-terminal kinase pathways, modulates the apoptotic response of hepatocytes. *Mol Cell Biol* 2003; 23:3052–3066.
- 23 Goldberg AA, Richard VR, Kyryakov P, Bourque SD, Beach A, Burstein MT, Glebov A, Koupaki O, Boukh-Viner T, Gregg C, Juneau M, English AM, Thomas DY, Titorenko VI: Chemical genetic screen identifies lithocholic acid as an anti-aging compound that extends yeast chronological life span in a TOR-independent manner, by modulating housekeeping longevity assurance processes. *Aging (Albany NY)* 2010;2:393–414.
- 24 Pierce KL, Premont RT, Lefkowitz RJ: Seven-transmembrane receptors. *Nat Rev Mol Cell Biol* 2002;3:639–650.
- 25 Raufman JP, Zimniak P, Bartoszko-Malik A: Lithocholytaurine interacts with cholinergic receptors on dispersed chief cells from guinea pig stomach. *Am J Physiol* 1998; 274:G997–G1004.
- 26 Raufman JP, Chen Y, Zimniak P, Cheng K: Deoxycholic acid conjugates are muscarinic cholinergic receptor antagonists. *Pharmacology* 2002;65:215–221.
- 27 Le Y, Murphy PM, Wang JM: Formyl-peptide receptors revisited. *Trends Immunol* 2002; 23:541–548.
- 28 Ferrari C, Macchiarulo A, Costantino G, Pellicciari R: Pharmacophore model for bile acids recognition by the FPR receptor. *J Comput Aided Mol Des* 2006;20:295–303.
- 29 Maruyama T, Miyamoto Y, Nakamura T, Tamai Y, Okada H, Sugiyama E, Itadani H, Tanaka K: Identification of membrane-type receptor for bile acids (M-BAR). *Biochem Biophys Res Commun* 2002;298:714–719.
- 30 Kawamata Y, Fujii R, Hosoya M, Harada M, Yoshida H, Miwa M, Fukusumi S, Habata Y, Itoh T, Shintani Y, Hinuma S, Fujisawa Y, Fujino M: A G protein-coupled receptor responsive to bile acids. *J Biol Chem* 2003;278: 9435–9440.
- 31 Vassileva G, Golovko A, Markowitz L, Abbondanzo SJ, Zeng M, Yang S, Hoos L, Tetzloff G, Levitan D, Murgolo NJ, Keane K, Davis HR Jr, Hedrick J, Gustafson EL: Targeted deletion of Gpbar1 protects mice from cholesterol gallstone formation. *Biochem J* 2006; 398:423–430.
- 32 Keitel V, Cupisti K, Ullmer C, Knoefel WT, Kubitz R, Haussinger D: The membrane-bound bile acid receptor TGR5 is localized in the epithelium of human gallbladders. *Hepatology* 2009;50:861–870.
- 33 Maruyama T, Tanaka K, Suzuki J, Miyoshi H, Harada N, Nakamura T, Miyamoto Y, Kanatani A, Tamai Y: Targeted disruption of G protein-coupled bile acid receptor 1 (Gpbar1/M-Bar) in mice. *J Endocrinol* 2006; 191:197–205.
- 34 Keitel V, Reinehr R, Gatsios P, Rupprecht C, Gorg B, Selbach O, Haussinger D, Kubitz R: The G-protein coupled bile salt receptor TGR5 is expressed in liver sinusoidal endothelial cells. *Hepatology* 2007;45:695–704.
- 35 Keitel V, Donner M, Winandy S, Kubitz R, Haussinger D: Expression and function of the bile acid receptor TGR5 in Kupffer cells. *Biochem Biophys Res Commun* 2008;372: 78–84.
- 36 Watanabe M, Houten SM, Matakaki C, Christoffolete MA, Kim BW, Sato H, Messaddeq N, Harney JW, Ezaki O, Kodama T, Schoonjans K, Bianco AC, Auwerx J: Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature* 2006;439:484–489.
- 37 Poole DP, Godfrey C, Cattaruzza F, Cottrell GS, Kirkland JG, Pelayo JC, Bunnett NW, Corvera CU: Expression and function of the bile acid receptor GpBAR1 (TGR5) in the murine enteric nervous system. *Neurogastroenterol Motil* 2010;22:814–825, e227–e228.
- 38 Thomas C, Gioiello A, Noriega L, Strehle A, Oury J, Rizzo G, Macchiarulo A, Yamamoto H, Matakaki C, Pruzanski M, Pellicciari R, Auwerx J, Schoonjans K: TGR5-mediated bile acid sensing controls glucose homeostasis. *Cell Metab* 2009;10:167–177.

- 39 Vassileva G, Hu W, Hoos L, Tetzloff G, Yang S, Liu L, Kang L, Davis H, Hedrick J, Lan H, Kowalski T, Gustafson E: Gender-dependent effect of Gpbar1 genetic deletion on the metabolic profiles of diet-induced obese mice. *J Endocrinol* 2010;205:225–232.
- 40 Lavoie B, Balemba OB, Godfrey C, Watson CA, Vassileva G, Corvera CU, Nelson MT, Mawe GM: Hydrophobic bile salts inhibit gallbladder smooth muscle function via stimulation of GPBAR1 receptors and activation of KATP channels. *J Physiol* 2010;588:3295–3305.
- 41 Keitel V, Ullmer C, Haussinger D: The membrane-bound bile acid receptor TGR5 (Gpbar-1) is localized in the primary cilium of cholangiocytes. *Biol Chem* 2010;391:785–789.
- 42 Shaham O, Wei R, Wang TJ, Ricciardi C, Lewis GD, Vasani RS, Carr SA, Thadhani R, Gerszten RE, Mootha VK: Metabolic profiling of the human response to a glucose challenge reveals distinct axes of insulin sensitivity. *Mol Syst Biol* 2008;4:214.
- 43 Patti ME, Houten SM, Bianco AC, Bernier R, Larsen PR, Holst JJ, Badman MK, Maratos-Flier E, Mun EC, Pihlajamaki J, Auwerx J, Goldfine AB: Serum bile acids are higher in humans with prior gastric bypass: potential contribution to improved glucose and lipid metabolism. *Obesity (Silver Spring)* 2009;17:1671–1677.
- 44 Thomas C, Auwerx J, Schoonjans K: Bile acids and the membrane bile acid receptor TGR5 – connecting nutrition and metabolism. *Thyroid* 2008;18:167–174.
- 45 Katsuma S, Hirasawa A, Tsujimoto G: Bile acids promote glucagon-like peptide-1 secretion through TGR5 in a murine enteroendocrine cell line STC-1. *Biochem Biophys Res Commun* 2005;329:386–390.
- 46 Tolessa T, Gutniak M, Holst JJ, Efendic S, Hellstrom PM: Inhibitory effect of glucagon-like peptide-1 on small bowel motility. Fasting but not fed motility inhibited via nitric oxide independently of insulin and somatostatin. *J Clin Invest* 1998;102:764–774.
- 47 Sato H, Genet C, Strehle A, Thomas C, Lobstein A, Wagner A, Mioskowski C, Auwerx J, Saladin R: Anti-hyperglycemic activity of a TGR5 agonist isolated from *Olea europaea*. *Biochem Biophys Res Commun* 2007;362:793–798.
- 48 Olefsky JM, Glass CK: Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol* 2010;72:219–246.
- 49 Yoshimura T, Kurita C, Nagao T, Usami E, Nakao T, Watanabe S, Kobayashi J, Yamazaki F, Tanaka H, Inagaki N, Nagai H: Inhibition of tumor necrosis factor- α and interleukin-1 β production by beta-adrenergic agonists from lipopolysaccharide-stimulated human peripheral blood mononuclear cells. *Pharmacology* 1997;54:144–152.
- 50 Baffy G: Kupffer cells in non-alcoholic fatty liver disease: the emerging view. *J Hepatol* 2009;51:212–223.
- 51 Grundy SM, Ahrens EH Jr, Salen G: Interruption of the enterohepatic circulation of bile acids in man: comparative effects of cholestyramine and ileal exclusion on cholesterol metabolism. *J Lab Clin Med* 1971;78:94–121.
- 52 Nestel PJ, Poyser A: Changes in cholesterol synthesis and excretion when cholesterol intake is increased. *Metabolism* 1976;25:1591–1599.
- 53 Angelin B, Einarsson K, Hellstrom K: Effect of cholestyramine on bile acid kinetics in patients with portal cirrhosis of the liver. Evidence of a selective defect in the formation of cholic acid. *Am J Dig Dis* 1978;23:1115–1120.
- 54 Glicksman C, Pournaras DJ, Wright M, Roberts R, Mahon D, Welbourn R, Sherwood R, Alagband-Zadeh J, le Roux CW: Postprandial plasma bile acid responses in normal weight and obese subjects. *Ann Clin Biochem* 2010;47:482–484.
- 55 Mussig K, Staiger H, Machicao F, Machann J, Schick F, Schafer SA, Claussen CD, Holst JJ, Gallwitz B, Stefan N, Fritsche A, Haring HU: Preliminary report: genetic variation within the GPBAR1 gene is not associated with metabolic traits in white subjects at an increased risk for type 2 diabetes mellitus. *Metabolism* 2009;58:1809–1811.
- 56 Pellicciari R, Sato H, Gioiello A, Costantino G, Macchiarulo A, Sadeghpour BM, Giorgi G, Schoonjans K, Auwerx J: Nongenomic actions of bile acids. Synthesis and preliminary characterization of 23- and 6,23-alkyl-substituted bile acid derivatives as selective modulators for the G-protein coupled receptor TGR5. *J Med Chem* 2007;50:4265–4268.
- 57 Macchiarulo A, Gioiello A, Thomas C, Massarotti A, Nuti R, Rosatelli E, Sabbatini P, Schoonjans K, Auwerx J, Pellicciari R: Molecular field analysis and 3D-quantitative structure-activity relationship study (MFA 3D-QSAR) unveil novel features of bile acid recognition at TGR5. *J Chem Inf Model* 2008;48:1792–1801.
- 58 Pellicciari R, Gioiello A, Macchiarulo A, Thomas C, Rosatelli E, Natalini B, Sardella R, Pruzanski M, Roda A, Pastorini E, Schoonjans K, Auwerx J: Discovery of 6 α -ethyl-23(S)-methylcholic acid (SEMCA, INT-777) as a potent and selective agonist for the TGR5 receptor, a novel target for diabetes. *J Med Chem* 2009;52:7958–7961.
- 59 Sato H, Macchiarulo A, Thomas C, Gioiello A, Une M, Hofmann AF, Saladin R, Schoonjans K, Pellicciari R, Auwerx J: Novel potent and selective bile acid derivatives as TGR5 agonists: biological screening, structure-activity relationships, and molecular modeling studies. *J Med Chem* 2008;51:1831–1841.
- 60 Genet C, Strehle A, Schmidt C, Boudjelal G, Lobstein A, Schoonjans K, Souchet M, Auwerx J, Saladin R, Wagner A: Structure-activity relationship study of betulinic acid, a novel and selective TGR5 agonist, and its synthetic derivatives: potential impact in diabetes. *J Med Chem* 2010;53:178–190.
- 61 Keitel V, Gorg B, Bidmon HJ, Zemtsova I, Spomer L, Zilles K, Haussinger D: The bile acid receptor TGR5 (Gpbar-1) acts as a neurosteroid receptor in brain. *Glia* 2010;58:1794–1805.
- 62 Evans KA, Budzik BW, Ross SA, Wisnoski DD, Jin J, Rivero RA, Vimal M, Szewczyk GR, Jayawickreme C, Moncol DL, Rimele TJ, Armour SL, Weaver SP, Griffin RJ, Tadepalli SM, Jeune MR, Shearer TW, Chen ZB, Chen L, Anderson DL, Becherer JD, De Los Frailes M, Colilla FJ: Discovery of 3-aryl-4-isoxazolecarboxamides as TGR5 receptor agonists. *J Med Chem* 2009;52:7962–7965.
- 63 Yang JI, Yoon JH, Myung SJ, Gwak GY, Kim W, Chung GE, Lee SH, Lee SM, Kim CY, Lee HS: Bile acid-induced TGR5-dependent c-Jun-N terminal kinase activation leads to enhanced caspase 8 activation in hepatocytes. *Biochem Biophys Res Commun* 2007;361:156–161.
- 64 Yasuda H, Hirata S, Inoue K, Mashima H, Ohnishi H, Yoshida M: Involvement of membrane-type bile acid receptor M-BAR/TGR5 in bile acid-induced activation of epidermal growth factor receptor and mitogen-activated protein kinases in gastric carcinoma cells. *Biochem Biophys Res Commun* 2007;354:154–159.
- 65 Desai MS, Shabier Z, Taylor M, Lam F, Thevananthar S, Kosters A, Karpen SJ: Hypertrophic cardiomyopathy and dysregulation of cardiac energetics in a mouse model of biliary fibrosis. *Hepatology* 2010;51:2097–2107.
- 66 Perides G, Laukkanen JM, Vassileva G, Steer ML: Biliary acute pancreatitis in mice is mediated by the G-protein-coupled cell surface bile acid receptor Gpbar1. *Gastroenterology* 2010;138:715–725.
- 67 Motola DL, Cummins CL, Rottiers V, Sharma KK, Li T, Li Y, Suino-Powell K, Xu HE, Auchus RJ, Antebi A, Mangelsdorf DJ: Identification of ligands for DAF-12 that govern dauer formation and reproduction in *C. elegans*. *Cell* 2006;124:1209–1223.
- 68 Gerisch B, Rottiers V, Li D, Motola DL, Cummins CL, Lehrach H, Mangelsdorf DJ, Antebi A: A bile acid-like steroid modulates *Caenorhabditis elegans* lifespan through nuclear receptor signaling. *Proc Natl Acad Sci USA* 2007;104:5014–5019.
- 69 Houtkooper RH, Williams RW, Auwerx J: Metabolic networks of longevity. *Cell* 2010;142:9–14.