

The Biology of Blood-Sucking in Insects Second Edition

Blood-sucking insects transmit many of the most debilitating diseases in humans, including malaria, sleeping sickness, filariasis, leishmaniasis, dengue, typhus and plague. In addition, these insects cause major economic losses in agriculture both by direct damage to livestock and as a result of the veterinary diseases, such as the various trypanosomiases, that they transmit. The second edition of The Biology of Blood-Sucking in Insects is a unique, topicled commentary on the biological themes that are common in the lives of blood-sucking insects. To do this effectively it concentrates on those aspects of the biology of these fascinating insects that have been clearly modified in some way to suit the blood-sucking habit. The book opens with a brief outline of the medical, social and economic impact of blood-sucking insects. Further chapters cover the evolution of the blood-sucking habit, feeding preferences, host location, the ingestion of blood and the various physiological adaptations for dealing with the blood meal. Discussions on host-insect interactions and the transmission of parasites by blood-sucking insects are followed by the final chapter, which is designed as a useful quick-reference section covering the different groups of insects referred to in the text.

For this second edition, *The Biology of Blood-Sucking in Insects* has been fully updated since the first edition was published in 1991. It is written in a clear, concise fashion and is well illustrated throughout with a variety of specially prepared line illustrations and photographs. The text provides a summary of knowledge about this important group of insects and will be of interest to advanced undergraduate and to postgraduate students in medical and veterinary parasitology and entomology.

MIKE LEHANE is Professor of Molecular Entomology and Parasitology in the Liverpool School of Tropical Medicine.



The Biology of Blood-Sucking in Insects

SECOND EDITION

M. J. Lehane

Liverpool School of Tropical Medicine





CAMBRIDGE UNIVERSITY PRESS

Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo

Cambridge University Press

The Edinburgh Building, Cambridge CB2 2RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org

Information on this title: www.cambridge.org/9780521836081

© M. J. Lehane 2005

This book is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2005

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this book is available from the British Library

ISBN-13 978-0-521-83608-1 hardback ISBN-10 0-521-83608-5 hardback

ISBN-13 978-0-521-54395-8 paperback ISBN-10 0-521-54395-9 paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this book, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.



Contents

	List of table	5	page v11
	List of boxes	3	X
	Preface		xi
	Acknowleds	gements	xiii
1	The impor	tance of blood-sucking insects	1
2	The evolution of the blood-sucking habit		7
		onged close association with vertebrates	7
		phological pre-adaptation for piercing	13
3	Feeding preferences of blood-sucking insects		15
	3.1 Host	choice	15
	3.2 Host	choice and species complexes	24
4	Location o	f the host	27
	4.1 A bel	navioural framework for host location	27
	4.2 App€	etitive searching	29
	4.3 Activ	ration and orientation	32
	4.4 Attra	ction	49
	4.5 Move	ement between hosts	52
5	Ingestion of the blood meal		56
	5.1 Probi	ng stimulants	56
	5.2 Mout	:hparts	57
	5.3 Verte	brate haemostasis	64
	5.4 Host	pain	68
	5.5 Insec	t anti-haemostatic and anti-pain factors	
	in sal	- · · ·	69
		ostimulants	76
	5.7 Blood	l intake	78
6	Managing the blood meal		84
		gut anatomy	84
	6.2 The b	plood meal	87
	6.3 Gond	otrophic concordance	96



vi	Cont	eents	
	6.4	Nutrition	98
	6.5	Host hormones in the blood meal	103
	6.6	Partitioning of resources from the blood meal	10ϵ
	6.7	Autogeny	109
7	Hos	t-insect interactions	116
	7.1	Insect distribution on the surface of the host	117
	7.2	Morphological specializations for life on the host	121
	7.3	Host immune responses and insect salivary	
		secretions	12ϵ
	7.4	Behavioural defences of the host	134
	7.5	Density-dependent effects on feeding success	142
8	Tran	smission of parasites by blood-sucking insects	150
	8.1	Transmission routes	150
	8.2	Specificity in vector–parasite relationships	163
	8.3	Origin of vector–parasite relationships	167
	8.4	Parasite strategies for contacting a vector	170
	8.5	Parasite strategies for contacting a vertebrate host	177
	8.6	Vector pathology caused by parasites	179
	8.7	Vector immune mechanisms	184
9	The blood-sucking insect groups 20		202
	9.1	Insect classification	202
	9.2	Phthiraptera	204
	9.3	Hemiptera	208
	9.4	Siphonaptera	213
	9.5	Diptera	219
	9.6	Other groups	257
	Refe	rences	259
	Inde:		312



Tables

1.1	An outline of the early investigations that laid the	
	foundations of medical and veterinary entomology.	page 2
1.2	Rounded estimates for the prevalence of disease, the	
	number at risk and the disability adjusted life years	
	(DALYs) for major vector-borne diseases.	3
1.3	Estimated losses in agricultural production caused by	
	blood-sucking insects.	4
4.1	Generalized opportunities and constraints on host location	
	by blood-sucking insects feeding during the day or night.	32
4.2	Different blood-sucking insects respond in different ways	
	to spectral information.	45
5.1	Adaptations of mouthpart components for different	
	purposes in various haematophagous insect groups.	58
5.2	Blood-sucking insects produce a wide range of	
	anti-haemostatic factors in their salivary secretions. This	
	table gives some examples with a range of different	
	activities.	71
5.3	The size of red corpuscles varies widely in different	
	animals. Given that many blood-sucking insects have	
	mouthparts with a terminal diameter of around 10µm, this	
	may be a factor affecting the feeding efficiency of	
	blood-sucking insects on different host species.	80
6.1	The size of the red blood meal and the time taken in its	
	digestion are affected by a range of factors including	
	ambient temperature, age of the insect, mating status,	
	stage of the gonotrophic cycle, previous feeding history,	
	and source of the blood meal. The figures given here are a	
	rough guideline to the 'average' meal size and time for	
	digestion in a variety of haematophagous insects.	88
6.2	The major constituents of the blood are reasonably	
	uniform in most host animals. The exception is the high	
	levels of nucleic acids in the blood of birds and reptiles	
	because of their nucleated red blood cells. Proteins are far	
	and away the most abundant nutrients in blood, and	



viii	List of tables	
	nutrients are unevenly distributed between whole blood (B), red blood cells alone (E) and plasma alone (P).	90
6.3	Symbionts are common in insects relying on blood as the sole food source throughout their lives. An outline is given of their anatomical locations and the means of transmission from one generation to the next in different insect groups.	99
6.4	Three types of female <i>Aedes taeniorhynchus</i> have been identified in terms of egg development: autogenous females (1); females that are autogenous if mated (2); and anautogenous forms (3). This pattern is influenced by the feeding success of the larval stage, as illustrated in this table.	113
6.5	Some mosquitoes can use sugar meals (10% sucrose in this experiment) to increase the number of autogenously produced eggs.	113
7.1	The choice of feeding site of <i>Aedes triseriatus</i> on eastern chipmunks and grey squirrels is influenced by length and density of body hair. The different feeding patterns on the two hosts reflects the differences in hair cover between them.	119
7.2	The anti-mosquito behaviour of a range of ciconiiform birds, showing that different host species display various types and degrees of defensive behaviour against blood-sucking insects.	137
8.1	Some of the most important associations of disease-causing organisms carried to humans and other animals by blood-sucking insects: (a) viruses, (b) rickettsia and bacteria, (c) protozoa and (d) nematodes.	151
8.2	Blood-sucking insects commonly take meals that are only a small proportion of the total blood present in the host animal (the ratio between total blood in the host and size of the insect's blood meal is given). This minimizes the chances of the insect ingesting any individual parasite during feeding. One strategy adopted by insect-borne parasites to overcome this problem is to produce large numbers of infective stages which circulate in the blood of the host.	171
8.3	The microfilariae of many filarial worms display a pronounced periodicity, with microfilarial numbers in the peripheral blood coinciding with the peak biting time of	171
	locally abundant vector species.	173



	List of tables	ix
8.4	Tsetse flies infected with trypanosomes feed more readily and probe more often than uninfected flies, thereby	
	increasing the chances of parasite transmission.	178
8.5	Comparison of the rate of formation of the peritrophic	170
0.0	matrix among various mosquito species.	189
8.6	The melanization response to subsequent challenge of	
	infected and uninfected Aedes aegypti, as shown by the	
	intrathoracic injection of specific microfilariae (mff) which	
	normally induce a strong melanization reaction.	200
9.1	The groups of insect. Those groups containing	
	blood-sucking insects are shown in bold.	203
9.2	The geographical distribution of triatomine species, which	
	have become highly adapted to the domestic-peridomestic	
	environment of man and so represent a particular threat as	
	vectors of Chagas' disease.	211
9.3	The divisions of the order Diptera and the major families	
	in each division. Families containing blood-sucking species	
	are in bold type.	221



Boxes

3.1 The importance of rates of mosquitoes biting humans for		
	the transmission of malaria.	page 20
3.2	Identification of the source of a blood meal.	21
7.1	Histopathology of the various stages in the sequence of	
	host response to insect bites.	130
8.1	Four blood cell types characterized in Aedes aegypti are	
	compared to haemocytes described in previous studies on	
	a variety of insects.	197



Preface

Blood-sucking insects are the vectors of many of the most debilitating parasites of humans and their domesticated animals. In addition they are of considerable direct cost to the agricultural industry through losses in milk and meat yields, and through damage to hides, wool and other products. So, not surprisingly, many books of medical and veterinary entomology have been written. Most of these texts are organized taxonomically, giving details of the life cycles, bionomics, relationships to disease and economic importance of each of the insect groups in turn. I have taken a different approach. This book is topic-led and aims to discuss the biological themes common to the lives of blood-sucking insects. To do this I have concentrated on those aspects of the biology of these fascinating insects that have been clearly modified in some way to suit the blood-sucking habit. For example, I have discussed feeding and digestion in some detail because feeding on blood presents insects with special problems, but I have not discussed respiration because it is not affected in any particular way by haematophagy. To reflect this better I have made a slight adjustment to the title of the book in this second edition. Naturally there is a subjective element in the choice of topics for discussion and the weight given to each. I hope that I have not let my enthusiasm for the particular subjects get the better of me on too many occasions and that the subject material achieves an overall balance. The major changes in this second edition most often reflect the revolutionary influence that molecular biology has had on the subject in the past 12 years.

Although the book is not designed as a conventional text of medical and veterinary entomology, in Chapter 9 I have given a brief outline of each of the blood-sucking insect groups. This chapter is intended as a quick introduction for those entirely new to the subject, or as a refresher on particular groups for those already familiar with the divisions of blood-sucking insects. There are several introductory textbooks of medical and veterinary entomology available to those requiring more information.

The book is primarily intended for advanced undergraduate and for postgraduate students, but because it looks at topics that cut across the normal research boundaries of physiology and ecology, behaviour and cell biology, I hope it may also be useful for more established scientists who



xii Preface

want to look outside their own specialism. I have tried to distil this broad spectrum of information, much of which is not readily available to the non-specialist, into a brief synthesis. For those who want to look further into a particular area I have included some of the references I found most useful in writing the text, and these will provide an entry into the literature. Clearly the subjects covered by the book encompass a vast number of publications and I am sure to have missed many important and interesting references for which I apologize in advance both to the reader and my fellow scientists. Many of the topics discussed in the different chapters are interrelated. To avoid repetition, and still give the broadest picture possible, I have given cross-references in the text which I hope the reader will find useful.

From a comparative point of view it is an unfortunate fact that most of the work on blood-sucking insects has been carried out on a few species. Consequently, tsetse flies and mosquitoes pop up on every other page. In many instances it remains to be seen how widely the lessons we have learned from these well-studied models can be applied. Where possible I have tried to point to general patterns that fit whole groups of blood-sucking insects. To help me in this I have divided the blood-sucking insects into three convenient but artificial categories: temporary ectoparasites, permanent ectoparasites and periodic ectoparasites. These categories are based solely on the behaviour biology of the blood-feeding stadia in the lives of these insects. Temporary ectoparasites are considered to be those largely free-living insects, such as the tabanids, mosquitoes, blood-feeding bugs and blackflies, that visit the host only long enough to take a blood meal. I also include insects such as the tsetse here, even though the male may be found in swarms closely associated with the host for large parts of its life. Permanent ectoparasites are considered to be those insects that live almost constantly on the host, such as lice, the sheep ked and tungid fleas. Finally, periodic ectoparasites are considered to be those insects that spend considerably longer on the host than is required merely to obtain a blood meal, but that nevertheless spend a significant amount of time away from the host. Insects that fall into this category include many of the fleas and Pupipara. These categories are no more than a useful generalization in the text; I make no claims for their rigour and I realize that it could be argued in several instances that an insect will sit as easily in one category as another.



Acknowledgements

I gladly and gratefully acknowledge the help I have received from many people during the writing of this book or the previous edition, particularly P. Billingsley, A. Blackwell, J. Brady, H. Briegel, I. Burgess, E. Bursell, R. Dillon, J. D. Edman, R. Galun, A. G. Gatehouse, M. Gaunt, M. Gillies, R. H. Gooding, M. Greaves, C. Green, M. Hafner, J. Hogsette, H. Hurd, A. M. Jordan, K. C. Kim, J. Kingsolver, M. Klowden, A. M. Lackie, B. R. Laurence, E. Levashina, A. G. Marshall, P. Mellor, D. Molyneux, P. Morrison, W. A. Nelson, G. O'Meara, G. S. Paulson, G. Port, N. A. Ratcliffe, J. M. Ribeiro, P. Rossignol, M. Rothschild, W. Rudin, C. J. Schofield, M. W. Service, J. J. B. Smith, W. Takken, S. Torr, G. A. Vale and J. Waage. I also thank Paula Hynes, Maria Turton, Paula Dwyer and Dafydd Roberts for help with the illustrations. Finally, most thanks go to my family, particularly Stella, without whose encouragement, support and practical help this book would never have been finished.