

# The Boundary Between Decidability and Undecidability for Transitive-Closure Logics

N. Immerman<sup>1\*</sup>, A. Rabinovich<sup>2</sup>, T. Reps<sup>3\*\*</sup>, M. Sagiv<sup>2\*\*</sup>, and G. Yorsh<sup>2\*\*\*</sup>

<sup>1</sup> Dept. of Comp. Sci. Univ. of Massachusetts, immerman@cs.umass.edu

<sup>2</sup> School of Comp. Sci., Tel Aviv Univ.,

{rabinoa, msagiv, gretay}@post.tau.ac.il

<sup>3</sup> Comp. Sci. Dept., Univ. of Wisconsin, reps@cs.wisc.edu

**Abstract.** To reason effectively about programs, it is important to have some version of a transitive-closure operator so that we can describe such notions as the set of nodes reachable from a program's variables. On the other hand, with a few notable exceptions, adding transitive closure to even very tame logics makes them undecidable.

In this paper, we explore the boundary between decidability and undecidability for transitive-closure logics. Rabin proved that the monadic second-order theory of trees is decidable, although the complexity of the decision procedure is not elementary. If we go beyond trees, however, undecidability comes immediately.

We have identified a rather weak language called  $\exists\forall(\text{DTC}^+[E])$  that goes beyond trees, includes a version of transitive closure, and is decidable. We show that satisfiability of  $\exists\forall(\text{DTC}^+[E])$  is NEXPTIME complete. We furthermore show that essentially any reasonable extension of  $\exists\forall(\text{DTC}^+[E])$  is undecidable.

Our main contribution is to demonstrate these sharp divisions between decidable and undecidable. We also compare the complexity and expressibility of  $\exists\forall(\text{DTC}^+[E])$  with related decidable languages including MSO(trees) and guarded fixed point logics.

We mention possible applications to systems some of us are building that use decidable logics to reason about programs.

## 1 Introduction

To reason effectively about programs, it is important to have some version of a transitive-closure operator so that we can describe such notions as the set of nodes reachable from a program's variables. On the other hand, with a few notable exceptions, adding transitive closure to even very tame logics makes them undecidable.

In this paper, we explore the boundary between decidability and undecidability for transitive-closure logics. Rabin [13] proved that the monadic second-order theory of trees is decidable, although the complexity of the decision procedure is not elementary. If we go beyond trees, however, undecidability comes immediately.

Modal logics and their extension to the  $\mu$  calculus have proved quite useful. The  $\mu$  calculus has an EXPTIME-complete satisfiability problem [3] and the same has been shown true even for the more expressive guarded fixed-point logic, as long as the vocabulary remains of bounded arity [6]. Guarded fixed-point logic can express reachability from a specific constant, or from some point of a specific color, and it can restrict this reachability to be along paths specified, for example, by a regular expression. What it cannot express is a reachability relation between a pair of variables, i.e., that there is a path from  $u$  to  $v$ .

We have identified a rather weak language, called  $\exists\forall(\text{DTC}^+[E])$ , that goes beyond trees, includes a version of the latter sort of transitive closure, and is decidable. We show

\* Supported by NSF grant CCR-0207373 and a Guggenheim fellowship.

\*\* Supported by ONR contract N00014-01-1-0796 and the von Humboldt Foundation.

\*\*\* Supported by the Israel Science Foundation.

that satisfiability of  $\exists\forall(\text{DTC}^+[E])$  is NEXPTIME complete. We furthermore show that **essentially any reasonable extension of  $\exists\forall(\text{DTC}^+[E])$  is undecidable.**

The main contribution of this paper is to demonstrate the above sharp divisions between decidable and undecidable. We also compare the complexity and expressibility of  $\exists\forall(\text{DTC}^+[E])$  with related decidable languages, including MSO(trees) and guarded fixed-point logics.

The main application we have in mind is for the static-analysis methods that we are pursuing. Very generally, we model the properties of an infinite set of data structures that can be generated by the program we are analyzing, using a bounded set of first-order, three-valued structures [14]. In [15], it is shown that this modeling can be improved so that it computes the most precise possible transformation summarizing each program step, through the use of decidable logics.

Furthermore, in [9] we show that we can use a method we call “structure simulation” to significantly extend the sets of data structures that we can model with decidable logics over trees (monadic second-order logic) or graphs ( $\exists\forall(\text{DTC}^+[E])$ ). In the latter case, transitive-closure information must be restricted to deterministic paths.

The advantage of  $\exists\forall(\text{DTC}^+[E])$  compared with MSO(trees) is that while the latter is usually much more expressive, we can go beyond trees in the former. As an example, to express reachability in dynamic, undirected graphs, as in [2], we need not only a spanning forest, but a record of all the remaining edges in the undirected graph [9].

Fig. 1 summarizes results concerning the decidability and complexity of satisfiability for relevant logics. All the languages will be defined precisely in the next two sections. For previously known results we include a reference, and for results new to this paper we include the number of the relevant theorem.

Decidable	Complexity	Citation	Undecidable	Citation
$\mu$ calculus	EXPTIME complete	[3]	$\text{FO}^2(\text{TC})$	[5]
Guarded Fixed Point	EXPTIME complete	[6]	$\text{FO}^2(\text{DTC})$	[5]
MSO(trees)	non-elementary	[13]	$\forall(\text{TC}^+[E])$	Cor 9
$\text{FO}^2$	NEXPTIME complete	[11, 4]	$\forall(\text{DTC}^+)$	Th 8
$\exists\forall$	$\Sigma_2^p$ complete	[1]	$\forall(\text{DTC}^- [E])$	Th 13
$\exists\forall(\text{TC}^-)$	$\Sigma_2^p$ complete	Prop 2		
$\exists\forall(\text{DTC}^+[E])$	NEXPTIME complete	Th 4, 5		
$\exists\forall(\text{TC}, f)$	NEXPTIME complete	Cor 6		

**Fig. 1.** Summary of the decidability and complexity, and the undecidability of the logics we study. The arity of all relation symbols is bounded. The results are the same for  $\forall$  and  $\exists\forall$ , and they are the same for the satisfiability and finite-satisfiability problems.

## 2 Background and Tiling

As we have mentioned, being able to express reachability is crucial for our applications. However, adding a transitive-closure operator tends to make even very tame logics undecidable. We use  $\text{TC}_{u,u'}[\varphi]$  to denote the reflexive, transitive closure of binary relation  $\varphi(u, u')$  [8]. **Note:** In this paper, we confine our attention to applications of  $\text{TC}[\varphi]$  for

which  $\varphi$  is quantifier-free and TC-free. Furthermore, we assume throughout that the arity of all relation symbols is bounded.<sup>4</sup>

For example, consider the simple, decidable logic  $\text{FO}^2$ . This is first-order logic restricted to having only two variables,  $x, y$ . Grädel et al. [5] prove that if we add the transitive-closure operator (TC) to  $\text{FO}^2$  then the resulting logic is undecidable. In fact, they prove that even  $\text{FO}^2(\text{DTC})$  is undecidable. Here DTC — deterministic transitive closure — is the restriction of transitive closure to paths that have no choices. For the binary relation  $E(x, y)$ , define  $E_d(x, y)$  as follows:

$$E_d(x, y) \stackrel{\text{def}}{=} E(x, y) \wedge \forall z (E(x, z) \rightarrow z = y) .$$

That is, if vertex  $v$  has more than one outgoing  $E$ -edge, then it has no outgoing  $E_d$ -edges. Then define DTC as follows:  $\text{DTC}[E] \stackrel{\text{def}}{=} \text{TC}[E_d]$ .

It is surprising that  $\text{FO}^2(\text{DTC})$  is undecidable, but the proof is that even this seemingly very weak language is strong enough to express tilings.

**Definition 1** *Define a tiling problem,  $\mathcal{T} = \langle T, R, D \rangle$ , to consist of a finite list of tile types,  $T = [t_0, \dots, t_k]$ , together with horizontal and vertical adjacency relations,  $R, D \subseteq T^2$ . Here  $R(a, b)$  means that tiles of type  $b$  fit immediately to the right of tiles of type  $a$ , and  $D(a, b)$  means that tiles of type  $b$  fit one step down from those of type  $a$ . A solution to a tiling problem is an arrangement of instances of the tiles in a rectangular grid such that a  $t_0$  tile occurs in the top left position, and a  $t_k$  tile occurs in the bottom right position, and all adjacency relationships are respected.*

Given a Turing machine,  $M$ , and an input,  $w$ , we can build a tiling problem,  $\mathcal{T}$ , of size  $O(|M| + |w|)$ , such that  $\mathcal{T}$  has a solution iff  $M$  on input  $w$  eventually halts. Here any correct tiling solution would represent an accepting computation of  $M$  on input  $w$ . Think of  $t_0$  as representing the initial state and  $t_k$  as representing the final accepting state. Thus, as is well known, any logic that can express tilings has undecidable finite satisfiability – and general satisfiability – problems.

(Standard definitions of tiling problems only require  $t_0$  at the top left, and do not also ask for  $t_k$  at the lower right. This minor change does not affect the undecidability and complexity results, but makes some of our constructions slightly simpler.) See [1] for a nice treatment of tiling problems, as well as discussions of many relevant decidable and undecidable logics.

### 3 Decidability of $\exists\forall(\text{DTC}^+[E])$

We start with the first-order logic  $\exists\forall$ , consisting of first-order formulas in prenex form with all existential quantifiers preceding all universal quantifiers. The vocabulary has no function symbols. It is well known and easy to see that the satisfiability problem for  $\exists\forall$  is decidable: Let  $\varphi \in \exists\forall$ . Form the Skolemization,  $\varphi_S$ , by replacing the existential quantifiers,  $\exists x_1, \dots, x_k$ , by new constants,  $c_1, \dots, c_k$ . Suppose  $\mathcal{A} \models \varphi_S$ . Let  $\mathcal{C}$  be the substructure of  $\mathcal{A}$  whose universe consists of the constant symbols appearing in  $\varphi_S$ . Since  $\varphi_S$  is universal, we have that  $\mathcal{C} \models \varphi_S$ . Thus,  $\varphi$  has a model iff it has a small model, i.e., one of size less than  $|\varphi|$ . We say that  $\exists\forall$  has the *small-model property*, in this case with models of at most linear size. To test if a universal formula,  $\varphi_S$ , is

<sup>4</sup> For our intended applications, arity 2 is sufficient and arity 3 is a luxury. In theory, an unbounded arity can significantly increase some of the complexity bounds.

satisfiable, we would guess a structure,  $\mathcal{A}$ , of size at most  $n = |\varphi_S|$  and then check that  $\mathcal{A} \models \varphi_S$ . Testing whether a given structure satisfies an input universal first-order formula is co-NP complete. Thus satisfiability of  $\exists\forall$  formulas is in, and in fact complete for,  $\Sigma_2^P$ , the second-level of the polynomial-time hierarchy.

Since the existential quantifiers in  $\exists\forall$  formulas can be eliminated by adding constants, we limit our discussion to universal formulas. Let  $\forall(\text{DTC})$  consist of universal formulas in which DTC may occur. Unfortunately, as we will see, satisfiability of  $\forall(\text{DTC})$  and  $\forall(\text{TC})$  are undecidable (Theorem 8).

It is the positive occurrences of TC that cause the satisfiability of  $\forall(\text{TC})$  to be undecidable. Let  $\exists\forall(\text{TC}^-)$  consist of formulas in prenex form in which TC only occurs negatively.

**Proposition 2** *Satisfiability and finite satisfiability of  $\exists\forall(\text{TC}^-)$  are decidable with complexity complete for  $\Sigma_2^P$ .*

**Proof:** The above argument for  $\exists\forall$  continues to work. If  $\varphi \in \exists\forall(\text{TC}^-)$  is satisfiable, let  $\mathcal{A} \models \varphi_S$ , where  $\varphi_S$  is the Skolemization of  $\varphi$ . As above, let  $\mathcal{C}$  be the substructure of  $\mathcal{A}$  whose universe consists of the constant symbols appearing in  $\varphi_S$ . Then  $\mathcal{C} \models \varphi_S$  because if a path did not exist in  $\mathcal{A}$  then it still does not exist in  $\mathcal{C}$ . (Recall that we only apply TC to quantifier-free formulas.) Furthermore, we can test in polynomial time whether such a path exists in  $\mathcal{C}$ . Thus, the complexity of satisfiability remains  $\Sigma_2^P$  complete.  $\square$

**Definition 3** *Define  $\exists\forall(\text{DTC}^+[E])$  to be the restriction of  $\exists\forall(\text{DTC})$  in which the language has only one binary relation symbol,  $E$ , (plus unary relation symbols and constants), and all applications of DTC are positive occurrences of the form  $\text{DTC}[E]$ . In addition, we include in  $\exists\forall(\text{DTC}^+[E])$  **arbitrary negative occurrences** of  $\text{TC}[\varphi]$  for  $\varphi$  quantifier-free.<sup>5</sup> However, it is very important that there are **no negative occurrences of DTC**, for otherwise the language would become undecidable (Theorem 13).*

**Theorem 4**  *$\exists\forall(\text{DTC}^+[E])$  has the small-model property, with models of size at most  $2^{O(n^2)}$ , where  $n$  is the size of the formula. Thus, satisfiability and finite satisfiability of  $\exists\forall(\text{DTC}^+[E])$  are decidable, with complexity at most NEXPTIME.*

**Proof:** Using Skolemization, it suffices to prove these results for  $\forall(\text{DTC}[E])$ . Let  $\varphi \in \forall(\text{DTC}[E])$  be satisfiable and let  $\mathcal{A} \models \varphi$ . We will show that there exists a model  $\mathcal{B} \models \varphi$  such that  $\|\mathcal{B}\| \leq 2^{O(n^2)}$ . Here  $\|\mathcal{B}\|$  denotes the cardinality of the universe of the structure  $\mathcal{B}$ , and  $n = |\varphi|$ ,

Let  $c_1 \dots c_k$  be the constants occurring in  $\varphi$ . For each pair of constants,  $c_i, c_j$ , such that  $\mathcal{A} \models \text{DTC}[E](c_i, c_j)$ , there is a unique path  $p_{ij}$  from  $c_i$  to  $c_j$  in  $\mathcal{A}$ . Let  $\mathcal{A}'$  be the substructure of  $\mathcal{A}$  whose universe consists of the constants, plus all vertices that lie on any of the paths  $p_{ij}$ .

We claim that  $\mathcal{A}' \models \varphi$ . To see this, first observe that for any two elements  $a, b$  of the universe of  $\mathcal{A}'$  we have

$$\mathcal{A} \models \text{DTC}[E](a, b) \Rightarrow \mathcal{A}' \models \text{DTC}[E](a, b) \quad (1)$$


<sup>5</sup> A more accurate name for  $\exists\forall(\text{DTC}^+[E])$  would really be  $\exists\forall(\text{DTC}^+[E], \text{TC}^-)$ , but this is a mouthful, and all bounds remain the same whether or not the negative occurrences of TC are allowed.

(The proof of Theorem 12 exploits the fact that the converse need not hold.) Since  $a$  and  $b$  occur on paths  $p_{ij}$ , if  $\mathcal{A} \models \text{DTC}[E](a, b)$  then the path from  $a$  to  $b$  must be along the paths  $p_{ij}$ . Thus  $\mathcal{A}' \models \text{DTC}[E](a, b)$  holds as well.

Since  $\mathcal{A}'$  is a substructure of  $\mathcal{A}$  and  $\varphi$  is a universal formula with only positive occurrences of DTC, it follows from Equation (1) that  $\mathcal{A}' \models \varphi$ . (Note that the negative occurrences of  $\text{TC}[\varphi]$  with  $\varphi$  quantifier-free do not cause problems: since  $\mathcal{A}'$  is a substructure of  $\mathcal{A}$  it follows that if  $\mathcal{A} \models \neg\text{TC}[\varphi](a, b)$ , then  $\mathcal{A}' \models \neg\text{TC}[\varphi](a, b)$  as well.)

Structure  $\mathcal{A}'$  consists of a set of “trees” directed from leaf to root, all of whose leaves and roots are constants; however, (1) some of the “trees” may end in a cycle rather than a root; and (2) multiple edges may occur from some of the roots to other vertices. Note that if there is more than one edge from vertex  $v$ , then  $v$  does not occur on any DTC path, except perhaps as the last vertex. For this reason, if there are multiple edges in  $\mathcal{A}$  from constant  $c_i$ , then we can remove all such edges and replace them by a new unary relation symbol  $Q_i$  true of all the vertices that had edges from  $c_i$ ; as long as we modify  $\varphi$  accordingly. (In particular, we would change all occurrences of “ $E(x, y)$ ” to “ $E(x, y) \vee (x = c_i \wedge Q_i(y))$ ”.) Because we can eliminate issue (2), we henceforth assume that the graph  $\mathcal{A}'$  has outdegree at most one.

Note that some of the paths,  $p_{ij}, p_{i'j'}$  may intersect. If so, for simplicity we identify the first point of intersection for each pair of paths as a new constant. Observe that there are a total of at most  $k - 1$  such new constant symbols. Thus from now on we will only consider *direct paths*  $p_{ij}$  containing no intermediate constants. See Fig. 2 for an example graph where constants  $c_7, c_8$ , and  $c_9$  have been added.



**Fig. 2.** Example  $\mathcal{A}'$  from proof of Theorem 4 after constants  $c_7, c_8, c_9$  have been added.


After these normalization steps,  $\mathcal{A}'$  consists of  $k'$  constants and at most  $k'$  direct paths,  $p_{ij}$ , where  $k' \leq 2k - 1$ . Let  $r$  be the number of unary relation symbols, and  $m$  be the number of (universal) quantifiers in  $\varphi$ . We claim that no direct path  $p_{ij}$  need have length greater than  $2^{rm} + m + 1$ . Suppose on the contrary that the length of  $p_{12}$  is greater than  $2^{rm} + m + 1$ . Let the color of a vertex be the set of unary relation symbols that it satisfies. There are  $2^r$  possible colors and  $2^{rm}$  possible  $m$ -tuples of colors; consequently there must be at least two identically colored consecutive  $m$ -tuples,  $u_1, \dots, u_m$ , and  $v_1, \dots, v_m$ , in the interior of  $p_{12}$ . (By “consecutive” we mean the  $m$ -tuple is a path.) Form the structure  $\mathcal{B}$  from  $\mathcal{A}'$  by deleting vertices  $u_2$  through  $v_1$  and adding an edge from  $u_1$  to  $v_2$ .

We claim that  $\mathcal{B} \models \varphi$ . It suffices to show that for any  $m$ -tuple of vertices from  $\mathcal{B}$ ,  $b_1, b_2, \dots, b_m$ , there is a corresponding, isomorphic<sup>6</sup>  $m$ -tuple from  $\mathcal{A}'$ ,  $a_1, a_2, \dots, a_m$ .

<sup>6</sup> More explicitly, we mean that the map taking each  $b_i$  to  $a_i$  is an isomorphism of the induced substructures of  $\mathcal{B}$  and  $\mathcal{A}'$  generated by  $b_1, \dots, b_m$  and  $a_1, \dots, a_m$ , respectively. This may

Note that every vertex in  $\mathcal{B}$  is in  $\mathcal{A}'$ , and furthermore, the only difference between  $\mathcal{B}$  and  $\mathcal{A}'$  concerning these vertices is that  $E(u_1, v_2)$  holds in  $\mathcal{B}$  but not in  $\mathcal{A}'$ .

If any  $b_i$  is not on the path  $p_{12}$ , then we let  $a_i$  be the identical vertex in  $\mathcal{A}'$ . We may thus confine our attention to the most difficult case, namely, that  $b_1, b_2, \dots, b_m$  are all in the path  $p_{12}$ . Assume for simplicity that they occur in order. Our only problem is if for some  $\ell$ ,  $b_\ell = u_1$  and  $b_{\ell+1} = v_2$ . In this case, we let  $a_t = b_t$  for  $t \leq \ell$ , but we let  $a_{\ell+1} = u_2$ . Similarly, if  $b_{\ell+i-1} = v_i$  for all  $i \in \{2, \dots, s\}$ , then we must let  $a_{\ell+i-1} = u_i$ . Consider the first gap (if any), i.e.,  $b_i$  and  $b_{i+1}$  are not consecutive. We have that  $b_i = v_z$  and  $a_i = u_z$ , for some  $z$ . We can let  $a_j = b_j$  for  $j > i$ , see Fig. 3. Note that we have replaced some  $v_i$ 's by  $u_i$ 's but all unary relations, edge relations and connectivity have been preserved. Thus, as desired,  $a_1, a_2, \dots, a_m$  is isomorphic to  $b_1, b_2, \dots, b_m$ .



**Fig. 3.** Illustration of how for every  $m$ -tuple of vertices  $b_1, \dots, b_m$  from  $\mathcal{B}$  there is a corresponding isomorphic  $m$ -tuple of vertices  $a_1, \dots, a_m$  from  $\mathcal{A}'$ . In region (2) of  $\mathcal{B}$ ,  $b_1, \dots, b_i$  are assigned consecutive vertices; similarly, in region (2) of  $\mathcal{A}'$ ,  $a_1, \dots, a_i$  are assigned consecutive vertices. Because  $b_i$  and  $b_{i+1}$  are separated by two or more  $E$  edges in region (3) of  $\mathcal{B}$  (i.e., there a “gap”), the assignments for  $a_{i+1}, \dots, a_m$  in region (4) of  $\mathcal{A}'$  can match those for  $b_{i+1}, \dots, b_m$  in region (4) of  $\mathcal{B}$  exactly.

Thus  $\mathcal{B} \models \varphi$  as desired. We can continue shortening any remaining paths of length greater than  $2^{r^m} + m + 1$ . It follows that there is a model  $\mathcal{B}$  of  $\varphi$  and  $\|\mathcal{B}\| \leq (2k - 1)(2^{r^m} + m + 1) \leq 2^{|\varphi|^2}$ , as desired.  $\square$

It follows from Theorem 4 that the satisfiability of  $\exists\forall(\text{DTC}^+[E])$  formulas can be checked in NEXPTIME. We next show that this cannot be improved.

**Theorem 5** *The satisfiability of  $\exists\forall(\text{DTC}^+[E])$  formulas is NEXPTIME-complete.*

**Proof:** Let  $\mathcal{T}$  be a tiling problem as in Definition 1, and let  $n$  be a natural number. It is an NEXPTIME-complete problem to test on input  $(\mathcal{T}, 1^n)$  whether there is a  $\mathcal{T}$ -tiling of a square grid of size  $2^n$  by  $2^n$  [12].

We will define a formula  $\varphi_n$  that expresses exactly a solution to this tiling problem. There will be two constants:  $s$ , denoting the cell in the upper-left corner, and  $t$ , denoting the cell in the lower-right corner. The desired model will consist of  $2^{2n}$  tiles:

---

be thought of as an Ehrenfeucht-Fraïssé game in which the spoiler chooses the  $b_i$ 's and the duplicator answers with the  $a_i$ 's [8].



It is well known that the satisfiability and finite-satisfiability problems for monadic second-order logic with a single unary function symbol are decidable,<sup>8</sup> although their complexities are not elementary, even when restricted to first-order quantification [10, 13, 1, 7].

It is not hard to modify the proofs of Theorems 4 and 5 to apply to  $\forall(\text{TC}, f)$ . (For functions, the implication of Equation (1) is a biimplication, and thus the result goes through for positive and negative DTC's.)

**Corollary 6** *The finite satisfiability problem for  $\forall(\text{TC}, f)$  is NEXPTIME complete.<sup>9</sup>*

**Proof:** If a formula  $\varphi \in \forall(\text{TC}, f)$  has a finite model  $\mathcal{A}$ , then it must have a model of the form  $\mathcal{A}'$  as in the proof of Theorem 4. The only difference is that since  $f$  must be a total function, there are no roots; that is, all trees end in cycles. The size of the smallest model is still  $2^{O(n^2)}$ . The difference in counting is slight, namely, applications of the function symbol  $f$  can extend the apparent number of constant symbols:  $f(c_i)$  behaves like a new constant symbol  $c'_i$ , and  $f(x)$  behaves like a new universally quantified variable  $y$ , such that  $E(c_i, c'_i)$  and  $E(x, y)$ , respectively, must hold. Thus, the proof of Theorems 4 and 5 go through if we replace  $k$  and  $m$  by  $qk$  and  $qm$ , respectively, where  $q$  is the number of occurrences of  $f$  in  $\varphi$ .  $\square$

## 5 Undecidability of Related Logics

We next show that most reasonable extensions of the language  $\exists\forall(\text{DTC}^+[E])$  can express the solution to tiling problems, and thus are undecidable. In this section we show that any of the following changes cause undecidability: the use of TC; the presence of more than one binary relation symbol; or a single positive use of  $\text{DTC}[\sigma]$ , where  $\sigma$  is quantifier-free. In the next section, we show that  $\forall(\text{DTC}^-[E])$  is undecidable. To begin, we first show

**Theorem 7** *Satisfiability and finite satisfiability of  $\forall(\text{DTC}^+[V], \text{DTC}^+[H])$  — universal logic with two binary relations,  $V$  and  $H$ , and their positive deterministic transitive closure — are undecidable.*

**Proof:** Let  $\mathcal{T}$  be a tiling problem (Definition 1). We show how to write a formula  $\varphi \in \forall(\text{DTC}^+[V], \text{DTC}^+[H])$  such that  $\varphi$  is satisfiable iff  $\mathcal{T}$  has a solution.

Formula  $\varphi$  contains four constant symbols,  $a, b, c$ , and  $d$ , representing the four corners of the solution to  $\mathcal{T}$ ; see Fig. 4.


We assert that every element satisfies exactly one of the tile relations,  $T_0, \dots, T_k$ . We assert  $T_0(a) \wedge T_k(d)$ , i.e., the upper left tile is  $t_0$  and the lower right is  $t_k$ . We assert that  $H$  and  $V$  paths exist between the four corners:  $\text{DTC}[H](a, b) \wedge \text{DTC}[H](c, d) \wedge \text{DTC}[V](a, c) \wedge \text{DTC}[V](b, d)$ .

We add a unary predicate,  $Last$ , and assert the conjunction of the universal closure of the following formulas:  $Last(b)$ ,  $\neg V(x, b)$ ,  $V(x, y) \rightarrow (Last(x) \leftrightarrow Last(y))$ , and  $(H(x, y) \wedge \neg V(x, y)) \rightarrow \neg Last(x)$ . These assure that  $Last$  is true exactly of the tiles

<sup>8</sup> This is equivalent to the MSO theory of trees with multiple successor functions.

<sup>9</sup> This holds as well for the general satisfiability problem. For infinite structures there is a similar “small model” except that from some constants there is an infinite chain that intersects no other vertices of the structure. The infinite chain must repeat an  $m$ -tuple of colors and can from thereafter repeat exactly. Thus it has a representation of size  $2^{O(n^2)}$ .





**Fig. 4.** A tiling as expressed in Theorem 7.

in the rightmost column. In this column, we make the  $H$ -edges go down along the  $V$ -edges, i.e.,  $Last(x) \wedge Last(y) \rightarrow (H(x, y) \leftrightarrow V(x, y))$ . This allows us to express the fact that  $H$ -edges continue all the way to the right in every row, i.e., we assert:  $\forall x \text{DTC}[H](x, d)$ .

We assert that  $H$  and  $V$  edges satisfy the corresponding horizontal and vertical tiling constraints, using the formulas  $\text{Hor}$  and  $\text{Vert}$  as in Equation (2).  $\forall x, y((H(x, y) \wedge \neg Last(x) \rightarrow \text{Hor}(x, y)) \wedge (V(x, y) \rightarrow \text{Vert}(x, y)))$ .

We assert that the intermediate rows are filled in:  $\forall x, y, x', y'((H(x, y) \wedge V(x, x') \wedge V(y, y')) \rightarrow H(x', y'))$ .

Finally, we assert that the columns are filled in and line up:  $\forall x, y, x', y'((\neg Last(x) \wedge H(x, y) \wedge V(x, x') \wedge H(x', y')) \rightarrow V(y, y'))$ .


It is not hard to see that the conjunction of the above assertions is equivalent to the existence of a solution to the tiling problem,  $\mathcal{T}$ . Thus satisfiability of  $\forall(\text{DTC}^+[V], \text{DTC}^+[H])$  is undecidable.  $\square$

Theorem 7 shows that a second binary relation over which we can take DTC causes undecidability. We can modify the proof to show that even if there is only one (positive) occurrence of DTC, the logic is still undecidable if a second binary relation is allowed, or if DTC is allowed to be taken not just over the relation  $E$ , but over a formula that also involves unary relation symbols.

**Theorem 8** *Satisfiability and finite satisfiability of  $\forall(\text{DTC}^+)$  are undecidable. This holds even if there is only one occurrence of DTC and only one binary relation symbol. Also, if there is a second binary relation symbol, then the single occurrence of DTC can be restricted to the form  $\text{DTC}[E]$ .*

**Proof:** We modify the proof of Theorem 7 so that the path from  $a$  to  $d$  through the tiled rectangle is along a single snake-like path of the edge predicate,  $E$ , as in Fig. 5.

We do this by adding unary relation  $\text{First}$  denoting the first column of the tiling rectangle, plus the relation  $R$  true of the tiles in the odd-number rows. We then make the  $E$ -path go left-to-right on the rows satisfying  $R$  and right-to-left on the other rows.



**Fig. 5.** A tiling expressed with a single occurrence of DTC as in Theorem 8.

Define the edges along the snake-like path,  $\sigma(x, y) \equiv E(x, y) \wedge ((R(x) \leftrightarrow R(y)) \vee (First(x) \wedge \neg R(x) \wedge R(y)) \vee (Last(x) \wedge R(x) \wedge \neg R(y)))$ .

The single use of DTC is the assertion  $DTC[\sigma](a, d)$ . We also assert the completion of squares (see Fig. 5),

$(E(x, y) \wedge E(y, y') \wedge E(y', x') \wedge (R(x) \leftrightarrow R(y)) \wedge (R(x') \leftrightarrow R(y')) \wedge (R(y) \leftrightarrow \neg R(y'))) \rightarrow E(x, x')$ .

Finally, we add the following assertions, which together make sure that all models must be valid tilings:

1.  $T_0(a) \wedge T_k(d) \wedge First(a) \wedge Last(d) \wedge \neg(First(x) \wedge Last(x))$
2.  $\bigvee_{i=0}^k T_i(x) \wedge \bigwedge_{0 \leq i < j \leq k} \neg(T_i(x) \wedge T_j(x))$
3.  $(E(x, y) \wedge (R(x) \leftrightarrow \neg R(y))) \rightarrow ((First(x) \leftrightarrow First(y)) \wedge (Last(x) \leftrightarrow Last(y)))$
4.  $E(x, y) \rightarrow \neg((R(x) \wedge R(y) \wedge (Last(x) \vee First(y))) \vee (\neg R(x) \wedge \neg R(y) \wedge (Last(y) \vee First(x))))$
5.  $((E(x, y) \wedge R(x) \wedge R(y)) \vee (E(y, x) \wedge \neg R(x) \wedge \neg R(y))) \rightarrow Hor(x, y)$
6.  $(E(x, y) \wedge (R(x) \leftrightarrow \neg R(y))) \rightarrow Vert(x, y)$

Again formulas Hor and Vert are as in Equation (2). The conjunction of the universal closure of all the above assertions thus asserts a solution to the tiling problem,  $\mathcal{T}$ , as desired. To prove the last assertion in the statement of the theorem: with a second relation symbol,  $W$ , we can let  $E$  correspond to  $\sigma$ , and  $W$  correspond to  $E \wedge \neg\sigma$ .  $\square$

We remark that if in the proof of Theorem 8 we reverse the edges that are not  $\sigma$  edges, then we can use  $TC[E]$  in lieu of  $DTC[\sigma]$  and the proof goes through. Thus we have,

**Corollary 9** *Satisfiability and finite satisfiability of  $\forall(\text{TC}^+[E])$  are undecidable. This holds even if there is only a single occurrence of TC (it occurs as  $\text{TC}[E]$ ) and  $E$  is the only binary relation symbol.*

Note that the formulas in Theorems 7, 8, and Corollary 9 use only two variables except in the completion-of-squares formula. In fact, using an extra occurrence of TC, we can write equivalent formulas with only two variables. We do this by reversing the vertical edges in the even columns. We then assert that each non-boundary edge,  $\langle x, y \rangle$  is in an appropriate cycle, i.e.,  $\text{TC}[E](y, x)$  or  $\text{DTC}[\gamma](y, x)$  holds, for appropriate  $\gamma$ .

**Corollary 10** *If we allow a second occurrence of a transitive-closure operator, the undecidability results of Theorems 7, 8, and Corollary 9 all remain true for the corresponding languages with only two variables.*

## 6 Undecidability of $\forall(\text{DTC}^-[E])$

We were quite surprised to find that although  $\forall(\text{TC}^-)$  is decidable,  $\forall(\text{DTC}^-[E])$  is not. We give the somewhat subtle proof in this section. First we show that  $\forall(\text{DTC}^-[E])$  has an infinity axiom.

**Proposition 11** *There is a sentence in  $\forall(\text{DTC}^-[E])$  that is satisfiable, but only in an infinite model.*

**Proof:** The idea is that we know that if  $E(c_0, c_1)$  and  $\neg\text{DTC}[E](c_0, c_1)$  both hold, then there must be another edge from  $c_0$ . We can use this observation to write an infinity axiom that essentially expresses the existence of a successor function. We write the conjunction of the following formulas:

1.  $\forall v(v \neq c_1 \rightarrow (E(v, c_1) \wedge \neg\text{DTC}[E](v, c_1)))$
2.  $\forall v u_1 u_2(v \neq c_1 \wedge E(u_1, v) \wedge E(u_2, v) \rightarrow u_1 = u_2)$
3.  $c_0 \neq c_1 \wedge \forall v \neg E(v, c_0)$


(1) says that every vertex besides  $c_1$  has an edge to  $c_1$  but not a DTC path to  $c_1$ , so it must have outdegree greater than 1; (2) says that every vertex besides  $c_1$  has in-degree at most one; and (3) says that  $c_0$  has in-degree 0. Thus, there must be an infinite chain of edges starting at  $c_0$ .

These formulas are satisfied by a model that contains the natural numbers plus a new point called  $c_1$ , with edges  $E(n, c_1)$  and  $E(n, n + 1)$ , for  $n = 0, 1, \dots$   $\square$

**Theorem 12** *Satisfiability and finite satisfiability of  $\forall(\text{DTC}[E])$  are undecidable.*

**Proof:** We take as our starting point the undecidability proof of Theorem 8. Our new idea is to remove all of the non-boldface  $E$ 's in Fig. 5 and to replace them by a gadget of new green vertices, satisfying the unary relation symbol,  $G$ , and associated edges. The existence of the green vertices and their associated edges will be implied by the “not DTC trick” introduced in the proof of Proposition 11, together with some universal first-order statements that make sure that the vertical edges continue to be attached appropriately.

Just as in the proof of Theorem 8, we express the existence of a tiling. Since we have removed the non-boldface  $E$ 's, we can now simply express the path from the first tile to the last as  $\text{DTC}[E](a, d)$ .



**Fig. 6.** Gadget used in Theorem 12.

To define the gadget, we add two new constants,  $b$ , for the top rightmost tile, and  $c_1$  for the top rightmost green vertex, just below it. The green path proceeds in the opposite direction of the non-green, tile path directly above it, see Fig. 6.

We make the following assertions. These all concern the green row below each  $R$ , i.e., right-going, row of tiles. For simplicity, we skip the analogous case below each left-going row of tiles.

1.  $G(c_1) \wedge E(c_1, b) \wedge \forall ux(E(c_1, x) \wedge G(x) \wedge E(b, u) \rightarrow E(x, u))$
2.  $\forall x((\neg G(x) \leftrightarrow \text{DTC}[E](x, d)) \wedge (\neg G(x) \leftrightarrow \text{DTC}[E](a, x)))$
3.  $\forall xyz(G(x) \wedge E(x, y) \wedge E(x, z) \wedge y \neq z \rightarrow (G(y) \leftrightarrow \neg G(z)))$
4.  $\forall uvxyz(\neg G(u) \wedge \neg G(v) \wedge G(x) \wedge G(y) \wedge G(z) \wedge R(u) \wedge R(v) \wedge E(v, u) \wedge E(x, u) \wedge E(x, y) \wedge E(y, z) \rightarrow E(z, v))$
5.  $\forall uvxyz(\neg G(u) \wedge \neg G(v) \wedge G(x) \wedge G(y) \wedge G(z) \wedge \neg R(u) \wedge \neg R(v) \wedge E(u, v) \wedge E(x, u) \wedge E(x, y) \wedge E(y, z) \rightarrow E(z, v))$
6.  $\forall u, v, x, y(\neg G(u) \wedge \neg G(v) \wedge G(x) \wedge G(y) \wedge R(u) \wedge \neg R(v) \wedge E(x, u) \wedge E(x, y) \wedge E(y, v) \rightarrow \text{Vert}(u, v))$

(1) starts us out by saying that  $c_1$  is green, has an edge to  $b$ , and its green successor has an edge to the tile directly below  $b$ . (2) says that green vertices do not have DTC paths to  $d$ , but all non-green vertices do; it also says that all the non-green edges occur on the DTC-path from  $a$  to  $d$ . (3) says that if the outdegree of a green vertex is at least 2, then it has a green and a non-green successor. We will assure later, inductively, that each green vertex has an edge to a non-green vertex. Since the non-green vertex has a DTC-path to  $d$ , but the green vertex does not, this assures that the green vertex has outdegree 2. (4) is an inductive condition, which says that if  $x, y$ , and  $z$  are consecutive green nodes, and if  $x$  points up to a non-green node,  $u$ , then  $z$  points up to  $u$ 's predecessor,  $v$ . (5) is the similar condition for the edges going down.

Finally, condition (6) asserts that these green gadgets transmit the vertical information between the non-green, i.e., tile, nodes as desired.  $\square$

Theorem 12 leaves open the question of the decidability of  $\forall(\text{DTC}^-[E])$ . It would seem that the positive use of DTC was crucial in the statement  $\text{DTC}[E](a, d)$ . However, even this can be replaced by the “not DTC trick”. (The positive uses of DTC in formula (2) of the proof of Theorem 12 can easily be removed.) The conclusion is that  $\forall(\text{DTC}^-[E])$  is undecidable.

**Theorem 13** *Satisfiability and finite satisfiability of  $\forall(\text{DTC}^-[E])$  are undecidable.*

**Proof:** We modify the proof of Theorem 12 by removing the assertion  $\text{DTC}[E](a, d)$  and replacing it using the “not DTC trick”. More explicitly, we add another unary predicate  $B$  true of the tiles, and we add another constant,  $c_0$ . Then we make the following additional assertions:

1.  $B(a) \wedge \forall x(B(x) \wedge x \neq d \rightarrow E(x, c_0) \wedge \neg \text{DTC}[E](x, c_0))$
2.  $\forall xy(B(x) \wedge y \neq c_0 \wedge E(x, y) \rightarrow B(y))$
3. The in-degree for  $B$ -vertices from  $B$ -vertices is at most one, and it is zero for  $a$ .

(1) and (2) together assert that each  $B$ -vertex besides  $d$  has an edge to another  $B$ -vertex. It follows that either  $\text{DTC}[E](a, d)$  holds, or there is an infinite path. Thus, the formula is finitely satisfiable iff the corresponding tiling problem has a solution. (To show that the general satisfiability problem for  $\forall(\text{DTC}^- [E])$  is undecidable, we would modify the construction to assert that there is no node  $d$ , and thus an infinite path, so that the corresponding Turing machine, when started on blank tape, never halts. The tiling would have to be modified so that the first row has length one, and each successive row has one greater length. This is necessary so that an infinite path corresponds to an infinite computation rather than an infinitely long first row.)  $\square$

## 7 Complexity of the Decision Procedure

In this section, we study the complexity of the decision procedure for  $\exists\forall(\text{DTC}^+ [E])$ . The first thing we do is look more carefully at the proof of Theorem 5, and show that our lower bound is tight, matching the  $2^{O(n^2)}$  upper bound of Theorem 4.

**Lemma 14** *The formula  $\varphi_n$  used in the proof of Theorem 5 may be written in length  $O(n)$ .*

**Proof:** The only difficulty in keeping  $\varphi_n$  to total size  $O(n)$  is in writing the formulas  $\text{Suc}_h(x, y)$  and  $\text{Suc}_v(x, y)$ . These are nearly identical and we will restrict our attention to  $\text{Suc}_h(x, y)$ . Recall that  $\text{Suc}_h(x, y)$  means that the horizontal position of  $y$  is one greater than the horizontal position of  $x$ . (Our convention is that the bit positions are numbered 1 to  $n$ , with 1 being the high-order bit, and  $n$  the low-order bit.)  $\text{Suc}_h(x, y)$  can be written as follows:

$$\begin{aligned} \text{Suc}_h(x, y) \equiv & \bigvee_{i=1}^n \left[ \bigwedge_{j>i} (H_j(x) \wedge \neg H_j(y)) \wedge (\neg H_i(x) \wedge H_i(y)) \right. \\ & \left. \wedge \bigwedge_{j<i} (H_j(x) \leftrightarrow H_j(y)) \right] \end{aligned}$$

However, the length of the above formula is  $O(n^2)$ . We can decrease the size by keeping track of the position  $i$  in the above formula. We do this by adding  $2n$  more unary relation symbols,  $G_j, K_j, 1 \leq j \leq n$ . The intuitive meaning of  $K_i(x)$  is that it is bit  $i$  of the horizontal number that will be incremented as we go from  $x$  to its successor. This means that  $\neg H_i(x)$ , and for all  $j > i, H_j(x)$ ; i.e., there is a “0” in position  $i$ , and a “1” in each position to the right of  $i$ .

The intuitive meaning of  $G_j(x)$  is that  $j > i$  where  $K_i(x)$ . We also use the abbreviation  $L_j(x) \equiv \neg(K_j(x) \vee G_j(x))$ . (The mnemonic is that  $G$  holds for elements in positions “greater” than the  $K$  position;  $L$  holds for elements in “lesser” positions.)

The advantage of having these new relations is that we can now reduce the length of  $\text{Suc}_h(x, y)$  as follows:

$$\text{Suc}_n(x, y) \equiv \bigwedge_{j=1}^n \left[ (G_j(x) \wedge H_j(x) \wedge \neg H_j(y)) \vee (K_j(x) \wedge \neg H_j(x) \wedge H_j(y)) \vee (L_j(x) \wedge (H_j(x) \leftrightarrow H_j(y))) \right]$$

Finally, we must write down several more conditions. The conjunction of the following conditions assures that the new relations  $G_i$  and  $K_i$  are defined correctly.

1.  $\forall x (K_1(x) \vee K_2(x) \vee \dots \vee K_n(x) \vee (H_1(x) \wedge H_2(x) \dots H_n(x)))$
2.  $\forall x \left( \bigwedge_{i=1}^{n-1} (K_i(x) \rightarrow G_{i+1}(x)) \wedge \bigwedge_{i=1}^{n-1} (K_{i+1}(x) \rightarrow L_i(x)) \right)$
3.  $\forall x \left( \bigwedge_{i=1}^n (L_{i+1}(x) \rightarrow L_i(x)) \wedge \bigwedge_{i=1}^n (G_i(x) \rightarrow G_{i+1}(x)) \right)$
4.  $\forall x \left( \bigwedge_{i=1}^n \neg (K_i(x) \wedge G_i(x)) \wedge \bigwedge_{i=1}^n ((G_i(x) \rightarrow H_i(x)) \wedge (K_i(x) \rightarrow \neg H_i(x))) \right)$

□

It follows from Lemma 14 and the proof of Theorem 5 that we can write a sequence of formulas  $\varphi_n \in \exists\forall(\text{DTC}^+[E])$ ,  $n = 1, 2, \dots$  such that  $|\varphi_n| = O(n)$ ,  $\varphi_n$  has only two variables, and yet  $\varphi_n$ 's smallest model is of size  $2^{\Omega(n)}$ . This is the best possible with only two variables. To match the  $2^{O(n^2)}$  upper bound of Theorem 4, we need a formula with  $n$  variables.

We can count up to  $2^{n^2}$  using a sequence of  $n$  consecutive vertices, each with a number between 1 and  $2^n$ . We will add  $n$  more unary relation symbols,  $C_i$ ,  $1 \leq i \leq n$ . A tile will then be encoded by  $n$  vertices as follows:

$$\begin{array}{cccc} [C_1, h_1, v_1, t] & [C_2, h_2, v_2, t] & \dots & [C_n, h_n, v_n, t] \\ [C_1, h'_1, v'_1, t'] & [C_2, h'_2, v'_2, t'] & \dots & [C_n, h'_n, v'_n, t'] \end{array}$$

That is, the first  $n$  vertices hold tile  $t$  with its (collective) horizontal and vertical numbers  $\langle h_1, \dots, h_n \rangle$  and  $\langle v_1, \dots, v_n \rangle$  having values between 1 and  $2^{n^2}$ , the next  $n$  vertices hold tile  $t'$  with the successor number, etc. Using very similar ideas to the proof of Lemma 14 we can prove,

**Lemma 15** *Given any tiling problem,  $\mathcal{T}$ , we can write a sequence of formulas  $\varphi'_n$  of length  $O(n)$ ,  $n = 1, 2, \dots$ , such that  $\varphi'_n$  is satisfiable iff there is a solution to  $\mathcal{T}$  that is a  $2^{n^2}$  by  $2^{n^2}$  square.*

It follows that

**Corollary 16** *The  $2^{O(n^2)}$  upper bound of Theorem 4 is optimal.<sup>10</sup>*

<sup>10</sup> A referee pointed out that we are measuring the size of a formula as the number of symbols occurring in it. One could also consider that when there are  $n$  different relation symbols we need length  $\log n$  per symbol. Our formulation is simpler and the lower bound is tight because the upper bound used an even more lax measure of size, namely, the maximum of the number of constant symbols, relation symbols, and quantifiers.

## 8 Conclusions

We have introduced the language  $\exists\forall(\text{DTC}^+[E])$ , which is a decidable transitive-closure logic that goes beyond trees. We have shown that all the reasonable extensions of  $\exists\forall(\text{DTC}^+[E])$  that we could think of are undecidable. Uses of  $\exists\forall(\text{DTC}^+[E])$  exist, but how useful it might be remains to be seen. The following questions are worth considering:

- Unlike our other undecidability proofs, which only required two variables, our proof of the undecidability of  $\forall(\text{DTC}^-[E])$  used five variables. We suspect that this can be improved.
- We showed that the satisfiability of  $\exists\forall(\text{DTC}^+[E])$  is NEXPTIME complete. The lower bound depended on a formula that describes an exponentially long sequence of colors. We suspect that in practice the formulas one encounters would have much, much shorter sequences of color types. We suspect that techniques related to Ehrenfeucht-Fraïssé games can automatically find the relevant color sequences. These ideas might lead to a satisfiability algorithm that is feasible in practice.

## References

1. E. Börger, E. Grädel, and Y. Gurevich. *The Classical Decision Problem*. Springer-Verlag, 1996.
2. G. Dong and J. Su. Space-bounded foies. In *Principles of Database Systems*, pages 139–150. ACM Press, 1995.
3. E. Allen Emerson and Charanjit S. Jutla. The complexity of tree automata and logics of programs. In *Proc. 29th IEEE Symposium on Foundations of Computer Science*, pages 328–337. IEEE Computer Society Press, 1988.
4. E. Grädel, Ph. Kolaitis, and M. Vardi. On the decision problem for two-variable first-order logic. *Bulletin of Symbolic Logic*, 3:53–69, 1997.
5. E. Grädel, M. Otto, and E. Rosen. Undecidability results on two-variable logics. *Archive of Math. Logic*, 38:313–354, 1999.
6. E. Grädel and I. Walukiewicz. Guarded fixed point logic. In *Proc. 14th IEEE Symposium on Logic in Computer Science*, pages 45–54. IEEE Computer Society Press, 1999.
7. J.G. Henriksen, J. Jensen, M. Jørgensen, N. Klarlund, B. Paige, T. Rauhe, and A. Sandholm. Mona: Monadic second-order logic in practice. In *Tools and Algorithms for the Construction and Analysis of Systems, First International Workshop, TACAS '95, LNCS 1019*, 1995.
8. N. Immerman. *Descriptive Complexity*. Springer-Verlag, 1999.
9. N. Immerman, A. Rabinovich, T. Reps, M. Sagiv, and G. Yorsh. Verification via structure simulation. To appear in CAV'04, 2004.
10. Albert R. Meyer. Weak monadic second-order theory of successor is not elementary recursive. In *Logic Colloquium, (Proc. Symposium on Logic, Boston, 1972)*, pages 132–154, 1975.
11. M. Mortimer. On languages with two variables. *Zeitschr. f. math. Logik u. Grundlagen d. Math.*, 21:135–140, 1975.
12. C. Papadimitriou. *Computational Complexity*. Addison-Wesley, 1994.
13. M. Rabin. Decidability of second-order theories and automata on infinite trees. *Trans. Amer. Math. Soc.*, 141:1–35, 1969.
14. M. Sagiv, T. Reps, and R. Wilhelm. Parametric shape analysis via 3-valued logic. *Trans. on Prog. Lang. and Syst.*, pages 217–298, 2002.
15. G. Yorsh, T. Reps, and M. Sagiv. Symbolically computing most-precise abstract operations for shape analysis. In *TACAS*, pages 530–545, 2004.