

The cellular prion protein mediates neurotoxic signalling of β -sheet-rich conformers independent of prion replication

Ulrike K Resenberger¹, Anja Harmeier²,
Andreas C Woerner³, Jessica L Goodman⁴,
Veronika Müller¹, Rajaraman Krishnan⁴,
R Martin Vabulas³, Hans A Kretzschmar⁵,
Susan Lindquist⁴, F Ulrich Hartl³,
Gerd Multhaup², Konstanze F Winklhofer¹
and Jörg Tatzelt^{1,*}

¹Department of Metabolic Biochemistry, Neurobiochemistry, Adolf-Butenandt-Institute, Ludwig-Maximilians-University Munich, Munich, Germany, ²Institut für Chemie und Biochemie, Freie Universität Berlin, Berlin, Germany, ³Department of Cellular Biochemistry, Max Planck Institute of Biochemistry, Martinsried, Germany, ⁴Whitehead Institute for Biomedical Research and HHMI, Department of Biology, MIT, Cambridge, MA, USA and ⁵Center for Neuropathology and Prion Research, Ludwig-Maximilians-University Munich, Munich, Germany

Formation of aberrant protein conformers is a common pathological denominator of different neurodegenerative disorders, such as Alzheimer's disease or prion diseases. Moreover, increasing evidence indicates that soluble oligomers are associated with early pathological alterations and that oligomeric assemblies of different disease-associated proteins may share common structural features. Previous studies revealed that toxic effects of the scrapie prion protein (PrP^{Sc}), a β -sheet-rich isoform of the cellular PrP (PrP^C), are dependent on neuronal expression of PrP^C. In this study, we demonstrate that PrP^C has a more general effect in mediating neurotoxic signalling by sensitizing cells to toxic effects of various β -sheet-rich (β) conformers of completely different origins, formed by (i) heterologous PrP, (ii) amyloid β -peptide, (iii) yeast prion proteins or (iv) designed β -peptides. Toxic signalling via PrP^C requires the intrinsically disordered N-terminal domain (N-PrP) and the GPI anchor of PrP. We found that the N-terminal domain is important for mediating the interaction of PrP^C with β -conformers. Interestingly, a secreted version of N-PrP associated with β -conformers and antagonized their toxic signalling via PrP^C. Moreover, PrP^C-mediated toxic signalling could be blocked by an NMDA receptor antagonist or an oligomer-specific antibody. Our study indicates that PrP^C can mediate toxic signalling of various β -sheet-rich conformers independent of infectious prion propagation, suggesting a pathophysiological role of the prion protein beyond of prion diseases.

The EMBO Journal (2011) 30, 2057–2070. doi:10.1038/emboj.2011.86; Published online 25 March 2011

*Corresponding author. Department of Metabolic Biochemistry, Neurobiochemistry, Adolf-Butenandt-Institute, Ludwig-Maximilians-University Munich, Schillerstrasse 44, Munich D-80336, Germany. Tel.: +49 89 2180 75442; Fax: +49 89 2180 75415; E-mail: joerg.tatzelt@med.uni-muenchen.de

Received: 26 August 2010; accepted: 3 March 2011; published online: 25 March 2011

Subject Categories: proteins; neuroscience

Keywords: Alzheimer; intrinsically disordered; NMDA receptor; oligomer; prion

Introduction

Various approaches coming from neuropathology, genetics, animal modelling and biophysics have established a crucial role of protein misfolding in the pathogenesis of different neurodegenerative disorders, such as Alzheimer's disease (AD), Parkinson's disease, polyglutamine expansion diseases and prion diseases. However, there is an ongoing debate about the nature of the harmful proteinaceous species and how toxic conformers selectively damage neuronal populations (reviewed in Haass and Selkoe, 2007; Winklhofer *et al*, 2008; Ilieva *et al*, 2009).

The prion protein is best known for its crucial role in the pathogenesis of prion diseases, such as Creutzfeldt–Jakob disease and Gerstmann–Sträussler–Scheinker syndrome in humans, scrapie in sheep and goat, bovine spongiform encephalopathy in cattle and chronic wasting disease in free-ranging deer. In the disease, the cellular isoform of PrP (PrP^C) is converted into an aberrantly folded, β -sheet-rich isoform, designated scrapie prion protein (PrP^{Sc}). PrP^{Sc} is found in extracellular deposits in the diseased brain and has been shown to be the essential constituent of infectious prions (reviewed in Weissmann *et al*, 1996; Prusiner *et al*, 1998; Collinge, 2001; Chesebro, 2003). Notably, mice with a targeted disruption of the PrP gene (PRNP) are resistant to prion diseases and do not propagate infectious prions (Büeler *et al*, 1993). Moreover, based on transmission studies in laboratory animals, it has been proposed that productive propagation of infectious prions involves a direct interaction between PrP^C and PrP^{Sc}, which is highly dependent on the sequence homology between the two isoforms. As a consequence, PrP^{Sc} molecules derived from one species fail to or inefficiently transmit the disease to an organism expressing heterologous PrP^C molecules, a feature denoted as a species or transmission barrier (Prusiner *et al*, 1990; Moore *et al*, 2005; Collinge and Clarke, 2007). Interestingly, transgenic mouse models revealed that neuronal expression of GPI-anchored PrP^C is required to mediate neurotoxic effects of PrP^{Sc} (Brandner *et al*, 1996; Mallucci *et al*, 2003; Chesebro *et al*, 2005). We corroborated the important role of PrP^C as a mediator of PrP^{Sc}-induced toxicity and identified the intrinsically disordered N-terminal domain and the C-terminal GPI anchor of PrP^C as essential domains for this activity (Rambold *et al*, 2008). Recently, it was suggested that PrP^C may also have a role in the pathogenesis of AD. It was reported that amyloid β (A β)-induced inhibition of long-term potentiation (LTP) and memory impairment in transgenic mouse

models of AD requires PrP^C (Lauren *et al*, 2009; Gimbel *et al*, 2010). AD is the most common neurodegenerative disorder characterized by the deposition of A β -peptides in extracellular amyloid plaques and hyperphosphorylated τ in intraneuronal neurofibrillary tangles (reviewed in De Strooper, 2010; Palop and Mucke, 2010). While the exact mechanisms leading to progressive neurodegeneration in AD remain elusive, a variety of experimental studies indicated that soluble oligomeric species of A β contribute to synaptic dysfunction (Lambert *et al*, 1998; Walsh *et al*, 2002; Lacor *et al*, 2004; Cleary *et al*, 2005; Harmeier *et al*, 2009; Lauren *et al*, 2009). In addition, there is growing evidence that NMDA receptors are involved in mediating toxic effects of A β (reviewed in Ondrejcek *et al*, 2010). Notably, PrP^C can attenuate excitotoxicity by inhibiting NMDA receptors (Khosravani *et al*, 2008).

Herein, we show that cell surface localized PrP^C has the ability to interact with and mediate toxic signalling of β -sheet-rich conformers of different origin. In addition, we show that pro-apoptotic signalling via PrP^C induced by various β -sheet-rich conformers is independent of the replication of infectious prions and can be significantly reduced by an NMDA receptor antagonist.

Results

PrP^C mediates toxic signalling of homologous and heterologous PrP^{Sc}

We have recently established a cell culture assay to demonstrate that PrP^C localized at the cell surface is a mediator of pro-apoptotic signalling induced by PrP^{Sc} (Rambold *et al*, 2008). This assay is based on the co-cultivation of PrP^C-expressing cells with scrapie-infected cells that release PrP^{Sc} and infectious prions into the cell culture medium. As illustrated in Figure 1A, co-cultivation of SH-SY5Y cells expressing PrP^C with scrapie-infected mouse neuroblastoma (ScN2a) cells increased apoptotic cell death, as determined by activated caspase-3 (Figure 1A; Supplementary Figure S1A) or fragmented nuclei (Figure 1B) in cells expressing mouse PrP^C (moPrP). Control SH-SY5Y cells could be co-cultured with ScN2a cells without adverse effects (Figure 1A and B). Interestingly, PrP^{Sc} had also an adverse effect on mitochondrial morphology in PrP^C-expressing cells. A significant increase in the number of cells harbouring non-tubular/fragmented mitochondria was observed in PrP^C-expressing SH-SY5Y cells exposed to PrP^{Sc}, but not in cells co-cultivated with N2a cells (Figure 1C). Based on these and previously published studies in mice (Brandner *et al*, 1996; Mallucci *et al*, 2003; Chesebro *et al*, 2005), two plausible scenarios for the toxic effects of PrP^{Sc} can be envisaged. Neurotoxicity of PrP^{Sc} could be linked to its propagation in neuronal cells, which is dependent on the expression of PrP^C. We addressed this possibility, but found no evidence that PK-resistant PrP was formed in mouse PrP^C-expressing SH-SY5Y cells during co-cultivation with ScN2a cells (Supplementary Figure S1B). Alternatively, PrP^{Sc} elicits a deadly signal through a PrP^C-dependent signalling pathway. To experimentally address this scenario, we modified our cell culture model in order to minimize the possibility that propagation of infectious prions occurs. We therefore used SH-SY5Y cells in our co-cultivation assay expressing hamster, human, cervid or bovine PrP^C instead of mouse PrP^C. Based on previous

studies in transgenic mice and cell culture models (Scott *et al*, 1989, 1997), it is highly unlikely that mouse (mo) PrP^{Sc} released by ScN2a cells can induce efficient conversion of heterologous PrP^C molecules into PrP^{Sc}. Co-cultivation of SH-SY5Y expressing the respective PrP^C with uninfected N2a cells had no effect on cellular viability. However, co-cultivation with ScN2a cells revealed that heterologous PrP^C, be it of hamster, human, cervid or bovine origin, efficiently mediated toxic signalling of moPrP^{Sc} (Figure 1D). Thus, toxic signalling of PrP^{Sc} via PrP^C appears to be independent of PrP^{Sc} propagation.

PrP^C mediates toxic signalling of oligomeric A β secreted from transfected cells or prepared by chemical synthesis

Prompted by the observation that PrP^C can mediate toxic signalling of heterologous PrP^{Sc} molecules, we investigated whether A β could mediate toxic signalling via PrP^C. The rationale behind this approach was provided by studies showing that A β -induced blockage of LTP and memory impairment in transgenic mouse models of AD requires PrP^C (Lauren *et al*, 2009; Gimbel *et al*, 2010). We made use of a stably transfected Chinese hamster ovary cell line (CHO-7PA2) that expresses the familial AD mutation V717F in the amyloid precursor protein APP₇₅₁ and secretes A β (Podlisny *et al*, 1995). Importantly, the presence of secreted oligomeric A β in the medium of CHO-7PA2 cells has been shown to potently inhibit LTP *in vivo* (Walsh *et al*, 2002; Cleary *et al*, 2005). SH-SY5Y cells were grown on cover slips and transiently transfected with PrP^C. The cover slips were then placed into cell culture dishes with CHO-7PA2 or CHO control cells, and apoptosis of SH-SY5Y cells was analysed after 16 h of co-cultivation. Control SH-SY5Y cells expressing GPI-anchored GFP could be co-cultivated with CHO-7PA2 or CHO cells without adverse effects on cell viability (Figure 2A). Similarly, SH-SY5Y cells expressing PrP^C did not show increased apoptosis when co-cultured with control CHO cells. However, a significant increase in apoptotic cell death was observed when SH-SY5Y cells expressing PrP^C were co-cultivated with CHO-7PA2 cells (Figure 2A). Notably, the toxic effect of CHO-7PA2 cells was dependent on the generation of A β , as co-cultivation with CHO-7PA2 cells pre-treated with the γ -secretase inhibitor DAPT did not induce apoptotic cell death in PrP^C-expressing SH-SY5Y cells (Figure 2B). If PrP^{Sc} and A β mediate toxic signalling via different PrP^C-dependent pathways, one would assume that the exposure to both PrP^{Sc} and A β enhances toxicity. However, the rate of cell death in PrP^C-expressing SH-SY5Y co-cultivated with CHO-7PA2 or ScN2a cells was similar to those co-cultivated with both CHO-7PA2 and ScN2a cells (Figure 2C). To provide further evidence for a causal role of A β as toxic agent and to characterize the A β species mediating toxic signalling via PrP^C, we employed high- and low-molecular weight A β 42 aggregates obtained by size-exclusion chromatography (SEC) (Supplementary Figure S2) (Harmeier *et al*, 2009). Immediately after elution from the column, equal amounts of A β 42-peptides (500 nM final concentration) were added to SH-SY5Y cells expressing PrP^C. Consistent with our results from the co-cultivation assay, A β 42 was only toxic to cells expressing PrP^C. Moreover, only low-molecular weight oligomeric A β 42 (oligo) efficiently induced apoptotic cell death in PrP^C-expressing SH-SY5Y cells, while high-molecular weight aggregates had no




Figure 1 Scrapie prions induce apoptosis and interfere with mitochondrial integrity in cells expressing homologous and heterologous PrP^C. (A, B) SH-SY5Y cells expressing mouse (mo) PrP or GFP were co-cultured with ScN2a or N2a cells for 16 h. For quantification of apoptotic cell death, SH-SY5Y cells were fixed, permeabilized and stained for active caspase-3 (A) or fragmented nuclei (B). Expression of PrP was analysed by western blotting using the anti-PrP antibody 3F4 (middle panel). (C) Scrapie prions interfere with mitochondrial integrity. SH-SY5Y cells expressing moPrP were co-cultured with ScN2a or N2a cells for 16 h. Cells were stained with MitoTracker Red CMXRos to visualize mitochondria and analysed by fluorescence microscopy (right panel). Cells displaying an intact network of tubular mitochondria were classified as tubular. When this network was disrupted and mitochondria appeared predominantly spherical or rod-like, they were classified as fragmented. For quantification (left panel), the mitochondrial morphology of at least 500 cells per experiment was determined in a blinded manner. Quantifications were based on triplicates of at least three independent experiments. (D) PrP^C mediates toxic signalling of heterologous PrP^{Sc}. SH-SY5Y cells transiently expressing hamster (ha), human (hu), cervid (ce) or bovine (bo) PrP^C were co-cultured with ScN2a or N2a cells for 16 h. Shown is the percentage of apoptotic cells among transfected cells. Expression of transfected constructs was analysed by western blotting using the anti-PrP antibody 4H11 or 3F4. ***P*<0.005; ****P*<0.0005.

adverse effects on cell viability (Figure 2D). Notably, the same fraction composed of low-molecular weight oligomeric Aβ₄₂ was shown to inhibit LTP in mouse hippocampal slices (Harmeier *et al*, 2009). In addition, it has been shown that oligomers of Aβ₄₂ G33A, in which G33 of the central GXXXG motif is substituted by A (Munter *et al*, 2007, 2010), do not inhibit LTP (Harmeier *et al*, 2009). In line with this observation, oligomers of Aβ₄₂ G33A did not induce a significant increase in cell death in SH-SY5Y cells expressing PrP^C (Figure 2E).

The intrinsically disordered N-terminal domain and the C-terminal GPI anchor of PrP are required to mediate the toxic effects of Aβ

After having established an assay to study PrP^C-mediated toxicity of Aβ, we sought to define the domains of PrP^C required for the transmission of Aβ toxicity. Our previous study revealed that the intrinsically disordered N-terminal domain and the C-terminal GPI anchor of PrP^C are required to mediate toxic signalling of PrP^{Sc} (Rambold *et al*, 2008). Consequently, we tested the ability of Aβ to induce cell




Figure 2 Soluble oligomers of A β induce apoptosis in PrP^C-expressing cells. (A, B) SH-SY5Y cells expressing the constructs indicated were co-cultivated with CHO-7PA2 or CHO cells for 16 h and apoptotic cell death in SH-SY5Y cells was determined as described in Figure 1A. (A) Soluble oligomers of A β secreted by stably transfected cells induce apoptosis in PrP^C-expressing cells. GPI-anchored GFP was used as control. Expression of PrP^C in CHO-7PA2 or CHO cells was analysed by western blotting using the 3F4 antibody (lower panel). (B) Inhibition of γ -secretase interferes with toxic effects of CHO-7PA2 cells. SH-SY5Y cells expressing PrP^C were co-cultured with DAPT-treated (1 μ M) CHO-7PA2 or CHO cells. A β present in the conditioned medium of CHO-7PA2 or CHO cells was analysed by immunoprecipitation followed by western blotting (lower panel). (C) SH-SY5Y cells expressing moPrP^C were co-cultivated with the cell lines indicated. After 16 h of co-cultivation, apoptotic cell death in SH-SY5Y cells was determined as described in Figure 1A. Expression of PrP^C in co-cultured SH-SY5Y cells was analysed by western blotting using the 3F4 antibody (lower panel). (D) Synthetic A β 42 oligomers are toxic to cells expressing PrP^C. SH-SY5Y cells transiently expressing moPrP^C were incubated in the presence of either oligomers (oligo) or high-molecular weight aggregates (HMW) of A β 42 (500 nM each) for 12 h and apoptotic cell death was determined. Expression of PrP^C was analysed by western blotting using the 3F4 antibody (lower panel). (E) Decreased toxicity of oligomers formed by mutant A β 42 G33A. SH-SY5Y cells expressing moPrP^C were incubated in the presence of A β 42 G33A oligomers (500 nM) as described in (C). Expression of PrP^C was monitored by western blotting using the anti-PrP antibody 3F4 (lower panel). ** $P < 0.005$; *** $P < 0.0005$.

death in SH-SY5Y cells either expressing a PrP construct lacking amino acids 27–89 (PrP Δ N) or containing a heterologous C-terminal transmembrane domain instead of the GPI anchor (PrP-CD4). Of note, both PrP mutants are complex glycosylated and present at the outer side of the plasma membrane, similarly to wild-type GPI-anchored PrP^C (Winklhofer *et al*, 2003b). Co-cultivation of PrP-expressing SH-SY5Y cells with A β -secreting CHO-7PA2 cells revealed that PrP-CD4 was not able to mediate toxic effects of A β . Expression of PrP Δ N increased apoptotic cell death, but the effect was significantly reduced compared with full-length

PrP (Figure 3A). In support of the notion that the GPI anchor of PrP is essential to mediate toxic signalling of A β , we could show that oligomers of synthetic A β 42 did not increase apoptotic cell death in PrP-CD4-expressing cells (Figure 3B).

PrP^C mediates toxic signalling of β -sheet-rich conformers of a yeast prion protein

Proteins with prion-like properties have also been identified in fungi (Wickner, 1994). In *Saccharomyces cerevisiae*, the [PSI⁺] prion is formed by the essential translation termination factor Sup35 (Glover *et al*, 1997). The amyloidogenic




Figure 3 Toxic signalling of PrP^C is dependent on the intrinsically disordered N-terminal domain and the C-terminal GPI anchor. (A) Apoptotic activity of soluble A β oligomers secreted by stably transfected CHO-7PA2 cells is compromised in cells expressing PrP mutants. SH-SY5Y cells expressing the PrP constructs indicated were co-cultivated with CHO-7PA2 or CHO cells and apoptotic cell death in SH-SY5Y cells was determined as described in Figure 1A. Expression of PrP constructs was controlled by western blotting using the 3F4 antibody (right panel). (B) Synthetic A β 42 oligomers are not toxic to cells expressing PrP-CD4. SH-SY5Y cells expressing PrP-CD4 were incubated with either oligomers (oligo) or high-molecular weight aggregates (HMW) of A β 42 (500 nM each) for 12 h and apoptotic cell death was determined. ** $P < 0.005$; *** $P < 0.0005$; NS, non-significant. Expression of PrP-CD4 was analysed by western blotting using the 3F4 antibody (right panel).

N-terminal region (N) contains the essential prion-forming determinants, while the C-terminal domain of Sup35 does not contribute to prion behaviour. A charged middle region (M) enhances the solubility of Sup35 in the non-prion form and contributes to the stability of the prion form (i.e. the [PSI⁺] phenotype). Thus, the N and M regions of Sup35 (designated NM) can serve as a separable prion-forming module (Li and Lindquist, 2000). Similarly to PrP in mammals, prion proteins in fungi typically form highly structured β -sheet-rich fibrils, known as amyloids, upon conversion to the infectious prion form (Shorter and Lindquist, 2005). Consequently, we tested the ability of β -sheet-rich conformers of NM to mediate toxic signalling in SH-SY5Y cells via PrP^C. Low-molecular weight NM oligomers, containing dimers, trimers and tetramers, were added to SH-SY5Y cells transiently expressing PrP^C and apoptosis was determined after 7 h. PrP^C-expressing SH-SY5Y cells exhibited a significantly increased rate of apoptotic cell death, whereas the viability of cells expressing PrP-CD4 was not affected (Figure 4A).

PrP^C mediates toxic signalling of β -sheet-rich oligomers formed by designed peptides

The ability of PrP^C to mediate toxic signalling not only of PrP^{Sc}, but also of proteins with completely unrelated primary sequence, such as A β and yeast prion oligomers, raised the possibility that the interaction of PrP^C with the pathogenic protein conformers is primarily determined by structural features. We followed up this hypothesis by including designed peptides in our analysis that are in either an α -helical or β -sheet conformation. The sequences of these peptides were derived from a previous study, showing that binary

patterning of polar and non-polar residues arranged in alternating periodicity can direct proteins to form β -sheet secondary structure (West *et al*, 1999; Wei *et al*, 2003). The peptides (200 nM final concentration) were added to SH-SY5Y cells expressing GPI-anchored wild-type PrP^C (PrP), PrP with a heterologous C-terminal transmembrane domain (PrP-CD4) or PrP Δ N and apoptotic cell death was quantified after 12 h of treatment. Notably, an increase in cell death was only induced in cells expressing GPI-anchored PrP^C by a peptide with β -sheet conformation (β -peptide) (Figure 4B). Incubation with a peptide in α -helical conformation (α -peptide) had no adverse effect on cell viability.

Toxic signalling via PrP^C can be inhibited by a secreted version of the intrinsically disordered N-terminal domain of PrP

The experiments described above revealed that both PrP mutants, PrP-CD4 and PrP Δ N, are impaired in their activity to transduce toxic signalling of β -conformers. To test whether this was due to an impaired binding of the PrP constructs to the β -conformers, we performed indirect immunofluorescence studies and co-immunoprecipitation experiments. Cells expressing wild-type PrP^C (PrP), PrP-CD4 or PrP Δ N were incubated with the synthetic β -peptide and localization of PrP and the β -peptide was visualized by indirect immunofluorescence. These experiments revealed that full-length PrP efficiently mediates the interaction of the β -peptide with the cells, irrespective of whether PrP is linked via a GPI anchor or a transmembrane domain to the plasma membrane (Figure 5A). However, cells expressing PrP Δ N showed a significantly reduced binding to the β -peptide (Figure 5A).




Figure 4 PrP^C mediates toxic signalling of an oligomeric yeast prion protein and of β-sheet-rich oligomers formed by a designed peptide. (A, B) β-sheet-rich conformers of completely different origin induce apoptosis via PrP^C. SH-SY5Y cells expressing the PrP constructs indicated were incubated with either an oligomeric yeast prion protein (NM, 500 nM) for 7 h (A) or a synthetic β-peptide or α-peptide (200 nM each) for 12 h (B). Apoptotic cell death was determined as described in Figure 1A. Expression of the PrP constructs was controlled by western blotting using the 3F4 antibody. ***P* < 0.005; ****P* < 0.0005; NS, non-significant.

To test for a direct interaction between PrP and the β-peptide, we expressed various secreted PrP constructs devoid of a GPI anchor and analysed a possible interaction of secreted PrP with the β-peptide by co-immunoprecipitation experiments. Our previous studies revealed that the isolated N-terminal domain of PrP cannot be expressed as a secreted protein in mammalian cells, since ER import of such a C-terminally truncated PrP construct is abrogated (Heske *et al*, 2004; Miesbauer *et al*, 2009). We therefore generated fusion proteins composed of the isolated N- or C-terminal domains of PrP and the Fc portion of human IgG₁ (Figure 5B). Western blots illustrated that all PrP/Fc fusion proteins are efficiently expressed and secreted into the cell culture media (Figure 5C). The conditioned media from PrP/Fc-expressing cells were mixed with the β-peptide (50 nM) and incubated for 3 h at 4°C. PrP/Fc was purified with protein A-sepharose beads and the immunopellet was analysed by western blotting (Figure 5D). The β-peptide efficiently co-precipitated with PrPN/Fc and to a lesser extent with PrPC/Fc, but not with the Fc part alone (Figure 5D), supporting the conclusion that an interaction with the β-conformers is mainly mediated by the N-terminal domain of PrP.

Finally, we tested the ability of PrPN/Fc to interfere with toxic signalling of Aβ via PrP^C. To this end, we employed the co-cultivation assay of PrP^C-expressing SH-SY5Y cells with Aβ-secreting CHO-7PA2 cells. Indeed, co-expression of the secreted N-terminal domain of PrP (PrPN/Fc) significantly reduced toxic signalling of Aβ via PrP^C (Figure 5E).

Toxic signalling via PrP^C can be inhibited by an oligomer-specific antibody and an NMDA receptor antagonist

The experiments described above indicated that PrP^C is a potent mediator of toxic signalling of β-sheet-rich conformers formed by polypeptides of different origins. Most likely, these polypeptides share common structural features that are recognized or bound by PrP^C. This idea is supported by the finding that the conformation-dependent antibody A11 recognizes toxic oligomers formed by both Aβ- and PrP-peptides (Kayed *et al*, 2003). To test whether A11 could interfere with toxic signalling of Aβ and PrP^C, SH-SY5Y cells expressing GPI-anchored PrP^C were co-cultured with either ScN2a or CHO-7PA2 cells in the presence or absence of A11. This assay




Figure 5 Toxic signalling can be inhibited by a secreted version of the intrinsically disordered N-terminal domain of PrP. **(A)** The N-terminal domain of PrP mediates association with β -peptides. SH-SY5Y cells expressing the indicated PrP constructs were grown on cover slips and incubated for 2 h at 37°C with the β -peptide (200 nM). SH-SY5Y cells were fixed and stained with the polyclonal anti-PrP antibody A7 and the monoclonal anti-myc antibody 4A6, recognizing the β -peptide. Cell nuclei were visualized by DAPI. Scale bar, 10 μ m. **(B)** Schematic presentation of PrP and the PrP/Fc fusion constructs analysed. PrPC/Fc, C-terminal domain of PrP fused to the Fc portion of human IgG₁; PrPN/Fc, N-terminal domain of PrP fused to the Fc portion; SS/Fc, the ER signal sequence of PrP fused to the Fc portion; ER-SS, ER signal sequence; OR, octarepeat; HD, hydrophobic domain; α , α -helical region; β , β -strand; GPI-SS, GPI signal sequence. **(C–E)** A secreted version of the N-terminal domain of PrP associates with β -peptides and interferes with their toxic signalling. **(C)** PrP/Fc fusion proteins are efficiently expressed and secreted into the cell culture medium. N2a cells were transiently transfected with the PrP/Fc fusion constructs. Proteins present in the cell culture medium (M) and cell lysates (L) were analysed by western blotting using the anti-PrP antibody 3F4 or an anti-human IgG antibody. **(D)** The β -peptide co-precipitates with a secreted version of the intrinsically disordered N-terminal domain of PrP. Conditioned medium from PrP/Fc-expressing N2a cells was mixed with the β -peptide (50 nM) and incubated for 3 h at 4°C. PrP/Fc was purified with protein A-sepharose beads and the pellet was analysed by western blotting. Non-transfected cells (n.t.) were used as control. **(E)** PrPN/Fc interferes with toxic signalling of A β via PrP^C. SH-SY5Y cells expressing GPI-anchored PrP^C and the indicated constructs were co-cultivated with CHO-7PA2 or CHO cells and apoptotic cell death in SH-SY5Y cells was determined as described in Figure 1A. ***P* < 0.005; ****P* < 0.0005; NS, non-significant.

revealed that the addition of A11 to the cell culture medium significantly interfered with toxic effects of both A β and PrP^{Sc} on PrP^C-expressing cells (Figure 6A).

To define components downstream of PrP^C involved in toxic signalling of PrP^{Sc} and A β , we performed a co-culture

assay in the presence of memantine, an NMDA receptor antagonist. The rationale behind this approach is based on the findings that NMDA receptors are involved in mediating toxic effects of A β (reviewed in Ondrejcek *et al*, 2010), that NMDA receptor antagonists can block PrP^{Sc}-induced toxicity




Figure 6 An oligomer-specific antibody and an NMDA receptor antagonist prevent toxic signalling of A β oligomers, PrP^{Sc} and β -peptide. (A, B) SH-SY5Y cells expressing GPI-anchored PrP^C were co-cultivated with the cell lines indicated or exposed to β -peptide (200 nM). Apoptotic cell death in SH-SY5Y cells was analysed after 16 h. Cells were co-cultured in the presence of the oligomer-specific polyclonal antibody A11 (1 μ g/ml) (A) or the NMDA receptor antagonist memantine (10 μ M) (B). An unspecific rabbit antiserum (IgG) or water (untreated) served as controls. * P <0.05; ** P <0.005; *** P <0.0005.

(Muller *et al*, 1993) and that PrP^C can attenuate excitotoxicity by inhibiting NMDA receptors (Khosravani *et al*, 2008). PrP^C-expressing SH-SY5Y cells were pre-incubated with memantine for 1 h and either co-cultivated with ScN2a or CHO-7PA2 cells or treated with the β -peptide. Memantine treatment had no effect on the levels of GPI-anchored PrP^C at the plasma membrane or A β in the cell culture media (data not shown); however, the toxic effects of PrP^{Sc}, A β - and β -peptide on PrP^C-expressing cells were significantly reduced by a pharmacological blockage of NMDA receptor activity (Figure 6B).

Primary neurons devoid of PrP^C are less vulnerable to toxic effects of PrP^{Sc} and β -peptides

To validate our findings in mammalian neurons, we prepared primary neuronal cultures from the cortical region of mouse embryos at E14.5–E15.5 days of gestation. Primary neurons, derived either from Prnp^{0/0} mice or from the corresponding wild-type line expressing PrP^C, were co-cultivated with ScN2a cells or their uninfected counterpart (N2a) (Figure 7A). First, cell viability was determined after 4 and 5 days in co-culture. As shown in Figure 7B, the viability of PrP^C-expressing neurons was significantly impaired when co-cultured with ScN2a cells. On the other hand, no differences in viability were observed between PrP^{0/0} neurons cultivated with either ScN2a or N2a cells. In addition, a significant reduction in dendritic length was observed in PrP^C-expressing neurons co-cultivated with ScN2a cells when compared with neurons co-cultivated with N2a cells, while the exposure of PrP^{0/0} neurons to PrP^{Sc} had no impact on dendritic length (Figure 7C). Furthermore, we assessed the distribution of mitochondria in primary neurons.

Exposure of PrP^C-expressing neurons to PrP^{Sc} significantly increased the number of cells harbouring perinuclearly clustered mitochondria. Importantly, in PrP^{0/0} neurons, the number of cells with clustered perinuclear mitochondria was not increased upon co-culturing with ScN2a cells (Figure 7D). In another approach we analysed the toxic effects of the designed β -peptide added to the cell culture medium of primary PrP^{0/0} or PrP^{+/+} cortical neurons. Cortical neurons were cultivated on poly-L-lysine-coated cover slips and β -peptide (2 or 5 μ M) was added on days 4 and 5. At day 6 cells were fixed, permeabilized, stained for β 3 tubulin and examined by fluorescence microscopy. Remarkably, the β -peptide caused neuronal cell loss only in PrP^C-expressing neurons (Figure 7E). In conclusion, these experiments revealed that PrP^C sensitizes primary cortical neurons to toxic effects of β -sheet-rich conformers.

Discussion

Our work revealed that PrP^C localized at the cell surface interacts with and mediates toxic signalling of β -sheet-rich conformers of completely different origin (summarized in the model shown in Figure 8). Both the interaction of PrP^C with β -sheet-rich conformers and induction of pro-apoptotic signalling are dependent on the intrinsically disordered N-terminal domain of PrP^C and the C-terminal GPI anchor.

A critical step in the pathogenesis of prion diseases is a direct interaction of PrP^C with PrP^{Sc}, through which PrP^{Sc} forces PrP^C to adopt the pathogenic conformation. An efficient conformational transition of PrP^C into PrP^{Sc} is highly dependent on the sequence homology between the




Figure 7 Primary neurons lacking PrP^C are less vulnerable to toxic effects of PrP^{Sc} or β-peptide. (A) Schematic model of the co-cultivation assay. Primary neurons were plated on a coated cover slips located in a cell culture dish with either N2a or ScN2a cells. (B–D) Primary neuronal cultures prepared from cortices of PrP^{0/0} or PrP^{+/+} mouse embryos (E14.5–E15.5) were co-cultured with N2a or ScN2a cells for 4 or 5 days. (B) Viability of primary cortical neurons is impaired by co-cultivation with ScN2a cells dependent on PrP^C expression. To analyse neuronal viability, MAP2-positive cells were determined in an area of 1 mm² by fluorescence microscopy. Shown is the percentage of viable neurons co-cultured with ScN2a cells in comparison to primary neurons co-cultured with N2a cells. Viability of neurons co-cultured with N2a cells was set as 100%. (C) Dendritic lengths of PrP^C-expressing primary neurons are reduced after co-cultivation with ScN2a cells. After 4 days in co-culture, dendritic lengths of at least six MAP2-positive primary neurons were quantified using a Zeiss LSM Image program. Shown are relative alterations in dendritic length of primary neurons co-cultured with ScN2a cells in comparison to primary neurons co-cultured with N2a cells (set as 100%). (D) Co-cultivation with ScN2a cells induces perinuclear mitochondrial clustering in PrP^C-expressing primary cortical neurons. Co-cultured primary cortical neurons were stained at day 4 with MitoTracker Red CMXRos to visualize mitochondria and analysed by fluorescence microscopy (right panel). β3 Tubulin was used as a neuronal marker. The cell nuclei are indicated by dotted lines. (Left panel) Shown is the percentage of neurons displaying clustered perinuclear mitochondria. For quantification, mitochondria of at least 500 cells per experiment were determined in a blinded manner. Quantifications were based on triplicates of at least three independent experiments. (E) Treatment of primary cortical neurons with the designed β-peptide causes neuronal cell loss only in PrP^C-expressing neurons. Primary cortical neurons from PrP^{0/0} or PrP^{+/+} mice were incubated with β-peptide (2 or 5 μM) on days 4 and 5. At day 6 neurons were fixed, permeabilized and analysed by indirect immunofluorescence using an anti-β3 tubulin antibody. The experiment was performed in triplicate and fluorescence images of one representative experiment are shown. ***P* < 0.005; ****P* < 0.0005, NS, non-significant.




Figure 8 Putative model of oligomer-induced toxic signalling via PrP^C. PrP^C at the plasma membrane can physically interact with β-sheet-rich conformers of different origin, such as PrP^{Sc}, amyloid β (Aβ), yeast prion protein (Sup35) or designed peptides (β-peptide) (step 1). Interaction of PrP^C with the β-sheet-rich conformers can be inhibited by an oligomer-specific antibody (A11), or a secreted version of the intrinsically disordered N-terminal domain of PrP (PrPN/Fc). The PrP/β-sheet complex can then induce apoptotic signalling (step 2). Toxic signalling via PrP^C is dependent on the GPI anchor of PrP and can be inhibited by the NMDA receptor antagonist memantine. Since PrP^C has no direct contact to the cytosolic compartment, it is plausible to assume that intracellular signal transmission involves additional cellular factors, such as the NMDA receptor (NMDAR), or a different transmembrane protein (X), or a cytosolic protein associated with lipid rafts (Y).

two isoforms. As a consequence, an inter-species transmission of prion diseases is extremely difficult, a phenomenon denoted as species or transmission barrier (Prusiner *et al*, 1990). Remarkably, our data revealed that PrP^C can mediate toxic signalling of PrP^{Sc} without being converted into PrP^{Sc}: hamster, human, cervid or bovine PrP^C-mediated toxic signalling of mouse PrP^{Sc} as efficient as mouse PrP^C, whereas mouse PrP^{Sc} could convert these heterologous PrP^C species into PrP^{Sc}. We also showed that β-sheet-rich conformers of completely different origin, formed by the Aβ-peptide, a yeast prion protein or a designed peptide, can induce apoptotic signalling through PrP^C. Taken together, these results strongly suggest that PrP^C has an intrinsic ability to interact with β-sheet-rich conformers independent of their primary sequence. Notably, the oligomer-specific antibody A11 could block PrP^C-mediated toxic signalling of both PrP^{Sc} and Aβ, supporting the idea that different toxic β-conformers share common structural features, which are recognized and bound by PrP^C.

Employing different PrP mutants, such as PrP-CD4 and PrPAN, we were able to define domains of PrP^C required for signalling activity. What could be the role of the C-terminal GPI anchor and the intrinsically disordered N-terminal domain in toxic signal transduction via PrP^C? PrP-CD4, which is anchored to the plasma membrane by a heterologous C-terminal transmembrane domain, binds to β-peptides, but cannot mediate toxic effects. In contrast to GPI-anchored PrP^C, PrP-CD4 does not localize to detergent-insoluble microdomains at the plasma membrane (Rambold *et al*, 2008). These data suggest that targeting of PrP^C to detergent-insoluble microdomains is a pre-requisite to induce intracellular signalling pathways. PrPAN lacks amino acids 27–89 and thereby most of the intrinsically disordered domain. Notably, intrinsically disordered domains have been shown to be involved in protein–protein interactions (Tompa *et al*, 2009). Indeed, our experiments revealed that the interaction of β-conformers with PrP^C-expressing cells is reduced by deleting the N-terminal domain of PrP. Moreover, a secreted version of the N-terminal domain of PrP (PrPN/Fc) efficiently interacted with β-conformers and interfered with toxic signalling via PrP^C.

Interestingly, PrP^C can also activate neuroprotective signalling pathways (McLennan *et al*, 2004; Shyu *et al*, 2005; Spudich *et al*, 2005; Weise *et al*, 2006; Mitteregger *et al*, 2007), and both the C-terminal GPI anchor and the unstructured N-terminal domain are required for this physiological activity (Mitteregger *et al*, 2007; Rambold *et al*, 2008). Thus, PrP^C might act as a signalling molecule at the cell surface to promote stress-protective signalling under physiological conditions, which can be switched to toxic signalling through an interaction with β-sheet-rich conformers. Similarly to other GPI-anchored proteins involved in signal transduction, PrP^C most probably acts as a co-receptor in concert with a transmembrane protein to transduce the signal into the cell. Unfortunately, little is known about signalling pathways downstream of PrP^C. While different PrP^C-dependent pathways and -interacting proteins have been described (reviewed in Caughey and Baron, 2006), experimental evidence for a direct role of these components in PrP^{Sc}- or Aβ-induced toxicity is missing. Our study presents evidence that Aβ-, PrP^{Sc}- and β-peptide-induced toxicity is significantly suppressed by pharmacologically inhibiting NMDA-type glutamate receptor activity. PrP^C has previously been reported to attenuate excitotoxicity by inhibiting NMDA receptors (Khosravani *et al*, 2008). In addition, there is increasing evidence that excitotoxicity mediated by NMDA receptors has a crucial role in the pathogenesis of AD. For example, it has been shown that Aβ-mediated spine loss requires activity of NMDA-type glutamate receptors (Shankar *et al*, 2007). All in all, our data provide a scientific rationale for a beneficial effect of NMDA receptor antagonists in a more wider range of neurodegenerative disorders associated with the formation of β-sheet-rich conformers, given that they are applied at an early stage of the disease before irreversible neuronal loss has occurred.

Materials and methods

Plasmids, antibodies and reagents

All expression constructs have been described previously (Rambold *et al*, 2008). Amino acid numbers refer to mouse Prion Protein sequence (GenBank™ accession number NP 035300), hamster

Prion Protein sequence (GenBank™ accession number P04273), human Prion Protein sequence (GenBank™ accession number AAA60182), cervid Prion Protein sequence (GenBank™ accession number ABW79904) or bovine Prion Protein sequence (GenBank™ accession number AAQ64648). As transfection marker the EYFP-C1 vector (Clontech) was used. The following antibodies were used: mouse monoclonal anti-PrP 3F4 antibody (Kascsak *et al*, 1987), mouse monoclonal anti-PrP 4H11 antibody (Ertmer *et al*, 2004), rabbit polyclonal anti-PrP antibody A7 (Winklhofer *et al*, 2003a), mouse monoclonal anti-myc 4A6 antibody (Millipore), rabbit polyclonal anti-active caspase-3 antibody (Promega), mouse monoclonal anti-MAP2 antibody (Sigma-Aldrich), rabbit polyclonal anti-β3 tubulin antibody (Abcam), fluorescent dye-labelled anti-rabbit IgG antibody Alexa Flour 555 (Invitrogen), fluorescent dye-labelled anti-mouse IgG antibody Alexa Flour 555 (Invitrogen), fluorescent dye-labelled anti-mouse IgG antibody Alexa Flour 488 (Invitrogen), fluorescent dye-labelled anti-rabbit IgG antibody Alexa Flour 488 (Invitrogen), horseradish peroxidase-conjugated anti-mouse IgG antibody (Amersham, Promega), horseradish peroxidase-conjugated anti-human IgG antibody (Promega), A11 polyclonal antibody (Kayed *et al*, 2003), rat monoclonal anti-Aβ antibody 2D8 (Shirovani *et al*, 2007), rabbit polyclonal anti-Aβ antibody 3552 (Yamasaki *et al*, 2006). The following reagents were used: DAPT (*N*-[*N*-(3,5-difluorophenacetyl)-*L*-alaninyl]-*S*-phenylglycine *t*-butyl ester), memantine hydrochlorid (Sigma), MitoTracker Red CMXRos (Invitrogen), Proteinase K (Sigma-Aldrich), TO-PRO-3 iodide (642/661) (Invitrogen), poly-L-lysine hydrobromide (Sigma-Aldrich). The mounting medium Mowiol (Calbiochem) was supplemented with DAPI (4',6-diamidino-2-phenylindole; Sigma-Aldrich).

Cell lysis, immunoprecipitation and western blot analysis

As described earlier (Tatzelt *et al*, 1996), cells were washed twice with cold phosphate-buffered saline (PBS), scraped off the plate and lysed in cold detergent buffer (0.5% Triton X-100, 0.5% sodium deoxycholat in PBS). Post-nuclear supernatants or secreted and TCA-precipitated proteins were boiled with Laemmli sample buffer and analysed by western blotting as described previously (Winklhofer and Tatzelt, 2000). Aβ oligomers in conditioned medium of CHO-7PA2 were analysed by immunoprecipitation with the polyclonal antibody 3552 followed by western blotting using the monoclonal antibody 2D8.

Cell culture, transfection, co-cultivation, γ-secretase inhibitor experiments and NMDA receptor antagonist treatment

Cells were cultivated and transfected as described earlier (Winklhofer *et al*, 2003b). Stably transfected Chinese hamster ovary cells (CHO-7PA2) that express the familial AD mutation V717F in the amyloid precursor protein APP₇₅₁ and secrete Aβ were described earlier (Podlisny *et al*, 1995). The human SH-SY5Y cell line (DSMZ number ACC 209) is a subline of bone marrow biopsy-derived SK-N-SH cells. For co-cultivation experiments, SH-SY5Y cells were grown on glass cover slips and transfected with Lipofectamine (Invitrogen). Two hours after transfection, cover slips were transferred into dishes containing a 90% confluent cell layer of ScN2a, N2a, CHO-7PA2 or CHO cells. For co-cultivation, SH-SY5Y cells were cultivated in a combination of CHO-7PA2/ScN2a, CHO-7PA2/N2a, CHO/ScN2a or CHO/N2a cells. After 16 h of co-cultivation, either apoptotic cell death or mitochondrial morphology was analysed (see below). To block Aβ generation, CHO-7PA2 cells were pre-treated for 24 h with DAPT (1 μM) before co-cultivation. To interfere with NMDA receptor activation, transfected SH-SY5Y cells were pre-treated for 1 h with memantine (10 μM) before co-cultivation. Memantine (10 μM) was also present during the co-cultivation.

Aβ42-peptides, SEC

As described earlier (Harmeier *et al*, 2009), synthetic Aβ42 and Aβ42 G33A peptides (PSL, Peptide Speciality Laboratories GmbH) were monomerized in formic acid and subsequently re-dissolved in water containing 0.1% ammonia. The dissolved peptides were loaded onto a Superose 12 (10/300 HR) column (Amersham Bioscience), and 1 ml fractions were collected in PBS as running buffer at a flow rate of 0.5 ml/min. Peptide concentrations were determined using spectrophotometric methods, and the aggregation state was further determined by SDS-PAGE and western blotting. Equal amounts (final concentration 500 nM) of oligomers or high-molecular weight aggregates of Aβ42 were applied on transiently transfected SH-SY5Y cells, and apoptotic cell death was analysed after 12 h.

Yeast prion protein NM: generation and treatment conditions

The NM stock solution was prepared as described (Scheibel *et al*, 2001). For the preparation of ditryrosine-crosslinked NM oligomers, the NM stock solution was diluted to a final concentration of 15 μM in 20 mM Tris, 150 mM sodium chloride, 0.5 mM coppers and 0.5 mM zinc sulphate. To this solution, 0.1 mg/ml of superoxide dismutase (Sigma-Aldrich) was added. Using a Fluorolog fluorimeter (HORIBA Jobin Yvon), the prepared NM solution was crosslinked via excitation at 280 nm for 2 h at room temperature while stirring. Subsequently, an equivalent volume of 6 M guanidinium hydrochloride (Fluka) was added to the solution. For isolation of low-molecular weight crosslinked NM oligomers, the NM solution was first filtered through a 100-kDa Amicon Ultra Centrifugal Filter (Millipore) and the flow through was concentrated using a 3-kDa Amicon Ultra Centrifugal Filter (Millipore). Transiently transfected SH-SY5Y cells were treated with either the low-molecular weight crosslinked NM (final concentration 500 nM) or buffer (10 mM Tris, 75 mM sodium chloride, 3 M guanidinium-HCl, 250 μM copper sulphate, 250 μM zinc sulphate) as control, and apoptotic cell death was analysed after 7 h of treatment.

β- and α-peptides: generation and treatment conditions

The N-terminally myc-tagged β-sheet peptide (ATGTGTGAACAAA GCTTATTTCTGAAGAAGACTTGGGTATGCAGATCTCCATGGATTATGA GATCAAGTTCACGGTGTATGGTATAACTTCGACCTCAACCTCGACGAT TCTGGTGTGATCTTCAACTTCAGATCCGCGGCCGGGTGGCCGTGCC ACGTCCACATCCACAGTAGTAGTGTAAGGTTCGACTTCCACGTCAACAA CGACGGCGCGGATGTTGAAGTAAAATGCACTAG) coding for MCEQK LISEEDLGMQISMDYEIKFHGDDGDNFDLNLDDSGGDLQLQIRPGGRVH VHHSSSGKVDHFVNNDGGDVEVKMH and the α-helical peptide (ATGTGTGAACAAAAGCTTATTTCTGAAGAAGACTTGGGTATGTATGGC AAGTTGAACGACCTGCTGGAAGACTTGCAGAGGTGCTGGAAGAACCTC CACAAAACCTGGCACGGTGGCAAAGACAACCTGCACGACGTCGACAA CCAGTTGCAGAACGTCATCGAAGACATCCACGACTTCATGCAAGCGG TGGCAGCGCGGCAAGCTGCAAGAGATGATGAAAGAGTTCACACAGG TGTGGACGAACCAACAACCCTTGCAGCGGTAAACACACCGTGC ACCACATCGAACAAAACATCAAGAGACTTCCACCCTGGAAAGAGC TCGTCCACCGTTAG) coding for MCEQKLISEEDLGMQYKLNLDLEDL QEVVLKLNHKNWHGGKDNLDHVDNHLQNVIEDIHFDMQGGSGGKQLQ EMMKEFQVQLDELNNHLQGGKHVTVHHIEQNIKEIFHHLEELVHR were expressed in *Escherichia coli* BL21 (DE3) using the pTrcHis vector or the pProEx HTb, respectively. The β-sheet peptide was mainly present in inclusion bodies and purified using a MonoQ 10/100 HR 16/10 column (0–1 M NaCl gradient in 8 M urea) followed by SEC (Sephacryl S-300, HiPreP 26/60) in 8 M urea, 100 mM NaCl (pH 7.5). After dialysis against 100 mM NaCl, 25 mM sodium phosphate, refolded soluble oligomers were purified by SEC. The α-helical peptide was purified under non-denaturing conditions using Ni-NTA agarose (Qiagen). β- or α-peptides (200 nM each) were applied to transiently transfected SH-SY5Y cells and apoptotic cell death was analysed after 12 h. To interfere with NMDA receptor activation, transfected SH-SY5Y cells were pre-treated for 1 h with memantine (10 μM) before application of β-peptide. To investigate a possible direct interaction with PrP, the β- or α-peptides (final concentration 200 nM) were added to transiently transfected SH-SY5Y cells. After 2 h incubation, SH-SY5Y cells were fixed and stained with the polyclonal anti-PrP antibody A7 and the monoclonal anti-myc antibody 4A6 overnight at 4°C, followed by an incubation with the fluorescently labelled secondary antibodies Alexa Flour 555 and Alexa Flour 488 for 1 h at room temperature. Cells were then mounted onto glass slides and examined by fluorescence microscopy using a Zeiss Axiovert 200 M microscope (Carl Zeiss). Cell nuclei were visualized by DAPI. In addition, conditioned media from PrP/Fc-expressing N2a cells were mixed with the β-peptide (50 nM final concentration) and incubated for 3 h at 4°C. PrP/Fc was purified with protein A-sepharose beads and the pellet was analysed by western blotting using anti-myc 4A6 antibody.

Preparation, co-cultivation and treatment of primary cortical neurons

Prnp^{0/0} mice (Büeler *et al*, 1992) were on a C57-129Sv background or had been backcrossed on a C57Bl/6 background for over 10 generations. Primary cortical neurons were prepared from C57-129Sv/PrP^{+/+} or C57-129Sv/PrP^{0/0} mouse embryos at E14.5–E15.5 days of gestation. Briefly, cortical neurons were cultured in 3.5 cm dishes (1.4 million cells/dish) containing poly-L-lysine (100 μg/ml)-coated cover slips in Neurobasal medium (Invitrogen) supplemented with basic fibroblast growth factor (10 ng/ml), nerve

growth factor (10 ng/ml), B27 supplement minus AO 50 × (1 ×) and L-glutamin (0.5 mM). For co-cultivation experiments, cortical neurons were seeded on poly-L-lysine-coated cover slips placed into a dish with a 20% confluent cell layer of either ScN2a or N2a cells. After 4 or 5 days in co-culture, primary neurons on glass cover slips were fixed with 3.7% paraformaldehyde for 20 min, washed and permeabilized with 0.2% Triton X-100 in PBS for 10 min at room temperature. Fixed cells were incubated with an anti-MAP2 antibody overnight at 4°C, followed by an incubation with the fluorescently labelled secondary antibody Alexa Fluor 488 for 1 h at room temperature. Cell nuclei were visualized by TO-PRO-3 iodide (642/661). To analyse cell viability, MAP2-positive cells were examined by fluorescence microscopy using a Zeiss Axiovert 200 M microscope (Carl Zeiss). Dendritic lengths of at least six primary neurons co-cultured for 4 days were quantified in an area of 1 mm² using a Zeiss LSM Image program. To examine mitochondrial morphology, cortical neurons were first incubated with MitoTracker Red CMXRos (Invitrogen) (final concentration 250 nM) before fixation. The number of cells with clustered perinuclear mitochondria out of at least 500 cells was determined. β3 Tubulin was used as a neuronal marker. All experiments were performed in a blinded manner, and quantifications were based on at least three independent experiments. To analyse toxicity of β-peptides, cortical neurons were cultivated on poly-L-lysine-coated cover slips. β-Peptides (2 or 5 μM) were added at days 4 and 5. At day 6, cells were fixed, permeabilized and incubated with an anti-β3 tubulin antibody overnight at 4°C, followed by an incubation with the fluorescently labelled secondary antibody Alexa Fluor 488 for 1 h at room temperature. Cells were examined by fluorescence microscopy using a Zeiss Axiovert 200 M microscope (Carl Zeiss). Expression of PrP in neuronal cultures was analysed by immunofluorescence using the monoclonal anti-PrP antibody 4H11 and a polyclonal anti-β3 tubulin antibody as neuronal marker. In addition, western blotting was performed using the monoclonal anti-PrP antibody 4H11 (Supplementary Figure S3A).

Apoptosis assay and mitochondria morphology assay

In all, 16 h after co-cultivation, SH-SY5Y cells were fixed on glass cover slips with 3.7% paraformaldehyde for 20 min, washed and permeabilized with 0.2% Triton X-100 in PBS for 10 min at room temperature. Fixed cells were incubated with an anti-active caspase-3 antibody overnight at 4°C, followed by an incubation with the fluorescently labelled secondary antibody Alexa Fluor 555 for 1 h at room temperature. Cells were then mounted onto glass slides and examined by fluorescence microscopy using a Zeiss Axiovert 200 M microscope (Carl Zeiss). The numbers of cells positive for activated caspase-3 or cells with fragmented nuclei out of at least 1000 transfected cells were determined in a blinded manner. To analyse

mitochondrial morphology after co-cultivation, SH-SY5Y cells were incubated with MitoTracker Red CMXRos (Invitrogen) (final concentration 100 nM) before fixation. The number of cells with non-tubular/fragmented mitochondria out of at least 500 transfected cells was determined. All quantifications were based on at least three independent experiments.

PrP^{Sc} propagation assay

After co-cultivation, cells were scraped off the plate, the cell pellets were washed twice with cold PBS and lysed in cold detergent buffer (0.5% Triton X-100, 0.5% sodium deoxycholate in PBS). The lysates were centrifuged to generate detergent-soluble and -insoluble fractions. The insoluble fractions were digested with proteinase K (10 μg/ml final concentration) for 0.5 h at 37°C. The reaction was terminated by the addition of PMSF and then both fractions were adjusted to 0.5% Sarkosyl. PrP was immunoprecipitated using the monoclonal anti-PrP antibodies 3F4 (specific for the transfected PrP in the SH-SY5Y cells) or 4H11. Immunoprecipitated proteins were boiled in Laemmli sample buffer and analysed by western blotting using the monoclonal anti-PrP antibody 4H11.

Statistical analysis

Quantifications were based on at least three independent experiments. Data were shown as means ± s.e. Statistical analysis was performed using Student's *t*-test. *P*-values are as follows: **P* < 0.05; ***P* < 0.005; ****P* < 0.0005.

Supplementary data

Supplementary data are available at *The EMBO Journal* Online (<http://www.embojournal.org>).

Acknowledgements

This work was supported by grants from the Deutsche Forschungsgemeinschaft (SFB 596, GRK 1123), the Max Planck Society and the BMBF (BioDisc, DIP5.1), the HHMI (to SL) and the NIH (GM25874 to SL and NS067782 to JLG). JT gratefully acknowledges support by the German Center for Neurodegenerative Diseases (DZNE). We are grateful to Dominic Walsh and Dennis Selkoe for providing the CHO-7PA2 cells, Charlie Glabe for the A11 antibody and Christian Haass for the Aβ antibodies and DAPT.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Brandner S, Isenmann S, Raeber A, Fischer M, Sailer A, Kobayashi Y, Marino S, Weissmann C, Aguzzi A (1996) Normal host prion protein necessary for scrapie-induced neurotoxicity. *Nature* **379**: 339–343
- Büeler H, Aguzzi A, Sailer A, Greiner R-A, Autenried P, Aguet M, Weissmann C (1993) Mice devoid of PrP are resistant to scrapie. *Cell* **73**: 1339–1347
- Büeler H, Fischer M, Lang Y, Bluethmann H, Lipp H-P, DeArmond SJ, Prusiner SB, Aguet M, Weissmann C (1992) Normal development and behaviour of mice lacking the neuronal cell-surface PrP protein. *Nature* **356**: 577–582
- Caughey B, Baron GS (2006) Prions and their partners in crime. *Nature* **443**: 803–810
- Chesebro B (2003) Introduction to the transmissible spongiform encephalopathies or prion diseases. *Br Med Bull* **66**: 1–20
- Chesebro B, Trifilo M, Race R, Meade-White K, Teng C, LaCasse R, Raymond L, Favara C, Baron G, Priola S, Caughey B, Masliah E, Oldstone M (2005) Anchorless prion protein results in infectious amyloid disease without clinical scrapie. *Science* **308**: 1435–1439
- Cleary JP, Walsh DM, Hofmeister JJ, Shankar GM, Kuskowski MA, Selkoe DJ, Ashe KH (2005) Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function. *Nat Neurosci* **8**: 79–84
- Collinge J (2001) Prion diseases of humans and animals: their causes and molecular basis. *Annu Rev Neurosci* **24**: 519–550
- Collinge J, Clarke AR (2007) A general model of prion strains and their pathogenicity. *Science* **318**: 930–936
- De Strooper B (2010) Proteases and proteolysis in Alzheimer disease: a multifactorial view on the disease process. *Physiol Rev* **90**: 465–494
- Ertmer A, Gilch S, Yun SW, Flechsig E, Klebl B, Stein-Gerlach M, Klein MA, Schatzl HM (2004) The tyrosine kinase inhibitor STI571 induces cellular clearance of PrP^{Sc} in prion-infected cells. *J Biol Chem* **279**: 41918–41927
- Gimbel DA, Nygaard HB, Coffey EE, Gunther EC, Lauren J, Gimbel ZA, Strittmatter SM (2010) Memory impairment in transgenic Alzheimer mice requires cellular prion protein. *J Neurosci* **30**: 6367–6374
- Glover JR, Kowal AS, Schirmer EC, Patino MM, Liu JJ, Lindquist S (1997) Self-seeded fibers formed by Sup35, the protein determinant of [PSI⁺], a heritable prion-like factor of *S. cerevisiae*. *Cell* **89**: 811–819
- Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* **8**: 101–112
- Harmeier A, Wozny C, Rost BR, Munter LM, Hua H, Georgiev O, Beyermann M, Hildebrand PW, Weise C, Schaffner W, Schmitz D,

- Multhaup G (2009) Role of amyloid-beta glycine 33 in oligomerization, toxicity, and neuronal plasticity. *J Neurosci* **29**: 7582–7590
- Heske J, Heller U, Winklhofer KF, Tatzelt J (2004) The C-terminal domain of the prion protein is necessary and sufficient for import into the endoplasmic reticulum. *J Biol Chem* **279**: 5435–5443
- Ilieva H, Polymenidou M, Cleveland DW (2009) Non-cell autonomous toxicity in neurodegenerative disorders: ALS and beyond. *J Cell Biol* **187**: 761–772
- Kacsak RJ, Rubenstein R, Merz PA, Tonna-DeMasi M, Fersko R, Carp RI, Wisniewski HM, Diringer H (1987) Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins. *J Virol* **61**: 3688–3693
- Kayed R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG (2003) Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* **300**: 486–489
- Khosravani H, Zhang Y, Tsutsui S, Hameed S, Altier C, Hamid J, Chen L, Villemare M, Ali Z, Jirik FR, Zamponi GW (2008) Prion protein attenuates excitotoxicity by inhibiting NMDA receptors. *J Cell Biol* **181**: 551–565
- Lacor PN, Buniel MC, Chang L, Fernandez SJ, Gong Y, Viola KL, Lambert MP, Velasco PT, Bigio EH, Finch CE, Krafft GA, Klein WL (2004) Synaptic targeting by Alzheimer's-related amyloid beta oligomers. *J Neurosci* **24**: 10191–10200
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL (1998) Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* **95**: 6448–6453
- Lauren J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM (2009) Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* **457**: 1128–1132
- Li L, Lindquist S (2000) Creating a protein-based element of inheritance. *Science* **287**: 661–664
- Mallucci G, Dickinson A, Linehan J, Klohn PC, Brandner S, Collinge J (2003) Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* **302**: 871–874
- McLennan NF, Brennan PM, McNeill A, Davies I, Fotheringham A, Rennison KA, Ritchie D, Brannan F, Head MW, Ironside JW, Williams A, Bell JE (2004) Prion protein accumulation and neuroprotection in hypoxic brain damage. *Am J Pathol* **165**: 227–235
- Miesbauer M, Pfeiffer NV, Rambold AS, Muller V, Kiachopoulos S, Winklhofer KF, Tatzelt J (2009) Alpha-helical domains promote translocation of intrinsically disordered polypeptides into the endoplasmic reticulum. *J Biol Chem* **284**: 24384–24393
- Mitteregger G, Vosko M, Krebs B, Xiang W, Kohlmannspurger V, Nolting S, Hamann GF, Kretschmar HA (2007) The role of the octarepeat region in neuroprotective function of the cellular prion protein. *Brain Pathol* **17**: 174–183
- Moore RA, Vorberg I, Priola SA (2005) Species barriers in prion diseases—brief review. *Arch Virol Suppl* **187**: 202
- Muller WE, Ushijima H, Schroder HC, Forrest JM, Schatton WF, Rytik PG, Heffner-Laue M (1993) Cytoprotective effect of NMDA receptor antagonists on prion protein (PrionSc)-induced toxicity in rat cortical cell cultures. *Eur J Pharmacol* **246**: 261–267
- Munter LM, Botev A, Richter L, Hildebrand PW, Althoff V, Weise C, Kaden D, Multhaup G (2010) Aberrant amyloid precursor protein (APP) processing in hereditary forms of Alzheimer disease caused by APP familial Alzheimer disease mutations can be rescued by mutations in the APP GxxxG motif. *J Biol Chem* **285**: 21636–21643
- Munter LM, Voigt P, Harmeier A, Kaden D, Gottschalk KE, Weise C, Pipkorn R, Schaefer M, Langosch D, Multhaup G (2007) GxxxG motifs within the amyloid precursor protein transmembrane sequence are critical for the etiology of Abeta42. *EMBO J* **26**: 1702–1712
- Ondrejcek T, Klyubin I, Hu NW, Barry AE, Cullen WK, Rowan MJ (2010) Alzheimer's disease amyloid beta-protein and synaptic function. *Neuromolecular Med* **12**: 13–26
- Palop JJ, Mucke L (2010) Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat Neurosci* **13**: 812–818
- Podlisny MB, Ostaszewski BL, Squazzo SL, Koo EH, Rydell RE, Teplow DB, Selkoe DJ (1995) Aggregation of secreted amyloid beta-protein into sodium dodecyl sulfate-stable oligomers in cell culture. *J Biol Chem* **270**: 9564–9570
- Prusiner SB, Scott M, Foster D, Pan K-M, Groth D, Mirenda C, Torchia M, Yang S-L, Serban D, Carlson GA, Hoppe PC, Westaway D, DeArmond SJ (1990) Transgenic studies implicate interactions between homologous PrP isoforms in scrapie prion replication. *Cell* **63**: 673–686
- Prusiner SB, Scott MR, DeArmond SJ, Cohen FE (1998) Prion protein biology. *Cell* **93**: 337–348
- Rambold AS, Müller V, Ron U, Ben-Tal N, Winklhofer KF, Tatzelt J (2008) Stress-protective activity of prion protein is corrupted by scrapie-prions. *EMBO J* **27**: 1974–1984
- Scheibel T, Kowal AS, Bloom JD, Lindquist SL (2001) Bidirectional amyloid fiber growth for a yeast prion determinant. *Curr Biol* **11**: 366–369
- Scott M, Foster D, Mirenda C, Serban D, Coufal F, Walchli M, Torchia M, Groth D, Carlson G, DeArmond SJ, Westaway D, Prusiner SB (1989) Transgenic mice expressing hamster prion protein produce species-specific scrapie infectivity and amyloid plaques. *Cell* **59**: 847–857
- Scott MR, Safar J, Telling G, Nguyen O, Groth D, Torchia M, Koehler R, Tremblay P, Walther D, Cohen FE, DeArmond SJ, Prusiner SB (1997) Identification of a prion protein epitope modulating transmission of bovine spongiform encephalopathy prions to transgenic mice. *Proc Natl Acad Sci USA* **94**: 14279–14284
- Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL (2007) Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci* **27**: 2866–2875
- Shirovani K, Tomioka M, Kremmer E, Haass C, Steiner H (2007) Pathological activity of familial Alzheimer's disease-associated mutant presenilin can be executed by six different gamma-secretase complexes. *Neurobiol Dis* **27**: 102–107
- Shorter J, Lindquist S (2005) Prions as adaptive conduits of memory and inheritance. *Nat Rev Genet* **6**: 435–450
- Shyu WC, Lin SZ, Chiang MF, Ding DC, Li KW, Chen SF, Yang HI, Li H (2005) Overexpression of PrPC by adenovirus-mediated gene targeting reduces ischemic injury in a stroke rat model. *J Neurosci* **25**: 8967–8977
- Spudich A, Frigg R, Kilic E, Kilic U, Oesch B, Raeber A, Bassetti CL, Hermann DM (2005) Aggravation of ischemic brain injury by prion protein deficiency: role of ERK-1/-2 and STAT-1. *Neurobiol Dis* **20**: 442–449
- Tatzelt J, Prusiner SB, Welch WJ (1996) Chemical chaperones interfere with the formation of scrapie prion protein. *EMBO J* **15**: 6363–6373
- Tomba P, Fuxreiter M, Oldfield CJ, Simon I, Dunker AK, Uversky VN (2009) Close encounters of the third kind: disordered domains and the interactions of proteins. *Bioessays* **31**: 328–335
- Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ (2002) Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation *in vivo*. *Nature* **416**: 535–539
- Wei Y, Liu T, Sazinsky SL, Moffet DA, Pelczar I, Hecht MH (2003) Stably folded *de novo* proteins from a designed combinatorial library. *Protein Sci* **12**: 92–102
- Weise J, Sandau R, Schwarting S, Crome O, Wrede A, Schulz-Schaeffer W, Zerr I, Bahr M (2006) Deletion of cellular prion protein results in reduced Akt activation, enhanced postsynaptic caspase-3 activation, and exacerbation of ischemic brain injury. *Stroke* **37**: 1296–1300
- Weissmann C, Fischer M, Raeber A, Büeler H, Sailer A, Shmerling D, Rüllicke T, Brandner S, Aguzzi A (1996) The role of PrP in pathogenesis of experimental scrapie. *Cold Spring Harb Symp Quant Biol* **61**: 511–522
- West MW, Wang W, Patterson J, Mancias JD, Beasley JR, Hecht MH (1999) *De novo* amyloid proteins from designed combinatorial libraries. *Proc Natl Acad Sci USA* **96**: 11211–11216
- Wickner RB (1994) [URE3] as an altered URE2 protein: evidence for a prion analog in *Saccharomyces cerevisiae* [see comments]. *Science* **264**: 566–569
- Winklhofer KF, Heller U, Reintjes A, Tatzelt J (2003a) Inhibition of complex glycosylation increases formation of PrP^{Sc}. *Traffic* **4**: 313–322

- Winklhofer KF, Heske J, Heller U, Reintjes A, Muranji W, Moarefi I, Tatzelt J (2003b) Determinants of the *in vivo*-folding of the prion protein: a bipartite function of helix 1 in folding and aggregation. *J Biol Chem* **278**: 14961–14970
- Winklhofer KF, Tatzelt J (2000) Cationic lipopolyamines induce degradation of PrPSc in scrapie-infected mouse neuroblastoma cells. *Biol Chem* **381**: 463–469
- Winklhofer KF, Tatzelt J, Haass C (2008) The two faces of protein misfolding: gain and loss of function in neurodegenerative diseases. *EMBO J* **27**: 336–349
- Yamasaki A, Eimer S, Okochi M, Smailowska A, Kaether C, Baumeister R, Haass C, Steiner H (2006) The GxGD motif of presenilin contributes to catalytic function and substrate identification of gamma-secretase. *J Neurosci* **26**: 3821–3828