

The cockle *Cerastoderma edule* at Northeast Atlantic shores: genetic signatures of glacial refugia

Manuela Krakau · Sabine Jacobsen ·
Kurt Thomas Jensen · Karsten Reise

Received: 7 July 2011 / Accepted: 16 September 2011 / Published online: 11 October 2011
© Springer-Verlag 2011

Abstract The cockle *Cerastoderma (Cardium) edule* (L.) ranges from NW Africa to northern Scandinavia. Abundance in shallow coastal sediment is often high, and it attracts commercial harvest. In this study, a complex genetic pattern has been revealed by mitochondrial DNA in 383 individuals from 19 sampling sites. Parsimony network analysis of 79 haplotypes identified two dominant central haplotypes separated by low divergence. One is characteristic for a homogenous southwestern group of populations from Africa to the British Isles. The other is characteristic for a heterogeneous northern group with a deviant Arctic population. At the entrance of the Baltic Sea, a mixture zone of the dominant haplotypes was found. The estimated population expansion time for the northern haplotype group predates that of the southwestern one, suggesting northern glacial refugia and a subsequent southern expansion of *C. edule* populations.

Introduction

Quaternary glaciation cycles and corresponding extinctions, contractions into glacial refugia and interglacial expansions of individual species have laid the foundation for present species assemblages at temperate and high latitudes (Hewitt 2000; Martin and McKay 2004; Hoffmann and Sgrò 2011). Large-scale population genetic analysis may detect signatures of these developments. In the simplest scenario with a single glacial refugium below the glaciated area, a linear decrease in genetic diversity toward higher latitudes can be expected. In the Northeast Atlantic, there are, however, several examples where marine species do not conform to this. Two species of seagrass, for example, show highest allelic diversity in the North Sea region (Coyer et al. 2004; Olsen et al. 2004). Several species of benthic macroalgae and invertebrates seem to have survived glaciations in periglacial or cryptic northern refugia (Roman and Palumbi 2004; Hoarau et al. 2007; Maggs et al. 2008; Luttikhuisen et al. 2008; Olsen et al. 2010; Coyer et al. 2011).

In benthic bivalves, no clear evidence for northern glacial refugia has been found so far. However, trans-Arctic pathways from Pacific coasts and dispersal across the northern Atlantic have been inferred from genetic analyses for bivalves. Notable examples are the circumpolar *Macoma balthica* complex (Luttikhuisen et al. 2003; Nikula et al. 2007) and the closely related mussels *Mytilus edulis*, *M. trossulus* and *M. galloprovincialis* (Riginos and Henzler 2008; Brannock and Hilbish 2010; Väinölä and Strelkov 2011). Patterns in the latter may have been further complicated by human-mediated translocations. Such pathways have been genetically reconstructed for various introduced alien species (e.g., Darling et al. 2008; Reitzel et al. 2008; Reusch et al. 2010; Lejeune et al. 2011). In the lagoon

Communicated by S. Uthicke.

Electronic supplementary material The online version of this article (doi:10.1007/s00227-011-1802-8) contains supplementary material, which is available to authorized users.

M. Krakau · S. Jacobsen · K. Reise (✉)
Alfred Wegener Institute for Polar and Marine Research,
Wadden Sea Station Sylt, Hafenstr. 43, 25992 List, Germany
e-mail: karsten.reise@awi.de

K. T. Jensen
Marine Ecology, Department of Biological Sciences,
University of Aarhus, Ole Worms Allé 1,
Bld. 1135, 8000 Aarhus, Denmark

cockle *Cerastoderma glaucum*, scattered populations are separated into an eastern group with Ponto-Caspian and Aegean Sea haplotypes and a western group from the Mediterranean to the Baltic Sea (Nikula and Väinölä 2003), and peculiar long-distance relationships may have been caused by migrant birds carrying eggs or larvae (Tarnowska et al. 2010).

The common cockle, *Cerastoderma (Cardium) edule* (L.), on the other hand, occupies a continuous range in shallow water sediments from northern Africa along the eastern Atlantic coasts to northern Norway and Murmansk in the Arctic and is not found in the Mediterranean and Baltic Seas (Brock 1980; Hayward and Ryland 1995). Local populations tend to fluctuate and occasionally collapse due to severe winters, predation, parasites and cockle fishery (Reise 1985; Jensen 1992; Piersma et al. 2001; Strasser 2002; Thieltges 2006; Beukema and Dekker 2006; Kraan et al. 2007). Dispersal in *C. edule* may last up to 4 weeks, primarily by pelagic larvae and secondarily by drifting juveniles (Creek 1960; Armonies 1994; de Montaudouin et al. 2003). This may usually substitute local losses from nearby sites, leaving little traces in the genetic structure in contrast to large-scale glaciations.

The aim of the present study was to examine the mitochondrial DNA (mtDNA) variability of *C. edule* for signatures of its Pleistocene and postglacial population history. Unlike in mussels and oysters, transfers for culturing purposes between coastal regions have not been practiced to our knowledge. Considering the present distribution range of *C. edule* and the suspected high dispersal capacity (Creek 1960; Boyden 1971), we expected high-frequency southern haplotypes with no or few private haplotypes in the northern range, indicating a southward retreat during last glacial maximum and subsequent recolonization of the northern range. On the other hand, survival in cryptic refugia and periglacial zones during glaciations would be indicated by more complex patterns of the genetic structure. Clarifying the mtDNA phylogeography of *C. edule* may improve our understanding on how modern benthic communities came into existence in a coastal region that has experienced pronounced climatic and topographic change since the Pleistocene.

Materials and methods

Samples

Between 2005 and 2007, individuals of *C. edule* were collected from 18 European sampling sites plus one African site (Table 1). Sampling was conducted randomly at the mid-intertidal by hand and rake next to Lagos and Lisbon (Portugal), Dublin (Ireland), Sylt (Germany),

Kristineberg (Sweden) and Bergen (Norway). In addition, specimens from 13 other sites were kindly provided by colleagues following the same sampling protocol. Whole bivalves or parts of soft tissue were preserved immediately in 96–100% ethanol and were stored at $\leq 4^{\circ}\text{C}$ until further processing. The analyses comprised 383 individuals of *C. edule* (Table 1).

DNA extraction and PCR

We extracted total DNA from adductor muscle or foot tissue following the protocol of the QIAGEN[®] DNeasy Blood and Tissue Kit (Qiagen; Hilden, Germany). A DNA fragment of about 700 base pairs (bp) from the 5' part of the mitochondrial cytochrome oxidase I (COI) gene was amplified. Polymerase chain reaction (PCR) amplification was performed on the Biometra[®] Thermal Cycler T1 (Biometra; Göttingen, Germany) as described below.

After receiving the sequencing results of the first PCRs where the primers HCO1490 and HCO2198 from Folmer et al. (1994) were used, a new forward primer was designed: COI5Fc: 5'-CAT AAA GAT ATT GGG ACT TTA TAC-3'. This forward primer was combined with the published reverse primer (HCO2198: 5'-TAA ACT TCA GGG TGA CCA AAA AAT CA-3') for reliable amplifications. PCR was carried out in 50- μL reaction volume with ~ 150 ng template DNA, 0.2 mM of each dNTP, 0.6 μM of each primer, 1.5 mM of MgCl_2 and 2.5 U Taq polymerase (Qiagen; Hilden, Germany) in the buffer supplied by the manufacturer. The first denaturation at 94°C lasted 2 min, followed by 40 cycles with denaturation at 94°C for 60 s, annealing at 50°C for 60 s and extension at 72°C for 60 s. Amplification results were confirmed by 1% agarose gel electrophoresis. Fragments were purified using the QIAquick purification kit (Qiagen; Hilden, Germany). Direct sequencing of PCR products was done in one direction by MWG Biotech (Ebersberg, Germany). Over 25% of sequences representing singleton haplotypes were tested and all verified after repeating DNA extraction, PCR and additional reverse sequencing. Thus, sequencing errors cannot be entirely excluded but should have no major impact on the results.

Sequence and statistical analyses

Quality of sequencing results was manually checked with the free software CHROMAS Lite 2.01 (Technelysium Pty Ltd), and sequences were aligned using "CLUSTAL W" as implemented in BioEdit (Hall 1999). Haplotype and nucleotide diversity analyses were performed with the software package DnaSP 4.10 (Rozas et al. 2003).

Spatial population structuring was tested using SAMOVA 1.0 (Dupanloup et al. 2002) by maximizing the F_{CT}

Table 1 Sampling sites for *C. edule*

| Code | Location | Country | Latitude | Longitude | Sampling date | GG |
|------|-----------------|-----------------|-----------|-----------|----------------|-----------|
| MO | Merja Zerga | Morocco | 34° 50' N | 06° 14' W | August 2005 | Southwest |
| AL | PO Lagos | Portugal | 37° 07' N | 08° 37' W | September 2005 | Southwest |
| LN | PO Lisbon | Portugal | 38° 43' N | 09° 00' W | September 2005 | Southwest |
| AR | FR Arcachon | France | 44° 35' N | 01° 14' W | August 2005 | Southwest |
| RO | FR Roscoff | France | 48° 43' N | 03° 58' W | June 2007 | Southwest |
| TH | Southend | England (UK) | 51° 28' N | 00° 42' E | March 2007 | Southwest |
| DB | Dublin | Ireland | 53° 19' N | 06° 12' W | August 2005 | Southwest |
| SO | St. Andrews | Scotland (UK) | 56° 21' N | 02° 50' W | August 2006 | Scottish |
| SL | North Gluss | Scotland (UK) | 60° 28' N | 01° 21' W | September 2007 | North |
| TX | WS Texel | The Netherlands | 53° 13' N | 04° 94' E | August 2004 | North |
| LA | WS Langeoog | Germany | 53° 45' N | 07° 29' E | July 2007 | North |
| SY | Sylt | Germany | 55° 01' N | 08° 26' E | September 2005 | North |
| LF | Limfjord | Denmark | 56° 56' N | 09° 04' E | July 2007 | North |
| NM | SK Norsminde | Denmark | 56° 02' N | 10° 25' E | July 2007 | Skagerrak |
| KR | SK Kristineberg | Sweden | 58° 14' N | 11° 26' E | July 2006 | Skagerrak |
| FL | SK Arendal | Norway | 58° 26' N | 08° 48' E | August 2005 | Skagerrak |
| BN | Bergen | Norway | 60° 23' N | 05° 20' E | August 2005 | North |
| BO | Bodø | Norway | 67° 17' N | 14° 37' E | August 2005 | North |
| DZ | Murmansk | Russia | 69° 10' N | 36° 05' E | August 2007 | Arctic |

The two-letter code is subsequently used in graphs and text. In the second column, some sites have been grouped for presentation in Fig. 1. The last column indicates the genetic grouping (GG)

value, which is the proportion of total genetic variance due to differences between groups of populations. Significance level was determined based on 1,000 random permutations of the data. A slightly modified structure based on geographic vicinity of sampling sites was additionally tested with the analysis of molecular variance (AMOVA: Excoffier 1995), as implemented in the program Arlequin (Version 3.1; Excoffier et al. 2005).

Pairwise F_{ST} values were calculated in Arlequin using uncorrected pairwise distances. To assess whether gene flow correlates with geographic distance (“isolation-by-distance” model), the matrix of pairwise F_{ST} values to the geographic distance matrix was tested using a Mantel test implemented in Arlequin. The statistical significance of the correlation coefficient was determined by conducting Mantel tests on 5,000 random permutations of the data set. Geographic distances between populations as estimates of dispersal ranges were calculated in kilometers as shortest along-shoreline and across-water distance using ArcGIS 9.2 (ESRI; Kranzberg, Germany).

Since trees are less adequate to visualize intraspecific phylogenies with small genetic distances, a statistical parsimony haplotype network was constructed by the computer program TCS v1.21 (Clement et al. 2000). Neutrality tests and mismatch analyses were performed with the software package DnaSP 4.10. Obtained graphical outputs were used to illustrate demographic model fit. Additionally, the time frame of population expansion (t) was estimated after Rogers (1995) applying the demographic parameters τ (“tau”) given by DnaSP 4.10. The equation

$t = \tau/2 \mu k$ (see Rogers and Harpending 1992) was used with τ as moment estimator of time to expansion, μ as year probability of an individual nucleotide site to mutate and k as length of sequence investigated.

Results

Genetic richness

The aligned data set of COI sequences derived from 383 *Cerastoderma edule* individuals consisted of 582 nucleotide positions with 78 (13%) variable positions. The 79 COI haplotypes identified (Genbank Accession Numbers: FJ403245–FJ403323) displayed between 0.17 and 1.37% nucleotide differences. Haplotype and nucleotide diversity varied between sampling sites, with generally high haplotype diversity but low nucleotide diversity (Table 2).

The highest diversity index values were found at Bodø at the northern Norwegian coast (BO: $h = 0.931$, $\pi = 0.00646$; Table 2). On average, haplotype and nucleotide diversity were observed to be higher in northern edge populations (SL, BO: $h > 0.700$) than in the southern Atlantic sites (MO and AL: $h < 0.430$). The samples from Roscoff (RO) and Southend (TH) exhibited relatively high diversity values compared to more southern sites of the Atlantic region. The lowest genetic diversity value was found in the population from St. Andrews (SO: $h = 0.140$, Table 2), and this Scottish sample was without private haplotypes.

Table 2 Genetic diversity indices for cockle populations

| Code | <i>N</i> | H_t Number of all haplotypes | H_u Number of unique haplotypes | <i>S</i> Number of polymorphic sites | <i>h</i> Haplotype diversity (SD) | π Nucleotide diversity (SD) $\times 10^2$ |
|---------|----------|--------------------------------------|--|---|---|--|
| MO | 21 | 5 | 1 | 4 | 0.424 (0.131) | 0.105 (0.037) |
| AL | 21 | 6 | 2 | 6 | 0.429 (0.134) | 0.113 (0.043) |
| LN | 10 | 3 | 1 | 5 | 0.511 (0.164) | 0.225 (0.089) |
| AR | 13 | 3 | 1 | 4 | 0.410 (0.154) | 0.128 (0.067) |
| RO | 23 | 10 | 6 | 14 | 0.731 (0.099) | 0.264 (0.075) |
| DB | 23 | 8 | 6 | 9 | 0.526 (0.126) | 0.135 (0.046) |
| TH | 21 | 9 | 5 | 11 | 0.729 (0.102) | 0.263 (0.064) |
| SO | 28 | 3 | 0 | 3 | 0.140 (0.087) | 0.048 (0.034) |
| SL | 20 | 6 | 1 | 7 | 0.705 (0.086) | 0.336 (0.070) |
| TX | 20 | 6 | 2 | 6 | 0.447 (0.137) | 0.176 (0.066) |
| LA | 21 | 5 | 3 | 10 | 0.486 (0.124) | 0.191 (0.074) |
| SY | 19 | 12 | 7 | 17 | 0.901 (0.059) | 0.428 (0.074) |
| LF | 19 | 9 | 4 | 8 | 0.772 (0.094) | 0.350 (0.058) |
| NM | 20 | 8 | 2 | 11 | 0.863 (0.049) | 0.553 (0.053) |
| KR | 20 | 7 | 1 | 8 | 0.642 (0.118) | 0.352 (0.093) |
| FL | 20 | 8 | 2 | 13 | 0.868 (0.044) | 0.524 (0.063) |
| BN | 22 | 8 | 2 | 14 | 0.771 (0.080) | 0.402 (0.082) |
| BO | 22 | 11 | 2 | 15 | 0.931 (0.028) | 0.646 (0.072) |
| DZ | 20 | 8 | 7 | 13 | 0.589 (0.130) | 0.223 (0.083) |
| Average | 20.2 | 7.1 | 2.8 | 9.4 | 0.625 | 0.287 |
| ALL | 383 | 79 | 54 | 78 | 0.866 (0.012) | 0.511 (0.017) |

Standard deviation (SD) is given for *h* and π . Code for sites as in Table 1

N number of cockles investigated

Spatial grouping due to regional haplotype dominance

Of the 79 COI variants identified for *C. edule*, only 15 haplotypes occurred at least at two localities (=shared haplotypes, Table 3). Nine haplotypes repeated in several individuals at one sampling site only. Of all haplotypes, 70% were found only once at all. Thus, 55 haplotypes represented real singletons. Two haplotypes, named S and N, appeared at twelve and ten sampling sites, representing 28.5 and 20.4% of individuals in the total sample, respectively (Table 3). The dominance of the S-haplotype and the simultaneous absence of the N-type along the southern Atlantic coast including the Irish and the African sites were apparent when haplotype frequencies were visualized on a geographic map (Fig. 1). Atlantic sites showed high numbers of local haplotypes next to the English Channel (Fig. 1). The haplotype N occurred in all northern populations except at the northernmost site DZ in the Barents Sea. In the North Sea, Norwegian Sea and Arctic area, *C. edule* populations were characterized by several shared northern haplotypes and the absence of the S-type, while a mixture of northern and southern haplotypes was observed for three sites next to the Baltic Sea entrance (Table 3, NM, KR, FL, summarized as SK in Fig. 1).

Spatial AMOVA (SAMOVA) was used to statistically define groups of populations that are geographically

homogeneous and maximally differentiated from each other. Subdivisions were tested for two to seven geographic groups (see Table S1). The analysis confirmed the two groups representing several sampling sites (named “southwestern” and “northern”) that were mainly defined by the presence or absence of one dominant haplotype. In addition, two single sites (SO and DZ) were explicitly separated from all others (Table S1, F_{CT} : 0.514, $p < 0.001$). They contained one dominant haplotype that was shared with no more than two other sites (Table 3). A significant amount of variation occurred between the four statistically distinct groups (Table S2, AMOVA: 48.9%, $p < 0.001$), while within-groups variation was low (6.3%). The southwestern group comprised the African and the southwestern European sites from Portugal to the British Isles dominated by haplotype S (66.7%). Within the northern group, all sites northeast of the English Channel region, the haplotype N was the characteristic and most common haplotype (35.0%), except at SO and DZ (Fig. 1). The number of haplotypes found exclusively either in southwestern or in northern cockle populations was high (Table 3).

Isolation by distance among northern populations

Focusing on the relationship between geographic and genetic distances between all 19 sampling sites along the

Table 3 Frequency of shared haplotypes in *C. edule* populations

| Code | S | I | II | III | IV | V | N | VI | VII | VIII | IX | X | XI | XII | XIII | N _{SH} | N _{LH} |
|------|-----|---|----|-----|----|----|----|----|-----|------|----|---|----|-----|------|-----------------|-----------------|
| MO | 16 | 1 | - | 2 | 1 | - | - | - | - | - | - | - | - | - | - | 20 | 1 |
| AL | 16 | - | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | 19 | 2 |
| LN | 7 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 7 | 3 |
| AR | 10 | - | - | 2 | - | - | - | - | - | - | - | - | - | - | - | 12 | 1 |
| RO | 12 | - | - | 2 | - | - | - | 1 | - | - | - | - | - | - | - | 15 | 8 |
| TH | 11 | - | 2 | - | - | - | - | 1 | - | - | - | - | - | - | - | 14 | 7 |
| DB | 16 | 1 | - | - | - | - | - | - | - | - | - | - | - | - | - | 17 | 6 |
| SO | 1 | - | - | - | - | 26 | - | - | - | - | 1 | - | - | - | - | 28 | 0 |
| SL | 1 | - | - | - | - | - | 10 | 1 | 1 | - | - | - | - | - | 5 | 18 | 2 |
| TX | - | - | - | - | - | - | 15 | 1 | - | - | 1 | - | 1 | - | - | 18 | 2 |
| LA | - | - | - | - | - | - | 15 | 3 | - | - | - | - | - | - | - | 18 | 3 |
| SY | - | - | - | - | - | - | 6 | - | 1 | - | - | 2 | 1 | - | - | 10 | 9 |
| LF | - | - | - | 1 | - | - | 9 | 1 | - | - | 3 | 1 | - | - | - | 15 | 4 |
| NM | 2 | - | - | - | - | - | 4 | 2 | - | 6 | 3 | - | - | 1 | - | 18 | 2 |
| KR | 12 | - | - | - | - | - | 1 | 1 | - | 2 | 2 | 1 | - | - | - | 19 | 1 |
| FL | 5 | - | 2 | - | - | - | 5 | - | - | 2 | 3 | - | - | 1 | - | 18 | 2 |
| BN | - | - | - | - | - | 1 | 10 | 2 | 1 | 2 | - | - | - | - | - | 16 | 6 |
| BO | - | - | - | - | - | 2 | 3 | - | 2 | 1 | - | - | - | - | 4 | 12 | 10 |
| DZ | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 13 | 13 | 7 |
| All | 109 | 2 | 5 | 8 | 2 | 29 | 78 | 13 | 5 | 13 | 13 | 4 | 2 | 2 | 22 | 307 | 76 |

N_{SH} Number of individuals that share haplotypes with cockles from ≥2 sites and N_{LH} Number of individuals with local haplotypes. S, N, I–XIII: haplotype identifier

northeastern Atlantic shoreline, the matrix correlation coefficient indicated that about 20% of the variation in pairwise genetic distances of uncorrected *F*_{ST} values (Table 4) could be attributed to geographic distances between pairs of sites (Mantel test: *R* = 0.448, *p* < 0.001). Repeating the Mantel test with the subset of eleven northern populations, the correlation coefficient *R* increased to 0.620 (*p* < 0.05). This implies that 38% of variation may be explained by geographic differentiation and indicates isolation by distance (IBD) for northern sampling sites, while the analysis of southwestern populations lacked any significant support for IBD (*R* = 0.113, *p* = 0.300).

Phylogenetic and demographic approach


Further insights into the genetic population structure of *C. edule* were provided by the application of a parsimony network analysis of the 79 individual haplotypes (Fig. 2). Differentiation of the geographic groups on nucleotide level was not well pronounced, since the dominant haplotype S in Atlantic populations was separated from haplotype N by only three mutational steps (0.51%). However, the network demonstrated fundamental differences between the associated haplotype complexes. The genealogical relationships between northern haplotypes (gray, Fig. 2) were less well resolved than those of southwestern ones (white, Fig. 2). Within the northern cluster, most of the haplotypes of DZ (black, Fig. 2) form an Arctic mini-cluster around haplotype XIII. In contrast, the southwestern populations displayed shallow divergence among haplotypes. Most haplotypes differentiate only by a single-nucleotide substitution from the dominant haplotype S,

which consequently occupies a central position in a star-like pattern. Noticeable is a close relationship of the predominant Scottish haplotype V to the common southwestern haplotype S (Fig. 2).

The high number of singleton haplotypes (Table 3) and the star-like phylogenetic pattern (Fig. 2) indicate an expansion in the effective population size of the cockle *C. edule* in the recent past, which was supported by neutrality tests (Table 5). Tajima’s *D* and Fu’s *F*_s statistics were negative for the southwestern and the northern group and differed significantly from expectations under a neutral model of evolution assuming constant population size (Table 5). Testing the growth-decline model for the southwestern populations, the unimodal curve in Fig. 3 supported the assumption of a recent rapid population expansion suggested by the strong clustering of the southern haplotypes in a single star-like pattern (Fig. 2). When the populations FL, KR and NM of the Skagerrak area, which were conspicuous because of the occurrence of the S-haplotype, are pooled for Tajima’s *D* and mismatch distribution analysis, a non-significant, negative value (Table 5) and a multimodal topology (Fig. 3) indicate an equilibrium state or secondary contact zone. However, this local grouping is not supported by SAMOVA tests.

The demographic parameters of the southwestern (*τ* = 0.29) and northern group (*τ* = 1.84) are used to estimate the time since regional population expansion (Table 5). Considering the sequence length of 582 bp and a mutation rate *μ* of 2.5% per million years (see Nikula and Väinölä 2003), the start for an expansion for the southwestern group is set to about 10,000 years BP. The northern population expansion is calculated to about 63,000 years BP. Even when the peculiar Skagerrak

Fig. 1 COI haplotypes of *C. edule* along Northeast Atlantic shoreline. Adjacent sites with very similar haplotype occurrence are pooled for clarity of the graph (see Table 1). Haplotype S occurs predominantly at southwestern sites that are separated by a dashed line from northern sites with haplotype N. Non-colored parts of pies refer to local haplotypes. For haplotype identifiers N, S and I–XIII, see Table 3



populations, where both most abundant haplotypes occur, are excluded, the estimate for the northern expansion (N^t : $\tau = 1.22$) is still much longer ago (42,000 years BP) than for the southwestern one.

Discussion

The genetic structure of *Cerastoderma edule* with its northern and southwestern groups of populations along the Northeast Atlantic coast suggests two historically separated populations during low sea-level stands of the last glaciation. High genetic diversity detected at North Sea and Scandinavian sites in contrast to the low diversity and

differentiation in the southwestern cockle populations further supports a scenario of northern glacial refugia.

Our initial expectation of a simple phylogeographic pattern in *C. edule* with diverse southern populations and reduced genetic variety in the North, indicative of northward postglacial recolonization from southern refugia, was not met by the results. It can be inferred that at least one ancestral population has managed to survive somewhere in or near to the North Sea region throughout the last glaciation. Genetic heterogeneity in the northern group of populations suggests a metapopulation structure during the latest glacial period. This is supported by 38% of genetic variation between sampling sites in the northern group explained by isolation by distance, while

Table 4 Pairwise F_{ST} values between sample sites

| | MO | AL | LN | AR | RO | DB | TH | TX | LA | SY | LF | NM | KR | SO | SL | FL | BN | BO | DZ |
|----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|
| MO | | – | – | – | – | – | – | + | + | + | + | + | + | + | + | + | + | + | + |
| AL | –1.2 | | – | – | – | – | – | + | + | + | + | + | + | + | + | + | + | + | + |
| LN | 10.0 | 7.2 | | – | – | – | – | + | + | + | + | + | – | + | + | + | + | + | + |
| AR | –2.4 | –1.5 | 0.3 | | – | – | – | + | + | + | + | + | – | + | + | + | + | + | + |
| RO | 0.8 | 0.2 | 3.3 | –0.9 | | – | – | + | + | + | + | + | – | + | + | + | + | + | + |
| DB | 3.3 | 0.3 | 6.0 | 1.6 | 0.1 | | – | + | + | + | + | + | – | + | + | + | + | + | + |
| TH | 6.3 | 3.4 | 5.3 | 3.4 | 1.1 | 2.0 | | + | + | + | + | + | – | + | + | + | + | + | + |
| TX | 74.4 | 73.6 | 68.1 | 71.9 | 61.0 | 70.9 | 59.1 | | – | – | – | – | + | + | – | – | – | – | + |
| LA | 75.5 | 74.9 | 69.5 | 73.0 | 63.4 | 72.4 | 61.5 | –0.6 | | – | – | + | + | + | – | + | – | + | + |
| SY | 58.1 | 57.6 | 49.9 | 53.3 | 47.8 | 55.5 | 44.5 | 0.5 | 2.5 | | – | – | + | + | – | – | – | – | + |
| LF | 56.8 | 56.1 | 49.3 | 52.3 | 44.9 | 54.1 | 42.0 | 4.3 | 1.1 | 1.7 | | – | + | + | – | – | – | + | + |
| NM | 49.9 | 48.6 | 41.4 | 44.6 | 40.7 | 47.1 | 37.7 | 12.5 | 14.9 | 9.8 | 6.6 | | – | + | – | – | – | – | + |
| KR | 15.8 | 12.9 | 11.7 | 12.0 | 7.1 | 11.0 | 3.9 | 42.6 | 46.6 | 30.4 | 24.6 | 19.8 | | + | + | – | + | + | + |
| SO | 67.9 | 66.2 | 63.6 | 67.6 | 50.9 | 62.9 | 52.4 | 84.2 | 84.0 | 71.2 | 71.9 | 63.9 | 50.2 | | + | + | + | + | + |
| SL | 63.6 | 62.9 | 55.8 | 59.4 | 52.3 | 60.5 | 50.0 | 5.7 | 7.2 | 2.9 | 8.0 | 13.7 | 35.7 | 75.5 | | – | – | – | + |
| FL | 31.8 | 30.3 | 24.9 | 26.9 | 22.6 | 28.6 | 18.6 | 15.3 | 20.3 | 9.2 | 3.7 | 3.8 | 3.8 | 53.7 | 13.9 | | – | + | + |
| BN | 61.5 | 60.9 | 54.1 | 57.2 | 51.6 | 59.0 | 49.3 | 2.2 | 2.0 | 2.2 | 6.3 | 7.6 | 34.7 | 72.4 | 6.3 | 12.6 | | – | + |
| BO | 53.9 | 53.1 | 45.8 | 48.8 | 46.5 | 51.9 | 44.1 | 7.7 | 8.5 | 6.2 | 8.9 | 4.8 | 30.3 | 64.0 | 4.3 | 12.1 | 5.3 | | + |
| DZ | 83.3 | 82.9 | 78.5 | 81.3 | 75.4 | 81.6 | 74.6 | 60.9 | 59.9 | 47.5 | 53.9 | 50.2 | 66.5 | 88.7 | 38.1 | 53.0 | 50.5 | 34.1 | |

Values ($\times 10^{-2}$, below diagonal) with significance after Bonferroni correction indicated by + ($p_B \leq 0.003$, above diagonal). Code for sites as in Table 1




Fig. 2 Minimum spanning network for the 79 COI haplotypes of *C. edule*. Haplotypes shared between two or more sites are labeled (S, N and I–XIII). Each step along a lineage represents a single mutation. *Small open circles* indicate missing haplotypes. *Color codes* of haplotypes correspond to their occurrence in the four spatial groups (as shown)

southwestern populations do not display such geographical structuring.

A relatively high genetic diversity in the North Sea region is also known from two species of seagrass, *Zostera*

marina and *Z. noltii*, which like *C. edule* occur on intertidal and shallow subtidal sedimentary bottoms (Olsen et al. 2004; Coyer et al. 2004). The more recent origin and the genetic homogeneity of the southwestern group reject the possibility of a southward retreat of *C. edule* during glaciations. Instead, there might have been a periglacial refugium west of the English Channel from where a southward extension has commenced well after the last glacial maximum. This would fit with the observation that populations in this area display higher diversity indices than other populations within the southwestern group. Survival during the last glaciation somewhere between Ireland and Brittany has also been assumed for the macroalgae *Fucus serratus*, *Palmaria palmata* and *Ascophyllum nodosum* (Maggs et al. 2008; Olsen et al. 2010). However, these rocky shore populations do not share a common habitat with *C. edule* and their refugial vicinity might be a mere coincidence (Marko et al. 2010).

In conclusion, a simple scenario of southward retreat during glaciation, followed by a northward recolonization after the present interglacial commenced, can be rejected for *C. edule*. Although this is the most common pattern for terrestrial species (Hewitt 2000), marine benthos shows other patterns as well (Marko et al. 2010). Northern cryptic refugia and southern periglacial refugia during glaciation provide a better match with the phylogeography inferred from our analyses. However, as the southwestern group of *C. edule* seems younger than the northern group, northern


Table 5 Neutrality tests and population demographic parameters for spatial groups in the population size model

| Spatial groups | Number of sequences | Neutrality tests | | Demographic parameters | |
|-----------------|---------------------|------------------|------------|------------------------|------------|
| | | Tajima's D | Fu's F_s | τ | θ_i |
| Southwest | 132 | −2.62*** | −4.17 | 0.29 | 0.75 |
| SO | 28 | −1.53 n.s. | −1.06 | 0.00 | 0.58 |
| North | 203 | −1.85* | −39.89 | 1.84 | 0.79 |
| North' | 143 | −1.96* | −36.13 | 1.22 | 1.01 |
| Skagerrak | 60 | −1.97 n.s. | −1.97 | 1.94 | 1.94 |
| DZ | 20 | −2.34** | −3.86 | 0.11 | 1.19 |
| All populations | 383 | −2.21** | −94.38 | 1.99 | 0.97 |

τ , moment estimator of time to expansion; θ_i , mutation parameter for initial populations. To show the special case of the Skagerrak (FL, KR and NM), it was calculated separately from the other northern sites (North')

n.s. not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Fig. 3 Mismatch distribution analysis. The growth-decline model was applied for two geographic groups. The population of St. Andrews (SO) is excluded from the analyses. *Exp* expected values and *Obs* observed values. N number of individuals



survival combined with southern extinction during the last glacial cannot be rejected. The isolated occurrence of the central haplotype of the southwestern group mixed with haplotypes of the northern group at three sites in the Skagerrak/Kattegat region may be interpreted as a remnant of previously separated populations. The deviant population at Murmansk points to the possible existence of a distinct Arctic group of *C. edule*, but more sampling would be required to answer this question.

Besides the correspondence between *C. edule* and the two seagrass species with high genetic diversity in the North Sea, there is also some similarity with two other coastal bivalves. *Mytilus edulis* shows patterns of mtDNA which indicate survival in glacial refugia within its present range of the Atlantic North American coastline (Riginos and Henzler 2008). In Europe, the regions west and south of the English Channel are dominated by *M. galloprovincialis* and east and north of this channel by *M. edulis* (Smietanka et al. 2004). The boundary between the two groups of populations in *C. edule* also coincides with the English Channel. In *Macoma balthica rubra*, a southern and a northern group mix in the North Sea and Kattegat region and multiple glacial refugia may have existed in the northern group of this subspecies (Nikula et al. 2007). A common predator on coastal bivalve species, the crab

Carcinus maenas, is also subdivided into northern and southwestern populations along the Atlantic shores of Europe (Roman and Palumbi 2004; Darling et al. 2008).

In view of the dramatic drop in sea level during glaciations, when most of the North Sea region turned into land (Gaffney et al. 2009), the most parsimonious assumption is that coastal bivalves such as *C. edule* took refuge in ecologically productive polynyas (unfrozen areas surrounded by sea ice) between Scottish and Fennoscandian ice sheets. Another potential glacial refugial zone might have been the Celtic Sea southwest of the British ice sheet (Ménot et al. 2006). *C. edule* performs best in non-stagnant water between 10 and 20°C but survives a wide range of 0–43°C (e.g., Ansell et al. 1981; Wilson 1981; Strasser et al. 2001; Genelt-Yanovskiy et al. 2010). Therefore, temperature alone is unlikely to drive this species to extinction as long as partial populations can survive at greater depth subtidally to avoid extreme temperatures of the intertidal zone (e.g., Eisma 1966; Dörjes et al. 1986; Dekker 1989; Bazairi et al. 2003; Sousa et al. 2006). Gradients from shallow to deep sedimentary bottoms were probably at all times rather common along European coasts to provide habitat for persistent populations of *C. edule*.

Our findings show that the COI gene can be a potentially useful locus for population genetic studies. However, it has

to be mentioned that the utility of COI mtDNA markers for population genetic purposes is limited by potential deviation from neutrality, overestimation of genetic drift due to smaller effective population size and limited amounts of genetic variation (e.g., Ballard and Whitlock 2004; Held and Leese 2007). This may cause bias in selectively neutral population processes like migration and drift (reviewed by Zhang and Hewitt 2003), and analyses based on multilocus genotypes are highly desirable for cockle phylogeography in the future.

Conclusion

With *Cerastoderma edule* as a further example, we support the assumption that North Sea benthic species survived European glaciations in cryptic refugia or periglacial zones. We found no evidence for a species complex, like in the genera *Mytilus* and *Macoma*, or a deep separation of phylogeographic groups like in the sister species *C. glaucum*. However, a heterogeneous northern and a homogeneous southwestern group of populations are distinguished in *C. edule*. Their intergradation needs further research as does the possible existence of an Arctic group.

Acknowledgments For sampling support, we thank colleagues involved in the two European exchange projects, French National Coastal Environment Programme (PNEC) and European MarBEF Network of Excellence “Marine Biodiversity and Ecosystem Functioning”. MK appreciates the cordial invitation to the laboratories of H. Cabral, S. Mortensen and J. Wilson. MK likes to thank following people for support in laboratory and field work: H. Halliger, G. Czichy, J. Kochmann, V. Liebich, J. F. Marques, B. Regenfuß and A. Schlüssel. The study was partly supported by the MarBEF Network funded by the European Community’s Sixth Framework Programme (contract no. GOCE-CT-2003-505446). We thank Mathias Wegner and the anonymous reviewers whose comments substantially improved the manuscript.

References

- Ansell AD, Barnett PRO, Bodoy A, Massé H (1981) Upper temperature tolerances of some European molluscs. *Mar Biol* 65:177–183
- Armonies W (1994) Turnover of postlarval bivalves in sediments of tidal flats in Königshafen (German Wadden sea). *Helgol Mar Res* 48:291–297
- Ballard JW, Whitlock MC (2004) The incomplete natural history of mitochondria. *Mol Ecol* 13:729–744
- Bazairi H, Bayed A, Glemarec M, Hily C (2003) Spatial organisation of macrozoobenthic communities in response to environmental factors in a coastal lagoon of the NW African coast (Merja Zerga, Morocco). *Oceanol Acta* 26:457–471
- Beukema JJ, Dekker R (2006) Annual cockle *Cerastoderma edule* production in the Wadden Sea usually fails to sustain both wintering birds and commercial fishery. *Mar Ecol Prog Ser* 309:189–204
- Boyden CR (1971) A comparative study of the reproductive cycles of the cockles *Cerastoderma edule* and *C. glaucum*. *J Mar Biol Assoc UK* 51:605–622
- Brannock PM, Hilbish TJ (2010) Hybridization results in high levels of sterility and restricted introgression between invasive and endemic marine blue mussels. *Mar Ecol Prog Ser* 406:161–171
- Brock V (1980) The geographical distribution of *Cerastoderma (Cardium) edule* (L) and *C. lamarcki* (Reeve) in the Baltic and adjacent seas related to salinity and salinity fluctuations. *Ophelia* 19:207–214
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Mol Ecol* 9:1657–1659
- Coyer JA, Peters AF, Stam WT, Olsen JL (2004) Population genetics of *Zostera noltii* (dwarf eelgrass) throughout its biogeographic range. *Mar Ecol Prog Ser* 281:51–62
- Coyer JA, Hoarau G, Van Schaik J, Luijckx P, Olsen JL (2011) Trans-Pacific and trans-Arctic pathways of the intertidal macroalga *Fucus distichus* L. reveal multiple glacial refugia and colonizations from the North Pacific to the North Atlantic. *J Biogeogr* 38(4):756–771
- Creek GA (1960) The development of the lamellibranch *Cardium edule*. *Proc Zool Soc Lond* 135:243–260
- Darling JA, Bagley MJ, Roman J, Telpot CK, Geller JB (2008) Genetic patterns across multiple introductions of the globally invasive crab genus *Carcinus*. *Mol Ecol* 17:4992–5007
- de Montaudouin X, Bachelet G, Sauriau PG (2003) Secondary settlement of cockles *Cerastoderma edule* as a function of current velocity and substratum: a flume study with benthic juveniles. *Hydrobiologia* 503:103–116
- Dekker R (1989) The macrozoobenthos of the subtidal western Dutch Wadden Sea. I. Biomass and species richness. *Neth J Sea Res* 23:57–68
- Dörjes J, Michaelis H, Rhode B (1986) Long-term studies of macrozoobenthos in intertidal and shallow subtidal habitats near the island of Norderney (East Frisian coast, Germany). *Hydrobiologia* 142:217–232
- Dupanloup I, Schneider S, Excoffier L (2002) A simulated annealing approach to define the genetic structure of populations. *Mol Ecol* 11:2571–2581
- Eisma D (1966) The distribution of benthic marine molluscs off the main Dutch coast. *Neth J Sea Res* 3:107–163
- Excoffier L (1995) AMOVA Analysis of Molecular Variance, Version 1.55. University of Geneva, Switzerland
- Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47–50
- Folmer O, Black M, Hoeh R, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3:294–299
- Gaffney V, Fitch S, Smith D (2009) Europe’s lost world: the rediscovery of Doggerland. CBA research report 160, York
- Genelt-Yanovskiy EA, Poloskin A, Granovitch A, Nazarova S, Strelkov P (2010) Population structure and growth rates at biogeographic extremes: a case study of the common cockle. *Cerastoderma edule* (L.) in the Barents Sea. *Mar Poll Bull* 61:247–253. doi:10.1016/j.marpolbul.2010.02.021
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98
- Hayward PJ, Ryland JS (1995) Handbook of the marine fauna of north-west Europe. Oxford University Press, Oxford
- Held C, Leese F (2007) The utility of fast evolving molecular markers for studying speciation in the Antarctic benthos. *Pol Biol* 30:513–521

- Hewitt GM (2000) The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913
- Hoarau G, Coyer JA, Veldsink JH, Stam WT, Olsen JL (2007) Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Mol Ecol* 16:3606–3616
- Hoffmann AA, Sgrò CM (2011) Climate change and evolutionary adaptation. *Nature* 470:479–4485
- Jensen KT (1992) Macrozoobenthos on an intertidal mudflat in the Danish Wadden Sea—comparisons of survey in the 1930s, 1940s and 1980s. *Helgol Meeresunters* 46:363–376
- Kraan C, Piersma T, Dekinga A, Koolhaas A, van der Meer J (2007) Dredging for edible cockles (*Cerastoderma edule*) on intertidal flats: short-term consequences of fisher patch-choice decisions for target and non-target benthic fauna. *ICES J Mar Sci* 64:1735–1742
- Lejeune C, Bock DG, Theriault TW, MacIsaac HJ, Cristescu ME (2011) Comparative phylogeography of two ascidians reveals contrasting invasion histories in North America. *Biol Invasions* 13:635–650
- Luttikhuisen PC, Drent J, Baker AJ (2003) Disjunct distribution of highly diverged mitochondrial lineage clade and population subdivision in a marine bivalve with pelagic larvae. *Mol Ecol* 12:2215–2229
- Luttikhuisen PC, Campos J, van Bleiswijk J, Peijnenburg KTCA, van der Veer HW (2008) Phylogeography of the common shrimp, *Crangon crangon* (L) across its distribution range. *Mol Phylogenet Evol* 46:1015–1030
- Maggs CA, Castilho R, Foltz D et al (2008) Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology* 89:S108–S122
- Marko PB, Hoffman JM, Emme SA, McGovern TM, Keever CC, Cox LN (2010) The ‘Expansion-Contraction’ model of Pleistocene biogeography: rocky shores suffer a sea change? *Mol Ecol* 19:146–169
- Martin PR, McKay JK (2004) Latitudinal variation in genetic divergence of populations and the potential for future speciation. *Evolution* 58:938–945
- Ménot G, Bard E, Rostek F, Weijers JWH, Hopmans EC, Schouten S, Damsté JSS (2006) Early reactivation of European rivers during the last deglaciation. *Science* 313:1623–1625
- Nikula R, Väinölä R (2003) Phylogeography of *Cerastoderma glaucum* (Bivalvia: Cardiidae) across Europe: a major break in the Eastern Mediterranean. *Mar Biol* 143:339–350
- Nikula R, Strelkov P, Väinölä R (2007) Diversity and trans-arctic invasion history of mitochondrial lineages in the North Atlantic *Macoma balthica* complex (Bivalvia: Tellinidae). *Evolution* 61:928–941
- Olsen JL, Stam WT, Coyer JA et al (2004) North Atlantic phylogeography and large-scale population differentiation of the seagrass *Zostera marina* L. *Mol Ecol* 13:1923–1941
- Olsen JL, Zechman FW, Hoarau G, Coyer JA, Stam WT, Valero M, Åberg P (2010) The phylogeographic architecture of the fucoid seaweed *Ascophyllum nodosum*: an intertidal ‘marine tree’ and survivor of more than one glacial-interglacial cycle. *J Biogeogr* 37:842–856
- Piersma T, Koolhaas A, Dekinga A, Beukema JJ, Dekker R, Essink K (2001) Long-term indirect effects of mechanical cockle-dredging on intertidal bivalve stocks in the Wadden Sea. *J Appl Ecol* 38:976–990. doi:10.1046/j.1365-2664.2001.00652.x
- Reise K (1985) Tidal flat ecology. *Ecol Stud* 54, Springer, Berlin
- Reitzel AM, Darling JA, Sullivan JC, Finnerty JR (2008) Global population genetic structure of the starlet anemone *Nematostella vectensis*: multiple introductions and implications for conservation policy. *Biol Invasions* 10:1197–1213
- Reusch TBH, Bolte S, Sparwel M, Moss AG, Javidpour J (2010) Microsatellites reveal origin and genetic diversity of Eurasian invasions by one of the world’s most notorious marine invader, *Mnemiopsis leidyi* (Ctenophora). *Mol Ecol* 19:2690–2699. doi:10.1111/j.1365-294X.2010.04701.x
- Riginos C, Henzler CM (2008) Patterns of mtDNA diversity in North Atlantic populations of the mussel *Mytilus edulis*. *Mar Biol* 155:399–412
- Rogers AR (1995) Genetic evidence for a pleistocene population explosion. *Evolution* 49:608–615
- Rogers AR, Harpending H (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Mol Biol Evol* 9:552–569
- Roman J, Palumbi SR (2004) A global invader at home: population structure of the green crab, *Carcinus maenas*, in Europe. *Mol Ecol* 13:2891–2898
- Rozas J, Sánchez-Delbarrio JC, Messeguer X, Rozas R (2003) DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19:2496–2497
- Smietanka B, Zbawicka M, Wolowicz M, Wenne R (2004) Mitochondrial DNA lineages in the European populations of mussels (*Mytilus* spp.). *Mar Biol* 146:79–92
- Sousa R, Dias S, Antunes JC (2006) Spatial subtidal macrobenthic distribution in relation to abiotic conditions in the Lima estuary, NW of Portugal. *Hydrobiologia* 559:135–148
- Strasser M (2002) Reduced epibenthic predation on intertidal bivalves after a severe winter in the European Wadden Sea. *Mar Ecol Prog Ser* 241:113–123
- Strasser M, Reinwald T, Reise K (2001) Differential effects of the severe winter of 1995/96 on the intertidal bivalves *Mytilus edulis*, *Cerastoderma edule* and *Mya arenaria* in Northern Wadden Sea. *Helgol Mar Res* 55:190–197
- Tarnowska K, Chenuil A, Nikula R, Féral JP, Wołowicz M (2010) Complex genetic population structure of bivalve (*Cerastoderma glaucum*) living in highly fragmented lagoon habitat. *Mar Ecol Prog Ser* 406:173–184
- Thieltges DW (2006) Parasite induced summer mortality in the cockle *Cerastoderma edule* by the trematode *Gymnophallus choledochus*. *Hydrobiologia* 559:455–461
- Väinölä R, Strelkov P (2011) *Mytilus trossulus* in Northern Europe. *Mar Biol* 158:817–833
- Wilson J (1981) Temperature tolerance of circatidal bivalves in relation to their distribution. *J therm Biol* 6:279–286
- Zhang D, Hewitt GM (2003) Nuclear DNA analyses in genetic studies of populations: practice, problems and prospects. *Mol Ecol* 12:563–584