

The Common Sense Model of Self-Regulation: Meta-Analysis and Test of a Process Model

Martin S. Hagger

Curtin University and University of Jyväskylä

Severine Koch and Nikos L. D. Chatzisarantis

Curtin University

Sheina Orbell

University of Essex

©2017, American Psychological Association. This paper is not the copy of record and may not exactly replicate the final, authoritative version of the article. Please do not copy or cite without authors permission. The final article will be available, upon publication, via its DOI: 10.1037/bul0000118

Full citation: Hagger, M. S., Koch, S., Chatzisarantis, N. L. D., & Orbell, S. (2017). The common-sense model of self-regulation: Meta-analysis and test of a process model.

Psychological Bulletin. <http://dx.doi.org/10.1037/bul0000118>

Author Note

Martin S. Hagger, Health Psychology and Behavioral Medicine Research Group and Laboratory of Self-Regulation, School of Psychology and Speech Pathology, Perth and Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä;

Severine Koch, and Nikos L. D. Chatzisarantis, Health Psychology and Behavioral Medicine Research Group and Laboratory of Self-Regulation, School of Psychology and Speech Pathology, Perth; Sheina Orbell, Department of Psychology, University of Essex, Colchester.

Martin S. Hagger's contribution was supported by a Finland Distinguished Professor (FiDiPro) award from TEKES, the Finnish funding agency for innovation.

We thank Associate Editor Blair T. Johnson for his helpful comments on earlier drafts of this manuscript. We are grateful to Angela Carroll for her help with the literature search. We also thank the following authors who provided additional data for the current meta-analysis: Abdolaziz Aflakseir, Seher Arat, Vera Araújo-Soares, Yael Benyamini, Becky Bih-O Lee, Felicity Bishop, Tore Bonsaksen, Noel T. Brewer, Elizabeth Broadbent, Trudie Chalder, Joseph Chilcot, Jimmy Chong, Arden Corter, Erin Costanzo, Francis Creed, Nicola Dalbeth, Martin Dempster, Aleid de Rooij, Alex Federman, Maarten Fischer, Nadine Foster, Lana Fu, Milena Gandy, Rinie Geenen, David Gillanders, Prosenjit Giri, Lesley Glover, Christopher Graham, Konstadina Griva, AnnMarie Groarke, Brooks Gump, Jorine Hartman, John Harvey, Ankie Heerema-Poelman, Susan Hill, Alex Holmes, Joanna Hudson, Thomas Hyphantis, Egil Jonsbu, Rebecca Knibb, Deirdre Lane, Margaret Lau-Walker, Valerie Lawson, Yvonne Leung, Carrie Llewellyn, Sean Lynch, Julie MacInnes, Pamela McCabe, Elaine McColl, Aušra Mickevičienė, Gerry Molloy, Kevin Morgan, Rona Moss-Morris, Stanton Newman, Arie Nouwen, Ronan O'Carroll, Agnieszka Olchowska-Kotala, Catherine O'Leary, Patrícia Pinto, Heather Powell, Yolanda Quiles, Gwyneth Rees, Orna Reges, Jillian Riley, Tina Rochelle, Michael Rose, Reza Sadjadi, Gregory Sawicki, Margreet Scharloo, Stefanie Schroeder, Louise Sharpe, Timothy Skinner, Debbie Snell, Lesley Stafford, Susan Stott, Ilse Stuive, Elaine Thomas, Barbara Tomenson, Amy Turriff, Maureen Twiddy, Mike van der Have, Tina van der Velde, Firdous Var, Manja Vollmann, Ken Watkins, Robert Whittaker, Wendy Woith, and Urška Žugelj.

Correspondence concerning this article should be addressed to Martin S. Hagger, Health Psychology and Behavioral Medicine Research Group and Laboratory of Self-Regulation, School of Psychology and Speech Pathology, Faculty of Health Sciences, Curtin

University, GPO Box U1987, Perth, WA 6845, Australia, email:

martin.hagger@curtin.edu.au

Abstract

According to the common-sense model of self-regulation, individuals form lay representations of illnesses that guide coping procedures to manage illness threat. We meta-analyzed studies adopting the model to (a) examine the intercorrelations among illness representation dimensions, coping strategies, and illness outcomes; (b) test the sufficiency of a process model in which relations between illness representations and outcomes were mediated by coping strategies; and (c) test effects of moderators on model relations. Studies adopting the common-sense model in chronic illness ($k = 254$) were subjected to random-effects meta-analysis. The pattern of zero-order corrected correlations among illness representation dimensions (identity, consequences, timeline, perceived control, illness coherence, emotional representations), coping strategies (avoidance, cognitive reappraisal, emotion venting, problem-focused generic, problem-focused specific, seeking social support), and illness outcomes (disease state, distress, well-being, physical, role, and social functioning) was consistent with previous analyses. Meta-analytic path analyses supported a process model that included direct effects of illness representations on outcomes and indirect effects mediated by coping. Emotional representations and perceived control were consistently related to illness-related and functional outcomes via, respectively, lower and greater employment of coping strategies to deal with symptoms or manage treatment. Representations signaling threat (consequences, identity) had specific positive and negative indirect effects on outcomes through problem- and emotion-focused coping strategies. There was little evidence of moderation of model effects by study design, illness type and context, and study quality. A revised process model is proposed to guide future research which includes effects of moderators, individual differences, and beliefs about coping and treatment.

Keywords: Illness perceptions; illness cognition; coping procedures; parallel-processing model; chronic illness

Public Significance Statement

This review indicates that relations between patients' illness beliefs and important illness-related outcomes (reducing disease progression, improving functioning, promoting well-being, allaying distress) across studies are accounted for by sets of coping strategies (avoidance, cognitive reappraisal, emotion venting, problem-focused coping, seeking social support). Behavioral interventions aimed at changing illness outcomes should not only target change in the beliefs linked to adaptive outcomes, but also the coping strategies related to the beliefs.

The Common-Sense Model of Illness Representations: Meta-Analysis and Test of a Process Model

The identification and interpretation of a deviation from the normally-functioning physical and somatic self is fundamental to initiating self-regulatory processes to restore or protect a state of health, or even protect life itself. When an individual experiences a physical symptom, how does his or her particular conception of illness relate to efforts to manage symptoms, and what processes lead the individual to engage in long-term self-regulatory action that will be effective in changing the course of the illness rather than symptom management alone? Prominent among social psychological approaches to understanding the regulatory mechanisms involved in managing perceived illness-related threats (e.g., Carver, Scheier, & Weintraub, 1989; Rodin & Salovey, 1989) is the ‘common-sense’ model of self-regulation. The model outlines the processes underlying individuals’ lay management of health threats (Leventhal, Diefenbach, & Leventhal, 1992; Leventhal, Leventhal, & Breland, 2011; Leventhal, Meyer, & Nerenz, 1980; Leventhal, Phillips, & Burns, 2016).

According to the model, an understanding of adaptation to health threats originates in an examination of threat from the individual’s lay perspective. Specifically, it is necessary to understand the individual’s cognitive and emotional representations of the threat that initiate subsequent coping procedures directed at threat and emotion management. These two parallel processes together represent a self-regulatory system with significant implications for illness outcomes such as disease, functional state, and psychological distress that are independent of the pathological markers of illness. For example, severe emotional reactions to symptoms arising from fear that the illness may be life threatening may lead to management by avoidance coping. Similarly, a cognitive representation that misattributes the cause of symptoms may lead to ineffective attempts at self-directed treatment (Leventhal et al., 1992).

Whereas research on the common-sense model of self-regulation has indicated that cognitive and emotional representations of illness are related to coping procedures adopted to manage illness threat and distress, and to illness-related outcomes such as recovery or symptom management (Brown, Westbrook, & Challagalla, 2005; Whitmarsh, Koutantji, & Sodell, 2003), few studies have explicitly tested the processes involved. Based on the hypotheses derived from Leventhal et al.'s model, a *process model* has been proposed in which "relationships [between illness representations, coping, and outcomes] represent a mediational model, in which coping mediates the effect of illness representations on health outcomes" (Hagger & Orbell, 2003, p. 146). Although there have been previous tests of the process model in participants with particular chronic illnesses (e.g., Brewer, Chapman, Brownlee, & Leventhal, 2002; Gould, Brown, & Bramwell, 2010; Gray & Rutter, 2007; Rutter & Rutter, 2002), there has been no systematic test of a full process model adopting multiple representation, coping, and outcome variables based on a cumulative synthesis of data across research on the common-sense model in chronic illness.

We aim to conduct a meta-analytic synthesis of studies testing relations from the common-sense model in chronic illness and use the correlations from the analysis to test the proposed process model. Not only do we expect that quantifying the overall effects of illness representations on outcomes mediated by coping to be of interest, we also expect to identify specific indirect effects that will indicate the extent to which illness representations are positively associated with outcomes related to recovery and effective management, or undermine recovery such as poorer functioning and well-being, and elevated distress. The mediational model may be informative in resolving inconsistent patterns of effects observed in previous studies and may potentially inform practice by identifying viable targets for intervention. In addition, the synthesis enables us to assess the influence of moderator variables that may explain variability in relations among model constructs across studies.

The Common-Sense Model of Self-Regulation: Origins and Conceptualization

The common-sense model was developed to understand how patients' lay perceptions of illness threats guide coping strategies to deal with those threats (Leventhal et al., 1980). The model is illustrated in Figure 1. The substantive *content* of an illness representation, derived from multidimensional scaling studies (Bishop & Converse, 1986; Linz, Penrod, & Leventhal, 1982) and open-ended interviews (Baumann, Cameron, Zimmerman, & Leventhal, 1989; Lau & Hartman, 1983), and formalized in psychometric instruments such as the illness perception questionnaire (IPQ; Weinman, Petrie, Moss-Morris, & Horne, 1996), comprises five main dimensions of the cognitive representation of illness: identity, timeline, cause, consequences, and perceived control. The *identity* of the illness is characterized by its symptoms and label; *timeline* reflects the rate of onset, duration, and fluctuation in the illness; *cause* is the perceived causal antecedents of the illness (e.g., genetics, infection, diet, aging, or other exacerbating factors); *consequences* reflects beliefs in the extent to which the illness will seriously impact life events (e.g., work, family, personal relationships); and *perceived control* is defined as whether the illness is expected to be, or has previously been, responsive to self-initiated or medically-prescribed treatment. A sixth dimension, *illness coherence*, representing patients' comprehension of their illness, was included in the revised illness perceptions questionnaire (IPQ-R; Moss-Morris et al., 2002). Illnesses may be more or less coherently represented, depending on the extent to which the patient's current illness representation matches stored prototypes of the illness and illnesses with similar content. The representation is also informed and modified by communications from health providers and intra-personal factors such as personality and individual differences (Leventhal et al., 1992).

The model also identifies emotional representations that reflect individuals' affective responses to illness and may independently affect selection of coping procedures in parallel to cognitive representations (Figure 1). According to Martin, Leventhal, and Leventhal

(2003), “affective and cognitive responses to symptoms may or may not be compatible. For example, a person might understand that a symptom (e.g., bloody stools) could have serious consequences if left untreated. This belief might motivate care seeking in the interests of early detection. However, for some people, the fear of potentially receiving an ominous diagnosis (e.g., of cancer) actually might inhibit self-referral behavior” (p. 210). Emotional representations were also included as an additional component in the IPQ-R (Moss-Morris et al., 2002). The measure has enabled tests of relations between individuals’ reflections on their emotional responses to the illness and outcomes such quality of life, and emotional coping responses such as emotion venting, alongside cognitive illness representations, (e.g., Evans & Norman, 2009; Knibb & Horton, 2008; Rozema, Völlink, & Lechner, 2009).

Illness representations are conceptualized as memory structures and may be activated by a novel somatic change or salient information and lead to a search for a matching illness prototype stored in memory. This is the first stage in the process of recognizing that one is ill, and initiates self-evaluation (e.g., recognizing other symptoms, observing whether they go away), self-management of illness (e.g., attempting to control the symptoms), and consulting a professional (Leventhal et al., 2011). These stimuli are depicted to the left of Figure 1. The progression to professional treatment may sometimes bypass both self-evaluation and self-management, when, for example, an acute event provokes immediate hospitalization leading to diagnosis of illness, or when screening procedures identify diseases such as breast cancer without any prior symptomatic information indicating to the individual that they are ill.

Also important are the moderating effects of socio-cultural and personal variables, labelled the ‘self-system’ by Leventhal et al. (1992), that may influence interpretation of, and response to illness stimuli, depicted at the top of Figure 1. For example, an individual may perceive a headache and muscle ache as symptoms of a benign illness such as a common cold. However, if such symptoms occurred after visiting a region where malaria is prevalent,

1 they might activate representations of serious infectious disease associated with elevated
2 threat. Illness schema can be activated from sources which may be directly symptomatic and
3 concrete, but also arise independent of symptoms based on more abstract internal states that
4 interact with contextual knowledge (Orbell, Henderson, & Hagger, 2015).

5 Following the identification of illness and the activation of an illness schema
6 associated with an illness label (e.g., multiple sclerosis, diabetes, rheumatoid arthritis, cancer,
7 asthma), a dynamic self-regulatory process is initiated, comprising attempts to manage the
8 health threat and concomitant emotional reactions (Leventhal, Brissette, & Leventhal, 2003;
9 Leventhal et al., 1980). These self-regulatory efforts are labeled coping procedures in the
10 common-sense model and follow from cognitive and emotional representations. For example,
11 an individual with bowel cancer may view the illness as having painful symptoms, a chronic
12 timeline, and controllable through dietary change, and therefore view a change of diet as an
13 effective means to cope with the disease progression. An individual who believes the same
14 disease is uncontrollable and finds it a source of emotional distress may adopt a denial coping
15 response. The parallel arrowed pathways in Figure 1 depict links between the representation
16 dimensions and coping procedures. Importantly, the adopted coping procedures will make
17 ‘common sense’ to the individual in that they follow from their lay representations, regardless
18 of the type and effectiveness of the coping procedure adopted (Leventhal et al., 1992).

19 While the common-sense model does not specify particular coping procedures, a
20 number of theoretically-derived coping dimensions have been employed that broadly
21 correspond to problem- or threat-focused and emotion-focused coping categories (Carver et
22 al., 1989; Folkman & Lazarus, 1988). Consistent with the proposal that illness cognitions are
23 schematic in nature (Henderson, Hagger, & Orbell, 2007; Leventhal & Cleary, 1980;
24 Leventhal et al., 2011), studies have demonstrated that automatic activation of an illness
25 representation is associated with activation of relevant coping procedures (Henderson, Orbell,

1 & Hagger, 2009). Information regarding 'typical' coping responses may, therefore, be
2 represented in memory alongside information about the illness and activated and modified
3 accordingly when information regarding the illness (e.g., felt symptoms) is made salient.

4 The common-sense model posits that cognitive and emotional representations will
5 affect outcomes that indicate the progress of the illness (e.g., disease state), the individual's
6 functional capacity, and perceived health status. Representations guide the individual's
7 selection of an appropriate coping strategy, a 'common sense' solution to the illness threat. In
8 a process of appraisal depicted in Figure 1 an individual will experience if the strategy has led
9 to improvement, no change, or deterioration in symptoms and adjust his or her representation.
10 These 'feedback loops' characterize the model as a dynamic framework: It describes the
11 process by which individuals represent their illness, select coping procedures, and modify
12 their representations through an appraisal of the coping and subsequent experience of illness.

13 **A Critical Appraisal of Research on the Common-Sense Model of Self-Regulation**

14 Hagger and Orbell (2003) tested the hypothesized relations among common-sense
15 model constructs in a meta-analysis of studies applying the model in chronic illness. The
16 studies measured illness perceptions and specific coping and outcome measures that were
17 systematically classified into categories based on theory and research (Carver, Scheier, &
18 Pozo, 1992; Carver et al., 1989; Stewart et al., 1992; Veit & Ware, 1983) and corroborated by
19 expert consensus. The analysis revealed a consistent pattern of relations such that
20 representation dimensions that signal an elevated level of threat, namely, serious
21 consequences, illness identity, and a chronic timeline were positively associated with
22 emotion-focused coping procedures including expressing emotions and denial. Perceptions of
23 the illness as controllable were positively associated with problem-focused coping strategies,
24 and with cognitive reappraisal. Relations between the illness representation dimensions and
25 illness outcomes revealed positive associations of the threat dimensions with psychological

1 distress, and negative associations with outcomes such as physical and social functioning, and
2 psychological well-being. Perceived control was positively associated with functioning,
3 vitality and psychological well-being, and negatively related to disease state¹. The pattern of
4 effects identified by Hagger and Orbell (2003) have been subsequently corroborated in meta-
5 analyses of subsets of the research literature on the common-sense model (Brandes & Mullan,
6 2014; Broadbent et al., 2015; Dempster, Howell, & McCorry, 2015; French, Cooper, &
7 Weinman, 2006; Hudson, Bundy, Coventry, & Dickens, 2014; Mc Sharry et al., 2011).

8 The typical pattern of relations between representations, coping, and outcomes
9 derived from previous research neglects to account for the moderating influence of contextual
10 factors. For example, transactional models of stress and coping indicate that problem-focused
11 coping strategies may not be adaptive if adopted to manage illnesses that are not controllable
12 (Lazarus & Folkman, 1984). Emotion-focused strategies may be more effective in these
13 contexts as they help individuals manage negative emotional reactions caused by the
14 perceived illness threat. This was recognized by Leventhal et al.'s (1992) 'goodness-of-fit'
15 hypothesis, which indicates that the effectiveness of coping strategies on recovery and illness
16 progression is context dependent. However, current meta-analytic data do not account for the
17 effects of these contextual factors (e.g., Dempster et al., 2015; Hagger & Orbell, 2003).

18 From a theoretical perspective, meta-analyses of studies of the common-sense model
19 suggest that representations that signal illness threat do not guide individuals to adopt
20 problem-focused coping strategies as hypothesized by Leventhal et al. (1980) in their original
21 conceptualization of the model. The meta-analytic research indicates that these beliefs tend to

¹Hagger et al.'s (2003) analysis did not include the emotional representation and illness coherence dimensions as few studies at the time had adopted the revised IPQ and there were insufficient studies reporting effect sizes for these studies. Studies and quantitative syntheses have since demonstrated statistically significant and positive relations between emotional representations and outcomes related to increased illness progression and decreased functioning (Broadbent, Petrie, Main, & Weinman, 2006; Mc Sharry, Moss-Morris, & Kendrick, 2011), while illness coherence has been related to better functioning and well-being (Paddison, Alpass, & Stephens, 2008; Karademas, 2012).

1 be related to coping procedures that are emotion-focused, or, at least, not problem focused,
2 such as avoidance or denial. From a lay beliefs perspective, such coping responses make
3 common sense; the immediate management of emotional upheaval is an important first step
4 in active illness management because emotional distress is likely to interfere with
5 individuals' capacity to formulate and engage in problem-focused coping strategies
6 (Leventhal et al., 1980). This interpretation notwithstanding, threat perceptions are likely to
7 guide problem-focused coping when the individual has a clear, concrete "reality-bound
8 picture of a knowable threat and not a mysterious, remote, uncertain, and infinitely
9 threatening view of the illness process" (Leventhal et al., 1980, p. 20). There is empirical
10 support for this hypothesis, indicating that threat perceptions (e.g., high perceived
11 consequences) are related to treatment-seeking coping behaviors such as medication
12 adherence (Brewer et al., 2002), adherence to medical advice (Karademas, Karamvakalis, &
13 Zarogiannos, 2009), and self-care behaviors (MacInnes, 2013). Contextual factors such as
14 illness familiarity and coherence may determine whether threat perceptions lead to problem-
15 or emotion-focused coping. Current meta-analytic data, therefore, capture overall tendencies
16 to prioritize emotion management in response to the perceived threat posed by the illness, but
17 may not reflect instances when the threat informs problem-focused coping efforts.

18 From an empirical perspective, it is important to note that no prior meta-analytic
19 synthesis of relations among the constructs of the common-sense model has controlled for the
20 effects of the other representation, coping, and outcome dimensions since they rely solely on
21 zero-order relations. Given the consistently strong intercorrelations among representation
22 dimensions, particularly among the threat dimensions (Hagger & Orbell, 2003), it is likely
23 that there will be a considerable degree of shared variance among these constructs that will
24 lead to substantively attenuated effects of illness representation dimensions on coping and
25 outcome constructs in multivariate analyses relative to the zero-order correlations (e.g., Evans

1 & Norman, 2009; Griva, Jayasena, Davenport, Harrison, & Newman, 2009). These findings
2 highlight the importance of conducting multivariate tests when synthesizing findings using
3 meta-analysis. This will permit identification of the unique effects of each representation
4 dimension on coping and outcome constructs controlling for the other dimensions (cf.
5 Hagger, Chan, Protogerou, & Chatzisarantis, 2016; Hagger & Chatzisarantis, 2016).

6 **A Process Model of ‘Common-Sense’ Illness Representations**

7 Central to the common-sense model is the assertion that individuals’ cognitive and
8 emotional representations of an illness threat will motivate a coping response to mitigate the
9 threat and related distress. Conceptually, therefore, specific coping procedures are proposed
10 to mediate the representation-outcome relations in the model (Figure 2). The inclusion of
11 multiple representation dimensions along with numerous types of coping responses and
12 illness outcomes means that specific effects of each illness representation on illness outcomes
13 through coping strategies can be proposed and tested. From an analytic perspective, therefore,
14 illness representations “may exert effects [on illness outcomes] by eliciting or suppressing
15 both adaptive and maladaptive coping responses” (Gould et al., 2010, p. 635).

16 The accuracy of the process model can be evaluated on two levels. Taking a
17 generalized perspective, evaluating the extent to which coping mediates relations between
18 representation dimensions and illness outcomes will provide a test of the *sufficiency* of the
19 model. This would be a substantial advance on previous meta-analytic tests of the model
20 which have relied exclusively on zero-order correlations among constructs, a sub-optimal
21 approach to testing unique effects in social cognitive models (Hagger, Chan, et al., 2016).
22 Empirically, examination of the total indirect effects of all representation dimensions on
23 illness outcomes with all coping constructs as multiple mediators affords an assessment as to
24 the extent to which coping fully accounts for the impact of representations on outcomes. An
25 alternative hypothesis would be that a total indirect effect is present, but direct effects also

1 exist, such that representation dimensions have unique effects on the outcome that are not
2 explained by coping. However, if the total indirect effects are found to be trivial or zero in the
3 presence of the direct effects, it would raise serious questions as to the adequacy of the
4 process model and could be grounds for its rejection.

5 The process model also permits a more fine-grained evaluation of the specific coping
6 mediated pathways by which individual representation constructs relate to outcomes. At this
7 level, testing for specific indirect effects may provide an indication of pathways not evident
8 when observing the total indirect effects, and may be ‘missed’ in an evaluation of the model.
9 For example, illness representations signaling threat (e.g., beliefs that an illness is highly
10 symptomatic, will have serious consequences, and has a chronic timeline) may compel an
11 individual to take action to attenuate the threat, consistent with Leventhal et al.’s (1980)
12 original hypothesis. Such actions may lead to adaptive improvements in illness outcomes.
13 However, the same representations may also lead an individual to engage in coping strategies
14 to manage negative feelings evoked by the perceived threat, such as emotion venting or
15 avoidance. Such strategies may not lead to improvements in disease state and functioning,
16 although they may make the person feel better. Illness identity, consequences, and timeline
17 may have two sets of specific indirect effects on illness outcomes through different coping
18 strategies that are not identified by observation of the total indirect effects alone.

19 The specific mediated pathways by which illness perceptions relate to outcomes
20 through coping procedures have seldom been explored in research on the common-sense
21 model. A few previous tests provide some preliminary evidence for the mediation hypothesis
22 in the common-sense model. For example, studies have found that perceived consequences
23 has a positive indirect effect on outcomes like anxiety and depression through emotion
24 venting and acceptance coping strategies (Benyamini, Gozlan, & Kokia, 2004; Evans &
25 Norman, 2009; Rutter & Rutter, 2002), while other studies have found a negative indirect

effect on adaptive outcomes like improved mental health and better social functioning through avoidance (Heijmans, 1999). These mediation analyses corroborate the pattern of effects in correlational analyses. However, some mediation analyses revealed indirect effects that did not conform to this pattern, but are consistent with Leventhal et al.'s proposal that threat representations motivate problem-focused coping procedures and better illness outcomes. For example, Brewer et al. (2002) found indirect effects of perceived consequences on cholesterol levels of hypercholesterolemic patients mediated by medication adherence, a problem-focused coping strategy. Such findings illustrate the potential for mediation analyses to reveal process-related effects not evident when observing total indirect effects or zero-order relations among illness representations, coping strategies, and illness outcomes.

To date, tests of the process model are limited in that they were selective in the illness representation and coping dimensions included in their mediation analysis and were restricted to single illnesses and samples. A stronger evaluation of the process model needs to include multiple illness representation, coping strategy, and outcome measures simultaneously across conditions and samples, and adopt appropriate multivariate confirmatory analyses (e.g., path analysis, structural equation modeling). Such an analysis would permit a test of the nomological validity of the process model including an explicit test of multiple mediation effects, and their magnitude and sign, positive or negative (Bagozzi, 1981; Hagger, Chan, et al., 2016). Our current analysis will also address the limitations of previous tests of the process model by providing a comprehensive test of the mediational pathways that includes all illness representation dimensions and multiple coping and outcome variables based on a cumulative synthesis of findings from empirical research on the common-sense model in chronic illness.

The Present Review

The current research is timely given the piqued interest in the common-sense model (Law, Tolgyesi, & Howard, 2012; Leventhal et al., 2011; Leventhal et al., 2016; Petrie & Weinman, 2012; Wearden & Peters, 2008) and its processes (Benyamini et al., 2004; Brewer et al., 2002; Gould et al., 2010), and the proliferation of research testing its component relations. Based on the foregoing review and critique we aim to test a process model derived from Leventhal et al.'s common-sense model based on a meta-analytic synthesis of findings from studies measuring illness representations, coping strategies, and illness outcomes in patient groups with chronic illnesses or conditions². A critical mass of studies now exists making a model test based on a cumulative synthesis of the available literature feasible. Our analysis will extend previous research by employing a full matrix of correlations, providing a test of model sufficiency that takes account of shared variance among model constructs, identifying specific indirect effects, and evaluating contextual moderators of model relations.

Zero-order intercorrelations. The starting point of our analysis is to conduct a meta-analytic synthesis of studies on chronic conditions or illnesses. Across studies, we will examine the zero-order, univariate patterns of relations among the three sets of variables that constitute the process model: illness representation dimensions, coping strategies, and illness outcomes.

Testing the sufficiency of the process model. We will subsequently subject the corrected zero-order correlation matrix of the representation, coping, and outcome variables to meta-analytic path analysis (Hagger, Chan, et al., 2016) to test the proposed mediation effects in the process model. The hypothesized model is depicted in Appendix A (supplemental materials). For the sake of parsimony, six separate path analyses of the process

²Although there is no universal definition of a chronic illness or condition, there is general consensus on the characteristics that constitute chronic illness (Goodman, Posner, Huang, Perekh, & Koh, 2013). For the purposes of the current research, we define a chronic illness as a departure from a state of physical or mental well-being lasting more than three months (US Department of Health and Human Services, 2011) that likely requires ongoing medical care and limits activities of daily living (US Department of Health and Human Services, 2013).

model are proposed, one with each illness outcome (disease state, distress, physical functioning, psychological well-being, role functioning, social functioning) as the dependent variable. In each analysis, the illness representation dimensions will be set as predictors of the coping constructs, and each coping construct will be set as a predictor of the outcome. The coping variables will serve as multiple mediators of each illness representation dimension on the outcome. To test the sufficiency of the model, we propose a generalized hypothesis in which the illness representation dimensions have statistically significant non-zero total indirect effects on each illness outcome mediated by the coping constructs with no direct effects of the representations on outcomes. An alternative model is also envisaged in which representations are related to outcomes via two pathways: directly, and indirectly through coping. In this case, the direct relations between representations and outcomes would be attributable to mediators other than the identified coping constructs.

Testing model pathways. The analysis will also permit examination of specific indirect coping pathways which might be missed by examining the total indirect effects alone. In particular, we sought to address inconsistent evidence that threat representations have positive indirect effects on maladaptive outcomes mediated by emotion-focused coping (Benyamini et al., 2004; Gould et al., 2010; Rutter & Rutter, 2002), but have also been found to predict adaptive outcomes mediated by the adoption of problem-focused coping strategies (e.g., Brewer et al., 2002). Representation dimensions may have multiple effects on adaptive (e.g., reduced distress, better functioning) and maladaptive (e.g., greater distress, increased illness state) outcomes through specific coping strategies. These pathways may not have been observed previously because researchers have not tested comprehensive models in which the indirect effects of all representation dimensions on illness outcomes are specified through multiple coping constructs.

1 **Testing for moderators.** Previous meta-analyses have observed considerable
2 heterogeneity in relations among model constructs (Broadbent et al., 2015; French et al.,
3 2006; Hagger & Orbell, 2003). As a goal of meta-analysis is to identify and explain
4 heterogeneity unattributed to methodological artifacts such as sampling and measurement
5 error, we aim to examine the effects of five candidate moderators with methodological and
6 conceptual bases: study design, illness type, medically-explained versus medically-
7 unexplained illness symptoms, illness stage, and study methodological quality. We aim to test
8 whether the relations among model constructs differed across moderator groups and plan to
9 conduct sensitivity analyses testing whether the pattern of effects in the process model varied
10 substantially across levels of the moderators.

11 Study design is an important methodological moderator given that effects in social
12 cognitive models tend to be attenuated when variables identified as antecedents and
13 consequents are measured at different time points. For example, attenuation may occur due to
14 patients modifying their illness representations over the time course of the illness due to
15 experience with treatment, or changes in symptoms. A time lag between self-report measures
16 also reduces the impact of common method variance, which has the tendency to inflate
17 correlations (Podsakoff, MacKenzie, Lee, & Podsakoff, 2003). We therefore expect relations
18 between model constructs to be weaker in studies with prospective and longitudinal designs.
19 We also plan to make the distinction between correlational and experimental studies as
20 experimental designs also tend to exhibit weaker relations among constructs relative to
21 correlational designs. However, few studies have experimentally manipulated illness
22 representations so it might be unreasonable to expect this distinction to reveal moderation of
23 model effects. We expect no moderation of the intercorrelations by study design as illness
24 constructs are typically measured concurrently, regardless of design.

Contextual factors have been identified as key moderators of the effects of illness representations on coping strategies and illness outcomes (Leventhal et al., 1992; Leventhal et al., 1980). Illnesses vary in their symptomatology, impact on the patients' life, chronicity, and responsiveness to treatment. We aim to examine the effects of specific illness types on relations between illness representation dimensions and coping and outcomes. Cancer is an illness likely to evoke strong negative reactions given its general cultural association with serious consequences (Erblich, Montgomery, Valdimarsdottir, Cloyre, & Bovbjerg, 2003; Henderson et al., 2007). We expect studies of cancer patients to report stronger effects of emotional representations and threat-related cognitive representations on emotion-focused coping. We also predict stronger effects of cognitive representations on problem-focused coping given the seriousness of the condition and relatively well-stipulated management regimens for cancers. We also expect to observe similar patterns of effects in studies of patients with cardiovascular disease and diabetes given the high prevalence and well-known debilitating outcomes of these illnesses. These illnesses are also likely to evoke strong emotional responses and elevated threat beliefs with similar effects on coping and outcomes in the model (Aalto, Heijmans, Weinman, & Aro, 2005; Mc Sharry et al., 2011).

Another candidate moderator is whether the illness or condition has a medically-supported etiology with clear prognosis and treatment or whether the condition is medically unexplained (Deary, Chalder, & Sharpe, 2007). Although the existence of a medical explanation does not necessarily imply congruence between expert and lay representations of the illness, the presence of a medical explanation provides a greater likelihood that the illness will be understood by patients and may be manifested in greater perceived coherence and control with respect to the illness; perceptions that may lead to the adoption of more adaptive coping strategies. In contrast, conditions with medically-unexplained symptoms (e.g., chronic fatigue syndrome, forms of chronic pain, fibromyalgia) are likely to be associated with less

1 coherence, greater perceptions of chronicity and emotional representations, increased
2 likelihood of attributing symptoms to the condition, and heightened concerns of the impact of
3 the illness on life. Such perceptions may lead to emotion-focused or avoidant coping
4 responses (e.g., catastrophizing thoughts, denial, hypervigilance), which likely lead to
5 outcomes that are maladaptive from a medical model perspective (Moss-Morris, Spence, &
6 Hou, 2011). We therefore expect illness representation dimensions signaling threat to be
7 related to emotion-focused coping strategies such as emotion venting among patients with
8 medically-unexplained symptoms. Similarly, we expect stronger links between these
9 dimensions and maladaptive outcomes such as increased distress and lower functioning in
10 patients with medically-unexplained symptoms.

11 Stage of illness is a further viable moderator. Consistent with model hypotheses, the
12 impact of representations on coping strategy selection is expected to vary in accordance with
13 whether the illness is in an early-stage diagnosis or whether the diagnosis is distal and
14 experience with the condition is more extensive. As threat perceptions and adverse emotional
15 responses are likely to be elevated on initial diagnosis, we expect stronger effects of threat
16 perceptions (consequences, timeline, and identity) and emotional representations on emotion-
17 focused coping strategies, and weaker effects on problem-focused coping strategies, in
18 studies of recently-diagnosed patients relative to those on patients for whom some time had
19 elapsed since diagnosis. As patients receive more information and have greater experience
20 with symptoms and treatment over time, they may adopt more problem-focused coping
21 strategies in response to threat-related and control representations. Testing for the moderation
22 effects of illness stage is therefore likely to capture a dynamic process in the model in which
23 coping responses change with illness and treatment experience (Leventhal et al., 2016).

24 Consistent with research highlighting the potential of inadequate methodological
25 quality to bias synthesized effects when conducting meta-analyses (Johnson, Low, &

MacDonald, 2014), we aim to conduct an assessment of the quality of the included studies. Quality assessments will be used in sensitivity analyses to evaluate whether conclusions based on the analysis are affected if studies with a high risk of bias were included.

Method

Literature Search and Inclusion Criteria

An electronic literature search was conducted using Web of Science (1980 to December, 2013) as the primary database with identical searches conducted in the following databases: Scopus, OVID PsycARTICLES, and PubMed. The search included the keywords *illness perception*, *illness cognition*, *illness representation*, *illness belief*, and *common-sense model*. In addition, a cited reference search was conducted to identify articles citing key theoretical overviews of the common-sense model (e.g., Leventhal et al., 1997; Leventhal et al., 2003; Leventhal et al., 1992; Leventhal, Leventhal, & Contrada, 1998; Leventhal et al., 1980). Furthermore, efforts were made to obtain any ‘fugitive literature’ (Rosenthal, 1994) in the form of missing correlations or unpublished datasets by contacting authors directly (see Appendix B, supplemental materials, for a flowchart of the search, selection, and inclusion process).

We developed a priori criteria for the inclusion of studies in the meta-analysis. Studies were considered eligible for inclusion if they reported quantitative relations between one or more illness representation dimensions from Leventhal et al.’s (1980) common-sense model and at least one coping or outcome variable directly related to a chronic illness or condition. Qualitative research (e.g., Bishop & Converse, 1986; Morris & Ogden, 2012), narrative reviews (e.g., DiMatteo, Haskard-Zolnieriek, & Martin, 2012), study protocols (e.g., Gray et al., 2012), theoretical articles (e.g., Rees, Fry, & Cull, 2001), non-English language articles (e.g., Balck, Preuss, Hendrichske, & Lippmann, 2012), studies on acute illnesses or conditions (e.g., Lau, Bernard, & Hartman, 1989; McCarthy, Lyons, Weinman, Talbot, &

Purnell, 2003), studies measuring illness representations but no coping or outcome variables or scale-development articles (e.g., Chilcot, Norton, Wellsted, & Farrington, 2012; Hagger & Orbell, 2005), studies measuring illness representations in healthy populations (e.g., Figueiras & Alves, 2007), studies using a composite illness representation score (e.g., Chilcot & Moss-Morris, 2013), studies on hypothetical reactions to illnesses (e.g., Karademas, Bati, Karkania, Georgiou, & Sofokleous, 2013), studies on screening behavior or other preventive measures in healthy samples (e.g., Anagnostopoulos et al., 2012), studies on conditions that are part of a normal variation in health state rather than a chronic illness or condition such as menopause (e.g., Hunter & O'Dea, 1999), studies involving representations of mental disorders (e.g., Lobban, Barrowclough, & Jones, 2005), and studies adopting proxy measures of illness representations in others (e.g., Janse, Sinnema, Uiterwaal, Kimpen, & Gemke, 2005) did not meet the inclusion criteria and were excluded.

Included Studies and Characteristics

The literature research identified 333 articles that met inclusion criteria on initial screening (Appendix C, supplemental materials). A substantial proportion of the eligible articles ($k = 172$) did not report the necessary data for the analysis. The corresponding authors of these articles were contacted to obtain additional data. We were unable to source the unreported data for 39 articles, and several articles either reported data from multiple samples within a single study or comprised overlapping samples (multiple studies using the same data). Details of articles providing multiple and overlapping samples are provided in Tables 1 and 2 in Appendix D (supplemental materials). The final sample comprised 254 studies with 270 independent samples and a total sample size of 52,599. Summary characteristics of the included studies including sample sizes, study design, demographic details, illness types encompassed by the studies, moderator coding, and additional variables are provided in Table 3 in Appendix E (supplemental materials). Studies focused on a diverse range of illnesses and

conditions with forms of cardiovascular disease ($k = 36$), diabetes ($k = 32$), cancers ($k = 22$), arthritis ($k = 16$), forms of chronic pain ($k = 12$), chronic obstructive pulmonary disease ($k = 9$), end-stage renal disease ($k = 9$), chronic fatigue ($k = 7$), multiple sclerosis ($k = 7$), irritable bowel syndrome ($k = 6$), psoriasis ($k = 6$), and hypertension ($k = 4$) the most frequently cited. Studies were largely focused on older samples (median of the average sample age reported in studies = 52 years) with most studies having approximately equal ratios of females and males. Single sex samples were usually due to studies targeting illnesses that generally affect, or occur exclusively in, females (e.g., cervical cancer) or males (e.g., prostate cancer). Other than measures of illness representations, coping strategies, and illness outcomes, a number of studies also included concurrent measures of belief-based, social cognitive, and individual difference constructs. Prominent among these constructs were beliefs about treatment, particularly beliefs about medicines measured by the beliefs about medicines questionnaire (BMQ; Horne, Weinman, & Hankins, 1999), and personality constructs, including constructs from the ‘big five’ personality traits (e.g., Skinner, Hampson, & Fife-Schaw, 2002), optimism and pessimism (e.g., Heijmans & De Ridder, 1998b), and positive and negative perfectionism (e.g., Moss-Morris et al., 2011). Studies tended to include these additional variables as predictors of target outcomes alongside constructs from the common-sense model and findings are summarized in Table 3 (Appendix E, supplemental materials).

Classification of Illness Representations

Studies were relatively consistent in the instruments employed to measure illness representations, relying largely on previously-validated and standardized generic questionnaires based on Leventhal et al.’s (1980) illness representation dimensions of cause,

control, consequences, identity, and timeline³. Specifically, 95% of the included studies used either the illness perception questionnaire (IPQ; Weinman et al., 1996), the revised illness perception questionnaire (IPQ-R; Moss-Morris et al., 2002), or the brief illness perceptions questionnaire (BIPQ; Broadbent et al., 2006) to measure illness representations. The latter two questionnaires included two additional illness representation dimensions, illness coherence and emotional representations, and effect sizes for these dimensions were also included in the present analysis. The illness representation dimensions of the IPQ, IPQ-R, and BIPQ have direct equivalence given that they are all derived from the same root scale, and the dimension structure of the three measures have exhibited adequate construct, predictive, and discriminant validity and satisfactory internal consistency (e.g., Broadbent et al., 2006; Hagger & Orbell, 2005; Moss-Morris et al., 2002). The remaining 5% of the studies employed either the diabetes illness representation questionnaire (DIRQ; Skinner et al., 2003), personal models of diabetes interview and questionnaire (Hampson, Glasgow, & Toobert, 1990), the implicit models of illness questionnaire (IMIQ; Turk, Rudy, & Salovey, 1986), the diabetes care profile (DCP; Fitzgerald et al., 1996), or non-validated measures of cognitive representations based on Leventhal et al.'s (1980) common-sense model (e.g., Eiser, Riazi, Eiser, Hammersley, & Tooke, 2001; Kemp, Morley, & Anderson, 1999).

In order to ensure consistency in the measures of illness representation dimensions across studies, we conducted a content analysis of all illness representation dimensions

³We noted considerable heterogeneity in measures of the cause dimension, which created difficulties in classifying its components into meaningful categories as it has been largely defined in terms of illness-specific causes or has been aggregated into dimensions that vary widely across studies. In line with Hagger and Orbell (2003), we excluded the cause dimension from the meta-analysis. It should also be noted that the revised illness perception questionnaire (IPQ-R; Moss-Morris, Weinman, Petrie, Horne, Cameron, & Buick, 2002) included an additional timeline dimension, timeline: cyclical. Due to the relatively small number of studies employing this subscale, we restricted our analysis to the timeline: acute/chronic dimension. Therefore, in the remainder of this article, timeline is synonymous with timeline: acute/chronic dimension. In addition, some instruments (e.g., IPQ-R, Moss-Morris et al., 2002; BIPQ, Broadbent, Petrie, Main, & Weinman, 2006) distinguish between personal control and treatment control while others employ a single control dimension (e.g., IPQ, Weinman, Petrie, Moss-Morris, & Horne, 1996). We adopted the classification with a single control dimension and therefore combined the personal and treatment control dimensions in studies measuring both components.

employed across studies, identical to the procedure described by Hagger and Orbell (2003). We coded the measures on the basis of content and label into the six illness representation dimensions: control, consequences, identity, timeline, emotional representations, and illness coherence. For instance, the seriousness, control, and symptoms dimensions from Hampson et al.'s (1990) personal models inventory were coded as equivalent to the consequences, control, and identity dimensions from the common-sense model. Similarly, the DIRQ (2003) perceived threat to health and perceived impact dimensions were coded as equivalent to the consequences dimension, and the effectiveness to prevent future complications and effectiveness to control diabetes dimensions were coded as equivalent to the control dimension. In order to provide external validity for our coding, the coding scheme was confirmed by three raters who independently conducted the classification procedure based on formal definitions of the six illness representation dimensions. Inter-rater reliability indicated perfect agreement among the raters (Cohen's $\kappa = 1.00$).

Many of the included studies used modified versions of the IPQ-R to assess the specific illness or condition of interest. In addition, the IPQ-R was translated, validated and standardized for various populations. Studies followed standard IPQ-R scoring procedures such that higher scores on the consequences, identity, timeline, and emotional representation dimensions reflected more serious consequences, greater symptom frequency, chronic duration, and heightened negative emotional reactions. In contrast, high scores on the control and coherence dimensions represented greater perceived control over, and a clearer understanding of, the illness.

Classification of Coping Strategies

The present sample of studies used a large number of instruments to measure coping strategies, many of which were based on cognitive-motivational-relational models of stress and coping (e.g., Carver et al., 1989; Lazarus & Folkman, 1984). A key task in the present

meta-analytic synthesis was to classify measures used to tap coping strategies in the current sample and classify them into conceptually-distinct categories based on theory. Many of the studies in the sample adopted generic, previously-validated questionnaires to assess coping strategies, such as the COPE inventory (Carver et al., 1989), the ways of coping checklist (WCCL; Vitaliano, Russo, Carr, Maiuro, & Becker, 1985), ways of coping questionnaire (WOCQ; Folkman & Lazarus, 1988), and the Utrecht coping list (UCL; Schreurs, Van Der Willige, Tellegen, & Brosschot, 1988).

Following Hagger and Orbell's (2003) classification procedure, we identified six distinct coping categories, five of which reflected the general scope and content of the pool of generic coping scales, namely avoidance, cognitive reappraisal, emotion venting, problem-focused coping (generic strategies), and seeking social support (see Appendix F, supplemental materials, for classification of constructs from generic coping measures into coping categories). A sixth coping category, labelled problem-focused specific coping strategies, encompassed active attempts to address the illness by means of specific illness-related coping behaviors, such as medication and dietary adherence, illness-related self-care behaviors, and attendance at illness-related medical appointments (see Appendix G, supplemental materials, for the classification of non-generic coping measures into coping strategies). We employed three independent raters with expertise in health psychology and theories of stress and coping to code measures of coping into the six a priori categories based on formal definitions of the six coping categories and the content of the source items of the scales used to tap the constructs (see Appendixes F and G, supplemental materials). Inter-rater reliability coefficients indicated good agreement between the raters for each coping dimension category (Fleiss-corrected $\kappa = .91$). Differences in classifications were discussed among the raters with respect to the category definition and item content with resolution based on consensus between all three raters.

Classification of Illness Outcomes

Given the range of measures used to tap illness-related outcomes in the current sample of studies, it was also important to categorize illness outcome measures into distinct categories. As with the coping strategies, we adopted Hagger and Orbell's (2003) procedure to code the outcome measures into relevant categories. We identified six distinct illness outcome categories a priori: disease state, physical functioning, psychological distress, psychological well-being, role functioning, and social functioning. Although Hagger and Orbell included a seventh category, vitality, this category was merged with the psychological well-being category in the present analysis due to considerable overlap in the content of measures used to tap constructs across these categories. While many of the studies adopted self-report psychometric instruments to tap the outcome variables such as the Medical Outcomes Short Form Health Survey (SF-36; Ware & Sherbourne, 1992) or the Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983), it is important to note that objective measures of outcomes were also coded, particularly for the disease state category, which reflects the degree of progression of the illness or condition (e.g., glycosylated hemoglobin (HbA1c) in diabetic patients, CD4 count in patients with HIV). Three raters coded the item content of the illness outcome measures from the source instruments with the operational definitions of the construct categories used in the current analysis with strong agreement across raters (Fleiss-corrected $\kappa = .92$). Differences in classifications were discussed among the raters with respect to the category definition and item content with resolution based on consensus between all three raters. Results of the classification of each outcome measure or questionnaire subscale into outcome categories and the individual studies that utilized each measure are provided in Appendix H (supplemental materials).

Meta-Analytic Strategy

1 The effect size metric of interest in the present meta-analysis was the zero-order
2 correlation coefficient. We conducted separate meta-analyses of intercorrelations among the
3 illness representation and coping strategy dimensions, and of correlations between the
4 representation, coping and outcome dimensions, resulting in 138 effect sizes⁴. Hunter and
5 Schmidt's (2004) formulas were adopted to correct the effect sizes for statistical artefacts.
6 The Hunter and Schmidt approach is equivalent to a random effects model for meta-analysis
7 and is considered optimal as it provides estimates that are generalizable to the population
8 rather than to the sample of included studies alone (Field, 2001; Hagger, 2006; Hunter &
9 Schmidt, 2000). We corrected the effect sizes for both sampling and measurement error and
10 utilized the zero-order correlation coefficient as the effect size metric. The meta-analyses
11 were conducted using the MetaQuick (Stauffer, 1996) and Comprehensive Meta-Analysis
12 Version 2 (Borenstein, 2011) statistical software. In the event that a particular category of our
13 key variables was represented by more than one construct in a specific study (e.g., the
14 positive reintegration and acceptance scales from the COPE inventory were both classified
15 into the cognitive reappraisal coping category), the average of the correlation coefficients was
16 taken to provide a single test of the expressed relationship for use in the meta-analysis
17 consistent with Hunter and Schmidt's (2004) methods. Studies reporting effect size data in
18 other metrics (e.g., standardized mean difference) were converted to correlation coefficients.
19 We corrected for measurement error using Cronbach alpha reliability coefficients of the
20 constructs used in each effect size calculation. Where reliability statistics were unavailable,
21 measurement error was inferred from available attenuation statistics using Stauffer's (1996)
22 formula.

23 The analyses provided key summary statistics of the effects among illness
24 representations, coping strategies and illness outcomes: the uncorrected averaged correlation

⁴Raw data files used to conduct the meta-analysis for each effect are available online at <https://osf.io/g48nt/>

coefficient for the effect (r), the averaged correlation corrected for sampling error only (r_+), the fully-corrected averaged correlation coefficient corrected for both sampling and measurement error (r_{++}), 95% confidence intervals (CI_{95}) of the fully-corrected averaged correlation which provided a formal test of statistical significance for each effect, and the variance in the fully-corrected averaged correlation accounted for by the statistical artefacts of sampling and measurement error. If the vast majority of the variance in the averaged effect size can be accounted for by statistical artifacts – Hunter and Schmidt (2004) recommend a 75% cut-off criterion – then the effect should be considered homogenous (i.e., free from bias other than sampling and measurement error). If the proportion of the variance falls below this criterion, then it is likely that there is substantial variance in the effect size across the studies that cannot be attributed to methodological artifacts and indicates the possibility that additional variance in the effect exists across the studies that can be attributed to extraneous or ‘moderator’ variables. We also report Cochran’s (1952) Q statistic, which provides a formal test of the hypothesis that variation in effect sizes across studies is greater than that expected on the basis of sampling error alone. Given that the number of studies (k) varies across meta-analyses, the Q -statistic cannot be compared across analyses. An alternative is offered by the I^2 statistic and its confidence interval (Higgins & Thompson, 2002). The I^2 is the proportion of the observed variance in the averaged effect size relative to the variance attributable to sampling error alone expressed as a percentage (Borenstein, Higgins, Hedges, & Rothstein, 2017). I^2 values of 25%, 50%, and 75% are said to represent low, medium, and high levels of heterogeneity in the averaged effect size, respectively (Higgins, Thompson, Deeks, & Altman, 2003). An I^2 value that exceeds 25% with large confidence intervals and a lower bound that includes the value of zero is indicative of substantial heterogeneity in the effect (Huedo-Medina, Sánchez-Meca, Marín-Martínez, & Botella, 2006) and should prompt a search for extraneous moderators of the effect. Reporting the between-study or

heterogeneity variance (τ^2) for each averaged effect size is also recommended. Small values for τ^2 are indicative of low heterogeneity in the effect size across studies. Finally, we evaluated the presence of small-study bias in the sample of effect sizes by computing statistics based on plots of the effect size from each study against study precision (usually the reciprocal of the study sample size). Asymmetry in the predicted ‘funnel’ shape of the plots is considered evidence of small study bias, that is, the tendency for studies included in the analysis to exhibit large effects relative to their size. This is often interpreted as a potential indicator of publication bias and the tendency for journals to favor statistically significant findings in underpowered studies. Specifically, we used Egger, Davey Smith, Schneider, and Minder’s (1997) regression test, which indicates the extent to which effect sizes are predicted by study precision and Duval and Tweedie’s (2000) ‘trim and fill’ technique, which identifies studies that deviate from the expected shape of the ‘funnel’ plot and adjusts the plot for missing studies to make it more symmetric.

Moderator Coding and Analyses

Assuming substantive, non-trivial variability in the effect sizes of the relations in the illness representation, coping, and outcome constructs unattributed to the statistical artifacts we corrected for in the model, we aimed to examine the effect of moderators of the model effects. A series of candidate moderator variables was identified and coded in the current analysis: study design, illness type, medically-explained versus medically-unexplained symptoms, illness stage, and study methodological quality. Moderator coding is presented in Table 3 (Appendix E, supplemental materials). We evaluated the effects of the moderator on correlations between the illness representation, coping, and outcome variables across studies by conducting separate meta-analyses for each level of the moderator⁵. Differences in the

⁵Raw data files used to conduct the moderator analyses for each effect including moderator coding are available online at <https://osf.io/g48nt/>

corrected correlations across moderator groups were evaluated by inspection of the 95% confidence intervals about each correlation with Welch's *t*-test providing a formal test of difference.

Study design. Study design was coded as cross-sectional, longitudinal, or intervention. Studies measuring all study variables at a single point in time were coded as cross-sectional. This included studies adopting longitudinal or intervention designs that only reported correlational data at a single time point or independent of the intervention. Studies adopting designs in which either coping or outcomes were measured at a distal point in time relative to the illness representation dimensions were coded as longitudinal. Studies including an intervention or experimental component targeting change in illness representations, and change in a subsequent behavior or illness-related outcome, were coded as intervention. However, these studies numbered very few (3.2%) and, in many cases, data were unavailable (1.5%), which precluded moderator comparisons between groups of studies adopting intervention or experimental designs and groups of studies with cross-sectional and longitudinal designs.

Illness type. Coding of illness type was based on the description of the illnesses reported in the study. Studies were classified as having cancer as the target illness if study participants had received a cancer diagnosis or were undergoing cancer screening. Studies were classified as having diabetes as the target illness if the patient group comprised either type I or type II diabetics. Studies were classified as targeting cardiovascular disease if the patient group was described as having experience with, or diagnosis for, a form of cardiovascular disease including acute coronary syndrome, angina, atrial fibrillation, cardiac chest pain, coronary artery/heart disease, heart failure, hypercholesterolemia, myocardial infarction, and stroke. In the case of each illness type moderator, the contrasting moderator group comprised the remaining studies in which the target illness was absent.

Medically-explained versus medically-unexplained symptoms. Illnesses and conditions were classified as those with medically-explained symptoms and medically-unexplained symptoms by two independent researchers with research experience in behavioral medicine. Studies describing the target illness or condition of the study as having unknown or uncertain etiology, or if the symptoms of the illness or condition are known to have unknown or uncertain causes, were classified as having medically-unexplained symptoms with the remaining illnesses classified as having medically-explained symptoms. Inter-rater reliability for the coding of studies for medically-explained and medically-unexplained illnesses resulted in good inter-rater reliability (intraclass correlation = .83; CI₉₅ = .78, .86). Inconsistencies were resolved through discussion and consultation with medical definitions of the reported conditions alongside the description of the illness and patient group reported in the studies.

Illness stage. Illness stage was coded as time since diagnosis. Studies on patients who received their diagnosis at four weeks or earlier prior to study data collection were classified as recently diagnosed while studies on patient groups who had received their diagnosis more than four weeks prior to data collection were classified as having a non-recent diagnosis. A third category was identified for samples comprising both patients who had received a recent diagnosis and patients who had not received their diagnosis recently.

Study methodological quality. Study quality was assessed using a methodological quality checklist developed for the current study. Given that the current sample comprised exclusively of studies adopting survey methods and largely adopted correlational designs, our checklist was based on those developed for these types of study. Checklist content was adapted from the National Institutes of Health Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies (National Institutes of Health, 2014) supplemented with items from other checklists developed for the assessment of studies with survey,

questionnaire, and cross-sectional designs (Husebø, Dyrstad, Søreide, & Bru, 2012; Jack, McLean, Moffett, & Gardiner, 2010; Oluka, Nie, & Sun, 2014). The checklist assessed 16 criteria, 13 of which were relevant to all studies while a further three (sample representativeness, loss to follow-up, follow-up measures) were only relevant to longitudinal or intervention studies. A full list of checklist criteria, item descriptions, and required quality standards is presented in Table 4 (Appendix I, supplemental materials). Studies meeting quality standards were assigned a score of 1 for each criterion and those not meeting the quality standard or provided insufficient information to evaluate the criterion were assigned a score of zero. Cross-sectional studies could achieve a maximum score of 13 and longitudinal studies a maximum score of 16. We weighted the scores for purposes of comparison by dividing raw scores by the number of relevant items and multiplying by 10.

Studies were scored on the checklist by a researcher with experience in the use of methodological quality checklists⁶. One study could not be scored because the relevant data were not available and was omitted from subsequent analyses. A subset of the sample of studies ($N = 20$) was independently scored by two other researchers with high agreement between raters across items (mean agreement = 94.17%) and inter-rater reliability (mean Fleiss' $\kappa = .82$). Inconsistencies were resolved through discussion and attributed to minor interpretation errors of the quality standard for two criteria. The criteria were subsequently revised on the basis of the discussion and applied to the coding of the entire sample. We coded two methodological quality categorical variables for use in our analysis. After a tertile division of the studies by checklist score, studies with scores in the upper third were classified as high quality and those in the lower third classified as low quality⁷. This strategy

⁶A spreadsheet of methodological quality scores for each study included in the analysis is available online at <https://osf.io/g48nt/>

⁷There were multiple studies with scores on the methodological quality checklist at the tertile cut points, so we adopted an inclusive strategy so that all studies with scores falling on the cut-points were classified in the upper and lower thirds. This led to slightly unequal numbers of studies in the upper (high quality studies; $k = 91$) and lower (low quality studies; $k = 74$) thirds.

allows comparisons of studies with extreme scores but also reduces the sample size. We therefore coded an additional moderator variable based on a cut-off checklist score (Husebø et al., 2012; Jack et al., 2010). Studies achieving weighted scores of 6 or greater on the checklist were classified as high quality and with scores below 6 classified as low quality.

Testing the Process Model

We used the meta-analytically derived corrected correlations to test hypotheses of the proposed process model in which the effects of illness representations on illness outcomes are mediated by coping strategies. Analyses were conducted using the MPlus v. 7.31 statistical software (Muthén & Muthén, 2012) with the corrected correlation matrices from the meta-analysis used as the input matrices⁸. In order to minimize complexity, we estimated separate path analyses for each of the six illness outcomes. Each model included the six illness representation dimensions as exogenous independent predictors of the illness outcome mediated by multiple pathways through the coping strategies. We tested two models for each outcome. The first (Model 1) assumed that the effects of the illness representations were fully mediated by the coping strategies, and, as such, no direct pathways were freed between the illness representation dimensions and illness outcome. This model assumes that the coping constructs are sufficient in accounting for relations between the representation dimensions and illness outcome. The model under consideration is a generalized process model, so we included all possible pathways between the representation and coping constructs and between the coping and outcome constructs. The second model (Model 2) included direct effects of the illness representation dimensions on the outcomes. This model assumed that the mediation effects were not sufficient in accounting for the representation-outcome relations and non-zero direct effects exist. Comparison of the fit of each model with the data provided an indication as to whether the coping constructs were adequate in accounting for the effects

⁸Full correlation matrices used as input for the path analyses are available online at <https://osf.io/g48nt/>

1 of illness representations on illness outcomes (Model 1) or whether substantive direct effects
2 of the illness representations remain in the presence of the mediated paths through coping
3 (Model 2). Comparisons of nested models are usually made on the basis of incremental fit
4 indexes based on the model χ^2 test, such as the comparative fit index (CFI). CFI values
5 exceeding .95 indicate a well-fitting model and CFI differences across models exceeding .01
6 indicate substantial model misspecification (Cheung & Rensvold, 2002). However, as the
7 model including both direct and indirect effects (Model 2) was a saturated model,
8 comparisons based on the goodness-of-fit χ^2 test and incremental fit indices would not permit
9 model comparisons given that fit for Model 2 would be perfect (CFI = 1.000). Nevertheless,
10 absolute CFI values for Model 1 give an indication of the levels of misspecification in the
11 model (CFI values < .95) when direct paths from illness representations to illness outcomes
12 are not set as free parameters. As an alternative, model comparisons were made on the basis
13 of Akaike's information criterion (AIC) based on the log likelihood ratio test, with the lowest
14 absolute values indicating the model with the best fit. The average study sample size was
15 used as the input sample size for each model, due to the variability in the number of studies
16 contributing to each cell in the corrected correlation matrix.

17 Hypothesized effects of the models were evaluated using parameter estimates
18 expressed as standardized regression coefficients (β) and their 95% confidence intervals. Two
19 types of indirect effect were estimated: (a) total indirect effects representing the overall
20 indirect effect of illness representation dimensions on each outcome through all coping
21 constructs, and (b) specific indirect effects representing the indirect effects of individual
22 representation dimensions on each outcome through each coping construct. For the model
23 incorporating both direct and indirect effects (Model 2), we also computed the total effects as
24 the sum of the direct effects of illness representation dimensions on outcomes and the total
25 indirect effects through all coping constructs.

As our analyses were based on relatively large sample sizes, many smaller effects were likely to achieve statistical significance (Seaton, Marsh, & Craven, 2010). Although the sample sizes used to estimate the models for each outcome variable were relatively conservative (N range = 10,574 to 12,594), and far short the total sample size ($N = 52,599$), the likelihood of small effects obtaining statistical significance was still high. Given the dependency of the statistical significance of our parameter estimates in our model tests on sample size, we focused on effect size when evaluating the effects in our models. Based on Seaton et al. (2010), standardized parameter coefficients (β) of .10 or larger were considered substantive and very small coefficients ($\beta < .075$) regarded as trivial even if they achieved statistical significance. Parameter estimates for indirect effects, however, tend to be much smaller and with smaller standard deviations. A suggested effect size statistic is the mediation proportion or relative indirect effect (P_M), which represents the proportion of the total effect mediated by the indirect effect (Ditlevsen, Christensen, Lynch, Damsgaard, & Keiding, 2005; Huang, Sivaganesan, Succop, & Goodman, 2004). In cases where both direct and indirect effects were present in the process model, such as when the illness representation dimensions predict outcomes directly and indirectly through coping constructs, the P_M statistic provides an indication of the relative contribution the indirect effect of interest makes to the overall effect.

In addition, we conducted sensitivity analyses to illustrate the extent to which the large sample sizes in the current meta-analysis affected the precision and statistical significance of the parameter estimates in our models, and whether these changes affected our inferences. In the sensitivity analyses, our models were re-estimated with substantively smaller sample sizes. Specifically, we used the largest sample size of the included studies ($n = 3,130$) and a sample size that approximated the average of the five next-largest sample sizes of the included studies ($n = 1,000$) as input sample sizes. Given that variability estimates and

1 significance tests associated with the effect sizes are influenced by sample size, we expected
2 the width of the 95% confidence intervals of the parameter estimates in the models to
3 progressively increase, and that smaller effects would be more likely to become statistically
4 non-significant, with declining sample size. In such cases, the estimated effect may be
5 rejected even if it is not trivial in size.

6 We also conducted additional sensitivity analyses comparing model effects across
7 levels of the candidate moderator variables. We estimated the proposed models (Model 1 and
8 Model 2) using the meta-analyzed mean correlation matrices in each moderator group. We
9 compared overall fit of model across levels of the moderator using the AIC to determine
10 whether the moderator affected our determination of which model exhibited best fit derived
11 from the full-sample analysis. We also report the CFI for each model, with absolute values
12 for Model 1 illustrating the level of misspecification in the model when direct effects were
13 fixed to zero. In addition, we tested whether the pattern of effects in each model varied across
14 moderator levels using multi-group path analysis. In these analyses, paths among the illness
15 representation, coping, and outcome variables were set to be invariant across the models
16 estimated at each level of the moderator by imposing a set of equality constraints. We
17 conducted the multi-group analysis for Model 1 and Model 2, with additional constraints
18 specified in Model 2 to test for invariance in the direct effects of the illness representation
19 dimensions on the outcome variable. Adequacy of the multi-group models was evaluated
20 using multiple recommended criteria for goodness-of-fit: the CFI, the normed fit index (NFI),
21 and the root mean square error of approximation (RMSEA), with values exceeding .95 for the
22 CFI and NFI and .050 or less for the RMSEA. Incorrectly-imposed constraints indicating
23 paths that are non-invariant across levels of the moderator will result in misspecification in
24 the model fit. As with the path models estimated on the full sample of studies, sensitivity

analyses were conducted separately for each of the six outcome variables (disease state, psychological distress, psychological well-being, and role, social, and physical functioning).

Results

Corrected correlations

Averaged correlation coefficients corrected for sampling and measurement error for all variables in the meta-analysis are appear in Table 5 in Appendix J (supplemental materials) with confidence intervals, variability and heterogeneity statistics, and analyses based on funnel plots. Intercorrelations among illness representation dimensions yielded a pattern consistent with results of previous meta-analytic findings (Hagger & Orbell, 2003). Specifically, the identity, serious consequences, timeline, and emotional representation dimensions were statistically significantly and positively related to each other, and significantly and negatively related to the perceived control and illness coherence dimensions. The only exception to this pattern was the coherence-timeline relationship, which was not statistically significant. Perceived control and illness coherence were significantly and positively correlated.

Relations between illness representations and coping strategies, and between coping strategies and illness outcomes, were also consistent with previous analyses. The identity, serious consequences, and timeline dimensions were statistically significantly and positively associated with emotion venting and avoidance coping strategies, and not significantly related to the cognitive reappraisal, problem-focused generic, and problem-focused specific coping strategies. Emotional representations were significantly and positively related to the emotion venting and avoidance coping strategies, and negatively related to cognitive reappraisal and problem-focused generic strategies. Analogously, the identity, consequences, timeline, and emotional representation dimensions were significantly and positively correlated with disease state and psychological distress, and significantly and negatively related to physical, role, and

1 social functioning, and psychological well-being. Perceived control and illness coherence
2 were significantly and positively related to cognitive reappraisal, problem-focused generic,
3 problem-focused specific, and seeking social support coping strategies, and significantly and
4 negatively related to avoidance, with the exception of the relation between coherence and
5 social support. Similarly, control and coherence were significantly and positively associated
6 with physical, role, and social functioning, and psychological well-being, with the exception
7 of the correlation between coherence and social functioning.

8 Focusing on the heterogeneity statistics, according to both Hunter and Schmidt's
9 (2004) 75% rule and Cochran's Q statistic, all but 13 of the 138 effects were found to have a
10 statistically significant degree of heterogeneity across studies, indicating the likely presence
11 of moderator variables acting on the observed relationships. This was corroborated by the I^2
12 statistic, which showed at least moderate heterogeneity for the majority of the effect sizes
13 (Huedo-Medina et al., 2006). Heterogeneity variance (τ^2) values were, however, relatively
14 small. Tests of bias indicated that 44% of effect sizes were subject to some degree of small-
15 study bias according to Duval and Tweedie's (2000) trim and fill technique, and 27% of
16 effect sizes according to Egger and colleagues' (1997) regression test. While these results
17 appear to point to deviations from the expected pattern in funnel plots and potential small
18 study bias for some of the effects, the high degree of heterogeneity is a cause for concern
19 given that problems have been identified with interpreting these statistics in the presence of
20 substantial heterogeneity (Gervais, 2015; Johnson & Eagly, 2014; Moreno et al., 2009).
21 Findings indicating substantial bias using these tests in the current analysis should, therefore,
22 be interpreted in light of this concern.

23 **Moderator Analyses**

24 We examined the effect of moderators on correlations among the illness
25 representation, coping strategies, and illness outcome constructs across studies by conducting

our meta-analysis at each level of the candidate moderators: study design, illness type, medically-explained and medically-unexplained symptoms, illness stage, and methodological quality. Averaged corrected correlations, confidence intervals, and heterogeneity statistics from the moderator analyses are presented in Table 6 in Appendix K (supplemental materials). In all cases, we found little evidence for moderator effects. Of the 138 effects⁹ tested in each analysis, fewer than seven effects per moderator were significantly different across levels of the moderator and there was substantive overlap in confidence intervals in the vast majority of the tests. In addition, the analyses did little to resolve the heterogeneity in the effect sizes with moderate-to-high heterogeneity according to I^2 values and significant values for Q observed for effects within each moderator group. Of the correlations that did exhibit statistically significant differences, many included moderator groups with low numbers of studies ($k < 10$); in such cases the confidence intervals, t -tests, and heterogeneity tests may not be reliable. Overall, there was little indication of systematic variation in effect sizes attributable to the candidate moderators in the current analyses.

Path Analyses of the Process Model

Sufficiency of the process model derived from the common-sense model (Appendix A, supplemental materials) was tested using a series of mediated path-analytic models using the averaged corrected correlation matrices among the illness representation, illness outcome,

⁹There were fewer than the full 138 tests in some of the moderator analyses because some moderator groups had insufficient studies ($k < 2$) to conduct the analysis.

and coping constructs as input¹⁰. Separate analyses for each illness outcome construct (disease state, psychological distress, physical functioning, psychological well-being, role functioning, and social functioning) were conducted to minimize model complexity.

Model sufficiency. We estimated two models to test the sufficiency hypothesis.

Model 1 specified direct effects of the illness perception dimensions (identity, consequences, timeline, control, illness coherence, and emotional representations) on coping constructs (avoidance, cognitive reappraisal, emotion venting, specific and generic forms of problem-focused coping, and seeking social support), direct effects of the coping constructs on the illness outcome, and indirect effects of all the illness perceptions constructs on the illness outcome mediated by the coping constructs. Model 2 was identical to Model 1 but also included direct effects of the illness representations dimensions on illness outcomes.

Goodness-of-fit statistics for each of the six models are provided in Table 7 (Appendix L, supplemental materials). The direct and indirect effects model (Model 2) had better fit with the data than the indirect effects only model (Model 1) as indicated by lower AIC values in every case. In addition, absolute CFI values for Model 1 indicated substantive misspecification in the model ($CFI < .95$) in all cases except the model for disease state. This provides converging evidence that the indirect effects only model did not account sufficiently for relations between the illness representation and outcome variables. We therefore rejected

¹⁰Correcting for methodological artifacts is known to increase the magnitude of effect sizes, perhaps more than is appropriate, which may lead to over- or under-estimation of the true effect sizes (see Johnson & Eagly, 2014; Köhler, Cortina, Kurtessis, & Gözl, 2015). For example, interdependency between the reliability estimates of predictor and criterion variables of a particular effect could effectively lead to ‘double correction’ for measurement error (Köhler et al., 2015). Such corrections may lead to an inflation of effect sizes and increase the probability of finding statistically significant effect sizes. In order to ascertain the extent to which interdependency of reliabilities was a problem, we followed the recommendations of Köhler et al. and computed correlations between the predictor and criterion reliability estimates used to correct our effect sizes for measurement error. Results revealed that the correlations were generally small (median = .25, inter-quartile range = .37), which indicates that there may be some bias attributable to correction for this artifact but it is unlikely to be substantive in most cases. We also reestimated our models using the uncorrected averaged correlations and compared the results to the analyses using the corrected correlations (see Table 5, Appendix J, supplemental materials). Results are presented in Tables 21 to 23 in Appendix O (supplemental materials). Unsurprisingly, the analysis demonstrated that the effects for the analyses using raw data tended to be smaller than those for the corrected data. However, the pattern of effects remained in both sets of analyses, indicating that the corrections tended to alter the strength rather than pattern of effects.

Model 1 in favor of Model 2. Subsequent examination of direct and indirect effects in Model 2 was warranted to evaluate the extent to which the coping variables mediated the illness representation-illness outcome relations. Parameter estimates for the direct (Table 8), indirect (Table 9), and total (combined direct and indirect effects; Table 10) effects for each model are provided in Appendix L (supplemental materials).

Direct effects. Direct effects of illness representation dimensions on coping strategies, of coping strategies on illness outcomes, and of illness representations on outcomes provided indication of the unique predictors of the coping and illness outcome variables while simultaneously accounting for the effects of other constructs in the model (see Table 8).

Focusing first on the direct effects of the illness representation dimensions on illness outcomes, we found statistically significant, non-trivial effects of perceived consequences and illness identity on all outcomes, with the exception of the consequences-distress effect which was small by comparison. Effects were positive for disease state and distress and negative for the functioning outcomes and well-being. Perceived control had positive non-trivial direct effects that were statistically significant and greater than .10 on physical, social, and role functioning, and well-being, while effects on disease state and distress were smaller and trivial in effects size. The pattern of direct effects for these representation dimensions on outcomes followed the same pattern as the zero-order correlations among these constructs. A notable exception was the direct effect of control on disease state which was positive in the models but had a negative zero-order correlation. Emotional representations had statistically significant non-trivial negative direct effects on well-being and role functioning, and positive effects on distress and social functioning. The timeline dimension had statistically significant non-trivial positive direct effects on role functioning and well-being. Effects for emotional representations and timeline deviated from the pattern of effects in the zero-order correlations: these dimensions were negatively correlated with functioning.

1 Focusing on the direct effects of the illness representation dimensions on coping, we
2 found statistically significant non-trivial effects of the representation dimensions on the
3 coping constructs. Dimensions with the largest effects were emotional representations which
4 positively predicted avoidance, emotion venting, and seeking social support strategies, and
5 negatively predicted problem-focused generic coping and cognitive reappraisal. Control had
6 statistically significant non-trivial positive direct effects on generic and specific forms of
7 problem-focused coping, cognitive reappraisal, and social support. There were also
8 statistically significant non-trivial positive direct effects of identity on avoidance, cognitive
9 reappraisal, and emotion venting. Other effects of the illness representation dimensions on
10 coping constructs were smaller and trivial by comparison.

11 Direct effects of coping strategies on illness outcomes tended to be larger than the
12 effects of the illness representation dimensions on coping, and were consistent with the
13 theory-derived predictions from the process model and previous analyses (Hagger & Orbell,
14 2003). We found statistically significant non-trivial direct effects of avoidance on all
15 outcomes, with positive effects on disease state and distress, and negative effects on physical,
16 role, and social functioning, and well-being. A similar pattern of direct effects was found for
17 emotion venting, the only exception was that emotion venting negatively predicted disease
18 state when the correlation between these variables was not significant, which may be
19 indicative of a suppressor effect. Problem-focused generic coping had statistically significant
20 non-trivial negative direct effects on disease state, distress, and role functioning, and positive
21 effects on physical functioning and well-being. Problem-focused specific coping had
22 statistically significant non-trivial positive direct effect on role functioning, with smaller
23 trivial effects on other outcomes.

24 Overall, findings of the direct effects indicate that illness representations had
25 statistically significant non-trivial direct effects on illness outcomes, consistent with the better

1 overall fit of the model that included direct effects (Model 2) relative to the model that
2 assumed no direct effects (Model 1). Effects for the coping strategies on illness outcomes
3 tended to be larger compared to the effects of the illness representation dimensions on coping
4 constructs.

5 **Total indirect effects.** While we rejected our hypothesis that the effects of illness
6 representations on illness outcomes would be completely mediated by coping strategies, this
7 conclusion did not rule out the possibility of indirect effects consistent with hypotheses of the
8 common-sense model. Examination of the total indirect effects to establish the extent to
9 which the effects of illness representation dimensions on illness outcomes were mediated by
10 the coping strategies was warranted (Leventhal et al., 1980). Total indirect effects are
11 presented in Table 9 (Appendix L, supplemental materials).

12 We found statistically significant non-trivial positive total indirect effects of perceived
13 control on physical functioning, role functioning, and psychological well-being, and negative
14 effects on psychological distress and disease state. There were statistically significant non-
15 trivial negative total indirect effects of identity on physical, role, and social functioning, and a
16 positive effect on distress. Chronic timeline had statistically significant non-trivial positive
17 indirect effects on psychological social functioning, physical functioning, and well-being, and
18 negative effects on distress and disease state. We found statistically significant non-trivial
19 positive total indirect effects of emotional representations on disease state and distress, and
20 negative total indirect effects on physical functioning, social functioning, and well-being.
21 Other indirect effects were small and trivial by comparison. Overall, these effects provided
22 support for the indirect, coping-mediated effects of illness representation dimensions on
23 illness outcomes consistent with the process model.

We also estimated the mediation proportion for each total indirect effect (P_M), which reflects the proportion of the total effect accounted for by the indirect effect¹¹. This is directly relevant to the evaluation of model sufficiency as it provides an indication of the extent to which the total indirect effects of illness representations on outcomes through coping contribute to the total effect. A substantial P_M value indicates that the mediated pathway makes a viable contribution to explaining the link between representations and outcomes, while a trivial value indicates that the mediated path was of little relevance relative to the direct effect. Results revealed that many of the indirect effects accounted for substantial proportions of the total effects of illness representations on illness outcomes¹². Prominent among these were the control, emotional representations, timeline, and coherence dimensions, for which the total indirect effects accounted for substantive proportions of the total effect. For the identity and serious consequences dimensions, the majority of the indirect effects accounted for a trivial proportion of the total effect, with the notable exception of the indirect effect of consequences on distress. Findings provide support for the process model, verifying that the indirect effects make a substantive contribution to the overall effects in the model.

Specific indirect effects. Our analyses also enabled us to isolate specific mediated effects involving each of the illness representation, coping, and outcome variables that constituted the total indirect effects. Specific indirect effects are presented in Table 9

¹¹One of the limitations of the P_M statistic is that it is difficult to interpret when total effect comprises negative and positive direct and indirect effects, such as the effect of perceived control on disease state in the current analysis which was made up of a positive direct effect ($\beta = .034$) and a negative total indirect effect ($\beta = -.079$). The total effect ($\beta = -.045$), therefore, represents the sum of the direct and total indirect effects. In this case, the positive direct effect has the effect of attenuating the negative indirect effect. However, as the total effect serves as the denominator for the P_M calculation, it may give misleading scores for the P_M when the total effects comprise negative and positive effects, as in the previous example. This may lead, for example, to the P_M exceeding unity such that it does not represent a true proportion (Preacher & Kelley, 2011). A solution was to estimate P_M using the modulus of the total effect in its calculation. The P_M , therefore, reflects the proportion of the total effect accounted for by the indirect effect regardless of whether the combination of the direct and indirect effects leads to an attenuation of the total effect due to the combination of negative and positive scores.

¹²Although no published guidelines exist, we considered P_M values $> .25$ to be of substantive value as it corresponds to a proportion of the total effect accounted for by the indirect effect above the 25th percentile.

(Appendix L, supplemental materials). An interesting trend in the current findings was that some of the total indirect effects of representation dimensions on outcomes were either zero or relatively trivial in size. However, in some cases the total indirect effects comprised both positive and negative specific indirect effects. These specific indirect effects were approximately equal in magnitude but opposite in sign (i.e., positive and negative) leading to a null or very and trivial small total indirect effect. A consideration of the pattern of specific indirect effects, therefore, may reveal important information on the pattern of effects of the illness perception variables on outcomes that cannot be gained from observing the total indirect effects alone. The specific indirect effects may facilitate interpretation as to whether the indirect effects of illness representation dimensions have effects on illness outcomes considered adaptive (e.g., reduced distress, better functioning) or maladaptive (e.g., increased distress and disease state), through coping procedures, or whether both adaptive and maladaptive patterns of indirect effects through coping are present. We evaluate the pattern of the specific indirect effects for each illness perception dimension on each illness outcome in turn.

Perceived consequences had zero total indirect effects on disease state and physical and role functioning, and the effect on well-being although statistically significant was very small ($\beta < .01$). This was unexpected given that primary studies and meta-analyses have consistently reported statistically significant, substantive negative correlations between perceived consequences and functioning and well-being, and positive correlations with distress and disease state. However, theory suggests that perceived consequences may also have positive effects on adaptive illness outcomes by motivating individuals to take action to mitigate the threat and there is previous evidence to support this (e.g., Brewer et al., 2002). Based on this evidence, it is possible that both positive and negative specific indirect effects would be present, and that these effects would amount to null or relatively small total indirect

1 effects. Close inspection of the specific indirect effects revealed consistency in the pattern of
2 effects of consequences on each outcome via the coping constructs as mediators. Specifically,
3 consequences tended to predict maladaptive outcomes through avoidance coping and
4 adaptive outcomes through problem-focused coping. For example, consequences had
5 statistically significant and positive effects on disease state and distress through avoidance,
6 and significant negative effects on these outcomes through problem-focused generic coping.

7 The effects for consequences were corroborated by the mediation proportion statistics.
8 The specific indirect effects for consequences through avoidance and problem-focused
9 coping accounted for a substantial proportion of the overall indirect effect. There were also
10 significant negative effects of consequences on functioning and well-being through avoidance
11 and significant positive effects on these outcomes through problem-focused generic coping.
12 Together these effects were similar in magnitude and opposite in sign leading to the null or
13 relatively small total indirect effects of consequences on outcomes. The effects for the
14 consequences dimension, therefore, comprised both positive and negative indirect
15 associations with adaptive and maladaptive outcomes through coping. This is consistent with
16 the view of consequences as a representation of threat, which may motivate patients to take
17 action to deal with the threat or to engage in procedures to manage the concomitant emotional
18 upheaval. The specific indirect effects, therefore, reveal the multiple pathways by which
19 consequences impact on outcomes not indicated by the total indirect effect.

20 Similar positive and negative patterns of specific indirect effects were found for the
21 identity dimension on illness outcomes. Total indirect effects revealed statistically significant
22 effects for identity on distress, significant negative effects on physical, role, and social
23 functioning, and null effects on disease state and well-being. Decomposition of the specific
24 indirect effects revealed that the total indirect effect comprised effects of opposing sign. For
25 example, identity had significant positive indirect effects on disease state and distress through

1 avoidance, and significant negative indirect effects on physical, role and social functioning,
2 and well-being through this mediator. Mediation proportion statistics indicated that a
3 substantial proportion of the indirect effect of identity on each outcome was through
4 avoidance. There were also statistically significant positive effects of identity on physical and
5 social functioning, and well-being mediated by problem-solving generic coping, although the
6 size of these effects was small and trivial by comparison. Analogously there were statistically
7 significant but small negative effects of identity on distress and disease state through
8 problem-focused generic coping. These oppositely-valanced effects had the effect of reducing
9 the size of the total indirect effects for this representation dimension. The total indirect effects
10 of identity on outcomes were, therefore, substantially weaker due to indirect effects of
11 opposing sign. Again, this is consistent with the view of identity as representing illness threat
12 and its potential to affect both adaptive and maladaptive outcomes through multiple
13 pathways.

14 Specific indirect effects of perceived control on outcomes were consistent with the
15 pattern of the zero-order corrected correlations, and the direct and total indirect effects, for
16 this construct. Control has been identified as having a pivotal role in driving the adoption of
17 problem-focused coping strategies, so the effects on outcomes were expected to be mediated
18 by the generic and specific forms of problem-focused coping. Perceived control over the
19 illness exhibited statistically significant and positive total indirect effects on physical and role
20 functioning and psychological well-being. This was corroborated by the specific indirect
21 effects, which indicated significant positive effects of perceived control on these outcomes
22 mediated by generic and specific forms of problem-focused coping. Mediation proportion
23 statistics corroborated this pattern, with substantive proportions of the total indirect effect of
24 perceived control on each outcome accounted for by the specific indirect effects through
25 problem-focused coping. We also found significant negative specific indirect effects of

1 control on disease state and psychological distress mediated by problem-focused coping and
2 cognitive reappraisal, effects that were in keeping with the predicted pattern. Overall,
3 perceived control was related to adaptive outcomes including improvements in functioning
4 and well-being and reductions in disease state and distress.

5 Illness coherence had statistically significant positive total indirect effects on role
6 functioning and well-being and negative indirect effects on distress and disease state. Such
7 effects indicate that individuals with a clearer understanding of the illness may be better
8 equipped to identify relevant coping strategies as they are likely to have relevant information
9 regarding which strategies may be more effective. Problem-focused coping and cognitive
10 reappraisal were, therefore, expected to be key mediators of effects of coherence on
11 outcomes. However, observing the specific indirect effects revealed that the effect of
12 coherence on outcomes was not mediated by problem-focused coping, with the exception of
13 the significant positive effect of coherence on role functioning through problem-focused
14 specific coping. Instead, the specific indirect effects of coherence on outcomes were mediated
15 by avoidance and emotion venting in most cases, and substantive proportions of the indirect
16 effect were accounted for by the specific effects through these mediators. For example,
17 coherence was statistically significantly and positively related to physical, role, and social
18 functioning through emotion venting and avoidance, and significantly and negatively related
19 to distress through these variables. These effects are notable because emotion venting and
20 avoidance are typically involved in mediating effects of illness representations on
21 maladaptive outcomes. These specific indirect effects illustrate that, in some cases, illness
22 representations and the adoption of emotion-focused coping strategies may lead to functional
23 improvements and adaptive outcomes.

24 The specific indirect effects for timeline on outcomes tended to be consistent with the
25 total indirect effects. There were consistent, albeit small, statistically significant and positive

specific indirect effects of chronic timeline on physical functioning, social functioning, and well-being mediated by problem-focused generic coping, and significant negative effects on disease state and distress through problem-focused generic coping. Mediation proportion statistics also indicated that effects through problem-focused generic coping accounted for the largest proportion of the total indirect effect of timeline on all outcomes except role functioning. It is important to note that this pattern of effects was inconsistent with the zero-order correlations between timeline and outcomes. Timeline denotes chronicity and has been associated with poorer functioning and well-being, and elevated distress in a previous meta-analysis (Hagger & Orbell, 2003). The correlations, however, may not reflect the unique effects when other illness representation dimensions are taken into account and when decomposing overall effects into specific indirect effects through coping strategies.

The specific indirect effects for emotional representations on illness outcomes tended to closely mirror the direct and total indirect effects for this variable and indicated that this dimension was generally associated with maladaptive outcomes. Specifically, emotional representations were statistically significantly and positively related to distress, mediated by avoidance and emotion venting, and significantly and negatively related to psychological well-being mediated by avoidance. Emotional representations were also significantly and negatively related to social, role, and physical functioning mediated by avoidance and emotion venting, and significantly and negatively related to well-being and physical functioning through problem-focused generic coping strategies. Mediation proportion statistics revealed that the indirect effects of emotional representations through avoidance and emotion venting accounted for substantive proportions of the total indirect effects, and effects through other mediators were trivial by comparison. These findings indicate that emotional representations tend to be related to maladaptive outcomes such as poorer functioning,

greater psychological distress, and increased disease state through adoption of emotion-focused coping strategies and lower engagement in problem-focused coping strategies.

Sensitivity analysis for sample size. Results of our sensitivity analyses in which we estimated our proposed model for each outcome variable using smaller sample sizes ($n = 3,130$ and $n = 1,000$) are presented in Tables 11 to 16 (Appendix M, supplemental materials). As expected, 95% confidence intervals about the parameter estimates were progressively wider with decreasing sample size. This meant that the confidence intervals for the smaller parameter estimates were more likely to encompass zero as a possible value. The attenuation effect notwithstanding, non-trivial parameter estimates remained statistically significant according to adopted criteria even in models tested with the smallest sample size. These findings corroborate the imperative of a focus on effect size rather than statistical significance alone when interpreting results from path analyses based on correlations from a meta-analysis. Overall, the sensitivity analyses did not alter our interpretation of the pattern of main effects among model constructs.

Sensitivity Analyses of Model Effects

Although the moderator analysis demonstrated few statistically significant differences in the individual averaged corrected correlations across moderator groups, we conducted sensitivity analyses to test whether the pattern of effects among constructs in our proposed process model was dependent on levels of the study design, illness type, medically-explained vs. medically-unexplained symptoms, and methodological quality moderators^{13,14}. Given we

¹³We did not conduct sensitivity analyses for the time from diagnosis moderator due to large numbers of correlations missing in the matrices for the recently diagnosed moderator group. For example, for correlations among the coping and outcome constructs there was only sufficient data to compute two of the thirty six possible correlations for the recently diagnosed (RD) moderator group.

¹⁴Scores on the methodological quality checklist did not result in substantive variation in the magnitude of the corrected correlations across studies. This was the case for the analyses coded according to checklist score tertiles and a cut-off score of six. Both coding methods revealed extremely similar patterns of effects, although there were many instances where an effect size could not be computed due to insufficient studies for the analysis based on tertiles. We opted to use the coding based on the cut-off scores in subsequent sensitivity analyses to maximize the sample size at each level of the moderator.

found that the model including direct and indirect effects of illness representations on outcomes (Model 2) was superior for each outcome variable in full sample analyses, a key purpose of our sensitivity analyses was to test whether this was the case in the moderator groups. We also tested whether the pattern of effects in the proposed models differed across moderator groups by conducting a set of multi-group path analyses of the models constraining each path to be invariant across levels of the moderator using a set of equality constraints. Models were estimated using the averaged corrected correlation matrices as input for the path analyses for each moderator group¹⁵. Goodness-of-fit estimates for the single sample and multi-group path analyses of both models for each outcome variable and at all levels of each moderator are presented in Tables 17 to 20 (Appendix N, supplemental materials). For the single-sample analyses, lower AIC values were observed for Model 2 relative to Model 1 within each moderator group in all analyses corroborating the observed superiority in fit of this model in the full sample analysis. In addition, CFI values for Model 1 were below acceptable cutoff criteria, indicating substantive misspecification as a result of omitting the direct effects. For the multi-group models, the indirect effects only model (Model 1) exhibited fit statistics which fell short of accepted criteria for a well-fitting model in virtually every case. Conversely, goodness-of-fit indexes for Model 2 exceeded adopted criteria for good fit, and were superior to Model 1, in all analyses. These data not only provided additional confirmatory evidence for the superiority of Model 2, but also indicated that constraining model parameters to equality across moderator levels resulted in very little misspecification. This suggested that the proposed pattern of effects among the illness representation, coping, and outcome constructs in Model 2 provided a robust representation of observed relations among these constructs in the meta-analyzed data. Consistent with the

¹⁵Some of the effects in the moderator groups were tested by fewer than two studies so an averaged corrected effect size could not be computed (see Table 6, Appendix K, supplemental materials). These effects were substituted by the equivalent correlation from the full sample analysis to generate the complete correlation matrix required to run the path analyses.

moderator analysis of the correlations, these analyses indicate that the pattern of effects among the constructs in the models did not vary substantially across levels of the moderator.

Discussion

Our primary goal in the current analysis was to test a process model in which effects of illness representation dimensions on outcomes in patients with chronic illnesses or conditions were mediated by coping procedures as specified by Leventhal et al.'s (1980) common-sense model of self-regulation. We expect our analysis to contribute to a better understanding of the illness process by (a) demonstrating the unique effects of illness representation and coping variables on illness outcomes in a comprehensive, fully-inclusive model accounting for all dimensions of the common-sense model; (b) confirming the sufficiency of the process model in which coping strategies account for the effects of illness representations on illness outcomes; (c) providing information on the specific indirect pathways by which illness representations relate to illness outcomes through coping procedures; and (d) assessing the impact of candidate moderator variables on relations among the common-sense model constructs. Our analysis informs theory development by testing the sufficiency of the process model, a central but often neglected premise of the common-sense model. It also has potential implications beyond the model as relations between beliefs about threat, emotional distress and coping are key tenets of other theories of stress and coping (e.g., Lazarus & Folkman, 1984). In the next sections we discuss the findings and implications of the present analysis in each area of contribution.

Sufficiency of the Model

The present research is the first to produce a full meta-analytic inter-correlation matrix among the representation, coping, and outcome variables across studies adopting the common-sense model. Not only did this permit us to ascertain the unique effects of

1 representation and coping dimensions on illness outcomes in multivariate analyses, but to
2 also test the proposed process model based on cumulative evidence from multiple studies.

3 Examining the unique effects in the current analysis, a key finding was that the
4 process model that assumed coping fully mediated effects of illness representations on
5 outcomes was not sufficient, and direct representation-outcome relations were present. This
6 suggests that the coping procedures did not fully explain the effects of illness representations
7 on outcomes. This finding is unique because previous meta-analyses were not able to test the
8 sufficiency of the model. However, given that we found total indirect effects of the illness
9 representation dimensions constructs on illness outcomes through coping means that the
10 mediation hypothesis should not be rejected; instead both direct and indirect effects exist. Our
11 findings suggest that coping partially accounts for the effects of cognitive and emotional
12 representation dimensions on outcomes in chronic illness, but the representations also have
13 unique effects on outcomes independent of the coping constructs included in these studies.

14 The total effects from the models enabled identification of the representation
15 dimensions that, overall, contribute most to explaining variance in illness outcomes. The
16 consequences and identity dimensions emerged as consistent positive predictors of
17 maladaptive outcomes, that is, outcomes related to increased illness progression, greater
18 distress, and poorer well-being, and perceived control as a positive predictor of adaptive
19 outcomes such as better functioning and well-being, and reduced distress and disease
20 progression. Individuals interpreting their illness as having less impact on their life,
21 attributing fewer symptoms to the illness, and perceiving the illness as under control and
22 treatable are likely to experience less distress, better functioning and well-being, and reduced
23 disease state. In addition, emotional representations had a strong positive total effect on
24 distress, a strong negative effect on well-being, and weaker negative effects on physical and
25 role functioning. Individuals who are able to downplay their emotional response to the illness

1 are less likely to experience deleterious emotional outcomes and more likely to report better
2 psychological well-being. By comparison, effects of emotional representations on functioning
3 and illness progression were weaker, suggesting that reduced emotional representations may
4 not have a strong effect on outcomes related to adaptive function and illness recovery. These
5 findings illustrate the overall unique effects of the cognitive and emotional illness
6 representation dimensions on outcomes based on the current sample of studies.

7 One explanation for the insufficiency of the full mediation model is that the coping
8 measures in the included studies may not have been optimally effective in testing mediation.
9 Many studies in the present sample adopted coping measures that tap generalized coping
10 procedures rather than illness- or behavior-specific measures that precisely capture means to
11 cope with the illness. The use of generalized measures is likely to result in weaker relations
12 of the coping measures with illness representations and outcomes, a problem that has been
13 noted elsewhere (e.g., Hagger & Orbell, 2003; Heijmans, 1999). Generalized measures are
14 unlikely to capture all possible coping procedures relevant to the illness. One solution
15 employed in the current analysis was to specify a separate problem-focused specific coping
16 category, which encompassed illness-specific coping procedures. Illness-specific coping
17 strategies were expected to have closer correspondence with the representation and outcome
18 measures. However, few studies in the current sample included such measures and the
19 measures inevitably captured only one specific coping strategy when a range of specific
20 strategies may have been relevant. There is, therefore, scope to improve prediction in tests of
21 the common-sense model by moving away from generic coping scales and adopting coping
22 measures specific to the target illness.

23 An alternative interpretation for the insufficiency of the model is that the coping
24 constructs included in the current sample of studies did not adequately explain relations
25 between illness representations and outcomes. Other coping strategies or unmeasured

extraneous variables may have served to mediate the direct effects of representations on outcomes. For example, research has identified ‘all or nothing’ coping procedures in which patients with chronic illness follow cycles of full engagement in a behavioral coping strategy to manage their illness followed by complete disengagement. This pattern was not accounted for in the current model as it has only been tested in a few studies (e.g., Chilcot & Moss-Morris, 2013; Moss-Morris et al., 2011). Such patterns are likely to be adopted by patients who are high on certain traits, such as perfectionism, and prone to distress (c.f., Limburg, Watson, Hagger, & Egan, 2016). Coping in the common-sense model may, therefore, only account for some of the effects on illness beliefs on outcomes in chronic illness. Finally, the direct effects may reflect effects of representations on outcomes that are independent of consciously-accessible coping strategies. Representations of the illness may be implicitly linked with typically-adopted coping strategies in memory (Henderson et al., 2009; Orbell et al., 2015). The strategies may become salient when the illness ‘schema’ is activated and serve to mediate effects of representations on subsequent illness outcomes beyond the individual’s awareness.

The Function of Coping in the Process Model

The total indirect effects of the cognitive and emotional representation dimensions on illness outcomes were consistent with the general expected pattern of effects in many cases. However, our current analysis also revealed some important effects which were not evident on inspection of the total indirect or total effects of the illness representation constructs on outcomes. For example, zero-order correlations from the current and previous meta-analyses (Dempster et al., 2015; Hagger & Orbell, 2003; Mc Sharry et al., 2011) have revealed consistent positive relations between the consequences dimension and maladaptive outcomes (increased disease state and distress) and negative relations with adaptive outcomes (greater well-being, better functioning). In contrast, the total indirect effects from our analysis

revealed null or relatively small effects of this construct on outcomes. Furthermore, examination of the specific indirect effects revealed patterns of effect for illness representation dimensions on outcomes that could not be ascertained from the total indirect effects alone. Prominent among these were the specific indirect effects for the perceived consequences dimension which had consistent negative effects on adaptive outcomes through avoidance, and positive effects on adaptive outcomes through problem-focused coping. The presence of both positive and negative specific indirect effects of approximately equal magnitude equated to a zero total indirect effect for consequences on all but one of the illness outcomes in the current analysis. Evaluating the effects of consequences on outcomes through coping based on the total indirect effects alone would lead to an erroneous conclusion that representing the illness as having serious consequences has no indirect effect on outcomes. The specific effects reveal otherwise and indicate that illness consequences can lead to individuals selecting coping strategies that lead to both adaptive and maladaptive outcomes.

The presence of effects of illness representation dimensions on both adaptive and maladaptive outcomes mediated by coping has important ramifications for theory. These findings suggest a pattern of effects among constructs in the common-sense model that is more complex than that found in previous research syntheses. Much of the research on the common-sense model has consistently demonstrated that beliefs indicating increased threat, i.e. viewing an illness as having serious consequences, highly symptomatic, and chronic, will lead to emotion-focused coping strategies and poorer outcomes including greater disease progression, lower functioning and well-being, and greater distress (e.g., Dempster et al., 2015; Hagger & Orbell, 2003; Mc Sharry et al., 2011). However, these findings have been exclusively derived from zero-order correlations among constructs and regression models examining effects of representation dimensions on outcomes in the absence of other

1 representation dimensions and coping strategies. The current analysis indicates that these
2 overall patterns, and corresponding conclusions, may be misleading because bivariate
3 analyses do not test the multiple pathways by which representation dimensions relate to
4 outcomes through coping strategies.

5 We have demonstrated that the overall effects of representations on outcomes
6 comprise sets of specific indirect effects that are opposite in sign (i.e., positive and negative).
7 These patterns are, in fact, consistent with theory, but have seldom been shown empirically.
8 For example, while zero-order correlations indicate largely negative effects of consequences
9 on functioning and well-being, and positive effects on distress, the emergent pattern of
10 indirect effects from the mediation model indicates that consequences was positively related
11 to functioning and well-being through problem-focused coping. These findings are consistent
12 with Leventhal et al.'s (1980) original proposal that beliefs indicating elevated seriousness of
13 the illness will motivate individuals to search for coping strategies to manage the illness and
14 down-regulate the felt dissonance between health state and beliefs. There are also similar
15 specific indirect effects for timeline which show positive effects on functioning and well-
16 being, and negative effects on disease state, through problem-focused coping, while zero-
17 order correlations indicate the opposite. Although rare, previous research has also found
18 negative mediated effects of consequences on disease state and distress through problem-
19 focused coping and self-nurturing coping procedures (Benyamini et al., 2004; Brewer et al.,
20 2002). Current findings indicate that representation dimensions relate to multiple coping
21 strategies which have both positive and negative effects on outcomes related to illness
22 recovery. This pattern suggests that representation dimensions might motivate adoption of
23 different coping strategies under different circumstances. Identifying the circumstances that
24 determine the specific pattern of effects is, therefore, critical for the accurate prediction of
25 coping responses and concomitant outcomes.

Role of Context and Moderators in the Process Model

High levels of heterogeneity were observed in the majority of the effect sizes in the current analysis. This means that after correcting for methodological artifacts there was still substantive variation in the size of the relations among the common-sense model constructs across studies. Given that these effect sizes were used as input for our meta-analytic path analysis of the process model, results must be interpreted in light of the potential of the coefficients involved in the analysis to vary and for that variability to affect the strength of the effects in the process model. The effects reported in the process model, to some extent, reflect a generalized, ‘ideal’ case of relations in chronic illness, which may be indicative of potential pathways that may operate in the model, but the pathways would be dependent on extraneous moderating variables that determine whether the pathway will be present or absent. A search for moderators was, therefore, warranted.

We contended, consistent with theory and previous research on the common-sense model (Horne & Weinman, 2002; Leventhal et al., 1980; Moss-Morris et al., 2011), that contextual factors will moderate effects within the common-sense model. The factors may assist in resolving the high heterogeneity observed in the zero-order correlations among model constructs and explaining the patterns of effects in the process model. We tested the effects of contextual factors (illness type), illness characteristics (medically-explained and medically-unexplained symptoms), illness stage, and methodological artifacts (study design, methodological quality) as moderators. Contrary to expectations, our analysis did not reveal consistent moderation effects or lead to a resolution of the high levels of heterogeneity observed in relations among model constructs. Furthermore, moderator analysis reproducing our models in each moderator group corroborated the consistency of the stipulated pattern of effects across levels of the moderator. Results indicated that the model that specified both direct and indirect effects of representation dimensions on outcomes (Model 2) exhibited

1 superior fit to the indirect effects only model (Model 1) in all moderator groups. In addition,
2 constraining the paths in Model 2 to be equal across levels of the moderator resulted in well-
3 fitting models with few misspecifications in every case.

4 It is important to note that our moderator analyses were limited due to substantial
5 heterogeneity remaining in the effect sizes within moderator groups, as well as the small
6 sample sizes in many of the moderator groups. For example, studies on cancer and CVD
7 included a number of different variants of the illness that likely introduced additional within-
8 group heterogeneity. More primary research is needed that systematically tests the proposed
9 mediation effects in the process model in the presence and absence of the moderators.

10 Resolving the unique moderating effects of these characteristics may require systematic
11 comparisons of the direct and indirect effects of illness representation dimensions on
12 outcomes through coping at different levels of the moderator. For example, primary research
13 examining the moderating effect of illness type would necessitate testing relations among
14 process model constructs using appropriate measures in patients of two illnesses, preferably
15 matching samples and taking measures at a similar time from diagnosis, or, at the very least,
16 controlling for these factors. Similar procedures could be adopted in the design of studies
17 testing the moderating effects of illnesses with medically-explained or medically-unexplained
18 symptoms or a definitive treatment protocol. Tests of the moderating effects of personality
19 and dispositional factors such as optimism and perfectionism would also require systematic
20 treatment and controls, and preferably confining analyses to particular illness types or
21 controlling for their effects.

22 Tests of the moderating effects of illness stage on model pathways could also be
23 conducted in a similar way. Data on illness representations captured in very close proximity
24 to actual diagnosis, coupled with matched group or, preferably, longitudinal comparison data
25 on coping and outcomes for the same illness, would be invaluable in providing an indication

as to how links between illness representations, coping procedure selection, and illness progression and other outcomes change over time. Although challenging to collect, such data may assist in providing empirical support for the dynamic processes proposed in the common-sense model (Leventhal et al., 1980; Leventhal et al., 2016). We call for the development of well-designed studies that systematically test moderator effects in the common-sense model and move the field beyond the static, correlational designs that predominate in the literature to date. Such research should test the effects of moderators in the model and explain changes in relations among representation, coping, and outcome constructs over time. The research will provide better evidence for the processes by which beliefs about illnesses affect coping strategies and outcomes in chronic disease.

Implications for Practice

The identification of specific indirect effects in the current analysis has implications for the application of the common sense model in practice. Behavioral interventions should not only target change in certain illness representation dimensions linked to adaptive outcomes (e.g., halting or reversing disease progression, improving functioning, promoting well-being, and allaying distress), but also target the coping strategies that these representations activate. In some cases, targeting change in a given representation dimension would be an appropriate strategy if the dimension was consistently related to adaptive outcomes through the model pathways. For example, the control representation dimension has consistent effects, both direct and indirect, on adaptive illness outcomes. Promoting positive change in this dimension, therefore, would likely lead to adaptive outcomes and it should be labelled as a priority target for intervention. However, in some cases targeting intervention efforts on a particular representation dimension may have effects on adaptive and maladaptive outcomes. For example, providing messages that highlight the serious consequences of an illness to patients may lead to adaptive outcomes by prompting adoption

1 of problem-focused coping strategies, but may, in turn, lead to maladaptive outcomes like
2 distress and disease progression through avoidance. A solution might be to adopt strategies
3 that link the representation with the desired coping strategy. The common-sense model
4 implies that coping strategies may be stored schematically alongside representation
5 dimensions (Leventhal et al., 2016), and there is research suggesting that activating the
6 representation leads to a concomitant activation of the coping strategy (Henderson et al.,
7 2009). Active reinforcement of this link in interventions may, therefore, lead illness sufferers
8 to associate their beliefs about the illness with an appropriate coping strategy. Returning to
9 the example of the consequences dimension, raising patients' awareness of the serious
10 consequences of the illness and simultaneously suggesting an illness-relevant problem-
11 focused strategy that will assist in managing the threat may increase the likelihood of
12 promoting adaptive outcomes.

13 Similarly, accounting for the contextual factors that affect the indirect effects in the
14 process model is a critical consideration when developing interventions to change beliefs and
15 affect subsequent change in illness outcomes. For example, patients with illnesses that have
16 clear treatment protocols and high levels of objective control may benefit from
17 communications highlighting the consequences of the illness and the consequences of not
18 complying with treatment. In contrast, patients with illnesses that have uncontrollable or
19 uncertain outcomes may not benefit from problem-focused coping efforts that are unlikely to
20 be effective, so focusing on representations such as reducing emotional representations to
21 promote emotion-focused coping may be a more realistic and beneficial coping strategy to
22 manage threat. So while current findings may provide guidance for interventions, it is
23 important to consider such suggestions in light of the heterogeneity of the effects identified
24 and the likely context-dependency of the indirect effects. Primary research to confirm the

moderators of the indirect effects would be important additions to the formative literature to provide stronger evidence on which to base recommendations for practice.

Revising the Common-Sense Model

Our meta-analytic test of the process model has been instrumental in identifying the prominence of particular illness representation dimensions in predicting illness outcomes directly and indirectly through coping strategies. For some representation dimensions, the indirect effects exhibited a consistent pattern corroborating findings from previous meta-analyses and primary research. An example of such a pattern was the positive indirect effects of control beliefs on adaptive outcomes (e.g., better functioning and well-being) through problem-focused-coping. However, there were also representation dimensions which had both positive and negative indirect effects on outcomes. Prominent among these were positive and negative indirect effects of perceived consequences on adaptive outcomes mediated by problem-focused coping and avoidance, respectively. The presence of these paths prompts suggestions that the processes reflected in the different pathways may be dependent on contextual factors that moderate their effects. Similarly, research has identified the potential of other variables in the common-sense model, including beliefs about coping behaviors and beliefs about treatment, in accounting for variance in coping and outcomes alongside cognitive and emotional representations (e.g., French, Wade, & Farmer, 2013; Hagger, Hardcastle, et al., 2016; Orbell, Hagger, Brown, & Tidy, 2006). These findings, together with extant theory, have provided the impetus to propose a revised common-sense process model that incorporates the moderation processes and additional constructs derived from theory and previous research. Next, we outline the conceptual bases, operationalization, and specific examples of our revised model.

Based on theory and research on the proposed effects of socio-contextual and self-systems on model relations (Martin et al., 2003; Moss-Morris et al., 2011; Petrie & Weinman,

2012), we propose three key moderators of representation effects on outcomes: illness characteristics (e.g., illness type and severity, whether the illness is symptomatic or asymptomatic, and whether the illness has medically-explained or medically-unexplained symptoms), personality and individual differences (e.g., optimism, perfectionism), and emotional representations. The proposed moderating effects are illustrated in Figure 3, which outlines the generalized version of the revised model. The basic mediation effects of the process model tested in the current meta-analysis are depicted in the central section of Figure 3. In the revised model, these mediated pathways are proposed to be moderated by three sets of factors, represented in the diagram by the effects of illness type, dispositions, and emotional representations on the mediated pathways (see broken lines in Figure 3). In its generalized form, the model indicates the potential for upward and downward moderating effects on the mediated pathways involving representation dimensions, coping strategies, and outcomes (Figure 3). The revised model extends previous theory by providing a formal operationalization of Leventhal et al.'s (1992; 1980) proposal that socio-cultural and self-system constructs will impact on relations between illness representations, coping strategies, and outcomes (see upper section of Figure 1).

We provide specific examples of the moderation effects and illustrate them in Figures 4 and 5¹⁶. The moderating effect of an illness characteristic, the extent to which the illness is treatable, on model pathways is presented in Figure 4. Illnesses and conditions that are unlikely to respond to treatment (e.g., chronic pain) may mean that problem-focused coping efforts will be viewed as less effective, while emotional focused-coping such as venting or even avoidance may be more appropriate. Lower illness treatability may, therefore, moderate the effects of threat and control representations on outcomes (e.g., functioning) through

¹⁶Constructs in Figures 4 and 5 are represented such that larger scores represent higher levels of the construct. For example, high levels of identity, consequences, and timeline cognitive illness representations reflect greater threat perceptions, high levels of functioning and well-being indicate adaptive outcomes, and high levels of distress and disease state indicate maladaptive outcomes.

1 problem-focused coping downwards, and effects of threat representations on outcomes (e.g.,
2 distress, disease state) through emotion-focused coping upwards.

3 The moderating effect of emotional representations on model relations is presented in
4 Figure 5. Viewing an illness as highly emotionally distressing may lead individuals to
5 prioritize the selection of emotion-focused coping strategies aimed at managing the increased
6 distress over more problem-focused strategies. Emotional representations are, therefore,
7 depicted as moderating the positive effects of perceived consequences on increased disease
8 state and distress through emotion-focused coping upward, and the positive effects of threat
9 and control perceptions on better functioning and reduced disease state downwards. These
10 moderation effects were implied by Leventhal et al. in the initial specification of the
11 common-sense model, suggesting that emotional representations have both independent and
12 interactive effects on coping and outcomes (see Leventhal et al., 1992; Martin et al., 2003),
13 and were depicted as bidirectional arrows between representation, coping, and outcome
14 dimensions in schematics of the model (see Figure 1). However, these effects have, to date,
15 not been formally operationalized or tested empirically. This aspect of the revised process
16 model extends Leventhal et al.'s original proposals by identifying and operationalizing
17 specific, testable hypotheses regarding the potential moderation of mediated pathways.
18 Empirical verification of these pathways should be considered a priority for future research.

19 In addition to incorporating moderating factors, our revised common-sense process
20 model also incorporates the independent effects of beliefs with respect to engaging in specific
21 coping behaviors and beliefs about treatment. The inclusion of these beliefs follows theory
22 and research that has focused on integrating the common-sense model with other social
23 cognitive approaches to understanding coping and illness outcomes. For example, behavioral
24 coping with an illness can be conceptualized as a function of beliefs about the illness (e.g.,
25 threat perceptions, perceived control, and emotional responses), as indicated in the common-

sense model. It can also be conceived as a function of beliefs about engaging in the coping behavior itself, such beliefs that the behavior will result in desired outcomes (attitudes), beliefs in personal capacity to perform the behavior (self-efficacy), and beliefs in capacity to cope with difficulties or setbacks in managing the illness (coping self-efficacy), as indicated in social cognitive approaches to behavior (e.g., Bandura, 1977; Fishbein & Ajzen, 2009; Schwarzer, 2008). The modifications provide a more comprehensive perspective on the factors that are related to problem-focused coping responses to manage illness. For example, an asthmatic patient's decision to engage in a problem-focused coping procedure to manage attacks (e.g., prophylactic use of an inhaler) is not only likely to be a response to beliefs that an attack is sufficiently serious but controllable, but also beliefs that the inhaler will be effective and that they can use it appropriately. The parallel influence of cognitive illness representations and social cognitive beliefs on illness outcomes through problem-focused coping strategies is illustrated in Figure 3. Studies incorporating both sets of beliefs in tests of the common-sense model have indicated that social cognitive beliefs tend to have stronger effects and usually attenuate or extinguish the effects of cognitive representations (e.g., French et al., 2013; Hagger, Hardcastle, et al., 2016; Orbell et al., 2006).

Leventhal et al. (1992; 2016) have also suggested that beliefs about treatment, including its efficacy and perceived side effects, would affect selection of, and adherence to, treatment. Treatment can refer to numerous problem-focused coping behaviors including medication adherence and behaviors aimed at rehabilitation, recovery, and prevention of relapse (e.g., physical activity in patients with cardiovascular disease or osteoarthritis). Beliefs relating to treatment behaviors engagement can be accounted for in tests of the common sense model by incorporating constructs from social cognitive theories (e.g., attitudes, self-efficacy). However, there has also been interest in isolating patients' beliefs about medication as a specific form of treatment (Horne, 1997). Horne et al. (1999) contend

1 that medication adherence (e.g., taking anti-hypertension tablets to manage blood pressure) is
2 a function of specific beliefs about the effectiveness (e.g., taking tablets reduces blood
3 pressure at the next test) and drawbacks (e.g., debilitating side-effects of taking the tablets) of
4 medication. Research on beliefs about medication identified that patients held general beliefs
5 about the harmfulness and overuse of medication, and specific beliefs about the necessity of
6 the medication and concerns over its use. In the revised model, medication beliefs are
7 expected to explain unique variance in specific coping behaviors (i.e., medication adherence)
8 alongside illness representations and beliefs about other coping behaviors (Figure 3).

9 A growing body of research has incorporated measures of illness representations
10 alongside measures of medication beliefs (e.g., BMQ; Horne et al., 1999) in an extended
11 model of illness self-regulation (see Horne & Weinman, 2002) and examined their
12 independent effects on medication adherence, including a number of the studies included in
13 the present analysis ($n = 18$; Table 3, Appendix E, supplemental materials). Results have
14 indicated that individuals with strong beliefs in the necessity of their medication and fewer
15 concerns over its use tend to have better medication adherence independent of illness
16 perceptions (e.g., Byrne, Walsh, & Murphy, 2005; Horne & Weinman, 2002; Nicklas,
17 Dunbar, & Wild, 2010). The unique effects of beliefs about medication identified in this
18 research highlights the importance of including treatment beliefs in the revised common-
19 sense process model.

20 A specific example illustrating the effects of additional beliefs in the revised
21 common-sense model is presented in Figure 6. In this model, problem-focused coping
22 procedures are depicted as a response to illness beliefs, consistent with the original model.
23 Coping is also viewed as a response to beliefs about the coping response itself derived from
24 social cognitive theories (e.g., attitudes, self-efficacy, coping self-efficacy). Intentions are
25 also included as a mediator of the effects of illness and behavioral beliefs on coping

1 procedures to reflect the effort and motivation toward the coping response. The relative
2 contribution of each belief set advances theory by illustrating the specific pathways that
3 influence patients' decisions on the adoption of coping procedures. Testing these pathways
4 empirically within the revised model may provide formative evidence to assist in the
5 identification of the beliefs that should be targeted in interventions to promote participation in
6 appropriate coping strategies (Hagger, Hardcastle, et al., 2016; Orbell et al., 2006). The
7 model also provides the opportunity to explore potential interactions between the sets of
8 beliefs. Given research indicating that individuals are more likely to engage in health
9 behaviors in response to a perceived threat when motivation and self-efficacy are high (e.g.,
10 Peters, Rutter, & Kok, 2013), we have indicated that threat perceptions may moderate the
11 effect of intentions on problem-focused coping (see broken line in Figure 6).

12 We also incorporate action planning as an important component of the process
13 preceding the adoption of coping procedures. Leventhal et al. identified action plans as
14 critical to the implementation of coping strategies to manage illness outcomes. The plans
15 identify the specific coping response (e.g., taking an insulin injection), the context in which it
16 will be performed (e.g., in the morning 15 minutes before breakfast), and expectations of the
17 outcomes of the response (e.g., appropriately-managed blood glucose levels). In the model,
18 action plans are depicted as generated by representations of the illness threat and beliefs
19 about the behavior. Furthermore, action plans form a 'bridge' between intentions and the
20 enactment of the coping procedure, as illustrated by the mediation of the effect of intentions
21 on coping by action plans in Figure 6. Formation of action plans has been shown to be pivotal
22 for illness management by assisting individuals in the efficient enactment of an appropriate,
23 effective coping response to a threat representation (Leventhal et al., 2016). The concept of
24 action plans has also been adopted and applied as an important intervention technique in the
25 promotion of health behavior in multiple contexts (e.g., Hagger, Luszczynska, et al., 2016;

Orbell, Hodgkins, & Sheeran, 1997; Schwarzer, 2008). Action plans provide opportunities to intervene and promote better adherence to health behaviors. Whereas some patients with chronic illnesses form action plans independently, others need assistance from health professionals. Assisting patients in developing appropriate skills to generate their own action plans may provide important means to enhance coping self-regulation.

It is important to note that the proposed effects in our revised common-sense process model are speculative, based on a combination of the findings of the current research, theory, and evidence from primary research to provide potential explanations for effects identified in the current analysis. We expect the revised model to provide a starting point for future research examining the processes by which representations impact outcomes in the common-sense model and we have provided examples of some key hypotheses that might be tested. Such research will assist in further advancing the common-sense model and the processes involved in coping with chronic illness.

Strengths, Limitations and Recommendations

The current analysis has numerous strengths. It is the first test of the sufficiency of a process model, based on the common-sense model of self-regulation, in which coping responses mediate the effects of illness representations on illness outcomes across research in chronic illness using a quantitative cumulative synthesis. No previous research has tested a comprehensive model that accounts for multiple representation, coping, and outcome constructs simultaneously. Our data set comprising studies from multiple chronic illnesses allowed us to test the sufficiency of a full mediation model, in which coping strategies fully accounted for effects of representation dimensions on outcomes, compared to an alternative in which both direct and indirect effects were specified (viz. Bagozzi, 1981; Hagger, Chan, et al., 2016). In addition, our research also permitted tests of the unique effects of illness representation dimensions on coping strategies and illness outcomes that may contribute to

1 explaining coping selection and outcomes and the selection of potential targets for behavioral
2 interventions. A further strength of the current analysis is our systematic classification and
3 coding of measures of illness representation, coping, and outcome across research adopting
4 the common-sense model in chronic illness. The theory-based classification was also
5 essential to minimize potential shared variance between constructs attributable to conceptual
6 overlap. Building on a previous coding scheme developed by Hagger and Orbell (2003), we
7 developed a set of definitions of constructs and used expert raters to develop sets of
8 independent illness representation, coping, and outcome categories which accounted for all
9 measures adopted in studies eligible for inclusion in the current analysis. We have provided
10 this coding scheme in the online supplemental materials (see Appendixes F, G, and H,
11 supplemental materials) so that researchers may locate future analyses within the current one,
12 and to assist researchers in categorizing coping and outcome measures.

13 A key limitation of the present analysis is that the vast majority of included studies
14 were correlational in design, those adopting experimental or intervention designs numbered
15 relatively few by comparison. This has also been noted in previous reviews on research
16 adopting the common-sense model (Hagger & Orbell, 2003; Leventhal et al., 2016). Not only
17 does the preponderance of correlational designs limit the extent to which causal effects can be
18 inferred, it also neglects the proposed dynamic nature of the model in which experience with
19 coping strategies and changes in symptoms provide feedback for the individual to modify his
20 or her beliefs and subsequent coping procedures. While the introduction of a time lag
21 between representation and coping or outcome measures in research on the common-sense
22 model may provide some indication of proposed direction, the resulting data will be static,
23 and will not account for change over time.

24 Given the heavy dependence on correlational data, causation in the process model is
25 inferred from theory rather than data. While analytic methods such as the path analyses used

1 in the present study imply directional relations, alternative models which specify other
2 directional relations among the variables could be estimated and would be plausible from a
3 statistical and empirical perspective, even if they were not consistent with theory. The
4 correlational nature of the data may account for some of the patterns of effects among model
5 variables identified in the current analysis and in previous research. For example, similar to
6 other meta-analyses, we found negative effects of perceived consequences on problem-
7 focused coping and adaptive outcomes, when a theory-based expectation was that such
8 beliefs would serve as a stimulus for problem-focused action focused on treatment to manage
9 the threat. Given the correlational data on which this finding is based, one interpretation
10 would be that the beliefs about consequences is a result of adherence such that success with
11 problem-focused-coping behaviors may have led to better illness outcome and hence fewer
12 perceived consequences. The correlational data does not account for such dynamic processes
13 and cross-lagged panel designs in which previous experience with coping and past behavior
14 are modelled along with illness representations may provide a solution.

15 Findings from our current analysis should be interpreted with these limitations in
16 mind, and the current test does not provide strong evidence to support the directional nature
17 of the causal structure proposed in the process model. We advocate the adoption of
18 longitudinal studies adopting cross-lagged panel designs, preferably in close proximity to first
19 diagnosis, so that change in the representation, coping, and outcome constructs over time can
20 be modelled. Such analyses would better capture dynamic effects in the model proposed in its
21 original operationalization (Leventhal & Cleary, 1980; Martin et al., 2003). Similarly,
22 randomized controlled intervention designs and experimental studies in which illness beliefs
23 are manipulated and effects on subsequent coping behaviors and illness outcomes over time
24 should also be conducted. Such data would shed light on the causal and dynamic effects in
25 the model. To date, there have been few studies that have adopted experimental and

1 intervention designs that involve manipulation of common-sense model constructs and
2 examination of their effects on outcomes (e.g., Evans & Norman, 2009; Jonsbu, Martinsen,
3 Morken, Moum, & Dammen, 2013; Petrie, Cameron, Ellis, Buick, & Weinman, 2002). We
4 recognize the challenges inherent in collecting such data, but the contribution would be
5 substantive given the dearth of research testing causal and dynamic effects in the common-
6 sense model.

7 **Conclusion**

8 Our meta-analysis of the common-sense model of self-regulation advances previous
9 research by simultaneously testing the mechanisms by which illness representations relate to
10 outcomes mediated by coping strategies based on Leventhal et al.'s (1980) predictions.
11 Overall, we rejected a full mediation model in favor of a model that included both direct and
12 indirect effects of representation dimensions on illness outcomes through coping strategies.
13 We also identified specific mediated pathways which demonstrated that illness
14 representations that signal a health threat, such as perceived consequences, were related to
15 both adaptive and maladaptive outcomes through specific coping strategies. Identification of
16 these specific pathways is important for a full understanding of model effects and conclusions
17 based on zero-order relations or overall pathways could be misleading. Our tests of effects of
18 key moderators revealed few moderation effects and did not resolve the heterogeneity
19 identified in the effect sizes across studies on the model. Many of the studies in the current
20 analysis adopted self-report measures and correlational designs, and we call for research
21 adopting stronger designs, particularly intervention and experimental research, and research
22 using objective measures of specific, behavioral, problem-focused coping strategies and
23 illness outcomes. The research would be extremely informative in resolving some of the
24 relatively untested processes in the common-sense model, such as the dynamic process by
25 which patients' lay representations of illness relate to coping strategy selection, and,

1 subsequently, coping appraisals. We have also proposed a revised common-sense process
2 model that we hope will catalyze primary research testing the effects of moderators, beliefs
3 about coping, and treatment beliefs on coping behavior selection and illness outcomes. We
4 expect findings from the current analysis and revised model will stimulate future research and
5 theory development to advance knowledge on the processes by which illness beliefs affect
6 coping and outcomes in chronic disease.

7

References

References marked with an asterisk indicate studies included in the meta-analysis.

*Aalto, A.-M., Heijmans, M., Weinman, J., & Aro, A. R. (2005). Illness perceptions in coronary heart disease. Sociodemographic, illness-related, and psychosocial correlates. *Journal of Psychosomatic Research*, 58, 393-402. doi: 10.1016/j.jpsychores.2005.03.001

*Ackroyd, K., Fortune, D. G., Price, S., Howell, S., Sharrack, B., & Isaac, C. L. (2011). Adversarial growth in patients with multiple sclerosis and their partners: Relationships with illness perceptions, disability and distress. *Journal of Clinical Psychology in Medical Settings*, 18, 372-379. doi: 10.1007/s10880-011-9265-0

*Aflakseir, A. (2012). Role of illness and medication perceptions on adherence to medication in a group of Iranian patients with type 2 diabetes. *Journal of Diabetes*, 4, 243-247. doi: 10.1111/j.1753-0407.2012.00183.x

*Alsén, P., Brink, E., Persson, L.-O., Brändström, Y., & Karlson, B. W. (2010). Illness perceptions after myocardial infarction: Relations to fatigue, emotional distress, and health-related quality of life. *Journal of Cardiovascular Nursing*, 25, E1-E10. doi: 10.1097/JCN.0b013e3181c6dcfd

Anagnostopoulos, F., Dimitrakaki, C., Fitzsimmons, D., Potamianos, G., Niakas, D., & Tountas, Y. (2012). Health beliefs and illness perceptions as related to mammography uptake in randomly selected women in Greece. *Journal of Clinical Psychology in Medical Settings*, 19, 147-164. doi: 10.1007/s10880-011-9272-1

*Arat, S., Verschueren, P., De Langhe, E., Smith, V., Vanthuyne, M., Diya, L., . . . Westhovens, R. (2012). The association of illness perceptions with physical and mental health in systemic sclerosis patients: An exploratory study. *Musculoskeletal Care*, 10, 18-28. doi: 10.1002/msc.223

- 1 Bagozzi, R. P. (1981). An examination of the validity of two models of attitude. *Multivariate*
2 *Behavioral Research*, 16, 323-359. doi: 10.1207/s15327906mbr1603_4
- 3 Balck, F., Preuss, M., Hendrichske, A., & Lippmann, M. (2012). Change of illness
4 representations and quality of life during the course of a psychotherapeutic-
5 psychosomatic treatment. *Zeitschrift Fur Psychosomatische Medizin Und*
6 *Psychotherapie*, 58, 357-373. doi: 10.13109/zptm.2012.58.4.357
- 7 Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change.
8 *Psychological Review*, 84, 191-215. doi: 10.1037/0033-295X.84.2.191
- 9 *Barnes, L., Moss-Morris, R., & Kaufusi, M. (2004). Illness beliefs and adherence in diabetes
10 mellitus: A comparison between Tongan and European patients. *The New Zealand*
11 *Medical Journal*, 117, U743.
- 12 Baumann, L. J., Cameron, L. D., Zimmerman, R. S., & Leventhal, H. (1989). Illness
13 representations and matching labels with symptoms. *Health Psychology*, 8, 449-469.
14 doi: 10.1037/0278-6133.8.4.449
- 15 *Bean, D., Cundy, T., & Petrie, K. J. (2007). Ethnic differences in illness perceptions, self-
16 efficacy and diabetes self-care. *Psychology & Health*, 22, 787-811. doi:
17 10.1080/14768320600976240
- 18 *Benyamini, Y., Goner-Shilo, D., & Lazarov, A. (2012). Illness perception and quality of life
19 in patients with contact dermatitis. *Contact Dermatitis*, 67, 193-199. doi:
20 10.1111/j.1600-0536.2012.02071.x
- 21 *Benyamini, Y., Gozlan, M., & Kokia, E. (2004). On the self-regulation of a health threat:
22 Cognitions, coping, and emotions in women undergoing treatment for infertility.
23 *Cognitive Therapy and Research*, 28, 557-592. doi:
24 10.1023/B:COTR.0000045566.97966.22

- 1 *Benyamini, Y., Medalion, B., & Garfinkel, D. (2007). Patient and spouse perceptions of the
2 patient's heart disease and their associations with received and provided social support
3 and undermining. *Psychology & Health*, 22, 765-785. doi:
4 10.1080/14768320601070639
- 5 *Bishop, F. L., Yardley, L., & Lewith, G. T. (2006). Why do people use different forms of
6 complementary medicine? Multivariate associations between treatment and illness
7 beliefs and complementary medicine use. *Psychology & Health*, 21, 683-698. doi:
8 10.1080/14768320500444216
- 9 *Bishop, F. L., Yardley, L., & Lewith, G. T. (2008). Treatment appraisals and beliefs predict
10 adherence to complementary therapies: A prospective study using a dynamic extended
11 self-regulation model. *British Journal of Health Psychology*, 13, 701–718. doi:
12 10.1348/135910707X249570
- 13 Bishop, G. D., & Converse, S. A. (1986). Illness representations: A prototype approach.
14 *Health Psychology*, 5, 95-114. doi: 10.1037/0278-6133.5.2.95
- 15 *Bonsaksen, T., Haukeland-Parker, S., Lerdal, A., & Fagermoen, M. S. (2007). A 1-year
16 follow-up study exploring the associations between perception of illness and health-
17 related quality of life in persons with chronic obstructive pulmonary disease.
18 *International Journal of Chronic Obstructive Pulmonary Disease*, 9, 41-50. doi:
19 10.2147/COPD.S52700
- 20 *Bonsaksen, T., Lerdal, A., & Fagermoen, M. S. (2012). Factors associated with self-efficacy
21 in persons with chronic illness. *Scandinavian Journal of Psychology*, 53, 333-339.
22 doi: 10.1111/j.1467-9450.2012.00959.x
- 23 *Boot, C. R. L., Heijmans, M., van der Gulden, J. W. J., & Rijken, M. (2017). The role of
24 illness perceptions in labor participation of the chronically ill. *International Archives*

- 1 *of Occupational and Environmental Health*, 82, 13-20. doi: 10.1007/s00420-007-
2 0298-5
- 3 Borenstein, M. (2011). Comprehensive Meta-Analysis [Computer Program] (Version 2.2).
4 Englewood, NJ: Biostat.
- 5 Borenstein, M., Higgins, J. P. T., Hedges, L. V., & Rothstein, H. R. (2017). Basics of meta-
6 analysis: I2 is not an absolute measure of heterogeneity. *Research Synthesis Methods*,
7 8, 5-18. doi: 10.1002/jrsm.1230
- 8 Brandes, K., & Mullan, B. A. (2014). Can the common-sense model predict adherence in
9 chronically ill patients? A meta-analysis. *Health Psychology Review*, 8, 129-153. doi:
10 10.1080/17437199.2013.820986
- 11 *Brewer, N. T., Chapman, G. B., Brownlee, S., & Leventhal, E. (2002). Cholesterol control,
12 medication adherence and illness cognition. *British Journal of Health Psychology*, 7,
13 433-447. doi: 10.1348/135910702320645408
- 14 *Broadbent, E., Donkin, L., & Stroh, J. C. (2011). Illness and treatment perceptions are
15 associated with adherence to medications, diet, and exercise in diabetic patients.
16 *Diabetes Care*, 34, 338.
- 17 *Broadbent, E., Petrie, K. J., Main, J., & Weinman, J. (2006). The brief illness perception
18 questionnaire. *Journal of Psychosomatic Research*, 60, 631-637. doi:
19 10.1016/j.jpsychores.2005.10.020
- 20 Broadbent, E., Wilkes, C., Koschwanez, H., Weinman, J., Norton, S., & Petrie, K. J. (2015).
21 A systematic review and meta-analysis of the Brief Illness Perception Questionnaire.
22 *Psychology & Health*, 30, 1361-1385. doi: 10.1080/08870446.2015.1070851
- 23 Brown, S. P., Westbrook, R. A., & Challagalla, G. (2005). Good cope, bad cope: Adaptive
24 and maladaptive coping strategies following a critical negative work event. *Journal of*
25 *Applied Psychology*, 90, 792-798. doi: 10.1037/0021-9010.90.4.792

- 1 *Buck, F., Drinnan, M., Wilson, J., & Barnard, I. S. (2006). What are the illness perceptions
2 of people with dysphonia: a pilot study. *The Journal of Laryngology & Otology*, 121,
3 31-39. doi: 10.1017/S0022215106004002
- 4 *Bucks, R. S., Hawkins, K., Skinner, T. C., Horn, S., Seddon, P., & Horne, R. (2009).
5 Adherence to treatment in adolescents with cystic fibrosis: The role of illness
6 perceptions and treatment beliefs. *Journal of Pediatric Psychology*, 34, 893-902. doi:
7 10.1093/jpepsy/jsn135
- 8 *Byrne, M., Walsh, J., & Murphy, A. W. (2005). Secondary prevention of coronary heart
9 disease: Patient beliefs and health-related behaviour. *Journal of Psychosomatic*
10 *Research*, 58, 403-415. doi: 10.1016/j.jpsychores.2004.11.010
- 11 *Cameron, L. D., Booth, R. J., Schlatter, M., Ziginskis, D., Harman, J. E., & Benson, S. R.
12 C. (2005). Cognitive and affective determinants of decisions to attend a group
13 psychosocial support program for women with breast cancer. *Psychosomatic*
14 *Medicine*, 67, 584-589. doi: 10.1097/01.psy.0000170834.54970.f5
- 15 *Carlisle, A. C. S., John, A. M. H., Fife-Schaw, C., & Lloyd, M. (2005). The self-regulatory
16 model in women with rheumatoid arthritis: Relationships between illness
17 representations, coping strategies, and illness outcome. *British Journal of Health*
18 *Psychology*, 10, 571-587. doi: 10.1348/135910705X52309
- 19 *Cartwright, T., Endean, N., & Porter, A. (2009). Illness perceptions, coping and quality of
20 life in patients with alopecia. *British Journal of Dermatology*, 160, 1034-1039. doi:
21 10.1111/j.1365-2133.2008.09014.x
- 22 *Cartwright, T., & Lamb, R. (1999, September). *The self-regulatory model: A framework for*
23 *chronic illness*. Paper presented at the British Psychological Society Division of
24 Health Psychology Annual Conference 1999, Leeds, England.

1 Carver, C. S., Scheier, M. F., & Pozo, C. (1992). Conceptualizing the process of coping with
2 health problems. In H. S. Friedman (Ed.), *Hostility, coping, and health* (pp. 167-187).
3 Washington, DC: APA.

4 Carver, C. S., Scheier, M. F., & Weintraub, J. K. (1989). Assessing coping strategies: A
5 theoretically based approach. *Journal of Personality and Social Psychology*, 56, 267-
6 283. doi: 10.1037//0022-3514.56.2.267

7 *Chaboyer, W., Lee, B.-O., Wallis, M., Gillespie, B., & Jones, C. (2010). Illness
8 representations predict health-related quality of life 6 months after hospital discharge
9 in individuals with injury: A predictive survey. *Journal of Advanced Nursing*, 66,
10 2743-2750. doi: 10.1111/j.1365-2648.2010.05431.x

11 *Chalder, T., Godfrey, E., Ridsdale, L., King, M., & Wessely, S. (2003). Predictors of
12 outcome in a fatigued population in primary care following a randomized controlled
13 trial. *Psychological Medicine*, 33, 283-287. doi: 10.1017/S0033291702007006

14 *Chan, J. C. Y., Ong, J. C. Y., Avalos, G., Regan, P. J., McCann, J., Groarke, A., & Kelly, J.
15 L. (2009). Illness representations in patients with hand injury. *Journal of Plastic*,
16 *Reconstructive & Aesthetic Surgery*, 62, 927-932. doi: 10.1016/j.bjps.2007.11.057

17 *Chen, S.-L., Tsai, J.-C., & Chou, K.-R. (2011). Illness perceptions and adherence to
18 therapeutic regimens among patients with hypertension: A structural modeling
19 approach. *International Journal of Nursing Studies*, 48, 235-245. doi:
20 10.1016/j.ijnurstu.2010.07.005

21 *Chen, S.-L., Tsai, J.-C., & Lee, W.-L. (2009). The impact of illness perception on adherence
22 to therapeutic regimens of patients with hypertension in Taiwan. *Journal of Clinical*
23 *Nursing*, 18, 2234-2244. doi: 10.1111/j.1365-2702.2008.02706.x

- 1 Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing
2 measurement invariance. *Structural Equation Modeling*, 9, 233-255. doi:
3 10.1207/S15328007SEM0902_5
- 4 Chilcot, J., & Moss-Morris, R. (2013). Changes in illness-related cognitions rather than
5 distress mediate improvements in irritable bowel syndrome (IBS) symptoms and
6 disability following a brief cognitive behavioural therapy intervention. *Behaviour*
7 *Research and Therapy*, 51, 690-695. doi: 10.1016/j.brat.2013.07.007
- 8 Chilcot, J., Norton, S., Wellsted, D., & Farrington, K. (2012). The factor structure of the
9 revised illness perception questionnaire (IPQ-R) in end-stage renal disease patients.
10 *Psychology, Health and Medicine*, 17, 578-588. doi: 10.1080/13548506.2011.647702
- 11 *Chilcot, J., Norton, S., Wellsted, S., Davenport, A., Firth, J., & Farrington, K. (2013).
12 Distinct depression symptom trajectories over the first year of dialysis: Associations
13 with illness perceptions. *Annals of Behavioral Medicine*, 45, 78-88. doi:
14 10.1007/s12160-012-9410-5
- 15 *Chilcot, J., Wellsted, D., Davenport, A., & Farrington, K. (2011). Illness representations and
16 concurrent depression symptoms in haemodialysis patients. *Journal of Health*
17 *Psychology*, 16, 1127-1137. doi: 10.1177/1359105311401672
- 18 *Chilcot, J., Wellsted, D., & Farrington, K. (2010). Illness representations are associated with
19 fluid nonadherence among hemodialysis patients. *Journal of Psychosomatic*
20 *Research*, 68, 203-212. doi: 10.1016/j.jpsychores.2009.08.010
- 21 *Chilcot, J., Wellsted, D., & Farrington, K. (2011). Illness perceptions predict survival in
22 haemodialysis patients. *American Journal of Nephrology*, 33, 358-363. doi:
23 10.1159/000326752

- *Chong, J., Mackey, A. H., Broadbent, E., & Stott, N. S. (2012). Children's perceptions of their cerebral palsy and their impact on life satisfaction. *Disability and Rehabilitation*, 34, 2053-2060. doi: 10.3109/09638288.2012.669021
- Cochran, W. G. (1952). The χ^2 test of goodness of fit. *Annals of Mathematical Statistics*, 23, 315-345. doi: 10.1214/aoms/1177692778
- *Cooper, M., Stockford, K., & Turner, H. (2007). Stages of change in anorexic and bulimic disorders: The importance of illness representations. *Eating Behaviors*, 8, 474-484. doi: 10.1016/j.eatbeh.2007.01.003
- *Corter, A. L., Findlay, M., Broom, R., Porter, D., & Petrie, K. J. (2013). Beliefs about medicine and illness are associated with fear of cancer recurrence in women taking adjuvant endocrine therapy for breast cancer. *British Journal of Health Psychology*, 18, 168-181. doi: 10.1111/bjhp.12003
- *Costanzo, E. S., Lutgendorf, S. K., & Roeder, S. L. (2011). Common-sense beliefs about cancer and health practices among women completing treatment for breast cancer. *Psycho-Oncology*, 20, 53-61. doi: 10.1002/pon.1707
- *Covic, A., Seica, A., Gusbeth-Tatomir, P., Gavrilovici, O., & Goldsmith, D. J. A. (2004). Illness representations and quality of life scores in haemodialysis patients. *Nephrology Dialysis Transplantation*, 19, 2078-2083. doi: 10.1093/ndt/gfh254
- *Dalbeth, N., Petrie, K. J., House, M., Chong, J., Leung, W., Chegudi, R., . . . Taylor, W. J. (2011). Illness perceptions in patients with gout and the relationship with progression of musculoskeletal disability. *Arthritis Care & Research*, 63, 1605-1612. doi: 10.1002/acr.20570
- *Daleboudt, G. M. N., Broadbent, E., McQueen, F., & Kaptein, A. A. (2011). Intentional and unintentional treatment nonadherence in patients with systemic lupus erythematosus. *Arthritis Care & Research*, 63, 342-350. doi: 10.1002/acr.20411

- 1 *De Filippis, S., Erbuto, D., Gentili, F., Innamorati, M., Lester, D., Tatarelli, R., . . . Pompili,
2 M. (2008). Mental turmoil, suicide risk, illness perception, and temperament, and
3 their impact on quality of life in chronic daily headache. *The Journal of Headache*
4 *and Pain*, 9, 349-357. doi: 10.1007/s10194-008-0072-4
- 5 *de Rooij, A., van der Leeden, M., Roorda, L. D., Steultjens, M. P., & Dekker, J. (2013).
6 Predictors of outcome of multidisciplinary treatment in chronic widespread pain: an
7 observational study. *BMC Musculoskeletal Disorders*, 14, 133. doi: 10.1186/1471-
8 2474-14-133
- 9 Deary, V., Chalder, T., & Sharpe, M. (2007). The cognitive behavioural model of medically
10 unexplained symptoms: A theoretical and empirical review. *Clinical Psychology*
11 *Review*, 27, 781-797. doi: 10.1016/j.cpr.2007.07.002
- 12 *DeJong, H., Hillcoat, J., Perkins, S., Grover, M., & Schmidt, U. (2012). Illness perception in
13 bulimia nervosa. *Journal of Health Psychology*, 17, 399-408. doi:
14 10.1177/1359105311416874
- 15 Dempster, M., Howell, D., & McCorry, N. K. (2015). Illness perceptions and coping in
16 physical health conditions: A meta-analysis. *Journal of Psychosomatic Research*, 79,
17 506-513. doi: 10.1016/j.jpsychores.2015.10.006
- 18 *Dempster, M., McCorry, N. K., Brennan, E., Donnelly, M., Murray, L., & Johnston, B. T.
19 (2012). Psychological distress among survivors of esophageal cancer: The role of
20 illness cognitions and coping. *Diseases of the Esophagus*, 25, 222-227. doi:
21 10.1111/j.1442-2050.2011.01233.x
- 22 *Dempster, M., McCorry, N. K., Brennan, E., Donnelly, M., Murray, L. J., & Johnston, B. T.
23 (2011a). Do changes in illness perceptions predict changes in psychological distress
24 among oesophageal cancer survivors? *Journal of Health Psychology*, 16, 500-509.
25 doi: 10.1177/1359105310386633

- 1 *Dempster, M., McCorry, N. K., Brennan, E., Donnelly, M., Murray, L. J., & Johnston, B. T.
2 (2011b). Illness perceptions among carer-survivor dyads are related to psychological
3 distress among Oesophageal cancer survivors. *Journal of Psychosomatic Research*,
4 70, 432-439. doi: 10.1016/j.jpsychores.2010.07.007
- 5 *Dennison, L., Moss-Morris, R., Silber, E., Galea, I., & Chalder, T. (2010). Cognitive and
6 behavioural correlates of different domains of psychological adjustment in early-stage
7 multiple sclerosis. *Journal of Psychosomatic Research*, 69, 353-361. doi:
8 10.1016/j.jpsychores.2010.04.009
- 9 *Dickens, C., Cherrington, A., & McGowan, L. (2011). Do cognitive and behavioral factors
10 mediate the impact of depression on medical outcomes in people with coronary heart
11 disease? *Journal of Cardiopulmonary Rehabilitation and Prevention*, 31. doi:
12 10.1097/HCR.0b013e318207d32b
- 13 *Dickens, C., McGowan, L., Percival, C., Tomenson, B., Cotter, L., Heagerty, A., & Creed,
14 F. (2008). Negative illness perceptions are associated with new-onset depression
15 following myocardial infarction. *General Hospital Psychiatry*, 30, 414-420. doi:
16 10.1016/j.genhosppsy.2008.04.003
- 17 DiMatteo, M. R., Haskard-Zolnierrek, K. B., & Martin, L. R. (2012). Improving patient
18 adherence: A three-factor model to guide practice. *Health Psychology Review*, 6, 74-
19 91. doi: 10.1080/17437199.2010.537592
- 20 Ditlevsen, S., Christensen, U., Lynch, J., Damsgaard, M. T., & Keiding, N. (2005). The
21 mediation proportion: A structural equation approach for estimating the proportion of
22 exposure effect on outcome explained by an intermediate variable. *Epidemiology*, 16,
23 114-120. doi: 10.1097/01.ede.0000147107.76079.07

- 1 *Dorrian, A., Dempster, M., & Adair, P. (2009). Adjustment to inflammatory bowel disease:
2 the relative influence of illness perceptions and coping. *Inflammatory Bowel Diseases*,
3 15, 47-55. doi: 10.1002/ibd.20583
- 4 Duval, S., & Tweedie, R. L. (2000). Trim and fill: A simple funnel plot based method of
5 testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455–463.
6 doi: 10.1111/j.0006-341X.2000.00455.x
- 7 *Edgar, K. A., & Skinner, T. C. (2003). Illness representations and coping as predictors of
8 emotional well-being in adolescents with type 1 diabetes. *Journal of Pediatric*
9 *Psychology*, 28, 485-493. doi: 10.1093/jpepsy/jsg039
- 10 *Edwards, R., Suresh, R., Lynch, S., Clarkson, P., & Stanley, P. (2001). Illness perceptions
11 and mood in chronic fatigue syndrome. *Journal of Psychosomatic Research*, 50, 65-
12 68. doi: 10.1016/S0022-3999(00)00204-X
- 13 Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Meta-analyses: Separating
14 the good from the bad. *British Medical Journal*, 315, 629-634. doi:
15 10.1136/bmj.315.7109.629
- 16 *Eiser, J. R., Riazi, A., Eiser, C., Hammersley, S., & Tooke, J. E. (2001). Predictors of
17 psychological well-being in types 1 and 2 diabetes. *Psychology and Health*, 16, 99-
18 110. doi: 10.1080/08870440108405492
- 19 Erbllich, J., Montgomery, G. H., Valdimarsdottir, H. B., Cloitre, M., & Bovbjerg, D. (2003).
20 Biased cognitive processing of cancer-related information among women with family
21 histories of breast cancer: Evidence from a cancer Stroop task. *Health Psychology*, 22,
22 235-244. doi: 10.1037/0278-6133.22.3.235
- 23 *Evans, D., & Norman, P. (2009). Illness representations, coping and psychological
24 adjustment to Parkinson's disease. *Psychology & Health*, 24, 1181-1196. doi:
25 10.1080/08870440802398188

- 1 *Fabbri, S., Kapur, N., Wells, A., & Creed, F. (2001). Emotional, cognitive, and behavioral
2 characteristics of medical outpatients. *Psychosomatics*, 42, 74-77. doi:
3 10.1176/appi.psy.42.1.74
- 4 *Fan, S.-Y., Eiser, C., Ho, M.-C., & Lin, C.-Y. (2013). Health-related quality of life in
5 patients with hepatocellular carcinoma: the mediation effects of illness perceptions
6 and coping. *Psycho-Oncology*, 22, 1353-1360. doi: 10.1002/pon.3146
- 7 *Fennessy, M. M., DeVon, H. A., Ryan, C., Lopez, J. J., & Zerwic, J. J. (2013). Changing
8 illness perceptions and adherence to dual antiplatelet therapy in patients with stable
9 coronary disease. *Journal of Cardiovascular Nursing*, 28, 573-583. doi:
10 10.1097/JCN.0b013e31825d6060
- 11 Field, A. P. (2001). Meta-analysis of correlation coefficients: A Monte Carlo comparison of
12 fixed- and random- effects methods. *Psychological Methods*, 6, 161-180. doi:
13 10.1037/1082-989X.6.2.161
- 14 Figueiras, M. J., & Alves, N. C. (2007). Lay perceptions of serious illnesses: An adapted
15 version of the Revised Illness Perception Questionnaire (IPQ-R) for healthy people.
16 *Psychology and Health*, 22, 143-158. doi: 10.1080/14768320600774462
- 17 *Fischer, M. J., Scharloo, M., Abbink, J., van 't Hul, A. J., van Ranst, D., Rudolphus, A., . . .
18 Kaptein, A. A. (2010). The dynamics of illness perceptions: Testing assumptions of
19 Leventhal's common-sense model in a pulmonary rehabilitation setting. *British*
20 *Journal of Health Psychology*. doi: 10.1348/135910710X492693
- 21 *Fischer, M. J., Scharloo, M., Abbink, J. J., van 't Hul, A. J., van Ranst, D., Rudolphus, A., . . .
22 . Kaptein, A. A. (2009). Drop-out and attendance in pulmonary rehabilitation: The
23 role of clinical and psychosocial variables. *Respiratory Medicine*, 103, 1564-1571.
24 doi: 10.1016/j.rmed.2008.11.020

- *Fischer, M. J., Wiesenhaan, M. E., Heijer, A. D.-d., Kleijn, W. C., Nortier, J. W. R., & Kaptein, A. A. (2013). From despair to hope: A longitudinal study of illness perceptions and coping in a psycho-educational group intervention for women with breast cancer. *British Journal of Health Psychology*, *18*, 526-545. doi: 10.1111/j.2044-8287.2012.02100.x
- Fishbein, M., & Ajzen, I. (2009). *Predicting and changing behavior: The reasoned action approach*. New York, NY: Psychology Press.
- Fitzgerald, J. T., Davis, W. K., Connell, C. M., Hess, G. E., Funnell, M. M., & Hiss, R. G. (1996). Development and validation of the diabetes care profile. *Evaluation & the Health Professions*, *19*, 208-230. doi: 10.1177/016327879601900205
- Folkman, S., & Lazarus, R. S. (1988). *Ways of coping questionnaire: Research edition*. Palo Alto, CA: Consulting Psychologists Press Inc.
- *Ford, S., Howard, R., & Oyeboode, J. (2012). Psychosocial aspects of coeliac disease: A cross-sectional survey of a UK population. *British Journal of Health Psychology*, *17*, 743-757. doi: 10.1111/j.2044-8287.2012.02069.x
- *Fortune, D. G., Richards, H. L., Main, C. J., & Griffiths, C. E. M. (2000). Pathological worrying, illness perceptions and disease severity in patients with psoriasis. *British Journal of Health Psychology*, *5*, 71-82. doi: 10.1348/135910700168775
- *Foster, N. E., Bishop, A., Thomas, E., Main, C. J., Horne, R., Weinman, J., & Hay, E. (2008). Illness perceptions of low back pain patients in primary care: What are they, do they change and are they associated with outcome? *PAIN*, *136*, 177-187. doi: 10.1016/j.pain.2007.12.007
- *Foster, N. E., Thomas, E., Bishop, A., Dunn, K. M., & Main, C. J. (2010). Distinctiveness of psychological obstacles to recovery in low back pain patients in primary care. *PAIN*, *148*, 398-406. doi: 10.1016/j.pain.2009.11.002

- 1 *Fowler, C., & Baas, L. S. (2006). Illness representations in patients with chronic kidney
2 disease on maintenance hemodialysis. *Nephrology Nursing Journal*, 33, 173-187.
- 3 French, D. P., Cooper, A., & Weinman, J. (2006). Illness perceptions predict attendance at
4 cardiac rehabilitation following acute myocardial infarction: A systematic review with
5 meta-analysis. *Journal of Psychosomatic Research*, 61, 757-767. doi:
6 10.1016/j.jpsychores.2006.07.029
- 7 *French, D. P., Lewin, R. J. P., Watson, N., & Thompson, D. R. (2005). Do illness
8 perceptions predict attendance at cardiac rehabilitation and quality of life following
9 myocardial infarction? *Journal of Psychosomatic Research*, 59, 315-322. doi:
10 10.1016/j.jpsychores.2005.03.010
- 11 *French, D. P., Wade, A. N., & Farmer, A. J. (2013). Predicting self-care behaviours of
12 patients with type 2 diabetes: The importance of beliefs about behaviour, not just
13 beliefs about illness. *Journal of Psychosomatic Research*, 74, 327-333. doi:
14 10.1016/j.jpsychores.2012.12.008
- 15 *French, D. P., Wade, A. N., Yudkin, P., Neil, H. A. W., Kinmonth, A. L., & Farmer, A. J.
16 (2008). Self-monitoring of blood glucose changed non-insulin-treated Type 2 diabetes
17 patients' beliefs about diabetes and self-monitoring in a randomized trial. *Diabetic*
18 *Medicine*, 25, 1218-1228. doi: 10.1111/j.1464-5491.2008.02569.x
- 19 *Fu, L., Bundy, C., & Sadiq, S. A. (2011). Psychological distress in people with
20 disfigurement from facial palsy. *Eye*, 25, 1322-1326. doi: 10.1038/eye.2011.158
- 21 *Galli, U., Ettlin, D. A., Palla, S., Ehlert, U., & Gaab, J. (2010). Do illness perceptions
22 predict pain-related disability and mood in chronic orofacial pain patients? A 6-month
23 follow-up study. *European Journal of Pain*, 14, 550-558. doi:
24 10.1016/j.ejpain.2009.08.011

- *Gandy, M., Sharpe, L., Perry, K. N., Miller, L., Thayer, Z., Boserio, J., & Mohamed, A. (2013). The psychosocial correlates of depressive disorders and suicide risk in people with epilepsy. *Journal of Psychosomatic Research*, 74, 227-232. doi: 10.1016/j.jpsychores.2012.11.001
- *Gaston, A. M., Cottrell, D. J., & Fullen, T. (2012). An examination of how adolescent-caregiver dyad illness representations relate to adolescents' reported diabetes self-management. *Child: Care, Health and Development*, 38, 513-519. doi: 10.1111/j.1365-2214.2011.01269.x
- Gervais, W. (2015). Putting PET-PEESE to the test. Retrieved July 5, 2015, from <http://willgervais.com/blog/2015/6/25/putting-pet-peese-to-the-test-1>
- *Gillanders, D. T., Ferreira, N. B., Bose, S., & Esrich, T. (2013). The relationship between acceptance, catastrophizing and illness representations in chronic pain. *European Journal of Pain*, 17, 893-902. doi: 10.1002/j.1532-2149.2012.00248.x
- *Giri, P., Poole, J., Nightingale, P., & Robertson, A. (2009). Perceptions of illness and their impact on sickness absence. *Occupational Medicine*, 59, 550-555. doi: 10.1093/occmed/kqp123
- *Glasgow, R. E., Hampson, S. E., Stryker, L. A., & Ruggiero, L. (1997). Personal-model beliefs and social-environmental barriers related to diabetes self-management. *Diabetes Care*, 20, 556-561. doi: 10.2337/diacare.20.4.556
- *Glattacker, M., Opitz, U., & Jackel, W. H. (2011). Illness representations in women with fibromyalgia. *British Journal of Health Psychology*, 15, 367-387. doi: 10.1348/135910709x466315
- *Glover, I., Novakovic, A., & Hunter, M. S. (2002). An exploration of the nature and causes of distress in women attending gynecology outpatient clinics. *Journal of*

1 *Psychosomatic Obstetrics & Gynecology*, 23, 237-248. doi:

2 10.3109/01674820209074678

3 *Goldstein, L. H., Holland, L., Soteriou, H., & Mellers, J. D. C. (2005). Illness
4 representations, coping styles and mood in adults with epilepsy. *Epilepsy Research*,
5 67, 1-11. doi: 10.1016/j.eplepsyres.2005.06.008

6 *Goldstein, L. H., Seed, P. T., Clark, L. V., Dowson, A. J., Jenkins, L. M., & Ridsdale, L.
7 (2011). Predictors of outcome in patients consulting their general practitioners for
8 headache: A prospective study. *Psychology & Health*, 26, 751-764. doi:
9 10.1080/08870446.2010.493217

10 *Goodman, D., Morrissey, S., Graham, D., & Bossingham, D. (2012). The Application of
11 Cognitive—Behaviour Therapy in Altering Illness Representations of Systemic Lupus
12 Erythematosus. *Behaviour Change*, 22, 156-171. doi: 10.1375/behc.2005.22.3.156

13 Goodman, R. A., Posner, S. F., Huang, E. S., Perekh, A. K., & Koh, H. K. (2013). Defining
14 and measuring chronic conditions: Imperatives for research, policy, program, and
15 practice. *Preventing Chronic Disease*, 10, 120239. doi: 10.5888/pcd10.120239

16 *Gould, R. V., Brown, S. L., & Bramwell, R. (2010). Psychological adjustment to
17 gynaecological cancer: Patients' illness representations, coping strategies and mood
18 disturbance. *Psychology & Health*, 25, 633-646. doi: 10.1080/08870440902811163

19 *Gourounti, K., Anagnostopoulos, F., Potamianos, G., Lykeridou, K., Schmidt, L., &
20 Vaslamatzis, G. (2012). Perception of control, coping and psychological stress of
21 infertile women undergoing IVF. *Reproductive BioMedicine Online*, 24, 670-679. doi:
22 10.1016/j.rbmo.2012.03.002

23 *Grace, S. L., Krepostman, S., Brooks, D., Arthur, H., Scholey, P., Suskin, N., . . . Stewart,
24 D. E. (2005). Illness perceptions among cardiac patients: Relation to depressive

1 symptomatology and sex. *Journal of Psychosomatic Research*, 59, 153-160. doi:

2 10.1016/j.jpsychores.2005.05.005

3 *Graham, C. D., Weinman, J., Sadjadi, R., Chalder, T., Petty, R. E., Hanna, M. G., . . . Rose,

4 M. R. (2013). A multicentre postal survey investigating the contribution of illness

5 perceptions, coping and optimism to quality of life and mood in adults with muscle

6 disease. *Clinical Rehabilitation*, 28, 508-519. doi: 10.1177/0269215513511340

7 *Graves, H., Scott, D. L., Lempp, H., & Weinman, J. (2009). Illness beliefs predict disability
8 in rheumatoid arthritis. *Journal of Psychosomatic Research*, 67, 417-423. doi:

9 10.1016/j.jpsychores.2009.01.006

10 Gray, L. J., Khunti, K., Williams, S., Goldby, S., Troughton, J., Yates, T., . . . , L. s. P. D. C.

11 (2012). Let's prevent diabetes: Study protocol for a cluster randomised controlled trial

12 of an educational intervention in a multi-ethnic UK population with screen detected

13 impaired glucose regulation. *Cardiovascular Diabetology*, 11, 56. doi: 10.1186/1475-

14 2840-11-56

15 *Gray, S. E., & Rutter, D. R. (2007). Illness representations in young people with chronic
16 fatigue syndrome. *Psychology & Health*, 22, 159-174. doi:

17 10.1080/14768320600774595

18 *Grayson, P. C., Amudala, N. A., McAlear, C. A., Leduc, R. L., Shereff, D., Richesson, R., .

19 . . Merkel, P. A. (2013). Illness perceptions and fatigue in systemic vasculitis.

20 *Arthritis Care & Research*, 65, 1835-1843. doi: 10.1002/acr.22069

21 *Griva, K., Davenport, A., Harrison, M., & Newman, S. P. (2010). An evaluation of illness,
22 treatment perceptions, and depression in hospital- vs. home-based dialysis modalities.

23 *Journal of Psychosomatic Research*, 69, 363-370. doi:

24 10.1016/j.jpsychores.2010.04.008

- 1 *Griva, K., Davenport, A., Harrison, M., & Newman, S. P. (2012). The impact of treatment
2 transitions between dialysis and transplantation on illness cognitions and quality of
3 life – A prospective study. *British Journal of Health Psychology*, 17, 812-827. doi:
4 10.1111/j.2044-8287.2012.02076.x
- 5 *Griva, K., Jayasena, D., Davenport, A., Harrison, M., & Newman, S. P. (2009). Illness and
6 treatment cognitions and health related quality of life in end stage renal disease.
7 *British Journal of Health Psychology*, 14, 17-34. doi: 10.1348/135910708X292355
- 8 *Griva, K., Myers, L. B., & Newman, S. (2000). Illness perceptions and self-efficacy beliefs
9 in adolescents and young adults with insulin dependent diabetes mellitus. *Psychology*
10 *and Health*, 15, 733-750. doi: 10.1080/08870440008405578
- 11 *Groarke, A., Curtis, R., Coughlan, R., & Gsel, A. (2004). The role of perceived and actual
12 disease status in adjustment to rheumatoid arthritis. *Rheumatology*, 43, 1142-1149.
13 doi: 10.1093/rheumatology/keh262
- 14 *Groarke, A., Curtis, R., Coughlan, R., & Gsel, A. (2005). The impact of illness
15 representations and disease activity on adjustment in women with rheumatoid
16 arthritis: A longitudinal study. *Psychology & Health*, 20, 597-613. doi:
17 10.1080/14768320500094177
- 18 *Gump, B. B., Matthews, K. A., Scheier, M. F., Schulz, R., Bridges, M. W., & Magovern, G.
19 J. (2001). Illness representations according to age and effects on health behaviors
20 following coronary artery bypass graft surgery. *Journal of the American Geriatrics*
21 *Society*, 49, 284-289.
- 22 Hagger, M. S. (2006). Meta-analysis in sport and exercise research: Review, recent
23 developments, and recommendations. *European Journal of Sport Science*, 6, 103-115.
24 doi: 10.1080/17461390500528527

- 1 Hagger, M. S., & Chatzisarantis, N. L. D. (2016). The trans-contextual model of autonomous
2 motivation in education: Conceptual and empirical issues and meta-analysis. *Review*
3 *of Educational Research*, 86, 360-407. doi: 10.3102/0034654315585005
- 4 Hagger, M. S., Chan, D. K. C., Protogerou, C., & Chatzisarantis, N. L. D. (2016). Using
5 meta-analytic path analysis to test theoretical predictions in health behavior: An
6 illustration based on meta-analyses of the theory of planned behavior. *Preventive*
7 *Medicine*, 89, 154-161. doi: 10.1016/j.ypmed.2016.05.020
- 8 *Hagger, M. S., Chatzisarantis, N. L. D., Griffin, M., & Thatcher, J. (2005). Injury
9 representations, coping, emotions, and functional outcomes in athletes with sport-
10 related injuries: A test of self-regulation theory. *Journal of Applied Social*
11 *Psychology*, 35, 2345-2374. doi: 10.1111/j.1559-1816.2005.tb02106.x
- 12 Hagger, M. S., Hardcastle, S. J., Hingley, C., Strickland, E., Pang, J., & Watts, G. F. (2016).
13 Predicting self-management behaviors in familial hypercholesterolemia using an
14 integrated theoretical model: The impact of beliefs about illnesses and beliefs about
15 behaviors. *International Journal of Behavioral Medicine*, 23, 282-294. doi:
16 10.1007/s12529-015-9531-x
- 17 Hagger, M. S., Luszczynska, A., de Wit, J., Benyamini, Y., Burkert, S., Chamberland, P.-E., .
18 . . Gollwitzer, P. M. (2016). Implementation intention and planning interventions in
19 health psychology: Recommendations from the Synergy expert group for research and
20 practice. *Psychology & Health*, 31, 814–839. doi: 10.1080/08870446.2016.1146719
- 21 Hagger, M. S., & Orbell, S. (2003). A meta-analytic review of the common-sense model of
22 illness representations. *Psychology and Health*, 18, 141-184. doi:
23 10.1080/088704403100081321

- 1 Hagger, M. S., & Orbell, S. (2005). A confirmatory factor analysis of the revised illness
2 perception questionnaire (IPQ-R) in a cervical screening context. *Psychology and*
3 *Health, 20*, 161-173. doi: 10.1080/0887044042000334724
- 4 *Hagger, M. S., & Orbell, S. (2006). Illness representation and emotion in people with
5 abnormal screening results. *Psychology and Health, 21*, 183-209. doi:
6 10.1080/14768320500223339
- 7 *Hallas, C. N., Wray, J., Andreou, P., & Banner, N. R. (2011). Depression and perceptions
8 about heart failure predict quality of life in patients with advanced heart failure. *Heart*
9 *& Lung: The Journal of Acute and Critical Care, 40*, 111-121. doi:
10 10.1016/j.hrtlng.2009.12.008
- 11 *Hampson, S. E., Glasgow, R. E., & Foster, L. S. (1995). Personal models of diabetes among
12 older adults: Relationship to self-management and other variables. *The Diabetes*
13 *Educator, 21*, 300-306. doi: 10.1177/014572179502100407
- 14 *Hampson, S. E., Glasgow, R. E., & Stryker, L. A. (2000). Beliefs versus feelings: A
15 comparison of personal models and depression for predicting multiple outcomes in
16 diabetes. *British Journal of Health Psychology, 5*, 27-40. doi:
17 10.1348/135910700168748
- 18 Hampson, S. E., Glasgow, R. E., & Toobert, D. J. (1990). Personal models of diabetes and
19 their relations to self-care activities. *Health Psychology, 9*, 632-646. doi:
20 10.1037/0278-6133.9.5.632
- 21 *Hampson, S. E., Glasgow, R. E., & Zeiss, A. M. (1994). Personal models of osteoarthritis
22 and their relation to self-management activities and quality of life. *Journal of*
23 *Behavioral Medicine, 17*, 143-158. doi: 10.1007/BF01858102
- 24 *Han, S. W., McColl, E., Barton, J. R., James, P., Steen, I. N., & Welfare, M. R. (2005).
25 Predictors of quality of life in ulcerative colitis.: The importance of symptoms and

1 illness representations. *Inflammatory Bowel Diseases*, 11, 24-34. doi:

2 10.1097/00054725-200501000-00004

3 *Hartman, J. E., Boezen, H. M., de Greef, M. H., & ten Hacken, N. H. (2013). Physical and

4 psychosocial factors associated with physical activity in patients with chronic

5 obstructive pulmonary disease. *Archives of Physical Medicine and Rehabilitation*, 94,

6 2396-2402.e2397. doi: 10.1016/j.apmr.2013.06.029

7 *Heerema-Poelman, A., Stuive, I., & Wempe, J. B. (2013). Adherence to a maintenance

8 exercise program 1 year after pulmonary rehabilitation: What are the predictors of

9 dropout? *Journal of Cardiopulmonary Rehabilitation and Prevention*, 33, 419-426.

10 doi: 10.1097/HCR.0b013e3182a5274a

11 *Heijmans, M. (1998). Coping and adaptive outcome in chronic fatigue syndrome:

12 Importance of illness cognitions. *Journal of Psychosomatic Research*, 45, 39-51. doi:

13 10.1016/s0022-3999(97)00265-1

14 *Heijmans, M. (1999). The role of patients' illness representations in coping and functioning

15 with Addison's disease. *British Journal of Health Psychology*, 4, 137-149. doi:

16 10.1348/135910799168533

17 *Heijmans, M., & De Ridder, D. (1998a). Assessing illness representations of chronic illness:

18 Explorations of their disease-specific nature. *Journal of Behavioral Medicine*, 21,

19 485-503. doi: 10.1023/a:1018788427100

20 *Heijmans, M., & De Ridder, D. (1998b). Structure and determinants of illness

21 representations in chronic disease: A comparison of Addison's disease and chronic

22 fatigue syndrome. *Journal of Health Psychology*, 3, 523-537. doi:

23 10.1177/135910539800300406

24 *Helder, D. I., Kaptein, A. A., van Kempen, G. M. J., Weinman, J., van Houwelingen, H. C.,

25 & Roos, R. A. C. (2002). Living with Huntington's disease: Illness perceptions,

- 1 coping mechanisms, and patients' well-being. *British Journal of Health Psychology*,
2 7, 449-462. doi: 10.1348/135910702320645417
- 3 *Hemphill, R. C., Stephens, M. A. P., Rook, K. S., Franks, M. M., & Salem, J. K. (2013).
4 Older adults' beliefs about the timeline of type 2 diabetes and adherence to dietary
5 regimens. *Psychology & Health*, 28, 139-153. doi: 10.1080/08870446.2012.685740
- 6 Henderson, C. J., Hagger, M. S., & Orbell, S. (2007). Does priming a specific illness schema
7 result in an attentional information-processing bias for specific illnesses? *Health*
8 *Psychology*, 26, 165-173. doi: 10.1037/0278-6133.26.2.165
- 9 Henderson, C. J., Orbell, S., & Hagger, M. S. (2009). Illness schema activation and
10 attentional bias to coping procedures. *Health Psychology*, 28, 101-107. doi:
11 10.1037/a0013690
- 12 *Hermele, S., Olivo, E. L., Namerow, P., & Oz, M. C. (2007). Illness representations and
13 psychological distress in patients undergoing coronary artery bypass graft surgery.
14 *Psychology, Health & Medicine*, 12, 580-591. doi: 10.1080/13548500601162705
- 15 *Heyhoe, J., & Lawton, R. (2009). Distress in patients with interstitial cystitis: Do illness
16 representations have a role to play? *Psychology, Health & Medicine*, 14, 726-739. doi:
17 10.1080/13548500903431477
- 18 Higgins, J. P. T., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis.
19 *Statistics in Medicine*, 21, 1539-1558. doi: 10.1002/sim.1186
- 20 Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring
21 inconsistency in meta-analyses. *British Medical Journal*, 327, 557-560. doi:
22 10.1136/bmj.327.7414.557
- 23 *Hill, S., Dziedzic, K., Thomas, E., Baker, S. R., & Croft, P. (2007). The illness perceptions
24 associated with health and behavioural outcomes in people with musculoskeletal hand

- 1 problems: findings from the North Staffordshire Osteoarthritis Project (NorStOP).
2 *Rheumatology*, 46, 944-951. doi: 10.1093/rheumatology/kem015
- 3 Horne, R. (1997). Representations of medication and treatment: Advances in theory and
4 measurement. In K. J. Petrie & J. A. Weinman (Eds.), *Perceptions of health and*
5 *Illness*. London: Harwood Academic Publishers.
- 6 *Horne, R., Cooper, V., Fisher, M., Buick, D., & Weinman, J. (2001, September). *Predicting*
7 *acceptance of highly active retroviral treatment (HAART): The utility of an extended*
8 *self-regulatory model*. Paper presented at the European Health Psychology
9 Society/British Psychology Society Division of Health Psychology Annual
10 Conference, St. Andrews, Scotland.
- 11 *Horne, R., & Weinman, J. (2002). Self-regulation and self-management in asthma:
12 Exploring the role of illness perceptions and treatment beliefs in explaining non-
13 adherence to preventer medication. *Psychology & Health*, 17, 17-32. doi:
14 10.1080/08870440290001502
- 15 Horne, R., Weinman, J., & Hankins, M. (1999). The beliefs about medicines questionnaire:
16 The development and evaluation of a new method for assessing the cognitive
17 representation of medication. *Psychology and Health*, 14, 1-24. doi:
18 10.1080/08870449908407311
- 19 *Hou, R., Moss-Morris, R., Peveler, R., Mogg, K., Bradley, B. P., & Belli, A. (2012). When
20 a minor head injury results in enduring symptoms: a prospective investigation of risk
21 factors for postconcussional syndrome after mild traumatic brain injury. *Journal of*
22 *Neurology, Neurosurgery & Psychiatry*, 83, 217-223. doi: 10.1136/jnnp-2011-300767
- 23 *Howard, C., Hallas, C. N., Wray, J., & Carby, M. (2009). The relationship between illness
24 perceptions and panic in chronic obstructive pulmonary disease. *Behaviour Research*
25 *and Therapy*, 47, 71-76. doi: 10.1016/j.brat.2008.10.004

- 1 Huang, B., Sivaganesan, S., Succop, P., & Goodman, E. (2004). Statistical assessment of
2 mediational effects for logistic mediational models. *Statistics in Medicine*, 23, 2713-
3 2728. doi: 10.1002/sim.1847
- 4 *Hudson, J. (2013). *Do diabetes cognitions and poor emotional health predict adherence to*
5 *diabetes self-care? A longitudinal test of the common sense self-regulation model*
6 *using structural equation modelling*. Unpublished PhD thesis, University of
7 Manchester, Manchester, UK.
- 8 Hudson, J. L., Bundy, C., Coventry, P. A., & Dickens, C. (2014). Exploring the relationship
9 between cognitive illness representations and poor emotional health and their
10 combined association with diabetes self-care. A systematic review with meta-analysis.
11 *Journal of Psychosomatic Research*, 76, 265-274. doi:
12 10.1016/j.jpsychores.2014.02.004
- 13 Huedo-Medina, T. B., Sánchez-Meca, J., Marín-Martínez, F., & Botella, J. (2006). Assessing
14 heterogeneity in meta-analysis: Q Statistic or I^2 Index? *Psychological Methods*, 11,
15 193-206. doi: 10.1037/1082-989X.11.2.193
- 16 Hunter, J. E., & Schmidt, F. L. (2000). Fixed effects vs. random effects meta-analysis
17 models: Implications for cumulative research knowledge in psychology. *International*
18 *Journal of Selection and Assessment*, 8, 275-292. doi: 10.1111/1468-2389.00156
- 19 Hunter, J. E., & Schmidt, F. L. (2004). *Methods of meta-analysis: Correcting error and bias*
20 *in research findings* (2nd ed.). Newbury Park, CA: Sage.
- 21 Hunter, M. S., & O'Dea, I. (1999). An evaluation of a health education intervention for mid-
22 aged women: five year follow-up of effects upon knowledge, impact of menopause
23 and health. *Patient Education and Counseling*, 38, 249-255. doi: 10.1016/S0738-
24 3991(98)00143-8

- 1 *Husain, M. O., Dearman, S. P., Chaudhry, I. B., Rizvi, N., & Waheed, W. (2008). The
2 relationship between anxiety, depression and illness perception in tuberculosis patients
3 in Pakistan. *Clinical Practice and Epidemiology in Mental Health*, 4, 4. doi:
4 10.1186/1745-0179-4-4
- 5 Husebø, A. M. L., Dyrstad, S. M., Søreide, J. A., & Bru, E. (2012). Predicting exercise
6 adherence in cancer patients and survivors: A systematic review and meta-analysis of
7 motivational and behavioural factors. *Journal of Clinical Nursing*, 22, 4-21. doi:
8 10.1111/j.1365-2702.2012.04322.x
- 9 *Hyphantis, T., Kotsis, K., Tsifetaki, N., Creed, F., Drosos, A. A., Carvalho, A. F., &
10 Voulgari, P. V. (2013). The relationship between depressive symptoms, illness
11 perceptions and quality of life in ankylosing spondylitis in comparison to rheumatoid
12 arthritis. *Clinical Rheumatology*, 32, 635-644. doi: 10.1007/s10067-012-2162-6
- 13 *Ireland, J. A., Reid, M., Powell, R., & Petrie, K. J. (2005). The role of illness perceptions:
14 psychological distress and treatment-seeking delay in patients with genital warts.
15 *International Journal of STD & AIDS*, 16, 667-670. doi:
16 10.1258/095646205774357334
- 17 Jack, K., McLean, S. M., Moffett, J. K., & Gardiner, E. (2010). Barriers to treatment
18 adherence in physiotherapy outpatient clinics: A systematic review. *Manual Therapy*,
19 15, 220-228. doi: 0.1016/j.math.2009.12.004
- 20 *Jackson, J., Fiddler, M., Kapur, N., Wells, A., Tomenson, B., & Creed, F. (2006). Number
21 of bodily symptoms predicts outcome more accurately than health anxiety in patients
22 attending neurology, cardiology, and gastroenterology clinics. *Journal of*
23 *Psychosomatic Research*, 60, 357-363. doi: 10.1016/j.jpsychores.2006.02.006

- 1 Janse, A. J., Sinnema, G., Uiterwaal, C. S. P. M., Kimpfen, J. L. L., & Gemke, R. J. B. J.
2 (2005). Quality of life in chronic illness: Perceptions of parents and paediatricians.
3 *Archives of Disease in Childhood*, 90, 486-491. doi: 10.1136/adc.2004.051722
- 4 *Jansen, D. L., Grootendorst, D. C., Rijken, M., Heijmans, M., Kaptein, A. A., Boeschoten,
5 E. W., & Dekker, F. W. (2010). Pre-dialysis patients' perceived autonomy, self-
6 esteem and labor participation: associations with illness perceptions and treatment
7 perceptions. A cross-sectional study. *BMC Nephrology*, 11, 35. doi: 10.1186/1471-
8 2369-11-35
- 9 *Jessop, D. C., & Rutter, D. R. (2003). Adherence to asthma medication: The role of illness
10 representations. *Psychology & Health*, 18, 595-612. doi:
11 10.1080/0887044031000097009
- 12 Johnson, B. T., & Eagly, A. H. (2014). Meta-analysis of social-personality psychological
13 research. In H. T. Reis & C. M. Judd (Eds.), *Handbook of research methods in social
14 and personality psychology* (2nd ed., pp. 675-707). London, UK: Cambridge
15 University Press.
- 16 Johnson, B. T., Low, R. E., & MacDonald, H. V. (2014). Panning for the gold in health
17 research: Incorporating studies' methodological quality in meta-analysis. *Psychology
18 & Health*, 30, 135-152. doi: 10.1080/08870446.2014.953533
- 19 *Jonsbu, E., Martinsen, E. W., Morken, G., Moum, T., & Dammen, T. (2012). Illness
20 perception among patients with chest pain and palpitations before and after negative
21 cardiac evaluation. *BioPsychoSocial Medicine*, 6, 19. doi: 10.1186/1751-0759-6-19
- 22 *Jonsbu, E., Martinsen, E. W., Morken, G., Moum, T., & Dammen, T. (2013). Change and
23 impact of illness perceptions among patients with non-cardiac chest pain or benign
24 palpitations following three sessions of CBT. *Behavioural and Cognitive
25 Psychotherapy*, 41, 398-407. doi: 10.1017/S1352465813000179

- 1 *Jopson, N. M., & Moss-Morris, R. (2003). The role of illness severity and illness
2 representations in adjusting to multiple sclerosis. *Journal of Psychosomatic Research*,
3 54, 503-511. doi: 10.1016/S0022-3999(02)00455-5
- 4 *Juergens, M. C., Seekatz, B., Moosdorf, R. G., Petrie, K. J., & Rief, W. (2010). Illness
5 beliefs before cardiac surgery predict disability, quality of life, and depression 3
6 months later. *Journal of Psychosomatic Research*, 68, 553-560. doi:
7 10.1016/j.jpsychores.2009.10.004
- 8 *Kaptein, A. A., Helder, D. I., Scharloo, M., Van Kempen, G. M. J., Weinman, J., Van
9 Houwelingen, H. J. C., & Roos, R. A. C. (2006). Illness perceptions and coping
10 explain well-being in patients with Huntington's disease. *Psychology & Health*, 21,
11 431-446. doi: 10.1080/14768320500456947
- 12 *Kaptein, A. A., Scharloo, M., Helder, D. I., Snoei, L., van Kempen, G. M. J., Weinman, J., .
13 . . Roos, R. A. C. (2007). Quality of life in couples living with Huntington's disease:
14 the role of patients' and partners' illness perceptions. *Quality of Life Research*, 16,
15 793-801. doi: 10.1007/s11136-007-9194-4
- 16 *Karademas, E. C. (2012). Conditional indirect relations of cardiac patients' subjective health
17 to optimism through illness representations: A self-regulation circuit. *Journal of*
18 *Health Psychology*, 17, 36-45. doi: 10.1177/1359105311405554
- 19 Karademas, E. C., Bati, A., Karkania, V., Georgiou, V., & Sofokleous, S. (2013). The
20 association between Pandemic Influenza A (H1N1) public perceptions and reactions:
21 A prospective study. *Journal of Health Psychology*, 18, 419-428. doi:
22 10.1177/1359105312436765
- 23 *Karademas, E. C., & Giannousi, Z. (2013). Representations of control and psychological
24 symptoms in couples dealing with cancer: A dyadic-regulation approach. *Psychology*
25 *& Health*, 28, 67-83. doi: 10.1080/08870446.2012.713954

- 1 *Karademas, E. C., Karamvakalis, N., & Zarogiannos, A. (2009). Life context and the
2 experience of chronic illness: Is the stress of life associated with illness perceptions
3 and coping? *Stress & Health*, 25, 405–412. doi: 10.1002/smi.1252
- 4 *Karademas, E. C., Kynigopoulou, E., Aghathangelou, E., & Anestis, D. (2011). The relation
5 of illness representations to the ‘end-stage’ appraisal of outcomes through health
6 status, and the moderating role of optimism. *Psychology & Health*, 26, 567-583. doi:
7 10.1080/08870441003653488
- 8 *Karademas, E. C., Tsalikou, C., & Tallarou, M.-C. (2011). The Impact of Emotion
9 Regulation and Illness-focused Coping Strategies on the Relation of Illness-related
10 Negative Emotions to Subjective Health. *Journal of Health Psychology*, 16, 510-519.
11 doi: 10.1177/1359105310392093
- 12 *Keeling, M., Bambrough, J., & Simpson, J. (2013). Depression, anxiety and positive affect
13 in people diagnosed with low-grade tumours: the role of illness perceptions. *Psycho-*
14 *Oncology*, 22, 1421-1427. doi: 10.1002/pon.3158
- 15 *Kemp, S., Feely, M., Hay, A., Wild, H., & Cooper, C. (2007). Psychological factors and use
16 of antiepileptic drugs: Pilot work using an objective measure of adherence.
17 *Psychology, Health & Medicine*, 12, 107-113. doi: 10.1080/14622200500358937
- 18 *Kemp, S., Morley, S., & Anderson, E. (1999). Coping with epilepsy: Do illness
19 representations play a role? *British Journal of Clinical Psychology*, 38, 43-58. doi:
20 10.1348/014466599162656
- 21 *Kesler, A., Kliper, E., Goner-Shilo, D., & Benyamini, Y. (2009). Illness perceptions and
22 quality of life amongst women with pseudotumor cerebri. *European Journal of*
23 *Neurology*, 16, 931-936. doi: 10.1111/j.1468-1331.2009.02625.x
- 24 *Khunti, K., Skinner, T. C., Heller, S., Carey, M. E., Dallosso, H. M., & Davies, M. J.
25 (2008). Biomedical, lifestyle and psychosocial characteristics of people newly

1 diagnosed with Type 2 diabetes: baseline data from the DESMOND randomized
2 controlled trial. *Diabetic Medicine*, 25, 1454-1461. doi: 10.1111/j.1464-
3 5491.2008.02620.x

4 *Kiebles, J. L., Doerfler, B., & Keefer, L. (2010). Preliminary evidence supporting a
5 framework of psychological adjustment to inflammatory bowel disease†.
6 *Inflammatory Bowel Diseases*, 16, 1685-1695. doi: 10.1002/ibd.21215

7 *Kim, Y., & Evangelista, L. S. (2010). Relationship between illness perceptions, treatment
8 adherence, and clinical outcomes in patients on maintenance hemodialysis.
9 *Nephrology Nursing Journal*, 37, 271-281.

10 *Kim, Y., Evangelista, L. S., Phillips, L. R., Pavlish, C., & Kopple, J. D. (2012).
11 Racial/Ethnic differences in illness perceptions in minority patients undergoing
12 maintenance hemodialysis. *Nephrology Nursing Journal*, 39, 39-48.

13 *Kirby, B., Richards, H. L., Woo, P., Hindle, E., Main, C. J., & Griffiths, C. E. M. (2001).
14 Physical and psychologic measures are necessary to assess overall psoriasis severity.
15 *Journal of the American Academy of Dermatology*, 45, 72-76. doi:
16 10.1067/mjd.2001.114592

17 *Kirby, S. E., & Yardley, L. (2009). Cognitions associated with anxiety in Ménière's disease.
18 *Journal of Psychosomatic Research*, 66, 111-118. doi:
19 10.1016/j.jpsychores.2008.05.027

20 *Knibb, R. C., & Horton, S. L. (2008). Can illness perceptions and coping predict
21 psychological distress amongst allergy sufferers? *British Journal of Health*
22 *Psychology*, 13, 103-119. doi: 10.1348/135910706x173278

23 Köhler, T., Cortina, J. M., Kurtessis, J. N., & Gözl, M. (2015). Are we correcting correctly?
24 Interdependence of reliabilities in meta-analysis. *Organizational Research Methods*,
25 18, 355-428. doi: 10.1177/1094428114563617

- *Kotsis, K., Voulgari, P. V., Tsifetaki, N., Machado, M. O., Carvalho, A. F., Creed, F., . . .
Hyphantis, T. (2012). Anxiety and depressive symptoms and illness perceptions in
psoriatic arthritis and associations with physical health-related quality of life. *Arthritis
Care & Research*, 64, 1593-1601. doi: 10.1002/acr.21725
- *Lane, D. A., Langman, C. M., Lip, G. Y. H., & Nouwen, A. (2009). Illness perceptions,
affective response, and health-related quality of life in patients with atrial fibrillation.
Journal of Psychosomatic Research, 66, 203-210. doi:
10.1016/j.jpsychores.2008.10.007
- *Lau-Walker, M. (2004). Relationship between illness representation and self-efficacy.
Journal of Advanced Nursing, 48, 216-225. doi: 10.1111/j.1365-2648.2004.03190.x
- *Lau-Walker, M. (2006). Predicting self-efficacy using illness perception components: A
patient survey. *British Journal of Health Psychology*, 11, 643-661. doi:
10.1348/135910705X72802
- *Lau-Walker, M. (2007). Importance of illness beliefs and self-efficacy for patients with
coronary heart disease. *Journal of Advanced Nursing*, 60, 187-198. doi:
10.1111/j.1365-2648.2007.04398.x
- *Lau-Walker, M. O., Cowie, M. R., & Roughton, M. (2009). Coronary heart disease patients'
perception of their symptoms and sense of control are associated with their quality of
life three years following hospital discharge. *Journal of Clinical Nursing*, 18, 63-71.
doi: 10.1111/j.1365-2702.2008.02386.x
- Lau, R. R., Bernard, T. M., & Hartman, K. A. (1989). Further explorations of commonsense
representations of common illnesses. *Health Psychology*, 8, 195-219. doi:
10.1037/0278-6133.8.2.195
- Lau, R. R., & Hartman, K. A. (1983). Common sense representations of common illnesses.
Health Psychology, 2, 167-185. doi: 10.1037/0278-6133.2.2.167

- 1 Law, G. U., Tolgyesi, C. S., & Howard, R. A. (2012). Illness beliefs and self-management in
2 children and young people with chronic illness: A systematic review. *Health*
3 *Psychology Review*, 8, 362-380. doi: 10.1080/17437199.2012.747123
- 4 *Law, G. U., Walsh, J., Queral, V., & Nouwen, A. (2013). Adolescent and parent diabetes
5 distress in type 1 diabetes: The role of self-efficacy, perceived consequences, family
6 responsibility and adolescent–parent discrepancies. *Journal of Psychosomatic*
7 *Research*, 74, 334-339. doi: 10.1016/j.jpsychores.2012.12.009
- 8 *Lawson, V. L., Bundy, C., Belcher, J., & Harvey, J. N. (2010). Mediation by illness
9 perceptions of the effect of personality and health threat communication on coping
10 with the diagnosis of diabetes. *British Journal of Health Psychology*, 15, 623-642.
11 doi: 10.1348/135910709X478664
- 12 *Lawson, V. L., Bundy, C., Lyne, P. A., & Harvey, J. N. (2004). Using the IPQ and PMDI to
13 predict regular diabetes care-seeking among patients with Type 1 diabetes. *British*
14 *Journal of Health Psychology*, 9, 241-252. doi: 10.1348/135910704773891078
- 15 *Lawson, V. L., Lyne, P. A., Bundy, C., & Harvey, J. N. (2007). The role of illness
16 perceptions, coping and evaluation in care-seeking among people with type 1
17 diabetes. *Psychology & Health*, 22, 175-191. doi: 10.1080/14768320600774546
- 18 Lazarus, R. S., & Folkman, S. (1984). Coping and adaptation. In W. D. Gentry (Ed.),
19 *Handbook of behavioral medicine* (pp. 282-325). New York: Guildford Press.
- 20 *Lee, B.-O., Chaboyer, W., & Wallis, M. (2008). Predictors of health-related quality of life 3
21 months after traumatic injury. *Journal of Nursing Scholarship*, 40, 83-90. doi:
22 10.1111/j.1547-5069.2008.00210.x
- 23 *Lehto, R. H., & Cimprich, B. (2009). Worry and the formation of cognitive representations
24 of illness in individuals undergoing surgery for suspected lung Ccancer. *Cancer*
25 *Nursing*, 32, 2-10. doi: 10.1097/01.NCC.0000343363.75752.f1

- *Leung, Y. W., Gravely-Witte, S., Macpherson, A., Irvine, J., Stewart, D. E., & Grace, S. L. (2010). Post-traumatic growth among cardiac outpatients. *Journal of Health Psychology, 15*, 1049-1063. doi: 10.1177/1359105309360577
- Leventhal, H., Benyamini, Y., Brownlee, S., Diefenbach, M. A., Leventhal, E. A., Patrick-Miller, L., & Robitaille, C. (1997). Illness representations: Theoretical foundations. In K. J. Petrie & J. Weinman (Eds.), *Perceptions of health and illness: Current research and applications* (pp. 19-45). Amsterdam: Harwood Academic Publishers.
- Leventhal, H., Brissette, I., & Leventhal, E. A. (2003). The common-sense model of self-regulation of health and illness. In L. D. Cameron & H. Leventhal (Eds.), *The self-regulation of health and illness behaviour* (pp. 42-65). London, UK: Routledge.
- Leventhal, H., & Cleary, P. D. (1980). The smoking problem - A review of the research and theory in behavioral risk modification. *Psychological Bulletin, 88*, 370-405. doi: 10.1037/0033-2909.88.2.370
- Leventhal, H., Diefenbach, M. A., & Leventhal, E. A. (1992). Illness cognition: Using common sense to understand treatment adherence and affect cognition interactions. *Cognitive Therapy and Research, 16*, 143-163. doi: 10.1007/BF01173486
- Leventhal, H., Leventhal, E., & Contrada, R. J. (1998). Self-regulation, health, and behavior: A perceptual-cognitive approach. *Psychology and Health, 13*, 717-734. doi: 10.1080/08870449808407425
- Leventhal, H., Leventhal, E. A., & Breland, J. Y. (2011). Cognitive science speaks to the "common sense" of chronic illness management. *Annals of Behavioral Medicine, 41*, 152-163. doi: 10.1007/s12160-010-9246-9
- Leventhal, H., Meyer, D., & Nerenz, D. (1980). The common sense model of illness danger. In S. Rachman (Ed.), *Medical Psychology* (Vol. II, pp. 7-30). New York: Pergamon Press.

- 1 Leventhal, H., Phillips, L. A., & Burns, E. (2016). The common-sense model of self-
2 regulation (CSM): A dynamic framework for understanding illness self-management.
3 *Journal of Behavioral Medicine* 1-12. doi: 10.1007/s10865-016-9782-2
- 4 Limburg, K., Watson, H. J., Hagger, M. S., & Egan, S. J. (2016). The relationship between
5 perfectionism and psychopathology: A meta-analysis. *Journal of Clinical Psychology*.
6 doi: 10.1002/jclp.22435
- 7 Linz, D., Penrod, S., & Leventhal, H. (1982, August). *Cognitive organisation of disease*
8 *among laypersons*. Paper presented at the 20th International Congress of Applied
9 Psychology, Edinburgh, Scotland.
- 10 *Ljosaa, T. M., Mork, C., Stubhaug, A., Moum, T., & Wahl, A. K. (2012). Skin pain and skin
11 discomfort is associated with quality of life in patients with psoriasis. *Journal of the*
12 *European Academy of Dermatology and Venereology*, 26, 29-35. doi: 10.1111/j.1468-
13 3083.2011.04000.x
- 14 *Llewellyn, C. D., McGurk, M., & Weinman, J. (2006). Head and neck cancer: To what
15 extent can psychological factors explain differences between health-related quality of
16 life and individual quality of life? *British Journal of Oral and Maxillofacial Surgery*,
17 44, 351-357. doi: 10.1016/j.bjoms.2005.06.033
- 18 *Llewellyn, C. D., McGurk, M., & Weinman, J. (2007a). Illness and treatment beliefs in head
19 and neck cancer: Is Leventhal's common sense model a useful framework for
20 determining changes in outcomes over time? *Journal of Psychosomatic Research*, 63,
21 17-26. doi: 10.1016/j.jpsychores.2007.01.013
- 22 *Llewellyn, C. D., McGurk, M., & Weinman, J. (2007b). The relationship between the
23 Patient Generated Index (PGI) and measures of HR-QoL following diagnosis with
24 head and neck cancer: Are illness and treatment perceptions determinants of

1 judgment-based outcomes? *British Journal of Health Psychology*, 12, 421-437. doi:
2 10.1348/135910706X118422

3 *Llewellyn, C. D., Miners, A. H., Lee, C. A., Harrington, C., & Weinman, J. (2003). The
4 illness perceptions and treatment beliefs of individuals with severe haemophilia and
5 their role in adherence to home treatment. *Psychology & Health*, 18, 185-200. doi:
6 10.1080/0887044031000098198

7 *Llewellyn, C. D., Weinman, J., McGurk, M., & Humphris, G. (2008). Can we predict which
8 head and neck cancer survivors develop fears of recurrence? *Journal of*
9 *Psychosomatic Research*, 65, 525-532. doi: 10.1016/j.jpsychores.2008.03.014

10 Lobban, F., Barrowclough, C., & Jones, S. (2005). Assessing cognitive representations of
11 mental health problems. I. The illness perception questionnaire for schizophrenia.
12 *British Journal of Clinical Psychology*, 44, 147-162. doi: 10.1348/014466504X19497

13 *Luyckx, K., Vanhalst, J., I., S.-K., & Weets, I. (2010). A typology of coping with Type 1
14 diabetes in emerging adulthood: associations with demographic, psychological, and
15 clinical parameters. *Journal of Behavioral Medicine*, 33, 228-238. doi:
16 10.1007/s10865-010-9249-9

17 *MacInnes, J. (2013). Relationships between illness representations, treatment beliefs and the
18 performance of self-care in heart failure: a cross-sectional survey. *European Journal*
19 *of Cardiovascular Nursing*, 12, 536-543. doi: 10.1177/1474515112473872

20 *Marke, V., & Bennett, P. (2013). Predicting post-traumatic stress disorder following first
21 onset acute coronary syndrome: Testing a theoretical model. *British Journal of*
22 *Clinical Psychology*, 52, 70-81. doi: 10.1111/bjc.12001

23 *Marques, M., De Gucht, V., Leal, I., & Maes, S. (2013). A cross-cultural perspective on
24 psychological determinants of chronic fatigue syndrome: A comparison between a

- 1 Portuguese and a Dutch patient sample. *International Journal of Behavioral*
2 *Medicine*, 20, 229-238. doi: 10.1007/s12529-012-9265-y
- 3 Martin, R., Rothrock, N., Leventhal, H., & Leventhal, E. (2003). Common sense models of
4 illness: Implications for symptom perception and health-related behaviors. In J. Suls
5 & K. A. Wallston (Eds.), *Social Psychological Foundations of Health and Illness* (pp.
6 199-225): Blackwell Publishing Ltd.
- 7 Mc Sharry, J., Moss-Morris, R., & Kendrick, T. (2011). Illness perceptions and glycaemic
8 control in diabetes: A systematic review with meta-analysis. *Diabetic Medicine*, 28,
9 1300-1310. doi: 10.1111/j.1464-5491.2011.03298.x
- 10 *McCabe, P. J., & Barnason, S. A. (2012). Illness perceptions, coping strategies, and
11 symptoms contribute to psychological distress in patients with recurrent symptomatic
12 atrial fibrillation. *Journal of Cardiovascular Nursing*, 27, 431-444. doi:
13 10.1097/JCN.0b013e31821e7ab1
- 14 *McCabe, P. J., Barnason, S. A., & Houfek, J. (2011). Illness beliefs in patients with
15 recurrent symptomatic atrial fibrillation. *Pacing and Clinical Electrophysiology*, 34,
16 810-820. doi: 10.1111/j.1540-8159.2011.03105.x
- 17 McCarthy, S. C., Lyons, A. C., Weinman, J., Talbot, R., & Purnell, D. (2003). Do
18 expectations influence recovery from oral surgery? An illness representation
19 approach. *Psychology & Health*, 18, 109-126. doi: 10.1080/0887044031000080674
- 20 *McGrady, M. E., Cotton, S., Rosenthal, S. L., Roberts, Y. H., Britto, M., & Yi, M. S.
21 (2010). Anxiety and asthma symptoms in urban adolescents with asthma: The
22 mediating role of illness perceptions. *Journal of Clinical Psychology in Medical*
23 *Settings*, 17, 349-356. doi: 10.1007/s10880-010-9214-3

- 1 *Mickevičienė, A., Vanagas, G., Jievaltas, M., & Ulys, A. (2013). Does illness perception
2 explain quality of life of patients with prostate cancer? *Medicina-Lithuania*, 49, 235-
3 241.
- 4 *Miglioretti, M., Mazzini, L., Oggioni, G. D., Testa, L., & Monaco, F. (2008). Illness
5 perceptions, mood and health-related quality of life in patients with amyotrophic
6 lateral sclerosis. *Journal of Psychosomatic Research*, 65, 603-609. doi:
7 10.1016/j.jpsychores.2008.05.012
- 8 *Molloy, G. J., Gao, C., Johnston, D. W., Johnston, M., Witham, M. D., Struthers, A. D., &
9 McMurdo, M. E. T. (2009). Adherence to angiotensin-converting-enzyme inhibitors
10 and illness beliefs in older heart failure patients. *European Journal of Heart Failure*,
11 11, 715-720. doi: 10.1093/eurjhf/hfp059
- 12 Moreno, S., Sutton, A., Ades, A., Stanley, T., Abrams, K., Peters, J., & Cooper, N. (2009).
13 Assessment of regression-based methods to adjust for publication bias through a
14 comprehensive simulation study. *BMC Medical Research Methodology*, 9, 2. doi:
15 10.1186/1471-2288-9-2
- 16 *Morgan, K., Thompson, J., Dixon, S., Tomeny, M., & Mathers, N. (2003). Predicting
17 longer-term outcomes following psychological treatment for hypnotic-dependent
18 chronic insomnia. *Journal of Psychosomatic Research*, 54, 21-29. doi:
19 10.1016/S0022-3999(02)00569-X
- 20 Morris, A., & Ogden, J. (2012). Making sense of children's medically unexplained
21 symptoms: Managing ambiguity, authenticity and responsibility. *Psychology, Health*
22 *& Medicine*, 17, 285-294. doi: 10.1080/13548506.2011.608804
- 23 *Moss-Morris, R., Humphrey, K., Johnson, M. H., & Petrie, K. J. (2007). Patients'
24 perceptions of their pain condition across a multidisciplinary pain management

- 1 program: Do they change and if so does it matter? *The Clinical Journal of Pain*, 23,
2 558-564. doi: 10.1097/AJP.0b013e318093fcab
- 3 *Moss-Morris, R., Petrie, K., & Weinman, J. (1996). Functioning in chronic fatigue
4 syndrome: Do illness perceptions play a regulatory role. *British Journal of Health*
5 *Psychology*, 1, 15-25. doi: 10.1111/j.2044-8287.1996.tb00488.x
- 6 *Moss-Morris, R., Spence, M. J., & Hou, R. (2011). The pathway from glandular fever to
7 chronic fatigue syndrome: Can the cognitive behavioural model provide the map?
8 *Psychological Medicine*, 41, 1099-1107. doi: 10.1017/s003329171000139x
- 9 *Moss-Morris, R., Weinman, J., Petrie, K. J., Horne, R., Cameron, L. D., & Buick, L. (2002).
10 The revised illness perception questionnaire (IPQ-R). *Psychology and Health*, 17, 1-
11 16. doi: 10.1080/08870440290001494
- 12 *Mulligan, K., Mehta, P. A., Fteropoulli, T., Dubrey, S. W., McIntyre, H. F., McDonagh, T.
13 A., . . . Newman, S. (2012). Newly diagnosed heart failure: Change in quality of life,
14 mood, and illness beliefs in the first 6 months after diagnosis. *British Journal of*
15 *Health Psychology*, 17, 447-462. doi: 10.1111/j.2044-8287.2011.02047.x
- 16 *Murphy, H., Dickens, C., Creed, F., & Bernstein, R. (1999). Depression, illness perception
17 and coping in rheumatoid arthritis. *Journal of Psychosomatic Research*, 46, 155-164.
18 doi: 10.1016/S0022-3999(98)00073-7
- 19 Muthén, L. K., & Muthén, B. O. (2012). *MPlus statistical analysis with latent variables:*
20 *User's guide* (7th ed.). Los Angeles, CA: Muthén & Muthén.
- 21 *Naab, F., Brown, R., & Heidrich, S. (2013). Psychosocial health of infertile Ghanaian
22 women and their infertility beliefs. *Journal of Nursing Scholarship*, 45, 132-140. doi:
23 10.1111/jnu.12013
- 24 National Institutes of Health. (2014). Quality assessment tool for observational cohort and
25 cross-sectional studies. Retrieved November 14, 2016, from

1 [https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-](https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools/cohort)
2 [reduction/tools/cohort](https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools/cohort)

3 *Neter, E., Litvak, A., & Miller, A. (2009). Goal disengagement and goal re-engagement
4 among multiple sclerosis patients: Relationship to well-being and illness
5 representation. *Psychology & Health*, 24, 175-186. doi: 10.1080/08870440701668665

6 *Nicklas, L. B., Dunbar, M., & Wild, M. (2010). Adherence to pharmacological treatment of
7 non-malignant chronic pain: The role of illness perceptions and medication beliefs.
8 *Psychology & Health*, 25, 601-615. doi: 10.1080/08870440902783610

9 *Nouwen, A., Urquhart L., G., Hussain, S., McGovern, S., & Napier, H. (2009). Comparison
10 of the role of self-efficacy and illness representations in relation to dietary self-care
11 and diabetes distress in adolescents with type 1 diabetes. *Psychology & Health*, 24,
12 1071-1084. doi: 10.1080/08870440802254597

13 *O'Connor, S. M., Jardine, A. G., & Millar, K. (2008). The prediction of self-care behaviors
14 in end-stage renal disease patients using Leventhal's Self-Regulatory Model. *Journal*
15 *of Psychosomatic Research*, 65, 191-200. doi: 10.1016/j.jpsychores.2008.02.008

16 *O'Leary, C. J., Creamer, D., Higgins, E., & Weinman, J. (2004). Perceived stress, stress
17 attributions and psychological distress in psoriasis. *Journal of Psychosomatic*
18 *Research*, 57, 465-471. doi: 10.1016/j.jpsychores.2004.03.012

19 *O'Carroll, R. E., Whittaker, J., Hamilton, B., Johnston, M., Sudlow, C., & Dennis, M.
20 (2011). Predictors of adherence to secondary preventive medication in stroke patients.
21 *Annals of Behavioral Medicine*, 41, 383-390. doi: 10.1007/s12160-010-9257-6

22 *Olchowska-Kotala, A. (2013). Illness representations in individuals with rheumatoid
23 arthritis and the willingness to undergo acupuncture treatment. *European Journal of*
24 *Integrative Medicine*, 5, 347-351. doi: 10.1016/j.eujim.2013.03.008

- 1 *Olchowska-Kotala, A., & Szmyrka-Kaczmarek, M. (2010). Illness representation and self-
2 assessment of disability level in patients suffering with rheumatoid arthritis. *Advances*
3 *in Clinical and Experimental Medicine*, 19, 503-512.
- 4 Oluka, O. C., Nie, S., & Sun, Y. (2014). Quality assessment of TPB-based questionnaires: A
5 systematic review. *PLoS ONE*, 9, e94419. doi: 10.1371/journal.pone.0094419
- 6 *Orbell, S., Hagger, M. S., Brown, V., & Tidy, J. (2006). Comparing two theories of health
7 behavior: A prospective study of non-completion of treatment following cervical
8 cancer screening. *Health Psychology*, 25, 604-615. doi: 10.1037/0278-6133.25.5.604
- 9 Orbell, S., Henderson, C. J., & Hagger, M. S. (2015). Illness schema activation and the
10 effects of illness seasonality on accessibility of implicit illness-related information.
11 *Annals of Behavioral Medicine*, 49, 918-923. doi: 10.1007/s12160-015-9719-y
- 12 Orbell, S., Hodgkins, S., & Sheeran, P. (1997). Implementation intentions and the theory of
13 planned behavior. *Personality and Social Psychology Bulletin*, 23, 945-954. doi:
14 10.1177/0146167297239004
- 15 *Orbell, S., Johnston, M., Rowley, D., Espley, A., & Davey, P. (1998). Cognitive
16 representations of illness and functional and affective adjustment following surgery
17 for osteoarthritis. *Social Science and Medicine*, 47, 93-102. doi: 10.1016/s0277-
18 9536(97)10132-0
- 19 *Orbell, S., O'Sullivan, I., Parker, R., Steele, B., Campbell, C., & Weller, D. (2008). Illness
20 representations and coping following an abnormal colorectal cancer screening result.
21 *Social Science & Medicine*, 67, 1465-1474. doi: 10.1016/j.socscimed.2008.06.039
- 22 *Paddison, C. A. M., Alpass, F. M., & Stephens, C. V. (2008). Psychological factors account
23 for variation in metabolic control and perceived quality of life among people with
24 type 2 diabetes in New Zealand. *International Journal of Behavioral Medicine*, 15,
25 180-186. doi: 10.1080/10705500802222295

- 1 *Paschalides, C., Wearden, A. J., Dunkerley, R., Bundy, C., Davies, R., & Dickens, C. M.
2 The associations of anxiety, depression and personal illness representations with
3 glycaemic control and health-related quality of life in patients with type 2 diabetes
4 mellitus. *Journal of Psychosomatic Research*, 57, 557-564. doi:
5 10.1016/j.jpsychores.2004.03.006
- 6 *Paterson, J., Moss-Morris, R., & Butler, S. J. (1999). The effect of illness experience and
7 demographic factors on children's illness representations. *Psychology and Health*, 14,
8 117-129. doi: 10.1080/08870449908407318
- 9 Peters, G.-J. Y., Ruiter, R. A. C., & Kok, G. (2013). Threatening communication: A critical
10 re-analysis and a revised meta-analytic test of fear appeal theory. *Health Psychology*
11 *Review*, 7, S8-S31. doi: 10.1080/17437199.2012.703527
- 12 *Petriček, G., Vrcić-Keglević, M., Vuletić, G., Cerovečki, V., Ožvačić, Z., & Murgić, L.
13 (2009). Illness perception and cardiovascular risk factors in patients with type 2
14 diabetes: Cross-sectional questionnaire study. *Croatian Medical Journal*, 50, 583–
15 593. doi: 10.3325/cmj.2009.50.583
- 16 Petrie, K. J., Cameron, L., Ellis, C. J., Buick, D., & Weinman, J. (2002). Changing illness
17 perceptions after myocardial infarction: An early intervention randomized controlled
18 trial. *Psychosomatic Medicine*, 64, 580-586. doi: 10.1097/00006842-200207000-
19 00007
- 20 Petrie, K. J., & Weinman, J. (2012). Patients' perceptions of their illness: The dynamo of
21 volition in health care. *Current Directions in Psychological Science*, 21, 60-65. doi:
22 10.1177/0963721411429456
- 23 *Petrie, K. J., Weinman, J., Sharpe, N., & Buckley, J. (1996). Role of patients' view of their
24 illness in predicting return to work and functioning after myocardial infarction:

1 Longitudinal study. *British Medical Journal*, 312, 1191-1194. doi:

2 10.1136/bmj.312.7040.1191

3 *Philip, E. J., Lindner, H., & Lederman, L. (2009). Relationship of illness perceptions with
4 depression among individuals diagnosed with lupus. *Depression and Anxiety*, 26, 575-
5 582. doi: 10.1002/da.20451

6 *Pinto, P. R., McIntyre, T., Ferrero, R., Almeida, A., & Araújo-Soares, V. (2013). Risk
7 factors for moderate and severe persistent pain in patients undergoing total knee and
8 hip arthroplasty: A prospective predictive study. *PLoS ONE*, 8, e73917. doi:
9 10.1371/journal.pone.0073917

10 *Pinto, P. R., McIntyre, T., Nogueira-Silva, C., Almeida, A., & Araújo-Soares, V. (2012).
11 Risk factors for persistent postsurgical pain in women undergoing hysterectomy due
12 to benign causes: A prospective predictive study. *The Journal of Pain*, 13, 1045-1057.
13 doi: 10.1016/j.jpain.2012.07.014

14 Podsakoff, P. M., MacKenzie, S. B., Lee, J.-Y., & Podsakoff, N. P. (2003). Common method
15 biases in behavioral research: A critical review of the literature and recommended
16 remedies. *Journal of Applied Psychology*, 88, 879-903. doi: 10.1037/0021-
17 9010.88.5.879

18 *Pollak, L., Segal, P., Stryjer, R., & Stern, H. G. (2012). Beliefs and emotional reactions in
19 patients with benign paroxysmal positional vertigo: a longitudinal study. *American*
20 *Journal of Otolaryngology*, 33, 221-225. doi: 10.1016/j.amjoto.2011.07.002

21 *Powell, H., McCaffery, K., Murphy, V. E., Hensley, M., J., Clifton, V. L., Giles, W., &
22 Gibson, P. G. (2013). Psychosocial variables are related to future exacerbation risk
23 and perinatal outcomes in pregnant women with asthma. *Journal of Asthma*, 50, 383-
24 389. doi: 10.3109/02770903.2012.757777

- 1 Preacher, K. J., & Kelley, K. (2011). Effect size measures for mediation models: Quantitative
2 strategies for communicating indirect effects. *Psychological Methods, 16*, 93-115.
3 doi: 10.1037/a0022658
- 4 *Price, A., Goodwin, L., Rayner, L., Shaw, E., Hansford, P., Sykes, N., . . . Lee, W. (2012).
5 Illness perceptions, adjustment to illness, and depression in a palliative care
6 population. *Journal of Pain and Symptom Management, 43*, 819-832. doi:
7 10.1016/j.jpainsymman.2011.05.013
- 8 *Quiles, Y., & Terol, M. C. (2010). Using common sense model in eating disorders. *Journal*
9 *of Health Psychology, 16*, 208-216. doi: 10.1177/1359105310373411
- 10 Rees, G., Fry, A., & Cull, A. (2001). A family history of breast cancer: Women's experiences
11 from a theoretical perspective. *Social Science and Medicine, 52*, 1433-1440. doi:
12 10.1016/S0277-9536(00)00248-3
- 13 *Rees, G., Leong, O., Crowston, J. G., & Lamoureux, E. L. (2010). Intentional and
14 unintentional nonadherence to ocular hypotensive treatment in patients with
15 glaucoma. *Ophthalmology, 117*, 903-908. doi: 10.1016/j.ophtha.2009.10.038
- 16 *Rees, G., Sasongko, M. B., Fenwick, E. K., Nicolaou, T. E., Wong, T. Y., & Lamoureux, E.
17 L. (2012). Impact of diabetic retinopathy on patients' beliefs about diabetes. *Clinical*
18 *and Experimental Optometry, 95*, 371-376. doi: 10.1111/j.1444-0938.2012.00745.x
- 19 *Reges, O., Vilchinsky, N., Leibowitz, M., Khaskia, A., Mosseri, M., & Kark, J. D. (2013).
20 Illness cognition as a predictor of exercise habits and participation in cardiac
21 prevention and rehabilitation programs after acute coronary syndrome. *BMC Public*
22 *Health, 13*, 956. doi: 10.1186/1471-2458-13-956
- 23 *Richards, H. L., Herrick, A. L., Griffin, K., Gwilliam, P. D. H., Loukes, J., & Fortune, D. G.
24 (2003). Systemic sclerosis: Patients' perceptions of their condition. *Arthritis Care &*
25 *Research, 49*, 689-696. doi: 10.1002/art.11385

- *Riley, J. P., Habibi, H., Banya, W., Gatzoulis, M. A., Lau-Walker, M., & Cowie, M. R. (2012). Education and support needs of the older adult with congenital heart disease. *Journal of Advanced Nursing*, 68, 1050-1060. doi: 10.1111/j.1365-2648.2011.05809.x
- *Rochelle, T. L., & Fidler, H. (2012). The importance of illness perceptions, quality of life and psychological status in patients with ulcerative colitis and Crohn's disease. *Journal of Health Psychology*, 18, 972-983. doi: 10.1177/1359105312459094
- Rodin, J., & Salovey, P. (1989). Health psychology. *Annual Review of Psychology*, 40, 533-379. doi: 10.1146/annurev.ps.40.020189.002533
- *Rogan, C., Fortune, D. G., & Prentice, G. (2013). Post-traumatic growth, illness perceptions and coping in people with acquired brain injury. *Neuropsychological Rehabilitation*, 23, 639-657. doi: 10.1080/09602011.2013.799076
- *Rose, M. R., Sadjadi, R., Weinman, J., Akhtar, T., Pandya, S., Kissel, J. T., . . . the Muscle Study Group. (2012). Role of disease severity, illness perceptions, and mood on quality of life in muscle disease. *Muscle & Nerve*, 46, 351-359. doi: 10.1002/mus.23320
- Rosenthal, M. C. (1994). The fugitive literature. In H. Cooper & L. V. Hedges (Eds.), *The Handbook of Research Synthesis* (pp. 85-94). New York: Russell Sage Foundation.
- *Rozema, H., Völlink, T., & Lechner, L. (2009). The role of illness representations in coping and health of patients treated for breast cancer. *Psycho-Oncology*, 18, 849-857. doi: 10.1002/pon.1488
- *Rutter, C. L., & Rutter, D. R. (2001a). *First time clinic attendees' illness representation of IBS*. Unpublished manuscript, University of Kent at Canterbury, Canterbury, UK.
- *Rutter, C. L., & Rutter, D. R. (2001b). *A longitudinal investigation of illness representation, coping and outcome in primary care: Irritable bowel syndrome, inflammatory bowel*

disease and chronic fatigue syndrome patients. Unpublished manuscript, University of Kent at Canterbury, Canterbury, UK.

*Rutter, C. L., & Rutter, D. R. (2001c). *Stability of the illness representation upon joining a self-help organisation*. Unpublished manuscript, University of Kent at Canterbury, Canterbury, UK.

*Rutter, C. L., & Rutter, D. R. (2002). Illness representation, coping and outcome in irritable bowel syndrome (IBS). *British Journal of Health Psychology*, 7, 377-391. doi: 10.1348/135910702320645372

*Rutter, C. L., & Rutter, D. R. (2007). Longitudinal analysis of the illness representation model in patients with irritable bowel syndrome (IBS). *Journal of Health Psychology*, 12, 141-148. doi: 10.1177/1359105307071748

*Saltapidas, H., & Ponsford, J. (2012). The influence of cultural background on experiences and beliefs about traumatic brain injury and their association with outcome. *Brain Impairment*, 9, 1-13. doi: 10.1375/brim.9.1.1

*Sawicki, G. S., Sellers, D. E., & Robinson, W. M. (2011). Associations between illness perceptions and health-related quality of life in adults with cystic fibrosis. *Journal of Psychosomatic Research*, 70, 161-167. doi: 10.1016/j.jpsychores.2010.06.005

*Scharloo, M., Baatenburg de Jong, R. J., Langeveld, T. P. M., van Velzen-Verkaik, E., den Akker, M. M. D.-o., & Kaptein, A. A. (2010). Illness cognitions in head and neck squamous cell carcinoma: predicting quality of life outcome. *Supportive Care in Cancer*, 18, 1137-1145. doi: 10.1007/s00520-009-0728-x

*Scharloo, M., Baatenburg de Jong, R. J., Langeveld, T. P. M., van Velzen-Verkaik, E., Doorn-op den Akker, M. M., & Kaptein, A. A. (2005). Quality of life and illness perceptions in patients with recently diagnosed head and neck cancer. *Head & Neck*, 27, 857-863. doi: 10.1002/hed.20251

- 1 *Scharloo, M., Kaptein, A. A., Schlösser, M. A. G., Pouwels, H., Bel, E. H., Rabe, K. F., &
2 Wouters, E. F. M. (2007). Illness perceptions and quality of life in patients with
3 Chronic Obstructive Pulmonary Disease. *Journal of Asthma*, 44, 575-581. doi:
4 10.1080/02770900701537438
- 5 *Scharloo, M., Kaptein, A. A., Weinman, J., Bergman, W., Vermeer, B. J., & Rooijmans, H.
6 G. M. (2000). Patients' illness perceptions and coping as predictors of functional
7 status in psoriasis: a 1-year follow-up. *British Journal of Dermatology*, 142, 899-907.
8 doi: 10.1046/j.1365-2133.2000.03469.x
- 9 *Scharloo, M., Kaptein, A. A., Weinman, J., Hazes, J. M., Willems, L. N. A., Bergman, W.,
10 & Rooijmans, H. G. M. (1998). Illness perceptions, coping and functioning in patients
11 with rheumatoid arthritis, chronic obstructive pulmonary disease and psoriasis.
12 *Journal of Psychosomatic Research*, 44, 573-585. doi: 10.1016/S0022-
13 3999(97)00254-7
- 14 *Schiaffino, K. M., Shawaryn, M. A., & Blum, D. (1998). Examining the impact of illness
15 representations on psychological adjustment to chronic illness. *Health Psychology*,
16 17, 262-268. doi: 10.1037/0278-6133.17.3.262
- 17 Schreurs, P. J. G., Van Der Willige, G., Tellegen, B., & Brosschot, J. F. (1988). *The*
18 *Utrechtse Coping List - Manual*. Lisse: Swetz & Zeitlinger.
- 19 *Schroeder, S., Achenbach, S., Körber, S., Nowy, K., de Zwaan, M., & Martin, A. (2012).
20 Cognitive-perceptual factors in noncardiac chest pain and cardiac chest Ppain.
21 *Psychosomatic Medicine*, 74, 861-868. doi: 10.1097/PSY.0b013e31826ae4ae
- 22 Schwarzer, R. (2008). Modeling health behaviour change: How to predict and modify the
23 adoption and maintenance of health behaviors. *Applied Psychology: An International*
24 *Review*, 57, 1-29. doi: 10.1111/j.1464-0597.2007.00325.x

- 1 *Sciacchitano, L., Lindner, H., & McCracken, J. (2009). Secondary beliefs: A mediator
2 between illness representations and coping behavior in arthritis sufferers. *Journal of*
3 *Rational-Emotive & Cognitive-Behavior Therapy*, 27, 23-50. doi: 10.1007/s10942-
4 007-0062-5
- 5 *Searle, A., Norman, P., Thompson, R., & Vedhara, K. (2007). A prospective examination of
6 illness beliefs and coping in patients with type 2 diabetes. *British Journal of Health*
7 *Psychology*, 12, 621-638. doi: 10.1348/135910706X164935
- 8 Seaton, M., Marsh, H. W., & Craven, R. G. (2010). Big-Fish-Little-Pond Effect:
9 Generalizability and moderation—Two sides of the same coin. *American Educational*
10 *Research Journal*, 47, 390-433. doi: 10.3102/0002831209350493
- 11 *Sharpe, L., Sensky, T., & Allard, S. (2001). The course of depression in recent onset
12 rheumatoid arthritis: The predictive role of disability, illness perceptions, pain and
13 coping. *Journal of Psychosomatic Research*, 51. doi: 10.1016/s0022-3999(01)00266-
14 5
- 15 *Sharpe, L., Sensky, T., Timberlake, N., Ryan, B., Brewin, C. R., & Allard, S. (2001). A
16 blind, randomized, controlled trial of cognitive-behavioural intervention for patients
17 with recent onset rheumatoid arthritis: preventing psychological and physical
18 morbidity. *PAIN*, 89, 275-283. doi: 10.1016/s0304-3959(00)00379-1
- 19 *Shiloh, S., Rashuk-Rosenthal, D., & Benyamini, Y. (2001). *Illness causal attributions:*
20 *Their structure and associations with other illness cognitions and perceptions of*
21 *control*. Unpublished manuscript, Tel Aviv University, Tel Aviv, Israel, Tel Aviv,
22 Israel.
- 23 *Shiyanbola, O. O., & Nelson, J. (2011). Illness perceptions, beliefs in medicine and
24 medication non-adherence among South Dakota minority women with diabetes: A
25 pilot study. *South Dakota Medicine*, 64, 365-368.

- 1 *Silva, S. M., Moreira, H. C., & Canavarro, M. C. (2012). Examining the links between
2 perceived impact of breast cancer and psychosocial adjustment: the buffering role of
3 posttraumatic growth. *Psycho-Oncology*, 21, 409-418. doi: 10.1002/pon.1913
- 4 *Simpson, J., Lekwuwa, G., & Crawford, T. (2013). Illness beliefs and psychological
5 outcome in people with Parkinson's disease. *Chronic Illness*, 9, 165-176. doi:
6 10.1177/1742395313478219
- 7 *Skinner, T. C., Carey, M. E., Cradock, S., Daly, H., Davies, M. J., Doherty, Y., . . . on
8 behalf of the DESMOND Collaborative. (2006). Diabetes education and self-
9 management for ongoing and newly diagnosed (DESMOND): Process modelling of
10 pilot study. *Patient Education and Counseling*, 64, 369-377. doi:
11 10.1016/j.pec.2006.04.007
- 12 *Skinner, T. C., & Hampson, S. E. (1998). Social support and personal models of diabetes in
13 relation to self-care and well-being in adolescents with type I diabetes mellitus.
14 *Journal of Adolescence*, 21, 703-715. doi: 10.1006/jado.1998.0190
- 15 *Skinner, T. C., Hampson, S. E., & Fife-Schaw, C. (2002). Personality, personal model
16 beliefs, and self-care in adolescents and young adults with Type 1 diabetes. *Health*
17 *Psychology*, 21, 61-70. doi: 10.1037//0278-6133.21.1.61
- 18 Skinner, T. C., Howells, L., Greene, S., Edgar, K., McEvilly, A., & Johansson, A. (2003).
19 Development, reliability and validity of the diabetes illness representations
20 questionnaire: Four studies with adolescents. *Diabetic Medicine*, 20, 283-289. doi:
21 10.1046/j.1464-5491.2003.00923.x
- 22 *Skinner, T. C., McNeil, L., Olaithe, M., Eastwood, P., Hillman, D., Phang, J., . . . Bucks, R.
23 S. (2013). Predicting uptake of continuous positive airway pressure (CPAP) therapy
24 in obstructive sleep apnoea (OSA): a belief-based theoretical approach. *Sleep and*
25 *Breathing*, 17, 1229-1240. doi: 10.1007/s11325-013-0828-1

- 1 *Sluiter, J. K., & Frings-Dresen, M. H. W. (2008). Quality of life and illness perception in
2 working and sick-listed chronic RSI patients. *International Archives of Occupational*
3 *and Environmental Health*, 81, 495-501. doi: 10.1007/s00420-007-0222-z
- 4 *Snell, D. L., Hay-Smith, E. J. C., Surgenor, L. J., & Siegert, R. J. (2013). Examination of
5 outcome after mild traumatic brain injury: The contribution of injury beliefs and
6 Leventhal's Common Sense Model. *Neuropsychological Rehabilitation*, 23, 333-362.
7 doi: 10.1080/09658211.2012.758419
- 8 *Snell, D. L., Siegert, R. J., Hay-Smith, E. J. C., & Surgenor, L. J. (2011). Associations
9 between illness perceptions, coping styles and outcome after mild traumatic brain
10 injury: Preliminary results from a cohort study. *Brain Injury*, 25, 1126-1138. doi:
11 10.3109/02699052.2011.607786
- 12 *Sniehotta, F. F., Gorski, C., & Araújo-Soares, V. (2010). Adoption of community-based
13 cardiac rehabilitation programs and physical activity following phase III cardiac
14 rehabilitation in Scotland: A prospective and predictive study. *Psychology & Health*,
15 25, 839-854. doi: 10.1080/08870440902915915
- 16 *Sofianou, S., Martynenko, M., Wolf, M. S., Wisnivesky, J. P., Krauskopf, K., Wilson, E. A.
17 H., . . . Federman, A. D. (2013). Asthma beliefs are associated with medication
18 adherence in older asthmatics. *Journal of General Internal Medicine*, 28, 67-73. doi:
19 10.1007/s11606-012-2160-z
- 20 *Spain, L. A., Tubridy, N., Kilpatrick, T. J., Adams, S. J., & Holmes, A. C. N. (2007). Illness
21 perception and health-related quality of life in multiple sclerosis. *Acta Neurologica*
22 *Scandinavica*, 116, 293-299. doi: 10.1111/j.1600-0404.2007.00895.x
- 23 *Spence, M. J., & Moss-Morris, R. (2007). The cognitive behavioural model of irritable
24 bowel syndrome: a prospective investigation of patients with gastroenteritis. *Gut*, 56,
25 1066. doi: 10.1136/gut.2006.108811

- 1 *Stafford, L., Berk, M., & Jackson, H. J. (2009). Are illness perceptions about coronary
2 artery disease predictive of depression and quality of life outcomes? *Journal of*
3 *Psychosomatic Research*, 66, 211-220. doi: 10.1016/j.jpsychores.2008.09.005
- 4 *Stafford, L., Jackson, H. J., & Berk, M. (2008). Illness beliefs about heart disease and
5 adherence to secondary prevention regimens. *Psychosomatic Medicine*, 70, 942-948.
6 doi: 10.1097/PSY.0b013e318187e22e
- 7 *Stamogiannou, I., Grunfeld, E. A., Denison, K., & Muir, G. (2004). Beliefs about illness and
8 quality of life among men with erectile dysfunction. *Int J Impot Res*, 17, 142-147. doi:
9 10.1038/sj.ijir.3901220
- 10 Stauffer, J. M. (1996). A graphical user interface psychometric meta-analysis program for
11 DOS. *Educational and Psychological Measurement*, 56, 675-677. doi:
12 10.1177/0013164496056004009
- 13 *Steed, L., Newman, S. P., & Hardman, S. M. C. (1999). An examination of the self-
14 regulation model in atrial fibrillation. *British Journal of Health Psychology*, 4, 337-
15 347. doi: 10.1348/135910799168687
- 16 *Stein, M. J., McNicholas, J., & Collis, G. M. (2001, September). *Cognitive representations*
17 *of asthma, belief concordance and psychological well-being*. Paper presented at the
18 European Health Psychology Society/British Psychological Society Division of
19 Health Psychology Annual Conference, St. Andrews, Scotland.
- 20 Stewart, A. L., Sherbourne, C. D., Hays, R. D., Wells, K. B., Nelson, E. C., Kamberg, C. J., .
21 . . Ware, J. E. (1992). Summary and discussion of MOS measures. In A. L. Stewart &
22 J. E. Ware (Eds.), *Measuring functioning and well-being: The medical outcomes study*
23 *approach* (pp. 345-371). London: Duke University Press.

*Stockford, K., Turner, H., & Cooper, M. (2007). Illness perception and its relationship to readiness to change in the eating disorders: A preliminary investigation. *British Journal of Clinical Psychology*, 46, 139-154. doi: 10.1348/014466506X115786

*Stuifbergen, A. K., Phillips, L. A., Voelmeck, W., & Browder, R. (2006). Illness perceptions and related outcomes among women with fibromyalgia syndrome. *Women's Health Issues*, 16, 353-360. doi: 10.1016/j.whi.2006.05.003

*Targosz, S. A., Kapur, N., & Creed, F. (2014). Medically unexplained symptoms, illness perception and childhood experience in neurology outpatients. *Irish Journal of Psychological Medicine*, 18, 16-20. doi: 10.1017/S0790966700006169

*Thuné-Boyle, I. C. V., Myers, L. B., & Newman, S. P. (2006). The role of illness beliefs, treatment beliefs, and perceived severity of symptoms in explaining distress in cancer patients during chemotherapy treatment. *Behavioral Medicine*, 32, 19-29. doi: 10.3200/BMED.32.1.19-29

*Tiemensma, J., Kaptein, A. A., Pereira, A. M., Smit, J. W. A., Romijn, J. A., & Biermasz, N. R. (2011). Affected Illness Perceptions and the Association with Impaired Quality of Life in Patients with Long-Term Remission of Acromegaly. *The Journal of Clinical Endocrinology & Metabolism*, 96, 3550-3558. doi: 10.1210/jc.2011-1645

*Traeger, L., Penedo, F. J., Gonzalez, J. S., Dahn, J. R., Lechner, S., C., Schneiderman, N., & Antoni, M. H. (2009). Illness perceptions and emotional well-being in men treated for localized prostate cancer. *Journal of Psychosomatic Research*, 67, 389-397. doi: 10.1016/j.jpsychores.2009.03.013

*Treharne, G. J., Kitas, G. D., Lyons, A. C., & Booth, D. A. (2005). Well-being in rheumatoid arthritis: The effects of disease duration and psychosocial factors. *Journal of Health Psychology*, 10, 457-474. doi: 10.1177/1359105305051416

- 1 *Treharne, G. J., Lyons, A. C., Hale, E. D., Douglas, K. M. J., Goodchild, C. E., Booth, D.
2 A., & Kitas, G. D. (2007). Sleep disruption frequency in rheumatoid arthritis:
3 Perceived stress predicts poor outcome over one year. *Musculoskeletal Care*, 5, 51-
4 64. doi: 10.1002/msc.99
- 5 *Treharne, G. J., Lyons, A. C., Hale, E. D., Goodchild, C. E., Booth, D. A., & Kitas, G. D.
6 (2008). Predictors of fatigue over 1 year among people with rheumatoid arthritis.
7 *Psychology, Health & Medicine*, 13, 494-504. doi: 10.1080/13548500701796931
- 8 *Trovato, G. M., Pace, P., Tamburino, C., Garufi, G., Martines, G. F., Pirri, C., . . . Catalano,
9 D. (2010). Elective coronary stent patients: preinterventional functional status and
10 clinical-instrumental assessment. *Heart and Vessels*, 25, 82-86. doi: 10.1007/s00380-
11 009-1163-0
- 12 Turk, D. C., Rudy, T. E., & Salovey, P. (1986). Implicit models of illness. *Journal of*
13 *Behavioral Medicine*, 9, 453-474. doi: 10.1007/BF00845133
- 14 *Turriff, A., Levy, H. P., & Biesecker, B. (2011). Prevalence and psychosocial correlates of
15 depressive symptoms among adolescents and adults with Klinefelter syndrome. *Genet*
16 *Med*, 13, 966-972. doi: 10.1097/GIM.0b013e3182227576
- 17 *Twiddy, M., House, A., & Jones, F. (2012). The association between discrepancy in illness
18 representations on distress in stroke patients and carers. *Journal of Psychosomatic*
19 *Research*, 72, 220-225. doi: 10.1016/j.jpsychores.2011.12.004
- 20 US Department of Health and Human Services. (2011). *Health, United States, 2010: With*
21 *special feature on death and dying*. Hyattsville, MD: Centers for Disease Control and
22 Prevention.
- 23 US Department of Health and Human Services. (2013). Multiple chronic conditions - a
24 strategic framework: Optimum health and quality of life for individuals with multiple

- 1 chronic conditions. Retrieved June 20, 2016, from
2 http://www.hhs.gov/ash/initiatives/mcc/mcc_framework.pdf
- 3 *van der Have, M., Minderhoud, I. M., Kaptein, A. A., Leenders, M., Siersema, P. D., Fidler,
4 H. H., & Oldenburg, B. (2013). Substantial impact of illness perceptions on quality of
5 life in patients with Crohn's disease. *Journal of Crohn's and Colitis*, 7, e292-e301.
6 doi: 10.1016/j.crohns.2012.11.002
- 7 *van der Velde, J. L., Flokstra-de Blok, B. M. J., Hamp, A., Knibb, R. C., Duiverman, E. J.,
8 & Dubois, A. E. J. (2011). Adolescent–parent disagreement on health-related quality
9 of life of food-allergic adolescents: who makes the difference? *Allergy*, 66, 1580-
10 1589. doi: 10.1111/j.1398-9995.2011.02726.x
- 11 *van Os, S., Norton, S., Hughes, L. D., & Chilcot, J. (2012). Illness perceptions account for
12 variation in positive outlook as well as psychological distress in Rheumatoid Arthritis.
13 *Psychology, Health & Medicine*, 17, 427-439. doi: 10.1080/13548506.2011.626432
- 14 *van Wilgen, C. P., Kaptein, A. A., & Brink, M. S. (2010). Illness perceptions and mood
15 states are associated with injury-related outcomes in athletes. *Disability and*
16 *Rehabilitation*, 32, 1576-1585. doi: 10.3109/09638281003596857
- 17 *van Wilgen, C. P., van Ittersum, M. W., Kaptein, A. A., & van Wijhe, M. (2008). Illness
18 perceptions in patients with fibromyalgia and their relationship to quality of life and
19 catastrophizing. *Arthritis and Rheumatism*, 58, 3618-3626. doi: 10.1002/art.23959
- 20 *Var, F., & Rajeswaran, J. (2012). Perception of illness in patients with traumatic brain
21 injury. *Indian Journal of Psychological Medicine*, 34, 223-226. doi: 10.4103/0253-
22 7176.106014
- 23 *Vaughan, R., Morrison, L., & Miller, E. (2003). The illness representations of multiple
24 sclerosis and their relations to outcome. *British Journal of Health Psychology*, 8, 287-
25 301. doi: 10.1348/135910703322370860

- 1 Veit, C. T., & Ware, J. E. (1983). The structure of psychological distress and well-being in
2 general populations. *Journal of Consulting and Clinical Psychology*, 51, 730-742.
- 3 Vitaliano, P. P., Russo, J., Carr, J. E., Maiuro, R. D., & Becker, J. (1985). The ways of coping
4 checklist: Revision and psychometric properties. *Multivariate Behavioral Research*,
5 20, 3-26. doi: 10.1207/s15327906mbr2001_1
- 6 *Vollmann, M., Kalkouskaya, N., Langguth, B., & Scharloo, M. (2012). When the ringing in
7 the ears gets unbearable: Illness representations, self-instructions and adjustment to
8 tinnitus. *Journal of Psychosomatic Research*, 73, 108-111. doi:
9 10.1016/j.jpsychores.2012.04.009
- 10 *Vollmann, M., Scharloo, M., Langguth, B., Kalkouskaya, N., & Salewski, C. (2014). Illness
11 representations as mediators of the relationship between dispositional optimism and
12 depression in patients with chronic tinnitus: A cross-sectional study. *Psychology &*
13 *Health*, 29, 81-93. doi: 10.1080/08870446.2013.828294
- 14 *Vurnek Živković, M., Buljan, M., Blajić, I., & Šitum, M. (2008). Psychological status and
15 illness perceptions in patients with melanoma. *Collegium Antropologicum*, 32, 75-78.
- 16 *Walsh, J. C., Lynch, M., Murphy, A. W., & Daly, K. (2004). Factors influencing the
17 decision to seek treatment for symptoms of acute myocardial infarction. *Journal of*
18 *Psychosomatic Research*, 56, 67-73. doi: 10.1016/S0022-3999(03)00082-5
- 19 *Wamsteker, E. W., Geenen, R., Iestra, J., Larsen, J. K., Zelissen, P. M. J., & van Staveren,
20 W. A. (2005). Obesity-related beliefs predict weight loss after an 8-week low-calorie
21 diet. *Journal of the American Dietetic Association*, 105, 441-444. doi:
22 10.1016/j.jada.2004.12.031
- 23 Ware, J. E., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-
24 36): I. Conceptual framework and item selection. *Medical Care*, 30, 473-483.

- *Watkins, K. W., Connell, C. M., Fitzgerald, J. T., Klem, L., Hickey, T., & Ingersoll-Dayton, B. (2000). Effect of adults' self-regulation of diabetes on quality-of-life outcomes. *Diabetes Care*, 23, 1511.
- Wearden, A., & Peters, S. (2008). Therapeutic techniques for interventions based on Leventhal's common sense model. *British Journal of Health Psychology*, 13, 189-193. doi: 10.1348/135910708X295613
- *Weinman, J., Petrie, K. J., Moss-Morris, R., & Horne, R. (1996). The illness perception questionnaire: A new method for assessing the cognitive representation of illness. *Psychology and Health*, 11, 431-445. doi: 10.1080/08870449608400270
- *Wenninger, K., Helmes, A., Bengel, J., Lauten, M., Völkel, S., & Niemeyer, C. M. (2013). Coping in long-term survivors of childhood cancer: relations to psychological distress. *Psycho-Oncology*, 22, 854-861. doi: 10.1002/pon.3073
- Whitmarsh, A., Koutantji, M., & Sodell, K. (2003). Illness perceptions, mood and coping in predicting attendance at cardiac rehabilitation. *British Journal of Health Psychology*, 8, 209-221. doi: 10.1348/135910703321649178
- *Whittaker, R., Kemp, S., & House, A. (2007). Illness perceptions and outcome in mild head injury: a longitudinal study. *Journal of Neurology, Neurosurgery & Psychiatry*, 78, 644. doi: 10.1136/jnnp.2006.101105
- *Wikman, A., Molloy, G. J., Randall, G., & Steptoe, A. (2011). Cognitive predictors of posttraumatic stress symptoms six months following acute coronary syndrome. *Psychology & Health*, 26, 974-988. doi: 10.1080/08870446.2010.512663
- *Woith, W. M., & Larson, J. L. (2008). Delay in seeking treatment and adherence to tuberculosis medications in Russia: A survey of patients from two clinics. *International Journal of Nursing Studies*, 45, 1163-1174. doi: 10.1016/j.ijnurstu.2007.07.014

- 1 *Wu, L. M., Mohamed, N. E., Winkel, G., & Diefenbach, M. A. (2013). Patient and spouse
2 illness beliefs and quality of life in prostate cancer patients. *Psychology & Health*, 28,
3 355-368. doi: 10.1080/08870446.2012.722219
- 4 Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta*
5 *Psychiatrica Scandinavica*, 67, 361-370. doi: 10.1111/j.1600-0447.1983.tb09716.x
- 6 *Žugelj, U., Zupančič, M., Komidar, L., Kenda, R., Varda, N. M., & Gregorič, A. (2010).
7 Self-reported adherence behavior in adolescent hypertensive patients: The role of
8 illness representations and personality. *Journal of Pediatric Psychology*, 35, 1049-
9 1060. doi: 10.1093/jpepsy/jsq027

Figure 1. Schematic representation of common-sense model of illness representations based on Leventhal et al.'s (1992) illustration. The coping strategy and illness outcome categories are based on the classification procedure used in the present study developed by Hagger and Orbell (2003).





Figure 2. Basic process model derived from the common-sense model.

Figure 3. Path diagram of generalized effects in the revised common-sense process model. Solid lines represent hypothesized effects of beliefs on coping strategies and effects of coping strategies on illness outcomes, and broken lines represent moderating effects. Direct effects of cognitive and emotional representations on illness outcomes omitted for clarity.




Figure 4. Path diagram of moderating effects of illness treatability (higher scores represent greater treatability) on mediated relations of perceived consequences and control on illness outcomes mediated by problem- and emotion-focused coping strategies in the revised common-sense process model. Solid lines represent hypothesized effects of representations on coping strategies and effects of coping strategies on illness outcomes, and broken lines represent moderating effects. Direct effects of cognitive and emotional representations on illness outcomes omitted for clarity.




Figure 5. Path diagram of moderating effects of emotional representations (higher scores represent greater emotional responses) on mediated relations of perceived consequences and control on illness outcomes mediated by problem- and emotion-focused coping strategies in the revised common-sense process model. Solid lines represent hypothesized effects of representations on coping strategies and effects of coping strategies on illness outcomes, and broken lines represent moderating effects. Direct effects of cognitive and emotional representations on illness outcomes omitted for clarity.




Figure 6. Path diagram showing independent effects of beliefs about coping behavior, derived from social cognitive theories (Bandura, 1977; Fishbein & Ajzen, 2009), and beliefs about illness and action plans, derived from the common-sense model (Leventhal et al., 1980; Leventhal et al., 2016) on problem-focused coping behaviors in the revised common-sense process model. Direct effects of cognitive and emotional representations on illness outcomes omitted for clarity. The proposed model is a generalized one with constructs comprising multiple dimensions of constructs (e.g., behavioral beliefs, cognitive representations) that may have effects on outcomes with different signs.

