

EXTENDED GENOME REPORT

Open Access



The complete genome sequence of the rumen bacterium *Butyrivibrio hungatei* MB2003

Nikola Palevich^{1,2*} , William J. Kelly¹, Sinead C. Leahy¹, Eric Altermann¹, Jasna Rakonjac² and Graeme T. Attwood¹

Abstract

Butyrivibrio hungatei MB2003 was isolated from the plant-adherent fraction of rumen contents from a pasture-grazed New Zealand dairy cow, and was selected for genome sequencing in order to examine its ability to degrade plant polysaccharides. The genome of MB2003 is 3.39 Mb and consists of four replicons; a chromosome, a secondary chromosome or chromid, a megaplasmid and a small plasmid. The genome has an average G + C content of 39.7%, and encodes 2983 putative protein-coding genes. MB2003 is able to use a variety of monosaccharide substrates for growth, with acetate, butyrate and formate as the principal fermentation end-products, and the genes encoding these metabolic pathways have been identified. MB2003 is predicted to encode an extensive repertoire of CAZymes with 78 GHs, 7 CEs, 1 PL and 78 GTs. MB2003 is unable to grow on xylan or pectin, and its role in the rumen appears to be as a utilizer of monosaccharides, disaccharides and oligosaccharides made available by the degradative activities of other bacterial species.

Keywords: Rumen, Bacteria, Hemicellulose, Pectin, Degradation, *Butyrivibrio*, Genome

Introduction

Butyrivibrio are important rumen bacteria [1], and are among the small number of rumen genera capable of utilizing the complex plant structural polysaccharides xylan and pectin [2, 3]. They are classified as anaerobic, monotrichous, butyrate-producing, curved rods and have been isolated from the gastrointestinal tracts and feces of various ruminants, monogastric animals and humans [4, 5]. *Butyrivibrio* are metabolically versatile and are capable of growing on a range of carbohydrates, from simple mono- or oligosaccharides to complex plant polysaccharides such as pectins, mannans, starch and hemicelluloses [6]. Furthermore, xylans of diverse chemical and physical properties, from a range of forages are degraded by *Butyrivibrio* species [7]. Some *Butyrivibrio* species show strong proteolytic activity [8], and *Butyrivibrio* are thought to be the main butyrate producers in the rumen [9, 10]. The genus *Butyrivibrio*

is classified within the family *Lachnospiraceae*, order *Eubacteriales*, and is phylogenetically diverse. The *Butyrivibrio* genus originally consisted of only one species, *Butyrivibrio fibrisolvens* [2]. In addition to phenotypic characterisations [11, 12], studies have utilized DNA-DNA hybridization [13, 14], 16S rRNA gene sequencing [15, 16] and 16S rRNA-based hybridization probes [17], to differentiate these organisms. To accommodate the observed diversity amongst the newly discovered bacterial strains, a new genus, *Pseudobutyrvibrio*, was described [18]. Four species are currently recognized: *B. fibrisolvens*, *B. hungatei*, *B. proteoclasticus* and *B. crossotus* [6], although *B. crossotus* is more distantly related to the other three. *B. hungatei* are common anaerobic rumen bacteria found in domestic and wild ruminants and the type strain is JK615^T [19]. *Butyrivibrio hungatei* JK615^T is non-proteolytic and non-fibrolytic, but is able to utilize oligo- and monosaccharides as substrates for growth. Gaining an insight into the role of these secondary degrader species in microbial plant polysaccharide breakdown is important for understanding rumen function. Here we present the

* Correspondence: nik.palevich@agresearch.co.nz

¹Rumen Microbiology, Animal Science, AgResearch Limited, Grasslands Research Centre, Tennent Drive, Private Bag 11008, Palmerston North 4442, New Zealand

²Institute of Fundamental Sciences, Massey University, Palmerston North, New Zealand




Fig. 1 Morphology of *B. hungatei* MB2003. Micrograph of Gram stained *B. hungatei* MB2003 cells at 100× magnification. Bar represents 10 μm

complete genome sequence of *Butyrivibrio hungatei* MB2003, a strain isolated from a pasture-grazed dairy cow in New Zealand [20], and describe its comparison with genomes of closely related *B. hungatei* strains.

Organism information

Classification and features

MB2003 was isolated from the plant-adherent fraction of rumen contents from a New Zealand dairy cow grazing fresh forage [20, 21]. MB2003 cells are Gram positive, short rods, occurring singly or in pairs (Fig. 1). The morphological features of MB2003 cells were determined by electron microscopy of cells grown on RM02 medium [22], negatively stained with 1% phosphotungstic acid, mounted on Formvar-coated copper grids, and examined using a Philips model 201C electron microscope (Eindhoven, The Netherlands). MB2003 cells were observed to have a




Fig. 2 Transmission electron micrograph of *B. hungatei* MB2003. Micrograph of negatively stained *B. hungatei* MB2003 cells at 10,000× magnification

single polar flagellum (Fig. 2), although cells in growing cultures were non-motile. A phylogenetic analysis of the full-length 16S rRNA gene sequence placed MB2003 within the *B. hungatei* species, being 98% similar to the *Butyrivibrio hungatei* type strain JK615^T [19] (Fig. 3). Additional characteristics of *B. hungatei* MB2003 are shown in Table 1.

Strain MB2003 grew to highest optical density (OD) at pH values of 6.1 to 6.5 and at a temperature of 39 °C, conditions which are typical of its rumen environment. VFA production was determined from triplicate broth cultures grown overnight in RM02 medium with cellobiose as substrate and analysed for formate, acetate, propionate, *n*-butyrate, *iso*-valerate and lactate on a HP 6890 series GC (Hewlett Packard) with 2-ethylbutyric acid (Sigma-Aldrich, St. Louis, MO, USA) as the internal standard. To derivatize formic, lactic and succinic acids, samples were mixed with HCl ACS reagent (Sigma-Aldrich, St. Louis, MO, USA) and diethyl ether, with the addition of *N*-methyl-*N*-*t*-butyldimethylsilyltri-fluoroacetamide (MTBSTFA) (Sigma-Aldrich, St. Louis, MO, USA) [23]. Under these conditions MB2003 produced 16.4 mM formate, 3.6 mM acetate and 4.7 mM butyrate. MB2003 was able to grow in CO₂-containing media with various soluble carbon sources and the semi-soluble inulin (all tested at 0.5% *w/v* final concentration). Growth on soluble substrates was assessed as an increase in culture density OD_{600nm} compared to cultures without carbon source added, whereas total VFA production was used as an indicator of substrate utilization and growth for insoluble polymers (Table 2). All strains tested were net producers of formate, acetate and *n*-butyrate, which is characteristic of *Butyrivibrio*. Cellobiose and glucose supported the growth of MB2003, JK615^T and B316^T to high cell densities. Therefore, cellobiose was used to examine the growth of MB2003 over a 24 h period. The exponential phase of growth was between 4 and 8 h, with the maximum cell density reached at 8 to 10 h, and stationary phase between 10 to 24 h (Fig. 4).

Genome sequencing information

Genome project history

Butyrivibrio hungatei MB2003 was selected for genome sequencing as a NZ strain of *B. hungatei*. A summary of the genome project information is shown in Table 3 and in Additional file 1: Table S1.

Growth conditions and genomic DNA preparation

MB2003 was grown in RM02 medium [22] with 10 mM glucose and 0.1% yeast extract but without rumen fluid. Culture purity was confirmed by Gram




Fig. 3 Phylogenetic tree highlighting the relationship of *B. hungatei* MB2003 relative to the type strains of the other species within the genus *Butyrivibrio*. The evolutionary history was inferred using the Maximum Likelihood method based on the General Time Reversible model [55]. The tree with the highest log likelihood (-3712.3329) is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (10,000 replicates) is shown next to the branches [56]. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. A discrete Gamma distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter=0.3950)). The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The analysis involved six nucleotide sequences. All positions with less than 95% site coverage were eliminated. There were a total of 1509 positions in the final dataset. Evolutionary analyses were conducted in MEGA6 [55]. GenBank accession numbers of the 16S rRNA gene sequences are shown in parentheses. Bar, 0.02 nucleotide substitutions per site. ^T, indicates type strain. All the type strains have their genome sequencing projects registered in the Genomes Online Database (GOLD) [57]

stain and sequencing of the 16S rRNA gene. Genomic DNA was extracted from freshly grown cells by a modification of the standard cell lysis method of Saito and Miura [24], using lysozyme, proteinase K and sodium dodecyl sulphate, followed by phenol-chloroform extraction, and purification using the Qiagen Genomic-Tip 500 Maxi kit (Qiagen, Hilden, Germany). Genomic DNA was precipitated by the addition of a 0.7 volume of isopropanol, and collected by centrifugation at 12,000×g for 10 min at room temperature. The supernatant was removed, and the DNA pellet was washed in 70% ethanol, re-dissolved in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 7.5) and stored at -20 °C until required.

Genome sequencing and assembly

The complete genome sequence of MB2003 was determined by pyrosequencing 3 kb mate paired-end sequence libraries using the 454 GS FLX platform with Titanium chemistry (Macrogen, Korea). Pyrosequencing reads provided 234× coverage of the genome and were assembled using the Newbler assembler (version 2.7, Roche 454 Life Sciences, USA) which resulted in 31 contigs across 7 scaffolds. Gap closure was managed using the Staden package [25] and gaps were closed using additional Sanger sequencing by standard and inverse PCR techniques. In addition, MB2003 genomic DNA was sequenced using shotgun sequencing of 2 kb paired-end sequence libraries using the Illumina MiSeq platform (Macrogen, Korea) which provided 800-fold sequencing coverage. Illumina reads were analysed using the Galaxy web-based platform [26] and de novo assembly was performed using the

Velvet assembler, version 3.0 [27]. The Velvet assembled MB2003 genome MiSeq sequences were combined with the Newbler assembly using the Staden package and Geneious, version 8.1 [28]. Genome assembly was confirmed by pulsed-field gel electrophoresis.

Genome annotation

Annotation of the MB2003 genome was performed as described previously [29]. The MB2003 genome sequence was prepared for NCBI submission using Sequin [30], and the adenine residue of the start codon of the chromosomal replication initiator protein DnaA1 (bhn_I0001, bhn_RS00450) gene was chosen as the first base for the MB2003 genome.

Genome properties

The genome of *B. hungatei* MB2003 consists of four replicons [21, 31]; a single chromosome (3,143,784 bp, %G + C 39.91), a chromid or secondary chromosome (BhuII, 91,776 bp, %G + C 37.71), a megaplasmid (pNP144, 144,470 bp, %G + C 36.86) and a plasmid (pNP6, 6284 bp, %G + C 35.71). The total size of the closed genome is 3,386,314 bp with an overall %G + C content of 39.71%. A total of 3064 genes were predicted, of which 2983 (97.4%) were protein-coding genes. A putative function was assigned to 2225 of the protein-coding genes, while 775 protein coding genes were annotated as hypothetical proteins. The MB2003 chromosome encodes 2758 genes, and BhuII, pNP144 and pNP6 encode 89, 147 and 6 genes, respectively. The properties and statistics of the MB2003 genome are

Table 1 Classification and general features of the rumen bacterium *B. hungatei* MB2003 in accordance with the MIGS recommendations [58]

MIGS ID	Property	Term	Evidence code ^a
	Current classification	Domain: <i>Bacteria</i>	TAS [59]
		Phylum: <i>Firmicutes</i>	TAS [60, 61]
		Class: <i>Clostridia</i>	TAS [62]
		Order: <i>Eubacteriales</i>	TAS [63]
		Family: <i>Lachnospiraceae</i>	TAS [64]
		Genus: <i>Butyrivibrio</i>	TAS [4]
		Species: <i>hungatei</i>	TAS [19]
		Type strain: No	
		Strain: MB2003	TAS [20, 21]
	Gram stain	Positive	TAS [21, 31]
	Cell shape	Rod	TAS [21, 31]
	Motility	Non-motile	IDA
	Sporulation	Not reported	NAS
	Temperature range	37–39 °C	IDA
	Optimum temperature	39 °C	IDA
	pH range; Optimum	6.0–7.0; 6.4	IDA
	Carbon source	Variety of carbohydrates	IDA
	Energy metabolism	Fermentative metabolism	IDA
MIGS-6	Habitat	Bovine rumen	TAS [20]
MIGS-6.3	Salinity	Not reported	
MIGS-22	Oxygen requirement	Anaerobic	IDA
MIGS-15	Biotic relationship	Symbiont of ruminants	TAS [20]
MIGS-14	Pathogenicity	Non-pathogen	NAS
MIGS-4	Geographic location	Ruakura, Hamilton, New Zealand	TAS [20]
MIGS-5	Sample collection time	May 2009	TAS [20]
MIGS-4.1	Latitude	−37.77 (37°46′28″S)	IDA
MIGS-4.2	Longitude	+175.31 (175°18′31″E)	IDA
MIGS-4.4	Altitude	40 m	IDA

^aEvidence codes - IDA, Inferred from Direct Assay, NAS, Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). Evidence codes are from the Gene Ontology project [65]

summarized in Tables 4, 5 and 6. The nucleotide sequences of the MB2003 chromosome, chromid (BhuII), megaplasmid (pNP144) and plasmid (pNP6) have been deposited in Genbank under accession numbers CP017831, CP017830, CP017832 and CP017833. The genome atlas for *B. hungatei* MB2003 is shown in Fig. 5.

Insights from the genome sequence

Comparison of the MB2003, *B. hungatei* JK615^T, and *B. proteoclasticus* B316^T genomes

A comparison of the *B. hungatei* MB2003 genome with the draft genome of *B. hungatei* JK615^T [32] and the complete *B. proteoclasticus* B316^T genome is shown in Table 7. The MB2003 genome is 8633 bp smaller than

Table 2 Carbon source utilization of the *Butyrivibrio* strains

Substrate		MB2003	JK615 ^T	B316 ^T
Monosaccharides	Arabinose	++	++	++
	Fructose	-	-	++
	Galactose	++	-	++
	Glucose	++	++	++
	Mannose	-	++	++
	Rhamnose	-	-	++
	Ribose	-	-	-
	Xylose	++	++	++
Disaccharides	Cellobiose	++	++	++
	Lactose	++	++	++
	Maltose	++	++	++
	Melibiose	-	-	+
	Sucrose	++	++	++
Trisaccharides	Melezitose	-	-	++
	Raffinose	-	++	++
	Trehalose	-	-	++
Sugar Alcohols	myo-Inositol	-	-	-
	Mannitol	-	-	+
	Sorbitol	-	-	-
Glycosides	Amygdalin	+	-	++
	Esculin	-	++	++
	Rutin	-	++	++
	Salicin	++	++	++
Insoluble substrates	Cellulose	-	-	-
	Dextrin	-	-	++
	Inulin	+	-	++
	Starch	-	-	++
	Pectin	-	-	++
	Xylan	-	-	++

ΔOD_{600nm} readings of 0.5–1.0 were scored as ++, 0.2–0.5 scored as +, and 0–0.2 scored as -. Results for *B. hungatei* JK615^T and *B. proteoclasticus* B316^T are adapted from Kopečný et al. [19] and Moon et al. [6], respectively

JK615^T and contains 27 fewer protein-coding genes. Although several plasmid replication genes have been identified in the JK615^T draft genome, the presence of extrachromosomal elements requires experimental validation.

A novel feature of both the MB2003 and B316^T genomes is the presence of chromids or secondary chromosomes [33]. Chromids are replicons that have %G + C content similar to that of their main chromosome, but have plasmid-type maintenance and replication systems, are smaller than the chromosome, but are usually larger than any other plasmids present. Chromids contain genes essential for growth and maintenance of the organism along with several core

genus-specific genes that can be found on the chromosome in other species of bacteria [33]. The Bhu II replicon has most of these characteristics and therefore has been designated as a chromid of MB2003. In B316^T, almost 10% of the genes encoding enzymes that have a role in carbohydrate metabolism and transport are found on the chromid [29]. The Bhu II chromid of MB2003 also encodes genes with similar predicted functions (Table 9). Since the Bhu II chromid of MB2003 is smaller than the BPC2 chromid of B316^T (186,325 bp), it is now the smallest chromid reported for bacteria. Comparison of MB2003, JK615^T and B316^T genomes based on COG category (Table 8) and synteny analysis (Fig. 6), show that these




Fig. 4 Culture density achieved in 24 h by MB2003 growing in media with cellobiose as the sole substrate. Points indicate means of three replicates, and the error bars represent +/-one standard error

Butyrivibrio species and strains are genetically similar. Although the MB2003 and B316^T genome sizes differ, the basic metabolism of these two rumen bacterial species is indicated to be similar.

Butyrate production

For the production of butyrate and H₂ from glucose, the MB2003 genome possesses a pyruvate:ferredoxin oxidoreductase gene, *nifj* (bhn_I2528) required for pyruvate conversion to acetyl-CoA, as well as a butyryl-CoA dehydrogenase/electron transferring flavoprotein *bcd-etfAB* (bhn_I2225, bhn_I2221 and bhn_I2222) to generate ATP by classic substrate level phosphorylation. In addition, MB2003 possesses genes that encode all six subunits of the Rnf (*rnfA*, *rnfB*, *rnfC*, *rnfD*, *rnfE*, *rnfG*) and Ech (*echA*, *echB*, *echC*, *echD*, *echE*, *echF*) hydrogenases. These pathways

involve the transmembrane ion pumps Ech [34] or Rnf [35–38], that generate a transmembrane proton and/or sodium electrochemical potential from redox cofactors for ATP synthesis by ETP [34, 36]. The MB2003 genome does not possess genes for Por-ABDG, a pyruvate ferredoxin oxidoreductase similar in function to Nifj or genes for EhaA-R, EhbA-P, HydA-C, MbhLKJ, or MvhADG/HdrABC similar in function to the Fd-dependent Ech hydrogenase. In addition, an alternative pathway exists where formate is predicted to be the end product and involves the decarboxylation of acetyl-CoA by a pyruvate formate lyase *pflB* (bhn_I0124) instead of Nifj. It has been proposed that Ech and Rnf work in concert with Nifj and Bcd-Etf complex to drive ATP synthesis by ETP during glucose fermentation to butyrate [34, 36, 39]. Interestingly, the vast majority of anaerobic prokaryotes appear to possess either an Ech or Rnf but not both [40, 41]. However, a recent analysis of rumen prokaryotic genomes identified *Butyrivibrio* and *Pseudobutyrvibrio* as a rare group of bacteria that possess genes for both Ech and Rnf [42]. These findings warrant further biochemical investigation to determine the activity of Ech and Rnf in *Butyrivibrio*.

The MB2003 pathways for butyrate production presume the possession of a complete Embden-Meyerhof-Parnas glycolytic pathway. Enolase (*eno*, EC4.2.1.11), converts 2-phospho-D-glycerate to phosphoenolpyruvate in the second to last step of

Table 3 MB2003 genome project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	High-quality, closed genome
MIGS-28	Libraries used	454 3 kb mate paired-end library
MIGS-29	Sequencing platforms	454 GS FLX Titanium chemistry
MIGS-31.2	Fold coverage	234x
MIGS-30	Assemblers	Newbler version 2.3
MIGS-32	Gene calling method	Glimmer and BLASTX
	Locus Tag	bhn and bhn_RS
	Genbank ID	CP017830, CP017831, CP017832, CP017833
	Genbank Date of Release	31 October 2016
	GOLD ID	Ga0074201
	BIOPROJECT ID	PRJNA349214 and PRJNA224116
	BIOSAMPLE ID	SAMN05928573
MIGS-13	Source Material Identifier	<i>Butyrivibrio hungatei</i> MB2003
	Project relevance	Ruminant plant-fibre degradation

Table 4 Summary of MB2003 genome replicon features

Replicon type	Size (bp)	Topology	INSDC identifier	RefSeq ID
Chromosome	3,143,784	circular	CP017831	NZ_CP017831
Chromid_Bhull	91,776	circular	CP017830	NZ_CP017830
Megaplasmid_pNP144	144,470	circular	CP017832	NZ_CP017832
Plasmid_pNP6	6284	circular	CP017833	NZ_CP017833

Table 5 MB2003 genome statistics

Attribute	Value	% of total ^a
Genome size (bp)	3,386,314	100
DNA coding (bp)	3,064,986	90.51
DNA G + C (bp)	1,344,683	39.71
DNA scaffolds	4	100
Total genes	3064	100
Protein coding genes	2983	97.36
RNA genes	60	1.96
Pseudogenes	17	0.56
Genes in internal clusters	160	5.22
Genes with function predicted	2247	73.34
Genes assigned to COGs	1842	61.34
Genes with Pfam domains	2350	78.26
Genes with signal peptides	148	4.93
Genes with transmembrane helices	881	29.34
CRISPR repeats	2	

^aThe total is based on either the size of the genome in base pairs or the total number of genes or protein-coding genes in the annotated genome

the EMP pathway. Previous work has shown that B316^T lacks a detectable enolase [29], and the Methylglyoxal Shunt was proposed as a possible alternative to the EMP pathway. In this pathway the dihydroxyacetone phosphate is transformed to pyruvate via methylglyoxal and D-lactate dehydrogenase, encoded by *ldhA* [43]. The MB2003 genome possesses two methylglyoxal synthase genes, *mgsA* (bhn_I1328 and bhn_I1996), glyoxalase gene *gloA* (bhn_I1783) and an alternative L-lactate dehydrogenase, encoded by *ldh* (bhn_I0363). MB2003 has the same set of genes as B316^T for the production of butyrate, formate, acetate and lactate, but also is the only *B. hungatei* reported to date that lacks a detectable enolase gene. Genome sequences from a wider range of *B. hungatei* and *B. proteoclasticus* strains are required to determine if these are common features in these organisms.

Polysaccharide degradation

The Carbohydrate-Active enZymes database was used to identify glycoside hydrolases, glycosyl transferases, polysaccharide lyases, carbohydrate esterases and carbohydrate-binding protein module families within the MB2003 genome. MB2003 has a similar CAZyme profile to B316^T [21, 31], and analysis of the functional domains of enzymes involved in the breakdown or synthesis of complex carbohydrates, has revealed the polysaccharide-degrading potential of this rumen bacterium.


Approximately 3% of the MB2003 genome (90 CDSs) is predicted to encode either secreted or intracellular

proteins dedicated to polysaccharide degradation, similar to that found in B316^T. The MB2003 genome is predicted to encode 19 secreted (16 GHs, two CEs and one CBP) and 65 intracellular (59 GHs, 5 CEs and one PL) proteins involved in polysaccharide breakdown (Table 9). The enzymatic profiles of MB2003 and JK615^T are almost identical, as both possess the same genes encoding predicted secreted and intracellular CAZymes in their genomes (Table 9). Out of the 19 genes predicted to encode secreted polysaccharide degrading enzymes, only two, lysozyme *lyc25B* (bhn_III074) and feruloyl esterase *estIA* (bhn_III076), are encoded by the MB2003 chromid (Bhu II). MB2003 has no secreted enzyme larger than 1000 aa in size, with the average size secreted enzymes being 510 aa. The majority (59) of MB2003 genes involved in polysaccharide breakdown (excluding GTs), had corresponding homologues in B316^T and

Table 6 Number of genes associated with the general COG functional categories

Code	Value	% of total ^a	Description
J	194	9.52	Translation, ribosomal structure and biogenesis
A	0	0	RNA processing and modification
K	149	7.31	Transcription
L	88	4.32	Replication, recombination and repair
B	0	0	Chromatin structure and dynamics
D	32	1.57	Cell cycle control, Cell division, chromosome partitioning
V	65	3.19	Defense mechanisms
T	139	6.82	Signal transduction mechanisms
M	155	7.61	Cell wall/membrane biogenesis
N	61	2.99	Cell motility
U	22	1.08	Intracellular trafficking and secretion
O	78	3.83	Posttranslational modification, protein turnover, chaperones
C	69	3.39	Energy production and conversion
G	243	11.93	Carbohydrate transport and metabolism
E	177	8.69	Amino acid transport and metabolism
F	80	3.93	Nucleotide transport and metabolism
H	79	3.88	Coenzyme transport and metabolism
I	72	3.53	Lipid transport and metabolism
P	79	3.88	Inorganic ion transport and metabolism
Q	16	0.79	Secondary metabolites biosynthesis, transport and catabolism
R	158	7.76	General function prediction only
S	73	3.58	Function unknown
-	1245	40.33	Not in COGs

^aThe total is based on the total number of protein coding genes in the genome



JK615^T. Three of the genes encoding intracellular proteins were found in the Bhu II chromid: a β -glucosidase *bgl3A* (bhn_III062), a β -galactosidase *bga42A* (bhn_III010) and a polysaccharide deacetylase *est4A* (bhn_III070). The analysis of the Pfam domains from the most abundant GH families (GH2, GH31, GH3, GH13 and GH43) showed they did not contain signal sequences and hence were predicted to be located intracellularly. Similarly, CAZymes with predicted roles in xylan and pectin degradation, the GH8, GH28, GH51, GH67, GH88, GH105, GH115, CE2 and CE10 families were also predicted to be intracellular. Of these, MB2003 contains CAZymes with homologues in B316^T except for the α -L-arabinofuranosidase *arf51C* (bhn_I1509). These findings suggest that a variety of complex oligosaccharides

resulting from extracellular hydrolysis are metabolized within the cell.

Growth experiments showed MB2003 to be a metabolically versatile bacterium able to grow on a wide variety of monosaccharides, disaccharides and glycosides (Table 2). However, unlike B316^T, MB2003 and JK615^T were unable to utilize the insoluble substrates pectin and xylan for growth (Table 2). In addition, MB2003, JK615^T and B316^T are unable to degrade cellulose, however among these organisms, only B316^T is able to utilize a range of other insoluble plant polysaccharides. The ability of B316^T to breakdown pectin, starch and xylan is predicted to be based on nine large (>1000 aa) cell-associated proteins shown to be significantly up-regulated in B316^T cells grown on xylan [44]. These are: α -

Table 7 Genome statistics of MB2003, JK615^T and B316^T

Attribute	<i>B. hungatei</i> MB2003		<i>B. hungatei</i> JK615 ^{Tb}		<i>B. proteoclasticus</i> B316 ^T	
	Value	% of total ^a	Value	% of total ^a	Value	% of total ^a
Status	Complete		Draft		Complete	
Isolation source	Bovine rumen		Ovine rumen		Bovine rumen	
Genome size (bp)	3,386,314	100	3,394,947	100	4,404,886	100
DNA coding (bp)	3,064,986	90.51	3,108,180	91.55	3,954,077	89.77
DNA G + C (bp)	1,344,683	39.71	1,353,252	39.86	1,762,323	40.01
Number of replicons	4		NA		4	
DNA scaffolds	4	100	22	100	4	100
Total genes	3064	100	3104	100	3863	100
Protein coding genes	2983	97.36	2996	96.52	3739	96.79
RNA genes	60	1.96	55	1.78	68	1.75
rRNA operons	4		4		6	
tRNA genes	48	1.57	46	1.49	50	1.29
Pseudo genes	17	0.56	49		54	1.39
Genes in internal clusters	160	5.22	211	6.82	327	8.43
Genes with function prediction	2225	72.62	2314	74.55	2505	64.85
Genes assigned to COGs	1842	61.34	1861	60.17	2075	53.49
Genes with Pfam domains	2350	78.26	2407	77.82	2784	71.77
Genes with signal peptides	148	4.93	137	4.43	269	6.93
Genes with transmembrane helices	881	29.34	847	27.38	1061	27.35
CRISPR repeats	2		NA		NA	
Reference	This report		[32]		[29]	

^aThe total is based on either the size of the genome in base pairs or the total number of genes or protein-coding genes in the annotated genome. ^bIndicates draft genome sequence

amylase *amy13A* (bpr_I1087), arabinogalactan endo-1,4- β -galactosidase *agn53A* (bpr_I2041), carbohydrate esterase family 12 *est12B* (bpr_I1204), endo-1,3(4)- β -glucanase *lic16A* (bpr_I2326), pectate lyase *pel1A* (bpr_I2372), pectin methylesterase *pme8B* (bpr_I2473), xylosidase/arabinofuranosidase *xsa43J* (bpr_I2935), endo-1,4- β -xylanase *xyn10B* (bpr_I0026), and the cell wall binding domain-containing protein (bpr_I0264). These proteins contain multiple cell wall binding repeat domains (CW-binding domain, Pfam01473) at their C-termini that are predicted to anchor the protein to the peptidoglycan cell membrane. Among these secreted polysaccharidases, some contain single or combinations of catalytic activities: GH10 (endo-1, 4- β -xylanase, *xyn10B*), GH43 (xylosidase/arabinofuranosidase, *xsa43J*), PL1 (pectate lyase, *pel1A*), CE8 and PL9 (pectin methylesterase, *pme8B*) [45, 46]. Neither MB2003 nor JK615^T contain any genes encoding CW-binding domains and are thus markedly different from B316^T.

A curious feature of MB2003 was the presence of a single large (983 aa) carbohydrate binding protein (CBP, bhn_I1848), also present in JK615^T

(EJ23DRAFT_00192). The domain structures of bhn_I1848 and EJ23DRAFT_00192 are unusual, containing six CBM6 (Pfam03422) domains towards the N-terminus and a single C-terminal CBM2a (Pfam00553) domain. In contrast, B316^T encodes two CBPs (bpr_I0736 and bpr_I1599) where both contain two CBM2a domains, and bpr_I1599 also contains two CBM6 domains [29]. CBM6 non-catalytic modules characteristically bind xylose and are associated with xylanase activity with ligand specificity for xylan [47, 48]. CBM2 domains, are divided into two sub-families: 2a, that bind to crystalline cellulose even when associated with xylanases [49], and 2b, that bind to xylan [50]. Recent studies have shown that in discrete regions of plant cell walls, initial enzymatic attack of pectin increases the access of CBMs to cellulose [51], effectively loosening the polysaccharide interactions to expose the xylan and xyloglucan substrates [52, 53]. This initial stage in enzymatic saccharification of plant cell walls termed amorphogenesis [54], and is a possible role for such CBPs containing multiple non-catalytic domains. In the rumen, MB2003, B316^T and JK615^T may secrete these non-catalytic CBPs synergistically with

Table 8 Comparison of MB2003, JK615^T and B316^T protein coding gene percentages to COG functional categories

Code	% of total ^a			Description
	MB2003	JK615 ^T	B316 ^T	
J	9.52	9.33	8.96	Translation
A				RNA processing and modification
K	7.31	7.59	7.30	Transcription
L	4.32	4.59	4.63	Replication, recombination and repair
B				Chromatin structure and dynamics
D	1.57	1.60	1.44	Cell cycle control, mitosis and meiosis
V	3.19	2.90	3.19	Defense mechanisms
T	6.82	6.72	7.47	Signal transduction mechanisms
M	7.61	7.45	8.52	Cell wall/membrane biogenesis
N	2.99	3.29	2.75	Cell motility
U	1.08	1.26	1.14	Intracellular trafficking and secretion
O	3.83	3.63	3.89	Posttranslational modification, protein turnover, chaperones
C	3.39	3.63	3.72	Energy production and conversion
G	11.93	11.99	12.15	Carbohydrate transport and metabolism
E	8.69	8.85	7.91	Amino acid transport and metabolism
F	3.93	3.82	3.98	Nucleotide transport and metabolism
H	3.88	3.77	3.23	Coenzyme transport and metabolism
I	3.53	3.19	2.80	Lipid transport and metabolism
P	3.88	4.06	2.75	Inorganic ion transport and metabolism
Q	0.79	0.68	0.79	Secondary metabolites biosynthesis, transport and catabolism
R	7.76	6.91	7.43	General function prediction only
S	3.58	3.53	4.11	Function unknown
-	40.33	39.83	46.51	Not in COGs

^aThe percentage is based on the total number of protein coding genes in the genome

polysaccharide-active enzymes as a mechanism to disrupt the interface between polysaccharides to enhance the rate and extent of plant cell wall degradation.

Conclusion

The *B. hungatei* MB2003 genome sequence adds valuable information regarding the polysaccharide-degrading potential present in the genus *Butyrivibrio*. Genomic comparisons revealed that *B. hungatei* MB2003 shows a high level of similarity with *B. hungatei* JK615^T and *B. proteoclasticus* B316^T type strains, including genes involved in production of butyrate, formate, acetate and lactate. While MB2003 and JK615^T encode a large repertoire of enzymes predicted to metabolize insoluble polysaccharides such as xylan and pectin, they are unable to grow on these substrates and instead appear to be equipped to utilize mainly oligo- and mono-saccharides as substrates for growth. Although MB2003




Fig. 6 Genome synteny analysis. Alignment of the *B. hungatei* MB2003 genome against the draft genome of *B. hungatei* JK615^T (a) and the complete genome of *B. proteoclasticus* B316^T (b). Whenever the two sequences agree, a colored line or dot is plotted. If the two sequences were perfectly identical, a single line would go from the bottom left to the top right. Units displayed in base-pairs. Color codes: blue, forward sequence, red, reverse sequence

has similar phenotypic characteristics and occupies the same habitat as other *Butyrivibrio* species, its genome encodes fewer extracellular polysaccharide degrading enzymes, in particular, those that contain multiple cell wall binding repeat domains. The overall genome similarities, metabolic versatility and differences in the abundance of CAZymes observed in *B. proteoclasticus* and *B. hungatei* offers a new view of the genes required for polysaccharide degradation in the rumen. MB2003 appears to occupy a ruminal niche as a secondary degrader of oligosaccharides, in order to coexist with fibre-degrading organisms in this dynamic and competitive environment.

Additional file

Additional file 1: Table S1. Associated MIGS record for *B. hungatei* MB2003, which links to the SIGS supplementary content website. (DOCX 17 kb)

Table 9 Genes encoding predicted polysaccharide degrading enzymes in the MB2003 genome

Locus tag	Name	Annotation	Size (aa)	CAZy ^a	Binding domains
bhn_I2518	<i>bga2A</i>	β-galactosidase ^b	1034	GH2	
bhn_I0827	<i>bga2C</i>	β-galactosidase ^b	714	GH2	
bhn_I1587	<i>bga2B</i>	β-galactosidase ^b	825	GH2	
bhn_I0200	<i>gh2B</i>	glycoside hydrolase family 2 ^b	641	GH2	
bhn_I1127	<i>gh2A</i>	glycoside hydrolase family 2 ^b	912	GH2	
bhn_I1849	<i>gh2C</i>	glycoside hydrolase family 2 ^b	776	GH2	
bhn_III062	<i>bgl3A</i>	β-glucosidase ^b	803	GH3	
bhn_I0707	<i>bgl3B</i>	β-glucosidase ^b	808	GH3	
bhn_I0180	<i>bgl3C</i>	β-glucosidase ^b	671	GH3	
bhn_I0706	<i>bgl3D</i>	β-glucosidase^b	982	GH3	C-terminal TMH
bhn_I0189	<i>xyl3A</i>	β-xylosidase ^b	707	GH3	
bhn_I1640	<i>bhx3A</i>	β-N-acetylhexosaminidase^b	427	GH3	
bhn_I1693	<i>cel5C</i>	endo-1,4-β-glucanase^b	543	GH5	CBM2a
bhn_I0165	<i>cel5A</i>	endo-1,4-β-glucanase/xylanase^b	417	GH5	
bhn_I1756	<i>xyn8A</i>	reducing end xylose-releasing exo-oligoxylanase ^b	383	GH8	
bhn_I0834	<i>cel9B</i>	cellodextrinase ^b	552	GH9	CelD
bhn_I0568	<i>xyn10B</i>	endo-1,4-β-xylanase^b	425	GH10	
bhn_I0169	<i>xyn10A</i>	endo-1,4-β-xylanase^b	451	GH10	
bhn_I1458	<i>glgB2</i>	1,4-α-glucan branching enzyme ^b	824	GH13	CBM48
bhn_I0053	<i>glgB1</i>	1,4-α-glucan branching enzyme ^b	663	GH13	CBM48
bhn_I2702	<i>amy13A</i>	α-amylase ^b	697	GH13	CBM34
bhn_I0634	<i>amy13B</i>	α-amylase^b	536	GH13	
bhn_I1680	<i>amy13C</i>	α-amylase ^b	434	GH13	
bhn_I0669	<i>amy13D</i>	α-amylase ^b	511	GH13	
bhn_I1153	<i>glgX1</i>	glycogen debranching enzyme ^b	726	GH13	CBM48
bhn_I1315	<i>glgX2</i>	glycogen debranching enzyme ^b	648	GH13	
bhn_I0652	<i>suc13P</i>	sucrose phosphorylase ^b	553	GH13	
bhn_I2526	<i>chi18A</i>	chitinase^b	567	GH18	
bhn_I1254	<i>lyc25A</i>	lysozyme^b	362	GH25	
bhn_III074	<i>lyc25B</i>	lysozyme^b	515	GH25	
bhn_I0191	<i>lyc25C</i>	lysozyme^b	561	GH25	
bhn_I1763	<i>lyc25D</i>	lysozyme^b	242	GH25	
bhn_I0527	<i>lyc25E</i>	lysozyme ^b	1213	GH25	Big2 (x2)
bhn_I1287	<i>aga27A</i>	α-galactosidase ^b	577	GH27	
bhn_I0082	<i>gh27A</i>	glycoside hydrolase family 27 ^b	442	GH27	
bhn_I1952	<i>pgl28A</i>	polygalacturonase ^b	531	GH28	
bhn_I2679	<i>pgl28B</i>	polygalacturonase ^b	519	GH28	
bhn_I1087	<i>fuc29A</i>	α-L-fucosidase ^b	475	GH29	
bhn_I2734	<i>gh30A</i>	glycoside hydrolase family 30^b	575	GH30	
bhn_I1581	<i>gh31A</i>	glycoside hydrolase family 31 ^b	756	GH31	
bhn_I2191	<i>gh31C</i>	glycoside hydrolase family 31 ^b	674	GH31	
bhn_I0283	<i>gh31B</i>	glycoside hydrolase family 31 ^b	635	GH31	
bhn_I0582	<i>scr32A</i>	sucrose-6-phosphate hydrolase ^b	493	GH32	
bhn_I0826	<i>bga35A</i>	β-galactosidase ^b	622	GH35	

Table 9 Genes encoding predicted polysaccharide degrading enzymes in the MB2003 genome (*Continued*)

Locus tag	Name	Annotation	Size (aa)	CAZy ^a	Binding domains
bhn_I1817	<i>bga35B</i>	β-galactosidase ^b	735	GH35	
bhn_I0644	<i>aga36A</i>	α-galactosidase ^b	782	GH36	
bhn_I1583	<i>aga36B</i>	α-galactosidase ^b	620	GH36	
bhn_I1945	<i>aga36C</i>	α-galactosidase ^b	730	GH36	
bhn_I0086	<i>man38A</i>	α-mannosidase ^b	1053	GH38	
bhn_III010	<i>bga42A</i>	β-galactosidase ^b	673	GH42	
bhn_I0167	<i>xsa43A</i>	xylosidase/arabinofuranosidase^b	543	GH43	CBM6
bhn_I0981	<i>xsa43B</i>	xylosidase/arabinofuranosidase ^b	301	GH43	
bhn_I2037	<i>xsa43C</i>	xylosidase/arabinofuranosidase ^b	302	GH43	
bhn_I2111	<i>xsa43D</i>	xylosidase/arabinofuranosidase ^b	517	GH43	
bhn_I2735	<i>xsa43E</i>	xylosidase/arabinofuranosidase ^b	352	GH43	
bhn_I0032	<i>xsa43G</i>	xylosidase/arabinofuranosidase ^b	312	GH43	
bhn_I0164	<i>xsa43F</i>	xylosidase/arabinofuranosidase and esterase ^b	925	GH43	
bhn_I1509	<i>arf51C</i>	α-L-arabinofuranosidase ^b	630	GH51	
bhn_I2228	<i>arf51A</i>	α-L-arabinofuranosidase ^b	502	GH51	
bhn_I0010	<i>arf51B</i>	α-L-arabinofuranosidase ^b	504	GH51	
bhn_I0670	<i>agn53A</i>	arabinogalactan endo-1,4-β-galactosidase^b	439	GH53	
bhn_I0183	<i>agu67A</i>	α-D-glucuronidase ^b	662	GH67	
bhn_I2177	<i>mal77A</i>	4-α-glucanotransferase^b	506	GH77	
bhn_I0697	<i>ugl88A</i>	unsaturated glucuronyl hydrolase ^b	385	GH88	
bhn_I2381	<i>ugl88B</i>	unsaturated glucuronyl hydrolase ^b	383	GH88	
bhn_I2196	<i>cbp94A</i>	cellobiose phosphorylase ^b	814	GH94	
bhn_I1582	<i>gh95A</i>	glycoside hydrolase family 95 ^b	734	GH95	
bhn_I2548	<i>gh105A</i>	unsaturated rhamnogalacturonyl hydrolase ^b	349	GH105	
bhn_I0090	<i>gh105B</i>	unsaturated rhamnogalacturonyl hydrolase ^b	363	GH105	
bhn_I2549	<i>gnpA</i>	D-galactosyl-β-1-4-L-rhamnose phosphorylase ^b	722	GH112	
bhn_I0185	<i>gh115A</i>	α-glucuronidase ^b	947	GH115	
bhn_I1083	<i>xyl120A</i>	xylosidase ^b	861	GH120	
bhn_I1738	<i>xyl120B</i>	xylosidase ^b	664	GH120	
bhn_III076	<i>est1A</i>	feruloyl esterase^b	351	CE1	
bhn_I1244	<i>est2A</i>	acetyl-xylan esterase ^b	372	CE2	
bhn_III070	<i>est4A</i>	polysaccharide deacetylase ^b	207	CE4	
bhn_I0843	<i>est4C</i>	polysaccharide deacetylase^b	280	CE4	
bhn_I0666	<i>nagA</i>	N-acetylglucosamine-6-phosphate deacetylase ^b	371	CE9	
bhn_I1609	<i>est12A</i>	carbohydrate esterase family 12 ^b	584	CE12	
bhn_I1927	<i>est12B</i>	carbohydrate esterase family 12 ^b	244	CE12	
bhn_I1926	<i>pl11A</i>	polysaccharide lyase ^b	746	PL11	
bhn_I0657	<i>glgP1</i>	glycogen phosphorylase ^b	769	GT35	
bhn_I2673	<i>glgP2</i>	glycogen phosphorylase ^b	824	GT35	
bhn_I1848	-	carbohydrate binding protein^b	983		CBM2a (×1), CBM6 (×6)

^aCAZy descriptions and classifications compiled from the CAZy database [68]. ^bIndicates homologues in the *B. hungatei* JK615^T draft genome. Genes encoding predicted secreted polysaccharide degrading enzymes are in bold

Abbreviations

Bp: Base pair(s); CAZymes: Carbohydrate-Active enZymes; CBMs: Carbohydrate-Binding Module(s); CEs: Carbohydrate Esterase(s); Ech: *Escherichia coli* hydrogenase-3-type hydrogenase; Eha: Energy-converting hydrogenase A; Ehb: Energy-converting hydrogenase B; EMP: Embden-Meyerhof-Parnas; ETP: Electron transport phosphorylation; GHs: Glycoside Hydrolase(s); GTs: Glycosyl Transferase(s); Hyd: Ferredoxin hydrogenase; Mbh: Membrane-bound hydrogenase; Mvh/Hdr: Methyl viologen hydrogenase/heterodisulfide reductase; Pfl: Pyruvate formate lyase; PLs: Polysaccharide Lyase(s); Por: Pyruvate ferredoxin oxidoreductase; Rnf: *Rhodobacter* nitrogen fixation

Acknowledgements

The MB2003 genome sequencing project was funded by the New Zealand Ministry of Business, Innovation and Employment New Economy Research Fund programme: Accessing the uncultured rumen microbiome, contract number C10X0803. Electron microscopy was conducted with the assistance of the Manawātū Microscopy and Imaging Centre at Massey University, Palmerston North, New Zealand.

Authors' contributions

NP, WJK, GTA conceived and designed the experiments. NP performed the sequencing and assembly experiments. NP, WJK, SCL, EA performed the genome annotation and comparative studies. NP performed the bacterial growth studies and polysaccharide utilization profiling. NP, WJK, GTA wrote the manuscript. All authors commented on the manuscript before submission. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 25 July 2017 Accepted: 23 November 2017

Published online: 04 December 2017

References

- Henderson G, Cox F, Ganesh S, Jonker A, Young W, Collaborators GRC, Janssen PH. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci Rep*. 2015;5(14567):1–13.
- Bryant MP, Small N. The anaerobic monotrichous butyric acid-producing curved rod-shaped bacteria of the rumen. *J Bacteriol*. 1956;72(1):16–21.
- Hungate RE. The rumen and its microbes. New York: Academic Press; 1966.
- Willems A, Collins MD. *Butyrivibrio*. Bryant and Small 1956b, 18, emend. Moore, Johnson and Holdeman 1976, 241^{AL}. In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*. Volume 3. 2nd ed. New York: Springer; 2009. p. 927–37.
- Hespell RB. The genera *Butyrivibrio*, *Lachnospira*, and *Roseburia*. In: Balows A, Trüper HG, Dworkin M, Harder W, Schleifer K-H, editors. *The prokaryotes: a handbook on the biology of bacteria: ecophysiology, isolation, identification, applications*. New York: Springer; 1992. p. 2023–33.
- Moon CD, Pacheco DM, Kelly WJ, Leahy SC, Li D, Kopecny J, Attwood GT. Reclassification of *Clostridium proteoclasticum* as *Butyrivibrio proteoclasticus* comb. nov., a butyrate-producing ruminal bacterium. *Int J Syst Evol Microbiol*. 2008;58(9):2041–5.
- Hespell RB, Cotta MA. Degradation and utilization by *Butyrivibrio fibrisolvens* H17c of xylans with different chemical and physical properties. *Appl Environ Microbiol*. 1995;61(8):3042–50.
- Attwood GT, Reilly K. Characterization of proteolytic activities of rumen bacterial isolates from forage-fed cattle. *J Appl Bacteriol*. 1996;81(5):545–52.
- Diez-Gonzalez F, Bond DR, Jennings E, Russell JB. Alternative schemes of butyrate production in *Butyrivibrio fibrisolvens* and their relationship to acetate utilization, lactate production, and phylogeny. *Arch Microbiol*. 1999; 171(5):324–30.
- Paillard D, McKain N, Rincon MT, Shingfield KJ, Givens DJ, Wallace RJ. Quantification of ruminal *Clostridium proteoclasticum* by real-time PCR using a molecular beacon approach. *J Appl Microbiol*. 2007;103(4):1251–61.
- Shane BS, Gouws L, Kistner A. Cellulolytic bacteria occurring in the rumen of sheep conditioned to low-protein teff hay. *J Gen Microbiol*. 1969;55(3):445–57.
- Hazlewood GP, Theodorou MK, Hutchings A, Jordan DJ, Galfre G. Preparation and characterization of monoclonal antibodies to a *Butyrivibrio* sp. and their potential use in the identification of rumen *Butyrivibrios*, using an enzyme-linked immunosorbent assay. *J Gen Microbiol*. 1986;132(1):43–52.
- Mannarelli BM. Deoxyribonucleic acid relatedness among strains of the species *Butyrivibrio fibrisolvens*. *Int J Syst Bacteriol*. 1988;38(4):340–7.
- Mannarelli BM, Stack RJ, Lee D, Ericsson L. Taxonomic relatedness of *Butyrivibrio*, *Lachnospira*, *Roseburia*, and *Eubacterium* species as determined by DNA hybridization and extracellular-polysaccharide analysis. *Int J Syst Bacteriol*. 1990;40(4):370–8.
- Forster R, Teather R, Gong J, Deng S-J. 16S rDNA analysis of *Butyrivibrio fibrisolvens*: phylogenetic position and relation to butyrate-producing anaerobic bacteria from the rumen of white-tailed deer. *Lett Appl Microbiol*. 1996;23(4):218–22.
- Willems A, Amat-Marco M, Collins MD. Phylogenetic analysis of *Butyrivibrio* strains reveals three distinct groups of species within the *Clostridium subphylum* of the gram-positive bacteria. *Int J Syst Bacteriol*. 1996;46(1):195–9.
- Forster RJ, Gong J, Teather RM. Group-specific 16S rRNA hybridization probes for determinative and community structure studies of *Butyrivibrio fibrisolvens* in the rumen. *Appl Environ Microbiol*. 1997;63(4):1256–60.
- Van Gylswyk NO, Hippe H, Rainey FA. *Pseudobutyrvibrio ruminis* sp. nov., a butyrate-producing bacterium from the rumen that closely resembles *Butyrivibrio fibrisolvens* in phenotype. *Int J Syst Bacteriol*. 1996; 46(2):559–63.
- Kopecny J, Zorec M, Mrazek J, Kobayashi Y, Marinsek-Logar R. *Butyrivibrio hungatei* sp. nov. and *Pseudobutyrvibrio xylanivorans* sp. nov., butyrate-producing bacteria from the rumen. *Int J Syst Evol Microbiol*. 2003;53(1):201–9.
- Noel S. Cultivation and community composition analysis of plant-adherent rumen bacteria. PhD thesis. Massey University, Institute of Molecular Biosciences; 2013.
- Pavevich N. Genome sequencing of rumen bacteria involved in lignocellulose digestion. MSc thesis: Massey University, Institute of Molecular Biosciences; 2011.
- Kenters N, Henderson G, Jeyanathan J, Kittelmann S, Janssen PH. Isolation of previously uncultured rumen bacteria by dilution to extinction using a new liquid culture medium. *J Microbiol Methods*. 2011;84(1):52–60.
- Richardson A, Calder A, Stewart C, Smith A. Simultaneous determination of volatile and non-volatile acidic fermentation products of anaerobes by capillary gas chromatography. *Lett Appl Microbiol*. 1989;9(1):5–8.
- Saito H, Miura K-I. Preparation of transforming deoxyribonucleic acid by phenol treatment. *Biochim Biophys Acta*. 1963;72:619–29.
- Staden R, Beal KF, Bonfield JK. The staden package. In: Misener S, Krawetz SA, editors. *Bioinformatics methods and protocols*, vol. Volume 132. New York: Humana Press; 1999. p. 115–30.
- Goecks J, Nekrutenko A, Taylor J. Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. *Genome Biol*. 2010;11(8):R86.
- Zerbino DR, Birney E. Velvet: algorithms for *de novo* short read assembly using *De Bruijn* graphs. *Genome Res*. 2008;18(5):821–9.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 2012;28(12):1647–9.
- Kelly WJ, Leahy SC, Altermann E, Yeoman CJ, Dunne JC, Kong Z, Pacheco DM, Li D, Noel SJ, Moon CD, Cookson AL, Attwood GT. The glycobiome of the rumen bacterium *Butyrivibrio proteoclasticus* B316^T highlights adaptation to a polysaccharide-rich environment. *PLoS One*. 2010;5(8):e11942.
- Benson DA, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids Res*. 2015;43(D1):D30–5.
- Pavevich N. Comparative genomics of *Butyrivibrio* and *Pseudobutyrvibrio* from the rumen. PhD thesis. Massey University, Institute of Fundamental Sciences; 2016.
- Kyrpides NC, Hugenholtz P, Eisen JA, Woyke T, Göker M, Parker CT, Amann R, Beck BJ, Chain PS, Chun J. Genomic encyclopedia of bacteria and archaea: sequencing a myriad of type strains. *PLoS Biol*. 2014;12(8): e1001920.

33. Harrison PW, Lower RP, Kim NK, Young JP. Introducing the bacterial 'chromid': not a chromosome, not a plasmid. *Trends Microbiol.* 2010; 18(4):141–8.
34. Welte C, Krätzer C, Deppenmeier U. Involvement of Ech hydrogenase in energy conservation of *Methanosarcina mazei*. *FEBS J.* 2010;277(16): 3396–403.
35. Biegel E, Müller V. Bacterial Na⁺-translocating ferredoxin: NAD⁺ oxidoreductase. *Proc Natl Acad Sci.* 2010;107(42):18138–42.
36. Buckel W, Thauer RK. Energy conservation via electron bifurcating ferredoxin reduction and proton/Na⁺ translocating ferredoxin oxidation. *Biochim Biophys Acta Bioenerg.* 2013;1827(2):94–113.
37. Schlegel K, Welte C, Deppenmeier U, Müller V. Electron transport during acetoclastic methanogenesis by *Methanosarcina acetivorans* involves a sodium-translocating Rnf complex. *FEBS J.* 2012;279(24):4444–52.
38. Tremblay P-L, Zhang T, Dar SA, Leang C, Lovley DR. The Rnf complex of *Clostridium ljungdahlii* is a proton-translocating ferredoxin: NAD⁺ oxidoreductase essential for autotrophic growth. *MBio.* 2013;4(1): e00406–12.
39. Gutekunst K, Chen X, Schreiber K, Kaspar U, Makam S, Appel J. The bidirectional NiFe-hydrogenase in *Synechocystis* sp. PCC 6803 is reduced by flavodoxin and ferredoxin and is essential under mixotrophic, nitrate-limiting conditions. *J Biol Chem.* 2014;289(4):1930–7.
40. Pereira IAC, Ramos A, Grein F, Marques M, Da Silva S, Venceslau S. A comparative genomic analysis of energy metabolism in sulfate reducing bacteria and archaea. *Front Microbiol.* 2011;2(69).
41. Weghoff MC, Bertsch J, Müller V. A novel mode of lactate metabolism in strictly anaerobic bacteria. *Environ Microbiol.* 2015;17(3):670–7.
42. Hackmann TJ, Firkins JL. Electron transport phosphorylation in rumen butyrvibrions: unprecedented ATP yield for glucose fermentation to butyrate. *Front Microbiol.* 2015;6(622).
43. Cooper R. Metabolism of methylglyoxal in microorganisms. *Annu Rev Microbiol.* 1984;38(1):49–68.
44. Leahy SC, Kelly WJ, Altermann E, Ronimus RS, Yeoman CJ, Pacheco DM, Li D, Kong Z, McTavish S, Sang C. The genome sequence of the rumen methanogen *Methanobrevibacter ruminantium* reveals new possibilities for controlling ruminant methane emissions. *PLoS One.* 2010;5(11):e8926.
45. Dunne JC, Li D, Kelly WJ, Leahy SC, Bond JJ, Attwood GT, Jordan TW. Extracellular polysaccharide-degrading proteome of *Butyrivibrio proteoclasticus*. *J Proteome Res.* 2011;11(1):131–42.
46. Bond JJ, Dunne JC, Kwan FY, Li D, Zhang K, Leahy SC, Kelly WJ, Attwood GT, Jordan TW. Carbohydrate transporting membrane proteins of the rumen bacterium, *Butyrivibrio proteoclasticus*. *J Proteome.* 2012;75(11):3138–44.
47. McCartney L, Gilbert HJ, Bolam DN, Boraston AB, Knox JP. Glycoside hydrolase carbohydrate-binding modules as molecular probes for the analysis of plant cell wall polymers. *Anal Biochem.* 2004;326(1):49–54.
48. Gilbert HJ, Knox JP, Boraston AB. Advances in understanding the molecular basis of plant cell wall polysaccharide recognition by carbohydrate-binding modules. *Curr Opin Struct Biol.* 2013;23(5):669–77.
49. Gilkes NR, Jervis E, Henrissat B, Tekant B, Miller R, Warren R, Kilburn D. The adsorption of a bacterial cellulase and its two isolated domains to crystalline cellulose. *J Biol Chem.* 1992;267(10):6743–9.
50. Simpson PJ, Bolam DN, Cooper A, Ciruela A, Hazlewood GP, Gilbert HJ, Williamson MP. A family IIb xylan-binding domain has a similar secondary structure to a homologous family IIa cellulose-binding domain but different ligand specificity. *Structure.* 1999;7(7):853–64.
51. Blake AW, McCartney L, Flint JE, Bolam DN, Boraston AB, Gilbert HJ, Knox JP. Understanding the biological rationale for the diversity of cellulose-directed carbohydrate-binding modules in prokaryotic enzymes. *J Biol Chem.* 2006; 281(39):29321–9.
52. Marcus SE, Verhertbruggen Y, Hervé C, Ordaz-Ortiz JJ, Farkas V, Pedersen HL, Willats WG, Knox JP. Pectic homogalacturonan masks abundant sets of xyloglucan epitopes in plant cell walls. *BMC Plant Biol.* 2008;8(60):1–12.
53. Hervé C, Rogowski A, Gilbert HJ, Paul Knox J. Enzymatic treatments reveal differential capacities for xylan recognition and degradation in primary and secondary plant cell walls. *Plant J.* 2009;58(3):413–22.
54. Coughlan MP. The properties of fungal and bacterial cellulases with comment on their production and application. *Biotechnol Genet Eng Rev.* 1985;3(1):39–110.
55. Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 2013;30(12):2725–9.
56. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 1985:783–91.
57. Reddy T, Thomas AD, Stamatis D, Bertsch J, Isbandi M, Jansson J, Mallajosyula J, Pagani I, Lobos EA, Kyrpides NC. The genomes OnLine database (GOLD) v. 5: a metadata management system based on a four level (meta) genome project classification. *Nucleic Acids Res.* 2015;43(D1): D1099–106.
58. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol.* 2008;26(5):541–7.
59. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, bacteria, and Eucarya. *Proc Natl Acad Sci.* 1990;87(12):4576–9.
60. Rainey F. Phylum SK-H, XIII. *Firmicutes* gibbons and Murray 1978, 5 (*Fimacutes* [sic] gibbons and Murray 1978, 5). In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*, vol. Volume 3. 2nd ed. New York: Springer; 2009. p. 19–1317.
61. Ludwig W, Schleifer K-H, Whitman W. Revised road map to the phylum *Firmicutes*. In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*. Volume 3. 2nd ed. New York: Springer; 2009. p. 1–13.
62. Rainey F. Class II. *Clostridia* class. nov. In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*, vol. Volume 3. 2nd ed. New York: Springer; 2009. p. 736.
63. Rainey F. Order I. *Clostridiales* Prévot 1953. In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*, vol. Volume 3. 2nd ed. New York: Springer; 2009. p. 736.
64. Rainey F. Family V. *Lachnospiraceae* fam. nov. In: De Vos P, Garrity G, Jones D, Krieg N, Ludwig W, Rainey F, Schleifer K-H, Whitman W, editors. *Bergey's manual of systematic bacteriology*, vol. Volume 3. 2nd ed. New York: Springer; 2009. p. 921–68.
65. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT. Gene ontology: tool for the unification of biology. *Nat Genet.* 2000;25(1):25–9.
66. Jensen LJ, Friis C, Ussery DW. Three views of microbial genomes. *Res Microbiol.* 1999;150(9):773–7.
67. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST. PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 1997;25(17):3389–402.
68. Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. The carbohydrate-active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* 2009;37(D1):D233–8.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

