
The complete nucleotide sequence of the rat 18S ribosomal RNA gene and comparison with the respective yeast and frog genes

Richard Torczynski, Arthur P. Bollon and Motohiro Fuke

Department of Molecular Genetics, Wadley Institutes of Molecular Medicine, 900 Harry Hines Boulevard, Dallas, TX 75235, USA

Received 21 March 1983; Revised and Accepted 14 June 1983

ABSTRACT

The complete nucleotide sequence of the rat 18S ribosomal RNA gene has been determined. A comparison of the rat 18S ribosomal RNA gene sequence with the known sequences of yeast and frog revealed three conserved (stable) regions, two unstable regions, and three large inserts. (A,T)→(G,C) changes were more frequent than (G,C)→(A,T) changes for three comparisons (yeast→frog, frog→rat, and yeast→rat). GC pairs were inserted preferentially over AT pairs for the same three comparisons. These two factors contribute to the progressively higher GC content of 18S ribosomal RNA of yeast, frog, and rat.

INTRODUCTION

The ribosomal RNA (rRNA) genes are essential genes existing in all life forms from bacteria to man and they have been subjected to extensive analysis. Nucleotide sequence comparison of rRNA genes of different organisms should provide insight into evolutionary trends. There are two high molecular weight RNA components (18S rRNA and 25-28S rRNA) in eukaryotic ribosomes, and 18S rRNA sequences are known to be more highly conserved than the other high molecular weight rRNA sequences (1, 2).

Complete nucleotide sequences of 18S rRNA genes of *Saccharomyces cerevisiae* (3) and *Xenopus laevis* (4) have been reported recently. The present study reveals the first complete nucleotide sequence of a mammalian 18S rRNA gene (rDNA) and analyzes the rat sequence compared with the yeast and frog sequences.

MATERIALS AND METHODS

The characterization of rat rDNA cloned in lambda bacteriophages and the recloning of the rDNA in plasmid pBR322 were previously described (5). Plasmid pDF40 was used for sequencing the 3'-terminal region of the rat 18S rDNA (6) and plasmid pDF15 was used for sequencing the 5'-terminal region




Figure 1. Strategy of sequencing the central portion of the rat 18S rDNA. Numbers reflect the distances from the 5'-terminus. Only the restriction sites that were utilized for sequencing are shown. Arrows indicate the directions of sequencing.

(7). Plasmid pDF8, which contains the central region of the rat 18S rDNA, was used to complete the sequencing of the 18S rDNA. Restriction endonucleases were purchased from BRL (Rockville, MD). DNA digestions with restriction endonucleases and gel-electrophoresis were performed as described previously (8, 9). DNA was sequenced by the method developed by Maxam and Gilbert (10) with minor modifications. In addition to the 5'-terminal labeling, in some cases, the 3'-termini of the fragments digested with restriction enzymes were labeled with α -³²P-dNTP and reverse transcriptase. The strategy employed for sequencing the central portion of the rat 18S rDNA is represented in Fig. 1.

RESULTS

An example of the sequencing gels is shown in Fig. 2. The region in Fig. 2 includes large T1 RNase fragment 8 (UUUUCUUAUAUCAAG) that was sequenced previously using the rat 18S rRNA (11). The total length of the rat 18S rDNA is 1869 bp and the nucleotide sequence determined is shown in Fig. 3 together with the yeast and frog sequences for comparison.

All the 19 large T1 RNase fragments, of which sequences were previously reported (11), were located in the rat 18S rDNA sequence in Fig. 3. The nucleotide sequences determined for these RNA fragments agree with the DNA sequence except for fragment 15, the sequence of which was cited from the work of other investigators who used HeLa cell 18S rRNA (12). Fragment 15 contains a hypermodified nucleoside identified as 1-methyl-3- γ -(α -amino- α -carboxypopyl) pseudouridine (13). The nucleotide sequence of fragment 15




Figure 2. A typical sequencing gel. Numbers show the distances from the 5'-terminus (Fig. 3). The Ava II digested fragment was labeled with ATP γ P³² at the 5'-termini and secondarily digested with Hae III. The electrophoresis was performed in a 6% polyacrylamide gel.

(AC^{am}CAACACG) is conserved among yeast, frog, and rat (3, 4).

In Fig. 3, the three sequences are arranged to give the best homology and the nucleotides that are common to all the three species are in bold letters. The whole sequence was sectioned every 10 nucleotides and the numbers of the nucleotides conserved in every section were plotted in Fig. 4. The sequence comparison was done in two ways: between frog and rat (top) and among the three species (bottom). It was known that the nucleotide sequence of the 3'-terminal region is well conserved among many species (6, 14, 15). This study shows that two additional larger conserved regions are located inside of 18S rDNA (Fig. 4). The three completely conserved large regions are: region A (nucleotides 424-499 in Fig. 3, 76 nucleotides long), region B (nucleotides 600-665, 66 nucleotides long), and region C (nucleotides 1837-1885, 49 nucleotides long).

Region U1 (nucleotides 181-357 in Fig. 3, 177 nucleotides long and 42% conserved) and Region U2 (nucleotides 690-805, 116 nucleotides long and 33% conserved) were identified as two evolutionarily unstable regions which are larger than 100 nucleotides and are less than 50% conserved (Fig. 4). Salim and Maden (4) identified four variable regions from their comparison of yeast and frog sequences. Two of the four regions correspond to U1 and U2 in this study. Unstable Region U1 contains two large insertions indicated by arrows in Fig. 4. Another large insertion, 10 nucleotides long, is located from nucleotides 1053 to 1062 in Fig. 3 and is indicated by an arrow in Fig. 4.

Table 1 summarizes base composition data of the three 18S rRNA's. The total number of nucleotides increases in the order of yeast → frog → rat, and the difference between rat and frog (44 nucleotides) is larger than the difference between yeast and frog (36 nucleotides). The numbers of U and A in 18S rRNA decrease in the order of yeast → frog → rat and the numbers of C and G increase. Hence, GC content in 18S rRNA increases in the order of yeast → frog → rat. The GC content difference between rat and frog (1.8%) is much smaller than the difference between yeast and frog (8.8%).

A more detailed analysis of base changes is shown in Table 2. From the data in Fig. 3, the number of base changes from A or T to G or C was counted (line [a] in Table 2) and the number of opposite base changes from G or C to A or T was also counted (line [b]). There is a clear preference for (A,T) → (G,C) changes over (G,C) → (A,T) changes for the three comparisons (yeast → frog, frog → rat, and yeast → rat, line [c]). Four

Yeast	-10	-1	1	20	40	60	80
Frog	TTAAGATAGT	TATCTGGTTGATCCTGCCAGTAGTCATATGCTTGCTCAAAGATTAAAGCCATGCATGTC	TAAGTATAAGCA	TAATTTACAGTGAACCTGC			
Rat	GAAAGGTGGC	TACCTGGTTGATCCTGCCAGTAG-CATATGCTTGCTCAAAGATTAAAGCCATGCATGTC	TAAGTACGCAC	GGCCGGTACAGTGAACCTGC			
	TCGC6CTCT	TACCTGGTTGATCCTGCCAGTAG-CATATGCTTGCTCAAAGATTAAAGCCATGCATGTC	TAAGTACGCAC	GGCCGGTACAGTGAACCTGC			
Yeast	100	120	140	160	180		
Frog	GAATGGCTCATTTAAATCAGTTATCGTTTATTTGA	---TAGTTCC-TT TACTACATGGTATAACCGTGGTAAITCTAGAGCTAAATACATGCTTAAAAATCTC					
Rat	GAATGGCTCATTTAAATCAGTTATGGTTCCCTTTGA	---TGCTCCATCTGTACTTGG-ATAAATCGTGGTAAITCTAGAGCTAAATACATGCTCCGACGAGCGC					
	GAATGGCTCATTTAAATCAGTTATGGTTCCCTTTGTCGCTCGCTCC	-TCTCCTACTTGG-ATAAATCGTGGTAAITCTAGAGCTAAATACATGCTCCGACGGGCGGC					
Yeast	200	220	240	260	280		
Frog	-GACCC	---TT---T---GGAAGAGATGTTATTTAGATAAAAAATCAAT	---GTC	---	---	---	
Rat	TGACCCCC	---AGGATGCGTGCATTTATCAGACCAAAA	-CCAATCCGG	-GGCCCC	---GC	---G-CC	
	TGACCCCCCTTCCCGTGGGGGGACGCGTGCATTTATCAGATCAAAA	-CCAACCCGGTCAAGCCCCCTCCCGTCCGGCCGGGGGTCCGGCGCCGGACGGC					
Yeast	300	320	340	360	380		
Frog	TTTGATGATTCATAATAACTTTTCG	--AATCGCATGGCCCTT	-GTGCTGGCGATGGTTCAATTC	CAAAATTC	TGCCCCTATCAACTTTTCGATGGT	TAGGATAGTG	
Rat	TTTGGTACTCTAGATAACCTCGGGCCGATCGACGCTCC	-CCGTGACGGCGACGATACATTCGGATGCTGGCCCTATCAACTTTTCGATGGT	ACTTTCTGC				
	TTTGGTACTCTAGATAACCTCGGGCCGATCGACGCCCCCTCCGTGGCGGGCAGACCCCAATTCGAACGCTCGCCCTATCAACTTTTCGATGGT	AGTCTGCGCGT					
Yeast	400	420	440	460	480		
Frog	GCCTACCATGGTTTCAACGGGTAAACGGGGMAATAAGGGTT	CGATTCCGGAGGGAGCCCTGAGAAACGGCTACACACATCC	AAGGAAGGCAGCAGGGCGCGCA				
Rat	GCCTACCATGGTGACCACGGGTAAACGGGGMAATCAGGGTT	CGATTCCGGAGAGGGAGCCCTGAGAAACGGCTACACACATCC	AAGGAAGGCAGCAGGGCGCGCA				
	GCCTACCATGGTGACCACGGGTAAACGGGGMAATCAGGGTT	CGATTCCGGAGAGGGAGCCCTGAGAAACGGCTACACACATCC	AAGGAAGGCAGCAGGGCGCGCA				
Yeast	500	520	540	560	580		
Frog	AATTACCACATCC	AATTCAGGGAGGTAGTGACAAATAAATACGATACAGGGCCCATCCG	-GGTCTTGTAAITGGAAATGAGTACAAATGTAATACCTTAA				
Rat	AATTACCACATCCCGACGGGGAGGTAGTGACAAATAAATACGATACAGGGCT	TTTCGAGGCCCTGTAATGGAAATGAGTACACTTTAAATCCCTTTAA					
	AATTACCACATCCCGACCCCGGGAGGTAGTGACAAATAAATACGATACAGGGACT	TTTCGAGGCCCTGTAATGGAAATGAGTACACTTTAAATCCCTTTAA					
Yeast	600	620	640	660	680		
Frog	CGAGAACAAITGGAGGGCAAGTCTGGTGCCAGCAGCCGCGGT	AAITCCAGCTCCAAATAGCGTATATTTAAAGTTGTTGACGTTAAAAAGCTCGTAGTTGA					
Rat	CGAGGATCTATTGGAGGGCAAGTCTGGTGCCAGCAGCCGCGGT	AAITCCAGCTCCAAATAGCGTATATTTAAAGTTGTTGACGTTAAAAAGCTCGTAGTTGG					
	CGAGGATCCATTGGAGGGCAAGTCTGGTGCCAGCAGCCGCGGT	AAITCCAGCTCCAAATAGCGTATATTTAAAGTTGTTGACGTTAAAAAGCTCGTAGTTGG					
Yeast	700	720	740	760	780		
Frog	ACTTTGGGCC	----GGTTGGCCGGTCCGATTTTTCGTGTAG	-TGGATTTCCAAAGGGGCCCTTCCCTCTCGGCTAAACCTT	TGAGTCCCTTGTGGCTTTG			
Rat	ATCTTGGGATCGAGCTGGCGGTCGCCCGAGGGCG	-GTACCCGCTGTCACAGCCCT	-G-CCTCTCGGGGCC	TCCCGATGCTCTTGACTGAGTGTCCC			
	ATCTTGGGAGGGGGCGGGTCCCGCCGCGAGT	CACCCGCCCCCTTG	-CCTCTCGGGCCGCCCTCGATGCTCTTAGCTGAGTGTCCC				

Yeast	800	820	840	860	880
Frog	GGGAACAGGAC -- -TTTAC TTTGAAA AATAGAGTGTCA AAGAGGC -- GTATTGCTCA ATATATTCAGCATGGAA TAATAGAA TAGAGACGTTTGGT				
Rat	G-GGGCCCGAAGCGTTTACTTTGAAA AATTAGAGTGTCC AAGCAGGC - GCGTGCGCTGGA TACTTCAGCTAGGAATAATGGA ATAGSAC - TCCGGT				
	900	920	940	960	980
Yeast	TCTATTTGTTGGTTCTAGG ACCATCGTAATGATTAA TAGGGACGGTCTGGGGGCATCGGTATTC AATTG - TCGAGGTGAAA TTC TTGGATTTATTTAAAG				
Frog	TCTATTTGTTGGTTCTCGGA AC TGGGCCATGATTAA GAGGGACGGCCGGGGGCATTCG TATTGTGCCCTTAGAGGTGAAATTC TTGGACCGGCCAAG				
Rat	TCTATTTGTTGGTTTTCCGGA AC TGAGCCATGATTAA GAGGGACGGCCGGGGGCATTCG TATTGTGCCCGCTAGATACCGCAGCTAGGAA TAATGGAATAGSAC - CGCGGT				
Yeast	1000	1020	1040	1060	1080
Frog	ACTAACTACTCGGA AAGCGTTTGC AAGGACGTTTTCGTTAATCA GACGAAAGTTGAGGG ----- ATCTGATACC GTCTGAGCTTA AACCAT				
Rat	ACGAACCAAAGCGAAAGCATTTGCC AAGAAATTTTTCAATCA GAAACGAAAGTCGAGGTTTCAAGACGATCA GATACCGTCTGATTTCCGACCATA				
	1100	1120	1140	1160	1180
Yeast	AACTATGCCGACTAG -- ATCGGGTGGTGT TTTTAAATGACC CACTCGGTACTTACGAGAAATCAA AGTCTTTGGGTTCTGGGGGGAGTATGGTTCGCAA				
Frog	AACGATGCCGACTAGCGATCCGG CCGGTTATCCCATGACCCGCCGAGCAGCTTCCGGGA AACC AAAGTCTTTGGGTTCCGGGGGGAGTATGGTTGC AA				
Rat	AACGATGCCGACTGGCGATCGGG CCGGTTATCCCATGACCCGCCGCGCAGCTTCCGGGA AACCAAAGTCTTTGGGTTCCGGGGGGAGTATGGTTGC AA				
Yeast	1200	1240	1260	1280	
Frog	AGGCTGAAACTTAAAGGAA TTGACGG AAGGGCCACCAC TAGGAGTGGAGCCTCGCGCT - AATTTGACTCA ACACGGGGAAACTCACCAGGTCCAGACACA A				
Rat	AG-CTGAACTTAAAGGAA TTGACGG AAGGGCCACCACCAGGAGTGGAGCTCGCGCTTAA TTTGACTCAACACGGGAAACTCACC CGCCCGGACACGG				
	1300	1320	1340	1360	1380
Yeast	TAAGGATTGACAGATTGAGAGCTCTTTC TGATTTTGTGGGTGGTGGATGGCCTTTC CAGTTGGTGGAGTGATTTGCTGTCTTAATTTCCGATAAC				
Frog	AAAGGATTGACAGATTGATAGCTCTTTC CGATTCGTGGGTGGTGGATGGCCGT - TCTTAGTTGGTGGAGCGATTTGCTGGTTAA TTTCCGATAAC				
Rat	ACAGGATTGACAGATTGATAGCTCTTTC CGATTCGGTGGGTGGTGGATGGCCGT - TCTTAGTTGGTGGAGCGATTTGCTGGTTAA TTTCCGATAAC				
Yeast	1400	1440	1460	1480	
Frog	GAAAGAGAC -- CTT - AACCTA - CTAATA TAGTGGTCTAGCATTTT GCTGGTTATCC ---- ACTTCTTAGAGGGACTATCGGT - TTCAAGCCGATGGAAGTT				
Rat	GAAAGAGACTCTCCATCCTA ACTAGTTACGCGACCCCG - GCGGTCGCG - GTCC - -- AAC TCTTAGAGGGACAAAGTGGCGTTCA - GCCACACAGATC				
	1500	1520	1540	1560	1580
Yeast	TGAGGCAATAACAGGCTGTGATGCCCTTAGA ACGTCTGGGCGCACGCCGCTACACTGA - CGGAGCCAGCGAGTCT - AACCTTGGCCGAGAGGCTCT				
Frog	-GAG - CAATAACAGGCTGTGATGCCCTTAGA - TGTCCGGGGCTGCAGCGCGCTACACTGAA CCGGATC - ACGGTGTCTACCC TGGCCCGACAGGTG C				
Rat	-GAG - CAATAACAGGCTGTGATGCCCTTAGA - TGTCCGGGGCTGCAGCGCGCTACACTGAA CCGGATC - ACGGTGTCTACCC TGGCCCGACAGGTG C				




Figure 3. Nucleotide sequence of Sprague-Dawley rat 18S rDNA and adjacent transcribed spacers with sequences of yeast *Saccharomyces cerevisiae* (3, 28, 29) and frog *Xenopus laevis* (4, 30, 31) 18S rDNA and adjacent spacers. Bases were numbered from the 5'-terminus and the upstream transcribed spacer region was numbered with negative numbers. The 3'-terminus was at nucleotide 1885 (the total length of the rat 18S rDNA was 1869 bp without dashes in this figure) and the adjoining spacer region was numbered under the sequences. The three sequences were arranged to give the best homology and dashes were added to fill spaces. But, slight ambiguities remained. For example, locations of T bases of yeast sequence or locations of G and C bases of frog sequence between nucleotides 250 and 280 (Fig. 3) were arbitrary. Bases which were common to the three species were typed with bold letters. Four portions, in which six or more consecutive A or T bases of yeast sequence changed mainly to G or C bases for frog and rat sequences, were shown with frames.




Figure 4. Conserved sequence analysis of 18S rDNA between frog and rat (top) and among the three species, yeast, frog, and rat (bottom). Sequences in Fig. 3 were sectioned every 10 nucleotides from the 5'-terminus and the numbers of the conserved nucleotides were plotted along the sequence. The 3'-terminal 5 nucleotides, conserved completely, were plotted as fully conserved with a half width. Three long completely conserved regions (A, B, and C) were marked. Arrows show positions of large (10 nucleotides or more) inserts. U1 and U2 are two unstable regions.

Table 1. 18S rRNA base composition data

	No. of nucleotides	U	A	C	G	%G+C
Yeast	1789	509	475	347	458	45.0
Difference	36	98	43	120	57	8.8
Frog	1825	411	432	467	515	53.8
Difference	44	3	11	30	28	1.8
Rat	1869	408	421	497	543	55.6
Total Difference ¹	80	101	54	150	85	10.6

¹Total difference is the difference between yeast and rat.

Table 2. 18S rDNA base change data¹

	Yeast \triangleright Frog	Frog \triangleright Rat	Yeast \triangleright Rat
[a] (A,T) \rightarrow (G,C)	217	46	224
[b] (G,C) \rightarrow (A,T)	70	21	65
[c] [a]/[b]	3.1	2.2	3.4
[d] (G,C) insertion	46	35	81
[e] (A,T) insertion	17	12	14
[f] [d]/[e]	2.7	2.9	5.8
[g] (A,T) deletion	19	2	10
[h] (G,C) deletion	8	1	5
[i] [g]/[h]	2.4	2	2

¹Base changes were counted in Fig. 3.

portions of the genes where six or more consecutive A or T bases of yeast sequence are changed mostly to G or C bases are framed in Fig. 3. There is another clear preference for insertions of G or C over A or T for the three comparisons (lines [d], [e], [f]). GC rich insertions in the 5'-domain of rabbit 18S rRNA that are absent in both yeast and frog 18S rRNA were also pointed out by Lockard *et al.* (16). Though the numbers are small, there are also selective deletions of A or T over G or C (lines [g], [h], [i]).

DISCUSSION

In addition to other genetic systems that have been utilized for the analysis of evolutionary trends (17, 18), the 18S rDNA sequence is an excellent system to study evolutionary trends among various organisms. Because the complete sequence information of 18S rDNA is limited to only three species, it is not easy to discuss their evolutionary relationship. But, these sequences will serve as an introduction to a detailed phylogenetic analysis in the future. Recently, Eckenrode and Meagher (personal communication) have determined the complete 18S rDNA sequence of soybean. It is only 1807 nucleotides long and quite different from the rat sequence. The other high molecular weight rRNA, 25S rRNA for yeast and 28S rRNA for rat, is more variable than 18S rRNA and would be suitable to study evolutionary trends between closer species. Nucleotide sequences of 25S rDNA were compared between two yeasts, *S. cerevisiae* (19) and *S. carlsbergensis* (20), and 16 base differences were observed between the two species (21).

The present study shows that there are three completely conserved large regions in the 18S rRNA (Fig. 4). These conserved regions are considered to be functionally important. Possibilities of base-paired interactions between the 3'-terminal region C (Fig. 4) of 18S rRNA and 5S RNA and between the 3'-terminal region of 18S rRNA and the 5'-terminal region of many eukaryotic mRNAs have been suggested (14, 15). Functions for internal conserved regions A and B (Fig. 4) have not been suggested previously.

Two main factors are recognized which may have contributed to the evolution of 18S rRNA. The first is a preference of point mutations of (A,T) \longrightarrow (G,C) compared to mutations of (G,C) \longrightarrow (A,T). The second is preferential insertions of (G,C) over (A,T). Point mutations are more frequent in the yeast \longrightarrow frog comparison than in the frog \longrightarrow rat comparison, but insertions are introduced almost to the same degree in these two cases. The only large T1 RNase fragment found to be different between rat and human 18S rRNA is fragment 12 (ACCCCCUUCCCG) (22). This fragment is missing in human 18S rRNA and a large portion of this rat fragment (CUUCCCG) is located in the inserted region between frog and rat, nucleotides 199-205 (Fig. 3). We can expect some difference between rat and human 18S rRNA due to insertions. Fragment 12 is also missing in chicken 18S rRNA (11).

The closer relatedness of the rat and Xenopus 18S rDNA sequences compared with the Saccharomyces cerevisiae sequence is consistent with the contrast between the organization of rat and Xenopus rDNA repeat units, which have the 5S and 45S rRNA information separated (23), and the S. cerevisiae rDNA repeat unit, which contains both the 5S and 37S rRNA information together (21, 24). Nevertheless, the 5S rDNA is not a part of the repeat unit containing the 37S rDNA precursor in Neurospora crassa (25) and Schizosaccharomyces pombe (26).

Comparing the base compositions of the 18S rRNAs, the total differences of pyrimidines between yeast and rat were about twice the total differences of purines (Table 1). In structural genes, because of wobble of the third bases of triplets, pyrimidines have more freedom for change than purines without changing the corresponding amino acids. The fact that pyrimidines changed about twice more than purines between yeast and rat might lead to a speculation that 18S rRNA functions, or functioned, as mRNA at some stage before maturation or at some evolutionary stages. But, the present mature X. laevis 18S rRNA is considered to be unsuitable as a template for protein synthesis (27).

ACKNOWLEDGEMENTS

We thank Kenrick J. Dennis for making the initial restriction map of the central portion of the 18S rDNA, Carol Crumley for secretarial assistance and Cheryl Hendrix and Drs. Susan Berent and Paul W. Bragg for critical reading of this manuscript. This work was supported by a Meadows Foundation Grant to the Oree Meadows Perryman Laboratory for Cancer Research at Wadley Institutes and a National Institutes of Health Grant (GM-28090) to APB.

REFERENCES

1. Gerbi, S.A. (1976) *J. Molec. Biol.* 106, 791-816.
2. Fuke, M., Busch, H. and Rao, P.N. (1976) *Nucleic Acids Res.* 3, 2939-2957.
3. Rubtsov, P.M., Musakhanov, M.M., Zakharyev, V.M., Krayev, A.S., Skryabin, K.G. and Bayev, A.A. (1980) *Nucleic Acids Res.* 8, 5779-5794.
4. Salim, M. and Maden, B.E.H. (1981) *Nature* 291, 205-208.
5. Fuke, M., Dennis, K.J. and Busch, H. (1981) *Mol. Gen. Genet.* 182, 25-30.
6. Torczynski, R., Bollon, A.P. and Fuke, M. (1981) *Mol. Gen. Genet.* 184, 557-559.
7. Torczynski, R., Bollon, A.P. and Fuke, M. (1983) *Mol. Gen. Genet.*, in press.
8. Fuke, M. (1974) *Proc. Natl. Acad. Sci. USA* 71, 742-745.
9. Fuke, M. and Busch, H. (1979) *FEBS Lett.* 99, 136-140.
10. Maxam, A.M. and Gilbert, W. (1977) *Proc. Natl. Acad. Sci. USA* 74, 560-564.
11. Fuke, M. and Busch, H. (1979) *Nucleic Acids Res.* 7, 1131-1136.
12. Maden, B.E.H., Forbes, J., de Jonge, P. and Klootwijk, J. (1975) *FEBS Lett.* 59, 60-63.
13. Saponara, A.G. and Enger, M.D. (1974) *Biochem Biophys. Acta* 349, 61-77.
14. Hagenbuchle, O., Santer, M., Steitz, J.A. and Mans, R.J. (1978) *Cell* 13, 551-563.
15. Azad, A.A. and Deacon, N.J. (1980) *Nucleic Acids Res.* 8, 4365-4376.
16. Lockard, R.E., Connaughton, J.F. and Kumar, A. (1982) *Nucleic Acids Res.* 11, 3445-3457.
17. Bryson, V. and Vogel, H.J. (1965) *Evolving Genes and Proteins*, Academic Press, New York.
18. Kimura, M. (1982) *Molecular Evolution, Protein Polymorphism and the New Theory*, Springer-Verlog, New York.
19. Georgiev, O.I., Nikolaev, N., Hadjiolev, A.A., Skryabin, K.G., Zakharyev, J.M. and Bayev, A.A. (1981) *Nucleic Acids Res.* 9, 6953-6958.
20. Veldman, G.M., Klootwijk, J., de Regt, V.C.H.F., Planta, R.J., Branlant, C., Krol, A. and Ebel, J.-P. (1981) *Nucleic Acids Res.* 9, 6935-6952.
21. Bollon, A.P. (1982) in *The Cell Nucleus*, Vol. 10, Eds. Busch, H. and Rothblum, L., pp. 67-125, Academic Press, New York.
22. Fuke, M. and Busch, H. (1975) *J. Molec. Biol.* 99, 277-281.
23. Reeder, R.H. (1974) in *Ribosomes*, Eds. Nomura, M., Tissières, A. and Lengyel, P., 489-518, Cold Spring Harbor Laboratory, New York.
24. Nath, K. and Bollon, A.P. (1977) *J. Biol. Chem.* 252, 6562-6571.

25. Free, S.J., Rice, P.W. and Metzenberg, R.L. (1979) *J. Bacteriol.* 137, 1219-1226.
26. Mao, J., Appel, B., Schaack, J., Sharp, S., Yamada H. and Söll, D. (1982) *Nucleic Acids Res.* 10, 487-500.
27. Maden, B.E.H. (1982) in *The Cell Nucleus*, Vol. 10, Eds. Busch, H. and Rothblum, L., pp. 319-351, Academic Press, New York.
28. Skryabin, K.G., Kraev, A.S., Rubtsov, P.M. and Bayev, A.A.A. (1979) *Dokl. Acad. Nauk. USSR* 247, 761-765.
29. Skryabin, K.G., Zakharyev, V.M., Rubtsov, P.M. and Bayev, A.A.A. (1979) *Dokl. Acad. Nauk. USSR* 247, 1275-1277.
30. Hall, L.M.C. and Maden, B.E.H. (1980) *Nucleic Acids Res.* 8, 5993-6005.
31. Maden, B.E.H., Moss, M. and Salim, M. (1982) *Nucleic Acids Res.* 10, 2387-2398.