



# The Complexity of Piroplasms Life Cycles

Marie Jalovecka<sup>1,2,3\*</sup>, Ondrej Hajdusek<sup>2</sup>, Daniel Sojka<sup>2</sup>, Petr Kopacek<sup>2</sup> and Laurence Malandrin<sup>1</sup>

<sup>1</sup> BIOEPAR, INRA, Oniris, Université Bretagne Loire, Nantes, France, <sup>2</sup> Institute of Parasitology, Biology Centre of the Czech Academy of Sciences, České Budějovice, Czechia, <sup>3</sup> Faculty of Science, University of South Bohemia, České Budějovice, Czechia

Although apicomplexan parasites of the group Piroplasmida represent commonly identified global risks to both animals and humans, detailed knowledge of their life cycles is surprisingly limited. Such a discrepancy results from incomplete literature reports, nomenclature disunity and recently, from large numbers of newly described species. This review intends to collate and summarize current knowledge with respect to piroplasm phylogeny. Moreover, it provides a comprehensive view of developmental events of *Babesia*, *Theileria*, and *Cytauxzoon* representative species, focusing on uniform consensus of three consecutive phases: (i) schizogony and merogony, asexual multiplication in blood cells of the vertebrate host; (ii) gamogony, sexual reproduction inside the tick midgut, later followed by invasion of kinetes into the tick internal tissues; and (iii) sporogony, asexual proliferation in tick salivary glands resulting in the formation of sporozoites. However, many fundamental differences in this general consensus occur and this review identifies variables that should be analyzed prior to further development of specific anti-piroplasm strategies, including the attractive targeting of life cycle stages of *Babesia* or *Theileria* tick vectors.

**Keywords:** piroplasms, *Babesia*, *Theileria*, developmental cycle, merogony, gamogony, sporogony

## INTRODUCTION

The group Piroplasmida received its name after its pear-shaped (pyriform) intra-erythrocytic stages and refers to intracellular parasites transmitted exclusively by hard ticks (Ixodidae) (Mehlhorn and Schein, 1993; Votýpka et al., 2017). Piroplasms belong to the most common group of mammalian blood parasites and their impact economically, as well as on veterinary and medical care, is significant. Due to the worldwide distribution of tick vectors, babesiosis is the most common blood disease of free living animals (Homer et al., 2000; Hunfeld et al., 2008) and is considered an emergent zoonosis of humans (Homer et al., 2000; Kjemtrup and Conrad, 2000; Zintl et al., 2003; Hunfeld et al., 2008; Leiby, 2011). From the veterinary point of view, great attention is paid to bovine babesiosis, which is associated with mortalities, abortions, decreased meat, and milk production, but despite permanent epidemiological surveillance, most of the 1–2 billion cattle worldwide are still exposed to babesiosis and outbreaks occur frequently (Bock et al., 2004; Gohil et al., 2013).

## OPEN ACCESS

### Edited by:

Lilach Sheiner,  
Wellcome Trust Centre for Molecular  
Parasitology (WT), University of  
Glasgow, United Kingdom

### Reviewed by:

Julius Lukes,  
University of South Bohemia, Czechia  
Mattie Christine Pawlowic,  
University of Dundee,  
United Kingdom

### \*Correspondence:

Marie Jalovecka  
jalovecka@paru.cas.cz

**Received:** 03 May 2018

**Accepted:** 29 June 2018

**Published:** 23 July 2018

### Citation:

Jalovecka M, Hajdusek O, Sojka D,  
Kopacek P and Malandrin L (2018)  
The Complexity of Piroplasms Life  
Cycles.  
*Front. Cell. Infect. Microbiol.* 8:248.  
doi: 10.3389/fcimb.2018.00248

Humans are not natural hosts for any species of *Babesia* but serve as accidental hosts (reviewed in e.g., Yabsley and Shock, 2013). Despite this fact, the incidence of human babesiosis is on the rise and clinical cases have been reported recently from many countries worldwide (reviewed in e.g., Yabsley and Shock, 2013; Vannier et al., 2015).

Taxonomic classification places Piroplasmida species in the phylum Apicomplexa, as close relatives of the malarial disease agents, *Plasmodium* parasites (e.g., Burki et al., 2009; Janouskovec et al., 2010; Arisue and Hashimoto, 2014; Schreeg et al., 2016). Based on multi-gene analyses, the order Piroplasmida includes three genera, *Babesia*, *Theileria*, and *Cytauxzoon*. Piroplasmids share many morphological and developmental features such as apical complex organelles, merogony (asexual multiplication) within erythrocytes of vertebrate hosts and sexual multiplication followed by sporozoite

formation in invertebrate vectors, ticks (Homer et al., 2000). There are five evolutionary lineages recognized in the order Piroplasmida (Schreeg et al., 2016). All these lineages differ in particular developmental features and possess unique adaptations (Table 1). The lifecycle of piroplasmids is considered as only partially elucidated. There are many inconsistencies about crucial developmental events of piroplasmids, and relevant information are spread throughout many publications. The recent nomenclature changes and redescription of many species has also contributed to these misconceptions (e.g., Mehlhorn and Schein, 1998; Malandrin et al., 2010; Baneth et al., 2015).

In this review we provide a comprehensive overview of piroplasm lifecycle events, proposing uniform consensus and stressing unique developmental adaptations with respect to evolutionary lineages.

**TABLE 1** | Summarizing overview of characteristic life cycle events of five evolutionary lineages of the order Piroplasmida (based on Schreeg et al., 2016).

	<i>Babesia sensu stricto</i>	<i>Theileria sensu stricto</i>	<i>Theileria equi</i>	Western <i>Babesia</i> group	<i>Babesia microti</i> group
Reduced apical complex organelles	✓	✓	✓	✓	✓ <sup>a</sup>
Schizogony in nucleated blood cells	✗ <sup>b</sup>	✓	✓	✗ <sup>b</sup>	✗ <sup>b</sup>
Neoplastic transformation of the nucleated blood cells	✗	✓ <sup>c</sup>	✗	✗	✗
Merogony in red blood cells	✓ <sup>d</sup>	✓ <sup>d</sup>	✓ <sup>d</sup>	✓ <sup>d</sup>	✓ <sup>d</sup>
Motile sporozoites	✓	✗ <sup>e</sup>	✗ <sup>c</sup>	?	✓
Parasitophorous vacuole formation	✗	✗	✗	✗	✗
Piriform shape of merozoites	✓	✓	✓	✓ <sup>f</sup>	✓ <sup>f</sup>
Gametocytes in the host bloodstream	✓ <sup>g</sup>	✓ <sup>g</sup>	✓ <sup>g</sup>	?	✓ <sup>g</sup>
Strahlenkörpers/spiky-rayed gametes	✓ <sup>h</sup>	✓	✓	?	✓ <sup>h</sup>
Macro- and micro-gametes differentiation	✗	✓ <sup>i</sup>	✓ <sup>i</sup>	?	✗
Zygote formation	✓ <sup>j</sup>	✓ <sup>j</sup>	✓ <sup>j</sup>	?	✓ <sup>j</sup>
Primary kinetogenesis in epithelial cells	✓	✓	✓	?	✓
Invasion of primary kinetes directly to salivary glands	✗	✓	✓	?	✓ <sup>k</sup>
Secondary kinetogenesis in tick tissues	✓	✗	✗	?	✓
Kinetes invasion into ovaries, transovarial transmission	✓	✗	✗	?	✗
Sporogony in tick salivary glands	✓	✓	✓ <sup>l</sup>	?	✓
Hypertrophy of infected acini cells	✓ <sup>m</sup>	✓	✓	?	✓
Cymeres formation during sporoblast maturation	✓	✗	✗	?	✗
Polar rings formation in sporozoites	✓	✗	✓	?	✗
Asynchronous sporozoites release by budding process	✓ <sup>n</sup>	✓	✓	?	✓

✓, present; ✗, absent; ?, not elucidated yet; a, more reduced compared to other piroplasmids; b, absence of schizonts has not been convincingly demonstrated for several species in these groups yet; c, neoplastic transformation of the nucleated host blood cells has been reported for only few species; d, merogony is not synchronous, thus trophozoites and merozoites occur in the bloodstream simultaneously; e, sporozoites do not require apical-end first orientation to internalize into the host cell; f, more frequently ovoid and/or polymorphic shape; g, undistinguishable by light microscopy; h, two gametes (Strahlenkörper) populations which are undistinguishable by light microscopy; i, formation of macrogametes (ovoid shape) and microgametes (Strahlenkörper, spiky-rayed shape); j, zygote is motile and penetrates the peritrophic matrix to internalize into midgut epithelial cell; k, *B. microti* is able to directly invade tick salivary glands where the secondary kinetogenesis can take place; l, although generally the sporozoites maturation starts after attachment of molted tick stage onto host, *T. equi* sporozoites can mature prior tick molting; m, hypertrophy of infected acini cells was confirmed for some species (e.g., *B. bovis* and *B. canis*) but excluded for some other species (e.g., *B. ovis* and *B. caballi*); n, apart from *B. canis*, where sporozoite differentiation was described as a result of successive binary fissions. The table is composed based on references provided in the text.

## SCHIZOGONY AND MEROGONY: ASEXUAL MULTIPLICATION IN BLOOD CELLS OF THE VERTEBRATE HOST

All parasites of the group Piroplasmida reproduce asexually inside the blood cells of the vertebrate host (**Figure 1**, **Table 1**). The host infection is initiated by the invasion of sporozoites, transmitted through saliva secretion during the tick bite. The blood cells targeted by sporozoites differ according to the species of piroplasm (Shaw, 2003; Lobo et al., 2012). *Theileria* parasites are characterized by schizogony (**Box 1**) in nucleated blood cells—monocytes and lymphocytes—prior to red blood cell invasion (Schein et al., 1981; Moltmann et al., 1983a; Conrad et al., 1985; Webster et al., 1985; Dobbelaere and Heussler, 1999; Dobbelaere and Rottenberg, 2003; Shaw, 2003). *Babesia* parasites are believed to multiply exclusively in erythrocytes; so far a schizogony has never been convincingly confirmed (Mehlhorn and Schein, 1993; Lobo et al., 2012; Schreeg et al., 2016).

### Schizogony

Intra-leukocytic asexual reproduction (**Figure 1A**) occurs in the lifecycle of two evolutionary lineages, *Theileria sensu stricto* (including *Cytauxzoon* spp.) and its sister clade, represented by *Theileria equi* (**Table 1**) (Kappmeyer et al., 2012; Schnittger et al., 2012; Schreeg et al., 2016). Schizogony serves to aid rapid parasite multiplication and gives rise to schizonts, referred to as Koch's bodies (Mehlhorn and Schein, 1984; Mehlhorn and Schein, 1993). These *Theileria* intra-leukocytic schizonts are able to modulate the host's immune response, e.g., to block host cell apoptosis (Blouin et al., 1987; Kawai et al., 1989; Sato et al., 1994; Hagiwara et al., 1997; Ahmed et al., 1999; Dobbelaere and Heussler, 1999; Susta et al., 2009). Moreover, leukocyte infection by *Theileria* parasites could lead to a fundamental change in the infected host cell's ability to proliferate indefinitely (Mehlhorn and Schein, 1984; Mehlhorn and Schein, 1993; Ahmed et al., 1999; Dobbelaere and Heussler, 1999; Dobbelaere and Rottenberg, 2003). Although schizogony in nucleated blood cells characterizes all *Theileria* parasites, the neoplastic transformation of the host cell was reported only for *Theileria parva*, *Theileria annulata*, *Theileria lestoquardi*, *Theileria taurotragi*, and *Theileria* sp. (buffalo) (**Table 1**; Ahmed et al., 1999; Dobbelaere and Heussler, 1999; Dobbelaere and Küenzi, 2004; Zwegarth et al., 2009; Sivakumar et al., 2014; Bishop et al., 2015). Changes in the host cell have not been described for *T. equi* (Schein et al., 1981; Moltmann et al., 1983a; Ramsay et al., 2013), presumably due to the absence of homologs of the putative *Theileria* host cell transforming genes (Kappmeyer et al., 2012; Schreeg et al., 2016).


The sporozoite invasion into nucleated blood cells is a complex process requiring numerous alterations in the metabolism of invading sporozoites (Shaw, 1995, 1996a,b), complicity of the host system (Shaw et al., 1991, 1993; Shaw, 1996a,b) and the involvement of tick saliva (Shaw et al., 1993). In contrast to other apicomplexan parasites, including *Babesia*, *Theileria* sporozoites are immotile (**Table 1**; Shaw, 1999, 2003). The initial contact of the parasite and host cell membrane thus occurs randomly (Shaw, 2003). Sporozoite attachment and internalization into the host cell does not require apical-end first

orientation and the parasite enters in any orientation (Shaw, 1999, 2003). Thus, the proteins excreted by apical organelles are more involved in the establishment in the host cell cytoplasm rather than the entry process (Shaw et al., 1991). The invasion process comprises several consecutive stages and is completed within about 3 min: (i) recognition and attachment to the host cell membrane; (ii) formation of junctions between the parasite and the host cell membrane; (iii) a “zippering” process resulting in fully internalized parasite in the host cell, yet still surrounded by the host cell membrane; (iv–v) separation and progressive dissolution of the enclosing host cell membrane; and (vi) appearance of a microtubule network derived from the host cell and closely associated with the developing parasite (Fawcett et al., 1982c, 1984; Shaw et al., 1991; Shaw, 1997, 2003). During the invasion process the sporozoite sheds its coat (Dobbelaere et al., 1985; Webster et al., 1985; Shaw, 2003) and lies loosely in the host cell cytoplasm once it has escaped from the enclosing host membrane; no parasitophorous vacuole is formed (Shaw, 2003).

The internalized *Theileria* sporozoite undergoes a change into a multinucleate schizont (Shaw, 1997, 2003) and schizont-infected cells then circulate in the bloodstream (Dobbelaere and Heussler, 1999). The *Theileria* schizogony is accompanied by a series of ultrastructural changes affecting internal organelles as well as the outer surface of the parasite (Shaw and Tilney, 1992). The intra-leukocytic schizogony ends with the production of uninucleated merozoites released into the host bloodstream where they invade erythrocytes (Shaw and Tilney, 1992). The process of *Theileria* merozoite internalization into erythrocytes occurs in the same manner as sporozoite invasion of leukocytes (Shaw and Tilney, 1995). All *Theileria* parasites reproduce in erythrocytes but the process has been described in only a limited number of species (Conrad et al., 1986; Bishop et al., 2004).

### Merogony

Exclusive intra-erythrocytic multiplication (**Figures 1B,C**) represents the cognitive feature of *Babesia* parasites but the absence of schizogony has not yet been demonstrated for several species of all three evolutionary distinct *Babesia* lineages, *Babesia sensu stricto*, Western *Babesia* group, and *Babesia microti* group (**Table 1**; Rudzinska and Trager, 1977; Kjemtrup et al., 2006; Lobo et al., 2012; Schreeg et al., 2016). The first contact between *Babesia* invasive stages and the host cell occurs through several random collisions. Unlike *Theileria*, *Babesia*'s orientation of the apical (anterior) end establishes the junction between parasite and host cell membrane (Rudzinska et al., 1976; Montero et al., 2009; Asada et al., 2012; Lobo et al., 2012). Parasite orientation and penetration is mediated by proteins secreted from the apical secretory organelles (Dubremetz et al., 1998; Soldati et al., 2004), and thus is accompanied by apposition of apical organelles with the host cell membrane (Ward and Jack, 1981). Similarly with *Theileria*, *Babesia* internalizes within a few minutes without parasitophorous vacuole formation (**Table 1**), and thus the parasite lies freely within the host cell cytoplasm (Simpson et al., 1963; Rudzinska et al., 1976; Potgieter and Els, 1977b; Kawai et al., 1999a,b; Guimarães et al., 2003; Montero et al., 2009; Sun et al., 2011; Lobo et al., 2012). The invasion process does not differ for *Babesia sensu stricto* species (Montero et al., 2009;



**Box 1** | Subsequent phases of piroplasm development.

**Schizogony.** A process of asexual multiplication in nucleated blood cells (leukocytes) is typical only for two evolutionary lineages of piroplasms, *Theileria* sensu stricto and *Theileria equi*. Schizogony starts after sporozoite internalization into leukocytes and results in merozoite production, which further multiply by merogony. Schizogony can lead to neoplastic transformation of the nucleated host cells, which then proliferate indefinitely. **Merogony.** A process of asexual division in the red blood cell starts either with sporozoite (*Babesia* species) or merozoite (*Theileria* species) invasion of red blood cells. The internalized parasites develop into trophozoites, which further asexually divide into merozoites. Merozoites are then released by rupture of the host red blood cells and invade healthy erythrocytes. **Gamogony.** Sexual multiplication of the parasite starts by gametocytes appearing in the host red blood cells. During blood uptake by ticks, gametocytes develop into gametes that mature in the tick midgut lumen. Gamete fertilization then gives rise to a zygote that penetrates the tick peritrophic matrix to tick epithelial cells. Inside these, the zygote undergoes a meiotic division and results in the formation of kinetes, which are released to the haemolymph. The kinetes of *Theileria* species directly invade salivary glands (primary kinetes) but kinetes of *Babesia* parasites are subjected to two series of asexual multiplication in various tick tissues and subsequent secondary kinetes invade the tick salivary glands. **Sporogony.** Sporogony starts after kinete invasion of tick salivary glands, which form the sporont, a polymorphous syncytium. The sporont later evolves into a multinucleated meshwork referred as a sporoblast, which is dormant during tick ecdysis. Maturation of the parasite sporoblast starts after tick attachment to the host and results in sporozoites being released into tick saliva.

Asada et al., 2012) and *B. microti* (Rudzinska, 1976; Rudzinska et al., 1976). The internalized *Babesia* sporozoites develop into trophozoites (also described as ring stages), which further asexually divide and produce merozoites by a process referred to as merogony (**Box 1, Figure 1**) (Rudzinska and Trager, 1977; Fawcett et al., 1987; Montero et al., 2009; Lobo et al., 2012). Later, merozoites are released from ruptured cells and invade other intact and healthy erythrocytes. The merogony of piroplasms is asynchronous (**Table 1**), and thus trophozoites and merozoites occur in the bloodstream simultaneously (Jalovecka et al., 2016). The short-time residence of the parasite outside the host cell is characterized by the appearance of a fuzzy coat created from fibrillary material and hypervariable surface proteins. The coat occurs on the surface of both *Theileria* and *Babesia* free merozoites and is cut off during invasion of the host cell (Rudzinska et al., 1976; Shaw, 2003; Montero et al., 2009).

The size of merozoites varies according to the piroplasm species as well as the vertebrate host species. The merozoites of piroplasms are characterized by a piriform shape, forming pairs or tetrads (Potgieter and Els, 1977a,b; Lewis et al., 1980; Mehlhorn and Schein, 1984; Conrad et al., 1985, 1986; Kawai et al., 1986, 1999b; Gorenflot et al., 1991, 1992; Shaw and Tilney, 1995; Shaw, 2003; Wise et al., 2013; Del Carmen Terrón et al., 2016). However, divergently shaped merozoites were recorded for species of two early divergent lineages of piroplasms, the Western *Babesia* clade and the *Babesia microti* clade (**Table 1**) (Rudzinska, 1976; Kjemtrup et al., 2006; Clancey et al., 2010). They possess smaller merozoites of ovoid shape, which later become polymorphic and form numerous invaginations and pseudopods, twisting and coiling. Although all piroplasms exhibit a much reduced apical complex (e.g., absence of conoid) compared to the other apicomplexan parasites (Votypka et al., 2017), *B. microti* merozoites apical complex displays only a single large rhoptry and lack polar rings and a microtubular section (Rudzinska, 1976). The question of feeding mechanisms of piroplasms has not yet been answered. There is a general consensus that piroplasms phagocytose or pinocytose the host cytoplasm (Rudzinska and Trager, 1962; Conrad et al., 1985; Fawcett et al., 1987; Guimarães et al., 2003), but food vacuoles full of host cytoplasm were observed in both *Theileria* and *Babesia* merozoites, suggesting potential extracellular digestion of host cytoplasm (Rudzinska and Trager, 1962; Simpson et al., 1963;

Rudzinska, 1976; Rudzinska et al., 1976; Simpson and Neal, 1980; Conrad et al., 1985; Fawcett et al., 1987; Guimarães et al., 2003). If piroplasms can directly digest host hemoglobin still remains a question; this phenomenon was so far suggested only for some *Theileria* species but potential host hemoglobin digestion is not accompanied by pigment or other visible residues formation (Votypka et al., 2017).

## GAMOGONY: SEXUAL REPRODUCTION IN THE GUT OF THE TICK VECTOR

The first sexual stages of piroplasms are referred to as gametocytes (misinterpreted as gamonts in older studies) and appear in the host red blood cells (**Box 1, Figure 1, Table 1**) (Rudzinska et al., 1979; Mehlhorn and Schein, 1984; Mackenstedt et al., 1990; MacKenstedt et al., 1995; Mehlhorn and Schein, 1993; Gauer et al., 1995; Becker et al., 2010; Bastos et al., 2013; Jalovecka et al., 2016). Gametocytes are predetermined to further differentiate into gametes in the lumen of the tick gut (Rudzinska et al., 1979; Bishop et al., 2004; Becker et al., 2010, 2013), and thus mediate the ability to infect the tick vector (Uilenberg, 2006; Lobo et al., 2012; Becker et al., 2013). Unlike the normally growing and asexually reproducing merozoites, the gametocytes do not reproduce (Rudzinska et al., 1979; Mackenstedt et al., 1990; MacKenstedt et al., 1995; Lobo et al., 2012). They are believed to be larger and unusually shaped compared to asexual stages, however, light microscopy does not allow their reliable recognition (Rudzinska et al., 1979; Lobo et al., 2012). As documented for *B. microti* by electron microscopy, intra-erythrocytic gametocytes are characterized by undifferentiated cytoplasm, large nuclei and unusually twisted, convoluted or folded shapes (Rudzinska et al., 1979). The gametocytes persistence in the host bloodstream was documented for many species of *Babesia* sensu stricto (Mackenstedt et al., 1990; MacKenstedt et al., 1995; Becker et al., 2010, 2013; Bastos et al., 2013; Jalovecka et al., 2016) and *B. microti* (Rudzinska et al., 1979, 1983b). It is generally assumed that sexual commitment of *Theileria* is identical to *Babesia* and gametocytes occur in the circulating blood (Mehlhorn and Schein, 1984; Mehlhorn and Schein, 1993; Zapf and Schein, 1994b; Gauer et al., 1995; Shaw, 2003; Bishop et al., 2004; Uilenberg, 2006). This is supported

by recent descriptions of genes with expression specific to sexual commitment in intra-erythrocytic stages of *Theileria* sensu stricto species (Pieszko et al., 2015; Lempereur et al., 2017) and *T. equi* (Bastos et al., 2013). The gametocytes are taken up into the tick midgut together with the blood meal. Subsequently, ingested asexual intra-erythrocytic stages are rapidly destroyed in the gut lumen (Rudzinska et al., 1979; Mehlhorn and Schein, 1984; Bishop et al., 2004; Lobo et al., 2012). Still hidden inside the red blood cells, gametocytes of both *Theileria* and *Babesia* parasites start reorganizing the cytoplasm (Schein et al., 1977; Rudzinska et al., 1979; Mehlhorn et al., 1980; Zapf and Schein, 1994b). Gametocyte metamorphosis is asynchronous, presumably due to non-contemporary blood uptake (Rudzinska et al., 1984). The process is accompanied by microtubular reorganization and gametocytes became completely stretched out compared to previously folded intra-erythrocytic forms (Friedhoff and Büscher, 1976; Weber and Friedhoff, 1977; Rudzinska et al., 1979; Mehlhorn and Schein, 1984; Zapf and Schein, 1994b). As documented for *B. microti*, development of the gametocytes is completed outside of the already lysed erythrocytes in the lumen of the midgut (Rudzinska et al., 1979). Yet, in some cases the process can be completed inside the erythrocyte in the environment of the tick lumen (Rudzinska et al., 1984; Gough et al., 1998).

Metamorphosis of the gametocytes results in the formation of gametes (**Figure 1**), referred to as Strahlenkörper or spiky-rayed stages (**Table 1**) (Mehlhorn and Schein, 1993). It was suggested that gametes multiply to form large aggregates but once division is completed, haploid gametes are released to the tick midgut lumen (Warnecke et al., 1980; Mehlhorn and Schein, 1993; MacKenstedt et al., 1995; Gough et al., 1998; Bock et al., 2004). The appearance of piroplasm gametes is unique among apicomplexan parasites and characteristic structures—tail, arms, and arrowhead—begin to form in gametocytes (Rudzinska et al., 1984). Gametes of both *Theileria* and *Babesia* species are haploid (Mackenstedt et al., 1990; MacKenstedt et al., 1995; Gauer et al., 1995) and are considered to be anisogametes, although *Babesia* gametes appear as isogametes when examined by light microscopy (Mehlhorn and Schein, 1984; Gough et al., 1998). The gametes of *Theileria* sensu stricto and *T. equi* are clearly distinguishable by light microscopy as micro- and macro-gametes (**Table 1**). The characteristic ray bodies are considered to be micro-gametes and macro-gametes are spherically shaped without protrusions (Schein et al., 1977; Warnecke et al., 1980; Mehlhorn and Schein, 1984; Mehlhorn and Schein, 1993; Zapf and Schein, 1994b; Bishop et al., 2004; Uilenberg, 2006). Gametes of *Babesia* sensu stricto species as well as *B. microti* do not differentiate into macro- and micro-gametes but two gamete populations are formed (**Table 1**). These two types differ in the details of cytoplasm density and shape (Friedhoff and Büscher, 1976; Rudzinska et al., 1979, 1983b; Mehlhorn et al., 1981; Gough et al., 1998).

Fertilization of piroplasmids is induced by close contact between two gametes of different types and may occur at very early stages of gamete formation (Rudzinska et al., 1983b). Filamentous structures are formed between membranes of closely adjacent gametes. Subsequently, a finger-like protrusion of one gamete penetrates the opposite one (Mehlhorn et al., 1981; Rudzinska

et al., 1983b; Mehlhorn and Schein, 1984). Gamete fertilization results in the formation of a zygote (Schein et al., 1977; Mehlhorn et al., 1979; Warnecke et al., 1980; Rudzinska et al., 1983b, 1984; Mackenstedt et al., 1990; MacKenstedt et al., 1995; Higuchi et al., 1991a, 1999a,b; Zapf and Schein, 1994b; Gauer et al., 1995; Gough et al., 1998; Bishop et al., 2004). The zygote of Piroplasmida is a motile stage that is often referred to as an ookinete or kinete. However, such nomenclature is misleading since kinetes (often also called ookinetes or sporokinetes) represent haploid stages resulting from the meiotic division of a diploid zygote (Mehlhorn et al., 1978, 1979; Mehlhorn and Schein, 1984; Rudzinska et al., 1984). To further develop, the zygote penetrates the peritrophic matrix (**Figure 1, Table 1**) (Rudzinska et al., 1982), which appears temporarily during feeding at all tick stages and compartments the gut lumen into endo-peritrophic and ecto-peritrophic spaces (Sonenshine, 1991). Since the peritrophic matrix represents a strong mechanical barrier, zygote penetration is an active process accomplished by enzymes released from the arrowhead structure of the zygote (Rudzinska et al., 1982, 1984). Matrix penetration starts immediately after zygote formation (Rudzinska et al., 1982, 1983a,b). The arrowhead structure opens the way for the zygote body by release of enzymes. Subsequently, the zygote enters the ecto-peritrophic space and immediately invades gut epithelial cells. This cell invasion is triggered by the arrowhead structure of the zygote but the arrowhead does not pierce the cell membrane; the membrane remains intact. Once the zygote is internalized into the epithelial cell, the invagination membrane disappears. Thus, the zygote occurs loosely in the cytoplasm of the epithelial cell and is surrounded by cell organelles. Inside the epithelial cell, the zygote turns into a spherical shape. Simultaneously, the arrowhead structure loses its organized pattern and gradually disappears (Rudzinska et al., 1982, 1983a). Zygote penetration of the peritrophic matrix and internalization into epithelial cells has been described in detail only for *B. microti* species but is generally assumed to be consistent for all Piroplasmida (Mehlhorn and Schein, 1993). Once morphological changes are finalized, the zygote undergoes a meiotic division as evidenced by DNA measurements of *Theileria* and *Babesia* species (Gauer et al., 1995; MacKenstedt et al., 1995). Meiosis inside the epithelial cell results in the formation of kinetes as was documented for species of both *Theileria* sensu stricto and *Babesia* sensu stricto lineages, as well as for *T. equi* and *B. microti* (**Table 1**) (Potgieter et al., 1976; Potgieter and Els, 1977a; Mehlhorn et al., 1978; Warnecke et al., 1980; Rudzinska et al., 1984; Zapf and Schein, 1994b).

The kinetes are released from the gut epithelial cells into the tick haemolymph (**Figure 1**) (Potgieter et al., 1976; Potgieter and Els, 1977a; Mehlhorn et al., 1978, 1979; Schein and Friedhoff, 1978; Warnecke et al., 1980; Mehlhorn and Schein, 1984; Rudzinska et al., 1984; Karakashian et al., 1986; Higuchi et al., 1991b; Zapf and Schein, 1994b), where the motile kinetes are disseminated via the haemolymph throughout the whole tick body and invade internal tissues. Kinetes are primarily uni-nucleated but exceptionally, kinetes with more nuclei can occur in the haemolymph due to the early beginning of nuclear division (Mehlhorn and Schein, 1993). In the haemolymph, as other invasive stages of piroplasmids, kinetes are covered with a fuzzy coat created from fibrillary material and hypervariable

surface proteins (Karakashian et al., 1986). The kinetes of *Babesia* sensu stricto species are subjected to two cycles of asexual multiplication (**Table 1**) (Potgieter and Els, 1977a; Mehlhorn et al., 1980; Mehlhorn and Shein, 1984; Mehlhorn and Schein, 1998). In the first, the *Babesia* kinetes invade various tick tissues like haemocytes, muscle fibers, Malpighian tubules, peritracheal cells, and ovaries of adult females. Here, the kinetes undergo the second asexual multiplication (Potgieter and Els, 1977a; Moltmann et al., 1982b; Mehlhorn and Shein, 1984). Subsequently, the secondary kinetes invade the salivary glands where sporogony, the maturation of sporozoites, takes place (Christophers, 1907; Friedhoff et al., 1972; Potgieter and Els, 1976; Weber and Friedhoff, 1979; Moltmann et al., 1982a; Mosqueda et al., 2004). Species of the *Babesia* sensu stricto lineage possess a unique feature among all apicomplexan parasites; transovarial transmission (**Figure 1**, **Table 1**). This process is mediated by *Babesia* invasion into the ovarian cells and transmission via larval progeny to tick larvae (Joyner et al., 1963; Donnelly and Peirce, 1975; Lewis and Young, 1980; Moltmann et al., 1982b; Mehlhorn and Shein, 1984; Higuchi et al., 1993; Mehlhorn and Schein, 1993; Bonnet et al., 2007; Boldbaatar et al., 2008, 2010). No transovarial transmission occurs in the lifecycle of *B. microti* or *Theileria* species (**Table 1**). The kinetes of *B. microti* primarily invade fat body (nephrocytes) and salivary glands. Inside, the kinetes form the kinetoblast, which differentiates to produce secondary kinetes. Subsequently, secondary kinetes invade salivary glands to undergo sporogony (Karakashian et al., 1986). Kinetes of *Theileria* sensu stricto species and *T. equi* are believed to migrate directly to salivary glands (**Table 1**) as no kinete invasion of other tick tissues has been documented (Mehlhorn et al., 1979; Moltmann et al., 1983b; Mehlhorn and Shein, 1984; Mehlhorn and Schein, 1993, 1998; Zapf and Schein, 1994a; Uilenberg, 2006).

Remarkable differences have been documented in size and chronological order of piroplasm sexual development. Such divergence can be attributed to the variety of the piroplasm species and the wide spectrum of both vertebrate hosts and vectors. Gametes of *Babesia* parasites develop during tick feeding, appear before full tick engorgement and within ~3 days post tick repletion. Subsequently, the kinetes are found in the tick haemolymph from ~2 to ~6 days post repletion. *Theileria* sexual development seems to be a longer process; the first appearance of gametes was documented between ~1 and ~5 days post tick repletion and kinetes released to the haemolymph were first seen from ~13 to ~34 days post repletion. In general, the length of piroplasm sexual development correlates with the feeding duration of tick developmental stages (larvae vs. nymphs vs. adults), and tick developmental differences derived from the number of host species (one- vs. two- vs. three-host ticks) (Friedhoff and Büscher, 1976; Potgieter et al., 1976; Potgieter and Els, 1977a; Schein et al., 1977; Weber and Friedhoff, 1977; Mehlhorn et al., 1978, 1979, 1980; Warnecke et al., 1980; Rudzinska et al., 1982, 1983a,b, 1984; Mehlhorn and Shein, 1984; Mackenstedt et al., 1990; MacKenstedt et al., 1995; Higuchi et al., 1991a, 1992, 1999a,b; Mehlhorn and Schein, 1993; Zapf and Schein, 1994b; Gauer et al., 1995; Gough et al., 1998).

## SPOROLOGY: ASEXUAL REPRODUCTION IN THE SALIVARY GLAND OF THE TICK VECTOR

The kinetes of Piroplasmida parasites further develop in tick salivary glands to produce invasive stages referred to as sporozoites (**Figure 1**). The sporozoites mediate parasite transmission from the tick vector to the vertebrate host. Piroplasmids develop in acini of types II and III (Fawcett et al., 1982a,b), which represent the majority of acini in the typical grape-like structure of salivary glands (Coons and Roshdy, 1973; Binnington, 1978). Sporogony (**Box 1**) begins with invasion of tick salivary glands by piroplasm kinetes (Christophers, 1907; Holbrook et al., 1968; Purnell and Joyner, 1968; Friedhoff et al., 1972; Potgieter and Els, 1976; Schein et al., 1979; Weber and Friedhoff, 1979; Weber and Walter, 1980; Fawcett et al., 1982a,b; Moltmann et al., 1982a, 1983b; Karakashian et al., 1983; Blouin and van Rensburg, 1988; Blouin and De Waal, 1989; Higuchi et al., 1994; Zapf and Schein, 1994a; Guimarães et al., 1998a,b). Invading kinetes rapidly enlarge and transform into the polymorphous single-membrane syncytium referred to as a sporont. Later, the sporont evolves into a sporoblast, a multinucleated and relatively undifferentiated three-dimensional branching meshwork which has already developed before the beginning of tick ecdysis and molting. The formation of a sporoblast is associated with hypertrophy of the infected acini cells (**Table 1**), which is a common feature of sporogony in *Theileria* sensu stricto species (Purnell and Joyner, 1968; Schein and Friedhoff, 1978; Mehlhorn et al., 1979; Fawcett et al., 1982a,b), *T. equi* (Zapf and Schein, 1994a) and *B. microti* (Weber and Walter, 1980; Karakashian et al., 1983; Piesman et al., 1986; Yano et al., 2005). The same phenomena were documented for some *Babesia* sensu stricto species like *B. bovis* (Potgieter and Els, 1976) and *B. canis* (Schein et al., 1979) but not for *B. ovis* (Friedhoff et al., 1972; Moltmann et al., 1982a) or *B. caballi* (Blouin and De Waal, 1989). During tick ecdysis, the sporoblast appears to be dormant and its maturation starts when the molted tick attaches to the host. However, a unique formation of fully matured sporozoites prior to tick molting was documented for *T. equi* (**Table 1**). Sporozoites developed during parasite acquisition and were competent to transmit and expand the infection to the naïve host. Later, the secondary sporozoites developed after attachment of the molted tick stages (Zapf and Schein, 1994a).

Sporoblast maturation of *Babesia* sensu stricto species begins with the appearance of numerous cytomeres (**Table 1**) (Friedhoff et al., 1972; Potgieter and Els, 1977a; Schein et al., 1979; Moltmann et al., 1982a) but cytomere formation was absent in *Theileria* species (Fawcett et al., 1982a,b; Hazen-Karr et al., 1987) and *B. microti* (Karakashian et al., 1983). The structures of the apical complex appear subsequently, but prior to the sporozoites budding off the sporoblast (Karakashian et al., 1983). Some skeletal components of the apical complex—the conoid and microtubules—are absent in sporozoites of all piroplasmids (Friedhoff et al., 1972; Schein and Friedhoff, 1978; Mehlhorn et al., 1979; Schein et al., 1979; Fawcett et al., 1982a,b; Karakashian et al., 1983; Moltmann et al., 1983b; Piesman et al., 1986; Zapf

and Schein, 1994a). Polar rings are formed in sporozoites of *Babesia sensu stricto* species (Table 1) (Friedhoff et al., 1972; Schein et al., 1979; Weber and Friedhoff, 1979; Moltmann et al., 1982a; Blouin and van Rensburg, 1988; Blouin and De Waal, 1989) and *T. equi* (Moltmann et al., 1983b) but do not appear in sporozoites of *B. microti* (Karakashian et al., 1983; Yano et al., 2005) and *Theileria sensu stricto* species (Fawcett et al., 1982a,b). Thus, *B. microti* sporogony is more reminiscent of *Theileria sensu stricto* species with respect to sporoblast structure. The formation of sporozoites is attributed to the process of multiple fissions, referred to as budding (Table 1). Apart from *B. canis*, where sporozoite differentiation was described as a result of successive binary fissions (Schein et al., 1979), the budding process was documented for both *Theileria* and *Babesia* species (Holbrook et al., 1968; Friedhoff et al., 1972; Potgieter and Els, 1976; Schein and Friedhoff, 1978; Fawcett et al., 1982a; Moltmann et al., 1982a, 1983b; Hazen-Karr et al., 1987; Blouin and van Rensburg, 1988; Blouin and De Waal, 1989). Since the parasitophorous vacuole is not formed, the sporogony stages are in immediate contact with the host-cell cytoplasm (Friedhoff et al., 1972; Fawcett et al., 1982a; Moltmann et al., 1982a, 1983b). Piroplasm sporogony is asynchronous and the various developmental stages occur within individual acinar cells (Karakashian et al., 1983; Blouin and van Rensburg, 1988; Yano et al., 2005). This is attributed to the continuous release of sporozoites into the tick saliva and to the bloodstream of the vertebrate host during the several days of tick feeding (Yano et al., 2005). Sporogony ends with longish piriform sporozoites equipped with apical organelles that later mediate internalization to the host blood cell.

Sporogenic events and progress are assumed to be consistent for both *Babesia* and *Theileria* species (Christophers, 1907; Purnell and Joyner, 1968; Friedhoff et al., 1972; Potgieter and Els, 1976; Weber and Friedhoff, 1979; Fawcett et al., 1982a; Moltmann et al., 1982a, 1983b; Karakashian et al., 1983; Higuchi et al., 1994; Zapf and Schein, 1994a; Guimarães et al., 1998a,b; Mehlhorn and Schein, 1998; Mosqueda et al., 2004) but alteration in the process length and sporozoite size occurs. Such a discrepancy is attributed to the variety of tick species and their natural developmental characteristics such as the number of host species (one- vs. two- vs. three-host ticks). Generally, sporozoite maturation after tick attachment to the host lasts at least 48 h, as documented for representatives of lineages *Babesia sensu stricto*, *Theileria sensu stricto*, *T. equi* and *B. microti* (Mehlhorn et al., 1979; Karakashian et al., 1983; Takahashi et al., 1993; MacKenstedt et al., 1995; Guimarães et al., 1998a,b).

## CONCLUSION AND FUTURE PERSPECTIVES: THE ROLE OF THE LIFE CYCLE IN ANTI-PIROPLASM STRATEGIES

Species of the group Piroplasmida possess a characteristic lifecycle that significantly differs from other apicomplexan parasites. A uniform consensus describes three consecutive phases (Figure 1): (i) schizogony and merogony, asexual multiplication in blood cells of the vertebrate host; (ii) gamogony, sexual reproduction inside the tick midgut, later

followed by kinete invasion of the tick internal tissues; and (iii) sporogony, asexual proliferation in tick salivary glands resulting in the formation of sporozoites (Box 1). However, the order Piroplasmida includes many species spread into five evolutionary distinct lineages. Thus, in the lifecycles of piroplasmids many fundamental variations occur from the general consensus and these discrepancies need to be taken into account in the development of anti-piroplasm strategies.

To date, the majority of inventions in anti-piroplasm vaccine development exploits the vertebrate host stages: schizonts and/or merozoites (Florin-Christensen et al., 2014; Nene and Morrison, 2016). Trends in anti-babesial strategies operate particularly with subunit vaccines based on merozoite surface antigens since recently, many of these have been characterized (recently reviewed in Florin-Christensen et al., 2014). In general, these surface antigens exhibit high immunogenicity and antibodies against them are able to mediate inhibition of parasite intra-erythrocytic invasion and development (Florin-Christensen et al., 2014). Analogously, surface antigens of schizont-infected cells represent hot candidates for anti-theilerial vaccines (Nene and Morrison, 2016). Sporozoites, crucial piroplasm stages responsible for parasite transmission from the tick vector to the vertebrate host, are currently the center of interest for anti-piroplasm strategies (Florin-Christensen et al., 2014; Nene and Morrison, 2016; Nene et al., 2016) but this research is restricted by the absence of effective laboratory transmission models, particularly for species in lineages of *T. equi* and *B. microti*. So far, a few sporozoite surface antigens have been defined but they generally displayed a low level of immunogenicity (Florin-Christensen et al., 2014; Nene and Morrison, 2016; Nene et al., 2016). However, antibodies neutralizing sporozoite infectivity have been demonstrated in animals exposed to repeated sporozoite challenges (Nene and Morrison, 2016). On the contrary, very limited knowledge of intra-tick developmental stages restricts research on tick-pathogen interactions. Up to now, only few vaccine candidates were defined for species, particularly of the *Babesia sensu stricto* lineage, and partially also of the *B. microti* group (Hajdušek et al., 2013; de la Fuente et al., 2017) but there is lack of knowledge about *Theileria* and tick interactions. Although intra-tick development differs among all piroplasm evolutionary lineages, comparative bioinformatics analysis implies a high level of conservation of crucial regulatory domains responsible for piroplasm life cycle transitions (Alzan et al., 2016). Targeting key *Babesia* or *Theileria* developmental stages in tick tissues represents an attractive way toward transmission-blocking vaccines. However, this research requires in depth knowledge of parasite intra-tick development with a strong focus on conserved or divergent developmental features.

## AUTHOR CONTRIBUTIONS

MJ, OH, and LM conceived and designed the review. DS and PK contributed ideas and concepts. MJ did the literature search,



wrote the manuscript, and designed the figure. MJ, OH, DS, PK, and LM finalized the paper and figure.

## FUNDING

This work was primarily supported by the grants of Czech Science Foundation (GA CR) No. 17-27386S to OH and No. 17-14631S to DS. OH, DS, and PK were also supported by the

Centre for research of pathogenicity and virulence of parasites (No. CZ.02.1.01/0.0/0.0/16\_019/0000759) funded by European Regional Development Fund (ERDF) and Ministry of Education, Youth and Sport, Czech Republic (MEYS). MJ was additionally supported by GACR grant No. 17-27393S and by the Dual Czech-French PhD. program (Doctorat en co-tutelle) provided by the French Institute in Prague, Czech Republic. LM was supported by her lab's own resources.

## REFERENCES

- Ahmed, J. S., Schnittger, L., and Mehlhorn, H. (1999). Review: *Theileria* schizonts induce fundamental alterations in their host cells. *Parasitol. Res.* 85, 527–538. doi: 10.1007/s004360050592
- Alzan, H. F., Knowles, D. P., and Suarez, C. E. (2016). Comparative bioinformatics analysis of transcription factor genes indicates conservation of key regulatory domains among *Babesia bovis*, *Babesia microti*, and *Theileria equi*. *PLoS Negl. Trop. Dis.* 10:e0004983. doi: 10.1371/journal.pntd.0004983
- Arisue, N., and Hashimoto, T. (2014). Phylogeny and evolution of apicoplasts and apicomplexan parasites. *Parasitol. Int.* 64, 254–259. doi: 10.1016/j.parint.2014.10.005
- Asada, M., Goto, Y., Yahata, K., Yokoyama, N., Kawai, S., Inoue, N., et al. (2012). Gliding motility of *Babesia bovis* merozoites visualized by time-lapse video microscopy. *PLoS ONE* 7:e35227. doi: 10.1371/journal.pone.0035227
- Baneth, G., Florin-Christensen, M., Cardoso, L., and Schnittger, L. (2015). Reclassification of *Theileria annae* as *Babesia vulpes* sp. nov. *Parasit. Vectors* 8:207. doi: 10.1186/s13071-015-0830-5
- Bastos, R. G., Suarez, C. E., Laughery, J. M., Johnson, W. C., Ueti, M. W., and Knowles, D. P. (2013). Differential expression of three members of the multidomain adhesion CCp family in *Babesia bigemina*, *Babesia bovis* and *Theileria equi*. *PLoS ONE* 8:e67765. doi: 10.1371/journal.pone.0067765
- Becker, C. A., Malandrin, L., Depoix, D., Larcher, T., David, P. H., Chauvin, A., et al. (2010). Identification of three CCp genes in *Babesia divergens*: novel markers for sexual stages parasites. *Mol. Biochem. Parasitol.* 174, 36–43. doi: 10.1016/j.molbiopara.2010.06.011
- Becker, C. A., Malandrin, L., Larcher, T., Chauvin, A., Bischoff, E., and Bonnet, S. I. (2013). Validation of BdCCp2 as a marker for *Babesia divergens* sexual stages in ticks. *Exp. Parasitol.* 133, 51–56. doi: 10.1016/j.exppara.2012.10.007
- Binnington, K. C. (1978). Sequential changes in salivary gland structure during attachment and feeding of the cattle tick, *Boophilus microplus*. *Int. J. Parasitol.* 8, 97–115. doi: 10.1016/0020-7519(78)90004-8
- Bishop, R., Musoke, A., Morzaria, S., Gardner, M., and Nene, V. (2004). *Theileria*: intracellular protozoan parasites of wild and domestic ruminants transmitted by ixodid ticks. *Parasitology* 129, 271–283. doi: 10.1017/S0031182003004748
- Bishop, R. P., Hemmink, J. D., Morrison, W. I., Weir, W., Toye, P. G., Sitt, T., et al. (2015). The African buffalo parasite *Theileria* sp. (buffalo) can infect and immortalize cattle leukocytes and encodes divergent orthologues of *Theileria parva* antigen genes. *Int. J. Parasitol. Parasites Wildl.* 4, 333–342. doi: 10.1016/j.ijppaw.2015.08.006
- Blouin, E. F., and De Waal, D. T. (1989). The fine structure of developmental stages of *Babesia caballii* in the salivary glands of *Hyalomma truncatum*. *Onderstepoort J. Vet. Res.* 56, 189–193.
- Blouin, E. F., Kocan, A. A., Kocan, K. M., and Hair, J. (1987). Evidence of a limited schizogonous cycle for *Cytauxzoon felis* in bobcats following exposure to infected ticks. *J. Wildl. Dis.* 23, 499–501. doi: 10.7589/0090-3558-23.3.499
- Blouin, E. F., and van Rensburg, L. (1988). An ultrastructural study of the development of *Babesia occultans* in the salivary glands of adult *Hyalomma marginatum rufipes*. *Onderstepoort J. Vet. Res.* 55, 93–100.
- Bock, R., Jackson, L., de Vos, A., and Jorgensen, W. (2004). Babesiosis of cattle. *Parasitol.* 129(Suppl.) S247–S269. doi: 10.1017/S0031182004005190
- Boldbaatar, D., Battsetseg, B., Matsuo, T., Hatta, T., Umeyama-Shirafuji, R., Xuan, X., et al. (2008). Tick vitellogenin receptor reveals critical role in oocyte development and transovarial transmission of *Babesia* parasite. *Biochem. Cell Biol.* 86, 331–344. doi: 10.1139/O08-071
- Boldbaatar, D., Umeyama-Shirafuji, R., Liao, M., Tanaka, T., Xuan, X., and Fujisaki, K. (2010). Multiple vitellogenins from the *Haemaphysalis longicornis* tick are crucial for ovarian development. *J. Insect Physiol.* 56, 1587–1598. doi: 10.1016/j.jinsphys.2010.05.019
- Bonnet, S., Jouglin, M., Malandrin, L., Becker, C., Agoulon, A., Lhostis, M., et al. (2007). Transstadial and transovarial persistence of *Babesia divergens* DNA in *Ixodes ricinus* ticks fed on infected blood in a new skin-feeding technique. *Parasitology* 134, 197–207. doi: 10.1017/S0031182006001545
- Burki, F., Inagaki, Y., Bråte, J., Archibald, J. M., Keeling, P. J., Cavalier-Smith, T., et al. (2009). Large-scale phylogenomic analyses reveal that two enigmatic protist lineages, telonemia and centroheliozoa, are related to photosynthetic chromalveolates. *Genome Biol. Evol.* 1, 231–238. doi: 10.1093/gbe/evp022
- Christophers, S. R. (1907). Preliminary note on the development of *Piroplasma canis* in the tick. *Br. Med. J.* 1, 76–78. doi: 10.1136/bmj.1.2402.76
- Clancey, N., Horney, B., Burton, S., Birkenheuer, A., McBurney, S., and Tefft, K. (2010). *Babesia (Theileria) annae* in a red fox (*Vulpes vulpes*) from Prince Edward Island, Canada. *J. Wildl. Dis.* 46, 615–621. doi: 10.7589/0090-3558-46.2.615
- Conrad, P. A., Denham, D., and Brown, C. G. (1986). Intraerythrocytic multiplication of *Theileria parva* *in vitro*: an ultrastructural study. *Int. J. Parasitol.* 16, 223–229. doi: 10.1016/0020-7519(86)90047-0
- Conrad, P. A., Kelly, B. G., and Brown, C. G. (1985). Intraerythrocytic schizogony of *Theileria annulata*. *Parasitology* 91, 67–82. doi: 10.1017/S0031182000056523
- Coons, L. B., and Roshdy, M. A. (1973). Fine structure of the salivary glands of unfed male *Dermacentor variabilis* (Say) (Ixodoidea: Ixodidae). *J. Parasitol.* 59, 900–912. doi: 10.2307/3278433
- de la Fuente, J., Antunes, S., Bonnet, S., Cabezas-Cruz, A., Domingos, A. G., Estrada-Peña, A., et al. (2017). Tick-pathogen interactions and vector competence: identification of molecular drivers for tick-borne diseases. *Front. Cell. Infect. Microbiol.* 7:114. doi: 10.3389/fcimb.2017.00114
- Del Carmen Terrón, M., González-Camacho, F., González, L. M., Luque, D., and Montero, E. (2016). Ultrastructure of the *Babesia divergens* free merozoite. *Ticks Tick Borne Dis.* 7, 1274–1279. doi: 10.1016/j.ttbdis.2016.07.001
- Dobbelaere, D., and Heussler, V. (1999). Transformation of leukocytes by *Theileria parva* and *T. annulata*. *Annu. Rev. Microbiol.* 53, 1–42. doi: 10.1146/annurev.micro.53.1.1
- Dobbelaere, D. A., and Küenzi, P. (2004). The strategies of the *Theileria* parasite: a new twist in host-pathogen interactions. *Curr. Opin. Immunol.* 16, 524–530. doi: 10.1016/j.coi.2004.05.009
- Dobbelaere, D. A., and Rottenberg, S. (2003). *Theileria*-induced leukocyte transformation. *Curr. Opin. Microbiol.* 6, 377–382. doi: 10.1016/S1369-5274(03)00085-7
- Dobbelaere, D. A., Webster, P., Leitch, B. L., Voigt, W. P., and Irvin, A. D. (1985). *Theileria parva*: expression of a sporozoite surface coat antigen. *Exp. Parasitol.* 60, 90–100. doi: 10.1016/S0014-4894(85)80026-6
- Donnelly, J., and Peirce, M. A. (1975). Experiments on the transmission of *Babesia divergens* to cattle by the tick *Ixodes ricinus*. *Int. J. Parasitol.* 5, 363–367. doi: 10.1016/0020-7519(75)90085-5
- Dubremetz, J. F., Garcia-Réguet, N., Conseil, V., and Fourmaux, M. N. (1998). Apical organelles and host-cell invasion by Apicomplexa. *Int. J. Parasitol.* 28, 1007–1013. doi: 10.1016/S0020-7519(98)00076-9
- Fawcett, D., Musoke, A., and Voigt, W. (1984). Interaction of sporozoites of *Theileria parva* with bovine lymphocytes *in vitro*. I. Early events after invasion. *Tissue Cell* 16, 873–884. doi: 10.1016/0040-8166(84)90068-5

- Fawcett, D. W., Büscher, G., and Doxsey, S. (1982a). Salivary gland of the tick vector of East Coast fever. III. The ultrastructure of sporogony in *Theileria parva*. *Tissue Cell* 14, 183–206. doi: 10.1016/0040-8166(82)90017-9
- Fawcett, D. W., Büscher, G., and Doxsey, S. (1982b). Salivary gland of the tick vector of East Coast fever. IV. Cell type selectivity and host cell responses to *Theileria parva*. *Tissue Cell* 14, 397–414. doi: 10.1016/0040-8166(82)90035-0
- Fawcett, D. W., Conrad, P. A., Grootenhuys, J. G., and Morzaria, S. P. (1987). Ultrastructure of the intra-erythrocytic stage of *Theileria* species from cattle and waterbuck. *Tissue Cell* 19, 643–655. doi: 10.1016/0040-8166(87)90071-1
- Fawcett, D. W., Doxsey, S., Stagg, D. A., and Young, A. S. (1982c). The entry of sporozoites of *Theileria parva* into bovine lymphocytes *in vitro*. Electron microscopic observations. *Eur. J. Cell Biol.* 27, 10–21.
- Florin-Christensen, M., Suarez, C. E., Rodriguez, A. E., Flores, D. A., and Schnittger, L. (2014). Vaccines against bovine babesiosis: where we are now and possible roads ahead. *Parasitology* 28, 1–30. doi: 10.1017/S0031182014000961
- Friedhoff, K., Scholtyssek, E., and Weber, G. (1972). Fine structure of the merozoites of *Babesia ovis* in the salivary glands of female ticks. *Z. Parasitenkd.* 38, 132–140. doi: 10.1007/BF00329024
- Friedhoff, K. T., and Büscher, G. (1976). Rediscovery of Koch's "strahlenörper" of *Babesia bigemina*. *Z. Parasitenkd.* 50, 345–347. doi: 10.1007/BF02462979
- Gauer, M., Mackenstedt, U., Mehlhorn, H., Schein, E., Zapf, F., Njenga, E., et al. (1995). DNA measurements and ploidy determination of developmental stages in the life cycles of *Theileria annulata* and *T. parva*. *Parasitol. Res.* 81, 565–574. doi: 10.1007/BF00932023
- Gohil, S., Herrmann, S., Günther, S., and Cooke, B. M. (2013). Bovine babesiosis in the 21st century: advances in biology and functional genomics. *Int. J. Parasitol.* 43, 125–132. doi: 10.1016/j.ijpara.2012.09.008
- Gorenflot, A., Brasseur, P., Precigout, E., L'Hostis, M., Marchand, A., and Schrevel, J. (1991). Cytological and immunological responses to *Babesia divergens* in different hosts: ox, gerbil, man. *Parasitol. Res.* 77, 3–12. doi: 10.1007/BF00934377
- Gorenflot, A., Precigout, E., Valentin, A., Bissuel, G., Carcy, B., Brasseur, P., et al. (1992). *Babesia divergens* vaccine. *Mem. Inst. Oswaldo Cruz.* 87, 279–281. doi: 10.1590/S0074-02761992000700047
- Gough, J. M., Jorgensen, W. K., and Kemp, D. H. (1998). Development of tick gut forms of *Babesia bigemina* *in vitro*. *J. Eukaryot. Microbiol.* 45, 298–306. doi: 10.1111/j.1550-7408.1998.tb04540.x
- Guimarães, A. M., Lima, J. D., and Ribeiro, M. F. B. (1998a). Sporogony and experimental transmission of *Babesia equi* by *Boophilus microplus*. *Parasitol. Res.* 84, 323–327. doi: 10.1007/s004360050404
- Guimarães, A. M., Lima, J. D., Ribeiro, M. F. B., Camargos, E. R. S., and Bozzi, I. A. (1998b). Ultrastructure of sporogony in *Babesia equi* in salivary glands of adult female *Boophilus microplus* ticks. *Parasitol. Res.* 84, 69–74.
- Guimarães, A. M., Lima, J. D., and Ribeiro, M. F. B. (2003). Ultrastructure of *Babesia equi* trophozoites isolated in Minas Gerais, Brazil. *Pesq. Vet. Bras.* 23, 101–104. doi: 10.1590/S0100-736X2003000300002
- Hagiwara, K., Takahashi, K., Taniyama, H., Kawamoto, S., Kurosawa, T., Ikuta, K., et al. (1997). Detection of *Theileria sergenti* schizonts in bovine lymph node. *Int. J. Parasitol.* 27, 1375–1378. doi: 10.1016/S0020-7519(97)00092-1
- Hajdušek, O., Síma, R., Ayllón, N., Jalovecká, M., Perner, J., de la Fuente, J., et al. (2013). Interaction of the tick immune system with transmitted pathogens. *Front. Cell. Infect. Microbiol.* 3:26. doi: 10.3389/fcimb.2013.00026
- Hazen-Karr, C. G., Kocan, A. A., Kocan, K. M., and Hair, J. A. (1987). The ultrastructure of sporogony in *Theileria cervi* (Bettencourt et al., 1907) in salivary glands of female *Amblyomma americanum* (L.) ticks. *J. Parasitol.* 73, 1182–1188. doi: 10.2307/3282304
- Higuchi, S., Izumitani, M., Hoshi, H., Kawamura, S., and Yasuda, Y. (1999a). Development of *Babesia gibsoni* in the midgut of larval tick, *Rhipicephalus sanguineus*. *J. Vet. Med. Sci.* 61, 689–691. doi: 10.1292/jvms.61.689
- Higuchi, S., Konno, H., Hoshi, F., Kawamura, S., and Yasuda, Y. (1993). Observations of *Babesia gibsoni* in the ovary of the tick, *Haemaphysalis longicornis*. *Arch. Exp. Med.* 65, 153–158.
- Higuchi, S., Kuroda, H., Hoshi, H., Kawamura, S., and Yasuda, Y. (1999b). Development of *Babesia gibsoni* in the midgut of the nymphal stage of the tick, *Rhipicephalus sanguineus*. *J. Vet. Med. Sci.* 61, 697–699. doi: 10.1292/jvms.61.697
- Higuchi, S., Oya, H., Hoshi, F., Kawamura, S., and Yasuda, Y. (1992). Observations of *Babesia gibsoni* in midgut epithelial cells of the tick, *Haemaphysalis longicornis*. *Arch. Exp. Med.* 65, 143–147.
- Higuchi, S., Oya, H., Hoshi, F., Kawamura, S., and Yasuda, Y. (1994). Development of *Babesia ovata* in the salivary glands of the nymphal tick, *Haemaphysalis longicornis*. *J. Vet. Med. Sci.* 56, 207–209. doi: 10.1292/jvms.56.207
- Higuchi, S., Simomura, S., Yoshida, H., Hoshi, F., Kawamura, S., and Yasuda, Y. (1991a). Development of *Babesia gibsoni* in the gut epithelium of the tick, *Haemaphysalis longicornis*. *J. Vet. Med. Sci.* 53, 129–131. doi: 10.1292/jvms.53.129
- Higuchi, S., Simomura, S., Yoshida, H., Hoshi, F., Kawamura, S., and Yasuda, Y. (1991b). Development of *Babesia gibsoni* in the hemolymph of the vector tick, *Haemaphysalis longicornis*. *J. Vet. Med. Sci.* 53, 491–493. doi: 10.1292/jvms.53.491
- Holbrook, A. A., Anthony, D. W., and Johnson, A. J. (1968). Observations on the development of *Babesia caballi* (Nuttall) in the tropical horse tick *Dermacentor nitens* Neumann. *J. Protozool.* 15, 391–396. doi: 10.1111/j.1550-7408.1968.tb02143.x
- Homer, M. J., Aguilar-Delfin, I., Telford, S. R., Krause, P. J., and Persing, D. H. (2000). Babesiosis. *Clin. Microbiol. Rev.* 13, 451–469. doi: 10.1128/CMR.13.3.451-469.2000
- Hunfeld, K. P., Hildebrandt, A., and Gray, J. S. (2008). Babesiosis: recent insights into an ancient disease. *Int. J. Parasitol.* 38, 1219–1237. doi: 10.1016/j.ijpara.2008.03.001
- Jalovecka, M., Bonsergent, C., Hajdusek, O., Kopacek, P., and Malandrin, L. (2016). Stimulation and quantification of *Babesia divergens* gametocytogenesis. *Parasit. Vectors* 9:439. doi: 10.1186/s13071-016-1731-y
- Janouskovec, J., Horák, A., Obornik, M., Lukes, J., and Keeling, P. J. (2010). A common red algal origin of the apicomplexan, dinoflagellate, and heterokont plastids. *Proc. Natl. Acad. Sci. U.S.A.* 107, 10949–10954. doi: 10.1073/pnas.1003335107
- Joyner, L. P., Davies, S. F., and Kendall, S. B. (1963). The experimental transmission of *Babesia divergens* by *Ixodes ricinus*. *Exp. Parasitol.* 14, 367–373. doi: 10.1016/0014-4894(63)90044-4
- Kappmeyer, L. S., Thiagarajan, M., Herndon, D. R., Ramsay, J. D., Caler, E., Djikeng, A., et al. (2012). Comparative genomic analysis and phylogenetic position of *Theileria equi*. *BMC Genomics.* 13:603. doi: 10.1186/1471-2164-13-603
- Karakashian, S. J., Rudzinska, M. A., Spielman, A., Lewengrub, S., and Campbell, J. (1986). Primary and secondary ookinetes of *Babesia microti* in the larval and nymphal stages of the tick *Ixodes dammini*. *Can. J. Zool.* 64, 328–339. doi: 10.1139/z86-053
- Karakashian, S. J., Rudzinska, M. A., Spielman, A., Lewengrub, S., Piesman, J., and Shoukrey, N. (1983). Ultrastructural studies on sporogony of *Babesia microti* in salivary gland cells of the tick *Ixodes dammini*. *Cell Tissue Res.* 231, 275–287. doi: 10.1007/BF00222180
- Kawai, S., Igarashi, I., Abgaandorjiin, A., Ikadai, H., Omata, Y., Saito, A., et al. (1999a). Tubular structures associated with *Babesia caballi* in equine erythrocytes *in vitro*. *Parasitol. Res.* 85, 171–175. doi: 10.1007/s004360050530
- Kawai, S., Igarashi, I., Abgaandorjiin, A., Miyazawa, K., Ikadai, H., Nagasawa, et al. (1999b). Ultrastructural characteristics of *Babesia caballi* in equine erythrocytes *in vitro*. *Parasitol. Res.* 85, 794–799. doi: 10.1007/s004360050635
- Kawai, S., Takahashi, K., Kawamoto, S., Nagahara, A., Sonoda, M., Kurosawa, T., et al. (1989). Bar-structure in bovine erythrocytes infected with *Theileria sergenti*. *Nippon Juigaku Zasshi* 51, 1219–1225. doi: 10.1292/jvms1939.51.1219
- Kawai, S., Takahashi, K., Sonoda, M., and Kurosawa, T. (1986). Ultrastructure of intra-erythrocytic stages of *Babesia ovata*. *Nippon Juigaku Zasshi* 48, 943–949. doi: 10.1292/jvms1939.48.943
- Kjemtrup, A. M., and Conrad, P. A. (2000). Human babesiosis: an emerging tick-borne disease. *Int. J. Parasitol.* 30, 1323–1337. doi: 10.1016/S0020-7519(00)00137-5
- Kjemtrup, A. M., Wainwright, K., Miller, M., Penzhorn, B. L., and Carreno, R. A. (2006). *Babesia conradae*, sp. Nov., a small canine *Babesia* identified in California. *Vet. Parasitol.* 138, 103–111. doi: 10.1016/j.vetpar.2006.01.044
- Leiby, D. A. (2011). Transfusion-transmitted *Babesia* spp.: bull's-eye on *Babesia microti*. *Clin. Microbiol. Rev.* 24, 14–28. doi: 10.1128/CMR.00022-10
- Lempereur, L., Larcombe, S. D., Durrani, Z., Karagenc, T., Bilgic, H. B., Bakirci, S., et al. (2017). Identification of candidate transmission-blocking antigen genes

- in *Theileria annulata* and related vector-borne apicomplexan parasites. *BMC Genomics* 18:438. doi: 10.1186/s12864-017-3788-1
- Lewis, D., Purnell, R. E., and Shaw, S. R. (1980). The isolation and characterization of human and bovine strains of *Babesia divergens* from Drummedrochit, Scotland. *Parasitology* 81, 145–155. doi: 10.1017/S003118200055116
- Lewis, D., and Young, E. R. (1980). The transmission of a human strain of *Babesia divergens* by *Ixodes ricinus* ticks. *J. Parasitol.* 66, 359–360. doi: 10.2307/3280841
- Lobo, C. A., Rodriguez, M., and Cursino-Santos, J. R. (2012). *Babesia* and red cell invasion. *Curr. Opin. Hematol.* 19, 170–175. doi: 10.1097/MOH.0b013e328352245a
- MacKenstedt, U., Gauer, M., Fuchs, P., Zapf, F., Schein, E., and Mehlhorn, H. (1995). DNA measurements reveal differences in the life cycles of *Babesia bigemina* and *B. canis*, two typical members of the genus *Babesia*. *Parasitol. Res.* 81, 595–604. doi: 10.1007/BF00932027
- Mackenstedt, U., Gauer, M., Mehlhorn, H., Schein, E., and Hauschild, S. (1990). Sexual cycle of *Babesia divergens* confirmed by DNA measurements. *Parasitol. Res.* 76, 199–206. doi: 10.1007/BF00930815
- Malandrin, L., Jouglin, M., Sun, Y., Brisseau, N., and Chauvin, A. (2010). Redescription of *Babesia capreoli* (Enigk and Friedhoff, 1962) from roe deer (*Capreolus capreolus*): isolation, cultivation, host specificity, molecular characterisation and differentiation from *Babesia divergens*. *Int. J. Parasitol.* 40, 277–284. doi: 10.1016/j.ijpara.2009.08.008
- Mehlhorn, H., Moltmann, U., Schein, E., and Voigt, W. P. (1981). Fine structure of supposed gametes and syngamy of *Babesia canis* (Piroplasma) after *in-vitro* development. *Zentralbl. Bakteriol. Mikrobiol. Hyg. A.* 250, 248–255.
- Mehlhorn, H., and Schein, E. (1993). The piroplasms: “A long story in short” or “Robert Koch has seen it”. *Eur. J. Protistol.* 29, 279–293. doi: 10.1016/S0932-4739(11)80371-8
- Mehlhorn, H., and Schein, E. (1998). Redescription of *Babesia equi* Laveran, 1901 as *Theileria equi* Mehlhorn, Schein 1998. *Parasitol. Res.* 84, 467–475. doi: 10.1007/s004360050431
- Mehlhorn, H., Schein, E., and Voigt, W. P. (1980). Light and electron microscopic study on developmental stages of *Babesia canis* within the gut of the tick *Dermacentor reticulatus*. *J. Parasitol.* 66, 220–228. doi: 10.2307/3280808
- Mehlhorn, H., Schein, E., and Warnecke, M. (1978). Electron microscopic studies on the development of kinetes of *Theileria parva* Theiler, 1904 in the gut of the vector ticks *Rhipicephalus appendiculatus* Neumann, 1901. *Acta Trop.* 35, 123–136.
- Mehlhorn, H., Schein, E., and Warnecke, M. (1979). Electron-microscopic studies on *Theileria ovis* Rodhain, 1916: development of kinetics in the gut of the vector tick, *Rhipicephalus eversti eversti* Neumann, 1897, and their transformation within cells of the salivary glands. *J. Protozool.* 26, 377–385. doi: 10.1111/j.1550-7408.1979.tb04640.x
- Mehlhorn, H., and Schein, E. (1984). The piroplasms: life cycle and sexual stages. *Adv. Parasitol.* 23, 37–103. doi: 10.1016/S0065-308X(08)60285-7
- Moltmann, U. G., Mehlhorn, H., and Friedhoff, K. T. (1982a). Electron microscopic study on the development of *Babesia ovis* (Piroplasmia) in the salivary glands of the vector tick *Rhipicephalus bursa*. *Acta Trop.* 39, 29–40.
- Moltmann, U. G., Mehlhorn, H., and Friedhoff, K. T. (1982b). Ultrastructural study of the development of *Babesia ovis* (Piroplasmia) in the ovary of the vector tick *Rhipicephalus bursa*. *J. Protozool.* 29, 30–38. doi: 10.1111/j.1550-7408.1982.tb02877.x
- Moltmann, U. G., Mehlhorn, H., Schein, E., Rehbein, G., Voigt, W. P., and Zwegarth, E. (1983a). Fine structure of *Babesia equi* Laveran, 1901 within lymphocytes and erythrocytes of horses: an *in vivo* and *in vitro* study. *J. Parasitol.* 69, 111–120. doi: 10.2307/3281285
- Moltmann, U. G., Mehlhorn, H., Schein, E., Voigt, W. P., and Friedhoff, K. T. (1983b). Ultrastructural study on the development of *Babesia equi* (Coccidia: Piroplasmia) in the salivary glands of its vector ticks. *J. Protozool.* 30, 218–225. doi: 10.1111/j.1550-7408.1983.tb02907.x
- Montero, E., Rodriguez, M., Oksov, Y., and Lobo, C. A. (2009). *Babesia divergens* apical membrane antigen 1 and its interaction with the human red blood cell. *Infect. Immun.* 77, 4783–4793. doi: 10.1128/IAI.00969-08
- Mosqueda, J., Ramos, J. A., Falcon, A., Alvarez, J. A., Aragon, V., and Figueroa, J. V. (2004). *Babesia bigemina*: sporozoite isolation from *Boophilus microplus* nymphs and initial immunomolecular characterization. *Ann. N.Y. Acad. Sci.* 1026, 222–231. doi: 10.1196/annals.1307.034
- Nene, V., Kiara, H., Lacasta, A., Pelle, R., Svitek, N., and Steinaa, L. (2016). The biology of *Theileria parva* and control of East Coast fever - Current status and future trends. *Ticks Tick Borne Dis.* 7, 549–564. doi: 10.1016/j.ttbdis.2016.02.001
- Nene, V., and Morrison, W. I. (2016). Approaches to vaccination against *Theileria parva* and *Theileria annulata*. *Parasite Immunol.* 38, 724–734. doi: 10.1111/pim.12388
- Piesman, J., Karakashian, S. J., Lewengrub, S., Rudzinska, M. A., and Spielman, A. (1986). Development of *Babesia microti* sporozoites in adult *Ixodes dammini*. *Int. J. Parasitol.* 16, 381–385. doi: 10.1016/0020-7519(86)90118-9
- Piesko, M., Weir, W., Goodhead, I., Kinnaird, J., and Shiels, B. (2015). ApiAP2 Factors as Candidate regulators of stochastic commitment to merozoite production in *Theileria annulata*. *PLoS Negl. Trop. Dis.* 9:e0003933. doi: 10.1371/journal.pntd.0003933
- Potgieter, F. T., and Els, H. J. (1976). Light and electron microscopic observations on the development of small merozoites of *Babesia bovis* in *Boophilus microplus* larvae. *Onderstepoort J. Vet. Res.* 43, 123–128.
- Potgieter, F. T., and Els, H. J. (1977a). Light and electron microscopic observations on the development of *Babesia bigemina* in larvae, nymphae and non-replete females of *Boophilus decoloratus*. *Onderstepoort J. Vet. Res.* 44, 213–231.
- Potgieter, F. T., and Els, H. J. (1977b). The fine structure of intra-erythrocytic stages of *Babesia bigemina*. *Onderstepoort J. Vet. Res.* 44, 157–168.
- Potgieter, F. T., Els, H. J., and Vuuren, A. S. (1976). The fine structure of merozoites of *Babesia bovis* in the gut epithelium of *Boophilus microplus*. *Onderstepoort J. Vet. Res.* 43, 1–9.
- Purnell, R. E., and Joyner, L. P. (1968). The development of *Theileria parva* in the salivary glands of the tick, *Rhipicephalus appendiculatus*. *Parasitology.* 58, 725–732. doi: 10.1017/S0031182000029036
- Ramsay, J. D., Ueti, M. W., Johnson, W. C., Scoles, G. A., Knowles, D. P., and Mealey, R. H. (2013). Lymphocytes and macrophages are infected by *Theileria equi*, but T cells and B cells are not required to establish infection *in vivo*. *PLoS ONE* 8:e76996. doi: 10.1371/journal.pone.0076996
- Rudzinska, M. A. (1976). Ultrastructure of intraerythrocytic *Babesia microti* with emphasis on the feeding mechanism. *J. Protozool.* 23, 224–233. doi: 10.1111/j.1550-7408.1976.tb03759.x
- Rudzinska, M. A., Lewengrub, S., Spielman, A., and Piesman, J. (1983a). Invasion of *Babesia microti* into epithelial cells of the tick gut. *J. Protozool.* 30, 338–346. doi: 10.1111/j.1550-7408.1983.tb02927.x
- Rudzinska, M. A., Spielman, A., Lewengrub, S., Piesman, J., and Karakashian, S. (1982). Penetration of the peritrophic membrane of the tick by *Babesia microti*. *Cell Tissue Res.* 221, 471–481. doi: 10.1007/BF00215696
- Rudzinska, M. A., Spielman, A., Lewengrub, S., Piesman, J., and Karakashian, S. (1984). The sequence of developmental events of *Babesia microti* in the gut of *Ixodes dammini*. *Protistologica* 4, 649–663.
- Rudzinska, M. A., Spielman, A., Lewengrub, S., Trager, W., and Piesman, J. (1983b). Sexuality in piroplasms as revealed by electron microscopy in *Babesia microti*. *Proc. Natl. Acad. Sci. U.S.A.* 80, 2966–2970. doi: 10.1073/pnas.80.10.2966
- Rudzinska, M. A., Spielman, A., Riek, R. F., Lewengrub, S. J., and Piesman, J. (1979). Intraerythrocytic ‘gametocytes’ of *Babesia microti* and their maturation in ticks. *Can. J. Zool.* 57, 424–434. doi: 10.1139/z79-050
- Rudzinska, M. A., and Trager, W. (1962). Intracellular phagotrophy in *Babesia rodhaini* as revealed by electron microscopy. *J. Protozool.* 9, 279–288. doi: 10.1111/j.1550-7408.1962.tb02621.x
- Rudzinska, M. A., and Trager, W. (1977). Formation of merozoites in intraerythrocytic *Babesia microti*: an ultrastructural study. *Can. J. Zool.* 55, 928–938. doi: 10.1139/z77-121
- Rudzinska, M. A., Trager, W., Lewengrub, S. J., and Gubert, E. (1976). An electron microscopic study of *Babesia microti* invading erythrocytes. *Cell Tissue Res.* 169, 323–334. doi: 10.1007/BF00219605
- Sato, M., Kamio, T., Tanaka, S., Taniguchi, T., and Fujisaki, K. (1994). Development of *Theileria sergenti* schizonts in the lymph node of experimentally infected cattle. *J. Vet. Med. Sci.* 56, 715–722. doi: 10.1292/jvms.56.715
- Schein, E., and Friedhoff, K. T. (1978). Light microscopic studies on the development of *Theileria annulata* (Dschunkowsky and Luhs, 1904) in *Hyalomma anatolicum excavatum* (Koch, 1844): the development in haemolymph and salivary glands. *Z. Parasitenkd.* 56, 287–303. doi: 10.1007/BF00931721

- Schein, E., Mehlhorn, H., and Voigt, W. P. (1979). Electron microscopical studies on the development of *Babesia canis* (Sporozoa) in the salivary glands of the vector tick *Dermacentor reticulatus*. *Acta Trop.* 36, 229–241.
- Schein, E., Rehbein, G., Voigt, W. P., and Zweygarth, E. (1981). *Babesia equi* (Laveran 1901) development in horses and in lymphocyte culture. *Tropenmed. Parasitology* 32, 223–227.
- Schein, E., Warnecke, M., and Kirmse, P. (1977). Development of *Theileria parva* (Theiler, 1904) in the gut of *Rhipicephalus appendiculatus* (Neumann, 1901). *Parasitology* 75, 309–316. doi: 10.1017/S0031182000051854
- Schnittger, L., Rodriguez, A. E., Florin-Christensen, M., and Morrison, D. A. (2012). *Babesia*: a world emerging. *Infect. Genet. Evol.* 12, 1788–1809. doi: 10.1016/j.meegid.2012.07.004
- Schreeg, M. E., Marr, H. S., Tarigo, J. L., Cohn, L. A., Bird, D. M., Scholl, E. H., et al. (2016). Mitochondrial genome sequences and structures aid in the resolution of piroplasmida phylogeny. *PLoS ONE* 11:e0165702. doi: 10.1371/journal.pone.0165702
- Shaw, M. K. (1995). Mobilization of intrasporozoite Ca<sup>2+</sup> is essential for *Theileria parva* sporozoite invasion of bovine lymphocytes. *Eur. J. Cell Biol.* 68, 78–87.
- Shaw, M. K. (1996a). Characterization of the parasite-host cell interactions involved in *Theileria parva* sporozoite invasion of bovine lymphocytes. *Parasitology* 113, 267–277.
- Shaw, M. K. (1996b). *Theileria parva* sporozoite entry into bovine lymphocytes involves both parasite and host cell signal transduction processes. *Exp. Parasitol.* 84, 344–354. doi: 10.1006/expr.1996.0123
- Shaw, M. K. (1997). The same but different: the biology of *Theileria* sporozoite entry into bovine cells. *Int. J. Parasitol.* 27, 457–474. doi: 10.1016/S0020-7519(97)00015-5
- Shaw, M. K. (1999). *Theileria parva*: sporozoite entry into bovine lymphocytes is not dependent on the parasite cytoskeleton. *Exp. Parasitol.* 92, 24–31. doi: 10.1006/expr.1998.4393
- Shaw, M. K. (2003). Cell invasion by *Theileria* sporozoites. *Trends Parasitol.* 19, 2–6. doi: 10.1016/S1471-4922(02)00015-6
- Shaw, M. K., and Tilney, L. G. (1992). How individual cells develop from a syncytium: merogony in *Theileria parva* (Apicomplexa). *J. Cell. Sci.* 101, 109–123.
- Shaw, M. K., and Tilney, L. G. (1995). The entry of *Theileria parva* merozoites into bovine erythrocytes occurs by a process similar to sporozoite invasion of lymphocytes. *Parasitology* 111, 455–461. doi: 10.1017/S0031182000065951
- Shaw, M. K., Tilney, L. G., and McKeever, D. J. (1993). Tick salivary gland extract and interleukin-2 stimulation enhance susceptibility of lymphocytes to infection by *Theileria parva* sporozoites. *Infect. Immun.* 61, 1486–1495.
- Shaw, M. K., Tilney, L. G., and Musoke, A. J. (1991). The entry of *Theileria parva* sporozoites into bovine lymphocytes: evidence for MHC class I involvement. *J. Cell Biol.* 113, 87–101. doi: 10.1083/jcb.113.1.87
- Simpson, C. F., Bild, C. E., and Stolkier, H. E. (1963). Electron microscopy of canine and equine *Babesia*. *Am. J. Vet. Res.* 24, 408–414.
- Simpson, C. F., and Neal, F. C. (1980). Ultrastructure of *Babesia equi* in ponies treated with imidocarb. *Am. J. Vet. Res.* 41, 267–271.
- Sivakumar, T., Hayashida, K., Sugimoto, C., and Yokoyama, N. (2014). Evolution and genetic diversity of *Theileria*. *Infect. Genet. Evol.* 27, 250–263. doi: 10.1016/j.meegid.2014.07.013
- Soldati, D., Foth, B. J., and Cowman, A. F. (2004). Molecular and functional aspects of parasite invasion. *Trends Parasitol.* 20, 567–574. doi: 10.1016/j.pt.2004.09.009
- Sonenshine, D. E. (1991). *Biology of Ticks*. New York, NY: Oxford University Press.
- Sun, Y., Moreau, E., Chauvin, A., and Malandrin, L. (2011). The invasion process of bovine erythrocyte by *Babesia divergens*: knowledge from an *in vitro* assay. *Vet. Res.* 42:62. doi: 10.1186/1297-9716-42-62
- Susta, L., Torres-Velez, F., Zhang, J., and Brown, C. (2009). An *in situ* hybridization and immunohistochemical study of cytauxzoonosis in domestic cats. *Vet. Pathol.* 46, 1197–1204. doi: 10.1354/vp.08-VP-0132-B-FL
- Takahashi, K., Kawai, S., Yaehata, K., Kawamoto, S., Hagiwara, K., Kurosawa, T., et al. (1993). Sporogony of *Theileria sergenti* in the salivary glands of the tick vector *Haemaphysalis longicornis*. *Parasitol. Res.* 79, 1–7. doi: 10.1007/BF00931210
- Uilenberg, G. (2006). *Babesia*—a historical overview. *Vet. Parasitol.* 138, 3–10. doi: 10.1016/j.vetpar.2006.01.035
- Vannier, E. G., Diuk-Wasser, M. A., Ben Mamoun, C., and Krause, P. J. (2015). Babesiosis. *Infect. Dis. Clin. North Am.* 29, 357–370. doi: 10.1016/j.idc.2015.02.008
- Votycka, J., Modry, D., Obornik, M., Slapeta, J., and Lukes, J. (2017). “Apicomplexa,” in *Handbook of the Protists*, eds J. M. Archibald, A. G. B. Simpson, and C. H. Slamovits (Cham: Springer International Publishing AG), 1–58. doi: 10.1007/978-3-319-32669-6\_20-1
- Ward, P. A., and Jack, R. M. (1981). The entry process of *Babesia* merozoites into red cells. *Am. J. Pathol.* 102, 109–113.
- Warnecke, M., Schein, E., Voigt, W. P., Uilenberg, G., and Young, A. S. (1980). Development of *Theileria mutans* (Theiler, 1906) in the gut and the haemolymph of the tick *Amblyomma variegatum* (Fabricius, 1794). *Z. Parasitenkd.* 62, 119–125. doi: 10.1007/BF00927858
- Weber, G., and Friedhoff, K. (1979). Electron microscopic detection of initial and some subsequent developmental stages of *Babesia bigemina* in salivary glands of ticks. *Z. Parasitenkd.* 58, 191–194. doi: 10.1007/BF01951346
- Weber, G., and Friedhoff, K. T. (1977). Preliminary observations on the ultrastructure of supposed sexual stages of *Babesia bigemina* (Piroplasma). *Z. Parasitenkd.* 53, 83–92. doi: 10.1007/BF00383118
- Weber, G., and Walter, G. (1980). *Babesia microti* (Apicomplexa, Piroplasmida) - electron-microscope detection in salivary glands of the tick vector *Ixodes ricinus* (Ixodoidea, Ixodidae). *Z. Parasitenkd.* 64, 113–115. doi: 10.1007/BF00927061
- Webster, P., Dobbelaere, D. A., and Fawcett, D. W. (1985). The entry of sporozoites of *Theileria parva* into bovine lymphocytes *in vitro*. Immunoelectron microscopic observations. *Eur. J. Cell Biol.* 36, 157–162.
- Wise, L. N., Kappmeyer, L. S., Mealey, R. H., and Knowles, D. P. (2013). Review of equine piroplasmiasis. *J. Vet. Intern. Med.* 27, 1334–1346. doi: 10.1111/jvim.12168
- Yabsley, M. J., and Shock, B. C. (2013). Natural history of Zoonotic *Babesia*: role of wildlife reservoirs. *Int. J. Parasitol. Parasites Wildl.* 2, 18–31. doi: 10.1016/j.ijppaw.2012.11.003
- Yano, Y., Saito-Ito, A., Anchalee, D., and Takada, N. (2005). Japanese *Babesia microti* cytologically detected in salivary glands of naturally infected tick *Ixodes ovatus*. *Microbiol. Immunol.* 49, 891–897. doi: 10.1111/j.1348-0421.2005.tb03680.x
- Zapf, F., and Schein, E. (1994a). New findings in the development of *Babesia (Theileria) equi* (Laveran, 1901) in the salivary glands of the vector ticks, *Hyalomma* species. *Parasitol. Res.* 80, 543–548. doi: 10.1007/BF00933000
- Zapf, F., and Schein, E. (1994b). The development of *Babesia (Theileria) equi* (Laveran, 1901) in the gut and the haemolymph of the vector ticks, *Hyalomma* species. *Parasitol. Res.* 80, 297–302. doi: 10.1007/BF02351869
- Zintl, A., Mulcahy, G., Skerrett, H. E., Taylor, S. M., and Gray, J. S. (2003). *Babesia divergens*, a bovine blood parasite of veterinary and zoonotic importance. *Clin. Microbiol. Rev.* 16, 622–636. doi: 10.1128/CMR.16.4.622-636.2003
- Zweygarth, E., Koekemoer, O., Josemans, A. I., Rambritch, N., Pienaar, R., Putterill, J., et al. (2009). *Theileria*-infected cell line from an African buffalo (*Syncerus caffer*). *Parasitol. Res.* 105, 579–581. doi: 10.1007/s00436-009-1467-0

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer JL declared a shared affiliation, with no collaboration, with several of the authors, MJ, OH, DS, and PF, to the handling Editor.

Copyright © 2018 Jalovecka, Hajdusek, Sojka, Kopacek and Malandrin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.