

The concentration and speciation of hydrogen in feldspars using FTIR and ^1H MAS NMR spectroscopy

ELIZABETH A. JOHNSON* AND GEORGE R. ROSSMAN

Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California 91125-2500, U.S.A.

ABSTRACT

A universal absorption coefficient was determined for quantitative analysis of OH and H_2O in feldspars using infrared spectroscopy. ^1H MAS (magic-angle spinning) NMR spectroscopy was used to determine the H concentration in three alkali feldspars, and for the first time, eight plagioclase samples. To accurately measure structural H concentration in samples with low H (<1000 ppm H_2O) it was necessary to eliminate the signal due to adsorbed water in the powdered NMR sample. The pegmatitic and metamorphic albite samples are transparent, but contain variable (40–280 ppm H_2O) concentrations of microscopic to sub-microscopic fluid inclusions. The pegmatitic albites also have sharp bands in the mid-IR similar to the OH bands found in quartz. The other plagioclase samples used in the IR calibration have broad anisotropic bands around 3200 cm^{-1} in the mid-IR and weak combination stretch-bend bands near 4550 cm^{-1} in the near-IR, indicative of structural OH. The OH vector in plagioclase is preferentially aligned parallel to the crystallographic **a** axis. The concentration of structural OH in the plagioclase samples ranges from 210–510 ppm H_2O by weight. The microcline samples contain structural H_2O molecules (1000–1400 ppm H_2O) and the sanidine sample contains structural OH (170 ppm H_2O). An approximately linear trend is produced when the total integrated mid-IR absorbance is plotted vs. the concentration of structural H determined from NMR (OH and H_2O) for plagioclase and alkali feldspars. The integral absorption coefficient for the total mid-IR peak area is $15.3 \pm 0.7\text{ ppm}^{-1}\cdot\text{cm}^{-2}$ [$107000 \pm 5000\text{ L}/(\text{mol H}_2\text{O}\cdot\text{cm}^2)$] for natural feldspar samples that contain structural OH or H_2O . Measurements of band areas of unpolarized IR spectra on (001) cleavage fragments provide an estimate of H concentration for alkali feldspars, but this method does not work for most plagioclase samples.