

THE DEGREES OF THE STANDARD IMBEDDINGS OF R -SPACES

YOSHIHIRO OHNITA

(Received November 4, 1982)

1. Introduction. R -spaces constitute an important class of homogeneous submanifolds in the Euclidean spheres: they are the orbits of the isotropy representations of symmetric spaces of noncompact type (cf. Takeuchi and Kobayashi [6]). This class includes many examples appearing in differential geometry of submanifolds. For example, all homogeneous hypersurfaces and all parallel submanifolds in spheres are realized as R -spaces.

Ferus [1] showed that the standard imbeddings of symmetric R -spaces have the parallel second fundamental forms and exhaust all submanifolds in spheres with the parallel second fundamental forms. So the following arises as a natural problem:

Problem. Characterize the standard imbedding of each R -space in the sense of differential geometry.

The first step in answering the Problem is to find many differential geometric properties of the standard imbeddings of R -spaces. In Kitagawa and Ohnita [3] we showed that the standard imbedding of every R -space has the parallel mean curvature vector.

Let M^n be a compact rank one symmetric space, that is, one of the following: S^n , RP^n , CP^n , QP^n and $CayP^6$. Let f_k be the standard minimal isometric immersion of M^n into a sphere $S^{m(k)}$ induced by the k -th eigenfunctions of the Laplace-Beltrami operator of M^n (cf. Wallach [7]). If $k = 1$, the immersion f_k is just the standard imbedding of a compact symmetric R -space of rank one. It is called a generalized Veronese submanifold except when M^n is a sphere. Wallach used the notion of its degree in studying the rigidity of a minimal isometric immersion. The degree of f_k coincides with k (cf. Wallach [7]) if M^n is a sphere, and with $2k$ (cf. Mashimo [4], [5]) otherwise. In particular, the degree of a generalized Veronese submanifold is 2.

In this note we show the following theorems:

Let Φ be the proper standard imbedding (cf. §2) of an R -space K/L .

THEOREM A. *The degree of Φ is equal to 2.*

THEOREM B. *If Φ is regular (cf. §2), then there exists a normal*

frame field $\{\tilde{\xi}_1, \dots, \tilde{\xi}_p\}$ defined globally on K/L such that each $\tilde{\xi}_\alpha$ ($\alpha = 1, \dots, p$) is parallel with respect to the normal connection of Φ . In particular, the normal connection of Φ is flat.

The author thanks Professor H. Urakawa for valuable suggestion, and also thanks the referees for suggesting improvements.

2. Preliminaries. Let (\mathfrak{g}, θ) be an orthogonal symmetric Lie algebra of noncompact type (cf. Helgason [2]). Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition of \mathfrak{g} , where $\mathfrak{k} = \{X \in \mathfrak{g}; \theta(X) = X\}$ and $\mathfrak{p} = \{X \in \mathfrak{g}; \theta(X) = -X\}$. Let α be a maximal abelian subspace of \mathfrak{p} . Let K be the analytic subgroup of the inner automorphism group $\text{Int}(\mathfrak{g})$ corresponding to the Lie algebra $\text{ad}(\mathfrak{k})$. \mathfrak{p} is invariant under K . Let ψ denote the Killing form of \mathfrak{g} and we consider $(\mathfrak{p}, \langle, \rangle)$ as a Euclidean space, where $\langle, \rangle = \psi|_{\mathfrak{p}}$. Put $S = \{X \in \mathfrak{p}; \langle X, X \rangle = 1\}$ and also denote by the same \langle, \rangle the Riemannian metric on S induced by \langle, \rangle . We denote by ∇° and $\bar{\nabla}$ the Riemannian connection of \mathfrak{p} and S , respectively.

For an arbitrarily fixed element H_0 in $S \cap \alpha$, put $L = \{k \in K; k(H_0) = H_0\}$ and an imbedding $\Phi: K/L \rightarrow S$ by $\Phi(kL) = k(H_0)$. Then $M = K/L$ is called an R -space and Φ its standard imbedding. If $\text{rank}(\mathfrak{g}, \theta) = 1$, Φ is the identity map. Φ is said to be proper if $\text{rank}(\mathfrak{g}, \theta) \geq 2$ and $\Phi(M)$ is not a great sphere of S . We can show that if $\Phi(M)$ is a great sphere of S then there are two orthogonal symmetric Lie algebras $(\mathfrak{g}_1, \theta_1)$ and $(\mathfrak{g}_2, \theta_2)$ such that (1) $(\mathfrak{g}, \theta) = (\mathfrak{g}_1, \theta_1) \oplus (\mathfrak{g}_2, \theta_2)$, (2) $\text{rank}(\mathfrak{g}_1, \theta_1) = 1$, (3) $H_0 \in \mathfrak{p}_1 = \{X \in \mathfrak{g}_1; \theta_1(X) = -X\}$.

For an R -linear form λ on α , we put $\mathfrak{g}_\lambda = \{X \in \mathfrak{g}; (\text{ad } H)X = \lambda(H)X \text{ for all } H \in \alpha\}$. If $\mathfrak{g}_\lambda \neq \{0\}$, then λ is called a root of (\mathfrak{g}, θ) with respect to α . Let Δ be the set of all nonzero roots on α . We put $\psi_\theta(X, Y) = \psi(X, \theta(Y))$ for $X, Y \in \mathfrak{g}$. ψ_θ is a negative definite symmetric bilinear form on \mathfrak{g} . Then \mathfrak{g} has the following orthogonal direct sum with respect to ψ_θ : $\mathfrak{g} = \mathfrak{g}_0 + \sum_{\lambda \in \Delta} \mathfrak{g}_\lambda$. For an element $H \in \alpha$, we put $\Delta_H = \{\lambda \in \Delta; \lambda(H) \neq 0\}$. If $\Delta = \Delta_H$, then H is called a regular element in α .

The standard imbedding Φ is called regular if H_0 is a regular element.

We now explain the notion of the degree of an isometric imbedding Φ of a Riemannian manifold $(M, \Phi^*\langle, \rangle)$ into a Euclidean sphere S . We denote by B the second fundamental form of Φ and let A be the shape operator of Φ defined by $\langle A_\xi X, Y \rangle = \langle B(X, Y), \xi \rangle$ for $X, Y \in T_x(M)$ and $\xi \in T_x(M)^\perp$. For $x \in M$, put $O_x^2(M) = \text{span}_R\{B(X, Y); X, Y \in T_x(M)\}$ and let N_x be the orthogonal projection of $T_x(M)^\perp = O_x^2(M) \oplus (O_x^2(M))^\perp$ onto $(O_x^2(M))^\perp$, where $(O_x^2(M))^\perp$ is the orthogonal complement of $O_x^2(M)$ in $T_x(M)^\perp$. Let

$\mathcal{R}_1 = M$ and $\mathcal{R}_2 = \{x \in M; \dim O_x^2(M) \text{ is maximal in } \mathcal{R}_1\}$. We define a symmetric 3-tensor field B_3 on \mathcal{R}_2 by $(B_3)(X, Y, Z) = N_x((\nabla_x^* B)(Y, Z))$ for $X, Y, Z \in T_x(M)$, where $(\nabla_x^* B)(Y, Z) = \nabla_x^{\perp}(B(\tilde{Y}, \tilde{Z})) - B(\nabla_x \tilde{Y}, Z) - B(Y, \nabla_x \tilde{Z})$. Here ∇^{\perp} is the normal connection of Φ and \tilde{Y}, \tilde{Z} are vector fields defined locally around x with $(\tilde{Y})_x = Y, (\tilde{Z})_x = Z$. B_3 is called the third fundamental form of Φ . We can define $O_x^j(M), \mathcal{R}_j, B_j$ for $j = 2, 3, \dots$, recursively. We call B_j the j -th fundamental form of Φ . There exists a natural number d such that $B_d \neq 0$ on \mathcal{R}_d and $B_{d+1} \equiv 0$ on \mathcal{R}_d . We call d the degree of Φ .

For example, $d = 1$ means that Φ is totally geodesic. The degree of Φ is 2 if Φ has the parallel second fundamental form.

3. Proof of Theorems. Let \mathfrak{l} be the Lie algebra of L and \mathfrak{m} be the orthogonal complement of \mathfrak{l} in \mathfrak{k} with respect to ψ_{θ} . We identify the tangent space $T_o(M)$ at the origin $o = \{L\} \in M = K/L$ with \mathfrak{m} . Then the differential $\Phi_*: T_o(M) \rightarrow T_{\Phi(o)}(\mathfrak{p})$ is identified with the mapping $-\text{ad}(H_0): \mathfrak{m} \rightarrow \mathfrak{p}$. Put $\mathfrak{p}_0 = \Phi_*(T_o(M)) = (-\text{ad}(H_0))\mathfrak{m}$ and let \mathfrak{n} be the orthogonal complement of $\text{span}_{\mathbb{R}}\{H_0\} + \mathfrak{p}_0$ in \mathfrak{p} with respect to ψ_{θ} . Then the normal space $T_o(M)^{\perp}$ at the origin of M in S is identified with \mathfrak{n} . We have an orthogonal decomposition of \mathfrak{p} :

$$\mathfrak{p} = \text{span}_{\mathbb{R}}\{H_0\} + \mathfrak{p}_0 + \mathfrak{n}.$$

Now we put

$$\mathfrak{h} = \sum_{\lambda \in \mathcal{A}_{H_0}} \mathfrak{g}_{\lambda}, \quad \mathfrak{b} = \mathfrak{g}_0 + \sum_{\lambda \in \mathcal{A}^{-\mathcal{A}}_{H_0}} \mathfrak{g}_{\lambda}.$$

\mathfrak{h} and \mathfrak{b} are invariant by θ since $\theta(\mathfrak{g}_{\lambda}) = \mathfrak{g}_{-\lambda}$. Hence we have $\mathfrak{g} = (\mathfrak{h} \cap \mathfrak{k}) + (\mathfrak{h} \cap \mathfrak{p}) + (\mathfrak{b} \cap \mathfrak{k}) + (\mathfrak{b} \cap \mathfrak{p})$. Then it is easy to show the following:

$$(3.1) \quad \begin{aligned} \mathfrak{l} &= \mathfrak{b} \cap \mathfrak{k}, \quad \mathfrak{m} = \mathfrak{h} \cap \mathfrak{k}, \quad \mathfrak{p}_0 = \mathfrak{h} \cap \mathfrak{p}, \\ \text{span}_{\mathbb{R}}\{H_0\} + \mathfrak{n} &= \mathfrak{b} \cap \mathfrak{p}. \end{aligned}$$

By $[\mathfrak{k}, \mathfrak{p}] \subset \mathfrak{p}, [\mathfrak{h}, \mathfrak{b}] \subset \mathfrak{h}$ and (3.1), we have

$$(3.2) \quad (\text{ad}(\mathfrak{m})\mathfrak{n}) \subset \mathfrak{p}_0.$$

LEMMA. Let X be an element of \mathfrak{m} and ξ an element of \mathfrak{n} . We put $x_t = (\exp(tX)) \cdot o \in M$. If a normal vector field ξ_t along x_t is defined by $\xi_t = (\exp(tX)) \cdot \xi \in T_{x_t}(M)^{\perp}$ then we have $\nabla_t^{\perp} \xi_t = 0$.

PROOF. $\nabla_t^0 \xi_t = (d/dt)\xi_t = (\exp(tX)) \cdot [X, \xi]$. By (3.2) and $\bar{\nabla}_t \xi_t = \nabla_t^0 \xi_t + \langle \dot{x}_t, \xi_t \rangle x_t = \nabla_t^0 \xi_t$, we have $\bar{\nabla}_t \xi_t \in (\exp(tX)) \cdot \mathfrak{p}_0 = \Phi_*(T_{x_t}(M))$. By Weingarten's fomula we obtain $\nabla_t^{\perp} \xi_t = 0$. q.e.d.

PROOF OF THEOREM A. If Φ is proper, then $B \neq 0$. Since Φ is K -

equivariant, we have $\mathcal{R}_2 = M$. Let X, Y and Z be elements of $\mathfrak{m} = T_o(M)$. Put $x_t = (\exp(tX)) \cdot o, Y_t = (\exp(tX)) \cdot Y \in T_{x_t}(M)$ and $Z_t = (\exp(tX)) \cdot Z \in T_{x_t}(M)$. Then by the K -equivariance of Φ we have $B_{x_t}(Y_t, Z_t) = (\exp(tX)) \cdot B(Y, Z)$. Applying the Lemma to $\xi = B(X, Y) \in \mathfrak{n}$, we have $\nabla_t^\perp(B(Y_t, Z_t)) = 0$. Hence $(\nabla_x^* B)(Y, Z) = -B(\nabla_x Y_t, Z) - B(Y, \nabla_x Z_t) \in O_o^2(M)$. From the definition of the third fundamental form B_3 we have $(B_3)_o = 0$. B_3 vanishes everywhere by the homogeneity of M and the equivariance of Φ . q.e.d.

REMARK. If $(\mathfrak{k}, \mathfrak{l})$ is a symmetric Lie algebra, then Y_t and Z_t are parallel along x_t with respect to the Riemannian connection of M . From the above proof we have $\nabla^* B = 0$.

PROOF OF THEOREM B. It is known that the centralizer in K of a regular element of \mathfrak{a} coincides with the centralizer in K of \mathfrak{a} (for example, see Helgason [2, p. 289]). Thus L is the centralizer in K of \mathfrak{a} . Since $\mathfrak{A} = \mathfrak{A}_{H_0}$, we have $\mathfrak{b} = \mathfrak{g}_o$. By (3.1) and the maximality of \mathfrak{a} , we have $\text{span}_{\mathbb{R}}\{H_o\} + \mathfrak{n} = \mathfrak{b} \cap \mathfrak{p} = \mathfrak{a}$. Hence the action of L on $\mathfrak{n} = T_o(M)^\perp$ is trivial. Select an orthonormal basis $\{\xi_1, \dots, \xi_p\}$ of \mathfrak{n} . We can extend ξ_α to a K -invariant normal vector field $\tilde{\xi}_\alpha$ defined globally on M . By Lemma we have $\nabla^\perp \tilde{\xi}_\alpha = 0$. Thus $\{\tilde{\xi}_1, \dots, \tilde{\xi}_p\}$ is the desired normal frame field. q.e.d.

REFERENCES

- [1] D. FERUS, Symmetric submanifolds of Euclidean spaces, *Math. Ann.* 247 (1980), 81-93.
- [2] S. HELGASON, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York, San Francisco, London, 1978.
- [3] Y. KITAGAWA AND Y. OHNITA, On the mean curvatures of R -spaces, *Math. Ann.* 262 (1983), 239-243.
- [4] K. MASHIMO, Degree of the standard isometric minimal immersions of complex projective spaces into spheres, *Tsukuba J. Math.* 4 (1980), 133-145.
- [5] K. MASHIMO, Degree of the standard isometric minimal immersions of the symmetric spaces of rank one into spheres, *Tsukuba J. Math.* 5 (1981), 291-297.
- [6] M. TAKEUCHI AND S. KOBAYASHI, Minimal imbeddings of R -spaces, *J. Differential Geom.* 2 (1968), 203-215.
- [7] N. WALLACH, Minimal immersions of symmetric spaces into spheres, in *Symmetric Spaces* (W. B. Boothby and G. L. Weiss, eds.), Marcel Dekker, New York, 1972, 1-40.

MATHEMATICAL INSTITUTE
TÔHOKU UNIVERSITY
SENDAI, 980
JAPAN