

 Open access • Journal Article • DOI:10.1021/JA0772795

The Effect of Charge on CO Binding in Rhodium Carbonyls: From Bridging to Terminal CO — [Source link](#)

Ingmar Swart, Frank M. F. de Groot, Bert M. Weckhuysen, David M. Rayner ...+2 more authors

Institutions: Fritz Haber Institute of the Max Planck Society, Max Planck Society

Published on: 29 Jan 2008 - Journal of the American Chemical Society (American Chemical Society)

Topics: Rhodium and HOMO/LUMO

Related papers:

- [Size and charge effects on the binding of CO to late transition metal clusters.](#)
- [The adsorption of CO on transition metal clusters: A case study of cluster surface chemistry](#)
- [Infrared spectroscopy of mass-selected metal carbonyl cations](#)
- [Size and Charge Effects on the Binding of CO to Small Isolated Rhodium Clusters](#)
- [Spectroscopic and theoretical investigations of vibrational frequencies in binary unsaturated transition-metal carbonyl cations, neutrals, and anions.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-effect-of-charge-on-co-binding-in-rhodium-carbonyls-from-fghlemehk>


Communication

The Effect of Charge on CO Binding in Rhodium Carbonyls: From Bridging to Terminal CO

Ingmar Swart, Frank M. F. de Groot, Bert M. Weckhuysen, David M. Rayner, Gerard Meijer, and Andr Fielicke

J. Am. Chem. Soc., **2008**, 130 (7), 2126-2127 • DOI: 10.1021/ja0772795

Downloaded from <http://pubs.acs.org> on January 12, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
High quality. High impact.

The Effect of Charge on CO Binding in Rhodium Carbonyls: From Bridging to Terminal CO

Ingmar Swart, Frank M. F. de Groot,* Bert M. Weckhuysen, David M. Rayner,* Gerard Meijer, and André Fielicke*

Inorganic Chemistry and Catalysis, Faculty of Science, Utrecht University, Sorbonnelaan 16, 3584 CA, Utrecht, The Netherlands, Steacie Institute for Molecular Sciences, National Research Council, 100 Sussex Drive Ottawa, Ontario, K1A 0R6, Canada, and Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195 Berlin, Germany

Received September 20, 2007; E-mail: f.m.f.degroot@uu.nl; david.rayner@nrc-cnrc.gc.ca; fielicke@fhi-berlin.mpg.de

Changing the electron density of a catalyst can have a profound influence on its selectivity and activity. For this reason, electronic promoter materials are often added to a catalyst to tailor the electron density of the active particle and to obtain optimal performance.¹ Despite the widespread use of electronic promoters, the mechanisms responsible for the observed effect(s) are often poorly understood. For CO oxidation over a Pt catalyst, the activity increases with electron density.² In contrast, the hydrogenation of CO over cobalt- and rhodium-based catalysts is found to be more active and more selective upon reducing the electron density of the metal particles.^{3,4} This change in electron density also leads to a decrease in the ratio of bridge to linear-bound CO but the mechanism behind this destabilization of bridge-bound CO is unknown.^{3,5}

We address the issue of charge-induced changes in the binding of CO to metal centers by focusing on rhodium carbonyls as models. Rhodium-based catalysts have been used for many reactions involving CO.⁶ The structures of the neutral saturated carbonyl complexes containing two, four, and six Rh atoms are well-known,^{7–10} but little is known about the charged complexes. We establish the structures of the cations by infrared multiple photon dissociation (IR-MPD) spectroscopy. By comparing the structures of the cationic species with structures of the corresponding neutral complexes, information regarding the effect of charge on the binding geometry of CO is obtained.

The experimental procedures for IR-MPD spectroscopy have been described in detail elsewhere.^{11–13} Briefly, a molecular beam of cluster complexes, generated in a laser ablation source coupled with a reaction channel, is overlapped with a counter-propagating beam of IR photons delivered by the Free Electron Laser for Infrared eXperiments (FELIX).¹⁴ When the laser frequency is resonant with an IR-active vibration of a complex, it can absorb several photons and subsequently undergo fragmentation. The vibrational spectra are obtained by monitoring the changes in the cluster mass distribution as a function of IR frequency. The experimental spectra are complemented with density functional theory (DFT) calculations (B3LYP/TZVP, see Supporting Information) to obtain information on cluster geometries and electronic structure.¹⁵

Upon introduction of CO in the reactor channel, the Rh_n^+ clusters bind multiple CO molecules. The pressure in the reactor channel was increased until the resulting molecular beam distribution did not change anymore, that is, when the clusters were saturated with carbon monoxide.¹⁶ The complexes saturate as $\text{Rh}(\text{CO})_5^+$, $\text{Rh}_2(\text{CO})_8^+$, $\text{Rh}_3(\text{CO})_9^+$, $\text{Rh}_4(\text{CO})_{12}^+$, $\text{Rh}_5(\text{CO})_{14}^+$, and $\text{Rh}_6(\text{CO})_{16}^+$. The saturation numbers of the cationic rhodium dimer, tetramer, and hexamer are the same as those of the corresponding neutral




Figure 1. Experimental (black) and calculated (red) vibrational spectra of $\text{Rh}_n(\text{CO})_m^+$ complexes. The values n, m indicate the number of Rh atoms and CO molecules, respectively. The simulated depletion spectra are based on the calculated single photon absorption spectra. In the complex geometries shown next to the graph, C atoms of μ_1 CO are depicted in gray and C atoms of μ_2 and μ_3 CO are shown in yellow.

clusters. Binary tri- and pentanuclear rhodium cluster carbonyls have not been identified before.

The vibrational spectra in the C–O stretching region and the Rh–C stretch and deformation region for complexes with 1–6 metal atoms are shown in Figure 1. Terminally bound CO ligands are present for all complexes (absorption bands between 2060–2100 cm^{-1}) but bridging (μ_2 -CO) and/or face capping (μ_3 -CO) carbonyls (absorption bands between 1750 and 1900 cm^{-1}) are only present for $\text{Rh}_5(\text{CO})_{14}^+$ and $\text{Rh}_6(\text{CO})_{16}^+$. The cationic carbonyl complexes with 1–4 metal atoms therefore contain only terminally bound CO ligands.¹⁷ The DFT calculations confirm this assignment and also predict that vibrational modes associated with bridge bound CO in isomers of complexes with 2–4 metal atoms have significant IR intensities which should be observed experimentally. The cluster structures and the calculated spectra are shown together with the IR-MPD spectra in Figure 1. The calculated and experimental spectra are in good agreement in both the C–O stretching range, $\nu(\text{CO})$, as well as in the Rh–C stretch and deformation region,




Figure 2. Experimental vibrational spectra of $\text{Rh}_2(\text{CO})_8^{+/0}$, $\text{Rh}_4(\text{CO})_{12}^{+/0}$, and $\text{Rh}_6(\text{CO})_{16}^{+/0}$ (cations, black; neutrals, red; bottom to top) in the $\nu(\text{CO})$ range. Structures for the neutral complexes are shown on the left and for the cations on the right. In the complex geometries shown next to the graph, C atoms of μ_1 CO are depicted in gray and C atoms of μ_2 and μ_3 CO are shown in yellow.

$\nu(\delta(\text{RhC}))$. Only in the case of $\text{Rh}_4(\text{CO})_{12}^+$ we find a discrepancy, a band at $\sim 530 \text{ cm}^{-1}$ is missing in the simulated spectrum. A possible explanation for this discrepancy is that the complex is distorted from the high-symmetry structure (T_d) shown in Figure 1 or the presence of a second isomer. The lowest energy isomers identified by the calculations are also the ones whose vibrational spectra are in best agreement with the experimental data.

In Figure 2 we compare the vibrational spectra of the neutral (taken from ref 10) and cationic species. It is immediately clear that cationic and neutral $\text{Rh}_2(\text{CO})_8$ and $\text{Rh}_4(\text{CO})_{12}$ have different structures. The neutral complexes have two and three bridge-bound carbonyl ligands, respectively,^{7,10} whereas all CO molecules are terminally bound in $\text{Rh}_2(\text{CO})_8^+$ and $\text{Rh}_4(\text{CO})_{12}^+$. Removing an electron from the neutral complexes thus leads to a destabilization of the bridge bound carbonyl ligands. In contrast, $\text{Rh}_6(\text{CO})_{16}$ does not undergo a structural rearrangement. The metal cores of the cations retain the structure of the neutral compounds.

DFT calculations were performed for neutral complexes as well as for cations in the structure of the neutral clusters to compare the electronic structures. The calculations identify the correct ground state isomers of the neutral complexes. For $\text{Rh}_2(\text{CO})_8$, the neutral complex with only terminally bound CO molecules is just 0.09 eV higher in energy. The calculations show that the highest occupied molecular orbital (HOMO) of the complex with the bridging carbonyls is doubly occupied and can be thought of as a combination of rhodium d-orbitals with antibonding $2\pi^*$ orbitals of the bridging CO ligands, the classic back-donation of M–CO binding. The HOMO is bonding with respect to the Rh– μ_2 –CO bonds. Ionization removes an electron from the HOMO. Mulliken overlap population analysis shows that the electron density in a single Rh– μ_2 –CO bond is reduced by 12% upon removing one electron from the neutral cluster. The $\text{Rh}_2(\text{CO})_8^+$ complex is unstable in the structure of the neutral. It is a transition state that is +0.65 eV higher in energy than the isomer with only terminally bound CO. Consequently the complex will undergo a charge induced structural rearrangement. According to our calculations, the accompanying reorganization in electronic structure results in a lengthening of the Rh–Rh bond (see Supporting Information). A similar argument holds for $\text{Rh}_4(\text{CO})_{12}$. The HOMO is involved in the binding of the

three bridge-bonded CO molecules. Upon removing one electron, the electron density between the rhodium atoms and the bridging CO ligands is reduced by 38%. The destabilization of the bridge-bonded CO molecules leads to a complex with only terminally bound CO. As shown by the experimental and calculated vibrational spectra, the high symmetry (T_d) complex $\text{Rh}_6(\text{CO})_{16}$ does not undergo structural rearrangement upon removal of an electron. The HOMO of $\text{Rh}_6(\text{CO})_{16}$ is involved in the binding of the face-capping CO. It has t_2 symmetry and is fully occupied, that is, there are six electrons in three equivalent orbitals. This results in a very stable complex. Removing one electron from the HOMO reduces the electron density between the rhodium atoms and the face-capping carbonyl ligands by only 4%, and hence the complex does not undergo reorganization.

In conclusion, a complete series of saturated cationic rhodium carbonyl complexes $\text{Rh}_n(\text{CO})_m^+$ ($n = 1-6$) has been structurally characterized. It is shown that the removal of an electron from a metal carbonyl cluster can selectively destabilize bridge-bonded CO molecules. These results provide a possible explanation for effects of promoter materials that are observed for CO bonding on transition-metal catalysts,^{2,3} providing information that contributes to understanding the origin of electronic promoter effects in catalysis in general.

Acknowledgment. We gratefully acknowledge the support of the Stichting voor Fundamenteel Onderzoek der Materie (FOM) in providing beam time on FELIX. The authors thank the FELIX staff for their skillful assistance, in particular Dr. B. Redlich and Dr. A.F.G. van der Meer. I.S., F.M.F.d.G., and B.M.W. acknowledge NWO and NRSCC for financial support. Computational time was provided by the SARA supercomputer center in Amsterdam, The Netherlands.

Supporting Information Available: Description of the computational procedure and detailed results. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Ertl, G.; Knözinger, H.; Weitkamp, J., Eds. *Handbook of Heterogeneous Catalysis*; Wiley-VCH: Weinheim, Germany, 1997; p 2801.
- (2) Visser, T.; Nijhuis, T. A.; van der Eerden, A. M. J.; Jenken, K.; Ji, Y.; Bras, W.; Nikitenko, S.; Ikeda, Y.; Lepage, M.; Weckhuysen, B. M. *J. Phys. Chem. B* **2005**, *109*, 3822.
- (3) Morales, F.; de Smit, E.; de Groot, F. M. F.; Visser, T.; Weckhuysen, B. M. *J. Catal.* **2007**, *246*, 91.
- (4) Koerts, T.; van Santen, R. A. *J. Catal.* **1992**, *134*, 13.
- (5) Beutel, T.; Alekseev, O. S.; Ryndin, Y. A.; Likholobov, V. A.; Knozinger, H. *J. Catal.* **1997**, *169*, 132.
- (6) Dyson, P. J.; McIndoe, J. S. *Transition Metal Carbonyl Cluster Chemistry*; Gordon and Breach Science Publishers: Amsterdam, The Netherlands, 2000.
- (7) Wei, C. H. *Inorg. Chem.* **1969**, *8*, 2384.
- (8) Corey, E. R.; Dahl, L. F.; Beck, W. J. *Am. Chem. Soc.* **1963**, *85*, 1202.
- (9) Heaton, B. T.; Jacob, C.; Podkorytov, I. S.; Tunik, S. P. *Inorg. Chim. Acta* **2006**, *359*, 3557.
- (10) Allian, A. D.; Wang, Y. Z.; Saeyns, M.; Kuramshina, G. M.; Garland, M. *Vib. Spectrosc.* **2006**, *41*, 101.
- (11) Fielicke, A.; von Helden, G.; Meijer, G.; Pedersen, D. B.; Simard, B.; Rayner, D. M. *J. Phys. Chem. B* **2004**, *108*, 14591.
- (12) Swart, I.; Fielicke, A.; Redlich, B.; Meijer, G.; Weckhuysen, B. M.; de Groot, F. M. F. *J. Am. Chem. Soc.* **2007**, *129*, 2516.
- (13) Oomens, J.; Tielens, A.; Sartakov, B. G.; von Helden, G.; Meijer, G. *Astrophys. J.* **2003**, *591*, 968.
- (14) Oepke, D.; van der Meer, A. F. G.; van Amersfoort, P. W. *Infrared Phys. Technol.* **1995**, *36*, 297.
- (15) Ahlrichs, R.; Bär, M.; Häser, M.; Horn, H.; Kölmel, C. *Chem. Phys. Lett.* **1989**, *162*, 165.
- (16) Fayet, P.; McGlinchey, M. J.; Wöste, L. H. *J. Am. Chem. Soc.* **1987**, *109*, 1733.
- (17) The spectra exhibit saturation and broadening of the μ_1 band(s) when repeated at 20 times the IR fluence but no new absorptions. This confirms that noise does not mask low strength transitions. Saturation of the depletion via the μ_1 band(s) confirms that other isomers are not present in significant quantities.

JA0772795