

Published in final edited form as:

Eur J Neurosci. 2015 November ; 42(9): 2707–2715. doi:10.1111/ejn.13071.

The effect of heterotopic noxious conditioning stimulation on A δ -, C-, and A β -fiber brain responses in humans

Diana M. Torta¹, Maxim V. Churyukanov^{1,2}, Leon Plaghki¹, and André Mouraux¹

¹Institute of Neurosciences (IONS), Université catholique de Louvain, Belgium

²A. Ya. Kozhevnikov Clinic of Nervous Diseases, The I.M. Sechenov First Moscow State Medical University, Russia

Abstract

Human studies have shown that heterotopic nociceptive conditioning stimulation (HNCS) applied to a given body location reduce the percept and brain responses elicited by noxious test stimuli delivered at a remote body location. It remains unclear to what extent this effect of HNCS relies on the spinal-bulbar-spinal loop mediating the effect of diffuse noxious inhibitory controls (DNIC) described in animals, and/or on top-down cortical mechanisms modulating nociception. Importantly, some studies have examined the effects of HNCS on the brain responses to nociceptive input conveyed by A δ -fibers. In contrast, no studies have explored the effects of HNCS on the responses to selective nociceptive C-fiber input and non-nociceptive A β fiber input. In this study, we measured the intensity of perception and event related potentials (ERPs) to stimuli activating A δ -, C- and A β -fibers, before, during and after HNCS, obtained by immersing one foot in painful cold water. We observed that (i) the perceived intensity of nociceptive A δ - and C- stimuli was reduced during HNCS, (ii) the ERPs elicited by A δ - and A β -, and C- stimuli were also reduced during HNCS. Importantly, because A β -ERPs are related to primary afferents that ascend directly through the dorsal columns without being relayed at spinal level, the modulation of these responses may not be explained by an influence of descending projections modulating the transmission of nociceptive input at spinal level. Therefore, our results indicate that, in humans, HNCS should be used with caution as a direct measure of DNIC-related mechanisms.

Keywords

Nociception; pain; cold pressor test; CO₂ laser; laser-evoked potentials; event related potentials (ERPs); heterotopic noxious conditioning stimuli; HNCS; DNIC

Introduction

Pain can be modulated by inhibitory and facilitatory mechanisms acting both at spinal and supra-spinal levels (Millan, 2002; Porreca *et al.*, 2002). These modulatory mechanisms are increasingly considered to play an important role in the individual susceptibility to pain.

Corresponding author: Prof. André Mouraux, Laboratoire d'algologie – Institute of neuroscience IoNS, Université catholique de Louvain, 53, Avenue Mounier – UCL 53.75, B-1200 Bruxelles, Belgium, Phone: +32-2-764-5449, andre.mouraux@uclouvain.be.

The authors have no conflict of interest.

In animals, modulatory mechanism can involve a spinal-bulbar-spinal loop in which noxious stimuli applied to a given body part inhibit, at the level of the dorsal horn, transmission of nociceptive input originating from all other body parts (see (Le Bars, 2002) for a review). The neural substrates of this mechanism, termed “diffuse noxious inhibitory control” (DNIC), remain yet to be fully understood. Lesions of the periaqueductal grey (PAG) do not modify DNIC. In contrast, the mechanism is disrupted by lesions of the *subnucleus reticularis dorsalis* in the caudal medulla (Bouhassira *et al.*, 1992; Villanueva & Le Bars, 1995). Recent data also suggest that, contrary to what has been previously proposed, lesions of the rostral ventromedial medulla (RVM), including the *nucleus raphe magnus*, can affect DNIC (Chebbi *et al.*, 2014). Furthermore, the parabrachial nucleus (PB) would also be involved in triggering DNIC (Lapirot *et al.*, 2009). Finally, a role of descending dopaminergic controls has been recently identified (Lapirot *et al.*, 2011).

Human studies have shown that nociceptive conditioning stimuli applied to a given body location reduce the percept and brain responses elicited by test stimuli delivered at a remote location (De Broucker *et al.*, 1990; Willer *et al.*, 1990; Bouhassira *et al.*, 1993). Furthermore, human studies have shown that heterotopic noxious conditioning stimulation (HNCS) inhibits the spinal RIII reflex, suggesting that at least part of the effect of HNCS on pain perception may be due to descending inhibitory influences on spinal transmission (Willer *et al.*, 1984; Roby-Brami *et al.*, 1987; Willer *et al.*, 1989; De Broucker *et al.*, 1990; Danziger *et al.*, 1998; Scheuren *et al.*, 2014). For this reason, the effects of HNCS have often been considered as a direct human correlate for DNIC (see (Pud *et al.*, 2009)).

Importantly, descending modulatory effects are thought to operate predominantly at the level of Wide Dynamic Range (WDR) neurons, onto which nociceptive C-fibers, A δ -fibers and A β -fibers converge. Using laser-evoked potentials, studies have demonstrated that HNCS can reduce the magnitude of the brain responses to nociceptive input conveyed by A δ -fibers (Plaghki *et al.*, 1994; Watanabe *et al.*, 1996).

So far, no studies have examined whether HNCS affects the brain responses to inputs conveyed selectively by nociceptive C-fibers. One of the reasons for this lack of knowledge is that selective activation of C-fibers is difficult to achieve. However we (Jankovski *et al.*, 2013; van den Broeke & Mouraux, 2014a) and other groups (Magerl *et al.*, 1999) have shown that laser-evoked brain responses and percepts (Churyukanov *et al.*, 2012) related to the selective activation of C-fibers can be obtained using a temperature-controlled infrared CO₂ laser stimulator to selectively activate heat-sensitive C-fiber afferents that have a lower thermal activation threshold than A δ -fiber afferents (Jankovski *et al.*, 2013; van den Broeke & Mouraux, 2014a). Therefore, the first objective of our study was to assess the effects of HNCS on the perception and brain responses to selective C-fiber input.

Furthermore, no study has yet examined whether HNCS also affects the perception and event-related potentials (ERPs) to sensory input conveyed by non-nociceptive A β -fibers (Somatosensory evoked potentials, SEPs). Considering that these brain responses are related to A β -fiber primary afferents which are not relayed in the spinal cord but ascend directly through the dorsal columns to reach second-order neurons in the dorsal column nuclei, one would expect that these responses would not be under the influence of descending spinal

modulatory mechanisms. Therefore, the second objective of our study was to assess the effects of HNCS on the perception and brain responses to A β -fiber input, in order to examine whether the effects of HNCS can be attributed to changes occurring exclusively at spinal level and/or to changes also occurring at supra-spinal level.

Methods

Experiment 1

Participants—Experiment 1 was performed on eight healthy volunteers recruited among students and staff of the Université catholique de Louvain (3 female and 5 male; all right-handed, aged 23 to 35 years). The phase of the menstrual cycle in the main experiment was not taken into consideration in the participating women, although previous studies have suggested that during the ovulatory phase, HNCS may trigger a greater modulation of nociceptive responses (Tousignant-Laflamme & Marchand, 2009). Participants had no history of neurological, psychiatric, dermatological or chronic pain disorders, and no recent history of psychotropic or analgesic drug use. During a preliminary session, volunteers were given a fully detailed explanation concerning the experimental procedures and were familiarized with the experimental setup and task, the nociceptive stimuli and the rating procedures. Participants were comfortably seated in a chair with forearms resting on the armrests. Written informed consent was obtained from all participants. The study was approved by the Ethics Committee of the Université catholique de Louvain (B40320096449).

Experimental design—Each participant underwent three blocks: before, during and after the HNCS conditioning procedure. Within each block, behavioural (reaction times, intensity of perception) and electrophysiological (event-related brain potentials) responses to 20 infrared CO₂ laser stimuli activating nociceptive C- and A δ -fibers free nerve endings (A δ -stimulus, C-stimulus) and 20 transcutaneous electrical stimuli activating non-nociceptive A β -fibers (A β -stimulus) were recorded. The laser stimuli were applied to the right hand dorsum. The electrical stimuli were applied over the superficial branch of the right radial nerve. Within each block, the two types of stimuli were presented in separate runs, whose order was randomized across participants.

Thermal CO₂ laser stimulation (A δ -stimulus and C-stimulus)—Thermal stimuli were applied to the right hand dorsum using a CO₂ laser in which the power of the laser beam is regulated continuously during stimulation using an online measurement of skin temperature optically in line with the laser beam. This feedback control of power output allowed defining specific skin temperature heating profiles (LSD, SIFEC, Belgium). Because the CO₂ laser has a wavelength of 10.6 μ m, 99% of the delivered energy was confined within the most superficial layer of the skin (<100 μ m beneath the skin surface), i.e. where most of the free nerve endings of nociceptors terminate (Meyer *et al.*, 1976). The laser beam was conducted through a 10 m optical fiber. An optical fiber vibrator was used to introduce a small amount of variable lateral or angular displacement in the fiber, altering internal reflections and, thereby, resulting in a homogeneous spatial distribution of power

within the stimulated area. At the end of the fiber, optics collimated the beam in a 6 mm beam diameter at target site.

An approach similar to Magerl et al. (Magerl *et al.*, 1999) was used to temporally dissociate the activation of nociceptive C-fibers and A δ -fibers and, thereby, allow the recording of behavioral and electrophysiological responses related to the activation of each fiber type. The procedure is represented in Figure 1B. First, skin temperature was brought to 34°C using a 1-second heating ramp and maintained at that temperature for 3 seconds (baseline skin temperature). Second, the skin temperature was increased to 43°C in 10 ms. In a previous study, we showed that such temperatures are above the thermal activation threshold of C-fibers but below the thermal activation threshold of A δ -fibers (Churyukanov *et al.*, 2012). Hence, this first step increase allowed activating C-fibers selectively (C-stimulus). Finally, after maintaining the skin temperature at 43°C for 3 seconds, the skin temperature was increased to 49°C in 10 ms, and maintained at that temperature for 40 ms before interrupting laser power output. This second target temperature activated A δ -fibers. After each trial, the target of the laser was displaced to a random position on the hand dorsum, in order to avoid nociceptor sensitization and/or habituation. The inter-trial interval varied randomly between 5-10 seconds.

Transcutaneous electrical stimulation (A β -stimulus)—Transcutaneous electrical stimuli consisted of constant current square-wave electrical pulses (0.5 ms duration; DS7A, Digitimer Ltd, UK) delivered through a pair of feltpad skin electrodes (0.5 cm diameter, 2 cm inter-electrode distance) placed at the right wrist, over the superficial branch of the radial nerve. Prior to the experimental session, the intensity of the electrical stimulus was set to twice the detection threshold (2.30 ± 0.80 mA). The inter-trial interval varied randomly between 5-10 seconds.

Heterotopic noxious conditioning stimulation (HNCS)—During the first block (before HNCS), participants immersed their left foot into a water bath containing water at $30.9 \pm 0.8^\circ\text{C}$, that was perceived as neither warm nor cold. During the second block (during HNCS), participants immersed their foot in another water bath of circulating water whose temperature was set to 5°C (measured at $5.1 \pm 0.3^\circ\text{C}$ at the beginning of the block and $5.7 \pm 0.3^\circ\text{C}$ at the end of the block). This procedure elicited a perception of pain after about 10-15 sec, which peaked after about 30-45 sec and then stabilized. The intensity of the elicited pain was just below the limit of tolerance for most subjects. During the third block (after HNCS), participants re-immersed the foot in the water bath of the first block (temperature $30.0 \pm 0.8^\circ\text{C}$). In each block, recordings began 30-seconds after immersing the foot in the water bath. A 5-minutes rest period separated each block. The foot was dried and covered with a towel during the rest period.

Behavioral measures

Reaction times: During nociceptive stimulation, participants were asked to respond as quickly as possible by pressing a button held in the left hand when perceiving the first (43°C) and second (49°C) step-increase in skin temperature, corresponding to the C-stimulus and the A δ -stimulus, respectively. Reaction times were measured relative to the

onset of the step-increase. Based on the large difference between the conduction velocities of A δ - and C- fibers, responses <650 ms were considered related to the detection of A δ -fiber input, whereas responses \geq 650 ms were considered related to the detection of C-fiber input (Churyukanov *et al.*, 2012). Only trials with a reaction time between 650-2000 ms were included for the analysis of C-fiber ERPs. This represented 85 ± 3 % of the total number of trials. Only trials with a reaction time <650 ms were included for the analysis of A δ -fiber ERPs, representing 91 ± 5 % of the total number of trials. During non-nociceptive stimulation, participants were also asked to respond as quickly as possible by pressing the same button used to detect nociceptive stimuli.

Intensity of perception: After each stimulus, participants were asked to rate the intensity of the perception elicited by C-, A δ - and A β -fiber stimuli using a numerical rating scale (NRS) presented at eye level, approximately 1.5 m in front of the participant. The extremities of the scale were annotated with the words “no detection” (NRS = 0) and “maximum pain” (NRS = 100), respectively. An anchor at the middle of the scale (NRS = 50) defined the borderline between the non-painful and painful domains of perception. Participants provided a verbal report of the perceived intensity of perception. Only ratings to perceived stimuli (>0) were included in the analysis.

Electrophysiological measures—The EEG was recorded using 64 Ag-AgCl electrodes placed on the scalp according to the international 10-20 system (Waveguard64 cap, Cephalon A/S, Denmark). Scalp signals were recorded using an average reference. Impedance was kept below 10 k Ω . Ocular movements and eye blinks were recorded using two surface electrodes placed at the upper-left and lower-right sides of the right eye. All signals were amplified and digitized at a 1 kHz sampling rate (64-channel ASA-LAB EEG/ERP system, Advanced Neuro Technologies, The Netherlands).

Continuous EEG signals were band-pass filtered (0.5-30 Hz) using a Butterworth zero-phase filter and segmented into 3 s epochs ranging from -1 to +2 s relative to the onset of the stimulus (C-stimulus, A δ -stimulus and A β -stimulus). Artifacts produced by eye blinks or eye movements were subtracted using a validated method based on an Independent Component Analysis (Jung *et al.*, 2000). Baseline correction was performed by subtracting the average voltage of the prestimulus interval ranging from -1 to 0 s. In addition, epochs with amplitude values exceeding ± 100 μ V (i.e., epochs likely to be contaminated by an artifact) were rejected.

For each type of stimulus (C-stimulus, A δ -stimulus, A β -stimulus), separate average waveforms were computed for each block (before, during and after), resulting in three average waveforms for the C-stimulus (C-LEP), three average waveforms for the A δ -stimulus (A δ -LEP) and three average waveforms for the A β -stimulus (A β -SEP). In the C-LEP waveforms, no significant signal deflection could be identified, and the signals were not further analyzed using this approach (see following paragraphs). In the A δ -LEP waveforms, three distinct ERP components were identified as follows. The A δ -N2 and A δ -P2 waves were identified at electrode Cz (average reference) as the most negative and positive deflections occurring between 100-600 ms after stimulus onset or step-increase in skin temperature. The A δ -N1 wave was identified at electrode C3 re-referenced to electrode Fz,

as the first negative deflection preceding the A δ -N2 wave, which appears as a positive wave in this montage. In the A β -ERP waveforms, three distinct ERP components were identified as follows. The A β -N2 and A β -P2 waves were identified at electrode Cz (average reference) as the most negative and positive deflections occurring between 100-350 ms after stimulus onset. The A β -N1 was identified at electrode C3 re-referenced to electrode Fz and defined as the first negative deflection preceding the A β -N2.

Experiment 2

Considering that we could not obtain reliable C-fiber responses with the experimental paradigm used in experiment 1, we conducted a second experiment in which we tested selectively the effects of HNCS on C-LEPs. This experiment was performed on 8 healthy volunteers recruited among students and staff of the Université catholique de Louvain (2 female and 6 male; one left handed, aged 22 to 37 years). We recruited mainly men to avoid potentially confounding effects of the phase of the menstrual cycle. Participants had no history of neurological, psychiatric, dermatological or chronic pain disorders, and no recent history of psychotropic or analgesic drug use. During a preliminary session, volunteers were given a fully detailed explanation concerning the experimental procedures. Participants were comfortably seated in a chair with forearms resting on the armrests. Written informed consent was obtained from all participants. The study was approved by the Ethics Committee of the Université catholique de Louvain.

Experimental design—As in experiment 1, each participant underwent three blocks: before HNCS, during HNCS, and after HNCS. Water temperature was $30.5 \pm 0.9^\circ\text{C}$ before HNCS, $5.9 \pm 0.5^\circ\text{C}$ during HNCS and $30.3 \pm 0.8^\circ\text{C}$ after HNCS.

Thermal CO₂ laser stimulation (C-stimulus)—Thermal stimuli were applied using the same device as in experiment 1. The laser beam was 6 mm. The stimuli consisted in 100 ms of radiant heat, of which 10 ms corresponded to the rapid heating ramp during which the skin was brought to the target temperature, and 90 ms corresponded to the plateau of the target temperature. In order to increase the number of trials in which the thermal stimulus activated C-fibers selectively, we used the approach recently proposed by van den Broeke and Mouraux (van den Broeke & Mouraux, 2014a). At the beginning of the experiment, the thermal detection threshold of A δ - and C-fibers was determined using two staircase algorithms. Reaction times >650 ms were considered compatible with the detection of input conveyed by C-fibers, and reaction times ≤ 50 ms were considered compatible with the detection of input conveyed by faster A δ -fibers (Churyukanov *et al.*, 2012). Participants were asked to respond as soon as they perceived a stimulus. The staircase procedure used to estimate the C-fiber thermal detection threshold started at 41°C . The temperature was then increased by 1°C when the stimulus was not detected, and decreased by 1°C when the stimulus was detected. The staircase procedure used to estimate the A δ -fiber thermal detection threshold started at 46°C . The temperature was increased by 1°C when the stimulus was detected with a reaction time ≤ 50 ms, and decreased when the stimulus was detected with a reaction time ≥ 50 ms. The staircases were interrupted after the occurrence of 3 reversals. The target temperature of the test stimulus was defined as the average of the C- and A δ -fibers thresholds. This ensured that the test stimulus was above the threshold of

C-fibers, but below the threshold of A δ -fibers (i.e., that the test stimulus activated C-fibers selectively). In each block, the test stimuli were repeated until we obtained at least 20 trials in which the stimulus was detected with a reaction time >650 ms, with a maximum of 40 trials. After each trial, the target of the laser was displaced to a random position on the hand dorsum, in order to avoid nociceptor sensitization and/or habituation. The inter-trial interval varied randomly between 5-10 seconds.

Intensity of perception: After each stimulus, participants were asked to rate the intensity of the perception elicited by the stimulus using a Numerical Rating Scale (NRS). The extremities were described as “no detection” (NRS = 0) and “maximum pain” (NRS = 100), respectively. An anchor at the middle of the scale (NRS = 50) defined the transition between the non-painful and painful domains of perception. Such as in experiment 1, only ratings to perceived stimuli (>0) were included in the analysis.

Electrophysiological measures—Previous studies showed that N2 and P2 peaks of the C-LEP are maximal over the midline centro-parietal electrodes. Therefore, we recorded the EEG using 5 channels (Cz, Cpz, Pz, M1, M2) of a 64 Ag-AgCl cap (Waveguard64 cap, Cephalon A/S, Denmark). Impedance was kept below 10 k Ω . Ocular movements and eye blinks were recorded using two surface electrodes placed at the upper-left and lower-right sides of the right eye. All signals were amplified and digitized at a 4 kHz sampling rate (64-channel ASA-LAB EEG/ERP system, Advanced Neuro Technologies, The Netherlands).

Continuous EEG signals were band-pass filtered (0.5-30 Hz) using a Butterworth zero-phase filter and segmented into 3 s epochs ranging from -1 to +2 s relative to the onset of the stimulus. The Gratton-Coles method was used to correct eye-movements and blinks (Gratton *et al.*, 1983). Contrary to ICA, this method can be applied also with a limited number of channels. Baseline correction was performed by subtracting the average voltage of the prestimulus interval ranging from -1 to 0 s. In addition, epochs with amplitude values exceeding ± 70 μ V (i.e., epochs likely to be contaminated by an artifact) were rejected. The signal was rereferenced to the mastoids. The maximal peak was observed at Cz in 6 subjects and in Cpz in 2 subjects.

Statistical analyses: To assess possible differential effects of HNCS on the intensity of perception and ERPs elicited by each of the three types of stimuli, we used non-parametric repeated measures Friedman’s tests. Post-hoc analyses with Wilcoxon signed-rank tests were conducted to follow up significant effects. All statistical analyses were performed in SPSS 19 (IBM Armonk, NY).

Results

Experiment 1

Intensity of perception—The Friedman test revealed a significant effect of ‘block’ (before, during, and after HNCS) on the perception of A δ and C stimuli (A δ -NRS: $\chi^2=14.2$, $p=0.001$; C-NRS $\chi^2=12.2$, $p=0.002$) but not A β stimuli (A δ -NRS: $\chi^2=4.2$, $p=0.122$). Post-hoc analyses with Wilcoxon tests showed that A δ -NRS ratings were reduced during HNCS as compared to before HNCS ($z=-2.5$, $p=0.012$), but also as compared to after HNCS

($z=-2.5$, $p=0.012$) (Table 1). A δ -NRS ratings after HNCS were also significantly reduced as compared to before HNCS ($z=-2.3$, $p=0.017$). C-NRS ratings were reduced during HNCS as compared to before HNCS ($z=-2.5$, $p=0.012$), and returned to baseline level after HNCS (during vs. after HNCS: $z=-2.5$, $p=0.012$; after vs. before HNCS: $z=-1.2$, $p=0.208$) (Table 1).

ERPs elicited by C-, A δ - and A β -stimuli—Group-level average waveforms are shown in Figure 3. As explained in Methods, C-stimuli did not elicit consistent ERPs, and were not further analyzed.

N2 waves elicited by A δ - and A β -stimuli: There was a significant effect of ‘block’ on the vertex negative peak of the A δ -ERP (A δ -N2: $\chi^2=9.7$, $p=0.008$), which was reduced during HNCS as compared to before HNCS ($z=-2.3$, $p=0.017$), and remained reduced after HNCS (after vs. during HNCS: $z=-0.280$, $p=0.779$, after vs. before HNCS: $z=-2.5$, $p=0.012$) (Table 1). The magnitude of the A β -N2 was also influenced by ‘block’ ($\chi^2=7.7$, $p=0.021$). However, this effect was mainly driven by an enhancement of the negative peak after HNCS, rather than by a reduction during HNCS (before vs. during HNCS: $z=-1.1$, $p=0.263$; before vs. after HNCS: $z=-1.2$, $p=0.208$; during vs. after HNCS: $z=-2.5$, $p=0.012$).

P2 waves elicited by A δ - and A β -stimuli: No significant effects of ‘block’ were observed for the vertex positive peak of the A δ -ERP ($\chi^2=3.2$, $p=0.073$). In contrast, the magnitude of the A β -P2 was significantly modulated by ‘block’ ($\chi^2=7$, $p=0.030$). Post-hoc comparisons showed that the A β -P2 was significantly reduced during HNCS as compared to before HNCS ($z=-2.3$, $p=0.017$). Importantly, the amplitude of the A β -P2 returned to baseline levels after HNCS (during vs. after HNCS: $z=-2.1$, $p=0.036$; after vs. before HNCS: $z=-0.7$, $p=0.484$) (Table 1).

N1 waves elicited by A δ - and A β -stimuli: The amplitude of the A δ -N1 did not change as a function of ‘block’ ($\chi^2=4$, $p=0.135$). In contrast, the A β -N1 was significantly modulated by ‘block’ ($\chi^2=6.7$, $p=0.034$). However, and in line with the effect of ‘block’ on the magnitude of the negative vertex component, this effect was not driven by a significant reduction during HNCS, but by a significant enhancement after HNCS (before vs. during HNCS: $z=-1.6$, $p=0.093$; during vs. after HNCS: $z=-0.840$, $p=0.401$; before vs. after HNCS: $z=-2.5$, $p=0.012$).

Experiment 2

Intensity of perception—Such as in experiment 1, the Friedman test revealed a significant effect of ‘block’ (before, during, and after HNCS) on the perception of C- stimuli (C-NRS $\chi^2=7.750$, $p=0.021$). Post-hoc analyses with Wilcoxon tests showed that C-NRS ratings were reduced during HNCS as compared to before HNCS ($z=-2.1$, $p=0.036$), but remained on average lower after HNCS (during vs. after HNCS: $z=-1.2$, $p=0.208$; before vs. after HNCS: $z=-1.9$, $p=0.050$) (Table 1 and Figure 2).

ERPs elicited by C-stimuli—The group-level average waveforms obtained in experiment 2 are shown in Figure 5.

N2 waves elicited by C-stimuli: There was a significant effect of ‘block’ on the vertex negative peak of the C-LEP (C-N2: $\chi^2=6.750$, $p=0.034$), which was reduced during HNCS as compared to before HNCS ($z=-2.5$, $p=0.012$), and remained reduced after HNCS (after vs. during HNCS: $z=-0.28$, $p=0.779$, after vs. before HNCS: $z=-2.1$ $p=0.036$).

P2 waves elicited by C-stimuli: No significant effect of ‘block’ was observed for the vertex positive peak of the C-LEP ($\chi^2=1$, $p=0.607$).

Discussion

In this study, we investigated the effects of HNCS on the perception and ERPs elicited by the stimulation of nociceptive A δ - and C-fiber afferents, and non-nociceptive A β -fiber afferents.

We observed that HNCS reduced the perceived intensity of both A δ - and C-fiber nociceptive inputs. In addition, HNCS reduced the magnitude of the A δ -N2, the C-N2, and the A β -P2. Importantly, this latter effect could not be attributed only to habituation as the amplitude of the A β -P2 decreased during HNCS, but returned to baseline levels after HNCS.

Notably, the ERPs elicited by A β -stimuli are primarily related to inputs which are not relayed in the spinal cord but ascend directly through the dorsal columns to reach second-order neurons in the dorsal column nuclei. Therefore, by showing that HNCS modulates the brain responses to non-nociceptive A β -fiber inputs, our results indicate that the effects of HNCS should be cautiously solely attributed to descending projections modulating transmission at spinal level.

HNCS modulates perception of C- and A δ -stimuli

In human studies, it has been shown that HNCS is able to modulate consistently the perceived intensity of noxious test stimuli, irrespective of the nociceptive conditioning procedure and the nature of the nociceptive test stimulus (for reviews see (Pud *et al.*, 2009; van Wijk & Veldhuijzen, 2010)). This effect has been attributed to descending modulatory projections which are thought to operate predominantly at the level of wide dynamic range (WDR) neurons of the dorsal horn, onto which nociceptive C-fibers and A δ -fibers, as well as non-nociceptive A β -fibers converge (Le Bars, 2002).

We observed that HNCS reduced the pain evoked by both nociceptive C-fiber input and nociceptive A δ -fiber input. Such as in previous studies (Treister *et al.*, 2010), the reduction in perception was stronger during HNCS as compared to after HNCS, indicating that the reduction observed during HNCS was not solely due to perceptual habituation. Indeed, had it been the case, one would have expected the intensity of the percepts elicited after HNCS to be further reduced. Importantly, the finding that the intensity of the percept elicited by A δ -fiber and C-fiber input remained lower 5-10 minutes after HNCS could be due to perceptual habituation, but also to a long-lasting after-effect of HNCS. Indeed, Willer *et al.* (Willer *et al.*, 1990) showed that the RIII reflex can be suppressed up to 6-9 minutes after HNCS.

Contrasting with the effects of HNCS on the perception of nociceptive A δ - and C-fiber inputs, we observed no effect of HNCS on the perception of non-nociceptive A β -fiber input. This lack of modulation should be interpreted with caution, as it could be explained by different factors. Indeed, several studies including the present one have observed that experimental manipulations can modulate the brain responses elicited by transcutaneous stimulation of non-nociceptive A β -fibers without modulating subjective reports of intensity of perception (Torta *et al.*, 2013; Van Den Broeke & Mouraux, 2014b; Torta *et al.*, 2015). This could be explained by the fact that transcutaneous electrical stimulation elicits a robust percept, identical from trial to trial. Hence, the modulatory effect of HNCS on the processing of A β -fiber input could be insufficient to result in a noticeable change of perception.

HNCS modulates ERPs elicited by A δ , A β and C stimuli

In line with previous studies showing a reduction of the vertex negativity (A δ -N2) after CO₂ laser stimulation (Plaghki *et al.*, 1994) and of the N2-P2 vertex component elicited by contact-heat stimulation (Moont *et al.*, 2011), we found that HNCS reduces the magnitude of the ERPs elicited by nociceptive A δ -fiber input (experiment 1), as well as the ERPs elicited by C-fiber input (experiment 2). However, because this reduction remained present after HNCS, the observed reduction could be explained by a sustained effect of HNCS, or by an unspecific effect of habituation.

Most notably, we observed that HNCS also modulates the ERPs elicited by non-nociceptive A β fiber input. Indeed, the A β -P2 was significantly more reduced during HNCS as compared to after HNCS, indicating that the reduction of A β -P2 amplitude observed during HNCS was not simply due to response habituation (Figure 3).

The finding that HNCS exerts a similar effect on the brain responses to nociceptive inputs conveyed by A δ - and C-fibers and non-nociceptive input conveyed by A β -fibers is important. Indeed, the ERPs elicited non-nociceptive A β -fiber input reflect activity, which is not mainly relayed in the dorsal horn of the spinal cord, but ascends predominantly through the dorsal columns to reach second-order neurons in the dorsal column nuclei. Therefore, the effect of HNCS on the brain responses to A β -fiber input may not be explained by mechanisms involving descending spinal modulatory projections, in particular, the spinal-bulbar-spinal loop underlying DNIC. It has to be considered that WDR neurons of the dorsal horn also receive input from non-nociceptive A β -fiber afferents, and that the ascending A β information through the dorsal column also has an indirect component (postsynaptic dorsal column) with relay in the spinal cord. However, the contribution of these A β -fiber projections onto the spinothalamic tract to the middle-latency N1, N2 and P2 waves of A β -ERPs is probably negligible. This is demonstrated by the results of studies showing that lesions of the spinothalamic tract lead to the dissociated observation of (1) altered or absent A δ -ERPs and (2) preserved A β -ERPs (Kakigi *et al.*, 1991; Treede *et al.*, 1991; Iannetti *et al.*, 2001; Iannetti *et al.*, 2013).

What then are the mechanisms that could explain the effects of HNCS on the evoked brain responses to both nociceptive and non-nociceptive inputs? Using functional magnetic resonance imaging (fMRI), Sprenger *et al.* (Sprenger *et al.*, 2011) found that HNCS leads to

a tonic increase of BOLD signal in the anterior cingulate cortex, as well as to increased functional coupling between the cingulate cortex and brain stem regions hypothesized to be involved in the control of descending spinal pain-modulatory pathways. Interestingly, using fMRI of the spinal cord, Sprenger et al. (Sprenger *et al.*, 2012) also showed that mental distraction can reduce the BOLD response to nociceptive stimuli at the level of the dorsal horn. Therefore, the effects of HNCS on the brain responses to nociceptive and non-nociceptive inputs might be also explained by top-down cognitive-attentional mechanisms distinct from DNIC. Regarding nociceptive input, there is convincing evidence that at least part of these effects is related to a top-down modulation occurring already at the level of the first relay in the dorsal horn. Whether similar corticofugal mechanisms could modulate the transmission of non-nociceptive somatosensory input at the level of dorsal column nuclei should be envisaged. Supporting this hypothesis, it has been shown that corticofugal projections can modulate the responses of dorsal column nuclei to tactile stimuli (Nunez & Malmierca, 2007; Malmierca *et al.*, 2014). This effect has been suggested to contribute to the mechanisms of selective attention. Obviously, the fact that top-down mechanisms triggered by HNCS may modulate the transmission of nociceptive and non-nociceptive somatosensory inputs at the level of their first synaptic relay in the dorsal horn or brainstem does not preclude the fact that somatosensory processing may be further modulated at higher-order relays such as the thalamus and cortex (Dockstader *et al.*, 2010).

Brain responses related to the selective activation of C fibers

In the first experiment we were unable to identify consistent ERPs related to the activation of C-fibers, in contrast with the results obtained by Magerl et al. (Magerl *et al.*, 1999). This could be due to the fact that assessing the effects of HNCS on the responses elicited by A β -, A δ - and C-stimuli required collecting EEG responses within very limited time periods. Therefore, we were able to collect only a small number of trials, and this could explain why no reproducible C-LEPs were identified in the average waveforms (van den Broeke & Mouraux, 2014a). In order to overcome this issue, we conducted a second experiment, which focused exclusively on C-LEPs. In this second experiment, we were able to increase the number of trials in which the stimulus selectively activated C fibers, and we were able to collect more C-LEPs in a more reliable fashion.

Conclusion

Our results suggest that HNCS indifferently inhibits the responses to nociceptive A δ and C-fiber inputs, but also to non-nociceptive A β -fiber input. This finding has important clinical implications as it indicates that the effects of HNCS on the perception or brain responses to noxious stimuli should be used cautiously as a direct measure of DNIC mechanisms and their possible involvement in patients with chronic pain.

Acknowledgements

We thank Daniel Le Bars and Emanuel Van Den Broeke for their insightful comments. MC received support from bursary of President of the Russian Federation for an internship abroad. DMT is an Academie Universitaire de Louvain-Marie Curie Incoming Post-doc. AM received support from an ERC starting grant ("PROBING-PAIN").

References

- Bouhassira D, Le Bars D, Bolgert F, Laplane D, Willer JC. Diffuse noxious inhibitory controls in humans: a neurophysiological investigation of a patient with a form of Brown-Sequard syndrome. *Annals of neurology*. 1993; 34:536–543. [PubMed: 8215241]
- Bouhassira D, Villanueva L, Bing Z, le Bars D. Involvement of the subnucleus reticularis dorsalis in diffuse noxious inhibitory controls in the rat. *Brain research*. 1992; 595:353–357. [PubMed: 1467976]
- Chebbi R, Boyer N, Monconduit L, Artola A, Luccarini P, Dallel R. The nucleus raphe magnus OFF-cells are involved in diffuse noxious inhibitory controls. *Exp Neurol*. 2014; 256:39–45. [PubMed: 24681000]
- Churyukanov M, Plaghki L, Legrain V, Mouraux A. Thermal detection thresholds of Adelta- and C-fibre afferents activated by brief CO₂ laser pulses applied onto the human hairy skin. *PloS one*. 2012; 7:e35817. [PubMed: 22558230]
- Danziger N, Rozenberg S, Bourgeois P, Charpentier G, Willer JC. Depressive effects of segmental and heterotopic application of transcutaneous electrical nerve stimulation and piezo-electric current on lower limb nociceptive flexion reflex in human subjects. *Archives of physical medicine and rehabilitation*. 1998; 79:191–200. [PubMed: 9474003]
- De Broucker T, Cesaro P, Willer JC, Le Bars D. Diffuse noxious inhibitory controls in man. Involvement of the spinoreticular tract. *Brain : a journal of neurology*. 1990; 113(Pt 4):1223–1234. [PubMed: 2397390]
- Dockstader C, Cheyne D, Tannock R. Cortical dynamics of selective attention to somatosensory events. *NeuroImage*. 2010; 49:1777–1785. [PubMed: 19781649]
- Gratton G, Coles MG, Donchin E. A new method for off-line removal of ocular artifact. *Electroencephalography and clinical neurophysiology*. 1983; 55:468–484. [PubMed: 6187540]
- Iannetti GD, Baumgartner U, Tracey I, Treede RD, Magerl W. Pinprick-evoked brain potentials: a novel tool to assess central sensitization of nociceptive pathways in humans. *Journal of neurophysiology*. 2013; 110:1107–1116. [PubMed: 23678019]
- Iannetti GD, Truini A, Galeotti F, Romaniello A, Manfredi M, Cruccu G. Usefulness of dorsal laser evoked potentials in patients with spinal cord damage: report of two cases. *Journal of neurology, neurosurgery, and psychiatry*. 2001; 71:792–794.
- Jankovski A, Plaghki L, Mouraux A. Reliable EEG responses to the selective activation of C-fibre afferents using a temperature-controlled infrared laser stimulator in conjunction with an adaptive staircase algorithm. *Pain*. 2013; 154:1578–1587. [PubMed: 23707267]
- Kakigi R, Shibasaki H, Tanaka K, Ikeda T, Oda K, Endo C, Ikeda A, Neshige R, Kuroda Y, Miyata K, et al. CO₂ laser-induced pain-related somatosensory evoked potentials in peripheral neuropathies: correlation between electrophysiological and histopathological findings. *Muscle & nerve*. 1991; 14:441–450. [PubMed: 1651448]
- Lapirot O, Chebbi R, Monconduit L, Artola A, Dallel R, Luccarini P. NK1 receptor-expressing spinoparabrachial neurons trigger diffuse noxious inhibitory controls through lateral parabrachial activation in the male rat. *Pain*. 2009; 142:245–254. [PubMed: 19231081]
- Lapirot O, Melin C, Modolo A, Nicolas C, Messaoudi Y, Monconduit L, Artola A, Luccarini P, Dallel R. Tonic and phasic descending dopaminergic controls of nociceptive transmission in the medullary dorsal horn. *Pain*. 2011; 152:1821–1831. [PubMed: 21514054]
- Le Bars D. The whole body receptive field of dorsal horn multireceptive neurones. *Brain research. Brain research reviews*. 2002; 40:29–44. [PubMed: 12589904]
- Magerl W, Ali Z, Ellrich J, Meyer RA, Treede RD. C- and A delta-fiber components of heat-evoked cerebral potentials in healthy human subjects. *Pain*. 1999; 82:127–137. [PubMed: 10467918]
- Malmierca E, Chaves-Coira I, Rodrigo-Angulo M, Nunez A. Corticofugal projections induce long-lasting effects on somatosensory responses in the trigeminal complex of the rat. *Frontiers in systems neuroscience*. 2014; 8:100. [PubMed: 24904321]
- Meyer RA, Walker RE, Mountcastle VB Jr. A laser stimulator for the study of cutaneous thermal and pain sensations. *IEEE transactions on biomedical engineering*. 1976; 23:54–60. [PubMed: 1244317]

- Millan MJ. Descending control of pain. *Progress in neurobiology*. 2002; 66:355–474. [PubMed: 12034378]
- Moont R, Crispel Y, Lev R, Pud D, Yarnitsky D. Temporal changes in cortical activation during conditioned pain modulation (CPM), a LORETA study. *Pain*. 2011; 152:1469–1477. [PubMed: 21339052]
- Nunez A, Malmierca E. Corticofugal modulation of sensory information. *Advances in anatomy, embryology, and cell biology*. 2007; 187:1–74. 1 p following table of contents.
- Plaghki L, Delisle D, Godfraind JM. Heterotopic nociceptive conditioning stimuli and mental task modulate differently the perception and physiological correlates of short CO2 laser stimuli. *Pain*. 1994; 57:181–192. [PubMed: 8090515]
- Porreca F, Ossipov MH, Gebhart GF. Chronic pain and medullary descending facilitation. *Trends in neurosciences*. 2002; 25:319–325. [PubMed: 12086751]
- Pud D, Granovsky Y, Yarnitsky D. The methodology of experimentally induced diffuse noxious inhibitory control (DNIC)-like effect in humans. *Pain*. 2009; 144:16–19. [PubMed: 19359095]
- Roby-Brami A, Bussel B, Willer JC, Le Bars D. An electrophysiological investigation into the pain-relieving effects of heterotopic nociceptive stimuli. Probable involvement of a supraspinal loop. *Brain : a journal of neurology*. 1987; 110(Pt 6):1497–1508. [PubMed: 2448000]
- Scheuren R, Anton F, Erpelding N, Michaux G. Beep tones attenuate pain following Pavlovian conditioning of an endogenous pain control mechanism. *PloS one*. 2014; 9:e88710. [PubMed: 24551138]
- Sprenger C, Bingel U, Buchel C. Treating pain with pain: supraspinal mechanisms of endogenous analgesia elicited by heterotopic noxious conditioning stimulation. *Pain*. 2011; 152:428–439. [PubMed: 21196078]
- Sprenger C, Eippert F, Finsterbusch J, Bingel U, Rose M, Buchel C. Attention modulates spinal cord responses to pain. *Current biology : CB*. 2012; 22:1019–1022. [PubMed: 22608507]
- Torta DM, Legrain V, Algoet M, Olivier E, Duque J, Mouraux A. Theta burst stimulation applied over primary motor and somatosensory cortices produces analgesia unrelated to the changes in nociceptive event-related potentials. *PloS one*. 2013; 8:e73263. [PubMed: 23977382]
- Torta DM, Legrain V, Mouraux A. Looking at the hand modulates the brain responses to nociceptive and non-nociceptive somatosensory stimuli but does not necessarily modulate their perception. *Psychophysiology*. 2015
- Toussignant-Laflamme Y, Marchand S. Excitatory and inhibitory pain mechanisms during the menstrual cycle in healthy women. *Pain*. 2009; 146:47–55. [PubMed: 19592167]
- Treede RD, Lankers J, Frieling A, Zangemeister WH, Kunze K, Bromm B. Cerebral potentials evoked by painful, laser stimuli in patients with syringomyelia. *Brain : a journal of neurology*. 1991; 114(Pt 4):1595–1607. [PubMed: 1884168]
- Treister R, Eisenberg E, Gershon E, Haddad M, Pud D. Factors affecting - and relationships between-different modes of endogenous pain modulation in healthy volunteers. *European journal of pain*. 2010; 14:608–614. [PubMed: 19906552]
- van den Broeke EN, Mouraux A. Enhanced brain responses to C-fiber input in the area of secondary hyperalgesia induced by high-frequency electrical stimulation of the skin. *Journal of neurophysiology*. 2014a; 112:2059–2066. [PubMed: 25098966]
- Van Den Broeke EN, Mouraux A. High frequency electrical stimulation of human skin induces heterotopical mechanical and heat hyperalgesia and enhanced responses to vibrotactile input. *Journal of neurophysiology*. 2014b
- van Wijk G, Veldhuijzen DS. Perspective on diffuse noxious inhibitory controls as a model of endogenous pain modulation in clinical pain syndromes. *The journal of pain : official journal of the American Pain Society*. 2010; 11:408–419. [PubMed: 20075013]
- Villanueva L, Le Bars D. The activation of bulbo-spinal controls by peripheral nociceptive inputs: diffuse noxious inhibitory controls. *Biological research*. 1995; 28:113–125. [PubMed: 8728826]
- Watanabe S, Kakigi R, Hoshiyama M, Kitamura Y, Koyama S, Shimojo M. Effects of noxious cooling of the skin on pain perception in man. *Journal of the neurological sciences*. 1996; 135:68–73. [PubMed: 8926499]

- Willer JC, De Broucker T, Le Bars D. Encoding of nociceptive thermal stimuli by diffuse noxious inhibitory controls in humans. *Journal of neurophysiology*. 1989; 62:1028–1038. [PubMed: 2585037]
- Willer JC, Le Bars D, De Broucker T. Diffuse noxious inhibitory controls in man: involvement of an opioidergic link. *European journal of pharmacology*. 1990; 182:347–355. [PubMed: 2168836]
- Willer JC, Roby A, Le Bars D. Psychophysical and electrophysiological approaches to the pain-relieving effects of heterotopic nociceptive stimuli. *Brain : a journal of neurology*. 1984; 107(Pt 4):1095–1112. [PubMed: 6509310]




Figure 1. Experimental setup.

Left panel (A). Twenty thermal stimuli activating successively C- and A δ -fibers and twenty electrical stimuli activating A β fibers were applied on the right hand dorsum before, during and after applying a heterotopic noxious conditioning stimulus (HNCS) to the left foot (immersion in a 5°C circulating water bath). The order of administration of the test stimuli was counterbalanced across participants. Right panel (B) details of how activation of C- and A δ -fibers was obtained with the laser stimulator. The ordinate represents skin surface temperature, the abscissa represents time in seconds.




Figure 2. Perceptual ratings

Median intensity of perception ratings for C, A δ , and A β stimuli presented before, during and after HNCs. Bars represent the interquartile range. Asterisks indicate significant differences between conditions at $p < 0.05$.




Figure 3. Event-related potentials (N2-P2)

Grand-average waveforms of the vertex ERPs elicited by C-, A δ - and A β - stimuli. Note that the magnitude of both the A δ -N2 and A β -P2 evoked responses was reduced during HNCS as compared to before and after HNCS. In contrast, no clear ERP could be identified following C-fiber stimulation. The bar graphs represent the difference in amplitude between conditions. Negative changes in amplitude correspond to a decrease in amplitude as compared to before HNCS.




Figure 4. Event-related potentials (N1)

Grand-average waveforms of the ERPs recorded at the contralateral central electrode (C3) following A δ - and A β -stimulation (N1). Both stimuli elicited an early-latency negative wave (N1). The magnitude of the A β -N1, but not the A δ -N1 was reduced significantly during HNCS. The bar graphs represent the difference in amplitude between conditions. Negative changes in amplitude correspond to a decrease in amplitude as compared to before HNCS.




Figure 5. Event-related potentials elicited by selective C-fiber stimulation

Grand-average waveforms of the vertex ERPs elicited by selective C-stimuli. There was a significant reduction of the vertex negative peak of the C-LEP during and after HNCS as compared to before HNCS. No significant effect of ‘block’ was observed for the vertex positive peak of the C-ERP. The bar graphs represent the difference in amplitude between conditions. Negative changes in amplitude correspond to a decrease in amplitude as compared to before HNCS.

Table 1
Perceived intensity and ERPs magnitude

Statistical values of the non-parametric analysis on the perceived intensity and on the magnitude of the ERPs. Post-hoc comparisons (Wilcoxon test) are reported when the overall analysis (Friedman Test) achieved the significance threshold.

	Friedman Test	Post-hoc comparisons (Wilcoxon)		
		Before vs. During	During vs. After	Before vs. After
<i>Ratings</i>				
A δ	p=0.001	p=0.012	p=0.012	p=0.017
C (experiment 1)	p=0.002	p=0.012	p=0.012	p=0.208
C (experiment 2)	p=0.036	p=0.012	p=0.208	p=0.050
A β	n.s.			
<i>ERPs</i>				
A δ -N2	p=0.008	p=0.017	n.s.	p=0.012
A δ -P2	n.s.			
A β -N2	p=0.021	n.s.	0.012	n.s.
A β -P2	p=0.030	p=0.017	p=0.036	p=0.484
C-N2	p=0.034	p=0.012	n.s.	p=0.036
C-P2	n.s.			