

RESEARCH ARTICLE

Open Access

The effect of sunlight exposure on interleukin-6 levels in depressive and non-depressive subjects

Rosa Levandovski^{1,4}, Bianca Pfaffenseller², Alicia Carissimi^{1,4}, Clarissa S Gama² and Maria Paz Loayza Hidalgo^{1,3,4,5*}

Abstract

Background: The objective of this epidemiological study was to evaluate the effect of length of sunlight exposure on interleukin 6 (IL-6) levels in depressive and non-depressive subjects.

Methods: This was a cross-sectional study with 154 subjects (54 males, mean age: 43.5 ± 12.8 years) who were living in a rural area in south Brazil. Chronobiological and light parameters were assessed using the Munich Chronotype Questionnaire. Sleep quality was evaluated using the Pittsburgh Sleep Quality Index. Depressive symptoms were assessed with the Beck Depression Inventory. Plasma levels of inflammatory cytokines (IL-2, IL-4, IL-6, IL-10, tumor necrosis factor- α , and interferon) were collected during the daytime and measured.

Results: IL-6 levels showed a positive correlation with light exposure ($r = 0.257$; $p < 0.001$) and a negative correlation with the mid-sleep phase on work-free days ($r = -0.177$; $p = 0.028$). Multiple linear regression analysis showed that only the length of light exposure was an independent factor for predicting IL-6 levels ($\beta = 0.26$; $p = 0.002$). In non-depressed subjects, exposure to a different intensity of light did not affect IL-6 levels ($t = -1.6$; $p = 0.1$). However, when the two depressive groups with low and high light exposure were compared, the low light exposure group had lower levels of IL-6 compared with the high light exposure group ($t = -2.19$ and $p = 0.0037$).

Conclusions: The amount of time that participants are exposed to sunlight is directly related to their IL-6 levels. Additionally, depressed subjects differ in their IL-6 levels if they are exposed to light for differing amounts of time.

Keywords: Interleukin, Light, Depressive disorder, Munich Chronotype Questionnaire (MCTQ)

Background

The pro-inflammatory cytokine interleukin-6 (IL-6) is involved in the regulation of several physiological processes, particularly in the immune response, as well as in sleep regulation [1], metabolism [2,3], and mood disorders [4,5].

IL-6 is produced by a variety of cells, including mononuclear phagocytes, T cells, fibroblasts, astrocytes, and microglia cells [6]. Although the light/dark circadian rhythm has been shown to regulate IL-6, only a few studies have been performed in humans. Diurnal variation of IL-6 has been reported in healthy subjects [7,8] and also in some clinical conditions, such as rheumatoid

arthritis, myocardial infarction, and sleep deprivation [9-12]. Administration of a synthetic nuclear receptor involved in the circadian clockwork, the Rev-Erb ligand, modulates the production and release of IL-6 in human macrophages [13].

Diurnal variation of IL-6 to the light/dark rhythm is modulated by melatonin, which has been implicated in the production of IL-6 [14]. The suprachiasmatic nucleus (SCN) may be responsible for this modulation. The SCN sends information regarding the dark-light cycle to the paraventricular nucleus of the hypothalamus, to the lateral column of the spinal cord, and to the pineal gland to modulate melatonin release [15]. The melatonin rhythm plays the role of an endogenous synchronizer, which peaks at night and is able to entrain rhythms, such as temperature, hormone release, and the sleep-wake cycle via receptors in the SCN [15,16]. We have previously shown that, in depressed patients, excretion of the major melatonin urinary metabolite,

* Correspondence: mpaz@cpovo.net

¹Laboratório de Cronobiologia do Hospital de Clínicas de Porto Alegre (HCPA), da Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil

³Departamento de Psiquiatria e Medicina Legal da Faculdade de Medicina, da Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

Full list of author information is available at the end of the article

6-sulfatoxymelatonin (aMT6s), increases after antidepressant use. The higher the increment in aMT6s excretion, the better the antidepressant response of the drug, indicating an enhancement in pineal production of melatonin overnight [17]. Additionally, this effect may be related to changes in phase or poor sleep quality found in depressed patients [18] or subjects with greater levels of depressive symptoms [19].

Catecholamines also play an important role in the release of melatonin. Norepinephrine, which is an important neurotransmitter involved in mood disorders and pathophysiology, is involved in the regulation of the enzymatic activity that converts serotonin in N-acetylserotonin, the immediate precursor of melatonin [20]. Furthermore, catecholamines may exert a direct effect on the immunoinflammatory system, given that IL-6 release is stimulated by noradrenaline, whereas IL-6 inhibits the secretion of dopamine and serotonin [21,22].

Proinflammatory cytokines are thought to be involved in the pathogenesis of depression. However, previous studies have reached divergent conclusions about the relationship between depression and the levels of IL-6. Increased IL-6 levels have been observed during depression [4,23-28]. However, psychological symptoms, such as a negative mood, tension, and fatigue, are significantly associated with a greater nocturnal decrease of IL-6 levels [29]. IL-6, melatonin, cortisol, and depressive symptoms are not significantly altered by light therapy

[4,30]; however, natural light exposure improves depressive symptoms [30].

Based on this evidence, we hypothesized that IL-6 levels are directly correlated with sunlight exposure and not just with artificial light exposure. Therefore, this epidemiological study evaluated the effects of the duration of sunlight exposure on IL-6 levels in depressive and non-depressive subjects.

Methods

Study population

This was a cross-sectional study of an epidemiological sample (i.e., homogeneous in terms of culture and socioeconomic level), living in a rural area in the south of Brazil. Notably, this area was unaffected by the nocturnal light pollution found in urban areas. The sample comprised 154 subjects with a mean age of 43.5 ± 12.8 years, ranging from 18 to 65 years of age. All subjects with a Beck Depression Inventory (BDI) ≥ 10 were selected from a larger sample of 6505 interviewed subjects. A total of 124 healthy subjects were randomly selected for the current study from 1500 subjects without a personal history of disease (Figure 1).

Exclusion criteria included sleep disturbance, cancer-related, rheumatological, endocrine, or neurological disease, co-morbidities related to inflammation, as well as the use of anti-inflammatory, antihypertensive, and psychopharmacological drugs, and cortisol. The study was




Figure 1 Flowchart of the inclusion procedure.

conducted according to international ethical standards protocols [31] (ethics approval number: 08-087 GPPG/HCPA), and all participants provided informed consent.

Instruments

Chronobiological and light parameters were assessed using the Munich Chronotype Questionnaire (MCTQ). Sleep quality was evaluated using the Pittsburgh Sleep Quality Index (PSQI), and depressive symptoms were evaluated with the BDI.

Research assistants blinded to the objective of the study and previously trained to avoid bias in measurements collected our data. Interviews were held during the day in the participant's house. To guarantee blinding, the questionnaires contained several questions that were not related to depressive symptoms. Questionnaires were answered individually, following the instructions for each question.

The MCTQ

We used the validated Brazilian-Portuguese version of the MCTQ (<http://www.bioinfo.mpg.de/wepcronotipo/>) to assess the mid-sleep phase on work-free days (MSFsc), average sleep duration, and average light exposure. The MCTQ assesses actual sleep times separately for work and work-free days. The MSFsc was calculated as the point of the mid-sleep phase on a work-free day. Sleep duration was calculated by averaging the estimates of sleep duration for both work and work-free days, over the course of 1 week [32]. The questionnaires provided information regarding sleep times, such as bed and rise times, plus the clock time of becoming fully awake, as well as sleep latency and inertia. In addition, the instrument has questions related to the total time of exposition to natural light, during work, and work-free days.

Outdoor sunlight exposure measurements

Outdoor light exposure was assessed using the validated version of the MCTQ. The questionnaire includes questions related to the total time of exposure to sunlight, during work, and work-free days. The time of light exposure is self-reported using a standard question: "How long per day do you spend on average outside (really outside) exposed to day light? On work days: ___ h ___ min and on free days: ___ h ___ min."

The average phase of sunlight exposure was defined by the average amount of time an individual was exposed to outdoor light, and was calculated by averaging the information for work and work-free days throughout the week using the equation: number of work days per week \times number of hours reported on work days + number of work-free days per week \times number of hours reported on work-free days [33].

The PSQI

The PSQI [34] was used to measure sleep quality in this study. The PSQI version used was validated for Brazilian Portuguese language [35]. This index is composed of 15 multiple-choice items that inquire about the frequency of sleep disturbances, the subjective sleep quality, as well as four items that inquire about typical bedtime, time of awakening, sleep latency, and sleep duration. The PSQI generates seven subscores that correspond to the domains of the global score. Each component score ranges from 0 (indicating no difficulty) to 3 (indicating severe difficulty). The component scores are summed to obtain a global score (range of 0–21). A score > 5 is suggestive of significant sleep complaints.

The BDI

We chose the validated Brazilian Portuguese version of the BDI for the screening of depressive symptoms [36,37]. This is a self-report scale that assesses cognitive, affective, and somatic symptoms of depression. The inventory consists of 21 items, with each question assigned to a score ranging from 0 (no symptoms) to 3 (severe symptoms), with the total BDI score ranging from 0 to 63. A total score of 0–9 indicates no or minimal depressive symptoms, 10–17 indicates mild depressive symptoms, 18–29 indicates moderate depressive symptoms, and 30–63 indicates severe depressive symptoms [38].

Cytokine measurements

Inflammatory cytokine levels (IL-2, IL-4, IL-6, IL-10, tumor necrosis factor- α [TNF- α], and interferon) were assayed in plasma, which was collected during the day. Five milliliters of blood was collected by venipuncture into an EDTA vacuum tube. Immediately after withdrawal, the blood was centrifuged, and the plasma was aliquoted and stored at -80°C until later assay. Plasma cytokine levels were determined by flow cytometry using the BD™ Cytometric Bead Array (CBA) Human Th1/Th2/Th17 Cytokine Kit (BD Biosciences, San Diego, CA). The CBA kit allows for the discrimination of the following cytokines: IL-2, IL-4, IL-6, IL-10, interferon- γ , TNF- α , and interferon. Sample processing and data analysis were performed according to the manufacturer's instructions. Briefly, plasma samples were incubated with cytokine capture beads for 1.5 h, washed, and then incubated for 1.5 h with PE-conjugated detection antibodies. Both incubations were performed at room temperature and protected from the light. The samples were then washed, and sample data were acquired using a FACSCalibur flow cytometer (BD Biosciences). The results were generated in both graphical and tabular formats using the BD CBA Analysis Software FCAP Array™ (BD Biosciences).

Statistical analysis

Logarithmic transformations were completed for the following variables because they showed asymmetric distributions: IL-2, IL-4, IL-6, IL-10, TNF- α , interferon, and BDI. Spearman's test was used to analyze the correlation among ILs and sleep parameters. To control for potential confounding effects, variables were analyzed by logistic regression analysis.

Average light exposure was categorized into two groups based on the 75th percentile: low and high light exposure. Additionally, BDI scores were categorized into two groups: less than 10 (non-depressive) and equal to or greater than 10 (depressive). A two-tailed Student's *t*-test for independent samples was used to analyze the mean difference between groups. For all analyses, the statistical significance was set at $p < 0.05$. Data were analyzed using SPSS version 18.0 (SPSS, Chicago, IL).

Results

With regard to symptoms of depression, approximately 19.5% of the sample population scored ≥ 10 on the BDI scale. The maximum score was 34 points, corresponding to mild to moderate depressive symptoms. Demographics, sleep characteristics, and anthropometric data of all subjects included in this study ($n = 154$) are shown in Table 1. The depressive group and non-depressive group were not different in terms of age (44.0 ± 12.28 vs 43.4 ± 12.95 years; $t = -0.27$; $p = 0.78$) and body mass index (25.2 ± 3.9 vs 25.6 ± 4.7 ; $t = 0.42$; $p = 0.68$). Moreover, mean IL-6 levels were not significantly different between women and men (1.3 ± 1.1 vs 1.1 ± 1.0 pg/ml; $t = 1.14$; $p = 0.26$). Therefore, these variables were not included in multivariate analyses because they were not potential confounding factors.

Table 1 Sample characteristics (n = 154)

Variable	Mean \pm Sd/n(%)
Age	43.49 \pm 12.8
Sex (Male)	54 (34.8%)
BDI score	5.9 \pm 5.9
Body mass index	25.6 \pm 4.6
Formal education (years)	7.7 \pm 3.6
MSFsc	3.34 \pm 1.7
Average Light Exposure	8.70 \pm 2.7
PSQI score	5.12 \pm 4.1
Average Sleep Duration	7.84 \pm 1.3
<i>Beck Depression Inventory</i>	
BDI <10	124 (80.5%)
BDI >10	30 (19.5%)

Values are means and standard deviation; BDI, Beck Depression Inventory; MSFsc, mid-sleep phase on work-free days sleep corrected; PSQI, Pittsburgh Sleep Quality Index.

Upon examination of the entire sample, a positive correlation was found between IL-6 levels and the average amount of sunlight exposure time (Spearman's $r = 0.26$; $p = 0.001$; Figure 2), and a negative correlation was found between IL-6 levels and MSFsc (Spearman's $r = -0.177$; $p = 0.028$; Table 2). However, there were no correlations among the average sleep duration, quality of sleep based on PSQI scores, and cytokine parameters (Table 2).

To control for potential multicollinearity effects among variables, we analyzed the data with multivariate regression analysis, with covariate factors, including chronobiological, sleep, and light parameters, predicting the outcome variable, IL-6. The only dependent variable that was retained in the model was the average amount of sunlight exposure time. Therefore, MSFsc was considered a confounding variable ($\beta = -0.05$; $p = 0.57$). The average amount of sunlight exposure accounted for variance in IL-6 levels ($\beta = 0.26$; $p = 0.002$) ($F = 6.304$; R^2 adjusted = 0.065; $p = 0.002$).

IL-6 levels were not different among the depressive and non-depressive groups (Figure 3A). In the non-depressive group, comparison of IL-6 levels in subjects with high and low light exposure yielded no group differences ($t = -1.6$; $p = 0.1$; Figure 3B). However, when the low and high light exposure groups were compared in subjects with depressive symptoms, the low light exposure group showed lower levels of IL-6 compared with the high light exposure group ($t = -2.19$ and $p = 0.0037$; Figure 3C). This finding indicates that, when subjects are depressed, the length of sunlight exposure is related to IL-6 levels.

Discussion

In this epidemiological study, our findings showed that the amount of sunlight exposure time was directly related to plasma IL-6 levels. Moreover, depressive subjects with high amounts of light exposure had significantly greater mean IL-6 levels compared with depressive subjects with low light exposure. These results are in accordance with the notion that exposure to sunlight is one of the most powerful factors that affects multiple physiological processes in humans, including the immune system.

Exposure to natural light has been the focus of human studies to understand the physiological effects of various 24-h patterns on the light-dark cycle, in contrast to effects of artificial light [39]. Our results demonstrated that depressed subjects differed in their IL-6 levels based on differing amounts of sunlight exposure. Behavior promoting natural light exposure is a simple and low-cost therapeutic practice to alleviate depression [30]. Therefore, the mechanism that links the therapeutic effect of light in depression may be the immune system, specifically by IL-6.

Our study reassessed the hypothesis that the link between depressive symptoms and immune responses is




Figure 2 The logarithmic values of IL-6 levels were plotted against the average sunlight exposure. There was a positive correlation between IL-6 levels and average light exposure. IL-6 levels are shown in pg/ml. IL = interleukin.

related to functioning of the SCN [40] and immune-pineal-axis, which are highly sensitive to light [20,41]. Unfortunately, we did not assess melatonin levels in our study. Therefore, this is an important hypothesis to be tested in a clinical study. Recently, inflammatory signals have been linked to the pineal gland, which produces melatonin. The immune-pineal-axis is one of the regulatory mechanisms responsible for the regulation of the inflammatory system by melatonin production [20].

Humans are likely to differ in their response to light depending on impairment of their brain, and therefore, depressed patients may be more prone to being reactive to light. This relationship can be explained by findings from previous reports, such as the demonstration that IL-6 has a protective effect at lower concentrations in brain tissue, but a neurotoxic effect at high concentrations [42]. However, even though IL-6 is a proinflammatory cytokine, it is not always beneficial and it is involved in acute inflammatory disease. Patients with acute episodes and those with sepsis have intermediate levels of interleukins, showing a systemic toxicity with mood episodes, compared with healthy subjects [43,44]. In a clinical study, patients with depressive episodes demonstrated

an increase in IL-6 levels, whereas patients with manic episodes showed an increase in IL-2, IL-4, and IL-6 levels [45].


In vitro, IL-6 promotes neuronal survival [46], exerts neuroprotective properties in cells against cytotoxic insults [47], and influences synaptic plasticity, as suggested by the modulation of synaptic potentiation in the hippocampus [48], with subsequent promotion of memory formation [49]. We hypothesize that the regulatory process of IL-6, promoted by the SCN, does not occur in the same manner in depressed patients compared with controls.

We did not assess the intensity or the quality of light using an objective instrument. Nevertheless, our sample was composed of rural individuals who worked outside. Therefore, it is likely that they are exposed to sunlight for longer amounts of time and are exposed to a different quality of light compared with European populations. In Brazil (30°S; 51°W), the temperature is high and the photoperiod is long. Given that light exposure is mediated by numerous factors, evaluation of the quantity of sunlight exposure, as well as the intensity and quality of light in field work, should be performed in future studies.

Table 2 Correlations between cytokines, chronobiological, and sleep variables

	MSFsc	Average light exposure	Average sleep duration	PSQI score
	Spearman	Spearman	Spearman	Spearman
	(p value)	(p value)	(p value)	(p value)
IL-2*	0.063 (0.44)	-0.026 (0.75)	-0.040 (0.63)	-0.063 (0.44)
IL-4*	-0.030 (0.71)	0.0659 (0.43)	-0.039 (0.63)	-0.015 (0.86)
IL-6*	-0.177 (0.03)	0.256 (0.001)	-0.019 (0.82)	0.050 (0.54)
IL-10*	0.034 (0.68)	0.022 (0.79)	0.057 (0.48)	-0.060 (0.46)
TNF-α*	-0.071 (0.38)	0.076 (0.35)	-0.038 (0.64)	0.066 (0.42)
Interferon*	0.062 (0.44)	-0.011 (0.90)	-0.146 (0.07)	-0.022 (0.78)

MSFsc = mid-sleep phase on work-free days sleep corrected; PSQI = Pittsburgh Sleep Quality Index; IL = Interleukin. * Variables log transformed.



The divergent findings related to IL-6 levels in mood disorders in the literature may be a consequence of differences in the methods and samples used. In our study, we chose only one time point for blood collection to assess cytokine levels throughout the day, and this may have limited our conclusions because of IL-6 rhythmicity in peripheral blood, which may be altered by day length. Nevertheless, our results demonstrated that, based on a positive linear correlation, IL-6 levels increased proportionally to longer periods of sunlight exposure. In addition, our sample was composed of mild and moderately depressed subjects who had never undergone treatment. Therefore, there was no effect of medication on IL-6 in this sample. The potential confounding effects of quality and phase of sleep were also controlled in our study.

Conclusions

The amount of time that participants are exposed to sunlight is directly related to their IL-6 levels. Additionally, depressed subjects differ in their IL-6 levels based on the duration of sunlight exposure. To the best of our knowledge, this is the first epidemiological study examining the crosstalk between circadian variations in sunlight and immune system parameters in depressive and non-depressive subjects. These findings underscore the hypothesis that the beneficial effect of sunlight exposure on depressive disorders is attributable to protection by ILs. We hypothesize that differences in IL-6 levels in depressive subjects are a result of impairment of the central nervous system, more specifically, the SCN.

Abbreviations

IL-6: Interleukin 6; MCTQ: Munich chronotype questionnaire; BDI: Beck depression inventory; SCN: Suprachiasmatic nucleus; (aMT6s): 6-sulfatoxymelatonin; PSQI: Pittsburgh sleep quality index; MSFsc: Mid-sleep phase on work-free days sleep corrected; CBA: Cytometric bead array.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RL and MPLH designed the study, wrote the protocol, and performed literature searches and analyses. RL, AC, and MPLH wrote the first draft of the manuscript. BP and CSG performed cytokine measurements, quality control, and the final revision of the manuscript. All authors contributed to and have approved the final manuscript. RL (Rosa Levandovski); MPLH (Maria Paz Loayza Hidalgo); AC (Alicia Carissimi); BP (Bianca Pfaffenseller); CSG (Clarissa S Gama).

Acknowledgements

We thank the graduate students Alicia Deitos, Ana Claudia Souza, Fabiane Dresch, Gabriela Laste, Janaina da Silveira (UNIVATES), Andre Oliveira Marques, Cristiane Koplin, Diego Fraga, Fabiana Guarienti, Jane Cronst, and Manoela Jornada (UFRGS) for their assistance in data collection and data management. We thank Juliana Vieira and Giovana Dantas for their technical support, and Professors' Karla Allebrandt, Wolnei Caumo, Iraci Torres, and Luciana Fernandes for their advice. This work was supported by FIPE-HCPA/UFRGS; UNIVATES; CNPq; PNPD-CAPES/Brazil; PROBRAL-CAPES/DAAD (Programa Brasil-Alemanha [PROBRAL] between Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior [CAPES] - Brazil and Deutscher Aka-demischer Austausch Dienst [DAAD] - Germany).

Author details

¹Laboratório de Cronobiologia do Hospital de Clínicas de Porto Alegre (HCPA), da Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil. ²INCT for Translational Medicine, Laboratory of Molecular Psychiatry, HCPA/UFRGS, Porto Alegre, Brazil. ³Departamento de Psiquiatria e Medicina Legal da Faculdade de Medicina, da Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil. ⁴Programa de Pós-Graduação em Ciências Médicas: Psiquiatria, UFRGS, Porto Alegre, Brazil. ⁵Programa de Pós-Graduação em Medicina: Ciências Médicas, UFRGS, Porto Alegre, Brazil.

Received: 20 November 2012 Accepted: 25 February 2013

Published: 5 March 2013

References

1. Vgontzas AN, Bixler EO, Lin HM, Prolo P, Trakada G, Chrousos GP: **IL-6 and its circadian secretion in humans.** *Neuroimmunomodulation* 2005, **12**:131–140.
2. Wallenius V, Wallenius K, Ahren B, Rudling M, Carlsten H, Dickson SL, Ohlsson C, Jansson JO: **Interleukin-6-deficient mice develop mature-onset obesity.** *Nat Med* 2002, **8**:75–79.
3. Vanitallie TB: **Sleep and energy balance: Interactive homeostatic systems.** *Metabolism* 2006, **55**:S30–35.
4. Leu SJ, Shiah IS, Yatham LN, Cheu YM, Lam RW: **Immune-inflammatory markers in patients with seasonal affective disorder: effects of light therapy.** *J Affect Disord* 2001, **63**:27–34.
5. Monje FJ, Cabatic M, Divisch I, Kim EJ, Herkner KR, Binder BR, Pollak DD: **Constant darkness induces IL-6-dependent depression-like behavior through the NF- κ B signaling pathway.** *J Neurosci* 2011, **31**:9075–9083.
6. Feghali CA, Wright TM: **Cytokines in acute and chronic inflammation.** *Front Biosci* 1997, **1**:d12–26.
7. Haack M, Kraus T, Schuld A, Dalal M, Koethe D, Pollmächer T: **Diurnal variations of interleukin-6 plasma levels are confounded by blood drawing procedures.** *Psychoneuroendocrinology* 2002, **27**(8):921–31.
8. Miles MP, Andring JM, Pearson SD, Gordon LK, Kasper C, Depner CM, Kidd JR: **Diurnal variation, response to eccentric exercise, and association of inflammatory mediators with muscle damage variables.** *J Appl Physiol* 2008, **104**(2):451–8.
9. Crofford LJ, Kalogerias KT, Mastorakos G, Magiakou MA, Wells J, Kanik KS, Gold PW, Chrousos GP, Wilder RL: **Circadian relationships between interleukin (IL)-6 and hypothalamic-pituitary-adrenal axis hormones: failure of IL-6 to cause sustained hypercortisolism in patients with early untreated rheumatoid arthritis.** *J Clin Endocrinol Metab* 1997, **82**(4):1279–83.
10. Perry MG, Kirwan JR, Jessop DS, Hunt LP: **Overnight variations in cortisol, interleukin 6, tumour necrosis factor alpha and other cytokines in people with rheumatoid arthritis.** *Ann Rheum Dis* 2009, **68**(1):63–68.
11. Ridker PM, Rifai N, Stampfer MJ, Hennekens CH: **Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men.** *Circulation* 2000, **101**(15):1767–72.
12. Voderholzer U, Fiebich BL, Dersch R, Feige B, Piosczyk H, Kopasz M, Riemann D, Lieb K: **Effects of sleep deprivation on nocturnal cytokine concentrations in depressed patients and healthy control subjects.** *J Neuropsychiatry Clin Neurosci* 2012, **24**(3):354–66.
13. Gibbs JE, Blaikley J, Beesley S, Matthews L, Simpson KD, Boyce SH, Farrow SN, Else KJ, Singh D, Ray DW, Loudon AS: **The nuclear receptor REV-ERB.alpha; mediates circadian regulation of innate immunity through selective regulation of inflammatory cytokines.** *Proc Natl Acad Sci USA* 2012, **109**(2):582–587.
14. Garcia-Mauriño S, Gonzalez-Haba MG, Calvo JR, Rafii-El-Idrissi M, Sanchez-Margalet V, Goberna R, Guerrero JM: **Melatonin enhances IL-2, IL-6, and IFN-gamma production by human circulating CD4+ cells: a possible nuclear receptor-mediated mechanism involving T helper type 1 lymphocytes and monocytes.** *J Immunol* 1997, **159**(2):574–81.
15. Kalsbeek A, Yi CX, Cailotto C, la Fleur SE, Fliers E, Buijs RM: **Mammalian clock output mechanisms.** *Essays Biochem* 2011, **49**(1):137–51.
16. Claustrat B, Brun J, Chazot G: **The basic physiology and pathophysiology of melatonin.** *Sleep Med Rev* 2005, **9**(1):11–24.
17. Hidalgo MP, Caumo W, Dantas G, Franco DG, Torres IL, Pezzi J, Elisabethy E, Detanico BC, Piato A, Markus RP: **6--Sulfatoxymelatonin as a predictor of clinical outcome in depressive patients.** *Hum Psychopharmacol* 2011, **26**(3):252–7.
18. Hidalgo MP, Caumo W, Posser M, Coccaro SB, Camozzato AL, Chaves ML: **Relationship between depressive mood and chronotype in healthy subjects.** *Psychiatry Clin Neurosci* 2009, **63**(3):283–90.
19. Levandovski R, Dantas G, Fernandes LC, Caumo W, Torres I, Roenneberg T, Hidalgo MP, Allebrandt KV: **Depression scores associate with chronotype and social jetlag in a rural population.** *Chronobiol Int* 2011, **28**(9):771–8.
20. Couto-Moraes R, Palermo-Neto J, Markus RP: **The immune-pineal axis: stress as a modulator of pineal gland function.** *Ann N Y Acad Sci* 2009, **1153**:193–202.
21. Seres J, Herichová I, Roman O, Bornstein S, Jurcovicová J: **Evidence for daily rhythms of the expression of proopiomelanocortin, interleukin-1-beta and interleukin-6 in adenopituitaries of male long-evans rats: effect of adjuvant arthritis.** *Neuroimmunomodulation* 2004, **11**(5):316–322.
22. Miranda HC, Nunes SOB, Reiche EMV, Oda JMM, Watanabe MAE: **Higher than Normal Plasma Interleukin-6 Concentrations in Brazilian Patients with Mood Disorders.** *Braz Arch Biol Technol* 2011, **54**(4):717–722.
23. Berk M, Wade AA, Kuschke RH, O'Neill-Kerr A: **Acute phase proteins in major depression.** *J Psychosom Res* 1997, **43**:529–534.
24. Maes M, Meltzer HY, Bosmans E, Bergmans R, Vandoolaeghe E, Ranjan R, Desnyder R: **Increased plasma concentrations of interleukin-6, soluble interleukin-6, soluble interleukin-2 and transferrin receptor in major depression.** *J Affect Disord* 1995, **34**:301–309.
25. Maes M, Bosmans E, De Jongh R, Kenis G, Vandoolaeghe E, Neels H: **Increased serum IL-6 and IL-1 receptor antagonist concentrations in major depression and treatment resistant depression.** *Cytokine* 1997, **9**:853–858.
26. Zeugmann S, Quante A, Heuser I, Schwarzer R, Angheliescu I: **Inflammatory biomarkers in 70 depressed inpatients with and without the metabolic syndrome.** *J Clin Psychiatry* 2010, **71**:1007–1016.
27. Lehto SM, Niskanen L, Miettola J, Tolmunen T, Viinamäki H, Mäntyselkä P: **Serum anti-inflammatory markers in general population subjects with elevated depressive symptoms.** *Neurosci Lett* 2010, **484**:201–205.
28. Koutsouraki E, Hatzifilipou E, Michmizos D, Cotsavasiloglou C, Costa V, Baloyannis S: **Increase in interleukin-6 levels is related to depressive phenomena in the acute (relapsing) phase of multiple sclerosis.** *J Neuropsychiatry Clin Neurosci* 2011 Fall, **23**:442–448.
29. Rief W, Mills PJ, Ancoli-Israel S, Ziegler MG, Pung MA, Dimsdale JE: **Overnight changes of immune parameters and catecholamines are associated with mood and stress.** *Psychosom Med* 2010, **72**:755–762.
30. Wirz-Justice A, Graw P, Kräuchi K, Sarrafzadeh A, English J, Arendt J, Sand L: **'Natural' light treatment of seasonal affective disorder.** *J Affect Disord* 1996, **37**(2–3):109–120.
31. Portaluppi F, Smolensky MH, Touitou Y: **Ethics and methods for biological rhythm research on animals and human beings.** *Chronobiol Int* 2010, **27**:1911–1929.
32. Wittmann M, Dinich J, Mewes M, Roenneberg T: **Social jetlag: misalignment of biological and social time.** *Chronobiol Int* 2006, **23**:497–509.
33. Roenneberg T, Wirz-Justice A, Mewes M: **Life between clocks: daily temporal patterns of human chronotypes.** *J Biol Rhythms* 2003, **18**(1):80–90.
34. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ: **The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research.** *Psychiatr Res* 1989, **28**:193–213.
35. Bertolazi AN, Fagundes SC, Hoff LS, Dartora EG, Miozzo IC, de Barba ME, Barreto SS: **Validation of the Brazilian Portuguese version of the Pittsburgh Sleep Quality Index.** *Sleep Med* 2011, **12**:70–5.
36. Gorenstein C, Andrade L: **Validation of a Portuguese version of the Beck Depression Inventory and the State-Trait Anxiety Inventory in Brazilian subjects.** *Braz J Med Biol Res* 1996, **29**:453–457.
37. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J: **An inventory for measuring depression.** *Arch Gen Psychiatry* 1961, **4**:561–571.
38. Lasa L, Ayuso-Mateos JL, Vázquez-Barquero JL, Díez-Manrique FJ, Dowrick CF: **The use of the Beck Depression Inventory to screen for depression in the general population: a preliminary analysis.** *J Affect Disord* 2000, **57**:261–265.
39. Dumont M, Beaulieu C: **Light exposure in the natural environment: relevance to mood and sleep disorders.** *Sleep Med* 2007, **8**(6):557–65.

40. Kalsbeek A, van der Spek R, Lei J, Enderit E, Buijs RM, Fliers E: **Circadian rhythms in the hypothalamo-pituitary-adrenal (HPA) axis.** *Mol Cell Endocrinol* 2012, **349**(1):20–9.
41. da Silveira C-MS, Pinato L, Tamura EK, Carvalho-Sousa CE, Markus RP: **Glia-pinealocyte network: the paracrine modulation of melatonin synthesis by tumor necrosis factor (TNF).** *PLoS One* 2012, **7**(7):e40142.
42. Muñoz-Fernández MA, Fresno M: **The role of tumour necrosis factor, interleukin 6, interferon-gamma and inducible nitric oxide synthase in the development and pathology of the nervous system.** *Prog Neurobiol* 1998, **56**(3):307–40.
43. Kapczinski F, Dal-Pizzol F, Teixeira AL, Magalhaes PV, Kauer-Sant'Anna M, Klamt F, Pasquali MA, Quevedo J, Gama CS, Post R: **A systemic toxicity index developed to assess peripheral changes in mood episodes.** *Mol Psychiatry* 2010, **15**(8):784–6.
44. Kapczinski F, Dal-Pizzol F, Teixeira AL, Magalhaes PV, Kauer-Sant'Anna M, Klamt F, Moreira JC, de Bittencourt Pasquali MA, Fries GR, Quevedo J, Gama CS, Post R: **Peripheral biomarkers and illness activity in bipolar disorder.** *J Psychiatr Res* 2011, **45**(2):156–61.
45. Brietzke E, Stertz L, Fernandes BS, Kauer-Sant'anna M, Mascarenhas M, Escosteguy Vargas A, Chies JA, Kapczinski F: **Comparison of cytokine levels in depressed, manic and euthymic patients with bipolar disorder.** *J Affect Disord* 2009, **116**(3):214–7.
46. Li XZ, Bai LM, Yang YP, Luo WF, Hu WD, Chen JP, Mao CJ, Liu CF: **Effects of IL-6 secreted from astrocytes on the survival of dopaminergic neurons in lipopolysaccharide-induced inflammation.** *Neurosci Res* 2009, **65**(3):252–8.
47. Spittau B, Zhou X, Ming M, Kriegstein K: **IL6 Protects MN9D Cells and Midbrain Dopaminergic Neurons from MPP(+)-Induced Neurodegeneration.** *Neuromolecular Med* 2012, **14**(4):317–27.
48. Li AJ, Katafuchi T, Oda S, Hori T, Oomura Y: **Interleukin-6 inhibits long-term potentiation in rat hippocampal slices.** *Brain Res* 1997, **748**(1–2):30–8.
49. Bellinger FP, Madamba SG, Campbell IL, Siggins GR: **Reduced long-term potentiation in the dentate gyrus of transgenic mice with cerebral overexpression of interleukin-6.** *Neurosci Lett* 1995, **198**(2):95–8.

doi:10.1186/1471-244X-13-75

Cite this article as: Levandovski et al.: The effect of sunlight exposure on interleukin-6 levels in depressive and non-depressive subjects. *BMC Psychiatry* 2013 **13**:75.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

