

 Open access • Posted Content • DOI:10.1101/2020.10.15.341156

The Effects of Inter-Individual Biological Differences and Taphonomic Alteration on Human Bone Protein Profiles: Implications for the Development of PMI/AAD Estimation Methods — [Source link](#)

[Hayley L. Mickleburgh](#), [Hayley L. Mickleburgh](#), [Edward C. Schwalbe](#), [Haruka Mizukami](#) ...+5 more authors

Institutions: [Texas State University](#), [Linnaeus University](#), [Northumbria University](#), [University of Turin](#) ...+1 more institutions

Published on: 15 Oct 2020 - [bioRxiv](#) (Cold Spring Harbor Laboratory)

Topics: [Bone mineral](#)

Related papers:

- [Human Bone Proteomes before and after Decomposition: Investigating the Effects of Biological Variation and Taphonomic Alteration on Bone Protein Profiles and the Implications for Forensic Proteomics.](#)
- [Intra- and Interskeletal Proteome Variations in Fresh and Buried Bones](#)
- [Aquatic Decomposition of Mammalian Corpses: A Forensic Proteomic Approach.](#)
- [Usefulness of protein analysis for detecting pathologies in bone remains.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-effects-of-inter-individual-biological-differences-and-4r7imdow45>

1 **The Effects of Inter-Individual Biological Differences and Taphonomic** 2 **Alteration on Human Bone Protein Profiles: Implications for the** 3 **Development of PMI/AAD Estimation Methods**

4 Hayley L. Mickleburgh^{1,2}, Ed Schwalbe³, Haruka Mizukami³, Federica Sellitto³, Sefora Starace⁴,
5 Daniel J. Wescott², David O. Carter⁵, Noemi Procopio^{3*}

6
7 ¹ *Department of Cultural Sciences, Linnaeus University, Kalmar, Sweden*

8
9 ² *Forensic Anthropology Center, Texas State University, San Marcos, Texas*

10
11 ³ *The Forensic Science Unit, Faculty of Health and Life Sciences, Ellison Building, Northumbria*
12 *University, Northumbria University Newcastle, Newcastle Upon Tyne, NE1 8ST, United*
13 *Kingdom*

14
15 ⁴ *Dipartimento di Chimica, University of Turin, Via P. Giuria 7, 10125 Turin, Italy*

16
17 ⁵ *Forensic Sciences Unit, School of Natural Sciences and Mathematics, Chaminade University*
18 *of Honolulu, Hawaii*

19 20 21 **Abstract**

22 Bone proteomics studies using animal proxies and skeletonized human remains have
23 delivered encouraging results in the search for potential biomarkers for precise and accurate
24 post-mortem interval (PMI) and the age-at-death (AAD) estimation in medico-legal
25 investigations. At present, however, the effects of inter-individual biological differences and
26 taphonomic alteration on recovered human bone protein profiles are not well understood.
27 This study investigated the human bone proteome in four human body donors studied
28 throughout decomposition outdoors. The effects of ageing phenomena (*in vivo* and post-
29 mortem), and intrinsic and extrinsic variables on the variety and abundance of the bone
30 proteome were assessed. Results identified a new potential biomarker for PMI estimation, as
31 well as three potential biomarkers for AAD estimation. The results also suggest that bone
32 mineral density (BMD) may be an important variable affecting the survival and extraction of
33 proteins.

34 35 **Highlights**

- 36 - CO3, CO9, COBA2, CO3A1, MGP, PGS2 and TTHY are potential biomarkers for post-
37 mortem interval estimation in skeletonized human remains
- 38 - FETUA, ALBU and OLFL3 are potential biomarkers for age-at-death estimation in human
39 remains
- 40 - Taphonomic and biological variables play a significant role in survival and extraction rates
41 of proteins in bone
- 42 - Bone mineral density may affect survival of proteins in bone, probably due to the effects
43 of the mineral matrix on the movement of decomposer microbes
- 44 - Higher bone mineral density may affect the survival and the extraction rate of collagen
45 and mineral-binding proteins

47 **Key words**

48 Forensic proteomics, forensic taphonomy, bone mineral density, post-mortem interval
49 estimation, age-at-death estimation, decomposition, postmortem microbiology

50

51 **1. Introduction**

52 Estimations of the time elapsed since death (post-mortem interval, PMI) and the age-at-death
53 (AAD) are crucial in the forensic investigation of unidentified human remains. This
54 information is important to distinguish between historical remains (>100 years old) and
55 remains of medico-legal relevance (≤ 100 years old)^{1,2}, and to narrow the search of missing
56 persons for identification purposes^{3,4}. High precision, accuracy and objectivity of PMI and AAD
57 estimation methods are essential in order to be considered admissible in a court of law.

58 PMI estimation often relies on visual assessment of gross morphological changes of
59 the body during decomposition⁵⁻⁷, even though the rate of these changes is known to be
60 highly variable^{8,9}. Accuracy of the PMI estimation decreases as decomposition progresses,
61 and interobserver reliability differs depending on the method and the experience of the
62 researcher^{9,10}. Biochemical techniques have shown promising results in the search for a
63 precise and accurate method to estimate late PMI in human bone, however, these methods
64 are yet to be validated for use in forensic contexts¹¹⁻¹⁴.

65 Standard AAD estimation methods are based on the examination of the morphological
66 characteristics of the remains¹⁵, and require the evaluation of several different skeletal
67 elements¹⁶. Different methods are applied to juveniles and adults^{17,18}. Limitations of these
68 methods include a high inter-observer variability¹⁵, inter- and intra-population variability with
69 increasing AAD¹⁹, lack of consensus regarding the evaluation of the errors²⁰, poor precision in
70 adult aging in comparison with juvenile and adolescent aging⁴, and the requirement for
71 remains to be as complete as possible²⁰.

72 In recent years, bone proteomics methods have been demonstrated to be highly
73 promising for the development of precise, accurate and objective PMI and AAD estimation
74 methods, requiring only small samples of bone. Proteins are relatively stable in bone, and
75 have been successfully extracted from archaeological²¹⁻²⁵ and paleontological specimens²⁶⁻
76 ²⁹, making them a promising target for forensic applications³⁰. Studies conducted using animal
77 models (e.g., *Sus scrofa* and *Bos bovid*) focused on inter- and intra-individual comparisons and
78 monitored changes in the bone proteomes associated with progressing decomposition
79 stages. These studies revealed inter-skeletal proteomic variability³¹, and identified potential
80 biomarkers for AAD³¹ and PMI estimations³². In addition, burial environment was found to
81 affect the proteome recovered from archaeological specimens³³. However, the development
82 of bone proteomics methods for forensics remains impeded by the fact that it is unknown
83 how representative animal models are for human specimens. Moreover, it is largely unknown
84 how taphonomic processes and inter-individual variation (both *in vivo* and at the time of
85 death), including underlying health conditions, affect the survival and extraction of bone
86 protein profiles in humans.

87 A recent study conducted on human bones collected from a cemetery in Southeast
88 Spain, provided promising new insights on the estimation of broad PMI ranges (5-20 years) in
89 humans using protein biomarkers in proximal femoral bone³⁴. The study identified 32 proteins
90 which could be used in conjunction to discriminate between PMIs greater or smaller than 12
91 years³⁴. The sampled individuals were subjected to similar taphonomic conditions, and PMIs
92 were greater than 7 years in all but one case. While the study was conducted on a relatively
93 large sample (n=40), inter-individual and inter-skeletal comparison of bone protein profiles at

94 different stages of decomposition of the body were not possible as only one skeletal element
 95 was available per individual. For the further development, and ultimately validation, of
 96 forensic proteomics to estimate PMI, the study of changes in human bone protein profiles
 97 from the fresh stage of decomposition to the skeletonized stage is crucial.

98 In this study, we aimed to investigate the effects of taphonomy and biological
 99 variation on the recovery and variability of the human bone proteome, and evaluate potential
 100 avenues to develop a broadly applicable, standardized method of PMI and AAD estimation in
 101 human remains in advanced state of decomposition. The proteomes of anterior midshaft tibia
 102 and iliac crest samples from four body donors of known AAD (two buried and two placed in
 103 an open pit), taken shortly after death and upon complete skeletonization of the body, were
 104 analysed to investigate 1) whether the previously identified potential biomarkers for PMI and
 105 AAD are applicable to human bones with lower PMIs, 2) whether additional potential
 106 biomarkers for PMI/AAD estimation could be identified, 3) whether the human bone
 107 proteome is subject to inter-skeletal and inter-individual variability, and 4) the role
 108 depositional environment, season and taphonomy play in bone proteome recovery.

109

110 2. Results

111

112 Proteomic data

113 The proteome of both the midshaft tibia and the iliac crest of four human body donors
 114 sampled at “fresh” (PMI = 2-10 days) and at “skeletonized” (i.e., when bodies did not have
 115 any adhering/desiccated soft tissue) stages of decomposition (PMI variable, between ~5200
 116 and ~17800 ADD, depending on the season of placement, see Table 1), was analysed. Three
 117 replicate extractions were taken from each bone, totaling 48 proteomic analyses. After
 118 refining the Progenesis results based on the number of unique peptides and on the ion score
 119 (see Methods section), 133 quantifiable proteins were identified (Supplementary Data 1). The
 120 protein interaction network (Fig. 1) showed a significant enrichment of interactions (PPI
 121 enrichment $p < 1.0 \times 10^{-16}$) and functional enrichments of specific GO terms for biological
 122 processes, cellular components and molecular functions (Supplementary Data 2).

123


124 **Table 1.** Biological and bone sample data.

Donor	Age at death (years)	Sex	Sample ID	Lab ID (three replicates)	Depositional context	Placement date (dd-mm-yy)	Collection date (dd-mm-yy)	T in days*
1	91	F	B1A-2-iliac	NP1-2-3	Open pit		28-04-2015	-1
1	91	F	B1A-2-tibia	NP4-5-6	Open pit	28-04-2015	28-04-2015	-1
1	91	F	B1C-2-iliac	NP7-8-9	Open pit		03-12-2015	219
1	91	F	B1C-2-tibia	NP10-11-12	Open pit		03-12-2015	219
2	67	F	B2A-2-iliac	NP13-14-15	Burial		07-05-2015	0
2	67	F	B2A-2-tibia	NP16-17-18	Burial	07-05-2015	07-05-2015	0
2	67	F	B2C-2-iliac	NP19-21-21	Burial		17-08-2017	834
2	67	F	B2C-2-tibia	NP22-23-24	Burial		17-08-2017	834
3	61	F	B3A-2-iliac	NP25-26-27	Burial	24-06-2015	24-06-2015	0
3	61	F	B3A-2-tibia	NP28-29-30	Burial		24-06-2015	0

3	61	F	B3C-2-iliac	NP31-32-33	Burial	21-08-2017	790
3	61	F	B3C-2-tibia	NP34-35-36	Burial	21-08-2017	790
4	77	F	B4A-2-iliac	NP37-38-39	Open pit	19-10-2015	-1
4	77	F	B4A-2-tibia	NP40-41-42	Open pit	19-10-2015	-1
4	77	F	B4C-2-iliac	NP43-44-45	Open pit	19-10-2015	872
4	77	F	B4C-2-tibia	NP46-47-48	Open pit	09-03-2018	872

125 * T0 = day of burial/placement.

126



127

128

129

130

131

132

133

134

135

Human proteomic inter-skeletal and inter-individual variability

136

137

138

139


140

141


Fresh samples were found to have a significantly greater protein diversity than skeletonized samples (Fig. 2A), and fresh iliac samples were the richest samples analysed, both in terms of proteome diversity (average of 55 different proteins in iliac fresh samples versus 27 for tibia fresh, 15 for iliac skeletonized and 23 for tibia skeletonized, see Supplementary Data 3 for details) and protein relative abundances (Supplementary Data 4). In fact, among the 116 proteins with significantly different relative abundances between the various bone and

142 sampling types (e.g., fresh vs. skeletonized), 105 (90.5%) were more abundant in the fresh
143 iliac samples, eight (6.9%) in the skeletonized tibia samples, two (1.7%) in the fresh tibia
144 samples and one (0.9%) in the skeletonized iliac samples (Supplementary Data 4). When
145 comparing iliac fresh and skeletonized samples, 96 proteins had a significantly different
146 abundance in the two groups, all of which were more abundant in fresh than in skeletonized
147 samples (Fig. 3 and Supplementary Data 5). Comparison of the fresh and skeletonized tibia
148 samples revealed 23 proteins with significantly different expression in the two groups, of
149 which 19 were more abundant in the fresh samples and four in the skeletonized samples (Fig.
150 3 and Supplementary Data 5).

151 Comparison of inter-individual proteome variability of fresh bones only (to exclude
152 any difference caused by taphonomic phenomena) showed that samples collected from D2
153 had a richer proteome variety (average number 62 for D2 versus 34, 33 and 35 for D1, D3 and
154 D4 respectively), although this difference was not statistically significant (Fig. 2B). Comparing
155 inter-individual relative protein abundances (Supplementary Data 6), we found 41 proteins
156 that showed differences in their relative abundances among the donors. Of these, 36 (87.8%)
157 were more abundant in D2, two each in D1 and D4 (4.9%) and one in D3 (2.4%).




158 **Figure 2. A)** Number of proteins extracted from each sample. Samples were grouped according to bone
159 type. All bone types were significantly different each other (post-hoc pairwise Wilcoxon-test with
160 corrections for multiple testing, p value < 0.0002 = ****, p value < 0.002 = ***, p value < 0.01 = *). Outliers
161 are represented as pointed-dots in the plot (two outliers identified here, one for iliac fresh (sample NP14,
162 see Table 1 for details) and one for tibia iliac skeletonized group (sample NP45, see Table 1 for details)). **B)**
163 Number of proteins extracted from fresh samples. Samples were grouped according to the donor. None of
164 the donors resulted in being significantly different each other (post-hoc pairwise Wilcoxon-test with
165 corrections for multiple testing, p value > 0.05).
166
167



168
169 **Figure 3.** Venn diagram to represent STRING protein networks of proteins significantly more abundant in
170 fresh iliac samples (left) and fresh tibia samples (right) than in their skeletonized counterparts. Proteins
171 shared between the two categories are represented in the middle. Immunoglobulin proteins (gene names
172 IGHA1, IGHG2, IGHG3, IGKC, IGLC2) were not found with STRING and are not represented in the figure. In
173 the iliac crest category, red cluster represents plasma proteins, yellow cluster represent collagens and
174 bone-related proteins, and green cluster ubiquitous proteins. No obvious clusters were identified for the
175 shared proteins and for the ones belonging to the tibia category.

176 177 **The influence of environment on bone proteome**

178 Comparison of samples from the different depositional environments (open pits vs. shallow
179 burials) showed no significant differences in the number of extracted proteins ($p=0.3$; Fig. 4A).
180 Comparison of the relative protein abundances in these two groups revealed only four
181 proteins with a different mean abundance for the two environments (three proteins more
182 abundant in shallow burials, one protein more abundant in open pit placements,
183 Supplementary Data 7). A test for association between the number of recovered proteins and
184 the season of placement found no significant differences ($p=0.4$; Fig. 4B).




185
186 **Figure 4.** Number of proteins extracted from skeletonized samples, grouped by **A)** depositional
187 environment or **B)** placement season. No significant differences were detected (Wilcoxon and Kruskal-
188 Wallis p value > 0.05).

189

190 **Potential proteomic biomarkers for human PMI estimation**

191 No association was found between the number of extracted proteins and the PMI of the
192 samples. However, significant decreases in the abundance of collagen alpha-1(III) chain
193 (CO3A1; $p=0.0041$), complement C9 (CO9; $p=1.9e-06$), collagen alpha-2(XI) chain (COBA2;
194 0.00055), matrix Gla protein (MGP; $p=5.3e-05$), decorin (PGS2; $p=0.045$) and transthyretin
195 (TTHY; $p=0.035$) in iliac crest (Fig. 5A-F) and of complement C3 (CO3) in tibia ($p=0.023$; Fig.
196 5G) were observed when comparing the protein abundances of the skeletonized samples.




197
198 **Figure 5.** Abundance of **A)** CO3A1, **B)** CO9, **C)** COBA2, **D)** MGP, **E)** PGS2, **F)** TTHY protein in iliac crest
199 skeletonized samples and of **G)** CO3 in tibia skeletonised samples with increasing PMIs. Groups are
200 significantly different each other (ANOVA p value < 0.05).

201 **Potential proteomic biomarkers for human AAD estimation**

202 The relative abundance of fetuin-A was found to be negatively associated with AAD in fresh
203 tibia ($p=0.033$) and in skeletonized iliac samples ($p=0.013$). Skeletonized tibia samples showed
204 lower levels for the oldest donor and higher levels for the other, but no significant trend was
205 found ($p=0.34$). Iliac fresh samples showed similar levels in D2 and D3 and lower values for
206 D1 and D4 ($p=0.34$; Fig. 6A-D).

207 Significant differences in albumin abundance were found between different donors
208 for both fresh ($p=0.011$) and skeletonized ($p=0.016$) iliac samples (Fig. 6E-H). In particular,
209 fresh iliac samples showed a positive association with AAD, while fresh and skeletonized tibia
210 samples both showed a negative relationship with AAD, although these results were not
211 significant (Fig. 6; ($p=0.12$ and 0.35 , respectively).

212 Additionally, a significant increase in the abundance of olfactomedin-like protein 3
213 (OLF3) was observed in skeletonized iliac samples with increasing AAD (Fig. 6G; $p=0.031$).



214
 215 **Figure 6.** Relative abundance of fetuin-A in **A)** fresh tibia, **B)** skeletonized tibia, **C)** fresh iliac crest and **D)**
 216 skeletonized iliac crest samples, of albumin in **E)** fresh tibia, **F)** skeletonized tibia, **G)** fresh iliac crest and **H)**
 217 skeletonized iliac crest samples and of **I)** orfantomedin like-3 in skeletonized iliac crest samples, arranged
 218 by the chronological age of the donors. ANOVA p value was reported for each plot. Only A), D), G), H) and
 219 I) resulted in being statistically significant.

220

221 3. Discussion

222 In this study, we identified specific proteins that significantly decreased in abundance with
 223 increasing PMI: complement C3 for tibia and collagen alpha-1(III) chain, complement C9,
 224 collagen alpha-2(XI) chain, matrix Gla protein, decorin and transthyretin for iliac crest. Four
 225 of the identified proteins are classified as bone structural/functional proteins (CO3A1, COBA2,
 226 MGP and PGS2) and three are plasma proteins (CO3, CO9 and TTHY). Previous work,
 227 conducted on animal proxies (pigs) left to decompose for a maximum of six months³⁵,
 228 revealed a similar trend of consistently decreasing protein abundances over time, but for
 229 different proteins: haemoglobins, transferrins, triosephosphate isomerase, collagen alpha-
 230 2(V) chain and albumin. Both studies showed a reduction in the abundances of plasma and
 231 ubiquitous proteins, but reduction in bone structural/functional proteins was observed only
 232 in the current study. It is known that certain mineral-binding proteins (including structural
 233 ones) are susceptible to taphonomic processes of decay and diagenesis with prolonged
 234 PMIs³⁶. The difference in duration and in the depositional environment and local climate
 235 between the pig study and the present study, and the resulting longer exposure to
 236 taphonomic processes, could therefore explain the different trends in mineral-binding
 237 protein abundances reduction that we observed. Analysis of human femoral bones from a
 238 cemetery context by Prieto-Bonete and colleagues³⁴ also revealed a distinct reduction in the
 239 amount of structural and functional proteins in the highest PMI samples (13-20 years). Among
 240 the list of proteins identified in their study as biomarkers for prolonged PMIs, COBA2 was the
 241 only one that was also found here showing a similar inverse association with increasing PMIs

242 in the present study. Overall, these findings suggest that COBA2 could be a good candidate
243 for PMI estimation of human skeletonized remains, due to its durability over time and under
244 different taphonomical conditions.

245 Our study found that the abundance of OLFL3, an osteoblast secreted extracellular
246 matrix glycoprotein³⁷, was positively associated with AAD in skeletonized iliac samples,
247 adding a previously unreported protein to the list of potential biomarkers for AAD. Previous
248 studies on animal and archaeological human bones identified a negative correlation between
249 serum fetuin-A and AAD^{21,38}, and proposed fetuin-A as potential biomarker for AAD³⁹. The
250 present study identified a similar negative relationship in fresh tibia and skeletonized iliac
251 crest samples, but not in fresh iliac and skeletonized tibia. In addition to fetuin-A, several
252 studies showed that serum albumin concentration is negatively correlated with AAD^{40,41}. The
253 present study found a non-significant negative correlation both in fresh and skeletonized tibia
254 samples, but an opposite and significant trend in fresh iliac samples. Considering the relatively
255 small sample size, it is difficult to interpret these findings. Based on observations of bone
256 properties (hardness, weight, coloration) during sampling of the bone we postulate that inter-
257 individual differences in bone mineral density (BMD; discussed below) as well as the inter-
258 skeletal differences in BMD (i.e., iliac bone is less densely mineralised than tibia), may have
259 affected these results. BMD is known to vary between different parts of the skeleton⁴². With
260 regards to inter-individual BMD differences it is important to note that our observations do
261 not constitute quantified measurements of mass-to-volume ratio, but are observations of the
262 physical properties of bones during drilling. Our study used three biological replicates,
263 however using a larger number in future studies might clarify whether fetuin-A and albumin
264 are consistently negatively correlated with AAD in humans in different bone types and
265 therefore can be used as a biomarker for AAD.

266
267 The potential effects of BMD on the recovered protein profile are suggested by the
268 results of inter-individual comparisons. Fresh tibia samples from all four donors showed
269 greater inter-individual reproducibility than fresh iliac samples. Fresh iliac samples from D2
270 showed significantly increased protein variety and abundances. Observations during drilling
271 of the bone samples suggest that BMD in D2 may be relatively high. BMD could theoretically
272 affect the variety and abundance of specific non-collagenous proteins that can either bind the
273 calcium ions or the collagen in the mineral matrix²⁶, thereby affecting the overall protein
274 profile. In addition to non-collagenous proteins, many ubiquitous and plasma proteins were
275 identified predominantly in fresh iliac samples, particularly from D2. Due to the greater blood
276 irrigation of the iliac crest in comparison with the midshaft tibia, a greater variety of these
277 types of proteins could be expected in fresh iliac samples (discussed below). In addition to
278 the observations of bone hardness in D2 made during drilling, the medical information
279 available for D2 indicates certain conditions and treatments received in the years prior to
280 death that are associated with changes in BMD, including chemotherapy treatment for
281 cancer, prolonged consumption of calcium lactate⁴³ and possible use of probiotics as adjuvant
282 during cancer treatment^{44,45}.

283 The likely greater BMD of D2 may have allowed for a stronger in vivo embedding of
284 these proteins within the mineral matrix, resulting in the greater proteomic variety observed
285 here. This effect of BMD on protein linkage in bones would be analogous to positive
286 relationships observed between organic matter content and soil density, which is often a
287 function of clay content. In fact, clay particles tend to carry a negative charge to bind with

288 nutrient cations such as calcium and potassium, and these bonds can protect proteins from
289 decomposition and even from extreme environmental conditions such as autoclaving⁴⁶.

290 The results of inter-skeletal comparisons of the skeletonized samples suggest that
291 both BMD and taphonomic processes affected the preservation and extraction of the bone
292 proteome. The highly irrorated and less densely mineralized fresh iliac samples yielded
293 greater variety and abundance of proteins, including those expressed specifically in plasma.
294 The proteomes recovered from skeletonized iliac samples demonstrated significant protein
295 decay occurred in this bone. The denser and less irrorated fresh tibia samples yielded lower
296 protein variety and abundances by comparison to the fresh iliac samples. Comparison of the
297 fresh tibia samples with the skeletonized tibia samples showed protein decay also occurred
298 in this bone, but not to the degree observed in the iliac crest. These results suggest that
299 differences in BMD and blood irroration between the iliac crest and the midshaft anterior
300 tibia affected both the successful extraction of proteins from fresh samples, as well as the
301 preservation in and extraction of proteins from skeletonized samples. Taphonomic processes
302 of decomposition are known to affect BMD in humans, and can differentially affect skeletal
303 elements^{47,48}. Higher porosity of the iliac exposed this bone to significant deterioration as a
304 result of taphonomic processes over time, resulting in the reduced inter-individual
305 differences observed in the skeletonized iliac samples. Protein extraction from the dense and
306 poorly irrorated fresh anterior midshaft tibia was less successful – at least with the mild
307 extraction protocol used here – but the structure of this bone led to less taphonomic
308 deterioration over time.

309 The current results likely indicate the effects of the decomposer community and
310 physicochemical environment on the decomposition of human remains. A less dense matrix
311 would facilitate leaching while promoting the movement of decomposer microbes
312 throughout the bone. Microbial induced bioerosion, which is characterized by the chemical
313 dissolution of mineral components of bone followed by the microbial enzymatic attack of
314 organic components of bone, is thought to be one of the main causes of bone diagenesis⁴⁹.
315 The movement of decomposer microbes might be restricted to the external surfaces of more
316 densely mineralised bone. The effect of the decomposer microbial community may be further
317 influenced by the location of the bones. The iliac crest might be subject to greater
318 decomposition because it is located closer to the trunk, which contains a significant amount
319 of moisture and a large gastrointestinal microbial community that is known to translocate
320 during decomposition^{50,51}. The iliac crest, therefore, is likely located in a microhabitat that is
321 favourable for decomposition. In contrast, the tibia is located further from the trunk in limbs
322 that often desiccate during decomposition. The body position during decomposition of the
323 four donors was flexed, and allowed the anterior tibiae to remain elevated above
324 decomposition fluids excreted from the trunk. Desiccation of the soft tissues around the
325 anterior tibiae was observed early on during decomposition of both open pit placements.
326 Desiccated, densely mineralised bone is unlikely to be favourable for decomposition.

327 Comparison of samples from open pit placements with samples from burials, as well
328 as comparison of season of placement in this study found no significant differences. While
329 archaeological remains have revealed differences in protein recovery related to depositional
330 environment^{23,26,33}, it is possible that due to the relatively short duration of this experiment
331 such environmental effects were not measurable in this study. It is also possible that the two
332 depositional environments did not produce distinct enough conditions (Supplementary Table
333 1) to cause noticeable differences in the preservation of the biomolecules.

334 The preliminary indications from this study support previous findings that specific
335 proteins decay at different rates, strengthening the potential for developing bone proteomics
336 PMI estimation methods. COBA2 appears to be a good candidate for PMI estimation of
337 skeletonized remains, together with CO3A1, PGS2 and MGP. The blood proteins CO3, CO9
338 and TTHY may be good candidates for shorter PMI estimation (i.e., before the complete
339 degradation of blood proteins). Our study only partially supported previous studies
340 identifying fetuin-A and albumin as potential biomarkers for AAD estimation, and additionally
341 found OLFL3 being positively correlated with AAD. At the same time, our findings suggest that
342 taphonomic (e.g., microbial bioerosion) and biological (e.g., variation in BMD) variables play
343 a significant role in both the survival and extraction rate of proteins, due to their effects on
344 the protective mineral matrix.

345 While the sample size is relatively small, the findings point toward potentially significant
346 effects of inter-individual variation associated with health conditions, medical treatment, and
347 possibly food and supplement intake. The results of both inter-individual and inter-skeletal
348 comparisons in our study suggest that BMD may be an important variable affecting the
349 survival and extraction of proteins in the bone mineral matrix. Higher BMD may promote
350 attachment of a greater abundance and variety of mineral binding proteins, and in highly
351 irrorated bones may additionally help to preserve more plasma proteins within the mineral
352 matrix. Inter-skeletal differences in BMD appear to lead to distinct differences in the variety
353 and abundance of preserved (and extracted) proteins. The attachment of proteins within a
354 more densely mineralized bone matrix may protect them during microbial bioerosion. Based
355 on these indications, we recommend including standard measurement of BMD and targeting
356 a combination of different biomarkers (i.e., abundances of selected plasma proteins and
357 bone-specific proteins) in future work. Overall, these results emphasize the limitations of
358 developing methods and models based on animal proxies, since farmed animals rarely show
359 the degree of inter-individual dietary and activity related variation that humans do, and BMD
360 and degree of irroration of bones differ between species⁵². Moreover, these results
361 emphasize the importance of conducting replication studies in larger human samples
362 representing a broader range of PMIs and AAD, as well as sampling different bones, to better
363 understand how different types of proteins and different parts of the human skeleton are
364 affected by inter-individual variation and taphonomic processes. Finally, preliminary
365 evaluation of the inter-skeletal differences we observed suggests that for future development
366 of proteomics PMI estimation methods, the iliac crest bone may be a more suitable sampling
367 target for relatively fresh remains of forensic interest and for archaeological studies
368 specifically targeting the serum-proteins, due to the presence of greater protein variety of
369 bone-marrow proteins. Specific burial conditions, such as dry burial environments, anaerobic
370 environments, and certain post-mortem treatments of the body (such as embalming
371 procedures) can limit the amount of bone diagenesis^{53,54} thereby promoting the survival of
372 bone proteins across archaeological timeframes. In such circumstances the iliac crest may
373 provide better results than the tibia to detect pathologies and infections associated with the
374 bone marrow. The midshaft tibia may be a more suitable sampling target for skeletonized
375 remains or those in a state of advanced decomposition of forensic interest, due to the better
376 survival of collagen and mineral-related proteins that could be ultimately used for developing
377 new biomolecular methods for PMI/AAD estimation.

378

379 **4. Methods**

380

381 Body donations

382 The body donations of four females aged between 61 and 91 years old were placed unclothed
383 to decompose at the Forensic Anthropology Research Facility (FARF), the outdoor human
384 decomposition facility associated with the Forensic Anthropology Center at Texas State
385 University (FACTS), between April 2015 and March 2018. While the targeted bone proteins in
386 this study are not thought to differ between males and females, only post-menopausal female
387 individuals were included, in order to exclude biological sex and major hormonal differences
388 as a potential variable from the study. Two body donations (D2 and D3) were buried with soil
389 in shallow hand dug pits. Two body donations (D1 and D4) were placed in pits of similar
390 dimensions, which remained open throughout the experiment. Open pits were covered with
391 a metal cages to protect the remains from large scavengers. The sample size in this study
392 reflects general trends in human decomposition research, in which larger samples – like those
393 used in clinical studies – can be difficult to obtain for practical, logistical and ethical reasons.
394 While animal analogues such as pigs can be used to alleviate some limitations associated with
395 small sample sizes, the study of human cadavers is important due to biological differences
396 between humans and pigs, including anatomical differences in the digestive vasculature, and
397 molecular differences in adipose tissue⁵⁵.

398 Data on body decomposition and weather were collected throughout the experiment
399 and can be found in Supplementary Table 1. Additional information on FARF's environment
400 can be found in the Supplementary Information. Gross decomposition was quantified using
401 the total body score (TBS) method following Megyesi et al.⁵⁶. Accumulated degree-days (ADD)
402 were calculated using temperature data recorded on the facility.

403

404 Bone sample collection

405 Bone samples (ca. 1 x 1 cm) of the anterior midshaft tibia and iliac crest (left) were collected
406 prior to placement of the fresh body outside and upon retrieval of the completely
407 skeletonized remains (right). The total 16 bone samples were stored in sterile plastic bags,
408 and immediately transferred to a lockable freezer at -80°C. Samples were shipped overnight
409 on dry ice to the Forensic Science Unit at Northumbria University, U.K. Upon arrival,
410 the samples were immediately transferred to a lockable freezer at -18°C, adhering to the U.K.
411 Human Tissue Act under the license number 12495. The experiment was reviewed and
412 approved by the ethics committee at Northumbria University, with the reference code 11623.
413 All biological and bone sample data are provided in Table 1. Observations on bone condition
414 (density and colour) during sampling can be found in the Supplementary Information.

415

416 Sub-sampling and sample preparation

417 The 16 samples were defrosted prior to their analysis, then cleaned in deionized water for
418 three hours at room temperature, exchanging the water three times, once every hour. They
419 were then dried in a fume cupboard at room temperature until completely dry. Bone samples
420 were then secured in a table clamp for the sampling. Contamination between samples was
421 prevented by using a double layer of aluminium foil within the clamp (in contact with the
422 bone) and by using new foil double layers for each piece of bone sampled. The clamp was also
423 cleaned in between each sampling step using 50% sodium hypochlorite (Sigma-Aldrich, U.K.),
424 to further prevent contamination issues. Once the bone was secured in the clamp, Dentist's
425 Protaper Universal Hand Files (Henry Schein Minerva Dental, U.K.) were used to hand-drill
426 ~25mg of fine bone powder for three times (i.e. three samplings were performed on the same
427 bone fragment), in order to obtain three replicates for each of the bones analysed. By

428 sampling in different locations close together on the same bone, we obtained multiple
429 biological samples. Since it is known that bone proteins can vary throughout the human
430 skeleton and within individual bones, these biological replicates, in contrast to technical
431 replicates, allow us to assess the degree of intra-bone variability and to establish whether
432 inter-individual differences are greater than the intra-bone variability, as indicated in a
433 previous study using pigs as proxies³¹. Protaper files were changed between each sample, to
434 prevent contamination. When the bone samples were too porous to obtain a fine bone
435 powder (e.g., iliac crest samples), small bone fragments were cut using the Protaper files, and
436 ~25mg of bone fragments were collected for each of the three subsamples in order to have
437 three replicates.

438

439 Protein extraction

440 Overall, 48 samples were obtained from the 16 bone pieces, and were subjected to bone
441 protein extraction following the protocol of Procopio and Buckley⁵⁷.

442 Briefly, each sample was decalcified with 1 mL of 10 v/v% formic acid (Fisher Scientific, U.K.)
443 for 6 hours at 4 °C. After removing all the acid soluble fraction, the acid insoluble fraction was
444 incubated for 18 hours at 4°C with 500 µL of 6 M guanidine hydrochloride/100mM TRIS buffer
445 (pH 7.4, Sigma-Aldrich, U.K.). The buffer was exchanged into 100 µL of 50 mM ammonium
446 acetate (Scientific Laboratory Supplies, U.K.) with 10K molecular-weight cut off filters
447 (Vivaspin 500 polyethersulfone, 10kDa, Sartorius, Germany), and samples were then reduced
448 with 4.2 µL of 5 mM dithiothreitol (DTT) (Fluorochem, U.K.) for 40 min at room temperature
449 and alkylated with 16.8 µL of 15 mM iodoacetamide (Sigma-Aldrich, U.K.) for 45 min in the
450 dark at room temperature. Samples were then quenched with another 4.2 µL of 5 mM DTT,
451 then digested with 0.4 µg of trypsin (Promega, U.K.) for 5 hours at 37°C and finally frozen. By
452 adding 15 µL of 1 v/v% trifluoroacetic acid (TFA) (Fluorochem, U.K.), the digestion was
453 stopped and the samples were then desalted, concentrated and purified using OMIX C18
454 pipette tips (Agilent Technologies, U.S.A.) with 0.1 v/v% TFA as washing solution and 50 v/v%
455 acetonitrile (ACN) (Thermo Fisher Scientific, U.K.)/0.1 v/v% TFA as a conditioning solution.
456 Pipette tips were prepared with two volumes of 100 µL of 0.1 v/v% TFA and washed twice
457 with 100 µL of 50 v/v% ACN/0.1 v/v% TFA. The sample was then aspirated into the tip at least
458 ten times to efficiently bind peptides to the absorbent membrane. Finally, two washing steps
459 with 100µL of 0.1 v/v% TFA were performed, prior to peptides elution into 100 µL of 50 v/v%
460 ACN/0.1 v/v% TFA. Purified peptides were left in the fume cupboard at room temperature
461 with lids open to dry prior to their submission for LC-MS/MS analysis.

462

463 LC/MS-MS analysis

464 Samples resuspended in 5 v/v% ACN/0.1 v/v% TFA were analyzed by LC-MS/MS using an
465 Ultimate™ 3000 Rapid Separation LC (RSLC) nano LC system (Dionex Corporation, Sunnyvale,
466 CA, USA) coupled to a Q Exactive™ Plus Hybrid Quadrupole-Orbitrap Mass Spectrometer
467 (Thermo Fisher Scientific, Waltham, MA, U.S.A.). Peptides were separated on an EASY-Spray™
468 reverse phase LC Column (500 mm x 75 µm diameter (i.d.), 2 µm, Thermo Fisher Scientific,
469 Waltham, MA, USA) using a gradient from 96 v/v% A (0.1 v/v% FA in 5 v/v% ACN) and 4 v/v%
470 B (0.1 v/v% FA in 95 v/v% ACN) to 8 v/v%, 30 v/v% and 50% B at 14, 50, and 60 min,
471 respectively, at a flow rate of 300 nL min⁻¹. Acclaim™ PepMap™ 100 C18 LC Column (5 mm x
472 0.3 mm i.d., 5 µm, 100 Å, Thermo Fisher Scientific) was used as trap column at a flow rate of
473 25 µL min⁻¹ kept at 45 °C. The LC separation was followed by a cleaning cycle with an
474 additional 15 min of column equilibration time. Then, peptide ions were analyzed in full scan

475 MS scanning mode at 35,000 MS resolution with an automatic gain control (AGC) of 1e6,
476 injection time of 200 ms and scan range of 375-1,400 m/z. The top ten most abundant ions
477 were selected for data-dependent MS/MS analysis with a normalized collision energy (NCE)
478 level of 30 performed at 17,500 MS resolution with an AGC of 1e5 and maximum injection
479 time of 100 ms. The isolation window was set to 2.0 m/z, with an underfilled ratio of 0.4%,
480 dynamic exclusion was employed; thus, one repeat scan (i.e., two MS/MS scans in total) was
481 acquired in a 45 s repeat duration with the precursor being excluded for the subsequent 45
482 s.

483

484 Data analysis and statistical analysis

485 Peptide mass spectra were then searched against the SwissProt_2019_11 database (selected
486 for Homo sapiens, unknown version, 20368 entries) using the Mascot search engine (version
487 2.5.1; www.matrixscience.com) for matches to primary protein sequences. This search
488 included the fixed carbamidomethyl modification of cysteine as it results from addition of DTT
489 to proteins. Deamidation (asparagine and glutamine) and oxidation (lysine, methionine and
490 proline) were considered as variable modifications. The enzyme was set to trypsin with a
491 maximum of two missed cleavages allowed. Mass tolerances for precursor and fragmented
492 ions were set at 5 ppm and 0.5 Da, respectively. It was assumed that all spectra hold either
493 2+ or 3+ charged precursors. Scaffold (version Scaffold_4.10.0, Proteome Software Inc.,
494 Portland, OR) was used to validate MS/MS based peptide and protein identifications. Peptide
495 identifications were accepted if they could be established at greater than 95.0% probability
496 to maximise the reliability of the identifications. Peptide Probabilities from Mascot were
497 assigned by the Scaffold Local FDR algorithm and by the Peptide Prophet algorithm⁵⁸ with
498 Scaffold delta-mass correction. Protein identifications were accepted if they could be
499 established at greater than 90.0% probability and contained at least 2 identified peptides, in
500 order to filter for the most accurate matches. This resulted in having a calculated decoy FRD
501 of 0.06% for peptides and 1.9% for proteins. Protein probabilities were assigned by the
502 Protein Prophet algorithm⁵⁹. Proteins that contained similar peptides and could not be
503 differentiated based on MS/MS analysis alone were grouped to satisfy the principles of
504 parsimony. Proteins sharing significant peptide evidence were grouped into clusters.
505 Progenesis Qi for Proteomics (version 4.1; Nonlinear Dynamics, Newcastle, U.K.) was used to
506 perform relative quantitation calculations using the recorded ion intensities (area under the
507 curve, AUC) and averaging the N most abundant peptides for each protein (Hi-N method,
508 where N=3) and protein and post-translational modifications identifications. In order to
509 increase the reliability of the matches, peptide ions with a score of <28, which indicates
510 identity or extensive homology ($p < 0.05$), were excluded from the analysis based on the
511 Mascot evaluation of the peptide score distribution for the searched .mgf file originating from
512 Progenesis (combining all the samples in a single experiment). To further improve the
513 reliability of the findings we implemented an additional level of filtering, excluding proteins
514 with a peptide count of <2. Samples were grouped together using the between-subject design
515 scheme in Progenesis, in order to compare selected group of samples (e.g. skeletonised
516 versus fresh bones) and to calculate ANOVA p-values and maximum fold changes accordingly.
517 The use of three biological replicates per targeted bone sample provided a sufficiently large
518 dataset for comparative analysis using non-parametric statistical methods such as ANOVA,
519 posthoc pairwise comparison, and Kruskal-Wallis tests. To identify proteins of interest,
520 proteins were flagged up in order to highlight the ones that had an ANOVA p-value ≤ 0.05 and
521 a maximum fold change ≥ 2 . Common contaminants such as keratins were excluded from the

522 interpretation of the results. Plots were done using R studio version 1.2.5033 with packages
523 dplyr, ggplot2, ggpubr and patchwork packages. When plotting boxplots, for data following a
524 normal distribution student's t test and one-way ANOVA and post-hoc pairwise comparisons
525 were used to test mean differences, otherwise Wilcoxon rank sum test and Kruskal Wallis test
526 with post-hoc pairwise comparisons were used. STRING software version 11.0 was used to
527 visualize functional links between the extracted proteins⁶⁰. The confidence score required for
528 showing interactions was set to "high = 0.700". MCL clustering method was used to identify
529 the clusters, with inflation parameter = 1.5.

530

531 **Data availability**

532 The mass spectrometry proteomics data have been deposited to the ProteomeXchange
533 Consortium via the PRIDE⁶¹ partner repository with the dataset identifier PXD019693 and
534 10.6019/PXD019693".

535

536 **Reviewer account details:**

537 **Username: reviewer46348@ebi.ac.uk**

538 **Password: QkJJd6VZ**

539

540

541 **Acknowledgements**

542 The authors would like to acknowledge the Royal Society for funding a Research Grant (N.P.)
543 under grant RGS/R1/191371, the UKRI for funding part of the research through a Fellowship
544 (N.P.) under grant MR/S032878/1, as well as the European Research Council for funding part
545 of the research under grant 319209 and the Leiden University Fund for funding under Byvanck
546 grant 5604/30-4-2015/Byvanck. Dr. William Cheung at the NUOmic Facility is acknowledged
547 for conducting the LC-MS/MS runs. The authors also gratefully acknowledge the donors and
548 their next of kin for allowing the use of donated bodies to perform this research.

549

550 **Author Contributions**

551 H.L.M. and N.P. conceived the study and wrote the paper. Text editing by all co-authors. N.P.
552 was the lead on all proteomics experiments. H.L.M. was the lead on the human
553 decomposition experiments and bone sampling at FACTS. D.W. was the lead on
554 environmental data collection at FARF. H.L.M., N.P. and D.C. contributed to the interpretation
555 of the results. N.P. and S.S. executed protein extraction, N.P. and E.S. conducted data analysis
556 and F.S. and H.M. contributed to data analysis and to the creation of graphical outputs tables
557 and supplementary data.

558

559 **Competing interests**

560 The authors declare there are no financial or non-financial competing interests.

561

562 **References**

- 563 1. White, B. United Kingdom. in *The Routledge handbook of archaeological human*
564 *remains and legislation: An international guide to laws and practice in the excavation*
565 *and treatment of archaeological human remains* (eds. Grant, N. M. & Fibiger, L.) 205–
566 206 (Taylor & Francis, 2011).
- 567 2. Creamer, J. I. & Buck, A. M. The assaying of haemoglobin using luminol
568 chemiluminescence and its application to the dating of human skeletal remains.

- 569 *Luminescence* **24**, 311–316 (2009).
- 570 3. Christensen, A. M., Passalacqua, N. V & Bartelink, E. J. *Forensic anthropology: current*
571 *methods and practice*. (Elsevier Inc., 2019).
- 572 4. Ritz-Timme, S. *et al.* Age estimation: the state of the art in relation to the specific
573 demands of forensic practise. *Int. J. Legal Med.* **113**, 129–136 (2000).
- 574 5. Junod, C. A. & Pokines, J. T. Subaerial Weathering. in *Manual of Forensic Taphonomy*
575 (eds. Pokines, J. T. & Symes, S. A.) 287–314 (CRC Press, 2013).
- 576 6. Wilson-Taylor, R. J. & Dautartas, A. M. Time since death estimation and bone
577 weathering, the post mortem interval. in *Forensic anthropology: a comprehensive*
578 *introduction* 273–313 (CRC Press Boca Raton, 2017).
- 579 7. Huculak, M. A. & Rogers, T. L. Reconstructing the sequence of events surrounding
580 body disposition based on color staining of bone. *J. Forensic Sci.* **54**, 979–984 (2009).
- 581 8. Berg, S. Methods in Forensic Science. in *Interscience* vol. 2 (Wiley and Sons New York,
582 1963).
- 583 9. Wescott, D. J. Recent advances in forensic anthropology: decomposition research.
584 *Forensic Sci. Res.* **3**, 278–293 (2018).
- 585 10. Blau, S. & Ubelaker, D. H. *Handbook of forensic anthropology and archaeology*. vol.
586 534 (Left Coast Press Walnut Creek, 2009).
- 587 11. Kanz, F., Reiter, C. & Risser, D. U. Citrate content of bone for time since death
588 estimation: results from burials with different physical characteristics and known
589 PMI. *J. Forensic Sci.* **59**, 613–620 (2014).
- 590 12. Pérez-Martínez, C., Pérez-Cárceles, M. D., Legaz, I., Prieto-Bonete, G. & Luna, A.
591 Quantification of nitrogenous bases, DNA and Collagen type I for the estimation of
592 the postmortem interval in bone remains. *Forensic Sci. Int.* **281**, 106–112 (2017).
- 593 13. Sarabia, J., Pérez-Martínez, C., del Rincón, J. P. H. & Luna, A. Study of
594 chemiluminescence measured by luminometry and its application in the estimation of
595 postmortem interval of bone remains. *Leg. Med.* **33**, 32–35 (2018).
- 596 14. Brown, M. A. *et al.* Citrate Content of Bone as a Measure of Postmortem Interval: An
597 External Validation Study. *J. Forensic Sci.* **63**, 1479–1485 (2018).
- 598 15. Priya, E. Methods of skeletal age estimation used by forensic anthropologists in
599 adults: a review. *Forensic Res Criminol Int J* **4**, 104 (2017).
- 600 16. Franklin, D. Forensic age estimation in human skeletal remains: current concepts and
601 future directions. *Leg. Med.* **12**, 1–7 (2010).
- 602 17. Lewis, M. E. & Flavel, A. Age assessment of child skeletal remains in forensic contexts.
603 in *Forensic Anthropology and Medicine* (eds. Schmitt, A., Cunha, E. & Pinheiro, J.)
604 243–257 (Humana Press, 2006).
- 605 18. Baccino, E. & Schmitt, A. Determination of adult age at death in the forensic context.
606 in *Forensic Anthropology and Medicine* (eds. Schmitt, A., Cunha, E. & Pinheiro, J.)
607 259–280 (Humana Press, 2006).
- 608 19. Schmitt, A., Murail, P., Cunha, E. & Rougé, D. Variability of the pattern of aging on the
609 human skeleton: evidence from bone indicators and implications on age at death
610 estimation. *J. Forensic Sci.* **47**, 1203–1209 (2002).
- 611 20. Valsecchi, A., Olivares, J. I. & Mesejo, P. Age estimation in forensic anthropology:
612 methodological considerations about the validation studies of prediction models. *Int.*
613 *J. Legal Med.* **133**, 1915–1924 (2019).
- 614 21. Sawafuji, R. *et al.* Proteomic profiling of archaeological human bone. *R. Soc. Open Sci.*
615 **4**, 161004 (2017).

- 616 22. Devièse, T. *et al.* Direct dating of Neanderthal remains from the site of Vindija Cave
617 and implications for the Middle to Upper Paleolithic transition. *Proc. Natl. Acad. Sci.*
618 *U. S. A.* **114**, 10606–10611 (2017).
- 619 23. Wadsworth, C. *et al.* Comparing ancient DNA survival and proteome content in 69
620 archaeological cattle tooth and bone samples from multiple European sites. *J.*
621 *Proteomics* 1–8 (2017) doi:10.1016/j.jprot.2017.01.004.
- 622 24. Brown, S. *et al.* Identification of a new hominin bone from Denisova Cave, Siberia
623 using collagen fingerprinting and mitochondrial DNA analysis. *Sci. Rep.* **6**, 23559
624 (2016).
- 625 25. Mackie, M. *et al.* Preservation of the metaproteome: variability of protein
626 preservation in ancient dental calculus. *STAR Sci. Technol. Archaeol. Res.* **3**, 58–70
627 (2017).
- 628 26. Buckley, M. & Wadsworth, C. Proteome degradation in ancient bone: diagenesis and
629 phylogenetic potential. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **416**, 69–79 (2014).
- 630 27. Buckley, M. A molecular phylogeny of Plesiorcyteropus reassigns the extinct
631 mammalian order 'Bibymalagasias'. *PLoS One* **8**, e59614 (2013).
- 632 28. Dorado, G. *et al.* Genomics and proteomics in bioarchaeology. *Archaeobios* 4–17
633 (2013).
- 634 29. Harvey, V. L. *et al.* Preserved collagen reveals species identity in archaeological
635 marine turtle bones from Caribbean and Florida sites. *R. Soc. Open Sci.* **6**, 191137
636 (2019).
- 637 30. Buckley, M. Proteomics in the Analysis of Forensic, Archaeological, and
638 Paleontological Bone. in *Applications in Forensic Proteomics: Protein Identification*
639 *and Profiling* 125–141 (ACS Publications, 2019).
- 640 31. Procopio, N., Chamberlain, A. T. & Buckley, M. Intra- and Interskeletal Proteome
641 Variations in Fresh and Buried Bones. *J. Proteome Res.* **16**, (2017).
- 642 32. Procopio, N., Williams, A., Chamberlain, A. T. & Buckley, M. Forensic proteomics for
643 the evaluation of the post-mortem decay in bones. *J. Proteomics* **177**, 21–30 (2018).
- 644 33. Procopio, N., Chamberlain, A. T. & Buckley, M. Exploring Biological and Geological
645 Age-related Changes through Variations in Intra- and Intertooth Proteomes of
646 Ancient Dentine. *J. Proteome Res.* **17**, (2018).
- 647 34. Prieto-Bonete, G., Pérez-Cárceles, M. D., Maurandi-López, A., Pérez-Martínez, C. &
648 Luna, A. Association between protein profile and postmortem interval in human bone
649 remains. *J. Proteomics* **192**, 54–63 (2019).
- 650 35. Procopio, N., Williams, A., Chamberlain, A. T. & Buckley, M. Forensic proteomics for
651 the evaluation of the post-mortem decay in bones. *J. Proteomics* **177**, (2018).
- 652 36. Wadsworth, C. & Buckley, M. Proteome degradation in fossils: investigating the
653 longevity of protein survival in ancient bone. *Rapid Commun. Mass Spectrom.* **28**,
654 605–615 (2014).
- 655 37. Sanchez, C. *et al.* Comparison of secretome from osteoblasts derived from sclerotic
656 versus non-sclerotic subchondral bone in OA: A pilot study. *PLoS One* **13**, (2018).
- 657 38. Robinson, K. N. & Teran-Garcia, M. From infancy to aging: biological and behavioral
658 modifiers of fetuin-A. *Biochimie* **124**, 141–149 (2016).
- 659 39. Procopio, N., Chamberlain, A. T. & Buckley, M. Intra- and Interskeletal Proteome
660 Variations in Fresh and Buried Bones. *J. Proteome Res.* **16**, 2016–2029 (2017).
- 661 40. Veering, B. T., Burm, A. G., Souverijn, J. H., Serree, J. M. & Spierdijk, J. O. H. The effect
662 of age on serum concentrations of albumin and alpha 1-acid glycoprotein. *Br. J. Clin.*

- 663 *Pharmacol.* **29**, 201–206 (1990).
- 664 41. Cooper, J. K. & Gardner, C. Effect of aging on serum albumin. *J. Am. Geriatr. Soc.* **37**,
665 1039–1042 (1989).
- 666 42. Njeh, C. F. & Boivin, C. M. Variation in bone mineral density between different
667 anatomical sites in a normal local population. *Appl. Radiat. Isot.* **49**, 685–686 (1998).
- 668 43. Devine, A., Criddle, R. A., Dick, I. M., Kerr, D. A. & Prince, R. L. A longitudinal study of
669 the effect of sodium and calcium intakes on regional bone density in postmenopausal
670 women. *Am. J. Clin. Nutr.* **62**, 740–745 (1995).
- 671 44. Collins, F. L., Rios-Arce, N. D., Schepper, J. D., Parameswaran, N. & McCabe, L. R. The
672 potential of probiotics as a therapy for osteoporosis. *Bugs as Drugs Ther. Microbes*
673 *Prev. Treat. Dis.* 213–233 (2018).
- 674 45. McCabe, L., Britton, R. A. & Parameswaran, N. Prebiotic and probiotic regulation of
675 bone health: role of the intestine and its microbiome. *Curr. Osteoporos. Rep.* **13**, 363–
676 371 (2015).
- 677 46. Carter, D. O., Yellowlees, D. & Tibbett, M. Autoclaving kills soil microbes yet soil
678 enzymes remain active. *Pedobiologia (Jena)*. **51**, 295–299 (2007).
- 679 47. Hale, A. R. & Ross, A. H. Scanning skeletal remains for bone mineral density in
680 forensic contexts. *JoVE (Journal Vis. Exp.)* e56713 (2018).
- 681 48. Lyman, R. L. Bone density and bone attrition. *Man. forensic Taphon.* 51–72 (2014).
- 682 49. Collins, M. J. *et al.* The survival of organic matter in bone: a review. *Archaeometry* **44**,
683 383–394 (2002).
- 684 50. Damann, F. E. & Jans, M. M. E. Microbes, anthropology, and bones. in *Forensic*
685 *Microbiology* (eds. Carter, D. O., Tomberlin, J. K., Benbow, M. E. & Metcalf, J. L.) 319
686 (John Wiley & Sons, 2017).
- 687 51. Mesli, V., Neut, C. & Hedouin, V. Postmortem bacterial translocation. in *Forensic*
688 *microbiology* (eds. Carter, D. O., Tomberlin, J. K., Benbow, M. E. & Metcalf, J. L.) 192–
689 211 (John Wiley & Sons, 2017).
- 690 52. Aerssens, J., Boonen, S., Lowet, G. & Dequeker, J. Interspecies differences in bone
691 composition, density, and quality: potential implications for in vivo bone research.
692 *Endocrinology* **139**, 663–670 (1998).
- 693 53. Child, A. M. Microbial taphonomy of archaeological bone. *Stud. Conserv.* **40**, 19–30
694 (1995).
- 695 54. Booth, T. An Investigation into the relationship between bone diagenesis and
696 funerary treatment. (University of Sheffield, 2014).
- 697 55. Miles, K. L., Finaughty, D. A. & Gibbon, V. E. A review of experimental design in
698 forensic taphonomy: moving towards forensic realism. *Forensic Sci. Res.* 1–11 (2020).
- 699 56. Megyesi, M. S., Nawrocki, S. P. & Haskell, N. H. Using accumulated degree-days to
700 estimate the postmortem interval from decomposed human remains. *J. Forensic Sci.*
701 **50**, 1–9 (2005).
- 702 57. Procopio, N. & Buckley, M. Minimizing Laboratory-Induced Decay in Bone Proteomics.
703 *J. Proteome Res.* doi:10.1021/acs.jproteome.6b00564.
- 704 58. Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical statistical model to
705 estimate the accuracy of peptide identifications made by MS/MS and database
706 search. *Anal. Chem.* **74**, 5383–5392 (2002).
- 707 59. Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A statistical model for
708 identifying proteins by tandem mass spectrometry. *Anal. Chem.* **75**, 4646–4658
709 (2003).

- 710 60. Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased
711 coverage, supporting functional discovery in genome-wide experimental datasets.
712 *Nucleic Acids Res.* **47**, D607–D613 (2019).
713 61. Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019:
714 improving support for quantification data. *Nucleic Acids Res.* **47**, D442–D450 (2019).
715

716 **Supplementary information**

717
718 Supplementary Table 1:

719 Weather data over the course of the experiment period. Data were recorded by two HOBO
720 Micro Station data loggers located on FARF at 30-minute intervals.
721

722 Supplementary Table 2:

723 ADD data during collection of the bone samples and samples taken for proteomics.
724

725 Supplementary Data 1:

726 List of 133 Quantifiable proteins extracted from Progenesis and their localisation and binding
727 site according to Uniprot.
728

729 Supplementary Data 2:

730 Top 20 enriched GO terms for Biological Process (first sheet), Molecular Function (second
731 sheet) and Cellular Component (third sheet) of the proteins extracted from all the samples.
732

733 Supplementary Data 3:

734 List of proteins identified in the four bone types (iliac fresh and skeletonized, tibia fresh and
735 skeletonized).
736

737 Supplementary Data 4:

738 Relative abundances of proteins extracted from all samples only and grouped by bone type
739 (iliac fresh, tibia fresh, iliac skeletonized, tibia skeletonized).
740

741 Supplementary Data 5:

742 List of proteins more abundant in iliac fresh samples than in iliac skeletonized samples (first
743 sheet) and more abundant in tibia fresh samples than in tibia skeletonized samples (second
744 sheet).
745

746 Supplementary Data 6:

747 Relative abundances of proteins extracted from fresh samples only and grouped by donor
748 (D1, D2, D3, D4).
749

750 Supplementary Data 7:

751 Relative abundances of proteins extracted from skeletonized samples only and grouped by
752 deposition type (open pit vs. shallow burial).