



The effects of modern war and military activities on biodiversity and the environment

Journal:	<i>Environmental Reviews</i>
Manuscript ID	er-2015-0039.R1
Manuscript Type:	Review
Date Submitted by the Author:	06-Sep-2015
Complete List of Authors:	Lawrence, Michael; Carleton University, Biology Stemberger, Holly; Carleton University, Biology Zolderdo, Aaron; Carleton University, Biology Struthers, Daniel; Carleton University, Biology Cooke, Steven; Carleton University, Biology
Keyword:	war, biodiversity, ecosystem structure, conservation biology, military activities



1 **The effects of modern war and military activities on biodiversity and the environment**

2

3

4

5

In Prep for Environmental Reviews

6

7

8 Michael J. Lawrence^{1,*}, Holly L.J. Stemberger², Aaron J. Zolderdo¹, Daniel P. Struthers¹ and

9

Steven J. Cooke¹

10

11

12 ¹Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of
13 Environmental Science, Carleton University, Ottawa, ON, Canada K1S 5B6

14

15 ² Department of Biology, Carleton University, Ottawa, ON, Canada K1S 5B6

16

17

18

19 *Address for correspondence:

20 Fish Ecology and Conservation Physiology Laboratory, Carleton University, 1125 Colonel By

21 Dr., Ottawa, ON, Canada, K1S 5B6.

22 Email address: m_lawrence27@live.ca (Michael Lawrence)

23

24 Running Title: War and the environment

25

26

27 Keywords: war, biodiversity, ecosystem structure, conflict, military activities, environment,
28 conservation biology

29

30 **Summary**

31 War is an ever-present force that has the potential to alter the biosphere. Here we review
32 the potential consequences of modern war and military activities on ecosystem structure and
33 function. We focus on the effects of direct conflict, nuclear weapons, military training and
34 military produced contaminants. Overall, the aforementioned activities were found to have
35 overwhelmingly negative effects on ecosystem structure and function. Dramatic habitat
36 alteration, environmental pollution and disturbance contributed to population declines and
37 biodiversity losses arising from both acute and chronic effects in both terrestrial and aquatic
38 systems. In some instances, even in the face of massive alterations to ecosystem structure,
39 recovery was possible. Interestingly, military activity was beneficial under specific conditions
40 such as when an exclusion zone was generated that generally resulted in population increases
41 and/or population recovery; an observation noted in both terrestrial and aquatic systems.
42 Additionally, military technological advances (e.g. GPS technology, drone technology,
43 biotelemetry) have provided conservation scientists with novel tools for research. Because of the
44 challenges associated with conducting research in areas with military activities (e.g. restricted
45 access, hazardous conditions), information pertaining to military impacts on the environment are
46 relatively scarce and are often studied years after military activities have ceased and with no
47 knowledge of baseline conditions. Additional research would help to elucidate the environmental
48 consequences (positive and negative) and thus reveal opportunities for mitigating negative
49 effects while informing the development of optimal strategies for rehabilitation and recovery.

50

51

52

53 **Introduction**

54

55 Conflict has been an ever-present aspect of human civilization. Indeed, the manifestation
56 of conflict in direct combat and military engagements has continuously plagued the world
57 throughout the 20th century leading to more than 100 million human deaths across a number of
58 major and minor wars (Westing 1980; Penderson 2002; Leitenberg 2006; Sarkees et al. 2003).
59 Beyond war's rather obvious negative impacts on human populations (Penderson 2002; Machlis
60 and Hanson 2008), human warfare has also been documented as having a significant influence
61 on the biosphere across a range of ecological scales (Dudley et al. 2002; Machlis and Hanson
62 2008). The degree to which warfare can exert an impact upon an ecosystem and its constituent
63 populations rests entirely on the nature of the disturbance, the sensitivity of the biological system
64 (including resilience) and the timescale of the impacts (Westing 1971; Demarais et al. 1999;
65 Dudley et al. 2002; Warren and Buttner 2006; Warren et al. 2007). Consequently, human conflict
66 has the potential to impart a wide range of impacts on biodiversity and ecosystem structure and
67 function. Interestingly, although one might presume that all conflict is overwhelmingly
68 "negative" in an ecological context, in reality the consequences of warfare generate a continuum
69 of outcomes ranging from highly positive to highly detrimental.

70

71 While a large body of knowledge of the consequences of war on the ecological dynamics
72 of a variety of biological systems is known, a comprehensive assessment of these impacts has yet
73 to be conducted. Current reviews on the subject often frame ecological changes in the greater
74 context of socio-economic factors and human interactions which are often restricted to terrestrial
75 mammalian megafauna (e.g., Dudley et al. 2002; Machlis and Hanson 2008). Thus, the purpose

76 of this review will be to address the specific impacts of modern warfare (i.e. turn of the 20th
77 century) on ecosystem structure (especially biodiversity and the status of populations and
78 communities) and ecosystem function in a variety of systems (e.g. aquatic, terrestrial). For the
79 sake of simplicity, our analysis will be restricted to the following impacts of military activities:
80 direct armed conflict (between two or more factions), nuclear warfare, military training, and
81 military produced chemical and metals contamination. For the entirety of this review, the term
82 warfare will encompass the preparation (e.g. training, material development and testing),
83 mobilization, conflict and related activities of nations/factions involved in a military operation
84 against one another. This review will also limit its scope to include assessments of the impacts of
85 military activities on ecosystem structure and function during the “preparations for war”,
86 “violent conflict” and “post-war activities” phases as outlined in Machlis and Hanson (2008). As
87 such, any activity that directly relates to preparation and/or is a product of war, outside of
88 civilian operations, will be considered an aspect of warfare. Our assessment will encompass a
89 continuum-based approach whereby both the negative and positive impacts of the above factors
90 are highlighted appropriately.

91

92 **Active Armed Conflict**

93

94 Armed conflict is the act of war generated by two or more governmental, non-
95 governmental groups or international states that generally involves a combination of active
96 military actions, including aerial assaults, naval craft operations, or ground forces (ICRC 2008;
97 Machlis and Hanson 2008; Pearson 2012). Often, natural ecosystems are termed “terrain” in
98 military battlespace terminology (O’May et al. 2005; Visone 2005; Hieb et al. 2007), taking on

99 an anthropogenic rather than an eco-centric view of natural landscapes during periods of armed
100 conflict. As a result, ecosystem health and integrity are often neglected casualties of warfare
101 with little responsibility from involved factions in contributing to conservation efforts (Gangwar
102 2003; Clark and Jorgenson 2012). The consequences of active armed conflict range across a
103 spectrum of ecological scales and lead to unexpected and complex outcomes - either beneficial,
104 negative, or a combination of these two. This component of the paper highlights a number of
105 types of active warfare engagement forms including airborne, naval, and ground warfare
106 activities, which have demonstrable impacts on ecosystem structure and function.

107

108 *Aerial Assault:*

109

110 Aircraft (both rotary and fixed-wing) are commonly used in military operations that can
111 produce bursts of noise (e.g. sonic booms, jet afterburners, rotary pulses, etc). The auditory
112 system is more sensitive in many animals compared to that of humans (Manci et al. 1988; Larkin
113 et al. 1996) and thus aerial activities possess a significant source of noise pollution that is of
114 global concern for the wellbeing of wildlife (Dunnet 1977; Dufour 1980; Gladwin et al. 1988).
115 The production of noise from military aircraft has variable impacts on wildlife, which encompass
116 primary, secondary, and tertiary effects (Janssen 1980; reviewed in Manci et al. 1988). These
117 effects can occur over an acute or chronic timescale representing both sub-lethal and lethal
118 impacts that have the potential to cause permanent damage; a factor that is influenced by
119 acoustic duration, intensity and the biology of the specific species. Primary effects can include
120 eardrum rupture, shifts in hearing abilities (either temporary or permanent), and/or auditory
121 signal masking (e.g. unable to identify noises from prey, predators, or mates). Secondary effects

122 are related to physiological impacts (Manci et al. 1988), which can lead to impediments in
123 reproduction, foraging behaviour, and natural habitat use of wildlife residing in areas where
124 aircraft noise is prevalent (Francis 2011). Tertiary impacts consist of a combination of primary
125 and secondary effects that can lead to population declines, species extinction and habitat
126 degradation (Klein 1973; Bender 1977; Manci et al. 1988).

127

128 Ecosystem structure has been affected by means beyond noise pollution from military
129 aircraft. For example, during WWII, aircraft acted as a vector for the transportation of exotics
130 whereby weeds and cultivated species were brought to oceanic island ecosystems by way of
131 aircraft landing strips used for refueling and staging stations during operations in the Pacific
132 theatre (Stoddart 1968). Prior to the war, these isolated islands were home to a number of
133 sensitive and endemic species that had naturally dispersed to their current positions. However, in
134 the aftermath of aerial warfare events, large numbers of invasive species had become established
135 on these small islands which altered the evolutionary pathways of native species causing
136 competitive exclusion, predation, and extinction of endemic species (Mooney and Cleland 2001).
137 Aerial warfare also has had a great influence on altering population dynamics directly. Air-to-
138 ground assaults are known to cause elevations in wildlife mortality (Zahler and Graham 2001;
139 Gangwar 2003) and destroy natural habitat (Levy et al. 1997) both of which may contribute to a
140 localized population decline. Conventional aerial assault weapons are generally categorized into
141 4 groups which include: high explosive fragmentation, incendiary weapons, enhanced blast
142 munitions, and defoliants; all of which have potential to destroy wildlife and natural habitat in
143 different ways and with varying degrees of severity (reviewed in Majeed 2004). These impacts
144 have been illustrated in a number of species including Asian elephants (*Elephas maximus*;

145 Chadwick 1992; Dudley et al. 2002) and snow leopards (*Panthera uncia*; Zahler and Graham
146 2001) where aerial combat maneuvers were observed to decimate entire forest ecosystems
147 leaving behind stumps and craters, alongside contaminated and destabilized soils (Levy et al.
148 1997).

149

150 *Naval Operations:*

151

152 Naval conflict between foreign nations has a diverse range of effects on the marine
153 environment. Like aircraft, ships have been implicated in introducing foreign species to
154 otherwise un-colonisable regions under normal circumstances. This has been achieved through
155 the dumping of ballast waters (Apte et al. 2000) and the introduction of naval
156 structures/materials into the region (Tavares and De Melo 2004). As an example of the latter, the
157 brown tree snake (*Boiga irregularis*) was introduced into Guam in 1949 just after WWII, most
158 likely as a stowaway on boats salvaging materials from a port in New Guinea (Rodda and
159 Savidge 2007). This species has subsequently invaded all terrestrial ecosystems in Guam leading
160 to the extirpation of many bird and lizard species, as well as a number of other native
161 invertebrates thus having a measurable effect on the local biodiversity (Rodda and Savidge
162 2007).

163

164 Naval blasts and sonar operations, during active periods of warfare, have the potential to
165 interfere with the daily lives of many aquatic species. The acoustic frequency used by dolphins
166 and whales coincides with that used by naval sonar devices which can cause ear hemorrhaging
167 and beach stranding (Science Wire 2001; NRDC 2003). In addition to this, conventional naval

168 ordinance (e.g. depth charges, torpedoes) create substantial underwater blasts which can inflict
169 overpressure and fragmentation injury to invertebrates, fish, reptiles, birds, and marine mammals
170 in proximity of the blast radius (Gaspin 1975; Westing 1980; Ketten 1995; Reviewed in Keevin
171 and Hempen 1997; see Nuclear Warfare for more info on blast injury).

172

173 While there are a number of negative impacts associated with naval operations, marine
174 environments have profited from this activity in a number of ways. Fish populations greatly
175 benefited from the activities occurring in the North Atlantic during WWII whereby sensitive and
176 overexploited populations were given time to recover from anthropogenic disturbances and
177 fisheries exploitation (Beare et al. 2010) as fishing fleets were drastically reduced in size
178 resulting from their participation in naval operations including mine sweeping and shipping
179 supplies (Gulland 1968; Engelhard 2008). If not called to assist in military services, then fishing
180 vessels were often harboured and, therefore, excluded from fishing activity due to threats at sea
181 from naval/aerial strikes and subsurface mining (Beare et al. 2010). During this period of war,
182 large areas in the Atlantic Ocean functioned as Marine Protected Areas (MPAs) for several years
183 which allowed commercial fish populations to proliferate with a reduction in fishing effort
184 (Beare et al. 2010). During this time, it was observed that the reduction in fishing mortality
185 altered the age-structure dynamics of gadoid fisheries resulting in a larger proportion of
186 mature/larger fish which allowed populations to proliferate to a greater extent (Beare et al. 2010).
187 Additionally, opportunistic species (e.g. oceanic whitetip sharks, *Carcharhinus longimanus*)
188 have been reported as benefitting from naval ship wrecks as they provided a rich food source
189 during periods of warfare representing an acute “ecological bonanza”(Bass et al. 1973).

190 Indirectly, the occurrence of naval warfare allowed fisheries and other untargeted species to
191 rebound and proliferate which may not have otherwise occurred in its absence.

192

193 Naval conflicts, particularly during WWII, have also led to the creation of heterogeneous
194 habitats that would not exist otherwise. During World War II, there was a global expansion with
195 ocean-going vessels that navigated the coastal and pelagic waters of the Atlantic and South-
196 Pacific oceans to engage hostile countries. Although this led to devastating consequences for
197 human life, the resulting ship wrecks created a large number of artificial reefs where aquatic life
198 could colonize, utilize, and flourish (Hynes et al. 2004). While there are concerns regarding long
199 term contamination with sunken naval craft (Westing 1980; Martore et al. 1998; Ampleman et al.
200 2004; Barrett 2011; Monfils 2005; see Military Contamination), these vessels have proven to be
201 a source of new habitat for aquatic life in areas of the ocean that were largely void of structure
202 for animals to colonize (Hynes et al. 2004).

203

204 *Terrestrial Conflict:*

205

206 Ground warfare often takes place in sensitive and remote locations around the globe.
207 Indeed, a large number of biological hotspots have set the stage for major ground conflict events
208 (Kim 1997; Hart et al. 1997; Hanson et al. 2009). Furthermore, modern ground warfare has often
209 altered natural landscapes and impacted wildlife in a number of different ways. Often, soldiers
210 were positioned for on-ground battle within critical habitats of endemic and endangered species
211 (Shambaugh et al. 2001; Zahler and Graham 2001; Hanson et al. 2009; Lindsell 2011)
212 representing a potential threat to these organisms. As one might expect, armed conflict found

213 within terrestrial ecosystems often facilitates poaching by military forces (Shambaugh et al.
214 2001; Draulans and Krunkelsven 2002; Dudley et al. 2002) and can promote further destruction
215 of the landscape and wildlife populations by displaced refugees of war (Shambaugh et al. 2001;
216 McNeely 2003; Dudley et al. 2002; Dubey and Shreni 2008). In contrast, there are reports of
217 large adaptable predators, including Bengal tigers (*Panthera tigris tigris*) and grey wolves (*Canis*
218 *lupus*) becoming habituated to gunfire noise on the battlefields of WWII; they were often sighted
219 foraging on casualties in the aftermath of battles (Orians and Pfeiffer 1970; Westing 1980;
220 McNeely 2003) which may acutely benefit the species as in the case of marine predators
221 illustrated above.

222
223 The weapons employed by militaries probably pose the greatest hazard by terrestrial
224 conflicts to ecosystem structure. The numerous explosive techniques and tools at the disposal of
225 army forces during ground warfare have left a legacy on landscapes across the globe by leaving
226 large craters, shrapnel, and contamination, thus devastating many ecosystems across the
227 biosphere (Westing 1980; Hupy 2008; Certini et al. 2013). Landmines applied during active
228 ground warfare have left a lasting legacy on the environment and still remain a major threat to
229 biodiversity, even decades after being deployed (Westing 1985; Roberts and Williams 1995;
230 Reviewed in Berhe 2007). However, landmines may help ecosystems recuperate after heavy
231 impact from armed conflict by creating “no-mans-land” in an analogous manner to a game
232 reserve or park as seen in the case of the cranes in the demilitarized zone of the Korean Peninsula
233 (Box 1; Higuchi et al. 1996; Kim 1997; Dudley et al. 2002). Landmines do not differentiate
234 between soldiers and wildlife (especially large mammals) and therefore, many organisms have
235 been damaged or killed directly from landmine explosions (Shambaugh et al. 2001; Zahler and

236 Graham 2001; Westing 1996; Berhe 2007). Indeed, land mines have been responsible for
237 pushing at risk species closer to extinction (e.g. elephants in Africa, leopards in Afghanistan;
238 Troll 2000) and deteriorating ecosystem integrity by destroying vegetation and degrading soil
239 structure (Miller 1972; Berhe 2007).

240

241 Artillery fire also poses a risk to the environment. During WWI and II, artillery weapons
242 were positioned behind soldiers and were fired towards the opposing factions with the capability
243 of firing hundreds of shells per hour (Hupy 2008). Troops often found shelter or fought battles in
244 forested areas resulting in heavy artillery fire on these regions, devastating the local ecosystem
245 and associated biodiversity (Hupy 2008). Decades after WWII, craters in Verdun, France
246 produced by heavy artillery fire still remain void of vegetative growth; deep craters extending to
247 the water table cause hydric conditions, making them unsuitable for colonization by terrestrial
248 plant species (Hupy 2006). Thus, shelling can result in chronic legacy impacts in addition to
249 acute influences (e.g. instant mortality).

250

251 Terrestrial conflicts have been known to target military and civilian infrastructure to stifle
252 opposing factions. Ground forces, in the past, have used explosives to destroy hydropower dams
253 (Sweetman 1982; Gleick 1993; Clodfelter 2006) and dikes (Lacoste 1973) as a means to impede
254 the mobility of countering factions (Francis 2011). The abrupt removal of long established dams
255 can cause a number of ecological consequences such as siltation, mortality of fish and wildlife
256 populations situated above and below the dam (e.g. abrasion, suffocation, habitat loss), and
257 produce lasting physical, chemical, and biological legacies (Bednarek 2001; Stanley and Doyle
258 2003).

259

260

261 Nuclear Warfare

262

263 The development and use of nuclear warheads, in both times of peace and conflict, has
264 undoubtedly left a significant scar on the earth's surface. As of the late 1990s, more than 2000
265 nuclear weapons tests have been conducted around the world (Yang et al. 2003). The detonation
266 of a nuclear warhead represents a significant threat to local biodiversity as, unlike conventional
267 ordinance, the energy released is partitioned into three distinct categories including thermal (35
268 %), kinetic (50 %) and radioactive (15 %) energies (Glasstone 1964; Brode 1968; Nishiwaki
269 1995; Eisenbud and Gesell 1997). Here we will review the documented and potential effects of
270 each of these detonation impacts on ecosystem structure and function.

271

272 Thermal Impacts:

273

274 Thermal emissions from nuclear blasts can have a number of impacts on local
275 ecosystems. The immense release of thermal energy at the detonation's epicentre results in
276 temperatures far in excess of 3000°C (Brode 1968; Pinaev and Shcherbakov 1996). As such,
277 thermal emissions pose a lethal force to any life in the vicinity of the epicentre resulting from
278 incineration (Glasstone 1964; Lifton 1967) as seen in the bombings of Japan (Summary Report
279 1951; Silberner 1981; Ruhm et al. 2006; Ochiai 2014). Beyond the epicentre, an outward thermal
280 wave (100-1000°C) moves radially (a distance dependent on the bomb strength) (Brode 1968)
281 and is a serious risk to most life over its expansion. Here, local vegetation is burnt and defoliated,

282 often perishing through the extreme heat (Palumbo 1962; Shields and Wells 1962; Shields et al.
283 1963; Craft 1964) representing severe reductions in plant species richness and abundances
284 (Shields and Wells 1962; Shields et al. 1963; Palumbo 1962), not unlike an intense forest fire
285 (Noble and Slatyer 1980; Rowell and Moore 2000; Grace and Keeley 2006). The spatial extent to
286 which vegetation burning occurs is highly dependent on the status (e.g. moisture content) and
287 composition of the vegetative assemblages present in the blast area (Chandler et al. 1963; Craft
288 1964; Small and Bush 1985). Some have speculated that thermal emissions may indirectly
289 impact adjacent forests/vegetative regions, through the generation and spread of wildfires
290 (Chandler et al. 1963; Craft 1964) that may extend the immediate population/diversity reduction
291 outside of the blast area for both plants (Noble and Slatyer 1980; Rowell and Moore 2000; Grace
292 and Keeley 2006) and animals (Singer et al. 1989; Kaufman et al. 1990; Moreira and Russo
293 2007; Lindenmayer et al. 2008). In contrast to plant life, there is comparatively little research on
294 the effects of thermal impacts from nuclear blasts on animals, humans notwithstanding. Thermal
295 wave exposure has been reported to cause severe whole body burns on unprotected skin in
296 humans (Kajitani and Hatano 1953; Oughterson et al. 1951; Oughterson and Warren 1956;
297 Nishiwaki 1995). In the bombings of Japan, fatal burns and mild non-lethal burns were observed
298 within 1.2-2.5 km and 3-4 km from the epicentre, respectively, (Oughterson et al. 1951;
299 Oughterson and Warren 1956; Glasstone 1964; Nishiwaki 1995) with the former resulting in a
300 large proportion of the total deaths (~30%) during this event (Oughterson and Warren 1956;
301 Glasstone 1964; Nishiwaki 1995). Additionally, thermal radiation, along with high intensity
302 visible radiation, can also result in severe retinal burning in humans (Oyama and Sasaki 1946;
303 Rose et al. 1956; Glasstone 1964). There is no reason to assume that similar consequences would
304 not be observed among terrestrial wildlife, especially mammals.

305

306 Experimental tests of simulated and/or actual nuclear weapons produced thermal energy
307 exposure in rats (Alpen and Sheline 1954), dogs (Brooks et al. 1952; Richmond et al. 1959a),
308 rabbits (Byrnes et al. 1955; DuPont Guerry III et al. 1956; Ham et al. 1957) and swine (Baxter et
309 al. 1953; McDonnel et al. 1961; Hinshaw 1968) have generated analogous effects as seen in
310 humans suggesting that wild mammals may have a similar burn response during a nuclear
311 detonation. Severe burns were also reported in teleost fish that were in close proximity to the
312 detonation of the warhead in Bikini Atoll (Donaldson et al. 1997). Not surprisingly, in simulated
313 experiments, severe burns increased the rates of mammalian mortality, resulting from general
314 physiological disturbances and secondary infection occurring 0-2 weeks “post-blast” (Brooks et
315 al. 1952; Alpen and Sheline 1954; McDonnel et al. 1961). This effect was also amplified under a
316 combined thermal and radiation exposure resulting in a severely immunocompromised,
317 physiologically disturbed individual (Brooks et al. 1952; Baxter et al. 1953; Alpen and Sheline
318 1954; Valeriote and Baker 1964; Ledney et al. 1992) similar to what is believed to occur in
319 humans (Nishiwaki et al. 2000). Scaling these effects up, it would be highly likely that thermal
320 emission exposure would result in a large die-off event in the local animal life thereby reducing
321 local populations and, potentially, reducing local species richness over an acute time frame (0-2
322 weeks). It should be noted that the intensity of the burns is likely to be a product of the distance
323 from the epicentre as the thermal wave will gradually reduce in magnitude (Brooks et al. 1952;
324 McDonnel et al. 1961; Glasstone 1964), a factor that must be acknowledged when predicting the
325 expected impacts on animal populations. However, this effect would not be equal for all
326 creatures as rats on Bikini Atoll were able to avoid both thermal and kinetic emissions from
327 warhead testing even in close proximity to the blast as a product of their subterranean existence

328 (Donaldson et al. 1997). As such, we would expect that species occupying “sheltered” habitats
329 may not experience a large die-off as described earlier.

330

331 *Blast Effects:*

332

333 As mentioned earlier, in a nuclear warhead detonation, blast energy accounts for
334 approximately 50% of the total emitted energy that moves away from the epicentre in a radial
335 pattern (Randall 1961; Glasstone 1962; Eisenbud and Gesell 1997; Glasstone 1964). The large
336 amount of kinetic energy emanating from the blast (1-3500+ kPa) is especially damaging to
337 plants whereby the blast force is capable of denuding foliage as well as damaging branch
338 structure and uprooting vegetation from the soil (Shields and Wells 1962; Shields et al. 1963;
339 Palumbo 1962; Beatley 1966; Glasstone and Dolan 1977; Hunter 1991) effectively destroying a
340 large proportion of the surrounding plant life and primary production.

341

342 Animals caught within the blast wave can be impacted in a number of ways. Terrestrial
343 species are likely to experience damage resulting from overpressure injury. Using blast pressures
344 similar to what has been reported during nuclear explosions, rats experienced severe lung
345 damage as well as large degrees of hemorrhaging in various regions of the body (Jaffin et al.
346 1987). Similar effects have been noted in a number of other vertebrate species (Richmond et al.
347 1959a; 1959b; Goldizen et al. 1961; Richmond and White 1962; Candole 1967; Jaffin et al.
348 1987; Mayorga 1997) with the extent of physiological damage dependent upon the mass of the
349 animal (larger animals are less susceptible to injury; Richmond and White 1962; Jaffin et al.
350 1987) as well as the magnitude and duration of the over-pressure exposure (Candole

351 1967). Unsurprisingly, mortality in these trials was elevated (Richmond et al. 1959a; 1959b;
352 Richmond and White 1962; Jaffin et al. 1987) which, under an actual nuclear detonation, would
353 be expected to increase mortality rates in exposed populations. Further exacerbating these effects
354 would be the large amount of debris and shrapnel carried through the air by the blast causing
355 injury and death to animals in the surrounding area (Candole 1967; Mayorga 1997). This effect
356 has been directly observed during a nuclear detonation on both humans (Shaeffer 1957; Liebow
357 1983; Kishi 2000) and other mammalian species (McDonnel et al. 1961; Goldizen et al. 1961;
358 Masco 2004).

359

360 Aquatic organisms are particularly sensitive to the effects of a blast. While direct
361 evidence is rather limited in the literature, nuclear denotations in proximity to aquatic
362 environments have been shown to result in large fish population die offs (Kirkwood 1970;
363 Kirkwood and Fuller 1972; Merritt 1970; Merritt 1973; Planes et al. 2005) demonstrating similar
364 impacts to conventional ordinance explosion on fish mortality on a much larger scale (Govoni et
365 al. 2008; Popper and Hastings 2009). This is primarily a result of the anatomical design of teleost
366 fish having a gas filled swim bladder that is easily ruptured upon exposure to large pressure
367 differentials (Simenstad 1974; Yelverton et al. 1975; Baxter et al. 1982; Planes et al. 2005;
368 Popper and Hastings 2009). Marine mammals, given the presence of large gas filled lungs,
369 would also be expected to suffer high rates of mortality under a nuclear blast resulting from
370 severe lung damage in a manner similar to that of fish swim bladders (Baxter et al. 1982;
371 Goertner 1982). Marine mammals in proximity to a warhead detonation experienced severe lung
372 damage and elevated mortality (Kirkwood and Fuller 1972; Rausch 1973). This effect also
373 extended to diving birds (Kirwood and Fuller 1972; Rausch 1973). Interestingly, invertebrates

374 are not seemingly affected by pressure waves in aquatic systems (Isakason 1974; Baxter et al.
375 1982) and are unlikely to be impacted, in this manner, under a nuclear blast. However, not all
376 invertebrates are equal, in respect to kinetic energy disturbances, in that warhead detonation over
377 coral reefs lead to widespread coral death presumably through mechanical disruption from the
378 blast (Richards et al. 2008). While most of the coral community appears able to recover, highly
379 turbid conditions generated during blasts have led to the extinction of calm water specialist coral
380 species on some reefs (Richards et al. 2008).

381

382 Both thermal and kinetic impacts of a nuclear detonation occur over an acute timeframe
383 and would likely result in a great reduction in the abundances and diversity of local flora and
384 fauna. However, over a more chronic duration, these impacts are likely to be minimal as
385 populations and diversity could recover through dispersal to the area as well as contributions
386 from surviving organisms. Indeed, this has been observed in a number of plant (Palumbo 1962;
387 Shields and Wells 1962; Shields et al. 1963; Beatley 1966; Fosberg 1985; Hunter 1991; 1992)
388 and animal (Pinca et al. 2005; Planes et al. 2005; Richards et al. 2008; Houk and Musburger
389 2013; Jorgensen and Hayward 1965; O'Farrell 1984; Hunter 1992; Wills 2001; Kolesnikova et
390 al. 2005) communities from a diversity of testing site environments. In some instances, the
391 exclusion of human activity from test sites has been quite beneficial to the recovery and
392 prosperity of organisms found in these areas as in the case of the atolls of the Marshall Islands
393 (see Box 2; Davis 2007; Richards et al. 2008; Houk and Musburger 2013).

394

395 *Radiation Impacts:*

396

397 Nuclear weapons emit a portion of their energy as ionizing, radioactive emissions either
398 as electromagnetic radiation (e.g. gamma & X rays) or through radionuclides of various elements
399 (Aakrog 1988; Robison and Noshkin 1999; Whicker and Pinder 2002) which are accumulated
400 primarily through direct exposure or through consumption of producers, respectively (Donaldson
401 et al. 1997; Entry and Watrud 1998; Whicker and Pinder 2002). However, the effects of
402 radioactivity on life are variable. Over an acute timescale, provided sufficient activity (< 2 Gy),
403 radiation exposure in humans can result in the development of radiation poisoning that can
404 manifest itself as (depending on the dose) hemorrhaging, blood cell and tissue destruction, and
405 mortality in doses in excess of 6 Gy (Prosser et al. 1947; Ohkita 1975; Guskova et al. 2001;
406 Mettler 2001) thus accounting for the elevated mortality rate in the bombings of Japan (Ohkita
407 1975). Similar effects have been observed to occur in terrestrial mammals in both laboratory
408 experiments (Eldred and Throwbridge 1954; Brown et al. 1961; Zallinger and Tempel 1998) and
409 bomb exposed animals (Tullis et al. 1955; McDonnell et al. 1961; Zallinger and Tempel 1998)
410 resulting in considerable mortality. As previously mentioned, radiation and thermal energy
411 exposure can work synergistically to induce higher mortality rates (Brooks et al. 1952; Baxter et
412 al. 1953; Alpen and Sheline 1954; Valeriote and Baker 1964; Ledney et al. 1992). In plants,
413 acute radiation exposure results in tissue degradation and death under sufficiently high
414 radioactivity levels (Sparrow and Woodwell 1962; Shields et al. 1963; Rhoads and Platt 1971;
415 Rhoads et al. 1972). However, the extent of tissue damage in plants varies with development
416 state (Sparrow and Woodwell 1962; Shields et al. 1963; Rhoads and Platt 1971; Rhoads and
417 Ragsdale 1971). Together, these effects could represent a substantial source of mortality
418 following a weapon detonation on ecosystems on an acute time scale.

419

420 Radioactive exposure may also lend itself to more chronic impacts on animal
421 populations. In humans exposed to nuclear weapon emissions, there has been an observed
422 elevation in the rates (Bizzozero et al. 1966; Wanebo et al. 1968; Prentice et al. 1982; Darby et
423 al. 1988) and risk level (Pierce and Preston 2000) of developing a chronic disease such as
424 neoplasia. Assuming this effect occurred in a similar manner as in humans (Mole 1958), it would
425 be expected to significantly reduce life expectancies and survival in wild animals. Chronic
426 radiation effects may also result in the development of chromosomal and/or genetic aberrations
427 (Hatch et al. 1968; Bickham et al. 1988; Lamb et al. 1991; Sugg et al. 1995) in addition to altered
428 genetic structure of populations (Theodorakis and Shugart 1997; Theodorakis and Shugart 1998;
429 Theodorakis et al. 1998) in wild animals under radiation exposure from weapons test and
430 development sites. While extremely limited data exist, reduced reproductive capacities in wild
431 animals have been noted at detonation sites (Medica et al. 1973; Turner et al. 1971; Turner 1975;
432 Turner and Meica 1977) consistent with the expected effects of radiation's impacts on the
433 reproductive system (Reviewed in Real et al. 2004). However, this effect seems to be variable as
434 a few species at weapons test sites seem to have no genetic or macroscopic level impacts (Hatch
435 et al. 1970; Campbell et al. 1975; Theodorakis et al. 2001) with the sensitivity of reproductive
436 systems to radiation being non-ubiquitous among species (Barnthouse 1995; Mudie et al. 2007).
437 It is believed that in some cases the "null" effect of radiation may be the product of immigration
438 of non-affected individuals into the irradiated area (Theodorakis et al. 2001). The overall effect
439 of these long term impacts are relatively uncertain and could have variable consequences on a
440 given population depending on the strength and type of the effect. However, it should be noted
441 that because of the high degree of hazard (i.e. radiation) and security precautions associated with
442 nuclear weapons/production sites, many of these areas are devoid of human activity and thus

443 serve as important refuge sites for a variety of plant and animal species. Indeed, these areas have
444 been demonstrated in having quite diverse and thriving ecosystems that are often in a better
445 ecological state when compared to similar areas where routine human activity is present (see Box
446 2; Gray and Rickard 1989; Whicker et al. 2004; Davis 2007; Richards et al. 2008; Houk and
447 Musburger 2013). Thus, sites devoted to nuclear arms production and testing can still be
448 considered a positive feature in maintaining biodiversity despite the potential for chronic health
449 impacts in resident organisms.

450

451 **Military Infrastructure and Bases**

452

453 *Military Bases:*

454

455 The impacts of war on ecosystems are not limited to armed conflict events, but can be
456 connected to, and influenced by, the development and operational use of military training bases.
457 A military training base is a general designation applied to military facilities that house military
458 equipment and personnel, and facilitate training exercises and tactical operations (Kazmarek et
459 al. 2005; Zentelis and Lindenmayer, 2014). Military training bases can range from small outpost
460 sites to large military “cities” (Brady 1992). The variation in size and operational use of military
461 training bases leads to a broad spectrum of anthropogenic impacts, both in type and severity, on
462 the local ecosystem (Owens 1990; Rideout and Walsh 1990; Goldsmith 2010). These impacts
463 can be broken down into two broad categories: (1) the development of military training bases,
464 which includes the establishment and construction of the facility and site and (2) operations of
465 the military training base, which include the functional operation of the infrastructure itself and

466 the corresponding military activities designated for the specific site. In this section, we will focus
467 our discussion on the effects of development and operations of military training bases (including
468 air, naval, and terrestrial) on ecosystem structure and function.

469

470 *Environmental Impacts of Military Base Development:*

471

472 The environmental impacts associated with the construction of infrastructure projects are
473 site specific (Augenbroe and Pearce 1998; Tang et al. 2005; Gontier 2007; Mortberg et al. 2007).

474 For example, the development of naval ports and shipyards are more likely to have a greater
475 contamination risk of adjacent water bodies than the development of a terrestrial airstrip which
476 can be situated miles from water sources and surrounded by a natural vegetation buffer zone
477 (Tull 2006; Mortberg et al. 2007). Even the construction of similar base infrastructure, situated in
478 different locales, are subject to different environmental impacts based on the landscape and
479 ecosystem they are built within and thus impacts are highly site specific (Kazmarek et al. 2005;
480 Gontier 2007; Mortberg et al. 2007). Although construction projects are associated with site-
481 specific environmental impacts, the focus of this section is not to dissect these site-specific
482 characteristics, but to address some overarching impacts on ecosystems that are germane to most
483 military base development projects.

484

485 There are several generic impacts associated with the construction of most complex
486 infrastructure projects. Some of these impacts include habitat degradation, soil erosion, and
487 chemical contamination (Westing 1980; Tang et al. 2005; Xun et al. 2013). Initial site
488 development requires the clearing of vegetation and trees, followed by intensive soil excavation

489 and compaction. This process alters the natural landscape by the removal of existing vegetation
490 and the prevention of future vegetation growth (Kopel et al. 2015). The removal of vegetation
491 coupled with soil excavation increases the potential for soil erosion, and reduces water
492 infiltration rates, altering the landscape ecology by changing soil structure and chemistry, and
493 increasing water runoff rates (Tang et al. 2005). Chemical contamination of local water sources
494 can also occur from increased water runoff carrying sediments and chemicals associated with
495 waste dumping (e.g. hazardous building materials, paints, solvents, etc.), and accidental chemical
496 spills (e.g. fuel and oil) during the development stage (Brady 1992; Kazmarek et al. 2005;
497 Villoria Saez et al. 2014; Kopel et al. 2015). These pollutants can alter community structure
498 within the vicinity of the infrastructure (Meyer-Reil and Koster 2000; Beasley and Kneale 2002;
499 Edwards 2002; Osuji and Nwoye 2007).

500 However, the establishment of military training bases can also have beneficial impacts on
501 biodiversity at the local, regional, and global scale. For effective combat training in real-world
502 scenarios, military training bases need to be large and encompass a wide variety of environments
503 and climates (Stephenson et al. 1996; Doxford and Judd 2002; Smith et al. 2002). Depending on
504 the specific nature and use of military training areas, public and commercial access are usually
505 restricted due to safety and security issues. This creates great tracts of land largely devoid of
506 human contact and commercial development, preserving these wilderness areas which have been
507 lost to human development elsewhere (Rideout and Walsh 1990; Doxford and Judd 2002;
508 Zentelis and Lindenmayer 2014). Military training areas have been increasingly recognized as
509 areas of high biodiversity, and in particular, for harbouring endangered and at-risk species (Box
510 3). It has been estimated that in the United States alone, over 200 federally listed endangered
511 species inhabit military training areas; which is more endangered species per area within military

512 installations compared to other federally managed lands in the United States (Doxford and Judd
513 2002; Pekins 2006; Zentelis and Lindenmayer 2014). Aside from these training lands supporting
514 IUCN red-listed species, they also support highly diverse landscapes. The U.S. Army holds two
515 of their largest European training bases in Bavaria, Grafenwohr and Hohenfels, which are
516 situated on 22,855 and 16,175 ha of land, comprising 0.34 and 0.24% of the land area in Bavaria,
517 respectively (Warren et al. 2007). Despite the relatively small size of these training areas and
518 their exposure to intensive military training exercises, they contain approximately 27% of the
519 total plant species richness found in Bavaria (Schonfelder et al. 1990). Similarly, the military
520 training areas in the Netherlands comprise approximately 1% of the total available land area, but
521 have been reported to support approximately 53% of all vascular plant species, and 61% of all
522 bird species found within this nation (Gazenbeek 2005; Warren et al. 2007). It is also important
523 to recognize the significance of military training areas to provide key habitat for wide-ranging
524 megafauna species such as bears, ungulates, coyotes and wolves that require large tracts of land
525 for foraging and hunting (Gese et al. 1989; Stephenson et al. 1996; Telesco and Van Manen
526 2006). Globally, military training areas have been estimated to encompass approximately 6% of
527 the Earth's surface spanning a multitude of environments and ecosystems. This extended global
528 coverage makes military training lands important areas for biodiversity conservation and
529 preservation (Zentelis and Lindenmayer 2014), notwithstanding the fact that the type of activities
530 that occur on these sites could rapidly alter biodiversity. Recognizing the importance of military
531 facilities in conserving biodiversity, the USA has begun rehabilitating former training sites to
532 serve as nature preserves (Coates 2013; Havlick 2014). As of 2014, 15 of these areas have been
533 developed in an effort to promote and conserve the biodiversity of these regions (Havlick 2014).
534 In this way, military facilities are of great benefit to sustaining and conserving biodiversity.

535

536 *Operations of a Military Training Base:*

537

538 The environmental impacts associated with the upkeep of military infrastructure and
539 equipment has been a growing concern. Many military bases have been targeted for
540 environmental assessment and site remediation (Kazmarek et al. 2005; Goldsmith 2010).
541 Military infrastructure and equipment is subject to rigorous use, often under extreme conditions,
542 creating the need for constant maintenance and upkeep. This maintenance leads to the generation
543 of large quantities of hazardous wastes including heavy metals, solvents, corrosives, paints, fuel,
544 and oils (Brady 1992; Kazmarek et al. 2005). When these hazardous wastes are improperly
545 stored/disposed of, it can cause serious water contamination and habitat degradation issues,
546 which can directly affect biodiversity (Edwards 2002; Osuji and Nwoye 2007). There have even
547 been documented reports of military sites that dump hazardous wastes into open holding ponds,
548 evaporation ponds, mines, and wells (Brady 1992; see Military Contamination for more detail).
549 The Otis Air Base in the United States has received significant attention over the past few
550 decades due to the extensive contamination of groundwater caused from fuel spills and aircraft
551 maintenance (Kazmarek et al. 2005; Goldsmith 2010). Similarly, the Norton Air Force Base in
552 the US is under scrutiny for its poor approach of storing hazardous wastes in above and below
553 ground storage drums, which have begun to leak, causing environmental contamination issues
554 (Brady 1992). However, poor environmental planning at military bases appears to be a common
555 theme. The US Environmental Protection Agency has listed over 53 military bases on the
556 National Priorities List of sites that pose direct hazards to human health and the environment
557 (Brady 1992; Kazmarek et al. 2005; Goldsmith 2010). Unfortunately, the majority of literature

558 on the environmental impacts associated with the upkeep of military infrastructure and
559 equipment is focused mainly on the USA with, comparatively, little known about such issues in
560 other jurisdictions.

561

562 *Training Activities:*

563

564 Live-fire training has similar impacts on the environment as those discussed in the active
565 armed conflict section, with respect to local landscape alteration and vegetation destruction,
566 chemical and heavy metal contamination, and the incidental killing or maiming of wildlife.

567 However, there are also differences in environmental impacts of live-fire training that occur in
568 training facilities as opposed to actual armed conflict events (Owens 1990; Goldsmith 2010).

569 Training facilities are faced with the challenge of repeated use of live-fire training shooting
570 ranges, which leads to consistent site-specific degradation and contamination. The most common
571 and extensive live-fire training occurs on small arms ranges (Goldsmith 2010), which are
572 associated with extensive heavy metal contamination, with lead being the most notable
573 contaminant (Cao et al. 2003a; 2003b; Goldsmith 2010). The weathering and oxidation of lead
574 bullets leads to the contamination of soils, groundwater and surface water sources. It has been
575 noted that high lead concentration in soils can reduce vegetation growth and species richness
576 (Cao et al. 2003a; 2003b; Hardison et al. 2004; Goldsmith 2010).

577

578 Other forms of live-fire training involve the use of advanced high-power weaponry
579 including, but not limited to, artillery and mortars, multiple-launch rocket systems, hand
580 grenades, and anti-tank weapons (Rideout and Walsh 1990; Doxford and Judd 2002; Pekins

581 2006). These high-powered weapons require special training areas to safely contain the blast
582 radius and noise from civilian areas. This type of weapon training can create significant habitat
583 damage by cratering the terrain and altering the species composition within the area.
584 Specifically, these highly disturbed landscapes can suffer from degraded soil structure and
585 quality, and are reduced to disturbance-tolerant flora and fauna species (Fehmi et al. 2001; Smith
586 et al. 2002; Pekins 2006; Warren et al. 2007). Chemical contamination is also prevalent in these
587 training areas in the form of heavy metals, radiation (see Nuclear Warfare), and unused
588 propellants, all of which can directly impact community composition (Doxford and Judd 2002;
589 Edwards 2002; Garten et al. 2003). However, for most of these high-powered weapons, ‘dummy’
590 rounds (rounds containing less explosives and/or propellants) have been developed to lessen the
591 environmental impacts (Doxford and Judd 2002; Goldsmith 2010).

592
593 Armoured vehicles are considered all tracked and wheeled military vehicles used for
594 combat and transport (Johnson 1982) and are essential in most conflict situations due to their
595 long range firing capacity, protective armour, and all terrain maneuverability (Doxford and Judd
596 2002). These vehicles are generally outfitted with heavy armour and weaponry, making them
597 extremely heavy, with some vehicles weighing upwards of 60 metric tons. Due to the heavy
598 weight of these vehicles, terrain compaction is a significant issue which can have detrimental
599 impacts on the soil and vegetation communities (Lathrop 1983; Foster et al. 2006; Dickson et al.
600 2008). Armoured manoeuvre training is seen as being particularly damaging and persistent
601 (Doxford and Judd 2002), especially in fragile environments such as the Mojave Desert (Johnson
602 1982). The conditions for when armoured manoeuvre training occurs can also influence the
603 severity of the impact on the landscape; operations during wet spring conditions can cause

604 enlarged track ruts and higher rates of vegetation removal (Johnson 1983; Watts 1998; Dickson
605 et al. 2008). In frequently used landscapes, tracked vehicles have been noted to reduce total plant
606 and woody vegetation cover, and increase soil erosion rates (Johnson 1982; Wilson 1988).
607 Armoured manoeuvre training can also lead to changes in soil structure and chemistry with
608 frequently used sites having lower carbon to nitrogen ratios, as well as reduced soil carbon
609 content (Garten et al. 2003). Certain training exercises in wooded areas can be particularly
610 degrading on vegetation communities, as tracked vehicles can often be used as bulldozers to
611 clear paths and sight lines (Rideout and Walsh 1990). Armoured vehicle operations have also
612 been linked to incidentally hitting and killing wildlife during training exercises (Zakrajsek and
613 Bissonette 2005; Telesco and Van Manen 2006).

614
615 Aside from terrestrial armoured vehicle training, military training areas are intensively
616 used for fighter jet and helicopter training exercises (Black et al. 1984; Harrington and Veitch
617 1991; Conomy et al. 1998). The largest environmental impact associated with aviation exercises
618 is hitting and killing birds during flight manoeuvres (Richardson and West 2000; Civil Aviation
619 Authority 2001; Zakrajsek and Bissonette 2005). Bird-aircraft collisions are particularly serious
620 as they can often cause a loss of human life and damage/destruction to aircraft. From 1985-1998,
621 the United States Air Force (USAF) recorded an average of 2,700 aviation related bird strikes
622 each year, accumulating in excess of 35,000 bird-aircraft collisions over the 13 year period; an
623 average cost of \$35 million US dollars annually in aircraft repair and replacement to the USAF
624 (Zakrajsek and Bissonette 2005). The most vulnerable bird species to aircraft collisions noted by
625 the USAF included raptors, waterfowl, and passerines (Lovell and Dolbeer 1999; Zakrajsek and
626 Bissonette 2005). For all bird-aircraft collisions, it has been estimated that roughly 69% take

627 place below 305 m of altitude, which makes birds especially vulnerable to low flight training
628 exercises (Lovell and Dolbeer 1999; Civil Aviation Authority 2001; Zakrajsek and Bissonette
629 2005; Dukiya and Gahlot 2013). Due to the high risk of bird-aircraft collisions, special measures
630 have been taken at airstrips to reduce bird strike hazards. These precautionary measures include
631 reducing attractive installations near airfields (e.g. landfills or new water environments), altering
632 flight training routes, and using falconry to deter birds from the airfield vicinity (Cleary and
633 Dolbeer 1999; Lovell and Dolbeer 1999; Civil Aviation Authority 2001).

634

635 Naval military training exercises can have negative impacts on marine life. Unlike the
636 issues associated with over-pressure injuries from explosive detonations and live-fire operations
637 (see Nuclear Warfare, and Active Armed Conflict sections for further explanation), the main
638 impacts of naval training exercises are caused from the generation of excessive noise pollution
639 (Dolman et al. 2009). Noise pollution can be generated from a variety of sources including, but
640 not limited to, mechanical and propeller noise, gun discharges, explosives detonations, and the
641 use of sonar technologies (Parsons et al. 2000; Scott 2007; Dolman et al. 2009). The latter source
642 has received a lot of research attention and has been noted to negatively impact large marine
643 mammals in various ways (reviewed in Parsons et al. 2008). Active sonar systems range from
644 low-frequency levels, 1 Hz – 1 kHz, to mid-frequency levels, 1 – 10 kHz (Dolman et al. 2009).
645 When operational, both low- and mid-frequency systems emit high-intensity sound into the
646 ocean and listen for echoes that provide a sonic image of the ocean environment (Dolman et al.
647 2009). This type of imaging technology is highly useful for military operations but it can impact
648 the behaviour and survival of large marine mammals (Balcomb and Claridge 2001; Madsen
649 2005). Marine mammals rely on echolocation for most biological aspects of their lives, and the

650 use of sonar technologies has been linked to disrupting their signaling abilities. This can interfere
651 with foraging, reproduction, communication, and their predator detection abilities (Rendell and
652 Gordon 1999; Miller et al. 2000; Dolman et al. 2009). The use of sonar technology has also been
653 linked to mass stranding mortality events in cetacean species, most notably in beaked whales
654 (reviewed in Parsons et al. 2008) however, the causal mechanism of mortality from sonar is still
655 unknown (Dolman et al. 2009).

656

657 Dry troop training refers to dismounted infantry exercises and is widely practiced by
658 militaries around the world. This type of training can have a wide range of environmental
659 impacts determined by the size of the infantry and the nature of the exercise itself (Fehmi et al.
660 2001; Garten et al. 2003). Dismounted infantry can cause vegetation destruction, alter soil
661 structuring, and increase soil erosion from repetitive use of designated training areas
662 (Whitecotton et al. 2000; Warren et al. 2007). Realistic training requires infantry to dig defensive
663 positions for combat, and tent ditches for sleep and rest, further increasing soil erosion rates
664 (Trumbull et al. 1994; Fehmi et al. 2001). Dismounted infantry exercises can also negatively
665 affect wildlife distribution in active training areas where infantry presence can act to deter large
666 mammal species including black bears (*Ursus americanus*), mule deer (*Odocoileus hemionus*),
667 and coyotes (*Canis latrans*) (Stephenson et al. 1996; Telesco and Van Manen 2006). Although
668 wildlife avoidance of such activities reduces likelihood of direct mortality, the disturbance and
669 displacement can have sublethal consequences.

670

671 **Military Contamination**

672

673 Military conflict is associated with the testing, production, transportation, and
674 deployment of weapons. At each of these stages, there exists the potential for environmental
675 contamination (Dudley et al 2002; Machlis and Hanson 2008). In a warfare context, chemicals
676 can be manufactured for use in weapons to cause direct human mortality and/or to alter
677 landscapes to gain strategic tactical advantages that can expose the surrounding ecosystems to
678 potentially toxic compounds (Stellman et al. 2003; Ganesan et al. 2010; Westing 2013a).
679 Military activities also have the potential to indirectly contaminate the environment through
680 various by-products and spills associated with warfare as in the case of fuels and compounds
681 used in maintaining vehicle operation (Dudley et al 2002; Brady 1992; Machlis and Hanson
682 2008). Chemicals (in the broader sense), such as hydrocarbons and metals, can have immediate
683 destructive and toxic effects that may also persist for long periods of time in soil, water, and the
684 tissues of animals, all posing legacy issues. This section will aim to review how military actions
685 contribute to harmful chemical contamination at the different stages of warfare and their
686 subsequent effect on ecosystems with a particular focus on wildlife.

687

688 *Pre-War Contamination:*

689

690 Military chemical production and testing facilities require massive attention due to
691 hazardous waste accidents, spills, and dumping as the production of chemicals can be highly
692 volatile. These chemicals are required for the day-to-day operation of the military, as well as in
693 weapons development. In the United States, military training facilities and bases are responsible
694 for localized contamination from the dumping of chemicals directly into the environment causing
695 regional waterbodies, including drinking water sources, in the area to become toxic (Brady 1992;

696 Miller et al. 1998). Contaminated reservoirs on US army bases have caused the deaths of
697 thousands of waterfowl from drinking water on site (Lanier-Graham 1993). Similar pollution
698 conditions are present in Russia where dioxin pesticides have been disposed of improperly
699 resulting in soil and water contamination, thereby affecting the surrounding vegetation
700 negatively (Sidel 2000). Additionally, weapons testing, such as those done in Puerto Rico, Bikini
701 Atoll, and the United States, can result in significant soil, groundwater, and marine
702 contamination of chemicals and metals which may include mercury, iron, and plutonium. This
703 could have deleterious consequences to local vegetation and marine organisms in these regions
704 resulting in food chain disturbances (Donaldson 1997; Ortiz-Roque and Lopez-Rivera 2004;
705 Porter 2005; Machlis and Hanson 2008). All of these pre-war activities can lead to soil, water,
706 and vegetation contamination and have negative impacts on the wildlife that interacts with these
707 contaminated areas.

708

709 *Active Combat Contamination:*

710

711 Chemical warfare agents (CWAs) are weapons employed by the military to cause direct
712 human mortality (Ganesan et al. 2010). Many of the products developed as chemical warfare
713 agents have highly toxic and damaging properties intended for human targets, but may have
714 negative impacts on other species as well. These chemicals can fall under five main categories of
715 weapon effects: blistering agents that cause burning and blistering, nerve agents that target
716 neuron impulses, choking agents that affect the respiratory tract, blood agents that interrupt
717 oxygen absorption, and riot agents that cause immediate, short-term incapacitation (Ganesan et al
718 2010). Most chemical agents that can harm humans are toxic to other vertebrates and can injure

719 or kill some aquatic organisms at high concentrations. Often, these chemicals persist in plant
720 tissue resulting in developmental issues and can be potentially toxic to herbivores upon
721 consumption (Coppock 2009; Ganesan et al. 2010). Bullets and related debris (e.g. shell casings)
722 are often composed of materials that can be harmful to the ecosystem they are fired in. Lead, one
723 of the more commonly used metals in bullets and casings, has toxic properties that are highly
724 detrimental to a number of organ systems in vertebrates including the nervous system (Burger
725 and Gochfeld 2000; Papanikolaou et al. 2005). Leftover shells or fragments after combat can
726 result in accidental ingestion by many bird species who consume small particles inadvertently, or
727 as grit to aid in their digestion (Fisher et al. 2006). Depleted uranium shells/casings are also used
728 by some factions and can cause localized soil and sediment contamination (Haavisto et al. 2001;
729 Papastefanou 2002; Briner 2010). Uranium toxicity is of concern to exposed terrestrial and
730 freshwater plants, freshwater invertebrates and vertebrates, and mammals (Sheppard et al. 2005).
731 In mammals, uranium toxicity can be highly detrimental to development, brain chemistry,
732 behaviour and kidney function (Briner 2010).

733

734 Not all chemical warfare agents used are directly targeted at humans. Herbicides have
735 also been used, during combat operations, to alter landscapes and reduce foliage to enhance
736 visibility (Westing 1980; Stellman et al. 2003). Agent Orange, used during the Vietnam War
737 (1961-1971), was one of several types of dioxin-based herbicides sprayed by United States
738 forces to destroy crops and obstructing vegetation (Orians and Pfeiffer 1970; Westing 1980;
739 Westing 1984; Stellman et al. 2003). During this war, the landscapes in Vietnam, Cambodia, and
740 Laos were exposed to over 77 million litres of herbicides covering some 2600 million hectares of
741 land (Nguyen 2009). Over the past three to four decades, various studies have attempted to

742 evaluate environmental damage caused by these events and to assess their long term effects. In
743 doing so, it was apparent that the defoliation of the landscape resulted in immediate tree and
744 shrub mortality in addition to the local extirpation of many large mammals such as ungulates,
745 carnivores and elephants (Westing 1980; Oriands and Pfeiffer 1970; Westing 1985).

746

747 The application of large quantities of concentrated herbicides can alter the local
748 community structure as well. In Vietnam, forested and mangrove dominated habitats have
749 become scrubby grasslands, greatly changing the community assemblages (Dinh 1984; Westing
750 1989; Nguyen 2009). Surveys comparing un-impacted habitat with that inflicted with herbicide
751 found notably less species diversity (Westing 1989). However, one of the major limiting factors
752 in assessing and quantifying ecosystem changes is the lack of data in the region's baseline
753 ecological conditions (e.g. before war). In attempting to evaluate how biodiversity was affected,
754 researchers have made broad assumptions based off of limited observations and local indigenous
755 knowledge. Orians and Pfeiffer (1970) used these methods and suggested that regions of
756 Vietnam experienced a decline in bird species richness post conflict, specifically in those
757 consuming insects and fruit.

758

759 An additional long term problem associated with herbicide exposure is bioaccumulation
760 and the persistence of these chemicals in the environment. After the Vietnam War, high
761 concentrations of dioxins were found in the ovaries and livers of turtles (Schechter et al. 1989).
762 This effect was also demonstrated in tissues isolated from local pigs and chickens, likely
763 resulting from a combination of residual Agent Orange and other herbicidal exposure over the
764 past few decades (Schechter et al. 2006). More recently, the dioxin contamination still present in

765 the soils near the Bien Hoa Airbase (a “hotspot”) was discovered to fall within a high risk
766 category in terms of Canadian Environmental Quality Guidelines (Mai et al. 2007). The probable
767 effect level was 46x higher than the standard value for soil, even 30 years after the initial
768 chemical deployment illustrating a capacity of these chemicals to have chronic impacts on the
769 ecosystem.

770

771 Military activity is a highly mobile system occurring at multiple spatial scales (e.g.
772 nationally and internationally) that requires vast fuel and hydrocarbon resources that may
773 increase the possibility of oil and gas contamination. The Gulf War oil spill of 1991 resulted in
774 over 10 million m³ of oil and heavy metals intentionally dumped into the ocean (Box 4; Westing
775 2003) resulting in elevated bird mortalities and damage to important avian, mammalian and
776 reptilian migratory feeding habitats (Evans et al. 1993; Westing 2013b). Studies on benthic
777 invertebrates such as snails and clams immediately after the spill were found to have
778 significantly higher levels of Zn, Cu, and Ni in their tissues (Bu-Olayan and Subrahmanyam
779 1997). A decade after the spill, studies on the tissues of crabs showed high levels of Zn and Cu,
780 along with detectable levels of other heavy metals, demonstrating the persistence of these
781 compounds in the ecosystem (Al-Mohanna and Subrahmanyam 2001).

782

783 *Post-War Environmental Impacts Associated with Disposal:*

784

785 The long term effects of chemicals results from both their potential persistence and the
786 poor disposal programs of nations with stockpiled weapons. After WWII, CWAs such as
787 mustard gases, and arsenic poisons were packaged in barrels and directly disposed of in the

788 ocean (Chepesiuk 1997; Smith 2011); a common practice across the globe at the time. Disposal
789 of these vessels in the ocean runs the risk of the metal-based containers corroding and leaching
790 the chemical contents of the vessel into the ocean; an effect which could lead to a localized
791 exposure to the chemical as well as more widespread impacts via trophic movements (Long
792 2009). Sanderson et. al. (2010) modelled propagation of chemical warfare agents in the Baltic
793 Sea through the food chain in cod (*Gadus morhua*), herring (*Clupea harengus*), and sprat
794 (*Sprattus sprattus*). Adamsite, a component found in chemical weapons, was found to be
795 consistently present in the tissues of Atlantic cod demonstrating bioamplification and
796 accumulation of these substances in higher trophic levels. However, that study did not take into
797 account the number of buried munitions and containers on the seafloor that could reintroduce
798 high levels of chemicals to the surrounding area as the containers holding them degrade and
799 corrode. Regardless, this represents a pathway by which contaminants may be spread throughout
800 the various components of the ecosystem. Similarly, wreckages from naval ships pose certain
801 risks for the marine ecosystem in which they are found. Oil contamination in the Atlantic Ocean
802 due to World War II shipwrecks alone is estimated at over 15 million tonnes (Monfils 2005).
803 Much of the oil still resides within these wrecks and will pose future problems as the vessels
804 begin to degrade (Westing 1980; Monfils 2005). In much the same manner, during the conflict in
805 Kosovo, shelling of civilian infrastructure, namely manufacturing plants, resulted in a significant
806 but unintentional emission of industrial contaminants into the environment (Haavisto et al.
807 1999). Attention and care needs to be present during all stages of warfare, as contamination
808 events are common throughout training and active war with their effects persisting well after the
809 conflict has been resolved. Stringent policies are recognized as necessary to hold militaries
810 accountable for cleanup before training facilities can be returned to the public. Indeed, many

811 Western nations have adopted policies that require strict environmental management and concern
812 on home soil (Durrant 2007; Ramos et al. 2007). However, it should be noted that during war
813 outside of their respective countries, these policies are not necessarily followed.

814

815 **The Up-Shot: Technology**

816

817 One undeniable benefit that environmental and conservation science has reaped from
818 military research and development is the ability to utilize and refine resulting technological
819 advances. Military research and development teams share a common interest with ecological and
820 environmental researchers in needing to collect meaningful information more efficiently. An
821 exhaustive list of military developments used in everyday applications would include everything
822 from computing systems and the internet to nylon material that makes field equipment durable
823 and light-weight (Alic et al. 1992). However, there are a few notable technologies that have been
824 crucial to shaping modern ecological research. Satellites emerged over the course of the Cold-
825 War tensions between the United States and the Soviet Union (Alic et al. 1992; Slotten 2002)
826 and were followed closely by the creation of Global Positioning Systems (GPS) which allows for
827 high precision and accurate navigation (Parkinson 1996). Today, satellite imagery has paved the
828 way for the development of GIS spatial analyses, the backbone of evaluating large scale spatial
829 patterns and trends (Goodchild 2000). GIS permits investigators to relate spatially organized data
830 to other variables such as weather, animal abundance, or natural resource quantities. Remote
831 sensing technologies have military roots as well, and involve either passive or active gathering of
832 energy to help locate and identify objects (Turner et al. 2003). Electromagnetic energy, detected
833 by satellites, is commonly used to accomplish tasks such as assessing wildlife spatial distribution

834 and calculating species diversity (Turner et al. 2003). Similarly, RADAR technology actively
835 uses radio-waves to locate objects and obstructions with system advances being developed
836 between the British and American forces during the 1940s (Science News Letter 1945).
837 Currently, RADAR is one of the best methods for monitoring migratory bird species
838 (Gauthreaux and Belsier 2003). Remote Operated Vehicles (ROVs) including aerial drones,
839 marine vehicles, and terrestrial vehicles were originally developed by military organizations for
840 training operations, bomb recovery, and hostile terrain observations (Springer 2013). Now, aerial
841 ROVs are used in conservation to film and survey overhead parks, to monitor wildlife and to
842 look for illegal activity such as poaching and unauthorized logging (Sutherland et al. 2013;
843 Schiffman 2014). Marine ROVs have been employed to monitor marine life as well as being
844 used as a potential tool in gaining valuable insight into the system in question (Cohen 1995;
845 Jones 2009; Moura et al. 2013; Van Dover et al. 2014). Integration of autonomous technology
846 has huge advantages in ecological studies that are often limited by man-power and overwhelmed
847 by spatial scales; ROVs can efficiently extend work periods and area covered without human
848 intervention. Lastly, advances in telemetry technology by the military have greatly improved
849 conservation research through the miniaturization of tag components (e.g. batteries, transmitters,
850 etc.) for use in a wide number of biotelemetry projects which are useful in animal tracking and
851 monitoring (Cooke 2008; Benson 2010). Not only has this greatly improved the performance,
852 operation and capabilities of many of these devices (Cooke 2008) but it has permitted them to be
853 deployed on a greater number of species/weight classes as the mass of the tag has often been a
854 limiting factor in determining the lower size limit of their use (Olival and Higuchi 2006; Bridge
855 et al. 2011; Cooke et al. 2013).

856 **Research Gaps**

857 War is a perilous activity that makes for a poor research environment. Conflict zones,
858 restricted access military bases and warfare-induced hazardous material zones are often out of
859 reach of researchers attempting to assess war's impact on ecological functioning resulting in a
860 significant knowledge gap for current and post-conflict field sites. Additionally, because of the
861 stochastic nature of war (e.g. unknown when/where conflict and battles will occur), the
862 battlefield sites may not have pre-conflict information available thereby complicating before-
863 after impact analysis. To the extent possible, conducting research of military activities in a
864 before-after-control-impact framework would help to elucidate the environmental consequences
865 and thus reveal opportunities for mitigating negative effects while informing the development of
866 optimal strategies for rehabilitation and recovery. There also exists an apparent taxonomic bias in
867 the literature presented here whereby mammals, fish, and plants are often the studied
868 components of the impacted system presumably as a result of some perceived importance and/or
869 ease of access. Given that war is unlikely to be eliminated from society, the literature should
870 further expand to include other taxonomic representations and/or focus on species that are vital
871 to ecological functioning (e.g. keystone species, ecosystem engineers, etc.) in warfare impact
872 assessment. This could allow for the potential to expand our knowledge base significantly while
873 providing a potentially more streamlined approach that may be able to infer how overall
874 ecosystem functioning may be impaired/impacted under "conflict stress". It may be of relevance
875 and use to develop a mesocosm model system whereby the various impacts of warfare could be
876 modelled in a controlled environment to aid in developing an impact model at the whole
877 ecosystem level under a variety of climatic and environmental scenarios.

878 **Conclusions**

879

880 Given the information presented here, it is quite evident that warfare's impacts on
881 ecosystem functioning are indeed overwhelmingly deleterious. The impacts of conflict, nuclear
882 weapons, training operations and chemical contaminations all contribute to both reductions in the
883 populations of local flora and fauna as well as reducing species diversity in the affected
884 ecosystems. Impacts were demonstrated in a number of environments with a diversity of
885 taxonomic groups represented with war resulting in both acute and chronic impacts on the
886 ecosystem. A general overview of the impacts induced by the various aspects of war can be
887 found in Fig. 1. In some instances, warfare is a positive force in ecosystem functioning whereby
888 unintentional human exclusion provides refugia for a variety of species and, in some cases,
889 provides suitable habitat for endangered/threatened species. Some of these beneficial impacts are
890 illustrated in the Boxes 1-4. Additionally, research into developing military technology has
891 benefitted ecosystem functioning, indirectly, through providing a wide diversity of technological
892 tools and devices that are employed by many researchers involved with conservation and
893 ecological sciences. However, because of the inherent dangers of warfare and its seemingly
894 stochastic nature, research and assessments of military activities' impacts on the environment are
895 difficult to conduct and as such, the literature is limited in its scope. Moreover, new technologies
896 and militarily-unique substances continue to be developed and deployed such that the threats are
897 dynamic. With humanity continually engaging in war, the biosphere is likely to continue to
898 suffer. As such, this area of research should be continually pursued to attempt to better
899 understand war's impact on ecosystem structure and assist with developing potential mitigation
900 strategies to minimize negative consequences and implementing effective rehabilitation and
901 restoration approaches.

902 There seems to be little evidence that military strategists consider environmental
903 consequences of military activities when planning or executing military actions related to
904 conflict. We submit that there is much scope for proactive efforts to consider the environment
905 and biodiversity in formulating military plans. Yet, we also recognize that at the end of the day,
906 battlefield supremacy and achieving military objectives will likely continue to trump any and all
907 concerns related to the environment during active conflict (Westing 1986). The situation is
908 somewhat different for training facilities or other military installations during the preparatory and
909 readiness phases, at least in developed countries, where there is legal obligation to address
910 environmental concerns (e.g., contamination, endangered species; see Durant 2007).
911 Unfortunately, most warfare occurs in developing countries that tend to have unstable
912 governance structures where there is limited capacity for developing environmental policy or
913 addressing environmental issues that arise following conflict (Westing 1986). The policy
914 implications of warfare are beyond the scope of this article, but nonetheless, we wish to
915 emphasize greater need for global consideration of the environmental consequences of warfare
916 with efforts to develop policy instruments and agreements that consider the environment (*sensu*
917 Gasser 1995; Mrema et al. 2009) in that same way that attempts are made to consider aspects of
918 human welfare (e.g., war crimes tribunals, Geneva convention; see Drumbl 1998). The findings
919 of the synthesis presented here are clear – the consequences of modern warfare are
920 overwhelmingly negative for the environment and biodiversity. Indeed, although not quantified,
921 it is not unreasonable to think that modern warfare is one of the major forces associated with
922 environmental issues and biodiversity declines in some regions.

923

924 **Acknowledgments**

925 The authors would like to thank Jacqueline Chapman and Vivian Nguyen for their insight into
926 military environmental policies and procedures that helped shape this paper. SJ Cooke is
927 supported by NSERC and the Canada Research Chairs program. We thank Sofia M.R. Jain-
928 Schlaepfer, James Monaghan, Tanya Prystay, Arthur H. Westing and an anonymous reviewer for
929 kindly commenting on drafts of this manuscript.

930

931 **Contributions**

932 All authors contributed equally to the design, research, writing and editing of this work.

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947 **References**

948

949 Aarkrog, A. 1988. The radiological impact of the Chernobyl debris compared with that from
950 nuclear weapons fallout. *J. Environ. Radioactiv.* **6**:151-162.

951

952 Alic, J.A., Branscomb, L.M., Brooks, H., Carter, A.B., and Epstein, G.L. 1992. Beyond spinoff:
953 military and commercial technologies in a changing world. Harvard Business Press,
954 Waterdown, M.A., USA.

955

956 Al-Mohanna, S.Y., and Subrahmanyam, M.N.V. 2001. Flux of heavy metal accumulation in
957 various organs of the intertidal marine blue crab, *Portunus pelagicus* (L.) from the Kuwait
958 coast after the Gulf War. *Environ. Int.* **27**(4):321-326.

959

960 Alpen, E.L., and Sheline, G.E. 1954. The combined effects of thermal burns and whole body X-
961 irradiation on survival time and mortality. *Ann. Surg.* **140**(1):113-118.

962

963 Ampleman, G., Baucher, D., Thiboutot, S., Hawari, J., and Monteil-Rivera, F. 2004. Evaluation
964 of underwater contamination by explosives and metals at Point Amour Labrador and in the
965 Halifax Harbour area. Available from Defence Research and Development Canada,
966 Valcartier, Q.C., CA. No. DRDC-V-TR-2004-125

967

968 Apte, S., Holland, B.S., Godwin, L.S., and Gardner, J.P. 2000. Jumping ship: a stepping stone
969 event mediating transfer of a non-indigenous species via a potentially unsuitable
970 environment. *Biol. Invasions* **2**:75-79.

- 971
- 972 Augenbroe, G.L.M., and Pearce, A.R. 1998. Sustainable construction in the USA; a perspective
973 to the year 2010. Available from the Sustainable Development and the Future of
974 Construction. Vol. report 225.
- 975
- 976 Balcomb, K.C., and Claridge, D.E. 2001. A mass stranding of cetaceans caused by naval sonar in
977 the Bahamas. *Bahamas J. Sci.* **8**:1–12.
- 978
- 979 Barnthouse, L.W. 1995. Effects of ionizing radiation on terrestrial plants and animals: a
980 workshop report. Available from Oak Ridge National Lab, T.N., USA. ORNL/TM-13141
- 981
- 982 Barrett, M.J. 2011. Potentially polluting shipwrecks. Doctoral dissertation, School of the
983 Environment, Duke University, Durham, N.C., USA.
- 984
- 985 Bass, A.J., D'Aubrey, J.D., and Kistnasamy, N. 1973. Sharks of the east coast of southern Africa.
986 The genus *Carcharhinus* (*Carcharhinidae*). Available from Invest. Rep. Oceanogr. Res.
987 Inst., Durban, SA. pp.1-168.
- 988
- 989 Baxter, H., Drummond, J.A., Stephens-Newsham, L.G., and Randall, R.G. 1953. Reduction of
990 mortality in swine from combined total body radiation and thermal burns by streptomycin.
991 *Ann. Surg.* **137**(4):450-455.
- 992

- 993 Baxter, L., Hays, E.E., Hampson, G.R., and Backus, R.H. 1982. Mortality of fish subjected to
994 explosive shock as applied to oil well severance on Georges Bank. Available from the
995 Woods Hole Oceanographic Institution, M.A., USA. Technical Report WHOI-82-54
996
- 997 Beare, D., Hölker, F., Engelhard, G.H., McKenzie, E., and Reid, D.G. 2010. An unintended
998 experiment in fisheries science: a marine area protected by war results in Mexican waves
999 in fish numbers-at-age. *Naturwissenschaften* **97**:797-808.
1000
- 1001 Beasley, G., and Kneale, P. 2002. Reviewing the impacts of metals and PAHs on
1002 macroinvertebrates in urban watercourses. *Prog. Phys. Geogr.* **26**(2):236-270.
1003
- 1004 Beatley, J.C. 1966. Winter annual vegetation following a nuclear detonation in the Northern
1005 Mojave Desert (Nevada Test Site). *Radiat. Bot.* **6**:69-82.
1006
- 1007 Bednarek, A.T. 2001. Undamming rivers: a review of the ecological impacts of dam removal.
1008 *Environ. Manage.* **27**:803-814.
1009
- 1010 Bender, A. 1977. Noise impact on wildlife: an environmental impact assessment. *In Proceedings*
1011 *of the 9th Conference of Space Simulation, NASA (P-2000,7), Los Angeles, C.A., USA,*
1012 *26-28 April 1977. pp. 155-165.*
1013

- 1014 Benson, E. 2010. Wired wilderness: technologies of tracking and the making of modern wildlife.
1015 John Hopkins University Press, Baltimore, M.D., USA.
1016
- 1017 Berhe, A.A. 2007. The contribution of landmines to land degradation. *Land Degrad. Dev.* **18**:1-
1018 15.
1019
- 1020 Bickham, J.W., Hanks, B.G., Smolen, M.J., Lamb, T., and Gibbons, J.W. 1988. Flow cytometric
1021 analysis of the effects of low-level radiation exposure on natural populations of slider
1022 turtles (*Pseudemys scripta*). *Arch. Environ. Contam. Toxicol.* **17**(6):837-841.
1023
- 1024 Bizzozero Jr, O.J., Johnson, K.G., Ciocco, C., Hoshino, T., Itoga, T., Toyoda, S., and Kawasaki,
1025 S. 1966. Radiation-related leukemia in Hiroshima and Nagasaki, 1946–1964: distribution,
1026 incidence and appearance time. *New Engl. J. Med.* **274**(20):1095-1101.
1027
- 1028 Black, B.B., Collopy, M.W., Percival, H.F., Tiller, A.A., and Bohall, P.G. 1984. Effects of low
1029 level military training flights on wading bird colonies in Florida. Available from the
1030 Florida Cooperative Fish and Wildlife Research Unit, University of Florida, Gainesville,
1031 F.L., USA. Technical Report No. 7.
1032
- 1033 Brady, M.T. 1992. Environmental review of military base closures: implications for affected
1034 governments. *Duke Environ. Policy Forum* **2**(1):1-15.
1035

- 1036 Bridge, E.S., Thorup, K., Bowlin, M.S., Chilson, P.B., Diehl, R.H., Fléron, R.W., Hartl, P.,
1037 Kelly, J.F., Robinson, D., and Wikelski, M. 2011. Technology on the move: recent and
1038 forthcoming innovations for tracking migratory birds. *BioScience* **61**(9):689-698.
- 1039
1040
- 1041 Briner, W. 2010. The toxicity of depleted uranium. *Int. J. Environ. Res. Publ. Health* **7**(1):303-
1042 313.
- 1043
- 1044 Brode, H.L. 1968. Review of nuclear weapons effects. *Annu. Rev. Nucl. Sci.* **18**(1):153-202.
- 1045
- 1046 Brooks, J.W., Evans, E.I., Ham, W.T., and Reid, J.D. 1952. The influence of external body
1047 radiation on mortality from thermal burns. *Ann. Surg.* **136**:533-545.
- 1048
- 1049 Brown, D.G., Thomas, R.E., Jones, L.P., Cross, F.H., and Sasmore, D.P. 1961. Lethal dose
1050 studies with cattle exposed to whole-body Co-60 gamma radiation. *Radiat. Res.* **15**(5):675-
1051 683.
- 1052
- 1053 Bu-Olayan, A.H., and Subrahmanyam, M.N.V. 1997. Accumulation of copper, nickel, lead and
1054 zinc by snail, *Lunella coronatus* and pearl oyster, *Pinctada radiata* from the Kuwait coast
1055 before and after the gulf war oil spill. *Sci. Total Environ.* **197**(1):161-165.
- 1056
- 1057 Burger, J., and Gochfeld, M. 2000. Effects of lead on birds (*Laridae*): a review of laboratory and
1058 field studies. *J. Toxicol. Environ. Health B* **3**(2):59-78.

- 1059
- 1060 Byrnes, V.A., Brwon, D., Rose H.W., and Cibis, P.A. 1955. Chorio-retinal burns produced by
1061 atomic flashes. *A.M.A. Arch. Ophthal.* **53**(3):351-364.
- 1062
- 1063 Campbell, C.A., Valentine, J.W., and Ayala, F.J. 1975. High genetic variability in a population
1064 of *Tridacna maxima* from the Great Barrier Reef. *Mar. Biol.* **33**(4): 341-345.
- 1065
- 1066 Candole, C.A. 1967. Blast injury. *Canad. Med. Ass. J.* **96**:207-214.
- 1067
- 1068 Cao, X., Ma, L.Q., Chen, M., Hardison Jr, D.W., and Harris, W.G. 2003a. Weathering of lead
1069 bullets and their environmental effects at outdoor shooting ranges. *J. Environ. Qual.*
1070 **Q32**(2):526-534.
- 1071
- 1072 Cao, X., Ma, L.Q., Chen, M., Hardison Jr, D.W., and Harris, W.Q. 2003b. Lead transformation
1073 and distribution in the soils of shooting ranges in Florida, USA. *Sci Total Environ.* **307**(1-
1074 3):179-189.
- 1075
- 1076 Certini, G., Scalenghe, R., Woods, W.I. 2013. The impact of warfare on the soil environment.
1077 *Earth Sci. Rev.* **127**:1-15.
- 1078
- 1079 Chadwick, D. 1992. The fate of the elephant. Available from the Sierra Club, Washington, D.C.,
1080 USA.

- 1081
- 1082 Chandler, C.C., Storey, T.G., and Tangren, C.D. 1963. Prediction of forest fire spread following
1083 a nuclear explosion. Available from U.S. Forest Service Pacific Southwest Forest and
1084 Range Experiment Station, Albany, C.A., USA. Research Paper PSW-5.
- 1085
- 1086 Chepesuik, R. 1997. A Sea of Trouble? *B. Atom. Sci.* **53**(5):40-44.
- 1087
- 1088 Civil Aviation Authority. 2001. Large flocking birds: an international conflict between
1089 conservation and air safety [Online]. Available from
1090 http://www.caa.co.uk/docs/224/srg_dps_flockingbirds.pdf [Accessed April 15 2015].
- 1091
- 1092 Clark, B., and Jorgenson, A.K. 2012. The treadmill of destruction and the environmental impacts
1093 of Militaries. *Sociol. Compass* **6**:557-569.
- 1094
- 1095 Cleary, E.C., and Dolbeer, R.A. 1999. Wildlife hazard management at airports. Available from
1096 the Federal Aviation Administration, Office of Airport Safety and Standards, Washington
1097 D.C., USA.
- 1098
- 1099 Clodfelter, M. 2006. *The limits of air power: the American bombing of North Vietnam.*
1100 University of Nebraska Press, Lincoln, N.E., USA.
- 1101 Coates, P. 2013. From hazard to habitat (or hazardous habitat): the lively and lethal afterlife of
1102 Rocky Flats, Colorado. *Prog. Phys. Geogr.* **XX**(X):1–15.

- 1103
1104 Cohen, A.S. 1995. Paleoecological approaches to the conservation biology of benthos in ancient
1105 lakes: a case study from Lake Tanganyika. *J. N. Am. Benthol. Soc.* **14**(4):654-668.
1106
- 1107 Conomy, J.T., Collazo, J.A., Dubovsky, J.A., and Fleming, W.J. 1998. Dabbling duck behavior
1108 and aircraft activity in coastal North Carolina. *J. Wildlife Manage.* **62**(3):1127-1134.
1109
- 1110 Cooke, S.J. 2008. Biotelemetry and biologging in endangered species research and animal
1111 conservation: relevance to regional, national, and IUCN Red List threat assessments.
1112 *Endanger. Species Res.* **4**:165-185.
1113
- 1114 Cooke, S.J., Midwood, J.D., Thiem, J.D., Klimley, P., Lucas, M.C., Thorstad, E.B., Eiler, J.,
1115 Holbrook, C., and Ebner, B.C. 2013. Tracking animals in freshwater with electronic tags:
1116 past, present and future. *Anim. Biotelem.* **1**(5):1-19.
1117
- 1118 Coppock, R.W. 2009. Threats to wildlife by chemical warfare agents. *In Handbook of*
1119 *Toxicology of Chemical Warfare Agents. Edited by R.C. Gupta. Academic Press,*
1120 *Waltham, M.E., USA. pp. 747-751.*
1121
- 1122 Craft, T.F. 1964. Effects of nuclear explosions on watersheds. *J. Am. Water Works Assoc.*
1123 **56**(7):846-852.
1124

- 1125 Darby, S.C., Kendall, G.M., Fell, T.P., O'Hagan, J.A., Muirhead, C.R., Ennis, J.R., Ball, A.M.,
1126 Dennis, J.A., and Doll, R. 1988. A summary of mortality and incidence of cancer in men
1127 from the United Kingdom who participated in the United Kingdom's atmospheric nuclear
1128 weapon tests and experimental programmes. *Brit. Med. J.* **296**(6618):332-338.
1129
- 1130 Davis, J.S. 2007. Scales of Eden: conservation and pristine devastation on Bikini Atoll. *Environ.*
1131 *Plan. D* **25**(2):213-235.
1132
- 1133 Demarais, S., Tazik, D. J., Guertin, P. J., and Jorgensen, E. E. 1999. Disturbance associated with
1134 military exercises. *In Ecosystems of Disturbed Ground. Edited by L.R. Walker.* Elsevier
1135 Science B.V., Amsterdam, NL. pp. 385-396.
- 1136 Dickson, T.L., Wilsey, B.J., Busby, R.R., and Gebhart, D.L. 2008. Grassland plant composition
1137 alters vehicular disturbance effects in Kansas, USA. *Environ. Manage.* **41**:676-684.
1138
- 1139 Dinh, H. 1984. Long-term changes in dense inland forest following herbicidal attack. *In*
1140 *Herbicides in war: the long term ecological and human consequences. Edited by A.H.*
1141 *Westing.* Taylor & Francis, London, UK. pp. 31-32.
1142
- 1143 Dolman, S.J., Weir, C.R., and Jasny, M. 2009. Comparative review of marine mammal guidance
1144 implemented during naval exercises. *Mar. Pollut. Bull.* **58**:465-477.
1145

- 1146 Donaldson, L.R., Seymour, A.H., and Nevissi, A.E. 1997. University of Washington's
1147 radioecological studies in the Marshall Islands, 1946-1977. *Health Phys.* **73**(1):214-222.
1148
- 1149 Downing N., and Roberts C. 1993. Has the Gulf War affected coral reefs of the Northwestern
1150 Gulf? *Mar. Pollut. Bull.* **27**:149-156.
1151
- 1152 Doxford, D., and Judd, A. 2002. Army training: the environmental gains resulting from the
1153 adoption of alternatives to traditional training methods. *J. Environ. Plann. Man.* **45**(2):245-
1154 265.
1155
- 1156 Draulans, D., Van Krunkelsven, E. 2002. The impact of wars on the Democratic Republic of
1157 Congo. *Oryx* **36**:35-40.
- 1158 Drumbl, M.A. 1998. Waging war against the world: the need to move from war crimes to
1159 environmental crimes. *Fordham Int'l L. J.* **22**:122-153.
1160
- 1161 Dubey, A., and Shreni, P.D. 2008. War and environment: an overview. *J. Environ. Res. Dev.*
1162 **4**:968-976.
1163
- 1164 Dudley, J.P., Ginsberg, J.R., Plumptre, A.J., Hart, J.A., and Campos, L.C. 2002. Effects of war
1165 and civil strife on wildlife and habitats. *Cons. Biol.* **16**(2): 319-329.
1166

- 1167 Dufour, P.A. 1980. Effects of noise on wildlife and other animals: review of research since 1971.
1168 Available from the U.S. Environmental Protection Agency. EPA 550/9-80-100:1-97.
1169
- 1170 Dukiya, J.J., and Gahlot, V. 2013. An evaluation of the effect of bird strikes on flight safety
1171 operations at international airports. *Int. J. Traffic Transport Eng.* **3**(1):16-33.
1172
- 1173 Dunnet, G.M. 1977. Observations on the effects of low-flying aircraft at seabird colonies on the
1174 coast of Aberdeenshire, Scotland. *Biol. Conserv.* **12**:55-63.
1175
- 1176 DuPont Guerry III, W.T., Ham, W.T., Wiesinger, H., Schmidt, F.H., Williams, R.C., Ruffin,
1177 R.S., and Shaffer, M.C. 1956. Experimental production of flash burns in the rabbit
1178 retina. *T. Am. Ophthalmol. Soc.* **54**:259-273.
1179
- 1180 Durant, R.F. 2007. The greening of the US military: environmental policy, national security, and
1181 organizational change. *Edited by* B.A. Radin. Georgetown University Press, Washington,
1182 D.C., USA.
1183
- 1184 Edwards, C.A. 2002. Assessing the effects of environmental pollutants on soil organisms,
1185 communities, processes and ecosystems. *Euro. J. Soil Biol.* **38**(3-4):225-231.
1186
- 1187 Eisenbud, M., and Gesell, T.F. 1997. Nuclear weapons. *In* Environmental Radioactivity from
1188 Natural, Industrial & Military Sources. Academic Press, San Diego, C.A., USA, pp. 271-
1189 314.

1190

1191 Eldred, E., and Trowbridge, W.V. 1954. Radiation Sickness in the monkey 1. Radiology
1192 **62**(1):65-73.

1193

1194 Engelhard, G.H. 2008. One hundred and twenty years of change in fishing power of English
1195 North Sea trawlers. *In* Advances in fisheries science 50 years on from Beverton and Holt.
1196 *Edited by* A. Payne, J. Cotter and T. Potter. Blackwell Publishing, Oxford, UK. pp.1-15.

1197

1198 Entry, J.A., and Watrud, L.S. 1998. Potential remediation of ¹³⁷Cs and ⁹⁰Sr contaminated soil by
1199 accumulation in alamo switchgrass. *Water Air Soil Poll.* **104**(3-4):39-352.

1200

1201 Evans, M.I., Symens, P., and Pilcher, C.W.T. 1993. Short-term damage to coastal bird
1202 populations in Saudi Arabia and Kuwait following the 1991 Gulf War marine pollution.
1203 *Mar. Pollut. Bull.* **27**:157–161.

1204

1205 Fehmi, J.S., Farmer, T., and Zimmerman, A. 2001. Impacts of military vehicle training activities
1206 on vegetation: bibliography with abstracts. Available from the US Army Engineer
1207 Research and Development Center, Vicksburg, MI, USA. pp 1-50.

1208

1209 Fisher, I.J., Pain, D.J., and Thomas, V.G. 2006. A review of lead poisoning from ammunition
1210 sources in terrestrial birds. *Biol. Conserv.* **131**:421-432.

1211

1212 Fosberg, F.R. 1985. Vegetation of Bikini Atoll. *Atoll Res. Bull.* **315**:1-28.

- 1213
- 1214 Foster, J.R., Ayers, P.D., Lombardi-Przylowicz, A.M., and Simmons, K. 2006. Initial effects of
1215 light armored vehicle use on grassland vegetation at Fort Lewis, Washington. *J. Environ.*
1216 *Manage.* **81**(4):315-322.
- 1217
- 1218 Francis, R.A. 2011. The impacts of modern warfare on freshwater ecosystems. *Environ. Manage.*
1219 **48**:985-999.
- 1220
- 1221 Ganesan, K., Raza, S.K., and Vijayaraghavan, R. 2010. Chemical warfare agents. *J*
1222 *Pharm Bioallied Sci.* **2**(3):166-178.
- 1223
- 1224 Gangwar, A. 2003. Impact of war and landmines on environment. *In* Forum Proceedings:
1225 Landmines-Challenges to Humanity and Environment. Srinigar, IN, 20 April 2003.
- 1226
- 1227 Garten, C.T., Ashwood, T.L., and Dale, V.H. 2003. Effect of military training on indicators of
1228 soil quality at Fort Benning, Georgia. *Ecol. Indic.* **3**:171-179.
- 1229
- 1230 Gaspin, J.B. 1975. Experimental investigations of the effects of underwater explosions on
1231 swimbladder fish. I. 1973 Chesapeake Bay tests. Available from the Naval Surface
1232 Weapons Center, White Oak Lab, Silver Spring, M.D., USA. No. NSWC/WOL/TR-75-58.
- 1233
- 1234 Gasser, H.P. 1995. For better protection of the natural environment in armed conflict: a proposal
1235 for action. *Am. J. Int'l L.* **1995**: 637-644.

- 1236
- 1237 Gauthreaux, S.A. Jr., and Belsier, C.G. 2003. Radar ornithology and biological conservation. The
1238 *Auk* **120**(2):266-277.
- 1239
- 1240 Gazenbeek, A. 2005. LIFE, Natura 2000 and the military. Available from the European
1241 Commission, Environment Directorate General, Brussels, BE.
- 1242
- 1243 Gese, E.M., Rongstad, O.J., and Mytton, W.R. 1989. Changes in coyote movements due to
1244 military activity. *J. Wildlife Manage.* **53**(2):334-339.
- 1245
- 1246 Gladwin, D.N., Asherin, D.A., and Mancini, K.M. 1988. Effects of aircraft noise and sonic booms
1247 on fish and wildlife: results of a survey of U.S. fish and wildlife service endangered species
1248 and ecological services field offices, refuges, hatcheries, and research centers. Available
1249 from the US Departments of the Interior, Fish and Wildlife Services. NERC-88/30
- 1250
- 1251 Glasstone, G. 1962. The effects of nuclear weapons (Revised Edn.). U.S. Department of Defense
1252 and Energy, Washington D.C., USA.
- 1253
- 1254 Glasstone, G., and Dolan, P.J. 1977. The effects of nuclear weapons (3rd Edn.). U.S. Department
1255 of Defence and Energy, Washington D.C., USA.
- 1256
- 1257 Glasstone, S. 1964. The effects of nuclear weapons. Available from the US Department of
1258 Defense, Washington D.C., USA. OTSI identification 972902.

- 1259
- 1260 Gleick, P.H. 1993. Water and conflict: fresh water resources and international security. *Int.*
1261 *Secur.* **18**:79–112.
- 1262
- 1263 Goertner, J.F. 1982. Prediction of underwater explosion safe ranges for sea mammals. Available
1264 from the Naval Surface Weapons Center, Silver Spring, M.D., USA, Report No.
1265 NSWC/TR-82-188.
- 1266
- 1267 Goldizen, V.C., Richmond, D.R., Chiffelle, T.L., Bowen, I.G., and White, C.S. 1961. Missile
1268 studies with a biological target. Available from the Office of Technical Services,
1269 Department of Commerce, Washington, D.C., USA. USAEC Civil Effects Test Group
1270 Report WT-1470.
- 1271
- 1272 Goldsmith, G.S. 2010. Environmental impacts and military range use: an investigation and
1273 summary of what we have learned after 12 years at Massachusetts Military Reservation
1274 (MMR) and implications for the continued use of military ranges in the United States.
1275 Available from the Army Environmental Policy Institute. pp 1-36.
- 1276
- 1277 Gontier, M. 2007. Scale issues in the assessment of ecological impacts using a GIS-based habitat
1278 model - A case study for the Stockholm region. *Environ. Impact Assess. Rev.* **27**:440-459.
- 1279
- 1280 Goodchild, M.F. 2000. The current status of GIS and spatial analysis. *J. Geogr. Syst.* **2**:5-10.
- 1281

- 1282 Govoni J.J., West, M.A., Settle, L.R., Lynch, R.T., and Greene, M.D. 2008. Effects of
1283 underwater explosions on larval fish: implications for a coastal engineering project. J.
1284 Coast. Res. **24**(2B):228-233.
- 1285
- 1286 Grace, J.B., and Keeley, J.E. 2006. A structural equation model analysis of postfire plant
1287 diversity in California shrublands. Ecol. Appl. **16**(2):503-514.
- 1288
- 1289 Gray, R.H., and Rickard, W.H. 1989. The protected area of Hanford as a refugium for native
1290 plants and animals. Environ. Conserv. **16**(03):251-260.
- 1291
- 1292 Gulland, J.A. 1968. Recent changes in the North Sea plaice fishery. J. Cons. Int. Explor. Mer.
1293 **31**:305-322.
- 1294
- 1295 Guskova, A.K., Baranov, A.E., and Gusev, I.A. 2001. Acute radiation sickness: underlying
1296 principles and assessments. *In* Medical Management of Radiation Accidents (2nd Edn.).
1297 *Edited by* I. Gusev, A.K. Guskova and F.A. Mettler. CRC Press, Boca Raton, F.L., USA.
1298 pp 33-52.
- 1299
- 1300 Haavisto, P. et al. 1999. The Kosovo conflict: consequences for the environment & human
1301 settlements. Available from UN Environment Programme, Nairobi, KE.
- 1302

- 1303 Haavisto, P. et al. 2001. Depleted Uranium in Kosovo: Post-conflict Environmental Assessment.
1304 Available from UN Environment Programme, Nairobi, KE.
1305
- 1306 Ham, W.T., Wiesinger, H., Guerry, D., Schmidt, F.H., Williams, R.C., Ruffin, R.S., and Shaffer,
1307 M.C. 1957. Experimental production of flash burns in the rabbit retina. *Am. J. Ophthalmol.*
1308 **43**(5):711-718.
1309
- 1310 Hanson, T., Brooks, T.M., Da Fonseca, G.A., Hoffmann, M., Lamoreux, J.F., Machlis, G., and
1311 Pilgrim, J.D. 2009. Warfare in biodiversity hotspots. *Conserv. Biol.* **23**:578-587.
1312
- 1313 Hardison Jr, D.W., Ma, L.Q., Luongo, T., and Harris, W.G. 2004. Lead contamination in
1314 shooting range soils from abrasion of lead bullets and subsequent weathering. *Sci. Total*
1315 *Environ.* **328**(1-3):175-183.
1316
- 1317 Harrington, F.H., and Veitch, A.M. 1991. Short-term impacts of low-level jet fighter training on
1318 caribou in Labrador. *Arctic* **44**(4):318-327.
1319
- 1320 Hart, T., Hart, J., Fimbel, C., Fimbel, R., Laurance, W.F., Oren, C., Struhsaker, T.T.,
1321 Rosenbaum, H.C., Walsh, P.D., Razafindrakoto, Y., Vely, M., and DeSalle, P. 1997.
1322 Conservation and civil strife: two perspectives from Central Africa Conservation and civil
1323 strife. *Conserv. Biol.* **11**:308-314.
1324

- 1325 Hatch, F.T., Mazarimas, J.A., Greenway, G.G., Koranda, J.J., and Moore, J.L. 1968. Studies on
1326 liver DNA in tritiated kangaroo rats living at Sedan Crater. Available from California
1327 University, Lawrence Radiation Lab, Livermore, C.A., USA. Technical Report No. UCRL-
1328 50481.
- 1329
- 1330 Hatch, T., Mazrimas, J.A., Koranda, J.J., and Martin, J.R. 1970. Ecology and radiation exposure
1331 of kangaroo rats living in a tritiated environment. *Radiat. Res.* **44**(1):97-107.
- 1332
- 1333 Havlick, D.G. 2014. Opportunistic conservation at former military sites in the United States.
1334 *Prog. Phys. Geogr.* **38**(3): 71-285.
- 1335
- 1336 Hieb, M.R., Mackay, S., Powers, M.W., Yu, H., Kleiner, M., and Pullen, J.M. 2007. Geospatial
1337 challenges in a net centric environment: actionable information technology, design, and
1338 implementation. *In* Defense and Security Symposium (657816), International Society for
1339 Optics and Photonics. pp. 1-12.
- 1340
- 1341 Higuchi, H., Ozaki, K., Fujita, G., Minton, J., Ueta, M., Soma, M., and Mita, N. 1996. Satellite
1342 tracking of white-naped crane migration and the importance of the Korean demilitarized
1343 zone. *Conserv. Biol.* **10**:806-812.
- 1344
- 1345 Hinshaw, J.R. 1968. Early changes in the depth of burns. *Ann. N.Y. Acad. Sci.* **150**(3):548-553.
- 1346

- 1347 Houk, P., and Musburger, C. 2013. Trophic interactions and ecological stability across coral
1348 reefs in the Marshal Islands. *Mar. Ecol. Prog. Ser.* **488**:23-34.
1349
- 1350 Hunter, R. 1991. *Bromus* invasions on the Nevada Test Site: present status of *B. rubens* and *B.*
1351 *tectorum* with notes on their relationship to disturbance and altitude. *Great Basin Nat.*
1352 **51**(2):176-182.
1353
- 1354 Hunter, R.B. 1992. Status of flora and fauna on the Nevada test site, 1988. Available from the
1355 United States Department of Energy Nevada Field Office, N.V., USA. Report No.
1356 DOE/NV/10630-29.
1357
- 1358 Hupy, J.P. 2006. The long-term effects of explosive munitions on the WWI battlefield surface of
1359 Verdun, France. *Scot. Geogr. Mag.* **122**:167-184.
1360
- 1361 Hupy, J.P. 2008. The environmental footprint of war. *Environ. Hist.* 405-421.
1362
- 1363 Hynes, M.V., Peter, J.E., and Rushworth, D. 2004. Artificial Reefs: A disposal option for Navy
1364 and MARAD Ships. Available from the Rand National Defense Research Institute, Santa
1365 Monica, C.A., USA. No. Rand/DB-391-Navy
1366
- 1367 ICRC. 2008. How is the term “armed conflict” defined in international humanitarian law?
1368 Available from the International Committee of the Red Cross. Opinion paper. pp.1-5.

1369

1370 Isakason, J.S. 1974. Biological effects of underground nuclear testing on marine organisms. II.

1371 Observed effects of Amchitka Island, Alaska, tests on marine fauna. *In* Proceedings of the

1372 First Conference on the Environmental Effects of Explosives and Explosions (Technical

1373 Report 73-223). Naval Ordnance Laboratory, White Oak, M.D., USA. pp. 86-97.

1374

1375 Jaffin, J.H., McKinney, L., Kinney, R.C., Cunningham, J.A., Mortiz, D.M., Kraimer, J.M.,

1376 Graeber, G.M., Moe, J.B., Salander, J.M., and Harmon, J.W. 1987. A laboratory model for

1377 studying blast overpressure injury. *J. Trauma* **27**(4):349-356.

1378

1379 Janssen, R. 1980. Future scientific activities in effects of noise on animals. *In* Proceedings of the

1380 Third International Congress on Noise as a Public Health Problem, American Speech-

1381 Language-Hearing Association Rockville, M.D., USA, 1980. *Edited by* J.V. Tobias, G.

1382 Jansen, and W.D. Ward. pp.632-637.

1383

1384 Johnson, F.L. 1982. Effects of tank training activities on botanical features at Fort Hood, Texas.

1385 *Southwest. Nat.* **27**(3):309-314.

1386

1387 Jones, D.O. 2009. Using existing industrial remotely operated vehicles for deep-sea science.

1388 *Zool. Scr.* **38**(s1):41-47.

1389

1390 Jorgensen, C.D., and Hayward, C.L. 1965. Mammals of the Nevada test site. Brigham Young

1391 *Univ. Sci. Bull.* **6**(3):1-81.

- 1392
- 1393 Kajitani, N., and Hatano, S. 1953. A-bomb disaster investigation report (Hiroshima) first and
1394 second investigations. *In* Research on the Effects of the Atomic Bomb, Part 1. *Edited by*
1395 the Japan Science Council. pp. 522-601.
- 1396
- 1397 Kaufman, D.W., Finck, E.J., and Kaufman, G.A. 1990. Small mammals and grassland fires. *In*
1398 Fire in North American Tallgrass Prairies. *Edited by* S.L. Collins and L.L. Wallace.
1399 University of Oklahoma Press, Norman, Oklahoma, USA, pp. 46-80.
- 1400
- 1401 Kazmarek, S.E.A., Gray, P., and Booher, A. 2005. Environmental issues at military installations:
1402 what they are and how they're handled. Available from the Environmental Advisory. pp 1-
1403 9.
- 1404
- 1405 Ketten, D.R. 1995. Estimates of blast injury and acoustic trauma zones for marine mammals
1406 from underwater explosions. *In* Sensory Systems of Aquatic Mammals. *Edited by* R.A.
1407 Kastelein, J.A. Thomas and P.E. Nachtigall. De Spil Publishers, Woerden, NE. pp 391-
1408 407.
- 1409
- 1410 Keevin, T.M., and Hempen, G.L. 1997. The environmental effects of underwater explosions with
1411 methods to mitigate impacts. Available from the US. Department of the Army Corps of
1412 Engineers St. Louis, MO, USA
- 1413

- 1414 Kim, K.C. 1997. Preserving biodiversity in Korea's demilitarized zone. *Science* **278**:242-243.
- 1415
- 1416 Kirkwood, J. B. 1970. Bioenvironmental safety studies, Amchitka Island, Alaska. Milrow D+2
1417 Months Report. Amchitka Bioenvironmental Program. Available from the Battelle
1418 Memorial Institute, Columbus Laboratories, Columbus, O.H., USA. Report No. BHI-171-
1419 126.
- 1420
- 1421 Kirkwood, J.B., and Fuller, R.G. 1972. Amchitka Bioenvironmental Program: Bioenvironmental
1422 Safety Studies, Amchitka Island, Alaska: Cannikin D+2 Months Report. Available from
1423 the Battelle Memorial Institute, Columbus Laboratories, Columbus, O.H., USA. Report
1424 No. BMI-171-147,
- 1425
- 1426 Kishi, H. 2000. Effects of the “special bomb”: recollections of a neurosurgeon in Hiroshima,
1427 August 8-15, 1945. *Neurosurgery* **47**(2):441-446.
- 1428
- 1429 Klein, D.R. 1973. The reaction of some northern mammals to aircraft disturbance. *In* 11th
1430 International Congress of Game Biology, Stockholm, SE, 3-7 September 1973. pp. 377-
1431 383.
- 1432
- 1433 Kolesnikova, A.A., Taskaeva, A.A., Krivolutskii, and Taskaev, A.I. 2005. Condition of the soil
1434 fauna near the epicentre of an underground nuclear explosion in the Northern Urals. *Russ.*
1435 *J. Ecol.* **36**(3):150-157.
- 1436

- 1437 Kopel, D., Malkinson, D., and Wittenberg, L. 2015. Characterization of vegetation community
1438 dynamics in areas affected by construction waste along the urban fringe. *Urban Ecosys.*
1439 **18**:133-150.
- 1440
- 1441 Lacoste, Y.1973. An illustration of geographical warfare: bombing the dikes on the Red River,
1442 North Vietnam. *Antipode* **5**:1–13.
- 1443
- 1444 Lamb, T., Bickham, J.W., Gibbons, J.W., Smolen, M.J., and McDowell, S. 1991. Genetic
1445 damage in a population of slider turtles (*Trachemys scripta*) inhabiting a radioactive
1446 reservoir. *Arch. Environ. Contam. Toxicol.* **20**(1):138-142.
- 1447
- 1448 Lanier-Graham, S. 1993. The ecology of war: environmental impacts of weaponry and warfare.
1449 Walker and Company, New York, N.Y., USA.
- 1450
- 1451 Larkin, R.P., Pater, L.L., and Tazik, D.J. 1996. Effects of military noise on wildlife. A literature
1452 review. Available from the Construction Engineering Research Lab (Army), Champaign,
1453 I.L., USA. No. USACERL-TR-96/21
- 1454
- 1455 Lathrop, E.W. 1983. Recovery of perennial vegetation in military maneuver areas. *In*
1456 *Environmental Effects of Off-Road Vehicles: Impacts and Management in Arid Regions.*
1457 *Edited by* R.H. Webb and H.G. Wilshire. Springer, NL. pp 265-277.
- 1458

- 1459 Ledney, G.D., Elliot, T.B., and Moore, M.M. 1992. Modulation of mortality by tissue trauma and
1460 sepsis in mice after radiation injury. *In* The Biological Basis of Radiation Protection
1461 Practice. *Edited by* K.L. Mossan and W.A. Mills. Williams and Wilkens, Baltimore, MD,
1462 USA, pp. 202-217.
- 1463
- 1464 Leitenberg, M. 2006. Deaths in Wars and Conflicts in the 20th Century. 3rd ed. Available from
1465 the Cornell University Peace Studies Program, Ithaca, N.Y., USA. ISSN 1075-4857
- 1466 Levy, B.S., Lee, C., and Shahi, B.S. 1997. The environmental consequences of war. *In* War and
1467 Public Health. *Edited by* B.S. Levy and V.W. Sidel. Oxford University Press New York,
1468 N.Y., USA. pp. 51-62.
- 1469
- 1470 Liebow, A.A. 1983. Encounter with disaster: a medical diary of Hiroshima, 1945. *Yale J. Biol.*
1471 *Med.* **56**:23-38.
- 1472
- 1473 Lifton, R.J. 1967. Death in life: survivors of Hiroshima. Random House, New York, NY, USA.
- 1474
- 1475 Lindenmayer, D.B., Wood, J.T., Cunningham, R.B., MacGregor, C., Crane, M., Michael, D.,
1476 Montague-Drake, R., Brown, D., Muntz, R., and Gill, A.M. 2008. Testing hypotheses
1477 associated with bird response to wildfire. *Ecol. Appl.* **18**(8):1967-1983.
- 1478
- 1479 Lindsell, J. A., Klop, E., and Siaka, A. M. 2011. The impact of civil war on forest wildlife in
1480 West Africa: mammals in Gola Forest, Sierra Leone. *Oryx* **45**(01):69-77.

- 1481
- 1482 Long, T. P. 2009. A global prospective on underwater munitions. *Mar. Technol. Soc. J.* **43**(4):
- 1483 5-10.
- 1484
- 1485 Lovell, C.D., and Dolbeer, R.A. 1999. Validation of the United States air force bird avoidance
- 1486 model. *Wildlife Soc. Bull.* **27**(1):167-171.
- 1487
- 1488 Machlis, G.E., and Hanson, T. 2008. Warfare ecology. *BioScience* **58**:729–736.
- 1489
- 1490 Madsen, P.T. 2005. Marine mammals and noise: problems with root mean square sound pressure
- 1491 levels. *J. Acoust, Soc. Am.* **117**: 3952–3957.
- 1492
- 1493 Mai, T.A., Doan, T.V., Tarradellas, J., de Alecastra, L.F., and Gradnjean, D. 2007. Dioxin
- 1494 contamination in soils of Southern Vietnam. *Chemosphere* **67**:1802-1807.
- 1495
- 1496 Majeed, A. 2004. The impact of militarism on the environment: an overview of direct and
- 1497 indirect effects. Available from the Physicians for Global Survival, Ottawa, O.N., CA.
- 1498 ISBN 0-9735916-0-9
- 1499
- 1500 Mancini, K.M., Gladwin, D.N., Villella, R., and Cavendish, M.G. 1988. Effects of aircraft noise
- 1501 and sonic booms on domestic animals and wildlife: a literature synthesis. Available from
- 1502 National Ecology Research Center, Fort Collins, C.O., USA. No. NERC 88/29
- 1503

- 1504 Martore, R.M., Mathews, T.D., and Bell, M. 1998. Levels of PCBs and heavy metals in biota
1505 found on ex-military ships used as artificial reefs. Available from the South Carolina
1506 Department of Natural Resources Charleston, S.C., USA.
1507
- 1508 Masco, J. 2004. Mutant ecologies: radioactive life in post-cold war New Mexico. *Cult.*
1509 *Anthropol.* **19**(4):517-550.
1510
1511
- 1512 Mayorga, M.A. 1997. The pathology of primary blast overpressure injury. *Toxicology* **121**: 17-
1513 28.
1514
- 1515 McDonnel, G.M., Crosby, W.H., Tessmer, C.F., Moncrief, W.H., Baker, H.J., Goldstein, J.D.,
1516 Woodward, K., Shively, J.N., Daniell, H.W., Horava, A., and Claypool, H.A. 1961. Effects
1517 of nuclear detonations on a large biological specimen (swine). Available from the Defense
1518 Atomic Support Agency, Sandia Base, Albuquerque, N.M., USA. Operation Plumbbob,
1519 Project 4.1 Report WT-1428
1520
- 1521 McNeely, J.A. 2003. Conserving forest biodiversity in times of violent conflict. *Oryx* **37**:142-
1522 152.
1523
- 1524 Medica, P. A., Turner, F.B., and Smith, D.D. 1973. Effects of radiation on a fenced population of
1525 horned lizards (*Phrynosoma platyrhinos*) in southern Nevada. *J. Herpetol.* **7**:79-85.
1526

- 1527 Merritt, M. 1970. Physical and biological effects: Milrow Event. Available from the U.S. Atomic
1528 Energy Commission, National Technical Information Service, U.S. Dep. Of Commerce,
1529 Springfield, V.A., USA. Report No. NVO-79.
1530
- 1531 Merritt, M. 1973. Physical and biological effects: Cannikin. Available from the U.S. Atomic
1532 Energy Commission, National Technical Information Service, U.S. Dep. of Commerce,
1533 Springfield, V.A., USA. Report No. NVO-123
1534
- 1535 Mettler, F.A. 2001. Direct effects of radiation on specific tissues. *In* Medical Management of
1536 Radiation Accidents (2nd Edn.). *Edited by* I. Gusev, A.K. Guskova and F.A. Mettler. CRC
1537 Press, Boca Raton, F.L., USA. pp 69-132.
1538
- 1539 Meyer-Reil, L., and Köster, M. 2000. Eutrophication of marine water: effects on benthic
1540 microbial communities. *Mar. Pollut. Bull.* **41**(1-6):255-263.
1541
- 1542 Miller, J.W. 1972. Forest fighting on the Eastern Front in World War II. *Geogr. Rev.* 186-202.
1543
- 1544 Miller, P.J.O., Biassoni, N., Samuels, A., and Tyack, P.L. 2000. Whale songs lengthen in
1545 response to sonar. *Nature* **405**(6789):903-903.
1546
- 1547 Miller, T.A., Yaghoobian, A., Stuckert, B.J., Goudey, J.S., and McAndless, J.M. 1998. The use
1548 of mapping techniques in an ecological risk assessment of sites contaminated with
1549 chemical warfare agents. *J. Hazard. Mater.* **61**:31-36.

- 1550
- 1551 Mole, R.H. 1958. The development of leukaemia in irradiated animals. *Brit Med. Bull.*
- 1552 **14**(2):174-177.
- 1553
- 1554 Monfils, R. 2005. The global risk of marine pollution from WWII shipwrecks: examples from
- 1555 the seven seas. *In International Oil Spill Conference, Miami Beach, FL, USA, 15-19 May*
- 1556 *2005. pp. 1049-1054.*
- 1557
- 1558 Mooney, H.A., and Cleland, E.E. 2001. The evolutionary impact of invasive species. *Proc. Nat.*
- 1559 *Acad. Sci.* **98**:5446-5451.
- 1560
- 1561 Moreira, F., and Russo, D. 2007. Modelling the impact of agricultural abandonment and
- 1562 wildfires on vertebrate diversity in Mediterranean Europe. *Landscape Ecol.* **22**:1461-1476.
- 1563
- 1564 Mortberg, U.M., Balfors, B., and Knol, W.C. 2007. Landscape ecological assessment: a tool for
- 1565 integrating biodiversity issues in strategic environmental assessment and planning. *J.*
- 1566 *Environ. Manage.* **82**(4):457-470.
- 1567
- 1568 Moura, R.L., Secchin, N.A., Amado-Filho, G.M., Francini-Filho, R.B., Freitas, M.O., Minte-
- 1569 Vera, C.V.Teixeira, J.B., Thompson, F.L., Dutr, G.F., Sumida, P.Y.G., Guth, A.Z., Lopes,
- 1570 R.M., and Bastos, A.C. 2013. Spatial patterns of benthic megahabitats and conservation
- 1571 planning in the Abrolhos Bank. *Cont. Shelf Res.* **70**:109-117.
- 1572

- 1573 Mrema, E., Bruch, C.E., and Diamond, J. 2009. Protecting the environment during armed
1574 conflict: an inventory and analysis of international law. UNEP/Earthprint.
1575
- 1576 Mudie, N.Y., Gusev, B. I., Pivina, L.M., Schoemaker, M.J., Rijinkova, O.N., Apsalikov, K.N.,
1577 and Swerdlow, A. J. 2007. Sex ratio in the offspring of parents with chronic radiation
1578 exposure from nuclear testing in Kazakhstan. *Radiat. Res.* **168**(5):600-607.
1579
- 1580 Nguyen, T.T. 2009. Vietnam and the environment: problems and solutions. *Int. J. Environ. Stud.*
1581 **66**(1):1-8.
1582
- 1583 Nishiwaki, Y. 1995. Fifty years after Hiroshima and Nagasaki (Invited Lecture). *In Proceedings*
1584 *Radiation Protection in Neighbouring Countries in Central Europe, Portoroz, SI, 4-8 Sept*
1585 *1995. Edited by D. Glavic-Cindro. Portoroz, SI. pp. 17-26.*
1586
- 1587 Nishiwaki, Y., Kawai, H., Shono, N., Fujita, S., Matsuoka, H., Fujiwara, S., and Hosoda, T.
1588 2000. Uncertainties under emergency conditions in Hiroshima and Nagasaki in 1945 and
1589 Bikini Accident in 1954. *In 10th International Congress of The International Radiation*
1590 *Protection Association. Hiroshima, JP, May 14-18 2000. pp 1-10*
1591
- 1592 Noble, I.R., and Slatyer, R.O. 1980. The use of vital attributes to predict successional changes in
1593 plants communities subject to recurrent disturbances. *Vegetatio* **4**(3):5-21.
1594

- 1595 NRDC. 2003. Navy Sonar System Threatens Whales [online]. Available from
1596 <http://www.nrdc.org/wildlife/marine> [Accessed 15 April 2015].
1597
- 1598 O'Farrell, T.P. 1984. Populations of small mammals inhabiting sites 22 to 32 years after nuclear
1599 events. *In* Joint international Meeting of the Australian and the American Society of
1600 Mammalogists, Sydney, AU, 9-13 July 1984. Report No. CONF-840770-1
1601
- 1602 Ochiai, E. 2014. Hiroshima to Fukushima biohazards of radiation. Springer-Verlag Berlin
1603 Heidelberg, Berlin, DE.
1604
- 1605 Ohkita, T. 1975. A. Acute Effects. *J. Radiat. Res.* **16**(Suppl 1):49-66.
1606
- 1607 Olival, K. J., and Higuchi, H. 2006. Monitoring the long-distance movement of wildlife in Asia
1608 using satellite telemetry. *In* Conservation Biology in Asia. *Edited by* J. McNeely, T.
1609 McCarthy, A. Smith, L. Whittaker and E. Wikramnayake. Society for Conservation
1610 Biology Asia Section and Resources Himalaya Foundation, Kathmandu, NP. pp 319-339.
1611
- 1612 O'May, J.F., Hansen, C.E., Heilman, E.G., Kaste, R.C., and Neiderer, A.M. 2005. Battlespace
1613 terrain ownership: a new situation awareness tool. Available from the Army Research Lab
1614 Aberdeen Proving Ground, M.D., USA.
1615

- 1616 Orians, G.H., and Pfeiffer, E.W. 1970. Ecological effects of the war in Vietnam. *Science*
1617 **168**:544-554.
- 1618
- 1619 Ortiz-Roque, C., and Lopez-Riviera, Y. 2004. Mercury contamination in reproductive age
1620 women in a Caribbean island: Vieques. *J. Epidemiol. Commun. Health* **58**: 756-757.
- 1621
- 1622 Osuji, L.C., and Nwoye, I. 2007. An appraisal of the impact of petroleum hydrocarbons on soil
1623 fertility: the Owaza experience. *Afr. J. Agricult. Res.* **2**(7):318-324.
- 1624
- 1625 Oughterson, A.W., and Warren, S. 1956. Medical effects of the atomic bomb in Japan (Vol 8).
1626 McGraw-Hill, New York, NY, USA.
- 1627
- 1628 Oughterson, A.W., LeRoy, G.V., Liebow, A.A., Hammond, E.C., Barnett, H.L., Rosenbaum,
1629 J.D., and Schneider, B.A. 1951. Statistical analysis of the medical effects of the atomic
1630 bombs. Available from the Report of the Joint Commission for the Investigation of the
1631 Effects of the Atomic Bomb in Japan, Army Institute of Pathology. Report No. TID-5252
- 1632
- 1633 Owens, S. 1990. Defense and the environment: the impacts of military live firing in national
1634 parks. *Cambridge J. Econ.* **14**:497-505.
- 1635
- 1636 Oyama, A., and Sasaki, T. 1946. A case of burn of the cornea and retina by atomic bomb. *Ganka*
1637 *Rinsho Iho* **40**:177.

- 1638
- 1639 Palumbo, R.F. 1962. Recovery of the land plants at Eniwetok Atoll following a nuclear
1640 detonation. *Radiat. Bot.* **1**:182-189.
- 1641
- 1642 Papanikolaou, N.C., Hatzidaki, E.G., Belivanis, S., Tzanakakis, G.N., and Tsatsakis, A.M. 2005.
1643 Lead toxicity update. A brief review. *Med. Sci. Monit.* **11**(10):RA329-RA336.
- 1644
- 1645 Papastefanou, C. 2002. Depleted uranium in military conflicts and the impact on the
1646 environment. *Health phys.* **83**(2):280-282.
- 1647
- 1648 Parkinson, B.W. 1996. Progress in astronautics and aeronautics: global positioning system:
1649 theory and applications, volume 2. AIAA, Reston, V.A., USA.
- 1650
- 1651 Parsons, E.C.M., Birks, I., Evans, P.G.H., Gordon, J.G., Shrimpton, J.H., and Pooley, S. 2000.
1652 The possible impacts of military activity on cetaceans in West Scotland. *Eur. Res.*
1653 *Cetaceans* **14**: 185-190.
- 1654
- 1655 Parsons, E.C.M., Dolman, S.J., Wright, A.J, Rose, N.A., and Burns, W.C.G. 2008. Navy Sonar
1656 and cetaceans: just how much does the gun need to smoke before we act? *Mar. Pollut. Bull.*
1657 **56**:1248-1257.
- 1658
- 1659 Pearson, C. 2012. Researching militarized landscapes: a literature review on war and the
1660 militarization of the environment. *Landscape Res.* **37**:115-133.
- 1661

- 1662 Pekins, C.E. 2006. Armored military training and endangered species restrictions at Fort Hood,
1663 Texas. *Fed. Facil. Environ. J.* **17**(1):37-50.
1664
- 1665 Pendersen, D. 2002. Political violence, ethnic conflict, and contemporary wars: broad
1666 implications for health and social well-being. *Soc. Sci. Med.* **55**:175-190.
1667
- 1668 Pierce, D.A., and Preston, D.L. 2000. Radiation-related cancer risks at low doses among atomic
1669 bomb survivors. *Radiat. Res.* **154**(2):178-186.
1670
- 1671 Pinaev, V.S., and Shcherbakov, V.A. 1996. Fires caused by nuclear explosions and their
1672 consequences. *Combust. Explo. Shock+*. **32**(5):572-576.
1673
- 1674 Pinca, S., Beger, M., Jacobson, D., and Keju, T. 2005. The state of coral reef ecosystems of the
1675 Marshall Islands. *In* *The State of Coral Reef Ecosystems of the United States and Pacific*
1676 *Freely Associated States. Edited by J. Waddell. NOAA Technical Memorandum NOS*
1677 *NCCOS 11, Silver Spring, M.D., USA. pp. 373–386.*
1678
- 1679 Planes, S., Galzin, R., Bablet, J.P., and Sale, P.F. 2005. Stability of coral reef fish assemblages
1680 impacted by nuclear tests. *Ecology* **86**(10):2578-2585.
1681

- 1682 Popper, A.N., and Hastings, M.C. 2009. The effects of human-generated sound on fish. *Integ.*
1683 *Zool.* **4**(1):43-52.
- 1684
- 1685 Porter W. 2005. Movement of toxic materials through the Vieques marine ecosystem: The
1686 effects of naval bombardment on a Puerto Rican coral reef. *In* Paper presented at
1687 Ecological Society of America Annual Meeting, Montreal, Q.C., CA, 7-12 August 2005.
- 1688
- 1689 Prentice, R.L., Kato, H., Yoshimoto, K., and Mason, M. 1982. Radiation exposure and thyroid
1690 cancer incidence among Hiroshima and Nagasaki residents. *Natl. Cancer I. Monogr.*
1691 **62**:207-212.
- 1692
- 1693 Prosser, C.L., Painter, E.E., Lisco, H., Brues, A.M., Jacobson, L.O., and Swift, M.N. 1947. The
1694 clinical sequence of physiological effects of ionizing radiation in animals 1. *Radiology*
1695 **49**(3):299-313.
- 1696
- 1697 Ramos, T.B., Alves, I., Subtil, R., and de Melo, J.J. 2007. Environmental performance policy
1698 indicators for the public sector: The case of the defence sector. *J. Environ. Manage.*
1699 **82**(4):410-432.
- 1700
- 1701 Randall, P.A., 1961. Damage to conventional and special types of residences exposed to nuclear
1702 effects. Available from the US Atomic Energy Commission, Washington, D.C., USA.
1703 Report WT-1194.

- 1704
- 1705 Rausch, R.L. 1973. Post mortem findings in some marine mammals and birds following the
1706 Cannikin test on Amchitka Island. Available from U.S. AEC, Nevada Operations Office,
1707 Las Vegas, N.V., USA. Report No. HVO-130.
- 1708
- 1709 Real, A., Sundell-Bergman, S., Knowles, J.F., Woodhead, D.S., and Zinger, I. 2004. Effects of
1710 ionising radiation exposure on plants, fish and mammals: relevant data for environmental
1711 radiation protection. *J. Radiol. Prot.* **24**(4A):A123-A137.
- 1712
- 1713 Rendell, L.E., and Gordon, J.C.D. 1999. Vocal response of long-finned pilot whales
1714 (*Globicephala melas*) to military sonar in the Ligurian Sea. *Mar. Mammal Sci.* **15**:198–
1715 204.
- 1716
- 1717 Rhoads, W.A., and Platt, R.B. 1971. Beta radiation damage to vegetation from close-in fallout
1718 from two nuclear detonations. *BioScience* **21**(15):1121-1125.
- 1719
- 1720 Rhoads, W.A., and Ragsdale, H.L. 1971. *Artemisia* shrub size and radiation damage to *Artemisiu*
1721 from local fallout from Project Schooner. *In* Pro. Nat. Symp. Radioecology, 3rd. *Edited by*
1722 D.J. Nelson. National Technical Information Service, U.S. Department of Commerce,
1723 Springfield, V.A., USA, pp. 953-960.
- 1724
- 1725 Rhoads, W.A., Asher D. K., and Ragsdale, H.L. 1972. Ecological and environmental effects
1726 from local fallout from schooner. 2. The beta and gamma radiation effects from close in

- 1727 fallout. Available from the US Atomic Energy Commission, Washington, D.C., USA.
- 1728 Technical Report No. PNE--529.
- 1729
- 1730 Richards, Z.T., Beger, M., Pinca, S., and Wallace, C.C. 2008. Bikini Atoll coral biodiversity
- 1731 resilience five decades after nuclear testing. *Mar. Pollut. Bull.* **56**:503-515.
- 1732
- 1733 Richardson, J.W., and West, T. 2000. Serious birdstrike accidents to military aircraft: updated
- 1734 list and summary. *IBSC Proc.* **25**(1):67-97.
- 1735
- 1736 Richmond, D.R., and White C.S. 1962. A tentative estimation of mans tolerance to overpressures
- 1737 from air blast. Available from the Defense Atomic Support Agency, Department of
- 1738 Defense, Washington, D.C., USA. Technical Progress Report No. DASA 1335.
- 1739
- 1740 Richmond, D.R., Taboreili, Bowen, I.G., Chiffelle, T.L., Hirsch, F.G., Longwell, B.B., Riley,
- 1741 J.G., White, C.S., Sherping, F., Goldizen, V.C., Ward, J.D., Wetherbe, M.B., Clare, V.R.,
- 1742 Kuhn, M.L., and Sanchez, R.T. 1959a. Blast biology a study of the primary and tertiary
- 1743 effects of blast in open underground protective shelters. Available from the Office of
- 1744 Technical Services, Department of Commerce, Washington D.C. USAEC Civil Effects
- 1745 Test Group Report WT-1467.
- 1746
- 1747 Richmond, D.R., Wetherbe, M.B., Taborelli, R.V., Sanchez,R.T., Sherping, F., Goldizen, V.C.,
- 1748 and White, C.S. 1959b. Shock tube studies of the effects of sharp-rising, long-duration

- 1749 overpressures on biological systems. Available from the Office of Technical Services,
1750 Department of Commerce, Washington, D.C., USA. USAEC Technical Report TID-6056.
1751
- 1752 Rideout, G. and Walsh, B.W. 1990. War games and multiple use: is it mission impossible to train
1753 combat troops and manage natural resources on the same forested acres? *Am. Forests*
1754 **96**:11-21.
1755
- 1756 Roberts S., and Williams J. 1995. After the guns fall silent: the enduring legacy of landmines.
1757 Available from the Vietnam Veterans of America Foundation, Washington, D.C., USA.
1758
- 1759 Robison, W.L., and Noshkin, V.E. 1999. Radionuclide characterization and associated dose from
1760 long-lived radionuclides in close-in fallout delivered to the marine environment at Bikini
1761 and Enewetak Atolls. *Sci. Tot. Environ.* **237**:311-327.
1762
- 1763 Rodda, G.H., and Savidge, J.A. 2007. Biology and impacts of Pacific Island invasive species.
1764 *Boiga irregularis*, the brown tree snake (Reptilia: *Colubridae*). *Pac. Sci.* **61**:307-324.
1765
- 1766 Rose, H.W., Brown, D.V.L., Byrnes, V.A., Cibis, P.A. 1956. Human chorioretinal burns from
1767 atomic fireballs. *Arch. Ophth.* **55**:205-210.
1768
- 1769 Rowell A., and Moore, P.F. 2000. Global review of forest fires. Available from Forests for Life
1770 Programme Unit, WWF International.

- 1771
- 1772 Ruhm, W., Walsh, L., and Nekolla, E.A. 2006. The cohort of the atomic bomb survivors-major
1773 basis of radiation safety regulations. *In* CAS - CERN Accelerator School and KVI:
1774 Specialised CAS Course on Small Accelerators. CERN, Geneva, CH. pp. 319-329.
- 1775
- 1776 Sanderson, H., Fauser, P., Thomsen, M., Vanninen, P., Soderstrom, M., Savin, Y., Khalikov, I.,
1777 Hirvonen, A., Niiranen, S., Missiaen, T., Gress, A., Borodin, P., Medvedeva, N., Polyak,
1778 Y., Paka, V., Zhurbas, V., and Feller P. 2010. Environmental hazards of sea-dumped
1779 chemical weapons. *Environ. Sci. and Technol.* **44**(12):4389-4394.
- 1780
- 1781 Sarkees, M.R., Wayman, F.W., and Singer, J.D. 2003. Inter-state, intra-state, and extra-state
1782 wars: a comprehensive look at their distribution over time, 1816–1997. *Int. Stud.*
1783 *Q.* **47**(1):49-70.
- 1784
- 1785 Schechter, A., Birnbaum, L., Ryan, J.J., and Constable, J.D. 2006. Dioxins: an overview. *Environ.*
1786 *Res.* **101**(3):419-428.
- 1787
- 1788 Schechter, A., Kooke, R., Serne, P., Olie, K., Huy, D.Q., Hue, N., and Constable, J. 1989.
1789 Chlorinated Dioxin and dibenzofuran levels in food samples collected between 1985-87 in
1790 the north and south of Vietnam. *Chemosphere.* **18**:627-634.
- 1791

- 1792 Schiffman, R. 2014. Drones flying high as new tool for field biologists. *Science* **344**(6183):459-
1793 459.
1794
- 1795 Schonfelder, P., Bresinsky, A., Garnweidner, E., Krach, E., Linhard, H., Mergenthaler, O.,
1796 Nezdal, W., and Wirth, V. 1990. *Verbreitungatlas der Farn- und Blütenpflanzen Bayerns*.
1797 Eugen Ulmer, Stuttgart, DE.
1798
- 1799 Science News Letter. 1945. Magnetron tube. *Sci. News* **48**(26):403.
1800
- 1801 Science Wire. 2001. How Does Sonar Work? [Online]. Available from
1802 <http://www.exploratorium.edu/theworld/sonar/sonar.html> [Accessed 16 April 2015].
1803
- 1804 Scott, K.N. 2007. Sound and cetaceans: a regional response to regulating acoustic marine
1805 pollution. *J. Int. Wildlife Law. Policy.* **10**:175-199.
1806
- 1807 Shaeffer, J.R. 1957. Radiation injuries, with notes on Hiroshima and Nagasaki. *Am. J. Surg.*
1808 **93**(4):641-643.
1809
- 1810 Shambaugh, J., Oglethorpe, J., and Ham, R. 2001. The trampled grass: mitigating the impacts of
1811 armed conflict on the environment. Available from the Biodiversity Support Program,
1812 Washington, DC, USA. Report No. 333.72096 S528.
1813
- 1814 Sheppard, S.C., Sheppard, M.I., Gallerand, M., and Sanipelli, B. 2005. Deviation of ecotoxicity
1815 thresholds for uranium. *J. Environ. Radioact.* **79**(1):55-83.

1816

1817 Shields, L.M., and Wells, P.V. 1962. Effects of nuclear testing on desert vegetation. *Science*
1818 **135**(3497):38-40.

1819

1820 Shields, L.M., Wells, P.V., and Rickard, W.H. 1963. Vegetational recovery on atomic target
1821 areas in Nevada. *Ecology* **44**(4):697-705.

1822

1823 Sidel, V.W. 2000. The impact of military preparedness on health and the environment. *In* *The*
1824 *Environmental Consequences of War: Legal, Economic, and Scientific Perspectives.*
1825 *Edited by* J.E., Austin, and C.E., Bruch. Cambridge University Press, Cambridge, UK. pp.
1826 426-443.

1827

1828 Silberner, J. 1981. Hiroshima & Nagasaki: thirty six years later, the struggle continues. *Sci.*
1829 *News* **120**(18):284-285,287.

1830

1831 Simenstad, C.A. 1974. Biological effects of underground nuclear testing on marine organisms. I.
1832 Review of documented shock effects, discussion of mechanisms of damage, and
1833 predictions of Amchitka test effects. *In* *Proceedings of the First Conference on the*
1834 *Environmental Effects of Explosives and Explosions (Technical Report 73-223).* Naval
1835 Ordinance Laboratory, White Oak, M.D., USA. pp. 86-97.

1836

- 1837 Singer, F.J., Schreier, W., Oppenhiem, J., and Garton. E.O. 1989. Drought, fires and large
1838 mammals. *BioScience* **39**(10):716-722.
- 1839
- 1840 Slotten, H.R. 2002. Satellite communications, globalization, and the Cold War. *Technol. Cult.*
1841 **43**(2):315-350.
- 1842
- 1843 Small, R.D., and Bush, B.W. 1985. Smoke production from multiple nuclear explosions in
1844 nonurban areas. *Science* **229**(4712):465-469.
- 1845
- 1846 Smith, M.A., Turner, M.G., and Rusch, D.H. 2002. Lupine and the karner blue butterfly at Fort
1847 McCoy Wisconsin, USA. *Environ. Manage.* **29**(1):102-115.
- 1848
- 1849 Smith, S. L. 2011. Toxic legacy: mustard gas in the sea around us. *J. law Med. Ethics* **39**(1):34-
1850 40.
- 1851
- 1852 Sparrow, A.H., and Woodwell, G.M. 1962. Prediction of the sensitivity of plants to chronic
1853 gamma irradiation. *Radiat. Bot.* **2**(1):9-26.
- 1854
- 1855 Springer, P.J. 2013. *Military robots and drones: a reference handbook*. ABC-CLIO, Santa
1856 Barbara, CA, USA.
- 1857

- 1858 Stanley, E.H., and Doyle, M.W. 2003. Trading off: the ecological effects of dam removal. *Front.*
1859 *Ecol. Environ.* **1**:15-22.
1860
- 1861 Stein, B.A., Scott, C., and Benton, N. 2008. Federal lands and endangered species: the role of
1862 military and other federal lands in sustain biodiversity. *BioScience* **58**(4):339-347.
1863
- 1864 Stellman, J.M., Stellman, S.D., Christian, R., Weber, T., and Tomasallo, C. 2003. The extent and
1865 patterns of usage of Agent Orange and other herbicides in Vietnam. *Nature* **422**:681-687.
1866
- 1867 Stephenson, T.R., Vaughan, M.R., and Anderson, D.E. 1996. Mule deer movement in response
1868 to military activity in southeast Colorado. *J. Wildlife Manage.* **60**(4):777-787.
1869
- 1870 Stoddart, D.R. 1968. Catastrophic human interference with coral atoll ecosystems. *Geography*
1871 **53**(1):25-40.
1872
- 1873 Sugg, D.W., Chesser, R.K., Brooks, J.A., and Grasman, B.T. 1995. The association of DNA
1874 damage to concentrations of mercury and radiocesium in largemouth bass. *Environ.*
1875 *Toxicol. Chem.* **14**(4):661-668.
1876
- 1877 Summary Report of Research in the Effects of the Atomic Bomb. 1951. *Edited by* the Japan
1878 Science Council. Japan Society for the Promotion of Science.
1879

- 1880 Sutherland, W.J., Bardsley, S., Clout, M., Depledge, M.H., Dicks, L.V., Fellman, L., Fleishman,
1881 E., Gibbons, D.W., Keim, B., Lickorish, F., Margerson, C., Monk, K.A., Norris, K., Peck,
1882 L.S., Prior, S.V., Scharlemann, J.P.W., Spalding, M.D., and Watkinson, A.R. 2013. A
1883 horizon scan of global conservation issues for 2013. *Trends Ecol. Evol.* **28**(1):16-22.
1884
- 1885 Sweetman, J. 1982. *Operation chastise: the dams raid: epic or myth.* Jane's Publishing Company,
1886 London, UK.
1887
- 1888 Tang, Z., Engel, B.A., Pijanowski, B.C., and Lim, K.J. 2005. Forecasting land use change and its
1889 environmental impact at a watershed scale. *J. Environ. Manage.* **76**:35-45.
1890
- 1891 Tavares, M., and De Melo, G.A.S. 2004. Discovery of the first known benthic invasive species in
1892 the southern ocean: The north Atlantic spider crab *Hyas araneus* found in the Antarctic
1893 Peninsula. *Antarct. Sci.* **16**:129-131.
1894
- 1895 Telesco, D.J., and Van Manen, F.T. 2006. Do black bears respond to military weapons training?
1896 *J. Wildlife Manage.* **70**(1):222-230.
1897
- 1898 Theodorakis, C.W., and Shugart, L.R. 1997. Genetic ecotoxicology II: population genetic
1899 structure in mosquito fish exposed in situ to radionuclides. *Ecotoxicology* **6**:335-354.
1900

- 1901 Theodorakis, C.W., and Shugart, L.R. 1998. Genetic ecotoxicology III: the relationship between
1902 DNA strand breaks and genotype in mosquito fish exposed to radiation. *Ecotoxicology*
1903 *7(4):227-235.*
- 1904
- 1905 Theodorakis, C.W., Bickham, J.W., Elbl, T., Shugart, L.R., and Chesser, R. K. 1998. Genetics of
1906 radionuclide-contaminated mosquitofish populations and homology between *Gambusia*
1907 *affinis* and *G. holbrooki*. *Environ. Toxicol. Chem.* **17(10):1992-1998.**
- 1908
- 1909 Theodorakis, C.W., Bickham, J.W., Lamb, T., Medica, P.A., and Lyne, T.B. 2001. Integration of
1910 genotoxicity and population genetic analyses in kangaroo rats (*Dipodomys merriami*)
1911 exposed to radionuclide contamination at the Nevada Test Site, USA. *Environ. Toxicol.*
1912 *Chem.* **20(2):317-326.**
- 1913
- 1914 Troll, K. 2000. The impact of anti-personnel landmines on the environment. Available from
1915 United Nations Institute for Disarmament Research, Geneva, CH.
- 1916
- 1917 Trumbull, V.L., Dubois, P.C., and Brozka, R.J. 1994. Military camping impacts on vegetation
1918 and soils of the Ozark Plateau. *J. Environ. Manage.* **40:329-339.**
- 1919
- 1920 Tull, M. 2006. The environmental impact of ports: an Australian case study. *In* Paper presented
1921 at the XIV International Economics History Congress (session No. 58). Helsinki, FI.
- 1922

- 1923 Tullis, J.L., Lamson, B.G., and Madden, S.C. 1955. Pathology of swine exposed to total body
1924 gamma radiation from an atomic bomb source. *Am. J. Pathol.* **31**(1):41-71.
1925
- 1926 Turner, F.B. 1975. Effects of continuous irradiation on animal populations. *Adv. Radiat. Biol.*
1927 **5**:83-144.
1928
- 1929 Turner, F.B., and Medica, P.A. 1977. Sterility among female lizards (*Uta stansburiana*) exposed
1930 to continuous γ irradiation. *Radiat. Res.* **70**(1):154-163.
1931
- 1932 Turner, F.B., Licht, P., Thrasher, J.D., Medica, P.A., Lannom Jr, J.R., and Nelson, D. J. 1971.
1933 Radiation- induced sterility in natural populations of lizards (*Crotaphytus wislizenii* and
1934 *Cnemidophorus tigris*). In *Radionuclides in Ecosystems. Edited by D.J., Nelson.* U.S.
1935 Atomic Energy Commission Report CONF-710501-P2, pp. 1131-1143.
1936
- 1937 Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., and Steininger, M. 2003.
1938 Remote sensing for biodiversity science and conservation. *Trends Ecol. Evol.* **18**(6):306-
1939 314.
1940
- 1941 Valeriote, F.A., and Baker, D.G. 1964. The combined effects of thermal trauma and x-irradiation
1942 on early mortality. *Radiat. Res.* **22**(4):693-702.
1943

- 1944 Van Dover, C.L., Aronson, J., Pendleton, L., Smith, S., Arnaud-Haond, S., Moreno-Mateos, D.,
1945 Barbier, E., Billett, D., Bowers, K., Danovaro, R., Edwards, A., Kellert, S., Morato, T.,
1946 Pollard, E., Rogers, A., and Warner, R. 2014. Ecological restoration in the deep sea:
1947 Desiderata. *Mar. Policy* **44**:98-106.
- 1948
- 1949 Villoria Saez, P., del Rio Merino, M., Porrás-Amores, C., and Gonzalez, A.S.A. 2014. Assessing
1950 the accumulation of construction waste generation during residential building construction
1951 works. *Resour. Conserv. Recy.* **93**:67-74.
- 1952
- 1953 Visone, D.L. 2005. Battlespace terrain reasoning and awareness. *In* Proceedings of ESRI User
1954 Conference, 25-29 July 2005.
- 1955
- 1956 Vogt, H.P. 1995. Coral reefs in Saudi Arabia: 3.5 years after the Gulf War oil spill. *Coral Reefs*
1957 **14**:271-273.
- 1958
- 1959 Wanebo, C.K., Johnson, K.G., Sato, K., and Thorslund, T.W. 1968. Breast cancer after exposure
1960 to the atomic bombings of Hiroshima and Nagasaki. *New Engl. J. Med.* **279**(13):667-671.
- 1961
- 1962 Warren, S.D., and Büttner, R. 2006. Documentation of disturbance-dependent threatened and
1963 endangered species on US Army–Europe training areas in Bavaria. Available from the
1964 Center for Environmental Management of Military Lands, Fort Collins, C.O., USA.
1965 CEMML TPS, 06-05.

- 1966
- 1967 Warren, S.D., Holbrook, S.W., Dale, D.A., Whelan, N.L., Elyn, M., Grimm, W., and Jentsch, A.
- 1968 2007. Biodiversity and the heterogeneous disturbance regime on military training lands.
- 1969 Rest. Ecol. **15**(4):606-612.
- 1970
- 1971 Warren, S.D., Holbrook, S.W., Dale, D.A., Whelan, N.L., Elyn, M., Grimm, W., and Jentsch, A.
- 1972 2007. Biodiversity and the heterogeneous disturbance regime on military training lands.
- 1973 Restor. Ecol. **15**(4):606-612.
- 1974
- 1975 Watts, S.E. 1998. Short-term influence of tank tracks on vegetation and microphyticcrusts in
- 1976 shrubsteppe habitat. Environ. Manage. **22**(4):611-616.
- 1977
- 1978 Westing, A.H. 1971. Ecological effects of military defoliation on the forests of South Vietnam.
- 1979 BioScience **21**(17):893-898.
- 1980
- 1981 Westing, A.H. 1980. Warfare in a fragile world: military impact on the human environment.
- 1982 Taylor & Francis. London, UK.
- 1983
- 1984 Westing, A.H. 1984. Herbicides in war: past and present. Herbicides in war: the long term
- 1985 ecological and human consequence. Taylor & Francis. London, UK.
- 1986

- 1987 Westing, A.H. 1985. Explosive remnants of war: mitigating the environmental effects.
1988 Stockholm Peace Research Institute and United Nations. Environment Programme. Taylor
1989 and Francis, London, UK.
- 1990 Westing, A.H. 1986. Global resources and international conflict: environmental factors in
1991 strategic policy and action. Oxford University Press, UK.
- 1992
- 1993 Westing, A.H. 1989. Herbicides in warfare: the case of Indochina. *Ecotoxicology and Climate*.
1994 John Wiley & Sons Ltd, Hoboken, N.J., USA.
- 1995
- 1996 Westing, A.H. 1996. Explosive remnants of war in the human environment. *Environ. Conserv.*
1997 **23**(4):283-285.
- 1998
- 1999 Westing, A.H. 2003. Environmental dimension of the Gulf War of 1991. *In Security and*
2000 *Environment in the Mediterranean: Conceptualizing Security and Environmental Conflicts*
2001 *Edited by H.G., Brauch, Liotta, P.H., Marquina, A., Rogers, P.F., and Selim, M.E.* pp 523-
2002 534.
- 2003
- 2004 Westing, A.H. 2013a. The second Indochina War of 1961-1975: its environmental impact. *In*
2005 *Arthur H. Westing: pioneer on the environmental impact of war*. Springer, New York,
2006 N.Y., USA. pp 35-50.
- 2007
- 2008 Westing, A.H. 2013b. The Gulf War of 1991: its environmental impact. *In Arthur H. Westing:*
2009 *pioneer on the environmental impact of war*. Springer, New York, N.Y., USA. pp 50-76.

- 2010
- 2011 Whicker, F.W., Hinton, T.G., MacDonell, M.M., Pinder III, J.E., and Habegger, L.J. 2004.
- 2012 Avoiding destructive remediation at DOE sites. *Science* **303**(5664):1615-1616.
- 2013
- 2014 Whicker, F.W., and Pinder, J.E. 2002. Food chains and biogeochemical pathways: contributions
- 2015 of fallout and other radiotracers. *Health Physics* **82**(5):680-689.
- 2016
- 2017 Whitecotton, R.C.A., David, M.B., Darmody, R.G., and Price, D.L. 2000. Impact of foot traffic
- 2018 from military training on soil and vegetation properties. *Environ. Manage.* **26**(6):697-706.
- 2019
- 2020 Wills, J. 2001. “Welcome to the atomic park”: American nuclear landscapes and the ‘unnaturally
- 2021 unnatural’. *Environ. Hist.* **7**(4):449-472.
- 2022
- 2023 Wilson, S.D. 1988. The effects of military tank traffic on prairie: a management model. *Environ.*
- 2024 *Manage.* **12**(3):397-403.
- 2025
- 2026 Xun, Z., Cao, N., and Zhao, T. 2013. Evaluation of the risks of environment in the construction
- 2027 process on the basis of Grey System theory. *Am. Soc. Civil Eng.* 741-749.
- 2028
- 2029 Yang, X., North, R., Romney, C. 2003. Worldwide nuclear explosions. *Int. Geophys. Ser.*
- 2030 **81**(B):1595-1600.
- 2031

- 2032 Yelverton, J.T., Richmond, D.R., Hicks, W., Saunders, K., and Fletcher, E.R. 1975. The
2033 relationship between fish size and their response to underwater blast. Available from the
2034 Defense Nuclear Agency, Fort Belvoir, V.A., USA. Topical Report DNA 3677T.
2035
- 2036 Zahler, P., and Graham, P. 2001. War and Wildlife. The Afghanistan conflict and its effects on
2037 the environment. Available from International Snow Leopard Trust Special Report. pp. 1-
2038 13.
2039
- 2040 Zakrajsek, E.J., Bissonette, J.A. 2005. Ranking the risk of wildlife species hazardous to military
2041 aircraft. *Wildlife Soc. Bull.* **33**(1):258-264.
2042
- 2043 Zallinger, C., and Tempel, K. 1998. The physiologic response of domestic animals to ionizing
2044 radiation: a review. *Vet. Radiol. Ultrasoun.* **39**(6):495-503.
2045
- 2046 Zentelis, R., and Lindenmayer, D. 2014. Bombing for biodiversity – enhancing conservation
2047 values of military training areas. *Conserv. Lett.* **00**(0):1-7.
2048
2049
2050
2051
2052
2053
2054

2055 **Table 1:** Research gaps related to the effects of warfare on ecosystem structure and function.
2056

Warfare Aspect	Research Gaps
<i>Active Armed Conflict</i>	<ul style="list-style-type: none"> • Areas of conflict are often too hazardous for researchers to enter and gather data. • BACI experimental (Before-After Control-Impact) approach is usually not possible since conflicts occur without consultation with researchers. Armed conflicts often occur randomly, kept confidential, or in areas that are difficult to access by researchers from abroad (e.g. drone or aerial assaults). • Conservation priorities can be overlooked during war activities resulting in lack of pre- and post-war efforts to maintain and monitor ecological integrity and animal populations. • Lack of international capacity to monitor threats from armed conflict on ecosystems, particularly when armed conflicts occur between several nations and across large spatial scales.
<i>Nuclear Warfare</i>	<ul style="list-style-type: none"> • Effects of nuclear weapons had high anthropogenic focus (e.g. effects on human health, buildings, etc.), information on greater impacts on ecological functioning at the population and biodiversity level is relatively scarce; taxonomic representation relatively low. • Ecosystem diversity is under-represented as testing was generally restricted to a few select habitat types, mostly desert regions, that are typically low in biodiversity to begin with. • The long term impacts from radiation exposure from nuclear weapon produced fallout/radiation has been minimally documented in wildlife with little regard for potential fitness impact; timescale is an issue and impacts may be mitigated by immigration. • As nuclear weapons use is currently banned under international treaties, new research avenues into ecosystem structural impacts are potentially limited.

Military Training

- Research focused on military training facilities operated by the U.S Department of Defense, within North America. Data extrapolated to address the impacts of military training facilities located abroad; minimal investigation into whether these assessments address impacts in different geographic environments, under different training regimes, etc.
- Many military training lands and facilities are situated within biodiversity hotspots, which are home to numerous rare and endemic species; these operational training bases may be located in hostile developing nations where research access is restricted creating knowledge gaps in these unmanaged and unprotected areas.
- Research and environmental assessments pertaining to military training activities is relatively new, gaining importance within the past 50 years, which has created knowledge gaps in how certain training landscapes existed and functioned prior to military management.

Military Contamination

- Many surplus marine munitions and barrels after WWII were dumped at undisclosed locations, making it both difficult to estimate potential contamination and to initiate recovery.
- The broad spatial scales at which environment may be exposed to contamination can cross geographic and political borders, complicating accountability procedures.
- Chemicals, hydrocarbons, and metals can have legacy effects in soil, water, and plant and animal tissue. Immediate effects may not always be obvious.

2057

2058


2059

2060

2061

2062

2063



2064

2065

2066 **Figure 1:** Overview of the potential deleterious impacts of warfare on the environment including

2067 terrestrial, aerial and naval theatres of war.

2068

2069

2070

2071

Draft

2072 **Box 1:** The Korean Demilitarized Zone



2073

2074

2075 The Demilitarized Zone (DMZ) on the Korean peninsula serves as a protected area for many endemic and
2076 endangered species (Kim 1997; Healy 2007). The Korean DMZ is a 4 km wide by 250 km long strip of
2077 natural land which has separated North and South Korea since 1953 with the Armistice Agreement (Kim
2078 1997). This area is home to 3,514 species, which equates to 67% of the species diversity of the Korean
2079 Peninsula, most of which are endemic to this small plot of land (Healy 2007). One of the world's most
2080 endangered bird species, the White-Naped Crane (*Grus vipio*) relies on the Korean DMZ for critical
2081 overwintering habitat for 50% of the remaining population. As well, areas within the DMZ are also
2082 reported as important resting locations during the north-south migration for a large proportion of the crane
2083 population in addition to numerous other bird species (Higuchi et al. 1996; Healy 2007). Photo credit:
2084 Adrian Pingstone, Wikimedia Commons, 2006.

2085

2086

2087

2088 **Box 2:** The Marshall Islands Reef Recovery

2089


2090

2091

2092

2093

2094



2095

2096

2097

2098 During the decades following the Second World War, the testing of nuclear weapons by the United States
2099 military was well underway (above). The Marshall Islands were home to a great number of nuclear
2100 detonations comprising a total of 66 test blasts that left the surrounding environment devastated.
2101 However, due to the area having large degrees of residual radioactivity, human exclusion from many of
2102 the test site islands has generated a marine protected area of sorts alleviating anthropogenic stress from
2103 the region (Donaldson et al. 1997; Berger et al. 2008). As such, the system has been allowed to recover in
2104 isolation for the greater part of the last half century and has produced some interesting results. With the
2105 exception of a few specialized species, scleractinian coral diversity has rebounded on a number of reefs
2106 affected by nuclear testing (Richards et al. 2008). As well, the size-frequency distribution, an indicator of
2107 biomass, of many fish taxonomic groups within former blast sites have been observed to be much greater
2108 than that of the surrounding waters unaffected by nuclear testing (Houk and Musburger 2013). While
2109 nuclear testing is devastating on an acute timescale, it may prove to be beneficial to the local ecosystem

2110 over a more chronic duration through human exclusion. Photo Credit: United States Department of
2111 Defense, 1946.

2112

2113

2114

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

Draft

2130 **Box 3: Military Training Bases**

2131



2132

2133

2134 Military training bases have long been known as areas of high biodiversity and, as of late, these vast
2135 military training landscapes are becoming increasingly recognized as important refuge areas for IUCN
2136 red-listed species (Zentelis and Lindenmayer 2014). A case study examination conducted by Stein et al.
2137 (2008) evaluated the status of U.S. federally listed endangered species across the 264 million hectares of
2138 government owned and managed lands in the United States. This case study identified a significantly
2139 greater density and diversity of endangered and imperiled species inhabiting military training lands,
2140 compared to all other federally managed lands across the country (Right image: recreated from Stein et
2141 al., 2008). In addition to this finding, the greatest diversity of endangered and imperiled species were
2142 found inhabiting 4 training bases in the Hawaiian Islands, led by Oahu's Schofield Barracks Military
2143 Reservation supporting approximately 47 federally endangered species and 53 imperiled species (Stein et
2144 al. 2008). Overall, more than 34% of the U.S. federally listed endangered species are found within
2145 Hawaiian military training bases, which makes these areas particularly vulnerable to military training
2146 exercises; stressing the importance for conservative land-use practices and management techniques to

2147 protect these ecologically valuable landscapes (Zentelis and Lindenmayer, 2014; Stein et al., 2008).

2148 Photo Credit: Polihale Wikimedia Commons, 2004.

2149

2150

2151

2152 **Box 4:** Oil Contamination in Arabian Gulf

2153



2154

2155

2156 After the Gulf War oil spill in 1991, coral reef species demonstrated substantial resilience to sea water

2157 temperature decreases and toxic hydrocarbon fall-out (Downing and Roberts 1993, Vogt 1995). The

2158 Arabian Gulf is home to several diverse coral reef communities both inshore and offshore. From 1984

2159 until 1994, coral species cover was estimated at six different locations off the coasts of Kuwait and Saudia

2160 Arabia using video recordings and diving teams. The combined results of two independent studies found

2161 that (Downing and Roberts 1993; Vogt 1995), after an estimated 6-8 million barrels of oil were added to

2162 the Arabian Gulf, there was no observable declines to coral reef health. Instead, from 1992 to 1994, Vogt

2163 (1995) observed significantly increasing trends in coral cover. Today, coral reef health is subject to

2164 climate change and increasing marine ecosystem pollution, which makes modern coral surveys unable to
2165 tease out any direct evidence of long-term effects of the Gulf War on coral communities (Downing and
2166 Roberts 1993, Al-Cibahy et al. 2012). Photo Credit: Michael J. Lawrence, 2012.

2167

2168

2169

2170

Draft