

The effects of problem-based learning during medical school on physician competency: a systematic review

Gerald Choon-Huat Koh MD MMed, Hoon Eng Khoo PhD, Mee Lian Wong MD MPH, David Koh MD PhD

∞ See related article page 61

ABSTRACT

Background: Systematic reviews on the effects of problem-based learning have been limited to knowledge competency either during medical school or postgraduate training. We conducted a systematic review of evidence of the effects that problem-based learning during medical school had on physician competencies after graduation.

Methods: We searched MEDLINE, EMBASE, CINAHL, PsycINFO, Cochrane Databases, and the tables of contents of 5 major medical education journals from earliest available date through Oct. 31, 2006. We included studies in our review if they met the following criteria: problem-based learning was a teaching method in medical school, physician competencies were assessed after graduation and a control group of graduates of traditional curricula was used. We developed a scoring system to assess the quality of the studies, categorized competencies into 8 thematic dimensions and used a second system to determine the level of evidence for each competency assessed.

Results: Our search yielded 102 articles, of which 15 met inclusion criteria after full text review. Only 13 studies entered final systematic analysis because 2 studies reported their findings in 2 articles. According to self-assessments, 8 of 37 competencies had strong evidence in support of problem-based learning. Observed assessments had 7 competencies with strong evidence. In both groups, most of these competencies were in the social and cognitive dimensions. Only 4 competencies had moderate to strong levels of evidence in support of problem-based learning for both self- and observed assessments: coping with uncertainty (strong), appreciation of legal and ethical aspects of health care (strong), communication skills (moderate and strong respectively) and self-directed continuing learning (moderate).

Interpretation: Problem-based learning during medical school has positive effects on physician competency after graduation, mainly in social and cognitive dimensions.

Une version française de ce résumé est disponible à l'adresse www.cmaj.ca/cgi/content/full/178/1/34/DC1

CMAJ 2008;178(1):34-41

Problem-based learning curricula have been introduced in many medical schools around the world.^{1,2} However, their adoption was met with some concern, primarily because of the substantial manpower needed. For example, student contact hours are 3–4 times greater for educators in a problem-based learning curriculum than for educators in a traditional curriculum. As a consequence, the economic viability of problem-based learning becomes a major concern when class sizes exceed 100 students.³ Given the limited resources available,⁴ evidence-based evaluation of the effects of problem-based learning during medical school on improving physician competency would certainly strengthen any justification for the adoption of such programs.^{5–9}

Past reviews of problem-based learning focused only on its effects during medical school or postgraduate training.^{10–16} In addition, 4 of those reviews also studied student and educator preferences and indicated that medical students^{11,14,15} and educators¹⁶ generally prefer problem-based learning to traditional teaching methods. The reviews did not study the effects of problem-based learning during medical school on the competencies of practising physicians. Collier emphasized this lack of evidence when he cautioned that student satisfaction cannot be extrapolated as a predictor of physician competency.¹⁷ To our knowledge, there has been only one systematic review, published in 1993, that indirectly reported the effects of problem-based learning during medical school on physician competency after graduation. The study was based on a small sample of doctors in their early postgraduate years.¹⁶ Since then, many rigorous studies have evaluated the effects of problem-based learning during medical school up to 20 years after graduation. We performed a systematic review of controlled studies to determine whether problem-based learning during medical school leads to greater physician competencies after graduation.

Methods

In our study, we used Maudsley's definition of problem-based learning, which she defined as both a method and philosophy involving problem-first learning via work in small groups and independent study.¹⁸

From the Department of Community, Occupational and Family Medicine (G. Koh, Wong, D. Koh), the Medical Education Unit (Khoo, Wong), Yong Loo Lin School of Medicine, and the Centre for Development of Teaching and Learning (Khoo), National University of Singapore, Singapore

Search

To identify relevant studies, we searched each of the following databases, from the earliest available date through Oct. 31, 2006: MEDLINE (1966), EMBASE (1980), CINAHL (1982), PsycINFO (1967), and the Cochrane Database of Systematic Reviews, the Cochrane Database of Abstracts of Reviews of Effects and the Cochrane Central Register of Controlled Trials (1993). We also searched the tables of contents of 5 major medical education journals (*Medical Education*, *Academic Medicine*, *Medical Teacher*, *BioMed Central Medical Education* and *Advances in Health Sciences Education*) for papers published from date of first issue through Oct. 31, 2006. We used the search terms “problem-based learning,” “practice-based,” “self-directed,” “learner centred” and “active learning” to identify problem-based learning studies and combined them with other key terms such as “medical education,” “undergraduate,” “postgraduate,” “practising,” “physicians” and “doctors.” We did not search by subject heading because not all databases use “problem-based learning” as a subject heading. For example, MEDLINE introduced the term as a medical subject heading only in 1995. We also manually searched the references of retrieved articles to trace potentially relevant papers.

Selection

We selected studies if they met the following inclusion criteria: use of problem-based learning as a teaching method in medical school, assessment of study population after graduation from medical school (including first-year residents and pre-registration house-staff) and use of a control group¹⁹ from a traditional (non-problem-based learning) curriculum. We excluded studies that were qualitative, involved nonphysicians, assessed competency before graduation, used problem-based learning only during postgraduate medical education or reported combined outcomes for medical students and postgraduate trainees. Because of concerns related to contamination including curriculum changes and co-interventions such as increased community focus or emphasis on communication skills, we excluded studies in which all outcomes could be contaminated by concurrent curriculum changes. And although we included studies that contained some outcomes that were contaminated by co-interventions, we excluded these contaminated outcomes from the final analysis. We included nonrandomized controlled trials because there were very few randomized controlled trials in medical education; we felt that this approach was acceptable, given that most educational research has methodological limitations because purity of curricular change and random assignment of students are rarely possible.^{8,20–22} Medical schools in the United States are mainly entry-level graduate programs, whereas those in the United Kingdom traditionally accept medical students without a university degree. As such, age and a prior degree could be major confounding factors when comparing studies between these 2 medical education systems.²³ Thus, we included only studies in which the control group came from the same medical education system. No restrictions were placed on study design, language or country. If there

was overlap in study populations and reported outcomes, we considered combining the findings from those studies for our final analysis.

Data abstraction

One of us (G.C.H.K.) developed a standardized data form to extract information from the articles we included in our systematic analysis. We extracted the following information: sample size, source of control group (whether the control group came from the same or a different university), study design, whether random allocation was performed, years spent in problem-based learning curriculum, number and type of competencies assessed, type of assessment, response rate, whether adjustment for potential confounders was performed, and methodological inconsistencies. If such information was missing, we obtained it by looking up references in the paper that previously reported the information or, in the case of verifying the number of years spent in a problem-based learning curriculum, by consulting the medical schools' websites. One of us (G.C.H.K.) performed the data abstraction, and the rest of us (H.E.K., M.L.W. and D.K.) independently checked the data extracted. We did not perform a meta-analysis because of the wide variations in the constructs of outcome measures and study reporting, which made calculating the effect sizes impossible for more than half of the selected studies.

Validity assessment

Because most validated tools for assessing study quality were designed for clinical interventions, the only tool available to assess the quality of problem-based learning studies was the one that Smits and colleagues¹³ developed to study the effectiveness of problem-based learning in postgraduate medical education. However, of the 5 study-quality indicators they proposed, only 2 could be applied across the studies included in our review: response rate and statistical adjustment. As suggested by Reed and colleagues,²⁴ we therefore adapted Smits and colleagues' instrument by adding 8 study-quality indicators of our own developed on the basis of our experience in epidemiology and medical education research.

The study-quality indicators we added were developed to adhere as closely as possible to validated quality-assessment instruments for clinical studies, such as the Scottish Intercollegiate Guidelines Network 50 instrument for cohort studies, and to apply to the full spectrum of heterogeneous studies. Like Smits and colleagues did with their instrument, we assigned some quality indicators (randomization, sample size, objective assessment and response rate) greater weight than others because we felt that they were of greater importance. We acknowledge that this approach is subjective, but our decisions were based largely on extensive discussion, reasonable assessment and consensus. We did not place restrictions on the timing of the post-graduation assessment because any differences due to the passage of time would be controlled by the comparison group.

All 4 of us independently assessed the quality of the studies using the adapted scoring model (Appendix 1, available at www.cmaj.ca/cgi/content/full/178/1/34/DC2). We sub-

sequently classified the studies into 2 levels of quality. Similar to the cut-off point used by Smits and colleagues, we considered studies to be high in quality if they scored above the mid-point of the scoring range.

Data synthesis

One of us (G.C.H.K.) developed a thematic categorization of physician competencies based on the 7 dimensions of physician competence described by Epstein and Hundert.²⁵ We combined 6 of their dimensions to form 3 for our study because of the lack of valid items within the 6 dimensions. The relationship and affective/moral dimensions were combined to form the social dimension, the context and habits-of-mind

dimensions became the managerial dimension, and the integrative dimension was subsumed under the cognitive dimension. We also extracted research, teaching and knowledge competencies as independent dimensions to explore the effects of problem-based learning in these 3 important areas. We created a new “overall” dimension because some studies included in our review reported global competency. We categorized the outcomes from studies that we entered into the final stage of our systematic analysis according to whether the competencies of graduates of problem-based learning curricula were better than, the same as or worse than those of graduates in the control group. This categorization of competencies was stratified according to study quality. We present the results for self- and observed assessments separately because medical education research suggests that students and physicians have a limited ability to assess themselves accurately.^{26,27} To determine the levels of evidence supporting the effect of problem-based learning during medical school on various physician competencies, we used a system adapted from Smits and colleagues in which the level of evidence is determined based on the number of studies in support of the data, the quality of those studies and the number of conflicting studies¹³ (Appendix 2, available online at www.cmaj.ca/cgi/content/full/178/1/34/DC2).

Results

Our selection process is outlined in Figure 1. Our keyword searches yielded 2675 articles. After applying the inclusion criteria to the abstracts of these articles, we excluded 2573. We obtained the full-text versions of the remaining articles and excluded a further 87 articles, which left 15 eligible articles for our final systematic analysis.²⁸⁻⁴²

There was overlap in study populations and reported outcomes among 4 of the 15 articles we retained for our final analysis. Although the studies by Mennin and colleagues³⁵ and Santos-Gomez and colleagues³⁶ were based on overlapping cohorts of graduates from the University of New Mexico, we treated these as 2 separate studies because both used different years of assessment and different types of assessment, and because only 4 of 17 self-assessed competencies overlapped. In 2 other cases, we opted to combine the findings from separate reports. In the first case Watmough and colleagues reported graduates' supervisor assessments and self-assessments in separate articles, which we treated as 1 study,^{32,33} and in the second case Woodard and colleagues' 1990 study included previous findings from a paper they published in 1981.^{41,42} As a result, our final analysis is based on 13 studies: 4 from the United States,^{28,29,35,36} 3 from Canada,^{38,40-42} 3 from Great Britain,^{30,32-34} 2 from the Netherlands^{37,39} and 1 from Australia³¹ (Table 1).

Study characteristics

The study populations ranged from first-year residents to physicians who had been practising for up to 20 years. Graduates of problem-based learning curricula attended medical schools that either offered problem-based learning in the first 2 years or throughout the duration (up to 5 years) of medical




Figure 1: Search and selection of studies for systematic review. *Reasons for exclusion do not add up to total because some articles were excluded for multiple reasons.

school. Types of assessment fell into 2 categories: self-assessment or observed assessment. Observed assessment was usually in the form of a supervisors' survey or an objective measure, such as a performance indicator database³⁴ or validated test.³⁸ One study included a nurses' survey.³⁶

Two studies had major methodological inconsistencies. Schmidt and colleagues³⁹ surveyed participants in the problem-based learning group in 1999 but surveyed participants in the control group 5 years later. Woodard⁴² used a different methodology to select controls between 2 graduating years of doctors. There was excellent agreement among all 4 of us for scoring study indicators (average chance-corrected [kappa] = 0.89). Table 2 classifies the study outcomes according to whether the competencies of graduates of problem-based learning curricula were better than (positive), the same as (no different) or worse than (negative) those of graduates in the control group (a more detailed version of Table 2 is in Appendix 3, available online at www.cmaj.ca/cgi/content/full/178/1/34/DC2). The outcomes of 6 studies^{28,31-35,37} could potentially be confounded by other re-

ported changes in problem-based-learning curricula (see remarks column of Table 2 for details). These outcomes were excluded when we assessed the levels of evidence.

Outcomes

We categorized the competencies assessed in the 13 studies into 8 dimensions: overall, technical, social, cognitive, managerial, research, teaching and knowledge competencies (Table 3) (more detailed versions of Table 3 are available in Appendices 4 [self-assessment] and 5 [observed assessment], available at www.cmaj.ca/cgi/content/full/178/1/34/DC2). Of the 8 dimensions, the social dimension showed the strongest evidence in support of problem-based learning. Within this dimension, there was strong evidence that problem-based learning was superior for 4 of 8 competencies: teamwork skills, appreciation of social and emotional aspects of health care, appreciation of legal and ethical aspects of health care, and appropriate attitudes toward personal health and well-being. Moreover, within the same dimension, there

Table 1: Characteristics of the 13 studies of problem-based learning included in the systematic analysis

Study	Country	n	Group; no. of graduates (class)		Timing of assessment post graduation, yr	Years of problem-based learning	No. of competencies assessed	Type of assessment (response rate)
			Problem-based learning	Control				
Peters et al ²⁸	United States	100	50 (1990)	50 (1989)	8-9	2	3	Self (80%)
Hoffman et al ²⁹	United States	1456¶	784¶ (1990-1996)	672¶ (1997-2003)	1	2	14	Supervisor (NR**)
Jones et al ³⁰	United Kingdom	261	138 (1999)	123 (1998)	1	5	19/13††	Self (61%-67%), supervisor (80%-82%)
Rolfe et al ^{31*†}	Australia	384	49 (1992)	335 (1992)	1	5	14	Supervisor (97.2%)
Watmough et al ^{32,33}	United Kingdom	639	316 (1999-2000)	323 (2001-2002)	1	2	18/13††	Self (52%-68%), supervisor (64%-78%)
Tamblyn et al ^{34*}	United Kingdom	165	128 (1991)	37 (1988-1990)	4-7	5	4	Administrative database (100%)
Mennin et al ^{35*}	United States	140	40 (1983-1986)	100 (1983-1986)	4-7	2	17	Self (83%-87%)
Santos-Gomez et al ³⁶	United States	130	44 (1985-1987)	86 (1985-1987)	1-3	2	8	Self (53%-83%), supervisor (60%-79%), nurse (58%-77%)
Prince et al ^{37†}	Netherlands	1159	239 (1999-2001)	920 (1999-2001)	2	5	12	Self (45%)
Shin et al ^{38*†}	Canada	96	48 (1974-1985)	48 (1974-1985)	5-16	5	1	Validated test (87%)
Schmidt et al ^{39††}	Netherlands	1441	820 (1980-1998)	621 (1980-2003)	1-23	5	18	Self (19%-39%)
Tolnai et al ^{40†}	Canada	342	156 (1974-1980)	186 (1974-1980)	9-15	5	1	Objective survey instrument (50%-58%)
Woodard ^{41,42†§}	Canada	152	105 (1978-1979)	47 (1978-1979)	1	5	8	Supervisor (70%-76%)‡‡

Note: NR = not reported.

*Authors adjusted for possible confounders.

†The problem-based learning group and the traditional group were from different universities.

‡The problem-based learning group was surveyed in 1999, and the traditional group was surveyed in 2004.

§Different methodology was used to select controls between the 1978 and 1979 cohort of graduates.

¶Although the exact number of residents included in the analysis was not stated, these numbers were inferred from data in the paper.

**Although the supervisor response rate was not reported, we assumed it to be > 80% because the authors stated that their school routinely solicited residency program directors' evaluations of graduates at the end of their first year of residency.

††No. of competencies/procedural skills, as categorized in original study.

‡‡Percentage reflects response rate among supervisors of problem-based learning graduates; the response rate among supervisors of graduates in the control group could not be calculated.

Table 2: Outcomes of problem-based learning versus traditional curricula (control) in the 13 studies

Study	Outcome of problem-based learning (type of assessment)			Remarks	Study quality
	Positive	No difference	Negative		
Peters et al ²⁸	Better than control group in humanism and social learning (self)	No difference in lifelong learning (self)		Changes in problem-based learning curriculum contaminated positive findings	High
Hoffman et al ²⁹	Better than control group in 12 competencies (supervisor)	No difference in 5 competencies (supervisor)		The significance level was set at $p < 0.01$ because of the study's large sample	High
Jones et al ³⁰	Better than control group in 11 competencies and 8 procedural skills (self) Better than control group in 5 competencies (supervisor)	No difference in 7 competencies and 5 procedural skills (self) No difference in 13 competencies or in any procedural skills (supervisor)	Control group better in understanding disease processes (self)		High
Rolfe et al ³¹	Better than control group in 4 competencies (supervisor)	No difference in 10 competencies (supervisor)		Curriculum emphasis on interpersonal skills contaminated 2 positive findings	High
Watmough et al ^{32,33}	Better than control group in 9 competencies and 6 procedural skills (self) Better than control group in 11 competencies (supervisor)	No difference in 8 competencies and 7 procedural skills (self) No difference in 6 competencies or in any procedural skills (supervisor)	Control group better in understanding disease processes (self)	Curriculum changes contaminated positive findings on 2 self-rated competencies, all procedural skills and 2 supervisor-rated competencies	High
Tamblyn et al ³⁴	Better than control group in mammography screening rates, no. of coordinated visits and diagnostic accuracy (database)	No difference in rate of prescribing for contraindicated drugs (database)		Curriculum emphasis on social, preventive and community care contaminated 2 positive findings Study surveyed only primary care doctors	High
Mennin et al ³⁵	Better than control group in 12 competencies (self)		Control group better in 5 competencies (self); however, differences were not significant after Bonferroni adjustment	Curriculum included community-based preceptorship, which contaminated 2 positive findings	High
Santos-Gomez et al ³⁶	Better than control group in communication with patients (self) Better than control group in attention to health care costs	No difference in 7 competencies (self, supervisor) No difference in any competencies (nurse)		Although the differences in knowledge between graduates of problem-based learning curricula and the control group were not significant, nurses felt the control group was more knowledgeable	Low
Prince et al ³⁷	Better than control group in 3 competencies (self)	No difference in 9 competencies (self)		Curriculum emphasis on communication skills contaminated 1 positive finding	Low
Shin et al ³⁸	Better than control group in adherence to hypertension management guidelines (test)			Multivariable analysis showed that only the type of curriculum was significant Study surveyed only primary care doctors	Low
Schmidt et al ³⁹	Better than control group in 14 competencies (self)	No difference in 2 competencies (self)	Control group better in knowledge and writing (self)		Low
Tolnai et al ⁴⁰		No difference in rates of participation in continuing medical education (objective)		Control group comprised more family physicians and fewer held postgraduate certifications	Low
Woodard ^{41,42}	Better than control group in global ratings and all competencies (supervisor)			Actual rating values and levels of significance were not reported	Low

*Further details of these studies appear in Appendix 3 (available at www.cmaj.ca/cgi/content/full/178/1/34/DC2)

was moderate evidence in support of problem-based learning for communication and interpersonal skills. In the technical dimension, continuity of care had a strong level of evidence in support of problem-based learning. Of the 12 competencies in the cognitive dimension, coping with uncertainty, use of computers or information resources and understanding of evidence-based medicine had strong levels of evidence for problem-based learning. In the knowledge dimension, there was strong evidence that problem-based learning graduates rated themselves as possessing less medical knowledge than graduates in the control group.

According to self-assessments, 8 competencies had a strong level of evidence in support of problem-based learning: continuity of care, teamwork, appreciation of social and emotional aspects of health care, appreciation of legal and ethical aspects of health care, attitudes toward personal

health and well-being, coping with uncertainty, use of computers or information resources and understanding evidence-based medicine (Table 3). According to the observed assessments, 7 competencies had a strong level of evidence in support of problem-based learning. All but 1 (appreciation of legal and ethical aspects of health care) were different: diagnostic skills, communication skills, appreciation of cultural aspects of health care, responsibility or reliability, and self- or peer appraisal (Table 3). These different findings based on type of assessment were not unexpected because medical education literature suggests physicians have limited self-assessment abilities.^{26,27}

Only 4 competencies had moderate to strong levels of evidence in support of problem-based learning according to both self- and observed assessments: coping with uncertainty (strong), appreciation of legal and ethical aspects of health

Table 3: Level of evidence in support of problem-based learning for self-assessed and observed competencies*

Competency	Level of evidence		Competency	Level of evidence	
	Self-assessed	Observed		Self-assessed	Observed
Overall/global	Weak	Moderate	Cognitive		
Technical			Self-directed or continuing learning	Moderate	Moderate
Clerking (e.g., history-taking, physical examination)	Weak	Moderate	Problem-solving	Weak	Weak
Selection of investigations	None	None	Independence, initiative	Weak	Weak
Diagnostic skills or accuracy	Weak	Strong	Responsibility, reliability	NA	Strong
Approach to management	None	None	Coping with uncertainty	Strong	Strong
Treatment (including prescribing)	None	Weak	Critical thinking	None	Weak
Procedural skills	Weak	None	Awareness of own limitations	None	None
Continuity of care	Strong	None	Ability to work under pressure	None	None
Social			Creativity	Weak	NA
Communication skills	Moderate	Strong	Clinical judgment, reasoning	Weak	None
Interpersonal skills	Moderate	Weak	Use of computers, information resources	Strong	Weak
Teamwork skills	Strong	Weak	Understanding evidence-based medicine	Strong	Weak
Appreciation of social and emotional aspects of health care	Strong	Weak	Managerial		
Appreciation of cultural aspects of health care	Weak	Strong	Time management and organizational skills	Weak	Weak
Appreciation of legal and ethical aspects of health care	Strong	Strong	Keeping accurate records	None	None
Preventive health care and health promotion	Weak	None	Self- or peer appraisal	Weak	Strong
Appropriate attitudes towards personal health and well-being	Strong	None	Attention to health care costs	None	Weak
Research			Knowledge		
Research and presentation skills	Weak	Weak	Possession of medical knowledge	Strong†	None
Writing reports or articles	Weak	NA	Use of medical knowledge	Weak	NA
Teaching					
Teaching medical students	NA	Weak			
Patient education	Weak	None			

Note: NA = not available.

*Further details about the levels of evidence appear in Appendix 4 (self-assessed competencies) and Appendix 5 (observed competencies), which are available online (www.cmaj.ca/cgi/content/full/178/1/34/DC2). Evidence for self-assessed competencies in references 28, 30, 32, 33, 35, 36, 38 and 39. Evidence for observed competencies in references 29-34, 36, 37 and 40-42.

†Level of evidence against graduates of problem-based learning curricula.

care (strong), communication skills (moderate and strong respectively) and self-directed continuing learning (moderate).

According to self-assessments, the knowledge dimension showed a strong level of evidence against problem-based learning in terms of graduates' possession of medical knowledge (Table 2 and Table 3).

Interpretation

Our study offers objective evidence that problem-based learning during medical school has positive effects on physician competencies after graduation, especially in the social and cognitive dimensions. The absence of evidence of positive effects of problem-based learning on physician competencies in the technical and teaching dimensions is likely due to these dimensions requiring psychomotor skills, which are more effectively instilled using deliberate practice methods including formal procedural skills training and preceptorships. Our findings indicated a strong level of evidence against problem-based learning in terms of graduates' self-assessments of their possession of medical knowledge. The meta-analysis by Dochy and colleagues⁴³ found nonrobust evidence that problem-based learning had negative effects on knowledge possession, but robust evidence that the method had positive effects on students' knowledge application. In our study, graduates of problem-based learning curricula assessed themselves as possessing less medical knowledge than graduates from the control group; however, supervisors generally found little difference between the 2 groups because, we suspect, they used knowledge application as their measure of physician knowledge. The fewer number of competencies with strong levels of evidence using observed assessments relative to self-assessment supports the findings by Davis and colleagues²⁷ that physicians have limited self-assessment abilities. We did not find any differences in outcomes for problem-based learning graduates based on the country of study; however, the number of studies from each country was too few to form the basis for any meaningful comparison.

Most studies and systematic reviews of problem-based learning focused on the effect of problem-based learning on knowledge but neglected other aspects of physician competency.^{9,14,43} Controversy has often surrounded the issue of whether problem-based learning results in a weaker knowledge base in students, especially of basic medical sciences.^{11,16} In our study, it is reassuring that, although graduates of problem-based learning curricula assessed themselves as possessing less medical knowledge than graduates of traditional curricula, the evidence from the objective assessments did not support this finding.

Our study has a number of limitations. First, the heterogeneity of problem-based learning is a challenge inherent in all problem-based learning research. The definition and implementation of problem-based learning varies widely among medical schools and educators,¹⁸ and a number of factors that are difficult to measure and control affect the success of problem-based learning curricula.^{44,45} Second, most of the studies used surveys as the main measurement to assess

graduates. Surveys are limited in reliability and validity, subjective and prone to rater biases.⁴⁶ Third, the study-quality scoring system, the evaluation system of levels of evidence and the thematic categorization we used were not validated; however, they were adapted from similar assessment tools used in previous research on problem-based learning^{13,14,24} and physician competency.²⁵ Furthermore, in our model we assigned greater weight to some quality indicators (randomization, sample size, objective assessment and response rate) than to others because we felt that they were of greater importance. We acknowledge that this approach is subjective, but our decisions were based largely on extensive discussion, reasonable assessment and consensus. Fourth, we did not perform a grey literature search, which may have contributed some information on publication bias. Fifth, with few exceptions, the introduction of problem-based learning into medical school curricula was accompanied by other changes such as increased community focus or emphasis on communication skills, which makes it impossible to fully dissociate the findings attributable to problem-based learning from those attributable to other curricula changes. Lastly, because we were unable to determine the adequacy of sample size, there may have been nonsignificant but true differences in outcomes because of small samples.

In summary, problem-based learning during medical school has positive effects on physician competencies, especially in the social and cognitive dimensions. Future research should look beyond knowledge competency and measure the effects of problem-based learning on other dimensions of physician competency.

This article has been peer reviewed.

Competing interests: None declared.

Contributors: All authors were involved in study conception, design and acquisition of data. All authors contributed to data analysis and interpretation, revised the article for important intellectual content and approved the final version for publication.

Acknowledgements: We acknowledge the support of the National University of Singapore Library in providing literature search support. We thank Goh Lee Gan MMed for his help in the early stages of research and Tai Bee Choo PhD for her statistical support. Both Goh Lee Gan and Tai Bee Choo are with the Community, Occupational and Family Medicine Department, Yong Loo Lin School of Medicine, National University of Singapore.

REFERENCES

1. Christopher DF, Harte K, George CF. The implementation of tomorrow's doctors. *Med Educ* 2002;36:282-8.
2. Kinkade S. A snapshot of the status of problem-based learning in US medical schools, 2003-04. *Acad Med* 2005;80:300-1.
3. Donner RS, Bickley H. Problem-based learning: an assessment of its feasibility and cost. *Hum Pathol* 1990;21:881-5.
4. Harden RM. Developments in outcome-based education. *Med Teach* 2002;24:117-20.
5. Mellon AF, Mellon J. Logical debate on problem-based learning. *BMJ* 2006;332:550-1.
6. Sanson-Fisher RW, Lynagh MC. Problem-based learning: A dissemination success story? *Med J Aust* 2005;183:258-60.
7. Williams G, Lau A. Reform of undergraduate medical teaching in the United Kingdom: a triumph of evangelism over common sense. *BMJ* 2004;329:92-4.
8. Farrow R, Norman GR. The effectiveness of PBL. The debate continues: Is meta-analysis helpful? *Med Educ* 2003;37:1131-2.
9. Colliver JA. Effectiveness of problem-based learning curricula: research and theory. *Acad Med* 2000;75:259-66.
10. Schmidt HG, Dauphinee WD, Patel VL. Comparing the effects of problem-based and conventional curricula in an international sample. *J Med Educ* 1987;62:305-15.

11. Vernon DT, Blake RL. Does problem-based learning work? A meta-analysis of evaluative research. *Acad Med* 1993;68:550-63.
12. Nandi PL, Chan JNF, Chan CPK, et al. Undergraduate medical education: comparison of problem-based learning and conventional teaching. *Hong Kong Med J* 2000;6:301-6.
13. Smits PBA, Verbeek JHAM, de Buissonjé CD. Problem based learning in continuing medical education: a review of controlled evaluation studies. *BMJ* 2002;324:153-6.
14. Newman M; Campbell Collaboration Systematic Review Group on the Effectiveness of Problem-based Learning. *A pilot systematic review and meta-analysis on the effectiveness of problem based learning*. Newcastle upon Tyne (UK): University of Newcastle upon Tyne; 2003.
15. Berkson L. Problem-based learning: Have the expectations been met? *Acad Med* 1993;68:S79-88.
16. Albanese MA, Mitchell S. Problem-based learning: a review of literature on its outcomes and implementation issues. *Acad Med* 1993;68:52-81.
17. Colliver JA. Educational theory and medical education practice: a cautionary note for medical school faculty. *Acad Med* 2002;77:1217-20.
18. Maudsley G. Do we all mean the same thing by "problem-based learning"? A review of the concepts and a formulation of the ground rules. *Acad Med* 1999;74:178-85.
19. Newman M, Van der Bossche P, Gijbels D, et al. Responses to the pilot systematic review of problem-based learning. *Med Educ* 2004;38:921-6.
20. Norman G. RCT = results confounded and trivial: the perils of grand educational experiments. *Med Educ* 2003;37:582-4.
21. Berliner DC. Educational research. The hardest science of all. *Educational Researcher* 2002;31:18-20. Available: www.aera.net/uploadedFiles/Journals_and_Publications/Journals/Educational_Researcher/3108/3108_CommentBerliner.pdf (accessed 2007 Nov. 19).
22. Norman GR, Schmidt HG. Effectiveness of problem-based learning curricula: theory, practice and paper darts. *Med Educ* 2000;34:721-8.
23. Wilkinson TJ, Wells JE, Bushnell JA. Are differences between graduates and undergraduates in a medical course due to age or prior degree? *Med Educ* 2004;38:1141-6.
24. Reed D, Price EG, Windish DM, et al. Challenges in systematic reviews of educational intervention studies. *Ann Intern Med* 2005;142:1080-9.
25. Epstein RM, Hundert EM. Defining and assessing professional competence. *JAMA* 2002;287:226-35.
26. Langendyk V. Not knowing that they do not know: self-assessment accuracy of third year medical students. *Med Educ* 2006;40:173-9.
27. Davis DA, Mazmanian PE, Fordis M, et al. Accuracy of physician self-assessment compared with observed measures of competence. A systematic review. *JAMA* 2006;296:1094-102.
28. Peters AS, Greenberger-Rosovsky R, Crowder C, et al. Long-term outcomes of the New Pathway Program at Harvard Medical School: A randomized controlled trial. *Acad Med* 2000;75:470-9.
29. Hoffman K, Hosokawa M, Blake R Jr, et al. Problem-based learning outcomes: Ten years of experience at the University of Missouri-Columbia School of Medicine. *Acad Med* 2006;81:617-25.
30. Jones A, McArdles PJ, O'Neill PAO. Perceptions of how well graduates are prepared for the role of pre-registration house officer: a comparison of outcomes from a traditional and an integrated PBL curriculum. *Med Educ* 2002;36:16-25.
31. Rolfe E, Andren JM, Pearson S, et al. Clinical competence of interns. *Med Educ* 1995;29:225-30.
32. Watmough S, Taylor DC, Garden A, et al. Educational supervisors' views on the competencies of preregistration house officers. *Br J Hosp Med (Lond)* 2006;67:92-5.
33. Watmough S, Ryland I, Taylor DC, et al. Preregistration house officer skill and competency assessment through questionnaires. *Br J Hosp Med (Lond)* 2006;67:487-90.
34. Tambllyn R, Abrahamowicz M, Dauphinee D, et al. Effect of a community oriented problem based learning curriculum on quality of primary care delivered by graduates: historical cohort comparison study. *BMJ* 2005;331:1002-8.
35. Mennin SP, Kalishman S, Friedman M, et al. A survey of graduates in practice from the University of New Mexico's conventional and community-oriented problem-based tracks. *Acad Med* 1996;71:1079-89.
36. Santos-Gomez L, Kalishman S, Rezler A, et al. Residency performance of graduates from a problem-based and a conventional curriculum. *Med Educ* 1990;24:366-75.
37. Prince KJAH, van Eijs PWL, Boshuizen HPA, van der Vleuten CPM, Scherpbier AJJA. General competencies of problem-based learning (PBL) and non-PBL graduates. *Med Educ* 2005;39:394-401.
38. Shin JH, Haynes RB, Johnston ME. Effect of problem-based, self-directed undergraduate education on life-long learning. *CMAJ* 1993;148:969-76.
39. Schmidt HG, Vermeulen L, van der Molen HT. Longterm effects of problem-based learning: a comparison of competencies acquired by graduates of a problem-based and a conventional medical school. *Med Educ* 2006;40:562-7.
40. Tolnai S. Continuing medical education and career choice among graduates of problem-based and traditional curricula. *Med Educ* 1991;25:414-20.
41. Woodard CA, McAuley RG, Ridge H. Unravelling the meaning of global comparative ratings of interns. In: *Research in Medical Education, 1981: Proceedings of the Twentieth Annual Conference*; Washington. Washington: Association of American Medical Colleges; 1981. p. 149-54.
42. Woodard CA. Monitoring an innovation in medical education: the McMaster experience. In: Nooman ZN, Schmidt HG, Ezzat ES, editors. *Innovation in Medical Education: An Evaluation of Its Present Status*. 1st ed. New York, Springer Publishing; 1990. p. 27-39.
43. Dochy F, Segers M, Bossche van den P, Gijbels D. Effects of PBL: a meta-analysis. *Learning and Instruction* 2003;13:533-68.
44. Wood DF. ABC of learning and teaching in medicine: problem based learning. *BMJ* 2003;326:328-30.
45. Harden R, Davis M. The continuum of problem-based learning. *Med Teacher* 1998;20:317-22.
46. Kern DE, Thomas PA, Howard D, et al. *Curriculum development for medical education: A six-step approach*. Baltimore (MD): Johns Hopkins University Press; 1998.

Correspondence to: Dr. Gerald Choon-Huat Koh, Level 3, Block MD3, 16 Medical Dr., Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597; fax 65 6779-1489; cofkohch@nus.edu.sg