



Published in final edited form as:

*Am J Psychiatry*. 2013 May 1; 170(5): 477–484. doi:10.1176/appi.ajp.2012.12081074.

## The Enigmatic Persistence of Anorexia Nervosa

**B. Timothy Walsh, M.D.**

Department of Psychiatry, Columbia University, and the New York State Psychiatric Institute, New York.

### Abstract

**Objective**—In this review, based on recent advances in cognitive neuroscience, the author presents a formulation in which the marked persistence of anorexia nervosa can be usefully understood as a well-ingrained maladaptive habit.

**Method**—The author reviewed the relevant literature on the development and course of anorexia nervosa and interpreted critical features in light of developments in cognitive neuroscience.

**Results**—Anorexia nervosa is a well characterized disorder with remarkable persistence both across history and among affected individuals. Food restriction, the salient behavioral feature of the disorder, often begins innocently but gradually takes on a life of its own. Over time, it becomes highly entrenched and resistant to change through either psychological or pharmacological treatment. Cognitive neuroscience has described two related but distinct processes that underlie the acquisition of new patterns of behavior, namely, action-outcome and stimulus-response learning. It is likely that both processes are engaged in the development of anorexia nervosa and that stimulus-response learning (that is, habit formation) is critical to the persistence of the dieting behavior.

**Conclusions**—The formulation of the dieting behavior characteristic of anorexia nervosa as a well-entrenched habit provides a basis for understanding the striking persistence of this disorder. This model helps explain the resistance of anorexia nervosa to interventions that have established efficacy in related disorders and implies that addressing the dieting behavior is critical, especially early in the course of the illness, before it has become ingrained.

---

For in truth habit is a violent and treacherous schoolmistress. She establishes in us, little by little, stealthily, the foothold of her authority; but having by this mild and humble beginning settled and planted it with the help of time, she soon uncovers to us a furious and tyrannical face against which we no longer have the liberty of even raising our eyes.

—Montaigne (1), cited by Graybiel (2)

Anorexia nervosa is a serious psychiatric disorder with remarkable persistence, both over centuries and among many individuals who develop it. Anorexia nervosa occurs primarily in cultures where food is plentiful and where, for some reason, special value is attached to

---

Address correspondence to Dr. Walsh (btw1@columbia.edu).

Presented in part at the 2012 International Conference on Eating Disorders, Austin, Tex., May 3–5, 2012, and at the 164th annual meeting of the American Psychiatric Association, Philadelphia, May 5–9, 2012.

diETING and weight loss. Its salient features—a relentless pursuit of thinness accompanied by an intense fear of weight gain despite being significantly underweight—have remained constant over time. The first known description of the disorder in the medical literature was provided by Richard Morton in 1689, and it has been speculated that some medieval saints may have suffered with anorexia nervosa (3, 4). In the late 19th century, Sir William Gull coined the term “anorexia nervosa” in his description of three young women who would clearly meet DSM-IV and the proposed DSM-5 criteria for the disorder (5).

The course of anorexia nervosa is also often remarkably persistent. Although the range of outcomes is wide, anorexia nervosa is one of the deadliest of psychiatric disorders, with an estimated mortality rate at least five times higher than that expected (6). Among the few factors associated with a more favorable outcome are adolescent onset (in contrast to adult onset) and a shorter duration of illness (7). The positive significance of an early adolescent onset of anorexia nervosa is in contrast to mood and psychotic disorders, in which early adolescent onset is associated with a more refractory course (8, 9).

Although a number of risk factors have been identified, including being female, being an adolescent, and having an obsessional style, they are not particularly specific, and they account for only a small fraction of the variation in occurrence (10). The overwhelming majority of adolescent girls, even those who tend to be obsessional, do not develop anorexia nervosa. Similarly, while dieting behavior is almost universal among adolescent girls in the United States, it develops into anorexia nervosa in very few. Most cases of anorexia nervosa begin unremarkably. Many young women begin to diet in hopes of losing weight and gaining attractiveness and self-esteem, often in response to one of the ordinary challenges of adolescence, such as beginning to date or leaving home for camp or school. In the weeks or months that follow, among the few who develop anorexia nervosa, the dieting takes on a life of its own and evolves into an unrelenting pursuit that becomes the individual’s primary focus. Early intervention appears to interrupt the increasingly restrictive dieting and growing social isolation. But in many individuals, dieting and weight loss dominate mental life for years and severely impair physical health and social and occupational development.

Efforts at improving outcomes through the development of specific treatment interventions have proven frustrating. One of the few significant advances in recent years is the articulation and promulgation of a family-based treatment approach developed at the Maudsley Hospital in London several decades ago (11, 12). In stark contrast to previous approaches, this method specifically charges the parents of the affected individual with a major role in intervention. Controlled studies support the utility of this approach (13), but it is focused on adolescents with anorexia nervosa, who are those most readily treated. Anorexia nervosa in adults, including those only in their 20s, is generally much more refractory to intervention. Treatments of established efficacy in related disorders, such as bulimia nervosa and mood and anxiety disorders, including cognitive-behavioral therapy and selective serotonin reuptake inhibitors, are surprisingly ineffective (14). Older individuals with anorexia nervosa can be helped to restore normal weight in structured treatment programs, and they may exhibit substantial improvement in both physical and psychological symptoms, but relapse rates after discharge are high (15). Thus, despite better recognition

and presumably more informed treatment approaches, the outcome of anorexia nervosa, particularly for adults, has probably not improved substantially in the past 50 years (7).

These salient facts pose a challenge to our understanding of this disorder. How is it that children and adolescents in generally good health develop an illness that is often chronic, severe, and possibly deadly but that may, especially if treated early, fully remit without any long-term consequences? In this review I extend an earlier discussion (16) and address one important facet of this enigma, namely, the remarkable persistence of anorexia nervosa once it has become established.

## **Eating Behavior and the Persistence of Anorexia Nervosa**

Individuals with anorexia nervosa achieve and maintain a significantly low body weight primarily by dieting and, in a significant fraction of individuals, by engaging in excessive physical activity. (Although in this review I focus on the dieting behavior for the sake of clarity of exposition, the discussion can readily be extended to include increased physical activity as well.)

Among the striking characteristics of anorexia nervosa is the homogeneity of disturbances in eating behavior across affected individuals. Individuals with the disorder uniformly restrict their caloric intake by consuming a limited variety of foods that are low in energy density. Once established, such behavior—which has been documented in laboratory settings—is highly resistant to change, persisting even after normal weight has been restored in intensive treatment programs through consumption of a varied diet typical of individuals without eating disorders (17–20). Recent work by our group strongly suggests that the persistence of such behavior is a major contributor to relapse (15, 21, 22).

Therefore, an improved understanding of why it is so difficult for individuals with anorexia nervosa to alter their eating behavior is critical to improving our understanding of the development of the disorder and to improving treatment interventions. The formulation I present here attempts to explain how such dieting behavior becomes established and persistent. Its central element is that this behavior has become habitual and is grounded in the neural mechanisms that underlie habit formation and persistence. This formulation is based on a substantial body of animal and human research, much of which is focused on the neural mechanisms underlying substance abuse.

## **Dieting Behavior in Anorexia Nervosa Is Habitual**

Graybiel (2) provided a useful working definition of habits from the perspective of cognitive neuroscience: habits are behaviors that are not innate, that are engaged in repeatedly and become fixed, that appear to occur without conscious effort, and that are elicited by a variety of stimuli. The dieting behavior characteristic of anorexia nervosa fulfills this definition. Although there are hints that some individuals who develop anorexia nervosa may have been picky eaters as children (23), there are no indications that eating behavior or rate of weight gain are abnormal prior to the onset of illness. In order to lead to significant weight loss, the dieting behavior must be engaged in repeatedly, over weeks to months, and the studies of eating behavior cited above demonstrate that it becomes markedly fixed (18, 20).

Individuals with anorexia nervosa appear to diet without obvious attempts to contemplate the specific elements of eating behavior; conversely, attempts to consume a more varied and calorically dense diet require substantial effort (24). The characteristic food choices are elicited by a variety of stimuli, including the presentation of foods at mealtimes (17, 18), and by emotional stress (25, 26).

## How Habits Develop

Substantial research over the past several decades has articulated two related but distinct processes responsible for the acquisition and persistence of behavior that is not innate: action-outcome and stimulus-response learning (2, 27–29). Much of this research has been carried out using laboratory animals, especially rodents, but the key findings have also been established in humans.

In action-outcome learning, also referred to as instrumental or operant conditioning, an animal detects that some action, such as pushing a lever, leads to a reward, such as the presentation of food, and the likelihood of the animal's engaging in the behavior is sensitive to the reward value of the outcome. Hence, if at some point the behavior no longer leads to the receipt of the reward, its frequency will diminish, and eventually it will occur only sporadically. This form of learning is critical to the acquisition of new behaviors, and it involves specific neural pathways, including the amygdala, the ventral striatum (the nucleus accumbens), and the orbitofrontal cortex.

While new behaviors are typically acquired through action-outcome learning, if they are repeated and continue to lead to the acquisition of rewards, the behaviors will eventually become relatively insensitive to the receipt of the reward. This process, referred to as stimulus-response learning, is the basis for habit formation. The key neural structures involved are neuroanatomically close to, but distinct from, those required for action-outcome learning: the dorsolateral striatum (the caudate/putamen) and the dorsolateral prefrontal cortex.

A critical element in stimulus-response learning is overtraining—the repetition of the behavior. A popular term referring to this phenomenon in athletic pursuits is the development of “muscle memory,” acquired, for example, by repeatedly practicing a baseball swing. “Muscle memory” is a misnomer, as muscles have no capacity for memory; it is the brain that (hopefully) learns a productive swing cued by the appearance of a baseball.

In the model presented here, which is based on these principles of learning, it is hypothesized that the dieting behaviors of individuals with anorexia nervosa begin as goal-directed actions that lead to weight loss, which is highly rewarding (action-outcome learning). Over time, the dieting behaviors are engaged in persistently and repeatedly and thereby become overtrained and habitual (stimulus-response learning). Once they have become established as habitual, the behaviors are highly resistant to change and are a critical element in the persistence of the disorder. The next sections describe critical elements in this learning process.

## Dieting Behavior Is Strongly Reinforced

By definition, individuals with anorexia nervosa diet effectively enough to lose weight (or, if still growing, to fail to gain weight as expected). In the current Western culture, successful weight loss is an often wished for and encouraged goal that is rarely achieved. Therefore, the initial weight loss of individuals with anorexia nervosa provides evidence of impressive self-control and personal accomplishment, leading to enhanced self-esteem. In addition, many individuals who develop anorexia nervosa describe receiving compliments at the beginning of their successful efforts at weight loss.

Such feelings about the emotional significance of weight loss are well known to clinicians experienced in the care of individuals with anorexia nervosa and have been documented in numerous descriptions of the disorder (30–32). Bruch (33), writing in the latter half of the 20th century, provided some of the most compelling accounts of the symptoms and internal motivations of individuals with anorexia nervosa. Describing the perspective of one individual, Bruch wrote, “That she could be so thin gave her a sense of pride, power, and accomplishment” (p. 2). Another individual “explained that losing weight was giving her power, that each pound lost was like a treasure that added to her power” (pp. 4–5). Garfinkel and Garner (34), in their important volume on anorexia nervosa, noted, “While anorexia nervosa patients have an overriding drive to be thin, they do not offer any explanation; they merely say they will feel better the thinner they are” (p. 3). Thus, the dieting behavior leads to weight loss, which is highly rewarding and is thereby positively reinforced.

Individuals with anorexia nervosa frequently report that dieting behavior also helps them cope with negative affect, and recent studies have begun to document this phenomenon experimentally. Engel et al. (25), using ecological momentary assessment, found that increased levels of negative affect were followed by restriction of food intake. Wildes et al. (26) found that induction of negative affect in a laboratory setting led to an increase in self-reported eating disorder symptoms, although intake of a test meal was not altered. These observations suggest that in individuals with anorexia nervosa, dieting behavior may be triggered by negative affect and serves as a method of modulating emotional distress. The relief of negative affect thus likely constitutes an additional, negative reinforcer of dieting behavior. In other words, dieting behavior is rewarded both by leading to weight loss and enhanced self-esteem and by helping individuals cope with emotional difficulties.

## Dieting Itself Becomes Rewarding

It is likely that in anorexia nervosa, dieting behavior itself becomes rewarding through a phenomenon called “conditioned reinforcement” (35). Conditioned reinforcement occurs when a stimulus or situation is regularly paired with an established reward. Such phenomena are viewed as a critical element in the development of the complex set of cues that, over time, become associated with craving for drugs of abuse and of the rituals associated with drug taking themselves becoming rewarding (36, 37). Since in anorexia nervosa dieting behavior is paired with a potent reward—weight loss—it is likely that, over time, dieting behavior itself takes on rewarding properties. Such a phenomenon helps to explain the distinction between “good” (low-calorie) and “bad” (high-calorie) foods described with

remarkable consistency by individuals with anorexia nervosa and the difficulties they experience in making even minor changes in their dieting practices.

## Rewards Are More Rewarding During Adolescence

Most cases of anorexia nervosa begin during adolescence, a time of dramatic psychological and physical development. During the teens and into the 20s, substantial cortical development contributes importantly to emotional and cognitive maturation—for example, to the ability to postpone the receipt of rewards (38). Many psychiatric disorders emerge during this phase of life, undoubtedly reflecting a complex interplay of environmental, social, and internal factors (39). Adolescents are disproportionately sensitive to the receipt of reward (40). For example, the response of the ventral striatum (a site well established in the assessment of the potential reward value of stimuli) to rewarding stimuli has been shown to be greater in adolescents than in children and adults, suggesting that the same reward is assessed as being more rewarding at this stage of development (41). The attraction of rewarding stimuli, coupled with the relative immaturity of inhibitory neural systems, has been suggested as an important factor underlying the vulnerability of adolescents to developing substance abuse (42), and it may similarly contribute to the vulnerability to anorexia nervosa during this stage of development.

## The Dieting Behavior Becomes Habitual

The preceding sections have made the case that at the onset of anorexia nervosa, weight loss is highly rewarding. Thus, the acquisition of dieting behavior is a classic example of action/outcome learning. Other factors lead to these behaviors becoming habitual.

First, in order to lead to significant weight loss, dieting behavior must be engaged in persistently and repeatedly. The behavior becomes highly practiced and overtrained, thereby becoming more entrenched. This sequence is the classic technique producing habit learning (2).

Second, the rewards of dieting behavior are intermittent. Daily weight loss is not inevitable, and the receipt of social accolades for becoming thinner occurs only occasionally. Skinner's early work on instrumental conditioning established that an intermittent and unanticipated pattern of reinforcement encourages establishment of habitual behavior (43, 44).

Third, the onset of anorexia nervosa typically begins during a period of stress. Most of the stressors are those routinely encountered during adolescence, but they are nonetheless emotionally compelling at the time they are experienced. Both rodent and human studies have demonstrated that behaviors learned during a period of stress are more likely to become habitual than those learned at other times (45, 46).

Finally, by definition, anorexia nervosa involves significant weight loss. There is good evidence that substantial weight loss enhances the tendency to develop compulsive patterns of behavior. The Minnesota starvation study conducted during World War II (47) documented the development of irrational compulsive behaviors in previously healthy subjects during a period of starvation. Conversely, in anorexia nervosa, weight gain is

associated with improvement in symptoms of obsessionality (48). Furthermore, a substantial body of work by Carr et al. (49, 50) has established that weight loss enhances the rewarding effects of drugs of abuse and up-regulates mechanisms of synaptic plasticity in the nucleus accumbens. These findings suggest a mechanistic basis for the development of particularly well-entrenched habits, such as resistance of cocaine-conditioned behavior to extinction in food-restricted rodents (51).

## Relationship to Other Models

Many models have been suggested to account for the development and persistence of anorexia nervosa. Strober (52) hypothesized that individuals who develop anorexia nervosa have an underlying propensity to anxiety and to fear-learning and a greater resistance to its extinction than unaffected individuals. Like the model presented here, Strober's model is based on relatively recent advances in our understanding of how behavior is acquired and is shaped by internal and external stimuli. It suggests that negative emotion, specifically anxiety, about weight gain leads to enduring patterns of dieting. The model presented here similarly suggests that dieting and weight loss become learned techniques to relieve anxiety and other negative emotions but that this development is only one of several factors contributing to such behaviors becoming strongly rewarding. Both models suggest that once acquired, dieting behavior in anorexia nervosa is remarkably resistant to extinction, but the formulation I present proposes that this resistance, to a substantial degree, reflects the establishment of well-entrained stimulus-response learning—that is, habits. As hypothesized by Strober, there may also be an innate resistance to the extinction of learned behaviors.

Other recent models have focused on the roles of emotional regulation (53–55) and of cognitive and cognitive-behavioral disturbances (56, 57). While generally acknowledging that weight loss and dieting are highly reinforcing for individuals with anorexia nervosa and are useful in responding to negative emotion, these models do not grapple in detail with precisely how, in terms of fundamental processes of learning, such behaviors become so rewarding and persistent.

An array of CNS disturbances have been described in anorexia nervosa that may affect the processing of a range of stimuli, including ones linked to reward (58–60), but the models based on these findings do not articulate how such disturbances specifically lead to the persistence of dieting behavior.

The model proposed here does not directly challenge or contradict these other proposals. Rather, it places them in a broader context by suggesting specific mechanisms through which weight loss and dieting behavior become intensely rewarding and, eventually, habitual. Once acquired, these learned behaviors may well be utilized to cope with negative emotion and be sustained by innate or acquired biological disturbances affecting the CNS.

## Implications

The proposed model has a number of implications. The hypothesis that the core behaviors characteristic of anorexia nervosa are habitual helps account for the established observation that the disorder is much more readily treated in younger individuals with shorter durations

of illness (7); in such individuals, complete recovery is a routine outcome. With a shorter duration of illness, the individual has had less time to overtrain the dieting behavior, and the establishment of dieting behavior as rewarding in its own right is less secure.

This formulation also supports the recommendation that interventions focusing on interrupting the behavior and restoring weight to normal are crucial in achieving full recovery. The mechanisms described suggest that weight loss and dieting behavior have acquired rewarding value and are likely to persist unless directly addressed. It is notable that the first phase in family-based treatment, an intervention of documented utility for younger patients, emphasizes the need for parents to actively refeed their child, that is, to interrupt the dieting behavior (12). It may also be useful to help patients recognize that dieting behavior has become largely automatic and may be triggered by a wide range of external and internal cues, including negative affective states. For patients with long-standing anorexia nervosa, it may be important to enhance current interventions by adapting techniques employed in habit reversal therapy that have been found to be useful in addressing disorders such as trichotillomania and Tourette's disorder (61).

The suggested neural basis underlying the development of persistent dieting behavior in anorexia nervosa is fundamentally identical to that proposed to account for the development of persistent abuse of inherently rewarding substances such as cocaine (2). These neurobiological similarities provide a compelling explanation for the striking clinical similarities between the behavior of individuals with substance use disorders and that of individuals with anorexia nervosa. In both disorders, individuals become intensely focused on the desired result (acquisition of the abused substance or weight loss, respectively) to the exclusion of occupational and social activities, may engage in deceitful behavior to attain it (stealing money to buy drugs or lying about whether or what they have eaten), and sacrifice physical health in pursuit of the rewarding goal. Indeed, Overeaters Anonymous specifically includes individuals with anorexia nervosa among those who may be helped by its 12-step program (62). But the model proposed here in no way suggests that individuals with anorexia nervosa are "addicted" to a substance or a food. Rather, the basis of the similarity is that the same neural structures are engaged in the acquisition of dieting behavior as habitual and in the chronic abuse of rewarding substances.

Indirectly, the proposed model may offer at least a partial explanation for the surprising relative ineffectiveness of interventions such as cognitive-behavioral therapy, interpersonal therapy, and antidepressants for anorexia nervosa (63–65). Such interventions are, in general, quite effective in treating affective disorders and related eating disorders, such as bulimia nervosa. However, there is no reason to believe that they possess substantial utility in addressing well-entrenched habits of behavior. Similar considerations may help explain the relative ineffectiveness of such treatments for substance abuse.

The proposed model provides an explanation for the occurrence of anorexia nervosa over many centuries. The critical element is that dieting and weight loss are valued and therefore capable of providing a potent subjective reward. In current Western society, dieting and being thin are equated with health and, especially for women, with attractiveness. In the



middle ages, other rewards of food restriction were prominent, including as a route to asceticism and spiritual purity (4).

The proposed model suggests that distinct neural circuits are involved in the acquisition and perpetuation of the dieting behavior characteristic of anorexia nervosa. In the early stages of illness, as weight loss and dieting behavior are recognized and “coded” as rewarding, the amygdala, the ventral striatum, and the orbitofrontal cortex are engaged. Over time, however, as the dieting behavior becomes habitual, although these centers remain involved, the dorsal striatum and the dorsolateral prefrontal cortex play a more critical role. Given the increasing sophistication of experimental paradigms using functional MRI to identify the relative contributions of different neural circuits in supporting decisions, including about what to eat, it should be possible to examine these hypotheses in individuals restricting their food intake, ranging from healthy dieters to individuals with chronic and severe anorexia nervosa (66).

It is possible that the key elements of the proposed model might be extended to help us understand the persistence of maladaptive behaviors in other psychiatric conditions. If a behavioral pattern develops and is repetitively engaged in at a vulnerable and malleable time of life, such as during adolescence, and is initially associated with a powerful reward, such as relief from an intensely negative emotion, it may become habitual, especially if, at least intermittently, it continues to be associated with the receipt of a reward. The application of this notion is straightforward if the reward is obvious, such as taking a drug of abuse or consuming highly palatable food in bulimia nervosa or binge eating disorder. But perhaps this sequence also plays a role in the development and persistence of maladaptive interpersonal relationships and risky behaviors.

## Summary

The developmental and psychosocial context in which anorexia nervosa develops sets the stage for weight loss and, over time, the characteristic dieting behaviors themselves to become intensely rewarding. Several additional factors lead to dieting behavior becoming a well-established habit and very resistant to change. This formulation thereby provides a possible insight into the impressive persistence of anorexia nervosa once it has developed. However, other factors, both innate and acquired, undoubtedly also contribute to its chronicity, including biological, psychological, and environmental influences that reduce emotional and behavioral flexibility.

It is also important to emphasize that this proposal addresses only one facet of a complex disorder. For example, it does not directly address the difficult but critical issue of vulnerability: Why do only a very small number of individuals who begin to diet during adolescence go on to develop anorexia nervosa? The proposed model raises the possibility that among the risk factors is an innate ability to establish habitual behaviors or, conversely, a difficulty in extinguishing well-practiced behaviors. An association between anorexia nervosa and both obsessive-compulsive disorder and obsessive-compulsive personality disorder is well documented, and there is good evidence that the symptoms of individuals with anorexia nervosa extend beyond issues related to eating or weight (10, 67). Relevant to

the formulation proposed here, a recent study focusing on obsessive-compulsive disorder found evidence that affected individuals, relative to healthy comparison subjects, rely more on habitual patterns of response and are less able to adjust their behavior to changing contingencies (68). Vulnerability may also be increased by individuals' difficulties in managing some of the emotional challenges of adolescence and by environmental influences, such as the perceived advantages of strict dieting. Presumably, such innate and external factors set the stage for the development and persistence of relentless dieting.

The proposed model focuses on the development and persistence of the salient behavioral feature of anorexia nervosa—severe restriction of food intake—and does not specifically address other important features of the disorder, such as the emergence of binge eating, a phenomenon that, over time, occurs in approximately one-half of those with anorexia nervosa (69). It is important to recognize that despite the occurrence of episodes of binge eating, often followed by purging, these individuals continue to restrict their food intake in order to maintain their low weight (70). Since such individuals are prone to engage as well in impulsive behaviors other than binge eating, it seems reasonable to hypothesize that these phenomena are a manifestation of difficulties with inhibiting responses to naturally rewarding stimuli, including palatable food, after long periods of starvation. Such phenomena were well documented among the men who participated in the Minnesota starvation study (71).

Each of the elements underlying the proposed model has a basis in fact, but the resulting formulation requires elaboration and testing. The propensity of individuals with anorexia nervosa to develop and extinguish habits when underweight and after short-term and long-term recovery should be examined. The activity of neural centers involved in the recognition of rewards and the development of habits is certainly relevant. It may be useful to examine such phenomena in animal models. Finally, the articulation and testing of treatment interventions based on the hypothesized mechanisms would be of potential heuristic importance and clinical utility.

## Acknowledgments

Dr. Walsh receives research support from AstraZeneca.

Supported in part by NIMH grant MH79397.

The following individuals provided helpful comments on earlier versions of this manuscript: Evelyn Attia, Peter Balsam, Allegra Broft, Kenneth Carr, Michael Devlin, Rudolph Leibel, Margaret Martinez, Laurel Mayer, Daphna Shohamy, Joanna Steinglass, and Stephen Wonderlich.

## References

1. Montaigne, M.; Frame, DM. *The Complete Essays of Montaigne*. Stanford: Stanford University Press; 1958. Of Custom, and Not Easily Changing an Accepted Law; p. 77-90. Translated by
2. Graybiel AM. Habits, rituals, and the evaluative brain. *Annu Rev Neurosci*. 2008; 31:359–387. [PubMed: 18558860]
3. Pearce JM. Richard Morton: origins of anorexia nervosa. *Eur Neurol*. 2004; 52:191–192. [PubMed: 15539770]
4. Bell, RM. *Holy Anorexia*. Chicago: University of Chicago Press; 1985.

5. Gull WW. Anorexia nervosa (apepsia hysterica, anorexia hysterica), 1868. *Obes Res.* 1997; 5:498–502. [PubMed: 9385628]
6. Arcelus J, Mitchell AJ, Wales J, Nielsen S. Mortality rates in patients with anorexia nervosa and other eating disorders: a meta-analysis of 36 studies. *Arch Gen Psychiatry.* 2011; 68:724–731. [PubMed: 21727255]
7. Steinhausen HC. The outcome of anorexia nervosa in the 20th century. *Am J Psychiatry.* 2002; 159:1284–1293. [PubMed: 12153817]
8. Remschmidt H, Theisen F. Early-onset schizophrenia. *Neuropsychobiology.* 2012; 66:63–69. [PubMed: 22797279]
9. de Girolamo G, Dagani J, Purcell R, Cocchi A, McGorry PD. Age of onset of mental disorders and use of mental health services: needs, opportunities, and obstacles. *Epidemiol Psychiatr Sci.* 2012; 21:47–57. [PubMed: 22670412]
10. Jacobi C, Hayward C, de Zwaan M, Kraemer HC, Agras WS. Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychol Bull.* 2004; 130:19–65. [PubMed: 14717649]
11. Russell GF, Szmulker GI, Dare C, Eisler I. An evaluation of family therapy in anorexia nervosa and bulimia nervosa. *Arch Gen Psychiatry.* 1987; 44:1047–1056. [PubMed: 3318754]
12. Lock, J.; Le Grange, D.; Agras, WS.; Dare, C. *Treatment Manual for Anorexia Nervosa: A Family-Based Approach.* New York: Guilford; 2001.
13. Lock J, Le Grange D, Agras WS, Moya A, Bryson SW, Jo B. Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. *Arch Gen Psychiatry.* 2010; 67:1025–1032. [PubMed: 20921118]
14. Attia E. Anorexia nervosa: current status and future directions. *Annu Rev Med.* 2010; 61:425–435. [PubMed: 19719398]
15. Kaplan AS, Walsh BT, Olmsted M, Attia E, Carter JC, Devlin MJ, Pike KM, Woodside B, Rockert W, Roberto CA, Parides M. The slippery slope: prediction of successful weight maintenance in anorexia nervosa. *Psychol Med.* 2009; 39:1037–1045. [PubMed: 18845008]
16. Steinglass J, Walsh BT. Habit learning and anorexia nervosa: a cognitive neuroscience hypothesis. *Int J Eat Disord.* 2006; 39:267–275. [PubMed: 16523472]
17. Hadigan CM, Anderson EJ, Miller KK, Hubbard JL, Herzog DB, Klibanski A, Grinspoon SK. Assessment of macronutrient and micronutrient intake in women with anorexia nervosa. *Int J Eat Disord.* 2000; 28:284–292. [PubMed: 10942914]
18. Mayer LE, Schebendach J, Bodell LP, Shingleton RM, Walsh BT. Eating behavior in anorexia nervosa: before and after treatment. *Int J Eat Disord.* 2012; 45:290–293. [PubMed: 21495053]
19. Schebendach JE, Mayer LE, Devlin MJ, Attia E, Contento IR, Wolf RL, Walsh BT. Food choice and diet variety in weight-restored patients with anorexia nervosa. *J Am Diet Assoc.* 2011; 111:732–736. [PubMed: 21515121]
20. Sysko R, Walsh BT, Schebendach J, Wilson GT. Eating behavior among women with anorexia nervosa. *Am J Clin Nutr.* 2005; 82:296–301. [PubMed: 16087971]
21. Schebendach J, Mayer LE, Devlin MJ, Attia E, Walsh BT. Dietary energy density and diet variety as risk factors for relapse in anorexia nervosa: a replication. *Int J Eat Disord.* 2012; 45:79–84. [PubMed: 21448937]
22. Schebendach JE, Mayer LE, Devlin MJ, Attia E, Contento IR, Wolf RL, Walsh BT. Dietary energy density and diet variety as predictors of outcome in anorexia nervosa. *Am J Clin Nutr.* 2008; 87:810–816. [PubMed: 18400701]
23. Kotler LA, Cohen P, Davies M, Pine DS, Walsh BT. Longitudinal relationships between childhood, adolescent, and adult eating disorders. *J Am Acad Child Adolesc Psychiatry.* 2001; 40:1434–1440. [PubMed: 11765289]
24. Murphy R, Straebl S, Cooper Z, Fairburn CG. Cognitive behavioral therapy for eating disorders. *Psychiatr Clin North Am.* 2010; 33:611–627. [PubMed: 20599136]
25. Engel, SG.; Crosby, RD.; Wonderlich, SA.; Crow, S.; Le Grange, D.; Mitchell, JE.; Peterson, CB. Ecological momentary assessment of eating disorder behavior and affect in samples of AN and BN participants. Scientific Program and Abstracts, XVIth Annual Meeting of the Eating Disorders

- Research Society; October 7–9, 2010; Cambridge, Mass. Glenview, Ill: Eating Disorders Research Society; 2010.
26. Wildes JE, Marcus MD, Bright AC, Dapelo MM, Psychol MC. Emotion and eating disorder symptoms in patients with anorexia nervosa: an experimental study. *Int J Eat Disord.* 2012; 45:876–882. [PubMed: 22473650]
  27. Rolls, ET. *Emotion Explained.* Oxford, UK: Oxford University Press; 2005.
  28. Shohamy D. Learning and motivation in the human striatum. *Curr Opin Neurobiol.* 2011; 21:408–414. [PubMed: 21658933]
  29. Balleine BW, O’Doherty JP. Human and rodent homologues in action control: corticostriatal determinants of goal-directed and habitual action. *Neuropsychopharmacology.* 2010; 35:48–69. [PubMed: 19776734]
  30. Arkell J, Robinson P. A pilot case series using qualitative and quantitative methods: biological, psychological, and social outcome in severe and enduring eating disorder (anorexia nervosa). *Int J Eat Disord.* 2008; 41:650–656. [PubMed: 18446832]
  31. Nordbø RH, Espeset EM, Gulliksen KS, Skårderud F, Holte A. The meaning of self-starvation: qualitative study of patients’ perception of anorexia nervosa. *Int J Eat Disord.* 2006; 39:556–564. [PubMed: 16845678]
  32. Skårderud F. Shame and pride in anorexia nervosa: a qualitative descriptive study. *Eur Eat Disord Rev.* 2007; 15:81–97. [PubMed: 17676677]
  33. Bruch, H. *The Golden Cage: The Enigma of Anorexia Nervosa.* New York: Random House; 1978.
  34. Garfinkel, PE., Garner, DM. *Anorexia Nervosa: A Multidimensional Perspective.* New York: Brunner/Mazel; 1982.
  35. Everitt BJ, Robbins TW. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci.* 2005; 8:1481–1489. [PubMed: 16251991]
  36. Wikler A. Dynamics of drug dependence: implications of a conditioning theory for research and treatment. *Arch Gen Psychiatry.* 1973; 28:611–616. [PubMed: 4700675]
  37. O’Brien CP. Experimental analysis of conditioning factors in human narcotic addiction. *Pharmacol Rev.* 1975; 27:533–543. [PubMed: 1223916]
  38. Johnson SB, Blum RW, Giedd JN. Adolescent maturity and the brain: the promise and pitfalls of neuroscience research in adolescent health policy. *J Adolesc Health.* 2009; 45:216–221. [PubMed: 19699416]
  39. Paus T, Keshavan M, Giedd JN. Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci.* 2008; 9:947–957. [PubMed: 19002191]
  40. Steinberg L. Risk taking in adolescence: what changes, and why? *Ann NY Acad Sci.* 2004; 1021:51–58. [PubMed: 15251873]
  41. Somerville LH, Hare T, Casey BJ. Frontostriatal maturation predicts cognitive control failure to appetitive cues in adolescents. *J Cogn Neurosci.* 2011; 23:2123–2134. [PubMed: 20809855]
  42. Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *Am J Psychiatry.* 2003; 160:1041–1052. [PubMed: 12777258]
  43. Humphreys LG. The effect of random alternation of reinforcement on the acquisition and extinction of conditioned eyelid reactions. *J Exp Psychol.* 1939; 25:141–158.
  44. Skinner, BF. *The Behavior of Organisms: An Experimental Analysis.* New York: Appleton-Century; 1938.
  45. Dias-Ferreira E, Sousa JC, Melo I, Morgado P, Mesquita AR, Cerqueira JJ, Costa RM, Sousa N. Chronic stress causes frontostriatal reorganization and affects decision-making. *Science.* 2009; 325:621–625. [PubMed: 19644122]
  46. Schwabe L, Wolf OT. Stress prompts habit behavior in humans. *J Neurosci.* 2009; 29:7191–7198. [PubMed: 19494141]
  47. Keys, A. *The Biology of Human Starvation.* Minneapolis: University of Minnesota Press; 1950.
  48. Olatunji BO, Tart CD, Shewmaker S, Wall D, Smits JA. Mediation of symptom changes during inpatient treatment for eating disorders: the role of obsessive-compulsive features. *J Psychiatr Res.* 2010; 44:910–916. [PubMed: 20359715]

49. Carr KD. Chronic food restriction: enhancing effects on drug reward and striatal cell signaling. *Physiol Behav.* 2007; 91:459–472. [PubMed: 17081571]
50. Carr KD, Chau LS, Cabeza de Vaca S, Gustafson K, Stouffer M, Tukey DS, Restituto S, Ziff EB. AMPA receptor subunit GluR1 downstream of D-1 dopamine receptor stimulation in nucleus accumbens shell mediates increased drug reward magnitude in food-restricted rats. *Neuroscience.* 2010; 165:1074–1086. [PubMed: 19931598]
51. Zheng D, Cabeza de Vaca S, Carr KD. Food restriction increases acquisition, persistence, and drug prime-induced expression of a cocaine-conditioned place preference in rats. *Pharmacol Biochem Behav.* 2012; 100:538–544. [PubMed: 22074687]
52. Strober M. Pathologic fear conditioning and anorexia nervosa: on the search for novel paradigms. *Int J Eat Disord.* 2004; 35:504–508. [PubMed: 15101066]
53. Hatch A, Madden S, Kohn M, Clarke S, Touyz S, Williams LM. Anorexia nervosa: towards an integrative neuroscience model. *Eur Eat Disord Rev.* 2010; 18:165–179. [PubMed: 20443202]
54. Haynos AF, Fruzzetti AE. Anorexia nervosa as a disorder of emotion dysregulation: evidence and treatment implications. *Clin Psychol Sci Pract.* 2011; 18:183–202.
55. Wildes JE, Marcus MD. Development of emotion acceptance behavior therapy for anorexia nervosa: a case series. *Int J Eat Disord.* 2011; 44:421–427. [PubMed: 20721894]
56. Fairburn CG, Harrison PJ. Eating disorders. *Lancet.* 2003; 361:407–416. [PubMed: 12573387]
57. Schmidt U, Treasure J. Anorexia nervosa: valued and visible: a cognitive-interpersonal maintenance model and its implications for research and practice. *Br J Clin Psychol.* 2006; 45:343–366. [PubMed: 17147101]
58. Kaye WH, Fudge JL, Paulus M. New insights into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci.* 2009; 10:573–584. [PubMed: 19603056]
59. Keating C. Theoretical perspective on anorexia nervosa: the conflict of reward. *Neurosci Biobehav Rev.* 2010; 34:73–79. [PubMed: 19619579]
60. Frank GK, Reynolds JR, Shott ME, Jappe L, Yang TT, Tregellas JR, O'Reilly RC. Anorexia nervosa and obesity are associated with opposite brain reward response. *Neuropsychopharmacology.* 2012; 37:2031–2046. [PubMed: 22549118]
61. Bate KS, Malouff JM, Thorsteinsson ET, Bhullar N. The efficacy of habit reversal therapy for tics, habit disorders, and stuttering: a meta-analytic review. *Clin Psychol Rev.* 2011; 31:865–871. [PubMed: 21549664]
62. Overeaters Anonymous: OA Members Come in All Sizes. Rio Rancho, NM: Overeaters Anonymous; 2005.
63. McIntosh VV, Jordan J, Carter FA, Luty SE, McKenzie JM, Bulik CM, Frampton CM, Joyce PR. Three psychotherapies for anorexia nervosa: a randomized, controlled trial. *Am J Psychiatry.* 2005; 162:741–747. [PubMed: 15800147]
64. Walsh BT, Kaplan AS, Attia E, Olmsted M, Parides M, Carter JC, Pike KM, Devlin MJ, Woodside B, Roberto CA, Rockert W. Fluoxetine after weight restoration in anorexia nervosa: a randomized controlled trial. *JAMA.* 2006; 295:2605–2612. [PubMed: 16772623]
65. Attia E, Haiman C, Walsh BT, Flater SR. Does fluoxetine augment the inpatient treatment of anorexia nervosa? *Am J Psychiatry.* 1998; 155:548–551. [PubMed: 9546003]
66. Hare TA, Malmaud J, Rangel A. Focusing attention on the health aspects of foods changes value signals in vmPFC and improves dietary choice. *J Neurosci.* 2011; 31:11077–11087. [PubMed: 21795556]
67. Bastiani AM, Altemus M, Pigott TA, Rubenstein C, Weltzin TE, Kaye WH. Comparison of obsessions and compulsions in patients with anorexia nervosa and obsessive compulsive disorder. *Biol Psychiatry.* 1996; 39:966–969. [PubMed: 9162209]
68. Gillan CM, Pappmeyer M, Morein-Zamir S, Sahakian BJ, Fineberg NA, Robbins TW, de Wit S. Disruption in the balance between goal-directed behavior and habit learning in obsessive-compulsive disorder. *Am J Psychiatry.* 2011; 168:718–726. [PubMed: 21572165]
69. Eddy KT, Keel PK, Dorer DJ, Delinsky SS, Franko DL, Herzog DB. Longitudinal comparison of anorexia nervosa subtypes. *Int J Eat Disord.* 2002; 31:191–201. [PubMed: 11920980]
70. Crosby, R.; Steffen, KJ.; Crow, S.; Peterson, CB.; Le Grange, D.; Mitchell, JE. Binge-purge anorexics engage in more restriction than restricting anorexics. *Programs and Abstracts, XVIIIth*

Annual Meeting of the Eating Disorders Research Society; September 20–22, 2012; Porto, Portugal. Glenview, Ill: Eating Disorders Research Society; 2012.

71. Schiele BC, Brozek J. Experimental neurosis resulting from semistarvation in man. *Psychosom Med.* 1948; 10:31–50. [PubMed: 18905659]

**Clinical Guidance: Anorexia Nervosa Viewed as Habitual Behavior**

Many patients with anorexia nervosa report that dieting helps them cope with negative affect, but once the behavior is paired with a desired outcome, it may become rewarding itself. This formulation of anorexia nervosa by Walsh is consistent with findings that adolescent onset and shorter duration of illness are among the few factors associated with better outcome. Cognitive-behavioral therapy, interpersonal therapy, and antidepressants are ineffective, whereas interrupting the behavior and restoring normal weight are considered crucial. Younger patients often respond to family-based treatment, in which parents refeed the child to interrupt the dieting behavior. Patients with long-standing anorexia nervosa may be helped by adding habit-reversal techniques to current interventions.