

RESEARCH

Open Access

The existence of solutions for a nonlinear mixed problem of singular fractional differential equations

Dumitru Baleanu^{1,2,3*}, Hakimeh Mohammadi⁴ and Shahram Rezapour⁴

*Correspondence:

dumitru@cankaya.edu.tr

¹Department of Chemical and Materials Engineering, Faculty of Engineering, King Abdulaziz University, P.O. Box 80204, Jeddah, 21589, Saudi Arabia

²Department of Mathematics, Cankaya University, Ogretmenler Cad. 14, Balgat, Ankara, 06530, Turkey

Full list of author information is available at the end of the article

Abstract

By using fixed point results on cones, we study the existence of solutions for the singular nonlinear fractional boundary value problem

$${}^c D^\alpha u(t) = f(t, u(t), u'(t), {}^c D^\beta u(t)),$$
$$u(0) = au(1), \quad u'(0) = b {}^c D^\beta u(1), \quad u''(0) = u'''(0) = u^{(n-1)}(0) = 0,$$

where $n \geq 3$ is an integer, $\alpha \in (n-1, n)$, $0 < \beta < 1$, $0 < a < 1$, $0 < b < \Gamma(2-\beta)$, f is an L^q -Caratheodory function, $q > \frac{1}{\alpha-1}$ and $f(t, x, y, z)$ may be singular at value 0 in one dimension of its space variables x, y, z . Here, ${}^c D$ stands for the Caputo fractional derivative.

Keywords: boundary value problem; fixed point; fractional differential equation; Green function; regularization; singular

1 Introduction

Fractional differential equations (see, for example, [1–6] and references therein) started to play an important role in several branches of science and engineering. There are some works about existence of solutions for the nonlinear mixed problems of singular fractional boundary value problem (see, for example, [7–11] and [12]). Also, there are different methods for solving distinct fractional differential equations (see, for example, [13–18] and [19]). By using fixed point results on cones, we focus on the existence of positive solutions for a nonlinear mixed problem of singular fractional boundary value problem. For the convenience of the reader, we present some necessary definitions from fractional calculus theory (see, for example, [20]). The Caputo derivative of fractional order α for a function $f : [0, \infty) \rightarrow \mathbb{R}$ is defined by

$${}^c D^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} f^{(n)}(s) ds \quad (n-1 < \alpha < n, n = [\alpha] + 1).$$

Let $q \geq 1$. As you know, $L^q[0, 1]$ denotes the space of functions, whose q th powers of modulus are integrable on $[0, 1]$, equipped with the norm $\|x\|_q = (\int_0^1 |x(t)|^q dt)^{\frac{1}{q}}$. We consider the sup norm

$$\|x\| = \sup\{|x(t)| : t \in [0, 1]\}$$

on the space $C[0, 1]$. Also, $AC[0, 1]$ is the set of absolutely continuous functions on $[0, 1]$. Let B be a subset of \mathbb{R}^3 . A function $f : [0, 1] \times B \rightarrow \mathbb{R}$ is called an L^q -Caratheodory function whenever the real-valued function $f(\cdot, x, y, z)$ on $[0, 1]$ is measurable for all $(x, y, z) \in B$, the function $f(t, \cdot, \cdot, \cdot) : B \rightarrow \mathbb{R}$ is continuous for almost all $t \in (0, 1]$, and for each compact set $U \subset B$, there exists a function $\varphi_U \in L^q[0, 1]$ such that $|f(t, x, y, z)| \leq \varphi_U(t)$ for almost all $t \in [0, 1]$ and $(x, y, z) \in U$. Consider the nonlinear fractional boundary value problem

$$\begin{aligned} {}^c D^\alpha u(t) &= f(t, u(t), u'(t), {}^c D^\beta u(t)), \\ u(0) &= au(1), \quad u'(0) = b {}^c D^\beta u(1), \quad u''(0) = u'''(0) = u^{(n-1)}(0) = 0, \end{aligned} \tag{*}$$

where $n \geq 3$ is an integer, $\alpha \in (n-1, n)$, $0 < \beta < 1$, $0 < a < 1$, $0 < b < \Gamma(2-\beta)$ and $q > \frac{1}{\alpha-1}$. We say that the function $u : [0, 1] \rightarrow \mathbb{R}$ is a positive solution for the problem whenever $u > 0$ on $[0, 1]$, ${}^c D^\alpha u$ is a function in $L^q[0, 1]$, and u satisfies the boundary conditions almost everywhere on $[0, 1]$. In this paper, we suppose that f is an L^q -Caratheodory function on $[0, 1] \times B$, where $B = (0, \infty) \times (0, \infty) \times (0, \infty)$, there exists a positive constant m such that $m \leq f(t, x, y, z)$ for almost all $t \in [0, 1]$ and $(x, y, z) \in B$, f satisfies the estimate

$$f(t, x, y, z) \leq h(x) + r(|y|) + k(|z|) + \gamma(t)w(x, |y|, |z|),$$

where $h, r, k \in C(0, \infty)$ are positive and non-increasing, $\gamma \in L^q[0, 1]$ and $w \in C([0, \infty) \times [0, \infty) \times [0, \infty))$ are positive, w is non-decreasing in all its variables, $\int_0^1 h^q(s^\alpha) ds < \infty$, $\int_0^1 r^q(s^{\alpha-1}) ds < \infty$, $\int_0^1 k^q(s^{\alpha-\beta}) ds < \infty$, and $\lim_{x \rightarrow \infty} \frac{w(x, x, x)}{x} = 0$. Since we suppose that problem (*) is singular, that is, $f(t, x, y, z)$ may be singular at the value 0 of its space variables x, y, z , we use regularization and sequential techniques for the existence of positive solutions of the problem. In this way, for each natural number n define the function f_n by

$$f_n(t, x, y, z) = f(t, \chi_n^+(x), \chi_n^+(y), \chi_n^+(z))$$

for all $t \in [0, 1]$ and $(x, y, z) \in \mathbb{R}^3$, where

$$\chi_n^+(u) = \begin{cases} u, & u \geq \frac{1}{n}, \\ \frac{1}{n}, & u < \frac{1}{n}. \end{cases}$$

It is easy to see that each f_n is an L^q -Caratheodory function on $[0, 1] \times \mathbb{R}^3$, $m \leq f_n(t, x, y, z)$,

$$f_n(t, x, y, z) \leq h\left(\frac{1}{n}\right) + r\left(\frac{1}{n}\right) + k\left(\frac{1}{n}\right) + \gamma(t)w(1+x, 1+|y|, 1+|z|)$$

and

$$f_n(t, x, y, z) \leq h(x) + r(|y|) + k(|z|) + \gamma(t)w(1+x, 1+|y|, 1+|z|)$$

for almost all $t \in [0, 1]$ and all $(x, y, z) \in B$. In 2012, Agarwal *et al.* proved the following result.

Lemma 1.1 [7] *Let $\rho \in L^q[0, 1]$ and $0 \leq t_1 < t_2 \leq 1$. Then we have $|\int_0^t (t-s)^{\alpha-2} \rho(s) ds| \leq (\frac{t^d}{d})^{1/p} \|\rho\|_q$ for all $t \in [0, 1]$ and*

$$\left| \int_0^{t_2} (t_2-s)^{\alpha-2} \rho(s) ds - \int_0^{t_1} (t_1-s)^{\alpha-2} \rho(s) ds \right| \leq \left(\frac{t_1^d + (t_2-t_1)^d - t_2^d}{d} \right)^{1/p} \|\rho\|_q + \left(\frac{(t_2-t_1)^d}{d} \right)^{1/p} \|\rho\|_q,$$

where $d = (\alpha - 2)p + 1$.

2 Main results

Now, we are ready to investigate the problem in regular and singular cases. First, we give the following result.

Lemma 2.1 *Let $y \in C[0, 1]$. Then the boundary value problem*

$$\begin{aligned} {}^c D^\alpha u(t) &= y(t) \quad (t \in (0, 1)), \\ u(0) &= au(1), \quad u'(0) = b {}^c D^\beta u(1), \quad u''(0) = u'''(0) = u^{(n-1)}(0) = 0 \end{aligned}$$

is equivalent to the fractional integral equation $u(t) = \int_0^1 G(t,s)y(s) ds$, where

$$\begin{aligned} G(t,s) &= \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \\ &+ \frac{a\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)(1-s)^{\alpha-1} + b\Gamma(\alpha)\Gamma(2-\beta)(a+t-at)(1-s)^{\alpha-\beta-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \end{aligned}$$

whenever $0 \leq s \leq t \leq 1$ and

$$G(t,s) = \frac{a\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)(1-s)^{\alpha-1} + b\Gamma(\alpha)\Gamma(2-\beta)(a+t-at)(1-s)^{\alpha-\beta-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)}$$

whenever $0 \leq t \leq s \leq 1$.

Proof From ${}^c D^\alpha u(t) = y(t)$ and the boundary conditions, we obtain

$$\begin{aligned} u(t) &= I^\alpha y(t) + u(0) + u'(0)t + \frac{u''(0)}{2!}t^2 + \dots + \frac{u^{(n-1)}(0)}{(n-1)!}t^{n-1} \\ &= \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + u(0) + u'(0)t. \end{aligned}$$

By properties of the Caputo derivative, we get

$$\begin{aligned} {}^c D^\beta u(t) &= I^{\alpha-\beta} y(t) + {}^c D^\beta (u(0) + u'(0)t) \\ &= \frac{1}{\Gamma(\alpha-\beta)} \int_0^t (t-s)^{\alpha-\beta-1} y(s) ds + \frac{u'(0)t^{1-\beta}}{\Gamma(2-\beta)}. \end{aligned}$$

Thus, $u(1) = \frac{1}{\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-1} y(s) ds + u(0) + u'(0)$ and

$${}^c D^\beta u(1) = \frac{1}{\Gamma(\alpha - \beta)} \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds + \frac{u'(0)}{\Gamma(2 - \beta)}.$$

By using the boundary conditions $u(0) = au(1)$ and $u'(0) = b {}^c D^\beta u(1)$, we get $u(0) = a(\frac{1}{\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-1} y(s) ds + u(0) + u'(0))$ and

$$u'(0) = b \left(\frac{1}{\Gamma(\alpha - \beta)} \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds + \frac{u'(0)}{\Gamma(2 - \beta)} \right).$$

Hence, $u'(0) = \frac{b\Gamma(2-\beta)}{\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds$ and

$$u(0) = \frac{a}{(1-a)\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-1} y(s) ds + \frac{ab\Gamma(2-\beta)}{(1-a)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds.$$

Thus,

$$\begin{aligned} u(t) &= \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + u(0) + u'(0)t \\ &= \int_0^t \left(\frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} + \frac{a\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)(1-s)^{\alpha-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \right. \\ &\quad \left. + \frac{b\Gamma(\alpha)\Gamma(2-\beta)(a+t-at)(1-s)^{\alpha-\beta-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \right) y(s) ds \\ &\quad + \int_t^1 \left(\frac{a\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)(1-s)^{\alpha-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \right. \\ &\quad \left. + \frac{b\Gamma(\alpha)\Gamma(2-\beta)(a+t-at)(1-s)^{\alpha-\beta-1}}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)} \right) y(s) ds \\ &= \int_0^1 G(t,s)y(s) ds. \end{aligned}$$

This completes the proof. □

Put $k_1 = \frac{\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)+b\Gamma(\alpha)\Gamma(2-\beta)}{(1-a)\Gamma(\alpha)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)}$ and $k_2 = \frac{ab\Gamma(2-\beta)}{(1-a)\Gamma(\alpha-\beta)(\Gamma(2-\beta)-b)}$. It is easy to check that the Green function G in the last result belongs to $C([0,1] \times [0,1])$, $G(t,s) > 0$ for all $(t,s) \in [0,1] \times [0,1]$,

$$G(t,s) \leq k_1(1-s)^{\alpha-\beta-1} \leq 1 \quad \text{and} \quad G(t,s) \geq k_2(1-s)^{\alpha-\beta-1}$$

for all $(t,s) \in [0,1] \times [0,1]$. Consider the Banach space $X = C^1[0,1]$ with the norm $\|x\|_* = \max\{\|x\|, \|x'\|\}$ and the cone

$$P = \{x \in X : x(t) \geq 0 \text{ and } x'(t) \geq 0 \text{ for all } t \in [0,1]\}.$$

For each natural number n , define the operator Q_n on P by

$$(Q_n x)(t) = \int_0^1 G(t, s) f_n(s, u(s), u'(s), {}^c D^\beta u(s)) ds.$$

Now, we prove that Q_n is a completely continuous operator (see [2]).

Lemma 2.2 *The operator Q_n is a completely continuous operator.*

Proof Let $x \in P$. Then, ${}^c D^\beta x \in C[0, 1]$ and ${}^c D^\beta x \geq 0$. Now, define $\rho(t) = f_n(t, u(t), u'(t), {}^c D^\beta u(t))$ for almost all $t \in [0, 1]$. Then $\rho \in L^q[0, 1]$ and $\rho(t) \geq m$ for almost all $t \in [0, 1]$. By using the properties of fractional integral I^α , it is easy to see that $Q_n x \in C[0, 1]$, $(Q_n x)(t) \geq 0$ and

$$(Q_n x)'(t) = \frac{1}{\Gamma(\alpha - 1)} \int_0^t (t - s)^{\alpha - 2} \rho(s) ds$$

for all $t \in [0, 1]$. This implies that $(Q_n x)' \in C[0, 1]$ and $(Q_n x)' \geq 0$ on $[0, 1]$. Consequently, Q_n maps P into P . In order to prove that Q_n is a continuous operator, let x_m be a convergent sequence in P and $\lim_{m \rightarrow \infty} x_m = x$. Thus, $\lim_{m \rightarrow \infty} x_m^{(j)}(t) = x^{(j)}(t)$ uniformly on $[0, 1]$ for $j = 0, 1$. Since

$$\begin{aligned} {}^c D^\beta x(t) &= \frac{1}{\Gamma(1 - \beta)} \frac{d}{dt} \int_0^t (t - s)^{-\beta} (x(s) - x(0)) ds \\ &= \frac{1}{\Gamma(1 - \beta)} \int_0^t (t - s)^{-\beta} x'(s) ds, \end{aligned}$$

we get $|{}^c D^\beta x_m(t) - {}^c D^\beta x(t)| \leq \frac{\|x'_m - x'\|}{\Gamma(1 - \beta)} \int_0^t (t - s)^{-\beta} ds \leq \frac{\|x'_m - x'\|_*}{\Gamma(\beta)}$ and $\lim_{m \rightarrow \infty} {}^c D^\beta x_m(t) = {}^c D^\beta x(t)$ uniformly on $[0, 1]$. Also, we have $|{}^c D^\beta x_m(t)| \leq \frac{x'_m}{\Gamma(\beta)}$ on $[0, 1]$, and so $\|{}^c D^\beta x_m\| \leq \frac{\|x'_m\|}{\Gamma(\beta)}$. Now, put

$$\rho_m(t) = f_n(t, x_m(t), x'_m(t), {}^c D^\beta x_m(t)) \quad \text{and} \quad \rho(t) = f_n(t, x(t), x'(t), {}^c D^\beta x(t)).$$

Then, it is easy to see that $\lim_{m \rightarrow \infty} \rho_m(t) = \rho(t)$ for almost all $t \in [0, 1]$, and there exists $\beta \in L^q[0, 1]$ such that $0 \leq \rho_m(t) \leq \beta(t)$ for almost all $t \in [0, 1]$ and all $m \geq 1$. Since f_n is an L^q -Caratheodory function, $\{x_m\}$ is bounded in $C^1[0, 1]$, and $\{{}^c D^\beta x_m\}$ is bounded in $C[0, 1]$. Therefore, $\lim_{m \rightarrow \infty} (Q_n x_m)(t) = (Q_n x)(t)$ uniformly on $[0, 1]$. Since $\{\rho_m\}$ is L^q -convergent on $[0, 1]$,

$$\lim_{m \rightarrow \infty} (Q_n x_m)'(t) = \frac{1}{\Gamma(\alpha - 1)} \lim_{m \rightarrow \infty} \int_0^t (t - s)^{\alpha - 2} \rho_m(s) ds = (Q_n x)'(t)$$

uniformly on $[0, 1]$. Hence, Q_n is a continuous operator. Now, we have to show that for each bounded sequence $\{x_m\}$ in P , the sequence $\{Q_n x_m\}$ is relatively compact in $C[0, 1]$. Choose a positive constant k such that $\|x_m\| \leq k$ and $\|x'_m\| \leq k$ for all m . Note that $\|{}^c D^\beta x_m\| \leq \frac{k}{\Gamma(\beta)}$ and $|\int_0^t (t - s)^{\alpha - 2} \rho_m(s) ds| \leq (\int_0^t (t - s)^{(\alpha - 2)p} ds)^{\frac{1}{p}} (\int_0^t |\rho_m(s)|^q ds)^{\frac{1}{q}} \leq (\frac{t^d}{d})^{\frac{1}{p}} \|\rho_m\|_q$ for all m , where $d = (\alpha - 2)p + 1$. But we have

$$0 \leq (Q_n x_m)(t) = \int_0^1 G(t, s) \rho_m(s) ds \leq \int_0^1 G(t, s) \beta(s) ds \leq \frac{\|\beta\|_1}{\Gamma(\alpha)}$$

and

$$\begin{aligned} 0 \leq (Q_n x_m)'(t) &= \frac{1}{\Gamma(\alpha - 1)} \int_0^t (t - s)^{\alpha - 2} \rho_m(s) \, ds \\ &\leq \frac{1}{\Gamma(\alpha - 1)} \int_0^t (t - s)^{\alpha - 2} \beta(s) \, ds \leq \frac{1}{\Gamma(\alpha - 1)} \left(\frac{1}{(\alpha - 2)p + 1} \right)^{\frac{1}{p}} \|\beta\|_q \end{aligned}$$

for all $t \in [0, 1]$ and m . This implies that $\{Q_n x_m\}$ is bounded in $C^1[0, 1]$. Also, we have

$$\begin{aligned} &|(Q_n x_m)'(t_2) - (Q_n x_m)'(t_1)| \\ &= \frac{1}{\Gamma(\alpha - 1)} \left| \int_0^{t_2} (t_2 - s)^{\alpha - 2} \rho_m(s) \, ds - \int_0^{t_1} (t_1 - s)^{\alpha - 2} \rho_m(s) \, ds \right| \\ &\leq \frac{\|\rho_m\|_q}{\Gamma(\alpha - 1)} \left(\left(\frac{t_1^d + (t_2 - t_1)^d - t_2^d}{d} \right)^{\frac{1}{p}} + \left(\frac{(t_2 - t_1)^d}{d} \right)^{\frac{1}{p}} \right) \\ &\leq \frac{\|\beta\|_q}{\Gamma(\alpha - 1)} \left(\left(\frac{t_1^d + (t_2 - t_1)^d - t_2^d}{d} \right)^{\frac{1}{p}} + \left(\frac{(t_2 - t_1)^d}{d} \right)^{\frac{1}{p}} \right) \end{aligned}$$

for all $0 \leq t_1 \leq t_2 \leq 1$, where $d = (\alpha - 2)p + 1$. Hence, $\{(Q_n x_m)'\}$ is equicontinuous on $[0, 1]$. Thus, $\{Q_n x_m\}$ is relatively compact in $C^1[0, 1]$ by the Arzela-Ascoli theorem. Hence, Q_n is a completely continuous operator. \square

We need the following result (see [2] and [21]).

Lemma 2.3 [21] *Let Y be a Banach space, P a cone in Y and Ω_1 and Ω_2 bounded open balls in Y centered at the origin with $\bar{\Omega}_1 \subset \Omega_2$. Suppose that $T : P \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow P$ is a completely continuous operator such that $\|Tx\| \geq \|x\|$ for all $x \in P \cap \partial\Omega_1$ and $\|Tx\| \leq \|x\|$ for all $x \in P \cap \partial\Omega_2$. Then T has a fixed point in $P \cap (\bar{\Omega}_2 \setminus \Omega_1)$.*

Theorem 2.4 *For each natural number n , problem (*) has a solution $u_n \in P$ such that $u_n \geq \frac{mk_2}{\alpha - \beta}$, $u_n'(t) \geq \frac{mt^{\alpha - 1}}{\Gamma(\alpha)}$ and ${}^c D^\beta u_n(t) \geq \frac{mt^{\alpha - \beta}}{\Gamma(\alpha - \beta + 1)}$ for all $t \in [0, 1]$.*

Proof Let $n \geq 1$. It is sufficient to show that Q_n has a fixed point u_n in P with the desired conditions. In this way, note that

$$\begin{aligned} (Q_n x)(t) &= \int_0^1 G(t, s) f_n(s, x(s), x'(s), {}^c D^\beta x(s)) \, ds \\ &\geq m \int_0^1 G(t, s) \, ds \geq m \int_0^1 k_2 (1 - s)^{\alpha - \beta - 1} \, ds = \frac{mk_2}{\alpha - \beta}, \end{aligned}$$

and so $\|Q_n x\|_* \geq \|Q_n(x)\| \geq \frac{mk_2}{\alpha - \beta}$. Put $\Omega_1 = \{x \in X : \|x\|_* < \frac{mk_2}{\alpha - \beta}\}$. Then $\|Q_n x\|_* \geq \|x\|_*$ for all $x \in P \cap \partial\Omega_1$. If $v_n = h(\frac{1}{n}) + r(\frac{1}{n}) + k(\frac{1}{n})$, then

$$\begin{aligned} |(Q_n x)(t)| &\leq \left| \int_0^1 G(t, s) f_n(s, u(s), u'(s), {}^c D^\beta u(s)) \, ds \right| \\ &\leq \int_0^1 |G(t, s)| (v_n + \gamma(s)w(1 + |x(s)|, 1 + |x'(s)|, 1 + |{}^c D^\beta x(s)|)) \, ds \\ &\leq k_1 (v_n + w(1 + \|x\|, 1 + \|x'\|, 1 + \|{}^c D^\beta x\|)) \|\gamma\|_1 \end{aligned}$$

and

$$\begin{aligned} & |(Q_n x)'(t)| \\ &= \left| \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} f_n(s, x(s), x'(s), {}^c D^\beta x(s)) ds \right| \\ &\leq \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} (v_n + \gamma(s)w(1 + |x(s)|, 1 + |x'(s)|, 1 + |{}^c D^\beta x(s)|)) ds \\ &\leq \frac{1}{\Gamma(\alpha-1)} \left(\frac{v_n t^{\alpha-1}}{\alpha-1} + w(1 + \|x\|, 1 + \|x'\|, 1 + \|{}^c D^\beta x\|) \int_0^t (t-s)^{\alpha-2} \gamma(s) ds \right) \end{aligned}$$

for all $x \in P$ and $t \in [0, 1]$, because w is non-decreasing in all its variables. Since $\|x\| \leq \|x\|_*$, $\|x'\| \leq \|x\|_*$, $\|{}^c D^\beta x\| \leq \frac{\|x'\|}{\Gamma(\beta)} \leq \frac{\|x\|_*}{\Gamma(\beta)}$ and

$$\int_0^t (t-s)^{\alpha-2} \gamma(s) ds \leq \left(\frac{1}{d}\right)^{1/p} \|\gamma\|_q,$$

where $d = (\alpha - 2)p + 1$, we have

$$\|Q_n(x)\| \leq k_1 \left(v_n + w \left(1 + \|x\|_*, 1 + \|x\|_*, 1 + \frac{\|x\|_*}{\Gamma(\beta)} \right) \|\gamma\|_1 \right)$$

and

$$\begin{aligned} & \|(Q_n x)'\| \\ &\leq \frac{1}{\Gamma(\alpha-1)} \left(\frac{v_n}{\alpha-1} + w \left(1 + \|x\|_*, 1 + \|x\|_*, 1 + \frac{\|x\|_*}{\Gamma(\beta)} \right) (1/d)^{1/p} \|\gamma\|_q \right). \end{aligned}$$

Hence, $\|Q_n x\|_* \leq M \left(\frac{v_n}{\alpha-1} + Nw(1 + \|x\|_*, 1 + \|x\|_*, 1 + \frac{\|x\|_*}{\Gamma(\beta)}) \right)$, where $N = \max\{\|\gamma\|_1, (1/d)^{1/p} \times \|\gamma\|_q\}$ and $M = \max\{\frac{1}{\Gamma(\alpha-1)}, k_1\}$. Since

$$\lim_{v \rightarrow \infty} \frac{1}{v} w(1 + v, 1 + v, 1 + v) = 0,$$

there exists a positive constant L such that

$$M \left(\frac{v_n}{\alpha-1} + Nw \left(1 + v, 1 + v, \frac{v}{\Gamma(\beta)} \right) \right) < v$$

for all $v \geq L$. Thus, $\|Q_n x\|_* < \|x\|_*$ for all $x \in P$ with $\|x\|_* \geq L$. Put $\Omega_2 = \{x \in X : \|x\|_* < L\}$. Then $\|Q_n x\|_* < \|x\|_*$ for all $x \in P \cap \partial\Omega_2$. By using last result, Q_n has a fixed point u_n in $P \cap (\bar{\Omega}_2 \setminus \Omega_1)$. But $u_n = (Q_n u_n)(t) \geq \frac{mt^{\alpha-1}}{\alpha-\beta}$ and

$$\begin{aligned} (Q_n x)'(t) &= \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} f_n(s, x(s), x'(s), {}^c D^\beta x(s)) ds \\ &\geq \frac{m}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} ds = \frac{mt^{\alpha-1}}{\Gamma(\alpha)} \end{aligned}$$

for all $t \in [0, 1]$ and $x \in P$. Since $\int_0^t (t-s)^{-\beta} s^{\alpha-1} ds = \frac{\Gamma(\alpha)\Gamma(1-\beta)}{\Gamma(\alpha-\beta+1)} t^{\alpha-\beta}$,

$$\begin{aligned} {}^c D^\beta u_n(t) &= \frac{1}{\Gamma(1-\beta)} \int_0^t (t-s)^{-\beta} u'_n(s) ds \\ &\geq \frac{m}{\Gamma(\alpha)\Gamma(1-\beta)} \int_0^t (t-s)^{-\beta} s^{\alpha-1} ds = \frac{mt^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)} \end{aligned}$$

for all $t \in [0, 1]$. This completes the proof. \square

Now, we give our last result.

Theorem 2.5 *Problem (*) has a solution u such that $u(t) \geq \frac{mk_2}{\alpha-\beta}$, $u'(t) \geq \frac{mt^{\alpha-1}}{\Gamma(\alpha)}$ and ${}^c D^\beta u(t) \geq \frac{mt^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}$ for all $t \in [0, 1]$.*

Proof By using Theorem 2.4, one gets that for each natural number n , problem (*) has a solution $u_n \in P$ with the desired conditions. Thus, $h(u_n(t)) \leq h(\frac{mk_2}{\alpha-\beta})$, $r(|u'_n(t)|) \leq r(\frac{mt^{\alpha-1}}{\Gamma(\alpha)})$ and $k(|{}^c D^\beta u_n(t)|) \leq k(\frac{mt^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)})$ for all $t \in [0, 1]$ and n . Also, we have $\|{}^c D^\beta u_n\| \leq \frac{\|u'_n\|}{\Gamma(\beta)}$. Suppose that

$$S(t) = h\left(\frac{mk_2}{\alpha-\beta}\right) + r\left(\frac{mt^{\alpha-1}}{\Gamma(\alpha)}\right) + k\left(\frac{mt^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right).$$

Then

$$\begin{aligned} m &\leq f_n(t, u_n(t), u'_n(t), {}^c D^\beta u_n(t)) \\ &\leq S(t) + \gamma(t)w(1 + \|u_n\|, 1 + \|u'_n\|, 1 + \|{}^c D^\beta u_n\|) \\ &\leq S(t) + \gamma(t)w\left(1 + \|u_n\|_*, 1 + \|u'_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)}\right) \end{aligned}$$

for almost all $t \in [0, 1]$ and n . Since $0 \leq G(t, s) \leq k_1$, we get

$$\begin{aligned} 0 &\leq u_n(t) \\ &= \int_0^1 G(t, s) f_n(s, u_n(s), u'_n(s), {}^c D^\beta u_n(s)) ds \\ &\leq k_1 \left(\int_0^1 S(s) ds + w\left(1 + \|u_n\|_*, 1 + \|u'_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)}\right) \|\gamma\|_1 \right) \end{aligned}$$

and

$$\begin{aligned} 0 &\leq u'_n(t) \\ &\leq \frac{1}{\Gamma(\alpha-1)} \left(\int_0^t (t-s)^{\alpha-2} S(s) ds \right. \\ &\quad \left. + w\left(1 + \|u_n\|_*, 1 + \|u'_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)}\right) \int_0^t (t-s)^{\alpha-2} \gamma(s) ds \right). \end{aligned}$$

We show that $\int_0^t (t-s)^{\alpha-2} S(s) ds$ is bounded on $[0, 1]$. Let $d = (\alpha - 2)p + 1$. Note that

$$\begin{aligned} & \int_0^1 (t-s)^{\alpha-2} h\left(\frac{mk_2}{\alpha-\beta}\right) ds \\ &= h\left(\frac{mk_2}{\alpha-\beta}\right) \int_0^1 (t-s)^{\alpha-2} ds = \frac{1}{\alpha-1} h\left(\frac{mk_2}{\alpha-\beta}\right) =: \eta_1 < \infty, \\ & \int_0^t (t-s)^{\alpha-2} r\left(\frac{ms^{\alpha-1}}{\Gamma(\alpha)}\right) ds \\ &= \left(\frac{1}{d}\right)^{1/p} \left(\frac{\Gamma(\alpha)}{m}\right)^{\frac{1}{(\alpha-1)q}} \left(\int_0^{\left(\frac{m}{\Gamma(\alpha)}\right)^{\frac{1}{\alpha-1}}} r^q(s^{\alpha-1}) ds\right)^{1/q} =: \eta_2 < \infty \end{aligned}$$

and

$$\begin{aligned} & \int_0^t (t-s)^{\alpha-2} k\left(\frac{ms^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right) ds \\ &= \left(\frac{1}{d}\right)^{1/p} \left(\frac{\Gamma(\alpha-\beta+1)}{m}\right)^{\frac{1}{(\alpha-\beta)q}} \left(\int_0^{\left(\frac{m}{\Gamma(\alpha-\beta+1)}\right)^{\frac{1}{\alpha-\beta}}} k^q(s^{\alpha-\beta}) ds\right)^{1/q} =: \eta_3 < \infty. \end{aligned}$$

Thus, $\int_0^t (t-s)^{\alpha-2} S(s) ds \leq \eta$ for all $t \in [0, 1]$, where $\eta = \eta_1 + \eta_2 + \eta_3$. Also, we have

$$\begin{aligned} & \int_0^1 S(s) ds \\ &\leq \frac{1}{\alpha-1} h\left(\frac{mk_2}{\alpha-\beta}\right) + \left(\frac{\Gamma(\alpha)}{m}\right)^{\frac{1}{\alpha-1}} \int_0^{\left(\frac{m}{\Gamma(\alpha)}\right)^{\frac{1}{\alpha-1}}} r(s^{\alpha-1}) ds \\ &+ \left(\frac{\Gamma(\alpha-\beta+1)}{m}\right)^{\frac{1}{\alpha-\beta}} \int_0^{\left(\frac{m}{\Gamma(\alpha-\beta+1)}\right)^{\frac{1}{\alpha-\beta}}} k(s^{\alpha-\beta}) ds < \infty. \end{aligned}$$

Since

$$\|u_n\| = k_1 \left(\int_0^1 S(s) ds + w \left(1 + \|u_n\|_*, 1 + \|u_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)} \right) \|\gamma\|_1 \right)$$

and

$$\|u'_n\| \leq \frac{1}{\Gamma(\alpha-1)} \left(\eta + w \left(1 + \|u_n\|_*, 1 + \|u_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)} \right) \left(\frac{1}{d}\right)^{1/p} \|\gamma\|_q \right),$$

we get $\|u_n\|_* \leq M(\Phi + Kw(1 + \|u_n\|_*, 1 + \|u_n\|_*, 1 + \frac{\|u_n\|_*}{\Gamma(\beta)}))$ for all n , where $\Phi = \max\{\eta, \int_0^1 S(s) ds\}$, $K = \max\{\|\gamma\|_1, (\frac{1}{d})^{1/p} \|\gamma\|_q\}$ and also $M = \max\{k_1, \frac{1}{\Gamma(\alpha-1)}\}$. On the other hand, there exists a positive constant L such that $M(\Phi + Kw(1 + \nu, 1 + \nu, 1 + \frac{\nu}{\Gamma(\beta)})) < \nu$ for all $\nu \geq L$, and so $\|u_n\|_* < L$ for all n . Thus, for almost all $t \in [0, 1]$ and all n , we have $f_n(t, u_n(t), u'_n(t), {}^c D^\beta u_n(t)) \leq R(t)$, where

$$R(t) = S(t) + \gamma(t)w \left(1 + L, 1 + L, 1 + \frac{L}{\Gamma(\beta)} \right).$$

Note that $R \in L^q[0,1]$. We show that $\{u'_n\}$ is equicontinuous on $[0,1]$. Let $\rho_n(t) = f_n(t, u_n(t), u'_n(t), {}^cD^\beta u_n(t))$ and $0 \leq t_1 < t_2 \leq T$. Then

$$\begin{aligned} & |u'_n(t_2) - u'_n(t_1)| \\ &= \frac{1}{\Gamma(\alpha-1)} \left| \int_0^{t_2} (t_2-s)^{\alpha-2} \rho_n(s) ds - \int_0^{t_1} (t_1-s)^{\alpha-2} \rho_n(s) ds \right| \\ &\leq \frac{1}{\Gamma(\alpha-1)} \left(\int_0^{t_1} ((t_1-s)^{\alpha-2} - (t_2-s)^{\alpha-2}) \rho_n(s) ds + \int_{t_1}^{t_2} (t_2-s)^{\alpha-2} \rho_n(s) ds \right) \\ &\leq \frac{1}{\Gamma(\alpha-1)} \left(\int_0^{t_1} ((t_1-s)^{\alpha-2} - (t_2-s)^{\alpha-2}) R(s) ds + \int_{t_1}^{t_2} (t_2-s)^{\alpha-2} R(s) ds \right), \end{aligned}$$

and so

$$\begin{aligned} & |u'_n(t_2) - u'_n(t_1)| \\ &\leq \frac{\|R\|_q}{\Gamma(\alpha-1)} \left(\left(\frac{t_1^d + (t_2-t_1)^d - t_2^d}{d} \right)^{1/p} + \left(\frac{(t_2-t_1)^d}{d} \right)^{1/p} \right). \end{aligned}$$

Hence, $\{u'_n\}$ is equicontinuous on $[0,1]$. Since $\{u_n\}$ is a bounded sequence in $C[0,1]$, by using the Arzela-Ascoli theorem, without loss of generality, we can assume that $\{u_n\}$ is convergent in $C[0,1]$. Let $\lim_{n \rightarrow \infty} u_n = u$. Then, it is easy to see that ${}^cD^\beta u_n(t) = \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} u'_n(s) ds$, and ${}^cD^\beta u_n(t)$ uniformly converges to $\frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} u'(s) ds$ on $[0,1]$. Thus, ${}^cD^\beta u_n$ converges to ${}^cD^\beta u$ in $C[0,1]$. Hence,

$$\lim_{n \rightarrow \infty} f_n(t, u_n(t), u'_n(t), {}^cD^\beta u_n(t)) = f(t, u(t), u'(t), {}^cD^\beta u(t))$$

for almost all $t \in [0,1]$. Since $R \in L^q[0,1]$, by using the dominated convergence theorem on the relation

$$u_n(t) = \int_0^1 G(t,s) f_n(s, u_n(s), u'_n(s), {}^cD^\beta u_n(s)) ds,$$

we get $u(t) = \int_0^1 G(t,s) f(s, u(s), u'(s), {}^cD^\beta u(s))$ for all $t \in [0,1]$. This completes the proof. \square

2.1 Examples for the problem

Example 2.1 Let $\rho_1, \rho_2 \in L^q[0,1]$, $\rho_1(t) \geq m > 0$ for almost all t in $[0,1]$. Suppose that

$$f(t, x, y, z) = \rho_1(t) + \frac{1}{x^{\frac{1}{3}-\lambda}} + \frac{1}{y^{\frac{1}{4}}} + \frac{1}{z^{\frac{1}{4}}} + |\rho_2(t)| (x^{\frac{1}{3}} + y^{\frac{1}{4}} + z^{\frac{1}{4}})$$

on $[0,1] \times B$, $\lambda = (au(1))^{\frac{1}{3}}$, $h(x) = \frac{1}{x^{\frac{1}{3}-\lambda}}$ whenever $x^{\frac{1}{3}} - \lambda \geq 0$ and $h(x) = 0$ whenever $x^{\frac{1}{3}} - \lambda < 0$, $r(x) = \frac{1}{x^{\frac{1}{4}}}$, $k(x) = \frac{1}{x^{\frac{1}{4}}}$, $w(x, y, z) = (x^{\frac{1}{3}} + y^{\frac{1}{4}} + z^{\frac{1}{4}} + 1)$ and $\gamma(t) = \rho_1(t) + |\rho_2(t)|$. Then Theorem 2.5 guarantees that problem (*) has a positive solution.

Example 2.2 Consider the nonlinear mixed problem of singular fractional boundary value problem

$${}^c D^{\frac{7}{3}} u(t) = t + 1 + \frac{1}{u(t)^{\frac{1}{3}} - \rho} + \frac{1}{u'(t)^{\frac{1}{4}}} + \frac{1}{({}^c D^{\frac{1}{4}} u(t))^{\frac{1}{4}}} + 2(u(t)^{\frac{1}{3}} + u'(t)^{\frac{1}{4}} + ({}^c D^{\frac{1}{4}} u(t))^{\frac{1}{4}} + 1)$$

via boundary value conditions $u(0) = \frac{1}{4}u(1)$, $u'(0) = \frac{1}{3}({}^c D^{\frac{1}{4}})u(1)$ and $u''(0) = u'''(0) = \dots = u^{(n-1)}(0) = 0$, where $\rho = ((\frac{1}{4})u(1))^{\frac{1}{3}}$. Let

$$f(t, u(t), u'(t), {}^c D^{\frac{1}{4}}) = t + 1 + \frac{1}{u(t)^{\frac{1}{3}} - \rho} + \frac{1}{u'(t)^{\frac{1}{4}}} + \frac{1}{({}^c D^{\frac{1}{4}} u(t))^{\frac{1}{4}}} + 2(u(t)^{\frac{1}{3}} + u'(t)^{\frac{1}{4}} + ({}^c D^{\frac{1}{4}} u(t))^{\frac{1}{4}} + 1).$$

Then the map f is singular at $t = 0$, and f satisfies the desired conditions, where $h(x) = \frac{1}{x^{\frac{1}{3}} - \rho}$ whenever $x^{\frac{1}{3}} - \rho \geq 0$ and $h(x) = 0$ whenever $x^{\frac{1}{3}} - \rho < 0$, $r(x) = \frac{1}{x^{\frac{1}{4}}}$, $k(x) = \frac{1}{x^{\frac{1}{4}}}$, $w(x, y, z) = x^{\frac{1}{3}} + y^{\frac{1}{4}} + z^{\frac{1}{4}} + 1$, $\rho_1(t) = t + 1 > 1 = m$, $\rho_2(t) = 2$ and $\gamma(t) = \rho_1(t) + |\rho_2(t)|$. Then Theorem 2.5 guarantees that this problem has a positive solution.

3 Conclusions

One of the most interesting branches is obtaining solutions of singular fractional differential via boundary value problems. Having these things in mind, we study the existence of solutions for a singular nonlinear fractional boundary value problem. Two illustrative examples illustrate the applicability of the proposed method. It seems that the obtained results could be extended to more general functional spaces. Finally, note that all calculations in proofs of the results depend on the definition of the fractional derivative.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors have equal contributions. All authors read and approved the final manuscript.

Author details

¹Department of Chemical and Materials Engineering, Faculty of Engineering, King Abdulaziz University, P.O. Box 80204, Jeddah, 21589, Saudi Arabia. ²Department of Mathematics, Cankaya University, Ogretmenler Cad. 14, Balgat, Ankara, 06530, Turkey. ³Institute of Space Sciences, Magurele, Bucharest, Romania. ⁴Department of Mathematics, Azarbaijan Shahid Madani University, Azarshahr, Tabriz, Iran.

Acknowledgements

Research of the second and third authors was supported by Azarbaijan Shahid Madani University. Also, the authors express their gratitude to the referees for their helpful suggestions, which improved the final version of this paper.

Received: 27 May 2013 Accepted: 28 August 2013 Published: 11 Dec 2013

References

- Bai, Z, Lu, H: Positive solutions for boundary value problem of nonlinear fractional differential equation. *J. Math. Anal. Appl.* **311**, 495-505 (2005)
- Baleanu, D, Mohammadi, H, Rezapour, S: Positive solutions of a boundary value problem for nonlinear fractional differential equations. *Abstr. Appl. Anal.* **2012**, Article ID 837437 (2012)
- Baleanu, D, Mohammadi, H, Rezapour, S: Some existence results on nonlinear fractional differential equations. *Philos. Trans. R. Soc. Lond. A* **371**, 20120144 (2013)
- Baleanu, D, Mohammadi, H, Rezapour, S: On a nonlinear fractional differential equation on partially ordered metric spaces. *Adv. Differ. Equ.* **2013**, 83 (2013)

5. Kilbas, AA, Trujillo, JJ: Differential equations of fractional order: methods, results and problems. *Appl. Anal.* **81**(2), 435-493 (2002)
6. Zhang, SQ: Existence of positive solution for some class of nonlinear fractional differential equations. *J. Math. Anal. Appl.* **278**, 136-148 (2003)
7. Agarwal, RP, O'Regan, D, Stanek, S: Positive solutions for mixed problems of singular fractional differential equations. *Math. Nachr.* **285**(1), 27-41 (2012)
8. Cabrera, IJ, Harjani, J, Sadarangani, KB: Existence and uniqueness of positive solutions for a singular fractional three-point boundary value problem. *Abstr. Appl. Anal.* **2012**, Article ID 803417 (2012)
9. Cao, J, Chen, H: Positive solution of singular fractional differential equation in Banach space. *J. Appl. Math.* **2011**, Article ID 352341 (2011)
10. Liu, Y, Nieto, JJ, Otero-Zarraqinos, O: Existence results for a coupled system of nonlinear singular fractional differential equations with impulse effects. *Math. Probl. Eng.* **2013**, Article ID 498781 (2013)
11. Wei, X, Lu, X: The periodic solutions of the compound singular fractional differential system with delay. *Int. J. Differ. Equ.* **2010**, Article ID 509286 (2010)
12. Zhou, WX, Chu, YD, Baleanu, D: Uniqueness and existence of positive solutions for a multi-point boundary value problem of singular fractional differential equations. *Adv. Differ. Equ.* **2013**, 114 (2013)
13. Baleanu, D, Agarwal, RP, Mohammadi, H, Rezapour, S: Some existence results for a nonlinear fractional differential equation on partially ordered Banach spaces. *Bound. Value Probl.* **2013**, 112 (2013)
14. Baleanu, D, Bhrawy, AH, Taha, TM: Two efficient generalized Laguerre spectral algorithms for fractional initial value problems. *Abstr. Appl. Anal.* **2013**, Article ID 546502 (2013)
15. Baleanu, D, Diethelm, K, Scalas, E, Trujillo, JJ: *Fractional Calculus Models and Numerical Methods*. Series on Complexity, Nonlinearity and Chaos. World Scientific, Singapore (2012)
16. Bhrawy, AH, Alghamdi, MA: A shifted Jacobi-Gauss-Lobatto collocation method for solving nonlinear fractional Langevin equation involving two fractional orders in different intervals. *Bound. Value Probl.* **2012**, 62 (2012)
17. Bhrawy, AH, Alofi, AS: The operational matrix of fractional integration for shifted Chebyshev polynomials. *Appl. Math. Lett.* **26**, 25-31 (2013)
18. Bhrawy, AH, Al-Shomrani, MM: A shifted Legendre spectral method for fractional-order multi-point boundary value problems. *Adv. Differ. Equ.* **2012**, 8 (2012)
19. Bhrawy, AH, Tharwata, MM, Yildirim, A: A new formula for fractional integrals of Chebyshev polynomials: application for solving multi-term fractional differential equations. *Appl. Math. Model.* **37**, 4245-4252 (2013)
20. Podlubny, I: *Fractional Differential Equations*. Mathematics in Science and Engineering, vol. 198. Academic Press, San Diego (1999)
21. Krasnoselski, MA: *Positive Solutions of Operator Equations*. Noordhoff, Groningen (1964)

10.1186/1687-1847-2013-359

Cite this article as: Baleanu et al.: The existence of solutions for a nonlinear mixed problem of singular fractional differential equations. *Advances in Difference Equations* 2013, **2013**:359

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com