


The Expanded Very Large Array

Michael P. Rupen
NRAO/Socorro

Tenth Summer Synthesis Imaging Workshop
University of New Mexico, June 13-20, 2006




Why Expand the VLA? Unique Radio Capabilities


2

Magnetic Fields

Polarization
Faraday rotation
Zeeman splitting




Low Obscuration




- No dust
- no bias
- see inner cores, where the action is

Transient Sources

- Trace shocks and ejecta
- Observe 24/7, regardless of weather, Sun, etc.
- High resolution



The Evolving Universe



- Accretion ↔ outflow
- Trace both thermal and non-thermal (AGN, Hii, etc.)
- Key lines

Why Expand the VLA? The Art of the Possible

3

- The VLA is still the most flexible and sensitive radio telescope in the world. But...
 - it's **over 30 years old**: the first VLA antenna came on-line on 24 October 1975
 - major improvements are possible, at very little cost: **keep the infrastructure** (antennas, railroad track, buildings, ...), but **replace the electronics**

The EVLA: Order-of-Magnitude Improvements


4

Parameter	VLA	EVLA	Factor
Sensitivity (1σ , 12 hours)	10 μ Jy	1 μ Jy	10
Maximum BW per polarization	0.1 GHz	8 GHz	80
# of frequency channels at max. bandwidth	16	16,384	1024
Maximum number of frequency channels	512	4,194,304	8192
Coarsest frequency resolution	50 MHz	2 MHz	25
Finest frequency resolution	381 Hz	0.12 Hz	3180
(Log) Frequency Coverage (1 – 50 GHz)	22%	100%	5

- **EVLA cost is less than $\frac{1}{4}$ the VLA capital investment**
- **No increase in basic operations budget**

Point-Source Sensitivity Improvements : 1- σ , 12-hours

5




Red: Current VLA,

Black: EVLA Goals

Frequency - Resolution Coverage

6

- Continuous frequency coverage from 1 to 50 GHz
- match instrument to science, not science to instrument!
- Blue area shows current VLA frequency -resolution coverage.
- Green area shows future EVLA coverage.
- Yellow letters and bars show band names and boundaries.
- Two low frequency bands (74 and 327 MHz) omitted



Bandwidth and Spectral Capabilities

7

- Combination of 2:1 bandwidth ratios and huge number of spectral channels
 - instantaneous spectral indices, rotation measures, uv-coverage
 - instantaneous velocity coverage (53,300 km/s vs. current 666 km/sec at 45 GHz)
 - lines at arbitrary redshift
- Ridiculously flexible correlator
 - 128 independently tunable sub-bands, vs. 2 now
 - “zoom in” on the regions of interest, and leave one 2 GHz baseband for continuum

The Time Domain

8

- Dynamic scheduling
 - use weather efficiently
 - respond to transients
- Fast time recording: initially 100 msec; 2.6 msec possible
- Pulsars: 1000 phase bins of 200 μ sec width, 15 μ sec possible
 - pulsar searches, timing, etc. with an interferometer!


Molecular Studies of High-Redshift Star-Forming Galaxies

9

•Currently:

- 50 MHz (z range of 0.001 at 50 GHz!)
- 8 spectral channels

- No z searches
- Very poor spectral resolution
- Resolve out wide lines, and add noise to narrow ones
- Each line must be done independently (CO, HCN, HCO+, ...)



CO_{J=3-2} Carilli, Walter, & Lo
 Z = 6.42
 Peak ~ 0.6 mJy


Molecular Studies of High-Redshift Star-Forming Galaxies

10

•EVLA:


- 8 GHz (z=1.4 to 1.9 for CO J=1-0; z=3.8 to 4.8 for 2-1)
- 16384 spectral channels (1 MHz res'n= 5.0 km/s)

- 200 km/s galaxy is 40 channels
- Every line at once
- Interferometry:
 - spatial res'n
 - excellent spectral baselines



Quasar Absorption Line Surveys

11




Absorption lines at $z=0.88582$
towards PKS 1830-211 (C. Carilli)

- **Unbiased** line surveys:
 - no dust obscuration
 - lots of *random* background sources
- **HI, CO, HCN, HCO+, ...**
 - ⇒ evolution of cosmic neutral baryons from $z=0$ to 3
 - ⇒ large-scale structure
 - ⇒ estimates of CMB temperature

Star-Forming Galaxies at High Redshift

12


- Sensitive to:
 - **Synchrotron** emission: AGN, SNR
 - **Free-free** emission: Hii regions
 - Thermal **dust** emission
- Resolution **50 mas: 200 pc @ $z=10$**
- Imaging: **1 arcsec over 30 arcmin @ 1.5 GHz**
- EVLA+ALMA give **complete galaxy SED's**
 - 3 orders of magnitude of frequency
 - large range of redshift



Arp220 SED scaled to high redshifts.

Galaxies Closer to Home

13




3x8 hours on a typical spiral galaxy

In **one** observation of a galaxy:

- deepest **radio continuum** image yet made, with **spectral index** too
- image **all (UC) HII's & SNRs**
- map **HI emission & radio recombination lines**
- measure **magnetic field** orientation, Faraday rotation, and Faraday depth
- absorption measurements against 100s of **background sources**
 - also rotation measures!
- simultaneous **"blind" HI survey**

Strong Gravity and Black Hole Accretion: The Galactic Center

14



VLA: 1 cm (Zhao)



VLT / NACO 1.6-3.5 microns

Strong Gravity and Black Hole Accretion: The Galactic Center

15


EVLA: the radio view

- **100s of pulsars** with $P_{\text{orbit}} < 100$ yr
 - higher frequency to avoid dispersion due to ionized gas
 - image fidelity (SgrA*:pulsar = $1e6:1$)
 - 10's mas astrometry
 - millisecond pulsar timing
- **complete survey & monitoring of OH/IR stellar masers**
 - detailed rotation curve
- **3D motions of ionized gas**
 - free-free emission + radio recombination lines
- **magnetic field structures and strength**
- **Mass and *spin* of a supermassive black hole**
 - deviations from elliptical orbits
- **Extended dark matter distribution**
- **Tests of GR in ultra-strong regime**
- **Detailed accretion estimates**
- **Gas vs. stellar motions**


Magnetic Fields in Galaxy Clusters

16

with X-rays, map magnetic fields & electron density in detail across entire, individual clusters



Rotation measures towards Hydra A (G. Taylor)




Residual RM towards 22 Abell clusters (T. Clarke)

- unambiguous rotation measures
- >100 sources per beam (vs. current 1-2) for scattering & polarization studies
- >20 RRM per cluster for >80 clusters!
- much less depolarization

Galactic Black Holes: The Accretion/Outflow Connection


17

- Ubiquity of jets
- Monitoring
 - continuous multi-freq. coverage
 - work at 45 GHz → 50mas res'n
 - triggering VLBI
- Polarization
- Going deeper
 - faint source imaging
 - typical rather than 20σ sources
 - other disk states
 - other source types (e.g., ULXs, low-luminosity XRBS, NS, etc.)



Stars

18




Stars detected with the VLA (S. White)

- first detections of **ordinary stars** like the Sun
- track radio emission from **young stars** (10^6 to 10^7 years)
- flares in **pre-main-sequence stars**

- **Zeeman splitting** of H recombination lines directly measures ISM magnetic fields
- Splitting is weak – 2.8 Hz per μG \rightarrow stack multiple lines
- 2-4 GHz band: 31 recombination lines
 - Each typically 250 kHz wide \rightarrow $\sim 0.4\%$ of the total band.
 - Need 10 kHz resolution
- So, either 400,000 channels...or zoom in with WIDAR!

WIDAR Setup


- Each line individually targeted
- H, He, C lines all within 4 MHz sub-band
- Res'n 15.6 kHz (1.6 km/sec)
- Each of 62 spectra gets 256 channels
- Or: use 8 MHz sub-bands with 4192 channels \rightarrow 0.2 km/sec resolution



- EVLA resolution provides images of:
 - gas density,
 - temperature,
 - metallicity,
 - B-fields (Zeeman)
- Sensitivity (12 hr, 5σ):
 - $\Delta S_{\text{line}} \sim 0.1 \text{ mJy}$ (stacked, integral)
 - $\Delta B \sim 150 \mu\text{Gauss}$.
- Orion, W3, Gal. Center ...

- 414 lines (8 to 50 GHz)
- 38 species
- EVLA offers
 - Spatial resolution
 - Spectral baseline stability
 - Full polarization (Zeeman splitting!)
- EVLA can observe **8 GHz** at once – an average of **80 lines** at **10 km/s** velocity res'n (30 GHz)
- EVLA can “target” many (**~60**) lines at once

Sky Frequency Bands							
1-2	2-4	4-8	8-12	12-18	18-27	27-40	40-50
L	S	C	X	U	K	Ka	Q



TMC-1 (Nobeyama: Kaifu et al. 2004)

EVLA : Cost and Timescale

23

- Proposal (EVLA-I) submitted to NSF in 2000
 - Funding started in 2001 following NSB approval.
 - Completion by 2012
- A cooperative project:
 - \$57M from NSF, over eleven years
 - \$15M from Canada, (correlator, designed and built by HIA/DRAO)
 - \$2M from Mexico, and
 - \$8M from re-directed NRAO operational budget
- A second proposal (EVLA-II) was submitted in April 2004
 - Goal: to improve the spatial resolution by a factor 10
 - \$115M, over 7 years
 - The NSF recently (Dec 2005) declined to fund this proposal


EVLA Project Status

24

- Six antennas currently withdrawn from VLA service, and being outfitted with new electronics.
 - Two fully outfitted & available upon request
 - Two being outfitted with final electronics, and are being intensively tested. Available for astronomical use by late summer.
 - Two others in early stages of outfitting.
- Antennas will be cycled through the conversion process at a rate six per year, beginning in 2007.
- Except for special testing, no more than three antennas will be out of service at any one time during construction phase.

Challenges: Radio Frequency Interference

27



Challenges: Data Processing

28

- Data rates
 - peak from correlator backend: ~25 MB/s
 - 8-hour “peak” observation ~ 700 GB (average is factor 10 lower)
 - data for 1 year ~ 80 TB
- Analysis
 - data flagging
 - sources everywhere
 - full (wide!) bandwidth synthesis (must account for spectral index, pol'n, rotation measure, etc.)
 - high-fidelity imaging (10 mJy \Rightarrow 10⁴:1)

Challenges: Ease of Use

29

- Much more complex and capable system
 - correlator modes
 - “wide-open” bands
 - lots of data
- How do we make this power available to multi-wavelength users?
 - data volumes
 - “end-to-end” processing
 - imaging pipelines
 - readily accessible archive, NVO

EVLA Spin-offs

30

- Correlator for eMERLIN
- Renewed (international!) radio collaborations
 - common problems of data volume, deep imaging, etc.
- Centimeter/millimeter connection
 - similar timescales for EVLA & ALMA
 - similar techniques
 - comparable instruments, and complementary information on much shared science
- Opportunities as the VLA winds down
 - spectral line: e.g., deep HI images or surveys
 - time-dependent science: space telescopes, transient science, etc.
 - **Note Oct06 call for Large Proposals!**

Challenges: Looking Ahead

31

- Higher resolution: how can we tie in the VLBA?
 - bring high bandwidth (= sensitivity) to the world array
- Higher sensitivity: more collecting area for spectral line studies (the Square Kilometer Array)
 - requires economies of scale, for the antennas, the feeds & receivers, the correlator, etc. etc.
 - the EVLA as a pathfinder

Challenges: Strengthening the US Community

32

- NSF funds radio astronomy through grants
 - budget is very tight compared to NASA
 - no direct tie to telescopes
 - unhealthy perception of competition between instruments (esp. NRAO) and science
- ➔ Fabulous new instruments --- now we have to make sure they are used as fully as they can be!
 - international collaboration
 - obviously wonderful science
 - make it easier to use
 - more direct ties to space instrumentation (cf. Chandra)
 - innovative approaches within NRAO

- Staff support/collaboration
- These schools
- Travel support for US observers (NRAO and foreign telescopes)
- Page charges
- Paid sabbatical/summer visits
- Postdocs
 - Traveling & resident Jansky fellows
- Student support
 - GBT projects
 - grad students (2 mos.-2 years, full support)
 - undergraduates (Co-Op Program up to 1 semester/year; summer REU)
- Aggressively pursuing other innovative programs
- At last, we will be hiring!

- After a long dry spell, telescopes galore
 - GMRT, SMA, eVLBI
 - EVLA, ALMA, ATA, eMERLIN, LWA, LOFAR, Australian initiatives, LMT, ...
- Looming on the horizon: the Square Kilometer Array
- **This is the perfect time to be a graduate student!**
 - get in on the ground floor
 - influence “first science”, software design, how the arrays operate
 - a unique opportunity to mix technology, software, and science