



The role of health communication in the experiences of families who are invited to attend a community-based childhood obesity programme: Why do some families attend whilst others do not?

Journal:	<i>European Health Psychology Society</i>
Manuscript ID:	Draft
Manuscript Type:	Psychology and Health
Keywords:	childhood, obesity, health communication, attendance, qualitative

SCHOLARONE™
Manuscripts

Abstract

Objectives: Although numerous interventions are in place to attempt to reduce the high prevalence of childhood obesity, uptake and success of these interventions vary. Little previous research has investigated the reasons why families choose to attend (or not attend) recommended interventions; this study has therefore investigated the experiences which influence the decisions to attend or not to attend a childhood obesity programme.

Design: Eleven semi-structured interviews were employed to explore the experiences of families who had been invited onto a childhood obesity programme. Interviews were examined using interpretative phenomenological analysis.

Results: Data were grouped into four themes which highlighted the differences and similarities between attendees and non-attendees. Barriers to attendance included parents either not recognising the condition in their children or not acknowledging its impact on health or wellbeing; parents having anxieties about their own participation in the programme; parents' views on their children's identities and parents believing that they have little control over their environment.

Conclusions: It is suggested that obesity programmes must address preconceived negative perceptions regarding obesity, by carefully considering how the delivery of health information is presented, how it is understood, and how underlying identity may affect motivation to engage in new behaviours.

Key Words: childhood; obesity; health communication; attendance; qualitative

Introduction

Obesity is considered to be a 'lifestyle disease' with many contributing factors including genetic, economic, social, psychological, environmental and cultural elements (Butland et al., 2007). Childhood obesity poses many health risks; research has shown that 58% of children with a body mass index (BMI) above the 95th percentile have hypertension, hyperlipidaemia or insulin resistance and 25% have two or more of these (Allen et al., 2007; Roberts, 2007; Rudolf, 2004). It has also been found to affect psychological wellbeing, especially self-esteem (Dietz & Robinson, 2005). It is reported that most overweight or obese children become overweight or obese adults and it is projected that the current generation of children will have shorter life spans than their parents' generation (Ludwig, 2007; Butland et al., 2007).

Overweight and obesity are stated as 'probably the most widespread threat to health and wellbeing' in England (Department of Health [DoH], 2011a). The annual results of the 2010/11 National Child Measurement Programme found that by 10-11 years old, a third (33.4%) of children in England were classified as either overweight or obese (The National Health Service [NHS] Information Centre, 2011). The UK government recognise that obesity has negative consequences for individuals, local communities, and to the NHS (as direct costs which are estimated at £5bn each year) and has implications to wider society and economic development. The government's ambition is to ascertain 'a sustained downward trend in the level of excess weight in children by 2020' in England (DoH, 2011a).

Evidence suggests that obesity services should combine dietary, physical activity and behavioural elements (Luttikhuis et al., 2009) and interventions should be family based, with the participation of at least one parent (Dietz, & Robinson 2005; National Institute for Health and Clinical Excellence [NICE] 2006,). Thus, childhood obesity treatment can be classed as a 'complex intervention' (Campbell et al., 2007), built from numerous components,

1
2
3 implemented by a range of professionals, and requiring the family to make sustained lifestyle
4 changes. However, one of the biggest hurdles to the success of implementing these
5 programmes (complex interventions) is the role of client attendance and retention (Bower et
6 al., 2009). Non-attendance at appointments costs the health service time and money.
7 Moreover, non-attendance at programmes directly impacts upon those individuals who do not
8 attend; either as their 'condition' is not investigated/supported (or may get worse); or that
9 there are other unknown consequences of not attending.
10
11
12
13
14
15
16
17

18
19 Whilst some evidence shows that recruitment onto obesity programmes is very
20 successful (Rudolph, 2004) other clinics experience difficulties in recruitment (Haisman,
21 Matyka & Stanton, 2005) and retention (Cote et al., 2004; Luttikhuis et al., 2009; Zeller et al.,
22 2004). Research suggests that to increase uptake and retention, services should consider
23 improvements in access to care, improving quality of care, and reducing costs (NICE, 2007).
24 However, there is limited research which highlights reasons for non-attendance at childhood
25 obesity services; or more specifically which considers the experiences of families invited to
26 attend a childhood obesity intervention and living in deprived areas of England (Luttikhuis et
27 al. 2009). As recommended by Luttikhuis et al.'s (2009) systematic review, qualitative
28 research is needed to ascertain the views of participants; specifically highlighting why
29 childhood obesity interventions may be more or less successful and why families may choose
30 to attend or not attend recommended treatment programmes. If obesity rates in the UK are to
31 be reduced then there is a need to develop a greater understanding of reasons for non-
32 attendance and to consider ways to improve engagement in lifestyle interventions, a step
33 towards this development is to understand families' beliefs towards obesity and their
34 experiences of interventions.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 56 **Objective** 57 58 59 60

1
2
3 The aim of this research was to explore why families choose to attend or not attend a
4 childhood obesity programme, and to examine the beliefs and experiences which influenced
5 these decisions. The research objectives were to highlight possible improvements for the
6 'studied' service and to contribute to the local Public Health strategy suggesting how to
7 tackle prevalence and the treatment of obesity in the future.
8
9
10
11
12

13 14 15 16 **Methods**

17 *Design*

18 This study utilized qualitative interview methodology as it explored families'
19 perceptions of obesity and assessed the perceived appropriateness of the recommended
20 intervention.
21
22
23
24
25
26

27 *Participants*

28 *Ethical approval*

29 Approval was received from a NHS research ethics committee; NHS clinical governance
30 approval was granted and a full independent peer review and monitoring process was conducted
31 by a UK university.
32
33
34
35
36
37

38 *Participant criteria*

39 Families eligible to participate in this research were identified directly from referrals
40 made to the NHS service. Potential participants were required to meet the following
41 inclusion criteria: child to have been referred to the service and identified as clinically obese
42 (BMI percentile greater than 98th); child and family invited to attend (prescribed) a
43 community obesity treatment programme (see Figure 1 for service details).
44 Parent/Guardian(s) of the children were required to provide informed consent; children were
45 required to provide informed assent.
46
47
48
49
50
51
52
53
54
55

56 Figure 1 about here.
57
58
59
60

Participant Recruitment

Families were provided with an initial information sheet explaining the purpose and requirements of the study. Consent to pass participants' contact details onto the research team were ascertained during the initial clinical assessment. The researcher contacted the families and provided further information about the study, explaining the process and agreeing the location and time of the interview. Interviews were conducted at the participants' homes, an NHS clinic, or a local children's centre, depending on their preference. Due to the nature of the study, it was important that the participants felt comfortable expressing their experiences honestly and openly, and that they were given the opportunity to provide feedback as they deemed necessary. The study therefore concentrated on the responses of parents (as parents were the main decision makers), however children were asked if they would like to participate and contribute to the research investigation.

Research location: The health of people in the area studied was generally worse than the England average (DoH, 2011b): deprivation was higher than average (with 62.2% of residents living in the 'most deprived' quintile, compared to England average of 19.9%). The level of children living in poverty was significantly greater than the England average (32.8% vs. 20.9% respectively). The population of the area was predominately white Caucasian, with only 2% of the population from a minority ethnic group (Langton, 2011). Rates of obesity in the studied area were one of the highest in England, with prevalence of obesity in 10-11 year olds recorded at 24.4%; compared to the England average of 19% (DoH, 2011b).

Research sample: Upon attendance at clinical assessment, all families who met the programme criteria were systematically invited into the research study. Twenty-one families were invited to participate, of which fourteen families agreed to take part, and eleven participated (reasons for non-participation were: 1 incorrect phone number; 1 onset of family illness so withdrew; 1 no suitable date). All participants were white Caucasian; and included

1
2
3 working full/part-time and unemployed parents or guardians. All families lived in an area of
4 high deprivation. It was deemed that this cohort was a representative sample of the local
5 population.
6
7
8

9
10 For three interviews the guardian(s) was/were the child's grandparent(s) (families 1, 8
11 & 11). The children's ages ranged from 5 to 15 years old, with a mean age of 10 years old.
12 Four children participated in the research interviews (families 1, 2, 8 and 11). Of the seven
13 other families, four parents requested that the children were not interviewed and in three
14 cases the parents were happy for the child to participate; however, two children declined
15 participation and one child was not able to participate due to parenting arrangements after
16 school. Participants were categorised as either programme "attendees" (families 2, 3, 4, 5,
17 and 11) or "non-attendees" (families 1, 6, 7, 8, 9 and 10). Attendees were the families who
18 attended the programme for at least the first two sessions. Non-attendees were the families
19 who did not attend any intervention sessions.
20
21
22
23
24
25
26
27
28
29
30

31 **Materials**

32
33
34 Smith and Osborn (2008) report that the best data collection method for an
35 interpretative phenomenological analysis (IPA) study is via semi-structured interviewing;
36 therefore a semi-structured interview schedule was devised utilising open ended questions.
37 The interview was structured in two parts: firstly to introduce the families to the study and
38 gain their demographic information; secondly to ask questions about their experiences and
39 reasons for choosing to attend or not to attend. For example some of the questions included:
40
41
42
43
44
45
46
47 *"Can you tell me when you were invited/referred to the childhood obesity service - How did you*
48 *feel?"* *"Why did you decide to (not) attend the programme?"* *"What reasons do you think other*
49 *families might give for (not) attending?"* The interview process allowed for flexibility in
50 wording/ordering and expansion of questions depending on the interview flow. So to explore
51 the psychological world of the participants, the questions for the interview were loosely based
52
53
54
55
56
57
58
59
60

1
2
3 upon the Theory of Planned Behaviour (Ajzen, 1991). Structuring the interview in this way
4
5 aided the extraction of how participants' thoughts and beliefs regarding obesity and its
6
7 treatment affected their decisions to attend the intervention, and with the flexibility of the
8
9 semi-structured interview, this approach allowed the participants to tell their stories and for
10
11 the interview to uncover potentially novel areas for interpretation.
12

13 14 ***Procedure***

15
16 Digitally-recorded semi-structured interviews were conducted by the interviewer, who
17
18 was previously unknown to the families and was not part of the clinical intervention team.
19
20 The interviewer was therefore considered neutral and held an independent relationship with
21
22 participants, which would not impact in any way on their current or future healthcare services.
23
24 Participants were informed that the interviews aimed to seek their experiences of the services
25
26 received so far and to collect their views about attending the programme. Interviews lasted
27
28 an average of 57 minutes; the digital recordings were transcribed verbatim with all names
29
30 replaced by pseudonyms. Participants were offered the opportunity to read the transcribed
31
32 interviews to check for readability of data, and if required to clarify comments. No
33
34 participants requested to amend or clarify their original interview transcripts.
35
36
37

38 39 ***Analysis***

40
41 The data was examined using Interpretative Phenomenological Analysis (IPA) as
42
43 described by Smith, Flower, and Larkin (2009). The process was conducted by two authors
44
45 (*author* and *author*) who read the transcripts several times and initial comments and
46
47 interpretations were noted. Subsequently coding for meaning, similarities, differences, and
48
49 contradictions were highlighted in the left hand margin of each script. Once completed the
50
51 scripts were re-read with emerging themes (based upon these initial codes) noted in the right
52
53 hand column. The coding and themes were mapped, clustered and then discussed, reviewed
54
55 and defined by researchers before creating full definitions and interpretations. The process of
56
57
58
59
60

1
2
3 interpretation aimed to explore the responses of the participants to specific questions and to
4
5 look for emerging commonalities/differences. The construction and reliability of themes was
6
7 carefully discussed between the authors, so that any interpretation was representative and not
8
9 skewed by any underlying preconceived perceptions. Care was taken to ensure that emerging
10
11 themes fitted the data rather than forcing the data to fit into the themes. As a consequence the
12
13 descriptions were written based upon these analytical discussions. This process was
14
15 idiographic, looking in detail at one interview, before examining subsequent interviews.
16
17
18
19

20 21 **Results**

22
23 There were many factors which impacted families' decisions to attend, these were
24
25 categorised into four master themes: perceptions of childhood obesity; perceptions of the
26
27 programme; practical barriers and overcoming hurdles to attending; and availability and
28
29 suitability of local facilities. Within each theme the beliefs and experiences of 'attendees'
30
31 were compared for similarities and differences with those who were 'non-attendees'. The
32
33 following table summarises these themes highlighting the differences and similarities across
34
35 the participants. Each of these themes is subsequently described within this section.
36
37

38
39 Table 1 about here.

40 41 ***Theme 1: Perceptions of childhood obesity***

42
43 It was evident that the majority of parents, for both those who attended and those who
44
45 were non-attendees, did not perceive their child to be visually obese. Many referred to their
46
47 child as 'normal', and were shocked when, during their medical assessment, the clinician had
48
49 explained their child's diagnosis. The following extract provides insight into the families'
50
51 inabilities to recognise obesity based on appearance.
52
53
54
55
56
57
58
59
60

An extract of Family 1's (non-attendees) perception of obesity:

Grandparent 1, Jeff: '.... We know he is defined as obese 'cause that's what they (health professionals) told us...(pause) but look at him...(points over, Peter smiles) he's normal, just like other children, we know we need to cut out the puds (puddings)'

Child, Peter: 'Yeh I know [I] have a bit, like me dad, but it's just normal and I'm not bothered'

Grandparent 1, Jeff: ' Yeh, that's right, just like his dad, who has always been big...(pause)'

Child, Peter: 'Ha ha like not like him up the road!'

[laughs] (all of the family together expand their cheeks)

Grandparent 2, Gail : 'What we mean is and we don't know what his situation is, but up the road is a BIG boy the same age as Peter and you can see that fat all over him, he really needs to go to see someone...but it's not for us to comment on that is it, hey?'

Even though Family 1 (non-attendee) understood that their child was categorised as obese, they referred to him as 'just a tiny little bit over the line' [*reference to the BMI percentile charts*] and Peter (child), himself said 'I'm just a little bit over, but nothing to be worried about'. This family had identified another child as obese, by referring to his appearance and making reference to his behaviour. However, they did not make an association between the needs of their child (Peter) and this 'big boy'. The perceived vulnerability to health risks and the perceived severity of being defined as clinically obese was minimised by the review and comparison against another child.

There seemed to be a difference in the perception of obesity when related to health. Those parents or guardians who recognised that obesity was affecting the health of their child were more likely to attend the programme:

'I worry about Simon (*be*)cause of his health and (*paediatrician*) has said that (*be*)cause of his size over the 98th (*percentile*), I was shocked when I found that out, that he needs to do lots more exercise' (Tara, mother, family 2, attendee)

However, those parents/guardians who did not perceive their child's health to be affected by obesity were often non-attendees:

1
2
3 'He doesn't need obesity treatment, it's normal and he will grow bigger...anyway there is nothing
4 medically up with him is there?!' (Alan, father, family 7, non-attendee)
5

6
7 The above comment is particularly important as it illustrates two points: firstly, this parent's
8 perception is that childhood obesity is 'normal', and whilst this comment was made by a non-
9 attendee, this was a commonality across most of the families interviewed. Secondly, the
10 reference to medical diagnosis highlights contradictions in the interpretations of health
11 messages communicated by different professionals and this was a key concern for all parents
12 (attendee or not). All of the children who participated in the programme had been clinically
13 assessed for co-morbidities to identify potential medical causes of obesity. In all cases, no
14 medical causes of obesity were highlighted and the results of the medical assessment were
15 communicated as '*fit and healthy*' (father, family 7, non-attendee). This was perceived as
16 contradictory to the explanation given by the childhood obesity delivery team, who had
17 already explained that the child was clinically obese and required supportive intervention to
18 prevent the onset of any future health issues.
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 'We did say we would go but, but hmm, you know... (*pause*) it's not like cancer, if it was that bad
34 then he would have appointments anytime and have to come out of school, but they did it at night
35 when we have so many other things to do...' (Eileen, grandparent, family 8, non-attendee)
36
37

38
39 and

40
41 'We went along (*to the medical assessment*) like we were told to, and they said 'nothing
42 physically wrong with him', so he is normal and then they say 'come and do this'...why? They
43 should be on the peeps (*people*) that really need to get better' (Jane, mother, family 10, non-
44 attendee).
45
46
47

48
49 The process of communication seems to have created misunderstanding for some parents:
50 when parents could make the distinction between the medical assessment and the
51 consequences of obesity, they were more likely to attend. However, non-attendees tended to
52 interpret this medical communication as an over-ride for any previous messages delivered by
53 the obesity delivery team. Whilst most of the parents acknowledged that lifestyle changes
54
55
56
57
58
59
60

1
2
3 were required to reduce obesity, many of the parents (both attendees and non-attendees),
4
5 expressed confusion over the cause of their child's obesity, by linking the obesity to family
6
7 history; as a result of other illnesses; or as a result of taking medications (which may affect
8
9 metabolism).

10
11 'I was surprised when the nurse said that her [my child's] weight wasn't related to her asthma, and
12
13 taking medication, I mean like, I'm a nurse and we tell people on steroids that it can affect your
14
15 metabolism and things...so it must affect hers, but I was really shocked at that, I still think that, I
16
17 think I could have more information on that and it could be explained better, but we do need to
18
19 take on board more healthy lifestyle messages so that's why I said "okay" ' (Jessica, mother,
20
21 family 3, attendee)

22
23 Finally, it is important to note that perceptions of obesity expressed by other family
24
25 members or friends may have also influenced people's decisions to attend.

26
27 'Well dad (the child's father) thinks it's a waste of time, and mum (the child's grandmother)
28
29 knows, but they only see him once a week so they don't really have time to talk about it...they say
30
31 he is healthy and that's all that matters' (Tara, mother, family 2, attendee)

32
33 Whilst all interviewees described experiences of negative comments about the programme
34
35 from significant others (extended family/friends), attendance appeared to be linked to the
36
37 participants' abilities to actively manage any negative perceptions expressed by significant
38
39 others. Non-attendees appeared to conform with the views of significant others: when more
40
41 negative comments were expressed by close friends and family, they were less likely to
42
43 believe that the programme would have positive outcomes.

44
45
46 One of the parents summarises this theme well. As a health professional herself, she
47
48 presents some insight as to why other families do not attend the programme. She links the
49
50 negative perceptions of such a programme, the influence of significant others and the
51
52 consequences of obesity together, suggesting how this could be improved:

53
54
55 'I think that some people don't see that their child has a problem, and also don't think that it's
56
57 really based on the health, like her [her daughter Charlotte's] dad don't see it. I think that if there
58
59
60

1
2
3 was a session for the kids only and they played, and the parents had a session but just about the
4 health risks of obesity and what it is doin' to their bodies, with models and pictures and stuff, and
5 then they realise about it, and THEN tell them about the programme that it might make more
6 people go'. (Jessica, mother, family 3, attendee).
7
8
9

10 ***Theme 2: Perceptions of the programme***

11
12 Expectations and prior perceptions of what the programme would involve and who else
13 would be attending the programme, influenced the decision to attend. Those families who
14 predicted that the programme would have positive effects on their lives were more likely to
15 attend, and believed that the programme would: help parents seek support from other parents;
16 help children to make friends, play together and improve their social skills and self-
17 confidence; and facilitate both the parents and their children to gain new skills and
18 knowledge:
19
20
21
22
23
24
25
26

27
28 'Going to the programme will give me information and education on what to do better and where
29 to go locally afterwards' (Jessica, mother, family 3, attendee).
30

31 and

32
33
34 'It is good that we can go along and I can speak to other parents who are trying to change things,
35 also James (*child*) can make friends with other kids the same who [*may*] not be fantastic at
36 football and feel really bad, but they can play on the same level you know' (Ray, father, family 4,
37 attendee).
38
39
40

41 Non-attendees on the other hand, were more likely to highlight negative consequences of
42 attending the programme, such as viewing the opportunity for their child to mix with other
43 obese children as destructive, by suggesting that their child would subsequently identify with
44 other obese children and as a consequence, this belief acted as a barrier to attendance:
45
46
47
48
49

50 'The thing is, he doesn't look fat ...hmm... obese, so if we go and he thinks, 'do I look like the
51 others?' then that wouldn't be nice, don't you think?' (Eileen, grandparent, family 8, non-
52 attendee).
53
54
55
56
57
58
59
60

1
2
3 Some parents expressed concerns that they would be expected to participate in the
4 physical activities, and became concerned with how their own behaviour would be assessed.
5
6

7 This seemed to create some anxiety:
8

9 'We haven't exercised properly for a long time; I have never been to a gym, so telling us that we
10 have to do everything with the kids, it's embarrassing' (Hilary, mother, family 9, non-attendee)
11
12

13 Whereas the non-attendees who expressed that they were less able to participate in activities
14 may, as a result, have used this as an explanation for not attending, the parents of attendees
15 also expressed concerns, but recognised that they themselves may need support in changing
16 their own lifestyle behaviours. Most parents reported having some knowledge about healthy
17 eating and knew that physical activity was beneficial. Those who were non-attendees were
18 most likely to state that they were already eating healthily and did not need further
19 information, or that they knew who to ask for information if they needed it:
20
21
22
23
24
25
26
27

28 'Like we already know about healthy eating, and doing exercise, blah blah blah, the diet person is
29 just gonna (*going to*) tell us stuff we already know anyway, we are well, we don't eat pizza and
30 KFC (*Kentucky Fried Chicken*) all the time' (Liz, mother, family 6, non-attendee).
31
32
33

34 ***Theme 3: Practical barriers and overcoming hurdles to attending***

35
36 All families raised similar practical barriers that would stop them from attending the
37 programme. Time to attend the programme was problematic for all. However, although
38 attendees highlighted that time was an obstacle, they described that they tried to 'make time'
39 to go and if they had missed sessions due to work or family issues, that they felt 'guilty'
40 (mother, family 2, attendee). Whilst interviewees understood that the programme was
41 delivered so not to interrupt the school day and parents' working hours, some attendees
42 reflected on others' decisions not to attend, suggesting that the time of the programme could
43 change so that it would be convenient for more people:
44
45
46
47
48
49
50
51
52

53 'Maybe Saturday morning would be okay and maybe think about it, or could come out of school if
54 it were that important' (Sandra, mother, family 5, attendee).
55
56
57
58
59
60

1
2
3 Furthermore, even though the obesity service was able to offer payment for transport,
4
5 reference to transport and travel duration to and from the programme was deemed a
6
7 consideration for all.
8

9
10 'If we had to go...we haven't got room, only got a little car... if it's far away then it takes too
11
12 long, by the time he [child, Peter] got home from school and then we get back it's bed(*time*)...'
13 (Jeff, grandparent, family 1, non-attende).

14
15 and

16
17 'Well we have the car so that's okay and we can park. I probably wouldn't be able to do it on the
18
19 buses and walking and things 'cause (*because*) by the time you add it up it takes hours.' (Tara,
20
21 mother, family 2, attendee).
22

23 In addition to time and transport, the location and personal security for families was
24
25 a key influencing factor towards decisions to attend. If the families thought that
26
27 participating in the programme would negatively affect their safety then they would be
28
29 less likely to attend:
30

31
32 'We can get there dead easy and it's always at the school, so that's okay and it's good [*that*] it's
33
34 in a separate building, but if there were kids hangin' around outside or anythin' then it would put
35
36 me off a bit, like if you were going the hospital they would have security and stuff [*but at the*
37
38 *school they don't*]' (Tara, mother, family 2, attendee).
39

40
41 and

42
43 'Even if it was close, we wouldn't walk (*be*)cause it's not safe in the dark and in the summer it's
44
45 not fair (*be*)cause there are gangs of kids and that worries us, doesn't it?' (Jeff, grandparent,
46
47 family 1, non-attende).

48 ***Theme 4: Availability and suitability of local facilities***

49 All participants reported that within the local area there weren't enough activities for their
50
51 children to do. Parents were concerned about the suitability of the local clubs/activities they
52
53 found for their children. This was reflected in terms of their children's ability and skill level
54
55
56
57
58
59
60

1
2
3 to be able to participate in the activities, the acceptability of their children to the activity
4
5 (social group), and in terms of their children's safety attending activities away from home.
6

7 'You know we have all the information and it's really frustrating (*be*)cause there is nothing for
8 him to do, like the exercise. I was really cross (*be*)cause I was taking him to football club, but
9 (*be*)cause of how he is they just left him twisting round in the pitch on his own and he needs extra
10 support to encourage him'. (Tara, mother, family 2, attendee).
11
12

13
14 Attendees were more likely to seek out information on local clubs and activities,
15 however it was acknowledged that local clubs were geared more towards 'sporty and talented
16 children' rather than open to all (particularly those not already engaged in activity). Some
17 attendees reported that this obesity programme 'filled the gap'. However, of concern to non-
18 attendees was the short duration of phase 1 of the programme, highlighting that afterwards
19 they would be in the same situation (lack of appropriate clubs).
20
21
22
23
24
25
26

27
28 'Things will be the same at the end (*of the programme*). Where will he go? What will he do?'
29
30 (Alan, father, family 7, non-attendee).
31

32 Parents explained that ensuring that children continued to exercise after the programme was
33 very difficult, not just because of the lack of access to appropriate clubs and services, which
34 the whole family could do together, but due to lack of safe places for the children to play:
35
36
37

38 'We try and do more activity, but they can't play outside, it's not safe, and there isn't enough
39 room in here (*house*) to be jumping around' (Jeff, grandparent, family 1, non-attendee).
40
41

42 Non-attendees in particular viewed the programme as a quick fix to accessing activity
43 services, which wouldn't remain upon completion of the programme.
44
45
46
47

48 Discussion

49
50 It was evident in this study that most parents (or guardians) did not recognise, by
51 appearance, that their child was obese. This concurs with findings from previous research
52 that has demonstrated that even when a child is obese, few parents recognise this in their own
53 child (Jeffery, Voss, Metcalf, Alba & Wilkin, 2005; White, O'Brien, Houlihan, Darker &
54
55
56
57
58
59
60

1
2
3 O'Shea, 2012) and/or are able to recognise the health implications of their child's weight
4
5 (Baur, 2005; Carnell, Edwards, Croker, Boniface & Wardle, 2005; Etelson, Brand, Patrick et
6
7 al., 2003; Jain, Sherman, Chamberlin et al., 2001; Jeffery, Voss, Metcalf, et al., 2005). When
8
9 given the 'diagnosis' of obesity, attendees were more likely to recognise the impact that
10
11 obesity had on their child's overall health and wellbeing, whereas non-attendees were less
12
13 likely to believe that obesity affected health, or at least believed that the diagnosis was not as
14
15 serious as other illnesses such as cancer. This study adds to previous research, as not only did
16
17 parents not recognise obesity in their children, but once given the 'diagnosis' of obesity the
18
19 acceptance of the condition as a health problem was often rejected. Parents often appeared to
20
21 be in denial, referring to their own interpretation of health professionals' explanations. For
22
23 example the statement 'there is nothing medically wrong' provided by a nurse clinician to
24
25 suggest that the obesity was not caused by a bio-medical issue, was interpreted by families as
26
27 'their child was healthy', and therefore even if their child 'was a little overweight' it wasn't a
28
29 health concern. Some were also sensitive to the language and comments from the health
30
31 providers, which they may have perceived as negative and blaming towards them. This
32
33 suggests that the way that the 'diagnosis' of childhood obesity is initially communicated
34
35 between the health professionals and families is key. Ultimately, the understanding resulting
36
37 from this communication may not only influence a family's decision to attend a childhood
38
39 obesity programme such as this, but is also likely to impact future lifestyle choices and
40
41 behaviour.

42
43
44
45
46
47
48 Interestingly in this study, those who were attendees stated that they would like the
49
50 children to improve their social skills and become more confident. In contrast, those parents
51
52 who were non-attendees were likely to express negative psychological effects of attending the
53
54 programme, such as experiencing concerns that their child would subsequently associate
55
56 (identify) themselves with other obese children. One explanation of these findings could be
57
58
59
60

1
2
3 described in terms of Oyserman's (2007; 2009a; 2009b) Identity-Based Motivation (IBM)
4
5 model which suggests that when behaviour is linked to identity (identity-infused), engaging
6
7 in the behaviour should carry a positive tone of inclusion in the in-group. For those families
8
9 in this study whose in-group (family and friends) associate negative attributes towards obese
10
11 people, it may be more acceptable for them to deny the existence of childhood obesity.
12
13 Should these families respond to the invitation to join an obesity intervention, this may create
14
15 identity conflict (socially, culturally or individually, either consciously or not), because they
16
17 have accepted the classification of obesity, and therefore have labelled themselves with the
18
19 negative characteristics of being obese (Puhl & Heuer 2009). In essence for some families,
20
21 attending an obesity intervention may challenge social, cultural and individual-cognitive
22
23 identities (see Oyserman & Destin, 2010) and, moreover, attendance at the intervention itself
24
25 may promote a shift in identity, which may or may not initially be perceived as a positive
26
27 outcome.
28
29
30
31

32
33 Non-attendees were more likely to highlight feelings of anxiety associated with
34
35 concerns that parents would also be expected to participate in physical activity. Whilst the
36
37 obesity intervention aimed to create an environment to improve the self-efficacy of the child
38
39 (see Trost, Kerr, Ward & Pat, 2001) by, for example, allowing the obese children to observe
40
41 influential others (i.e. parents and peers perform physical activity), support should also be
42
43 provided to increase the self-efficacy of parents performing this behaviour (Smith et al.,
44
45 2010). In addition, this study found that the views and opinions of significant others (family,
46
47 friends, social network) influenced the family's initial decision to attend the intervention.
48
49 This reflects findings from previous research where it was found that grandmothers had a
50
51 strong influence on parents' feeding decisions for children (Black, Siegel, Abel, & Bentley,
52
53 2001). However, this study highlights that whilst most families experienced some negative
54
55 comments from others about the intervention and/or 'supposed diagnosis' of obesity,
56
57
58
59
60

1
2
3 attendees seemed more able to manage this conflict with their friends/family, whereas for
4
5 non-attendees this communication had a greater impact on attendance. It is therefore
6
7 important that views and beliefs of both direct and indirect family members are considered
8
9 during the initial communication of the programme.
10

11
12 The main environmental restrictors which influenced families' decisions to attend
13
14 were time and security. Lack of time emerged as a barrier for all parents, reflecting findings
15
16 from previous research by Chatterjee, Blakely and Barton (2005). It is important that service
17
18 providers address a barrier such as lack of time, because if clients cannot attend services, then
19
20 the service itself risks failing. Concerns about the environment, safe places for children to
21
22 play, clubs that were suitable for their child's ability (were not just for the 'elite' - those really
23
24 good at sport), and easy access to services were all highlighted in this study as barriers to
25
26 seeking treatment or changing lifestyle behaviour. Similar barriers have been reported in
27
28 other studies (Ellaway, Kirk, McIntyre, & Mutrie, 2007; Chatterjee et al, 2005). Lack of
29
30 time and concerns about security may highlight the importance of modifying social, cultural
31
32 and organisational environments to enable individuals to practice healthier behaviours
33
34 (Bassett & Perl, 2004).
35
36
37
38
39
40
41

42 **Limitations**

43
44 The sample was self-selected in nature; participants responded initially to the referral
45
46 to the childhood obesity programme and had attended their initial assessment. This self-
47
48 selection resulted in a sample that was fairly open to expressing their views and concerns
49
50 regarding their child's health and wellbeing. The study adopted interpretative qualitative
51
52 methodology and as such the results are not definitive cause/effect explanations for
53
54 attendance to obesity programmes. The results of this study were based upon IPA analysis on
55
56 a specific cohort of families living in a very deprived area of England. The findings of this
57
58
59
60

1
2
3 study may therefore not generalise across groups, but may be useful to consider the reasons
4
5 for non-attendance to similar groups and settings.
6
7

8 9 **Conclusion**

10
11 This study found that there are many reasons why individual families may or may not
12
13 engage in a childhood obesity programme. Some of the common barriers identified in this
14
15 investigation related to: parents either not recognising obesity in their children or not
16
17 acknowledging its impact on health or wellbeing; parents having anxieties about their own
18
19 participation in the programme; parents' concerns about their children's identities and parents
20
21 believing that they have little control over their environment. Changing eating and activity
22
23 behaviour is an active process in which people must exert considerable conscious effort to
24
25 change established habits. In order to be effective, obesity services and the local environment
26
27 must meet the needs of their clients and address any preconceived negative perceptions, by
28
29 carefully considering how the delivery of health information is presented, how it is
30
31 understood and how underlying identity may affect motivation to engage in new behaviours.
32
33
34
35
36
37

38 **Acknowledgements**

39
40 The authors would like to thank all the research participants, the service that assisted in the
41
42 recruitment process and to both X and X for reviewing an initial draft of this paper.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Ajzen, I. (1991). The Theory of Planned Behavior. *Organizational Behavior and Human Decision Processes*, 50, 179-211. doi:[10.1016/0749-5978\(91\)90020-T](https://doi.org/10.1016/0749-5978(91)90020-T)
- Allen, D. B., Nemeth, B. A., Clark, R., Peterson, S. E., Eickhoff, J., & Carrel, A. L. (2007). Fitness is stronger predictor of fasting insulin levels than fatness in overweight male middle-school children. *Journal of Pediatrics*, 150, 383-387. doi:dx.doi.org/10.1016/j.jpeds.2006.12.051
- Bassett, M., & Perl, S. (2004). Obesity: The public health challenge of our time. *American Journal of Public Health*, 94, 1477-1477. doi: 10.2105/AJPH.94.9.1477
- Baur, L. (2005). Childhood obesity: practically invisible. *International Journal of Obesity*, 29, 351-352. doi:10.1038/sj.ijo.0802931
- Butland, B., Jebb, S., Kopelman, P., McPherson, K., Thomas, S., Mardell, J., Parry, V. (2007). Foresight. Tackling Obesities: Future Choices- Project Report, 2nd Edition. London: Government Office for Science.
- Black, M.M., Siegel, E.H., Abel, Y., Bentley, M.E. (2001). Home and videotape intervention delays early complementary feeding among adolescent mothers. *Pediatrics*, 107, E67. doi: 10.1542/peds.107.5.e67
- Bower, P., Wallace, P., Ward, E., Graffy, J., Miller, J., Delaney, B., Kinmonth, A. L. (2009). Improving recruitment to health research in primary care. *Family Practice*, 2, 391-397. doi: 10.1093/fampra/cmp037
- Chatterjee, N., Blakely, D. E., Barton, C. (2005). Perspectives on Obesity and Barriers to Control From Workers at a Community Center Serving Low-Income Hispanic Children and Families. *Journal of Community Health nursing*, 22, 23-26. doi: 10.1207/s15327655jchn2201_3

- 1
2
3 Campbell, N.C., Murray, E., Darbyshire, J., Emery, J., Farmer, A., Griffiths, F., Guthrie, B.,
4
5 Lester, H., Wilson, P., Kinmonth, A. L. (2007). Designing and evaluating complex
6
7 interventions to improve health care. *British Medical Journal*, 334, 455-459. doi:
8
9 10.1136/bmj.39108.379965.BE
10
11
12 Carnell, S., Edwards, C., Croker, H., Boniface, D., & Wardle, J. (2005). Parental perceptions
13
14 of overweight in 3–5 y olds. *International Journal of Obesity*, 29, 353–5.
15
16 doi:10.1038/sj.ijo.0802889
17
18
19 Cote, M., Byczkowski, T., Kotagal, U., Kirk, S., Zeller, M., Daniels, S. (2004). Service
20
21 quality and attrition: an examination of a pediatric obesity program. *International Journal*
22
23 *for Quality in Health Care*, 16, 165–173. doi:10.1093/intqhc/mzh015
24
25
26 Department of Health (2011a). Healthy Lives, Healthy People: A call to action on obesity in
27
28 England. London: HM Government.
29
30 Department of Health (2011b). Health Profile 2011 Knowsley. Crown Copyright 2011.
31
32 Downloaded from:
33
34 www.apho.org.uk/default.aspx?QN=HP_METADATA&AreaID=50307
35
36
37 Dietz, W. H., & Robinson, T. N. (2005). Overweight children and adolescence. *The New*
38
39 *England Journal of Medicine*, 352, 2100-2009. doi 10.1056/NEJMcp043052
40
41
42 Ellaway, A., Kirk, A., McIntyre, S., & Mutrie, N. (2007). Nowhere to play? The relationship
43
44 between the location of outdoor play areas and deprivation in Glasgow. *Health & Place*,
45
46 13, 557-570. doi dx.doi.org/10.1016/j.healthplace.2006.03.005
47
48
49 Etelson, D., Brand, D. A., Patrick, P.A., & Shirali A. (2003). Childhood obesity: do parents
50
51 recognize this health risk? *Obesity Research*, 11, 1362–8. doi 10.1038/oby.2003.184
52
53
54 Haisman, N., Matyka, K. A., & Stanton, A. (2005). Lessons from an unsuccessful local
55
56 attempt to tackle childhood overweight and obesity. *Archives of Disease in Childhood*, 90,
57
58 984-985. doi:10.1136/adc.2005.073338
59
60

- 1
2
3 Jain, A., Sherman, S.N., Chamberlin, L.A., Carter, Y., Powers, S. W. & Whitaker, R.
4
5 C. (2001). Why don't low-income mothers worry about their preschoolers being
6
7 overweight? *Pediatrics*, 107, 1138–46. doi: 10.1542/peds.107.5.1138
8
9
10 Jeffery, A.N., Voss, L.D., Metcalf, B.S., Alba, S., & Wilkin, T. J. (2005). Parents' awareness
11
12 of overweight in themselves and their children: cross sectional study within a cohort
13
14 (EarlyBird 21). *British Medical Journal*, 330, 23–24. doi: 10.1136/bmj.38315.451539.F7
15
16
17 Langton, P. (2011). Public Health Annual Report, Statistics Compendium, 2010. Retrieved
18
19 from:
20
21 [www.knowsley.nhs.uk/assets/uploaded/documents/25329_PH%20Stats%20Compendium](http://www.knowsley.nhs.uk/assets/uploaded/documents/25329_PH%20Stats%20Compendium%20Complete%20v4.pdf)
22
23 [%20Complete%20v4.pdf](http://www.knowsley.nhs.uk/assets/uploaded/documents/25329_PH%20Stats%20Compendium%20Complete%20v4.pdf)
24
25
26 Ludwig, D. S. (2007). Childhood obesity: The shape of things to come. *New England*
27
28 *Journal of Medicine*, 357, 2325–2327. doi: 10.1056/NEJMp0706538
29
30
31 Luttikhuis, H.O., Baur, L., Jansen, H., Shrewsbury, V.A., O'Malley, C., Stolk, R.P.,
32
33 Summerbell, C.D. (2009). Interventions for treating obesity in children (Review). *The*
34
35 *Cochrane Library* 2009, Issue 1. The Cochrane Collaboration. Published by JohnWiley &
36
37 Sons, Ltd. doi: 10.1002/14651858.CD001872.pub2
38
39
40 National Institute for Health and Clinical Excellence. (2006). *Obesity guidance on the*
41
42 *prevention, identification, assessment and management of overweight and obesity in adults*
43
44 *and children*. Clinical Guideline, 43. London: NICE.
45
46
47 National institute for Health and Clinical Excellence (2007). *How to change practice.*
48
49 *Understand, identify and overcome barriers to change*. London, NICE.
50
51
52 NHS Knowsley (2008). *Childhood Obesity Service Development* (unpublished reports 1-3).
53
54
55 NHS, Northwest England.
56
57
58 Oyserman, D. (2007). Social identity and self-regulation. In Kruglanski, A., & Higgins, T.
59
60 (Eds.), *Handbook of social psychology*, (2nd ed., pp. 432–453). New York: Guilford Press

- 1
2
3 Oyserman, D. (2009a). Identity-based motivation: Implications for action-readiness,
4 procedural-readiness, and consumer behavior. *Journal of Consumer Psychology, 19*, 250-
5 260. doi 10.1016/j.jcps.2009.05.008
6
7
8
9
10 Oyserman, D. (2009b). Identity-based motivation and consumer behavior. *Journal of*
11 *Consumer Psychology, 19*, 276-279. doi 10.1016/j.jcps.2009.06.001
12
13
14 Oyserman, D., & Destin, M. (2010). Identity-Based Motivation: Implications for
15 Intervention. *The Counseling Psychologist 38*, 1001. doi 10.1177/0011000010374775
16
17
18 Puhl, R.M., & Heuer, C. A. (2009) The Stigma of Obesity: A Review and Update.
19 *Obesity, 17* 5, 941–964. doi 10.1038/oby.2008.636
20
21
22
23 Roberts, E. A. (2007). Pediatric non-alcoholic fatty liver disease (NAFLD): A “growing”
24 problem. *Journal of Hepatology, 46*, 1133-1142. doi:10.1016/j.jhep.2007.03.003
25
26
27 Rudolf, M. C. J. (2004). The obese child. *Archives of Disease in Childhood, 89*, 57-62.
28 doi:10.1136/adc.2004.059824
29
30
31
32 Smith, J. A., & Osborn, M. (2008). Interpretative Phenomenological Analysis. In J. A. Smith
33 (2008). *Qualitative Psychology, a practical guide to research methods*, 2nd ed (53-80).
34 London: Sage Publications Ltd
35
36
37
38 Smith, J.A., Flower, P., Larkin, M (2009). Interpretative Phenomenological Analysis: Theory,
39 Method and Research Sage Publications Ltd; Reprint edition
40
41
42
43 Smith, B. S., Grunseit, A., Hardy, L. L., King, L., Wolfenden, L., & Milat, A. (2010).
44 Parental influences on child physical activity and screen viewing time: a population based
45 study *BMC Public Health, 10*:593 doi 10.1186/1471-2458-10-593
46
47
48
49
50
51 Smolak, L., Levine, M. D., Schermer, F. (1999) Parental input and weight concerns among
52 elementary school children. *International Journal of Eating disorders vol. 25* (3) 263-271.
53 doi 10.1002/(SICI)1098-108X(199904)25:3<263::AID-EAT3>3.0.CO;2-V
54
55
56
57
58
59
60

1
2
3 The NHS Information centre (2011). National Child Measurement Programme: England,
4
5 2010/11 school year. The NHS Information Centre, Lifestyles Statistics Retrieved
6
7 http://www.ic.nhs.uk/webfiles/publications/003_Health_Lifestyles/ncmp%202010-
8
9 [11/NCMP_2010_11_Report.pdf](http://www.ic.nhs.uk/webfiles/publications/003_Health_Lifestyles/ncmp%202010-11/NCMP_2010_11_Report.pdf)

10
11
12 Trost, S. G., Kerr, L.M., Ward, D. S., Pate, R. R. (2001) Physical activity and determinants of
13
14 physical activity in obese and non-obese children. *International Journal of Obesity*, 25,
15
16 822-829. www.nature.com/ijo/journal/v25/n6/full/0801621a.html

17
18
19 White, A., O'Brien, B., Houlihan, T., Darker, C., O'Shea, B. (2012). Childhood obesity:
20
21 parents fail to recognise, general practitioners fail to act. *Irish Medical Journal*, 105, 10-3.
22
23 <http://www.imj.ie/ViewArticleDetails.aspx?ArticleID=7893>

24
25
26 Zeller, M., Kirk, S., Clayton, R., Khoury, P., Grieme, J., Santangeol, M. and Daniels, S.
27
28 (2004) Predictors of Attrition from a Pediatric Weight-management Program. *The Journal*
29
30 *of Pediatrics* 144, 466-70. doi:10.1016/j.jpeds.2003.12.031
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: NHS childhood obesity service outline; and procedures

Procedures from clinical team: 1) Referral to service received, indicates child as obese. 2) Letter sent to families to attend a clinical assessment. 3) Families attend a clinical assessment to assess physical health, aetiology of obesity, lifestyle and suitability for the obesity intervention. If they met inclusion criteria families were provided with a brief overview of the programme and a start date was agreed.

Programme inclusion criteria: Clinically obese children, aged 5-15years (BMI >98th percentile), to attend with their family (specifically at least one parent/guardian). Assessed to rule out co-morbidities or medical reasons for the onset of obesity (if a medical reason identified, alternative treatment offered as appropriate). Able to participate in group environments.

Programme: The structure of treatment was designed from clinical guidelines (NICE, 2006) and previous pilot research (unpublished, NHS Knowsley, 2008) into programme development. The service was provided in a community venue after school hours, if required the service would make payment for taxis to and from the programme. The programme was delivered using diverse, interactive and “fun” methods and provided advice and resources in reading-age/literacy appropriate material (for both adults and children).

It was a 1 year programme, split into three stages:

Stage 1: Intense 12 x weekly, 2-hour community sessions. Promoted lifestyle change through physical fitness, psychological support and healthy eating principles. **Stage 2:** Bi-monthly individual follow-up sessions at community clinic locations. **Stage 3:** After 1 year, the service supported families by developing a long-term practical action plan and signposting to other ‘universal’ based programmes.

Table 1: Summary of Similarities and Differences across Attendees and Non-Attendees

	Attendees	Non-attendees
Theme 1: Perceptions of childhood obesity	Did not recognise obesity in their child.	
	Childhood obesity is considered a 'norm' within local society.	
	Believed that the cause of obesity was linked to medical issues (genetics/ medication/ illness)	
	Understood the link between obesity and preventable ill-health.	Used the medical assessment outcome as an indicator that obesity wasn't a serious medical condition.
	Did not experience negative influences from significant others or, had the ability to deal with/avoid negative influences from others.	Influenced by social network (extended family/friends) negative attitudes and comments about obesity
Theme 2: Perceptions of the programme	Anxieties of parents' own ability to engage in the programme/ make personal changes.	
	Parents recognising the need for self-support, new knowledge and to change own behaviour (diet and exercise).	Parents 'embarrassed' about perceived inability to participate, alongside their belief of already being knowledgeable.
	Perception that the programme has a positive effect on the child's self-esteem, psychological wellbeing and social relationships.	Perception that the programme has a negative effect on the child's self-esteem, psychological wellbeing and social relationships.
Theme 3: Practical barriers and	Personal safety travelling to and from the venues.	
	Making time (feeling guilty when missing appointments).	No time

overcoming hurdles to attending	Recognition that transport to the venue can act as a barrier, although typically managed to overcome this hurdle (car being usual transportation).	Transport cited as a barrier to attending. Particularly if requesting travel across the borough to attend.
Theme 4: Availability and	Referenced the lack of local, safe and appropriate (fitness/skill level) activities for their children to do. Families felt that the local area did not enable children to be active.	
suitability of local facilities	The obesity programme 'filled the gap' and supported the children to develop new skills. Subsequently they hoped to have confidence to try local activities (which are targeted towards sporty children).	Reported that once the programme ended, there would still remain to be no activities that the children could do/access locally (or in a safe environment).
	The programme offered an opportunity for safe places to play.	