

# THE EXTENDED RELATIVITY THEORY IN CLIFFORD SPACES

C. Castro<sup>a</sup> and M. Pavšič<sup>b</sup>

January 5, 2004

<sup>a</sup>Center for Theoretical Studies of Physical Systems, Clark Atlanta University, Atlanta

<sup>b</sup>Jožef Stefan Institute, Jamova 39, SI-1000 Ljubljana, Slovenia;

Email: matej.pavsic@ijs.si

## Abstract

A brief review of some of the most important features of the Extended Relativity theory in Clifford-spaces ( $C$ -spaces) is presented whose "point" coordinates are noncommuting Clifford-valued quantities and which incorporate the lines, areas, volumes,.... degrees of freedom associated with the collective particle, string, membrane,... dynamics of  $p$ -loops (closed  $p$ -branes) living in target  $D$ -dimensional spacetime backgrounds.  $C$ -space Relativity naturally incorporates the ideas of an invariant length (Planck scale), maximal acceleration, noncommuting coordinates, supersymmetry, holography, higher derivative gravity with torsion and variable dimensions/signatures that allows to study the dynamics of all (closed)  $p$ -branes, for all values of  $p$ , on a unified footing. It resolves the ordering ambiguities in QFT and the problem of time in Cosmology. A discussion of the maximal-acceleration Relativity principle in phase-spaces follows along with the study of the invariance group of symmetry transformations in phase-space that allows to show why Planck areas are invariant under acceleration-boosts transformations and which seems to suggest that a maximal-string tension principle may be operating in Nature. We continue by pointing out how the relativity of signatures of the underlying  $n$ -dimensional spacetime results from taking different  $n$ -dimensional slices through  $C$ -space. The conformal group emerges as a natural subgroup of the Clifford group and Relativity in  $C$ -spaces involves natural scale changes in the sizes of physical objects without the introduction of forces nor Weyl's gauge field of dilations. We finalize by constructing the generalization of Maxwell theory of Electrodynamics of point charges to a theory in  $C$ -spaces that involves extended charges coupled to antisymmetric tensor fields of arbitrary rank. In the concluding remarks we outline briefly the current promising research programs and their plausible connections with  $C$ -space Relativity.

# 1 Introduction

In recent years it was argued that the underlying fundamental physical principle behind string theory, not unlike the principle of equivalence and general covariance in Einstein's general relativity, might well be related to the existence of an invariant minimal length scale (Planck scale) attainable in nature [8]. A theory involving spacetime *resolutions* was developed long ago by Nottale [23] where the Planck scale was postulated as the minimum observer independent invariant resolution [23] in Nature. Since "points" cannot be observed physically with an ultimate resolution, it is reasonable to postulate that they are smeared out into fuzzy balls. In refs.[8] it was assumed that those balls have the Planck radius and arbitrary dimension. For this reason it was argued in refs.[8] that one should construct a theory which includes all dimensions (and signatures) on the equal footing. In [8] this Extended Scale Relativity principle was applied to the quantum mechanics of  $p$ -branes which led to the construction of Clifford-space ( $C$ -space) where all  $p$ -branes were taken to be on the same footing, in the sense that the transformations in  $C$ -space reshuffled a string history for a five-brane history, a membrane history for a string history, for example.

Clifford algebras contained the appropriate algebraic-geometric features to implement this principle of polydimensional transformations [14]–[17]. In [14]–[16] it was proposed that every physical quantity is in fact a *polyvector*, that is, a Clifford number or a Clifford aggregate. Also, spinors are the members of left or right minimal ideals of Clifford algebra, which may provide the framework for a deeper understanding of supersymmetries, i.e., the transformations relating bosons and fermions. The Fock-Stueckelberg theory of a relativistic particle can be embedded in the Clifford algebra of spacetime [15, 16]. Many important aspects of Clifford algebra are described in [1],[6], [7], [3], [15, 16, 17], [5], [48]. It is our belief that this may lead to the proper formulation of string and M theory.

A geometric approach to the physics of the Standard Model in terms of Clifford algebras was advanced by [4]. It was realized in [43] that the  $Cl(8)$  Clifford algebra contains the 4 fundamental nontrivial representations of  $Spin(8)$  that accommodate the chiral fermions and gauge bosons of the Standard model and which also includes gravitons via the McDowell-Mansouri-Chamseddine-West formulation of gravity, which permits to construct locally, in  $D = 8$ , a geometric Lagrangian for the Standard Model plus Gravity. Furthermore, discrete Clifford-algebraic methods based on hyperdiamond-lattices have been instrumental in constructing  $E_8$  lattices and deriving the values of the force-strengths (coupling constants) and masses of the Standard model with remarkable precision by [43]. These results have recently been corroborated by [46] for Electromagnetism, and by [47], where all the Standard model parameters were obtained from first principles, despite the contrary orthodox belief that it is senseless to "derive" the values of the fundamental constants in Nature from first principles, from pure thought alone; i.e. one must invoke the Cosmological anthropic principle to explain why the constants of Nature have the values they have.

Using these methods the bosonic  $p$ -brane propagator, in the quenched mini superspace approximation, was constructed in [18, 19]; the logarithmic corrections to the black hole entropy based on the geometry of Clifford space (in short  $C$ -space) were obtained in [21];

The modified nonlinear de Broglie dispersion relations, the corresponding minimal-length stringy [11] and p-brane uncertainty relations also admitted a  $C$ -space interpretation [10], [19]. A generalization of Maxwell theory of electromagnetism in  $C$ -spaces comprised of extended charges coupled to antisymmetric tensor fields of arbitrary rank was attained recently in [75]. The resolution of the ordering ambiguities of QFT in curved spaces was resolved by using polyvectors, or Clifford-algebra valued objects [26]. One of the most remarkable features of the Extended Relativity in  $C$ -spaces is that a higher derivative Gravity with Torsion in ordinary spacetime follows naturally from the analog of the Einstein-Hilbert action in *curved*  $C$ -space [20].

In this new physical theory the arena for physics is no longer the ordinary spacetime, but a more general manifold of Clifford algebra valued objects, noncommuting polyvectors. Such a manifold has been called a pan-dimensional continuum [14] or  $C$ -space [8]. The latter describes on a unified basis the objects of various dimensionality: not only points, but also closed lines, surfaces, volumes,..., called 0-loops (points), 1-loops (closed strings) 2-loops (closed membranes), 3-loops, etc.. It is a sort of a *dimension* category, where the role of functorial maps is played by  $C$ -space transformations which reshuffles a  $p$ -brane history for a  $p'$ -brane history or a mixture of all of them, for example. The above geometric objects may be considered as corresponding to the well-known physical objects, namely closed  $p$ -branes. Technically those transformations in  $C$ -space that reshuffle objects of different dimensions are generalizations of the ordinary Lorentz transformations to  $C$ -space.

$C$ -space Relativity involves a generalization of Lorentz invariance (and not a deformation of such symmetry) involving superpositions of p-branes (p-loops) of all possible dimensions. The Planck scale is introduced as a natural parameter that allows us to bridge extended objects of different dimensionalities. Like the speed of light was need in Einstein Relativity to fuse space and time together in the Minkowski spacetime interval. Another important point is that the Conformal Group of four-dimensional spacetime is a consequence of the Clifford algebra in *four-dimensions* [25] and it emphasizes the fact why the natural dilations/contractions of objects in  $C$ -space is *not* the same physical phenomenon than what occurs in Weyl's geometry which requires introducing, by hand, a gauge field of dilations. Objects move dilationally, in the absence of forces, for a different physical reasoning than in Weyl's geometry: they move dilationally because of inertia. This was discussed long ago in refs.[27, 28].

This review is organized as follows: Section **2** is dedicated to extending ordinary Special Relativity theory, from Minkowski spacetime to  $C$ -spaces, where the introduction of the invariant Planck scale is required to bridge objects,  $p$ -branes, of different dimensionality.

The generalized dynamics of particles, fields and branes in  $C$ -space is studied in section **3**. This formalism allows us to construct for the first time, to our knowledge, a *unified* action which comprises the dynamics of *all* p-branes in  $C$ -spaces, for all values of  $p$ , in one single footing (see also [15]). In particular, the polyparticle dynamics in  $C$ -space, when reduced to 4-dimensional spacetime leads to the Stuckelberg formalism and the solution to the problem of time in Cosmology [15].

In section **4** we begin by discussing the geometric Clifford calculus that allows us to reproduce all the standard results in differential and projective geometry [41]. The resolution of the ordering ambiguities of QFT in curved spaces follows next when we

review how it can be resolved by using polyvectors, or Clifford-algebra valued objects [26]. Afterwards we construct the Generalized Gravitational Theories in Curved C-spaces, in particular it is shown how Higher derivative Gravity with Torsion in ordinary spacetime follows naturally from the Geometry of C-space [20].

In section 5 we discuss the Quantization program in C-spaces, and write the C-space Klein-Gordon and Dirac equations [15]. The corresponding bosonic/fermionic  $p$ -brane loop-wave equations were studied by [12], [13] without employing Clifford algebra and the concept of C-space.

In section 6 we review the Maximal-Acceleration Relativity in Phase-Spaces [119], starting with the construction of the submaximally-accelerated particle action of [53] using Clifford algebras in phase-spaces; the  $U(1, 3)$  invariance transformations [74] associated with an 8-dimensional phase space, and show why the minimal Planck-Scale areas are invariant under pure acceleration boosts which suggests that there could be a principle of maximal-tension (maximal acceleration) operating in string theory [68].

In section 7 we discuss the important point that the notion of spacetime signature is relative to a chosen  $n$ -dimensional subspace of  $2^n$ -dimensional Clifford space. Different subspaces  $V_n$ —different sections through C-space—have in general different signature [15] We show afterwards how the Conformal algebra of spacetime emerges from the Clifford algebra [25] and emphasize the physical *differences* between our model and the one based on Weyl geometry. At the end we show how Clifford algebraic methods permits one to generalize Maxwell theory of Electrodynamics (associated with ordinary point-charges) to a generalized Maxwell theory in Clifford spaces involving *extended* charges and  $p$ -forms of arbitrary rank [75]

In the concluding remarks, we briefly discuss the possible avenues of future research in the construction of QFT in C-spaces, Quantum Gravity, Noncommutative Geometry, and other lines of current promising research in the literature.

## 2 Extending Relativity from Minkowski Spacetime to C-space

We embark into the construction of the extended relativity theory in C-spaces by a natural generalization of the notion of a spacetime interval in Minkowski space to C-space [8, 14, 16, 15, 17]:

$$dX^2 = d\sigma^2 + dx_\mu dx^\mu + dx_{\mu\nu} dx^{\mu\nu} + \dots \quad (1)$$

where  $\mu_1 < \mu_2 < \dots$ . The Clifford valued polyvector:<sup>1</sup>

$$X = X^M E_M = \sigma \underline{1} + x^\mu \gamma_\mu + x^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + \dots x^{\mu_1 \mu_2 \dots \mu_D} \gamma_{\mu_1} \wedge \gamma_{\mu_2} \dots \wedge \gamma_{\mu_D}. \quad (2)$$

denotes the position of a point in a manifold, called Clifford space or C-space. The series of terms in (2) terminates at a *finite* grade depending on the dimension  $D$ . A Clifford

---

<sup>1</sup>If we do not restrict indices according to  $\mu_1 < \mu_2 < \mu_3 < \dots$ , then the factors  $1/2!$ ,  $1/3!$ , respectively, have to be included in front of every term in the expansion (1).

algebra  $Cl(r, q)$  with  $r + q = D$  has  $2^D$  basis elements. For simplicity, the gammas  $\gamma^\mu$  correspond to a Clifford algebra associated with a flat spacetime:

$$\frac{1}{2}\{\gamma^\mu, \gamma^\nu\} = \eta^{\mu\nu}. \quad (3)$$

but in general one could extend this formulation to curved spacetimes with metric  $g^{\mu\nu}$  (see section 4).

The connection to strings and p-branes can be seen as follows. In the case of a closed string (a 1-loop) embedded in a target flat spacetime background of  $D$ -dimensions, one represents the projections of the closed string (1-loop) onto the embedding spacetime coordinate-planes by the variables  $x^{\mu\nu}$ . These variables represent the respective *areas* enclosed by the projections of the closed string (1-loop) onto the corresponding embedding spacetime planes. Similarly, one can embed a closed membrane (a 2-loop) onto a  $D$ -dim flat spacetime, where the projections given by the antisymmetric variables  $x^{\mu\nu\rho}$  represent the corresponding *volumes* enclosed by the projections of the 2-loop along the hyperplanes of the flat target spacetime background.

This procedure can be carried to all closed p-branes ( $p$ -loops) where the values of  $p$  are  $p = 0, 1, 2, 3, \dots$ . The  $p = 0$  value represents the center of mass and the coordinates  $x^{\mu\nu}, x^{\mu\nu\rho}, \dots$  have been *coined* in the string-brane literature [24]. as the *holographic* areas, volumes, ... projections of the nested family of  $p$ -loops (closed  $p$ -branes) onto the embedding spacetime coordinate planes/hyperplanes. In ref.[17] they were interpreted as the generalized centre of mass coordinates of an extended object. Extended objects were thus modeled in  $C$ -space.

The scalar coordinate  $\sigma$  entering a polyvector  $X$  is a measure associated with the  $p$ -brane's world manifold  $V_{p+1}$  (e.g., the string's 2-dimensional worldsheet  $V_2$ ): it is proportional to the  $(p + 1)$ -dimensional area/volume of  $V_{p+1}$ . In other words,  $\sigma$  is proportional to the areal-time parameter of the Eguchi-Schild formulation of string dynamics [118, 37, 24].

We see in this generalized scheme the objects as observed in spacetime (which is a section through  $C$ -space) need not be infinitely extended along time-like directions. They need not be infinitely long world lines, world tubes. They can be finite world lines, world tubes. The  $\sigma$  coordinate measures how long are world lines, world tubes. During evolution they can become longer and longer or shorter and shorter.

If we take the differential  $dX$  of  $X$  and compute the scalar product among two polyvectors  $\langle dX^\dagger dX \rangle_0 \equiv dX^\dagger * dX \equiv |dX|^2$  we obtain the  $C$ -space extension of the particles proper time in Minkowski space. The symbol  $X^\dagger$  denotes the *reversion* operation and involves reversing the order of all the basis  $\gamma^\mu$  elements in the expansion of  $X$ . It is the analog of the transpose (Hermitian) conjugation. The  $C$ -space proper time associated with a polyparticle motion is then the expression (1) which can be written more explicitly as:

$$\begin{aligned} |dX|^2 &= G_{MN} dX^M dX^N = dS^2 \\ &= d\sigma^2 + L^{-2} dx_\mu dx^\mu + L^{-4} dx_{\mu\nu} dx^{\mu\nu} + \dots + L^{-2D} dx_{\mu_1 \dots \mu_D} dx^{\mu_1 \dots \mu_D} \end{aligned} \quad (4)$$

where  $G_{MN} = E_M^\dagger * E_N$  is the  $C$ -space metric.

Here we have introduced the Planck scale  $L$  since a length parameter is needed in order to tie objects of different dimensionality together: 0-loops, 1-loops,...,  $p$ -loops. Einstein introduced the speed of light as a universal absolute invariant in order to “unite” space with time (to match units) in the Minkowski space interval:

$$ds^2 = c^2 dt^2 + dx_i dx^i.$$

A similar unification is needed here to “unite” objects of different dimensions, such as  $x^\mu$ ,  $x^{\mu\nu}$ , etc... The Planck scale then emerges as another universal invariant in constructing an extended relativity theory in  $C$ -spaces [8].

Since the  $D$ -dimensional Planck scale is given explicitly in terms of the Newton constant:  $L_D = (G_N)^{1/(D-2)}$ , in natural units of  $\hbar = c = 1$ , one can see that when  $D = \infty$  the value of  $L_D$  is then  $L_\infty = G^0 = 1$  (assuming a finite value of  $G$ ). Hence in  $D = \infty$  the Planck scale has the natural value of unity. However, if one wishes to avoid any serious algebraic divergence problems in the series of terms appearing in the expansion of the analog of proper time in  $C$ -spaces, in the extreme case when  $D = \infty$ , from now on we shall focus solely on a *finite* value of  $D$ . In this fashion we avoid any serious algebraic convergence problems. We shall not be concerned in this work with the representations of Clifford algebras in different dimensions and with different signatures.

The line element  $dS$  as defined in (4) is *dimensionless*. Alternatively, one can define [8, 9] the line element whose dimension is that of the  $D$ -volume so that:

$$d\Sigma^2 = L^{2D} d\sigma^2 + L^{2D-2} dx_\mu dx^\mu + L^{2D-4} dx_{\mu\nu} dx^{\mu\nu} + \dots + dx_{\mu_1 \dots \mu_D} dx^{\mu_1 \dots \mu_D} \quad (5)$$

Let us use the relation

$$\gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_D} = \gamma \epsilon_{\mu_1 \dots \mu_D} \quad (6)$$

and write the *volume element* as

$$dx^{\mu_1 \dots \mu_D} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_D} \equiv \gamma d\tilde{\sigma} \quad (7)$$

where

$$d\tilde{\sigma} \equiv dx^{\mu_1 \dots \mu_D} \epsilon_{\mu_1 \dots \mu_D} \quad (8)$$

In all expressions we assume the ordering prescription  $\mu_1 < \mu_2 < \dots < \mu_r$ ,  $r = 1, 2, \dots, D$ . The line element can then be written in the form

$$d\Sigma^2 = L^{2D} d\sigma^2 + L^{2D-2} dx_\mu dx^\mu + L^{2D-4} dx_{\mu\nu} dx^{\mu\nu} + \dots + |\gamma|^2 d\tilde{\sigma}^2 \quad (9)$$

where

$$|\gamma|^2 \equiv \gamma^\dagger * \gamma \quad (10)$$

Here  $\gamma$  is the pseudoscalar basis element and can be written as  $\gamma_0 \wedge \gamma_1 \wedge \dots \wedge \gamma_{D-1}$ . In flat spacetime  $M_D$  we have that  $|\gamma|^2 = +1$  or  $-1$ , depending on dimension and signature. In  $M_4$  with signature  $(+ - - -)$  we have  $\gamma^\dagger * \gamma = \gamma^\dagger \gamma = \gamma^2 = -1$  ( $\gamma \equiv \gamma_5 = \gamma_0 \gamma_1 \gamma_2 \gamma_3$ ), whilst in  $M_5$  with signature  $(+ - - - -)$  it is  $\gamma^\dagger \gamma = 1$ .

The analog of Lorentz transformations in  $C$ -spaces which transform a polyvector  $X$  into another poly-vector  $X'$  is given by

$$X' = R X R^{-1} \quad (11)$$

with

$$R = e^{\theta^A E_A} = \exp [(\theta I + \theta^\mu \gamma_\mu + \theta^{\mu_1 \mu_2} \gamma_{\mu_1} \wedge \gamma_{\mu_2} \dots)]. \quad (12)$$

and

$$R^{-1} = e^{-\theta^A E_A} = \exp [-(\theta I + \theta^\nu \gamma_\nu + \theta^{\nu_1 \nu_2} \gamma_{\nu_1} \wedge \gamma_{\nu_2} \dots)]. \quad (13)$$

where the theta parameters in (12)(13) are the components of the Clifford-value parameter  $\Theta = \theta^M E_M$ :

$$\theta; \theta^\mu; \theta^{\mu\nu}; \dots \quad (14)$$

they are the C-space version of the Lorentz rotations/boosts parameters.

Since a Clifford algebra admits a matrix representation, one can write the norm of a poly-vectors in terms of the trace operation as:  $\|X\|^2 = \text{Trace } X^2$  Hence under C-space Lorentz transformation the norms of poly-vectors behave like follows:

$$\text{Trace } X'^2 = \text{Trace } [RX^2R^{-1}] = \text{Trace } [RR^{-1}X^2] = \text{Trace } X^2. \quad (15)$$

These norms are invariant under C-space Lorentz transformations due to the cyclic property of the trace operation and  $RR^{-1} = 1$ .

The above transformations are *active transformations* since the transformed Clifford number  $X'$  (polyvector) is different from the “original” Clifford number  $X$ . Considering the transformations of components we have

$$X' = X'^M E_M = L^M_N X^N E_M \quad (16)$$

If we compare (16) with (11) we find

$$L^M_N E_N = R E_N R^{-1} \quad (17)$$

from which it follows that

$$L^M_N = \langle E^M R E_N R^{-1} \rangle_0 \equiv E^M * (R E_N R^{-1}) = E^M * E'_N. \quad (18)$$

where we have labelled  $E'_N$  as new basis element since in the active interpretation one may perform either a change of the polyvector components or a change of the basis elements. The  $\langle \ \ \rangle_0$  means the scalar part of the expression and “\*” the scalar product. Eq(18) has been obtained after multiplying (17) from the left by  $E^J$ , taking into account that  $\langle E^J E_N \rangle_0 \equiv E^J * E_N = \delta^J_N$ , and renaming the index  $J$  into  $M$ .

### 3 Generalized Dynamics of Particles, Fields and Branes in C-space

An immediate application of this theory is that one may consider “strings” and “branes” in C-spaces as a unifying description of *all* branes of different dimensionality. As we have already indicated, since spinors are in left/right ideals of a Clifford algebra, a supersymmetry is then naturally incorporated into this approach as well. In particular, one can have world manifold and target space supersymmetry *simultaneously* [15]. We hope that the C-space “strings” and “branes” may lead us towards discovering the physical foundations of string and M-theory. For other alternatives to supersymmetry see the work by [50]. In particular,  $Z_3$  generalizations of supersymmetry based on ternary algebras and Clifford algebras have been proposed by Kerner [120] in what has been called Hypersymmetry.

### 3.1 The Polyparticle Dynamics in C-space

We will now review the theory [15, 17] in which an extended object is modeled by the components  $\sigma, x^\mu, x^{\mu\nu}, \dots$  of the Clifford valued polyvector (2). By assumption the extended objects, as observed from Minkowski spacetime, can in general be localized not only along space-like, but also along time-like directions [15, 17]. In particular, they can be “instantonic”  $p$ -loops with either space-like or time-like orientation. Or they may be long, but finite, tube-like objects. The theory that we consider here goes beyond the ordinary relativity in Minkowski spacetime, therefore such localized objects in Minkowski spacetime pose no problems. They are postulated to satisfy the dynamical principle which is formulated in  $C$ -space. All conservation laws hold in  $C$ -space where we have infinitely long world “lines” or Clifford lines. In Minkowski spacetime  $M_4$  –which is a subspace of  $C$ -space– we observe the intersections of Clifford lines with  $M_4$ . And those intersections appear as localized extended objects,  $p$ -loops, described above.

Let the motion of such an extended object be determined by the action principle

$$I = \kappa \int d\tau (\dot{X}^\dagger * \dot{X})^{1/2} = \kappa \int d\tau (\dot{X}^A \dot{X}_A)^{1/2} \quad (19)$$

where  $\kappa$  is a constant, playing the role of “mass” in  $C$ -space, and  $\tau$  is an arbitrary parameter. The  $C$ -space velocities  $\dot{X}^A = dX^A/d\tau = (\dot{\sigma}, \dot{x}^\mu, \dot{x}^{\mu\nu}, \dots)$  are also called “holographic” velocities.

The equation of motion resulting from (19) is

$$\frac{d}{d\tau} \left( \frac{\dot{X}^A}{\sqrt{\dot{X}^B \dot{X}_B}} \right) = 0 \quad (20)$$

Taking  $\dot{X}^B \dot{X}_B = \text{constant} \neq 0$  we have that  $\ddot{X}^A = 0$ , so that  $x^A(\tau)$  is a straight worldline in  $C$ -space. The components  $x^A$  then change linearly with the parameter  $\tau$ . This means that the extended object position  $x^\mu$ , effective area  $x^{\mu\nu}$ , 3-volume  $x^{\mu\nu\alpha}$ , 4-volume  $x^{\mu\nu\alpha\beta}$ , etc., they all change with time. That is, such object experiences a sort of generalized dilational motion [17].

We shall now review the procedure exposed in ref. [17] according to which in such a generalized dynamics an object may be accelerated to faster than light speeds as viewed from a 4-dimensional Minkowski space, which is a subspace of  $C$ -space. For a different explanation of superluminal propagation based on the modified nonlinear de Broglie dispersion relations see [68].

The canonical momentum belonging to the action (19) is

$$P_A = \frac{\kappa \dot{X}_A}{(\dot{X}^B \dot{X}_B)^{1/2}} \quad (21)$$

When the denominator in eq.(21) is zero the momentum becomes infinite. We shall now calculate the speed at which this happens. This will be the *maximum speed* that an object accelerating in  $C$ -space can reach. Although an initially slow object cannot accelerate beyond that speed limit, this does not automatically exclude the possibility



that fast objects traveling at a speed above that limit may exist. Such objects are  $C$ -space analog of tachyons [31, 32]. All the well known objections against tachyons should be reconsidered for the case of  $C$ -space before we could say for sure that  $C$ -space tachyons do not exist as freely propagating objects. We will leave aside this interesting possibility, and assume as a working hypothesis that there is no tachyons in  $C$ -space.

Vanishing of  $\dot{X}^B \dot{X}_B$  is equivalent to vanishing of the  $C$ -space line element

$$dX^A dX_A = d\sigma^2 + \left(\frac{dx^0}{L}\right)^2 - \left(\frac{dx^1}{L}\right)^2 + \left(\frac{dx^{12}}{L^2}\right)^2 - \left(\frac{dx^{123}}{L^3}\right)^2 - \left(\frac{dx^{0123}}{L^4}\right)^2 + \dots = 0 \quad (22)$$

where by “...” we mean the terms with the remaining components such as  $x^2$ ,  $x^{01}$ ,  $x^{23}$ , ...,  $x^{012}$ , etc.. The  $C$ -space line element is associated with a particular *choice* of  $C$ -space metric, namely  $G_{MN} = E_M^\dagger * E_N$ . If the basis  $E_M$ ,  $M = 1, 2, \dots, 2^D$  is generated by the flat space  $\gamma^\mu$  satisfying (3), then the  $C$ -space has the diagonal metric of eq. (22) with +, - signa. In general this is not necessarily so and the  $C$ -space metric is a more complicated expression. We take now dimension of spacetime being 4, so that  $x^{0123}$  is the highest grade coordinate. In eq. (22) we introduce a length parameter  $L$ . This is necessary, since  $x^0 = ct$  has dimension of length,  $x^{12}$  of length square,  $x^{123}$  of length to the third power, and  $x^{0123}$  of length to the fourth power. It is natural to assume that  $L$  is the *Planck length*, that is  $L = 1.6 \times 10^{-35}m$ .

Let us assume that the coordinate time  $t = x^0/c$  is the parameter with respect to which we define the speed  $V$  in  $C$ -space.

So we have

$$V^2 = - \left(L \frac{d\sigma}{dt}\right)^2 + \left(\frac{dx^1}{dt}\right)^2 - \left(\frac{1}{L} \frac{dx^{12}}{dt}\right)^2 + \left(\frac{1}{L^2} \frac{dx^{123}}{dt}\right)^2 + \left(\frac{1}{L^3} \frac{dx^{0123}}{dt}\right)^2 - \dots \quad (23)$$

From eqs. (22),(23) we find that the maximum speed is The maximum speed is given by

$$V^2 = c^2 \quad (24)$$

First, we see that the maximum speed squared  $V^2$  contains not only the components of the 1-vector velocity  $dx^1/dt$ , as it is the case in the ordinary relativity, but also the multivector components such as  $dx^{12}/dt$ ,  $dx^{123}/dt$ , etc..

The following special cases when only certain components of the velocity in  $C$ -space are different from zero, are of particular interest:

(i) Maximum 1-vector speed

$$\frac{dx^1}{dt} = c = 3.0 \times 10^8 m/s$$

(ii) Maximum 3-vector speed

$$\begin{aligned} \frac{dx^{123}}{dt} &= L^2 c = 7.7 \times 10^{-62} m^3/s \\ \frac{d\sqrt[3]{x^{123}}}{dt} &= 4.3 \times 10^{-21} m/s \quad (\text{diameter speed}) \end{aligned}$$

(iii) Maximum 4-vector speed

$$\begin{aligned}\frac{dx^{0123}}{dt} &= L^3 c = 1.2 \times 10^{-96} \text{m}^4/\text{s} \\ \frac{d\sqrt[4]{x^{0123}}}{dt} &= 1.05 \times 10^{-24} \text{m}/\text{s} \quad (\text{diameter speed})\end{aligned}$$

Above we have also calculated the corresponding diameter speeds for the illustration of how fast the object expands or contracts.

We see that the maximum multivector speeds are very small. The diameters of objects change very slowly. Therefore we normally do not observe the dilatational motion.

Because of the positive sign in front of the  $\sigma$  and  $x^{12}$ ,  $x^{012}$ , etc., terms in the quadratic form (22) there are no limits to corresponding 0-vector, 2-vector and 3-vector speeds. But if we calculate, for instance, the energy necessary to excite 2-vector motion we find that it is very high. Or equivalently, to the relatively modest energies (available at the surface of the Earth), the corresponding 2-vector speed is very small. This can be seen by calculating the energy

$$p^0 = \frac{\kappa c^2}{\sqrt{1 - \frac{V^2}{c^2}}} \quad (25)$$

(a) for the case of pure 1-vector motion by taking  $V = dx^1/dt$ , and

(b) for the case of pure 2-vector motion by taking  $V = dx^{12}/(Ldt)$ .

By equating the energies belonging to the cases (a) and (b) we have

$$p^0 = \frac{\kappa c^2}{\sqrt{1 - \left(\frac{1}{c} \frac{dx^1}{dt}\right)^2}} = \frac{\kappa c^2}{\sqrt{1 - \left(\frac{1}{Lc} \frac{dx^{12}}{dt}\right)^2}} \quad (26)$$

which gives

$$\frac{1}{c} \frac{dx^1}{dt} = \frac{1}{Lc} \frac{dx^{12}}{dt} = \sqrt{1 - \left(\frac{\kappa c^2}{p_0}\right)^2} \quad (27)$$

Thus to the energy of an object moving translationally at  $dx^1/dt = 1$  m/s, there corresponds the 2-vector speed  $dx^{12}/dt = L dx^1/dt = 1.6 \times 10^{-35}$  m<sup>2</sup>/s (diameter speed  $4 \times 10^{-18}$  m/s). This would be a typical 2-vector speed of a macroscopic object. For a microscopic object, such as the electron, which can be accelerated close to the speed of light, the corresponding 2-vector speed could be of the order of  $10^{-26}$  m<sup>2</sup>/s (diameter speed  $10^{-13}$  m/s). In the examples above we have provided rough estimations of possible 2-vector speeds. Exact calculations should treat concrete situations of collisions of two or more objects, assume that not only 1-vector, but also 2-vector, 3-vector and 4-vector motions are possible, and take into account the conservation of the polyvector momentum  $P_A$ .

*Maximum 1-vector speed, i.e., the usual speed, can exceed the speed of light when the holographic components such as  $d\sigma/dt$ ,  $dx^{12}/dt$ ,  $dx^{012}/dt$ , etc., are different from zero [17]. This can be immediately verified from eqs. (22),(23). The speed of light is no*

longer such a strict barrier as it appears in the ordinary theory of relativity in  $M_4$ . In  $C$ -space a particle has extra degrees of freedom, besides the translational degrees of freedom. The scalar,  $\sigma$ , the bivector,  $x^{12}$  (in general,  $x^{rs}$ ,  $r, s = 1, 2, 3$ ) and the threevector,  $x^{012}$  (in general,  $x^{0rs}$ ,  $r, s = 1, 2, 3$ ), contributions to the  $C$ -space quadratic form (22) have positive sign, which is just opposite to the contributions of other components, such as  $x^r$ ,  $x^{0r}$ ,  $x^{rst}$ ,  $x^{\mu\nu\rho\sigma}$ . Because the quadratic form has both  $+$  and  $-$  signs, the absolute value of the 3-velocity  $dx^r/dx^0$  can be greater than  $c$ .

In the usual theory of relativity faster than light motion implies causality violation, since one can arrange situations such that tachyons are sent into the past. The “past” means that the coordinate  $t' \equiv x'^0$  at which the second tachyons arrives is smaller than the coordinate  $t \equiv x^0$  at which the first tachyons is emitted, i.e.,  $t' < t$ ).

In the extended relativity, i.e., in the relativity in  $C$ -space, the dynamics refers to a larger space. Minkowski space is just a subspace of  $C$ -space. In  $C$ -space we have the *constrained action* (19), whilst in Minkowski space we have a reduced, *unconstrained action*, which was shown in ref.[16, 15, 17] to be equivalent to the well known Stueckelberg action [33, 34]. In Stueckelberg theory the parameter of evolution is not the coordinate  $t \equiv x^0$ . The evolution parameter is an extra parameter  $\tau$  which is invariant under Lorentz transformations in  $M_4$ . For the extra parameter we can take one of the extra coordinates that enter our action (19). One natural choice for the evolution parameter is the scalar component  $\sigma$  of a polyvector  $X$  (eq.(4)). Another choice is the pseudoscalar component  $x^{0123} \equiv \tilde{\sigma}$ . If a particle arrives at an earlier coordinate time  $t \equiv x^0$  this does not mean a violation of causality, provided that it arrives at later evolution time  $\tau$ . For a more complete discussion see refs. [29, 15, 17].

If nature indeed obeys the dynamics in Clifford space, then a particle, as observed from the 4-dimensional Minkowski space, can be accelerated beyond the speed of light [17], provided that its extra degrees of freedom  $x^{\mu\nu}$ ,  $x^{\mu\nu\alpha}$ , ..., are changing simultaneously with the ordinary position  $x^\mu$ . In  $C$ -space we thus retain all the nice features of relativity, but in the subspace  $M_4$  we have an unconstrained Stueckelberg theory in which tachyons are not paradoxical and are consistent with the quantum field theory as well [15].

**An alternative procedure** In ref. [9] an alternative factorization of the  $C$ -space line element has been undertaken. Starting from the line element  $d\Sigma$  of eq. (5), instead of factoring out the  $(dx^0)^2$  element, one may factor out the  $(d\Omega)^2 \equiv L^{2D} d\sigma^2$  element, giving rise to the generalized “holographic” velocities measured w.r.t the  $\Omega$  parameter, for example the areal-time parameter in the Eguchi-Schild formulation of string dynamics [118], [37], [24], instead of the  $x^0$  parameter (coordinate clock). One then obtains

$$d\Sigma^2 = d\Omega^2 \left[ 1 + L^{2D-2} \frac{dx_\mu}{d\Omega} \frac{dx^\mu}{d\Omega} + L^{2D-4} \frac{dx_{\mu\nu}}{d\Omega} \frac{dx^{\mu\nu}}{d\Omega} + \dots + |\gamma|^2 \left( \frac{d\tilde{\sigma}}{d\Omega} \right)^2 \right] \quad (28)$$

The idea of ref. [9] was to restrict the line element (28) to the non tachyonic values which imposes an upper limit on the holographic velocities. The motivation was to find a lower bound of length scale. This upper holographic-velocity bound does not necessarily translate into a lower bound on the values of lengths, areas, volumes...without the introduction of quantum mechanical considerations. One possibility could be that the upper limiting

speed of light and the upper bound of the momentum  $m_p c$  of a Planck-mass elementary particle (the so-called *Planckton* in the literature) generalizes now to an upper-bound in the  $p$ -loop holographic velocities and the  $p$ -loop holographic momenta associated with elementary closed  $p$ -branes whose tensions are given by powers of the Planck mass. And the latter upper bounds on the holographic  $p$ -loop momenta implies a lower-bound on the holographic areas, volumes,..., resulting from the string/brane uncertainty relations [11], [10],[19]. Thus, Quantum Mechanics is required to implement the postulated principle of minimal lengths, areas, volumes...and which cannot be derived from the classical geometry alone. The emergence of minimal Planck areas occurs also in the Loop Quantum Gravity program [103] where the expectation values of the Area operator are given by multiples of Planck area.

### 3.2 A Unified Theory of all p-Branes in C-Spaces

The generalization to C-spaces of string and p-brane actions as embeddings of world-manifolds onto target spacetime backgrounds involves the embeddings of polyvector-valued world-manifolds (of dimensions  $2^d$ ) onto polyvector-valued target spaces (of dimensions  $2^D$ ), given by the Clifford-valued maps  $X = X(\Sigma)$  (see [15]). These are maps from the Clifford-valued world-manifold, parametrized by the polyvector-valued variables  $\Sigma$ , onto the Clifford-valued target space parametrized by the polyvector-valued coordinates  $X$ . Physically one envisions these maps as taking an  $n$ -dimensional simplicial cell ( $n$ -loop) of the world-manifold onto an  $m$ -dimensional simplicial cell ( $m$ -loop) of the target C-space manifold ; i.e. maps from  $n$ -dim objects onto  $m$ -dim objects generalizing the old maps of taking points onto points. One is basically dealing with a dimension-category of objects. The size of the simplicial cells ( $p$ -loops), upon quantization of a generalized harmonic oscillator, for example, are given by multiples of the Planck scale, in area, volume, hypervolume units or Clifford-bits.

In compact multi-index notation  $X = X^M \Gamma_M$  one denotes for each one of the components of the target space polyvector  $X$ :

$$X^M \equiv X^{\mu_1 \mu_2 \dots \mu_r}, \quad \mu_1 < \mu_2 < \dots < \mu_r. \quad (29)$$

and for the world-manifold polyvector  $\Sigma = \Sigma^A E_A$ :

$$\Sigma^A \equiv \xi^{a_1 a_2 \dots a_s}, \quad a_1 < a_2 < \dots < a_s. \quad (30)$$

where  $\Gamma_M = (\underline{1}, \gamma_\mu, \gamma_{\mu\nu}, \dots)$  and  $E_A = (\underline{1}, e_a, e_{ab}, \dots)$  form the basis of the target manifold and world manifold Clifford algebra, respectively. It is very important to order the indices within each multi-index  $M$  and  $A$  as shown above. The above Clifford-valued coordinates  $X^M, \Sigma^A$  correspond to antisymmetric tensors of ranks  $r, s$  in the target spacetime background and in the world-manifold, respectively.

There are many different ways to construct C-space brane actions which are on-shell equivalent to the analogs of the Dirac-Nambu-Goto action for extended objects and that are given by the world-volume spanned by the branes in their motion through the target spacetime background.

One of these actions is the Polyakov-Howe-Tucker action:

$$I = \frac{T}{2} \int [D\Sigma] \sqrt{|H|} [H^{AB} \partial_A X^M \partial_B X^N G_{MN} + (2 - 2^d)]. \quad (31)$$

with the  $2^d$ -dim world-manifold measure:

$$[D\Sigma] = (d\xi)(d\xi^a)(d\xi^{a_1 a_2})(d\xi^{a_1 a_2 a_3})\dots \quad (32)$$

Upon the algebraic elimination of the auxiliary world-manifold metric  $H^{AB}$  from the action (31), via the equations of motion, yields for its on-shell solution the pullback of the target C-space metric onto the C-space world-manifold:

$$H_{AB}(\text{on-shell}) = G_{AB} = \partial_A X^M \partial_B X^N G_{MN} \quad (33)$$

upon inserting back the on-shell solutions (33) into (31) gives the Dirac-Nambu-Goto action for the C-space branes directly in terms of the C-space determinant, or measure, of the induced C-space world-manifold metric  $G_{AB}$ , as a result of the embedding:

$$I = T \int [D\Sigma] \sqrt{\text{Det}(\partial_A X^M \partial_B X^N G_{MN})}. \quad (34)$$

However in C-space, the Polyakov-Howe-Tucker action admits an even further generalization that is comprised of two terms  $S_1 + S_2$ . The first term is:

$$S_1 = \int [D\Sigma] |E| E^A E^B \partial_A X^M \partial_B X^N \Gamma_M \Gamma_N. \quad (35)$$

Notice that this is a generalized action which is written in terms of the C-space coordinates  $X^M(\Sigma)$  and the C-space analog of the target-spacetime vielbein/frame one-forms  $e^m = e^m{}_\mu dx^\mu$  given by the  $\Gamma^M$  variables. The auxiliary world-manifold vielbein variables  $e^a$ , are given now by the Clifford-valued frame  $E^A$  variables.

In the conventional Polyakov-Howe-Tucker action, the auxiliary world-manifold metric  $h^{ab}$  associated with the standard p-brane actions is given by the usual scalar product of the frame vectors  $e^a \cdot e^b = e^a{}_\mu e^b{}_\nu g^{\mu\nu} = h^{ab}$ . Hence, the C-space world-manifold metric  $H^{AB}$  appearing in (33) is given by scalar product  $\langle (E^A)^\dagger E^B \rangle_0 = H^{AB}$ , where  $(E^A)^\dagger$  denotes the reversal operation of  $E^A$  which requires reversing the ordering of the vectors present in the Clifford aggregate  $E^A$ .

Notice, however, that the form of the action (35) is far more general than the action in (31). In particular, the  $S_1$  itself can be decomposed further into two additional pieces by rewriting the Clifford product of two basis elements into a symmetric plus an antisymmetric piece, respectively:

$$E^A E^B = \frac{1}{2} \{E^A, E^B\} + \frac{1}{2} [E^A, E^B]. \quad (36)$$

$$\Gamma_M \Gamma_N = \frac{1}{2} \{\Gamma_M, \Gamma_N\} + \frac{1}{2} [\Gamma_M, \Gamma_N]. \quad (37)$$

In this fashion, the  $S_1$  component has *two* kinds of terms. The first term containing the symmetric combination is just the analog of the standard non-linear sigma model

action, and the second term is a Wess-Zumino-like term, containing the antisymmetric combination [15]. To extract the non-linear sigma model part of the generalized action above, we may simply take the scalar product of the vielbein-variables as follows:

$$(S_1)_{sigma} = \frac{T}{2} \int [D\Sigma] |E| \langle (E^A \partial_A X^M \Gamma_M)^\dagger (E^B \partial_B X^N \Gamma_N) \rangle_0. \quad (38)$$

where once again we have made use of the reversal operation (the analog of the hermitian adjoint) before contracting multi-indices. In this fashion we recover again the Clifford-scalar valued action given by [15].

Actions like the ones presented here in terms of derivatives with respect to quantities with multi-indices can be mapped to actions involving *higher* derivatives, in the same fashion that the C-space scalar curvature, the analog of the Einstein-Hilbert action, could be recast as a higher derivative gravity with torsion (reviewed in sec. 4). Higher derivatives actions are also related to theories of Higher spin fields [109] and  $W$ -geometry,  $W$ -algebras [108], [114]. For the role of Clifford algebras to higher spin theories see [51].

The  $S_2$  (scalar) component of the C-space brane action is the usual cosmological constant term given by the C-space determinant  $|E| = \det(H^{AB})$  based on the scalar part of the geometric product  $\langle (E^A)^\dagger E^B \rangle_0 = H^{AB}$

$$S_2 = \frac{T}{2} \int [D\Sigma] |E| (2 - 2^d) \quad (39)$$

where the C-space determinant  $|E| = \sqrt{|\det(H^{AB})|}$  of the  $2^d \times 2^d$  generalized world-manifold metric  $H^{AB}$  is given by:

$$\det(H^{AB}) = \frac{1}{(2^d)!} \epsilon_{A_1 A_2 \dots A_{2^d}} \epsilon_{B_1 B_2 \dots B_{2^d}} H^{A_1 B_1} H^{A_2 B_2} \dots H^{A_{2^d} B_{2^d}}. \quad (40)$$

The  $\epsilon_{A_1 A_2 \dots A_{2^d}}$  is the totally antisymmetric tensor density in C-space.

There are many different forms of  $p$ -brane actions, with and without a cosmological constant [115], and based on a new integration measure by recurring to auxiliary scalar fields [107], that one could have used to construct their C-space generalizations. Since all of them are on-shell equivalent to the Dirac-Nambu-Goto  $p$ -brane actions, we decided to focus solely on those actions having the Polyakov-Howe-Tucker form.

## 4 Generalized Gravitational Theories in Curved C-spaces: Higher Derivative Gravity and Torsion from the Geometry of C-Space

### 4.1 Ordinary space

#### 4.1.1 Clifford algebra based geometric calculus in curved space(time)

Clifford algebra is a very useful tool for description of geometry, especially of curved space  $V_n$ . Let us first review how it works in curved space(time). Later we will discuss a generalization to curved Clifford space [20].

Let  $\gamma_\mu$ ,  $\mu = 1, 2, \dots, n$  be a set of  $n$  independent basis vectors which satisfy the Clifford algebra relation

$$\gamma_\mu \cdot \gamma_\nu \equiv \frac{1}{2}(\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu) = g_{\mu\nu} \quad (41)$$

where  $g_{\mu\nu}$  is the metric of  $V_n$ . In curved space  $\gamma_\mu$  and  $g_{\mu\nu}$  cannot be constant but necessarily depend on position  $x^\mu$ . A arbitrary vector is a linear superposition [1]

$$a = a^\mu \gamma_\mu \quad (42)$$

where the components  $a^\mu$  are *scalars* from the geometric point of view, whilst  $\gamma_\mu$  are *vectors*.

Besides the basis  $\gamma_\mu$  we can introduce the dual basis  $\gamma^\mu$  satisfying

$$\gamma^\mu \cdot \gamma^\nu \equiv \frac{1}{2}(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu) = g^{\mu\nu} \quad (43)$$

where  $g^{\mu\nu}$  is the covariant metric tensor such that  $g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$ ,  $\gamma^\mu \gamma_\nu + \gamma_\nu \gamma^\mu = 2\delta^\mu_\nu$  and  $\gamma^\mu = g^{\mu\nu} \gamma_\nu$ .

Following ref.[1] (see also [15]) we consider the *vector derivative* or *gradient* is defined according to

$$\partial \equiv \gamma^\mu \partial_\mu \quad (44)$$

where  $\partial_\mu$  is an operator, that we call *geometric derivative*, whose action depends on the quantity it acts on [26].

Applying the vector derivative  $\partial$  on a *scalar*  $\phi$  we have

$$\partial\phi = \gamma^\mu \partial_\mu \phi \quad (45)$$

where  $\partial_\mu \phi \equiv (\partial/\partial x^\mu)\phi$  coincides with the partial derivative of  $\phi$ .

But if we apply it on a *vector*  $a$  we have

$$\partial a = \gamma^\mu \partial_\mu (a^\nu \gamma_\nu) = \gamma^\mu (\partial_\mu a^\nu \gamma_\nu + a^\nu \partial_\mu \gamma_\nu) \quad (46)$$

In general  $\gamma_\nu$  is not constant; it satisfies the relation [1, 15]

$$\partial_\mu \gamma_\nu = \Gamma_{\mu\nu}^\alpha \gamma_\alpha \quad (47)$$

where  $\Gamma_{\mu\nu}^\alpha$  is the *connection*. Similarly, for  $\gamma^\nu = g^{\nu\alpha} \gamma_\alpha$  we have

$$\partial_\mu \gamma^\nu = -\Gamma_{\mu\alpha}^\nu \gamma^\alpha \quad (48)$$

The *non commuting* operator  $\partial_\mu$  so defined determines the *parallel transport* of a basis vector  $\gamma^\nu$ . It should be distinguished from the ordinary—*commuting*—partial derivative  $\gamma^\nu_{,\mu}$  usually used in the literature on the Dirac equation in curved spacetime. Instead of the symbol  $\partial_\mu$  Hestenes uses  $\square_\mu$ , whilst Wheeler et. al. [36] use  $\nabla_\mu$  and call it “covariant derivative”. We find it convenient to keep the symbol  $\partial_\mu$  for components of the geometric operator  $\partial = \gamma^\mu \partial_\mu$ . When acting on a scalar field the geometric derivative  $\partial_\mu$  happens to be commuting and thus behaves as the ordinary partial derivative.

Using (47), eq.(46) becomes

$$\partial a = \gamma^\mu \gamma_\nu (\partial_\mu a^\nu + \Gamma_{\mu\alpha}^\nu a^\alpha) \equiv \gamma^\mu \gamma_\nu D_\mu a^\nu = \gamma^\mu \gamma^\nu D_\mu a_\nu \quad (49)$$

where  $D_\mu$  is the covariant derivative.

Decomposing the Clifford product  $\gamma^\mu \gamma^\nu$  into its symmetric and antisymmetric part [1]

$$\gamma^\mu \gamma^\nu = \gamma^\mu \cdot \gamma^\nu + \gamma^\mu \wedge \gamma^\nu \quad (50)$$

where

$$\gamma^\mu \cdot \gamma^\nu \equiv \frac{1}{2}(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu) = g^{\mu\nu} \quad (51)$$

is the *inner product* and

$$\gamma^\mu \wedge \gamma^\nu \equiv \frac{1}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) \quad (52)$$

the *outer product*, we can write eq.(49) as

$$\partial a = g^{\mu\nu} D_\mu a_\nu + \gamma^\mu \wedge \gamma^\nu D_\mu a_\nu = D_\mu a^\mu + \frac{1}{2} \gamma^\mu \wedge \gamma^\nu (D_\mu a_\nu - D_\nu a_\mu) \quad (53)$$

Without employing the expansion in terms of  $\gamma_\mu$  we have simply

$$\partial a = \partial \cdot a + \partial \wedge a \quad (54)$$

Acting twice on a vector by the operator  $\partial$  we have<sup>2</sup>

$$\begin{aligned} \partial \partial a &= \gamma^\mu \partial_\mu (\gamma^\nu \partial_\nu) (a^\alpha \gamma_\alpha) = \gamma^\mu \gamma^\nu \gamma_\alpha D_\mu D_\nu a^\alpha \\ &= \gamma_\alpha D_\mu D^\mu a^\alpha + \frac{1}{2} (\gamma^\mu \wedge \gamma^\nu) \gamma_\alpha [D_\mu, D_\nu] a^\alpha \\ &= \gamma_\alpha D_\mu D^\mu a^\alpha + \gamma^\mu (R_{\mu\rho} a^\rho + K_{\mu\alpha}{}^\rho D_\rho a^\alpha) \\ &\quad + \frac{1}{2} (\gamma^\mu \wedge \gamma^\nu \wedge \gamma_\alpha) (R_{\mu\nu\rho}{}^\alpha a^\rho + K_{\mu\nu}{}^\rho D_\rho a^\alpha) \end{aligned} \quad (55)$$

We have used

$$[D_\mu, D_\nu] a^\alpha = R_{\mu\nu\rho}{}^\alpha a^\rho + K_{\mu\nu}{}^\rho D_\rho a^\alpha \quad (56)$$

where

$$K_{\mu\nu}{}^\rho = \Gamma_{\mu\nu}^\rho - \Gamma_{\nu\mu}^\rho \quad (57)$$

is *torsion* and  $R_{\mu\nu\rho}{}^\alpha$  the *curvature tensor*. Using eq.(47) we find

$$[\partial_\alpha, \partial_\beta] \gamma_\mu = R_{\alpha\beta\mu}{}^\nu \gamma_\nu \quad (58)$$

from which we have

$$R_{\alpha\beta\mu}{}^\nu = ([[\partial_\alpha, \partial_\beta] \gamma_\mu] \cdot \gamma^\nu) \quad (59)$$

Thus in general the commutator of partial derivatives acting on a vector does not give zero, but is given by the curvature tensor.

---

<sup>2</sup>We use  $(a \wedge b)c = (a \wedge b) \cdot c + a \wedge b \wedge c$  [1] and  $(a \wedge b) \cdot c = (b \cdot c)a - (a \cdot c)b$ .



In general, for an  $r$ -vector  $A = a^{\alpha_1 \dots \alpha_r} \gamma_{\alpha_1} \gamma_{\alpha_2} \dots \gamma_{\alpha_r}$  we have

$$\begin{aligned} \partial \partial \dots \partial A &= (\gamma^{\mu_1} \partial_{\mu_1}) (\gamma^{\mu_2} \partial_{\mu_2}) \dots (\gamma^{\mu_k} \partial_{\mu_k}) (a^{\alpha_1 \dots \alpha_r} \gamma_{\alpha_1} \gamma_{\alpha_2} \dots \gamma_{\alpha_r}) \\ &= \gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_k} \gamma_{\alpha_1} \gamma_{\alpha_2} \dots \gamma_{\alpha_r} D_{\mu_1} D_{\mu_2} \dots D_{\mu_k} a^{\alpha_1 \dots \alpha_r} \end{aligned} \quad (60)$$

At this point it is convenient to define the covariant derivative as acting not only on the scalar components such as  $a^\alpha$ ,  $a^{\alpha\beta}$ , etc., but also on vectors, bivectors, etc.. In particular, when acting on a basis vector the covariant derivative —because of (47)— gives

$$D_\mu \gamma_\nu = \partial_\mu \gamma_\nu - \Gamma_{\mu\nu}^\rho \gamma_\rho = 0 \quad (61)$$

Therefore, for instance, the partial derivative of a vector is equal to the covariant derivative:

$$\partial_\mu a = \partial_\mu (a^\nu \gamma_\nu) = (D_\mu a^\nu) \gamma_\nu = D_\mu a \quad (62)$$

In general, for an arbitrary  $r$ -vector  $A$  we have

$$\partial_\mu A = \partial_\mu (a^{\alpha_1 \dots \alpha_r} \gamma_{\alpha_1} \dots \gamma_{\alpha_r}) = (D_\mu a^{\alpha_1 \dots \alpha_r}) \gamma_{\alpha_1} \dots \gamma_{\alpha_r} = D_\mu A \quad (63)$$

For the commutator of *partial derivatives* acting on a vector  $a = a^\rho \gamma_\rho$  we have

$$[\partial_\mu, \partial_\nu] a = a^\rho [\partial_\mu, \partial_\nu] \gamma_\rho = R_{\mu\nu\rho}{}^\sigma a^\rho \gamma_\sigma \quad (64)$$

whilst for the commutator of *covariant derivatives* we find

$$[D_\mu, D_\nu] a = \gamma_\rho [D_\mu, D_\nu] a^\rho = (R_{\mu\nu\rho}{}^\sigma a^\rho + K_{\mu\nu}{}^\alpha D_\alpha a^\sigma) \gamma_\sigma \quad (65)$$

#### 4.1.2 Clifford algebra based geometric calculus and resolution of the ordering ambiguity for the product of momentum operators

Clifford algebra is a very useful tool for description of geometry of curved space. Moreover, as shown in ref.[26] it provides a resolution of the long standing problem of the ordering ambiguity of quantum mechanics in curved space. Namely, eq.(44) for the vector derivative suggests that the momentum operator is given by

$$p = -i \partial = -i \gamma^\mu \partial_\mu \quad (66)$$

One can consider three distinct models:

- (i) *The non relativistic particle* moving in  $n$ dimensional curved space. Then,  $\mu = 1, 2, \dots, n$ , and signature is  $(+ + + + \dots)$ .
- (ii) *The relativistic particle* in curved spacetime, described by the *Schild action* [37]. Then,  $\mu = 0, 1, 2, \dots, n - 1$  and signature is  $(+ - - - \dots)$ .
- (iii) *The Stueckelberg unconstrained particle*. [33, 34, 35, 29].

In all three cases the classical action has the form

$$I[X^\mu] = \frac{1}{2\Lambda} \int d\tau g_{\mu\nu}(x) \dot{X}^\mu \dot{X}^\nu \quad (67)$$

and the corresponding Hamiltonian is

$$H = \frac{\Lambda}{2} g^{\mu\nu}(x) p_\mu p_\nu = \frac{\Lambda}{2} p^2 \quad (68)$$

If, upon quantization we take for the momentum operator  $p_\mu = -i \partial_\mu$ , then the ambiguity arises of how to write the quantum Hamilton operator. The problem occurs because the expressions  $g^{\mu\nu} p_\mu p_\nu$ ,  $p_\mu g^{\mu\nu} p_\nu$  and  $p_\mu p_\nu g^{\mu\nu}$  are not equivalent.

But, if we rewrite  $H$  as

$$H = \frac{\Lambda}{2} p^2 \quad (69)$$

where  $p = \gamma^\mu p_\mu$  is the *momentum vector* which upon quantization becomes the momentum vector operator (66), we find that there is no ambiguity in writing the square  $p^2$ . When acting with  $H$  on a *scalar* wave function  $\phi$  we obtain the unambiguous expression

$$H\phi = \frac{\Lambda}{2} p^2 \phi = \frac{\Lambda}{2} (-i)^2 (\gamma^\mu \partial_\mu) (\gamma^\nu \partial_\nu) \phi = -\frac{\Lambda}{2} D_\mu D^\mu \phi \quad (70)$$

in which there is no curvature term  $R$ . We expect that a term with  $R$  will arise upon acting with  $H$  on a *spinor* field  $\psi$ .

## 4.2 $C$ -space

Let us now consider  $C$ -space and review the procedure of ref. [20]. . A basis in  $C$ -space is given by

$$E_A = \{\gamma_\mu, \gamma_\mu \wedge \gamma_\nu, \gamma_\mu \wedge \gamma_\nu \wedge \gamma_\rho, \dots\} \quad (71)$$

where in an  $r$ -vector  $\gamma_{\mu_1} \wedge \gamma_{\mu_2} \wedge \dots \wedge \gamma_{\mu_r}$  we take the indices so that  $\mu_1 < \mu_2 < \dots < \mu_r$ . An element of  $C$ -space is a Clifford number, called also *Polyvector* or *Clifford aggregate* which we now write in the form

$$X = X^A E_A = s \underline{1} + x^\mu \gamma_\mu + x^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + \dots \quad (72)$$

A  $C$ -space is parametrized not only by 1-vector coordinates  $x^\mu$  but also by the 2-vector coordinates  $x^{\mu\nu}$ , 3-vector coordinates  $x^{\mu\nu\alpha}$ , etc., called also *holographic coordinates*, since they describe the holographic projections of 1-loops, 2-loops, 3-loops, etc., onto the coordinate planes. By  $p$ -loop we mean a closed  $p$ -brane; in particular, a 1-loop is closed string.

In order to avoid using the powers of the Planck scale length parameter  $L$  in the expansion of the polyvector  $X$  we use the dilatationally invariant units [15] in which  $L$  is set to 1. The dilation invariant physics was discussed from a different perspective also in refs. [23, 21].

In a flat  $C$ -space the basis vectors  $E^A$  are constants. In a curved  $C$ -space this is no longer true. Each  $E_A$  is a function of the  $C$ -space coordinates

$$X^A = \{s, x^\mu, x^{\mu\nu}, \dots\} \quad (73)$$

which include scalar, vector, bivector, ...,  $r$ -vector, ..., coordinates.

Now we define the connection  $\tilde{\Gamma}_{AB}^C$  in  $C$ -space according to

$$\partial_A E_B = \tilde{\Gamma}_{AB}^C E_C \quad (74)$$

where  $\partial_A \equiv \partial/\partial X^A$  is the partial derivative in  $C$ -space. This definition is analogous to the one in ordinary space. Let us therefore define the  $C$ -space curvature as

$$\mathcal{R}_{ABC}^D = ([\partial_A, \partial_B]E_C) * E^D \quad (75)$$

which is a straightforward generalization of the relation (59). The 'star' means the *scalar product* between two polyvectors  $A$  and  $B$ , defined as

$$A * B = \langle AB \rangle_S \quad (76)$$

where 'S' means 'the scalar part' of the geometric product  $AB$ .

In the following we shall explore the above relation for curvature and see how it is related to the curvature of the ordinary space. Before doing that we shall demonstrate that the derivative with respect to the bivector coordinate  $x^{\mu\nu}$  is equal to the commutator of the derivatives with respect to the vector coordinates  $x^\mu$ .

Returning now to eq.(74), the differential of a  $C$ -space basis vector is given by

$$dE_A = \frac{\partial E_A}{\partial X^B} dX^B = \Gamma_{AB}^C E_C dX^B \quad (77)$$

In particular, for  $A = \mu$  and  $E_A = \gamma_\mu$  we have

$$\begin{aligned} d\gamma_\mu &= \frac{\partial \gamma_\mu}{\partial X^\nu} dx^\nu + \frac{\partial \gamma_\mu}{\partial x^{\alpha\beta}} dx^{\alpha\beta} + \dots = \tilde{\Gamma}_{\nu\mu}^A E_A dx^\nu + \tilde{\Gamma}_{[\alpha\beta]\mu}^A E_A dx^{\alpha\beta} + \dots \\ &= (\tilde{\Gamma}_{\nu\mu}^\alpha \gamma_\alpha + \tilde{\Gamma}_{\nu\mu}^{[\rho\sigma]} \gamma_\rho \wedge \gamma_\sigma + \dots) dx^\nu \\ &\quad + (\tilde{\Gamma}_{[\alpha\beta]\mu}^\rho \gamma_\rho + \tilde{\Gamma}_{[\alpha\beta]\mu}^{[\rho\sigma]} \gamma_\rho \wedge \gamma_\sigma + \dots) dx^{\alpha\beta} + \dots \end{aligned} \quad (78)$$

We see that the differential  $d\gamma_\mu$  is in general a polyvector, i.e., a Clifford aggregate. In eq.(78) we have used

$$\frac{\partial \gamma_\mu}{\partial x^\nu} = \tilde{\Gamma}_{\nu\mu}^\alpha \gamma_\alpha + \tilde{\Gamma}_{\nu\mu}^{[\rho\sigma]} \gamma_\rho \wedge \gamma_\sigma + \dots \quad (79)$$

$$\frac{\partial \gamma_\mu}{\partial x^{\alpha\beta}} = \tilde{\Gamma}_{[\alpha\beta]\mu}^\rho \gamma_\rho + \tilde{\Gamma}_{[\alpha\beta]\mu}^{[\rho\sigma]} \gamma_\rho \wedge \gamma_\sigma + \dots \quad (80)$$

Let us now consider a *restricted* space in which the derivatives of  $\gamma_\mu$  with respect to  $x^\nu$  and  $x^{\alpha\beta}$  do not contain higher rank multivectors. Then eqs. (79),(80) become

$$\frac{\partial \gamma_\mu}{\partial x^\nu} = \tilde{\Gamma}_{\nu\mu}^\alpha \gamma_\alpha \quad (81)$$

$$\frac{\partial \gamma_\mu}{\partial x^{\alpha\beta}} = \tilde{\Gamma}_{[\alpha\beta]\mu}^\rho \gamma_\rho \quad (82)$$

Further we assume that

- (i) the components  $\tilde{\Gamma}_{\nu\mu}^\alpha$  of the  $C$ -space connection  $\tilde{\Gamma}_{AB}^C$  coincide with the connection  $\Gamma_{\nu\mu}^\alpha$  of an ordinary space.
- (ii) the components  $\tilde{\Gamma}_{[\alpha\beta]\mu}^\rho$  of the  $C$ -space connection coincide with the curvature tensor  $R_{\alpha\beta\mu}{}^\rho$  of an ordinary space.

Hence, eqs.(81),(82) read

$$\frac{\partial\gamma_\mu}{\partial x^\nu} = \Gamma_{\nu\mu}^\alpha \gamma_\alpha \quad (83)$$

$$\frac{\partial\gamma_\mu}{\partial x^{\alpha\beta}} = R_{\alpha\beta\mu}{}^\rho \gamma_\rho \quad (84)$$

and the differential (78) becomes

$$d\gamma_\mu = (\Gamma_{\alpha\mu}^\rho dx^\alpha + \frac{1}{2}R_{\alpha\beta\mu}{}^\rho dx^{\alpha\beta})\gamma_\rho \quad (85)$$

The same relation was obtained by Pezzaglia [14] by using a different method, namely by considering how polyvectors change with position. The above relation demonstrates that a geodesic in  $C$ -space is not a geodesic in ordinary spacetime. Namely, in ordinary spacetime we obtain Papapetrou's equation. This was previously pointed out by Pezzaglia [14].

Although a  $C$ -space connection does not transform like a  $C$ -space tensor, some of its components, i.e., those of eq. (82), may have the transformation properties of a tensor in an ordinary space.

Under a general coordinate transformation in  $C$ -space

$$X^A \rightarrow X'^A = X'^A(X^B) \quad (86)$$

the connection transforms according to<sup>3</sup>

$$\tilde{\Gamma}_{AB}^C = \frac{\partial X'^C}{\partial X^E} \frac{\partial X^J}{\partial X'^A} \frac{\partial X^K}{\partial X'^B} \tilde{\Gamma}_{JK}^E + \frac{\partial X'^C}{\partial X^J} \frac{\partial^2 X^J}{\partial X'^A \partial X'^B} \quad (87)$$

In particular, the components which contain the bivector index  $A = [\alpha\beta]$  transform as

$$\tilde{\Gamma}_{[\alpha\beta]\mu}^{\rho} = \frac{\partial X'^\rho}{\partial X^E} \frac{\partial X^J}{\partial \sigma'^{\alpha\beta}} \frac{\partial X^K}{\partial x'^\mu} \tilde{\Gamma}_{JK}^E + \frac{\partial x'^\rho}{\partial X^J} \frac{\partial^2 X^J}{\partial \sigma'^{\alpha\beta} \partial x'^\mu} \quad (88)$$

Let us now consider a particular class of coordinate transformations in  $C$ -space such that

$$\frac{\partial x'^\rho}{\partial x^{\mu\nu}} = 0, \quad \frac{\partial x'^{\mu\nu}}{\partial x'^\alpha} = 0 \quad (89)$$

---

<sup>3</sup>This can be derived from the relation

$$dE'_A = \frac{\partial E'_A}{\partial X'^B} dX'^B \quad \text{where} \quad E'_A = \frac{\partial X^D}{\partial X'^A} E_D \quad \text{and} \quad dX'^B = \frac{\partial X'^B}{\partial X^C} dX^C$$

Then the second term in eq.(88) vanishes and the transformation becomes

$$\tilde{\Gamma}'^{\rho}_{[\alpha\beta]\mu} = \frac{\partial X'^{\rho}}{\partial x^{\epsilon}} \frac{\partial x^{\rho\sigma}}{\partial \sigma'^{\alpha\beta}} \frac{\partial x^{\gamma}}{\partial x'^{\mu}} \tilde{\Gamma}^{\epsilon}_{[\rho\sigma]\gamma} \quad (90)$$

Now, for the bivector whose components are  $dx^{\alpha\beta}$  we have

$$d\sigma'^{\alpha\beta} \gamma'_{\alpha} \wedge \gamma'_{\beta} = dx^{\alpha\beta} \gamma_{\alpha} \wedge \gamma_{\beta} \quad (91)$$

Taking into account that in our particular case (89)  $\gamma_{\alpha}$  transforms as a basis vector in an ordinary space

$$\gamma'_{\alpha} = \frac{\partial x^{\mu}}{\partial x'^{\alpha}} \gamma_{\mu} \quad (92)$$

we find that (91) and (92) imply

$$d\sigma'^{\alpha\beta} \frac{\partial x^{\mu}}{\partial x'^{\alpha}} \frac{\partial x^{\nu}}{\partial x'^{\beta}} = dx^{\mu\nu} \quad (93)$$

which means that

$$\frac{\partial x^{\mu\nu}}{\partial \sigma'^{\alpha\beta}} = \frac{1}{2} \left( \frac{\partial x^{\mu}}{\partial x'^{\alpha}} \frac{\partial x^{\nu}}{\partial x'^{\beta}} - \frac{\partial x^{\nu}}{\partial x'^{\alpha}} \frac{\partial x^{\mu}}{\partial x'^{\beta}} \right) \equiv \frac{\partial x^{[\mu}}{\partial x'^{\alpha}} \frac{\partial x^{\nu]}{\partial x'^{\beta}} \quad (94)$$

The transformation of the bivector coordinate  $x^{\mu\nu}$  is thus determined by the transformation of the vector coordinates  $x^{\mu}$ . This is so because the basis bivectors are the wedge products of basis vectors  $\gamma_{\mu}$ .

From (90) and (94) we see that  $\tilde{\Gamma}^{\epsilon}_{[\rho\sigma]\gamma}$  transforms like a 4th-rank tensor in an ordinary space.

Comparing eq.(84) with the relation (58) we find

$$\frac{\partial \gamma_{\mu}}{\partial x^{\alpha\beta}} = [\partial_{\alpha}, \partial_{\beta}] \gamma_{\mu} \quad (95)$$

The partial derivative of a basis vector with respect to the bivector coordinates  $x^{\alpha\beta}$  is equal to the commutator of partial derivatives with respect to the vector coordinates  $x^{\alpha}$ .

The above relation (95) holds for the basis vectors  $\gamma_{\mu}$ . For an arbitrary polyvector

$$A = A^A E_A = s + a^{\alpha} \gamma_{\alpha} + a^{\alpha\beta} \gamma_{\alpha} \wedge \gamma_{\beta} + \dots \quad (96)$$

we have

$$\frac{DA}{Dx^{\mu\nu}} = [D_{\mu}, D_{\nu}]A \quad (97)$$

where  $D/Dx^{\mu\nu}$  is the covariant derivative, defined in analogous way as in eqs. (49),(61):

$$\frac{DE^A}{DX^B} = 0, \quad \frac{DA^A}{DX^B} = \frac{\partial A^A}{\partial X^B} + \tilde{\Gamma}^A_{BC} A^C \quad (98)$$

In general, thus, we employ the commutator of the covariant derivatives.

From (96) and (97) we obtain, after using (65), that

$$\begin{aligned} \frac{Ds}{Dx^{\mu\nu}} + \frac{Da^\alpha}{Dx^{\mu\nu}}\gamma_\alpha + \frac{Da^{\alpha\beta}}{Dx^{\mu\nu}}\gamma_\alpha \wedge \gamma_\beta + \dots \\ = [D_\mu, D_\nu]s + [D_\mu, D_\nu]a^\alpha\gamma_\alpha + [D_\mu, D_\nu]a^{\alpha\beta}\gamma_\alpha \wedge \gamma_\beta + \dots \end{aligned} \quad (99)$$

From the latter polyvector equation we obtain

$$\frac{Ds}{Dx^{\mu\nu}} = [D_\mu, D_\nu]s = K_{\mu\nu}{}^\rho \partial_\rho s \quad (100)$$

$$\frac{Da^\alpha}{Dx^{\mu\nu}} = [D_\mu, D_\nu]a^\alpha = R_{\mu\nu\rho}{}^\alpha a^\rho + K_{\mu\nu}{}^\rho D_\rho a^\alpha \quad (101)$$

Using (98) we have that

$$\frac{Ds}{Dx^{\mu\nu}} = \frac{\partial s}{\partial x^{\mu\nu}} \quad (102)$$

and

$$\frac{Da^\alpha}{Dx^{\mu\nu}} = \frac{\partial a^\alpha}{\partial x^{\mu\nu}} + \tilde{\Gamma}_{[\mu\nu]\rho}^\alpha a^\rho = \frac{\partial a^\alpha}{\partial x^{\mu\nu}} + R_{\mu\nu\rho}{}^\alpha a^\rho \quad (103)$$

where, according to (ii),  $\tilde{\Gamma}_{[\mu\nu]\rho}^\alpha$  has been identified with curvature. So we obtain, after inserting (102),(103) into (100),(101) that

- (a) the partial derivative of the coefficients  $s$  and  $a^\alpha$ , which are Clifford scalars<sup>4</sup>, with respect to  $x^{\mu\nu}$  are related to *torsion*:

$$\frac{\partial s}{\partial x^{\mu\nu}} = K_{\mu\nu}{}^\rho \partial_\rho s \quad (104)$$

$$\frac{\partial a^\alpha}{\partial x^{\mu\nu}} = K_{\mu\nu}{}^\rho D_\rho a^\alpha \quad (105)$$

- (b) whilst those of the basis vectors are related to *curvature*:

$$\frac{\partial \gamma_\alpha}{\partial x^{\mu\nu}} = R_{\mu\nu\alpha}{}^\beta \gamma_\beta \quad (106)$$

In other words, the dependence of coefficients  $s$  and  $a^\alpha$  on  $x^{\mu\nu}$  indicates the presence of torsion. On the contrary, when basis vectors  $\gamma_\alpha$  depend on  $x^{\mu\nu}$  this indicates that the corresponding vector space has non vanishing curvature.

---

<sup>4</sup>In the geometric calculus based on Clifford algebra, the coefficients such as  $s$ ,  $a^\alpha$ ,  $a^{\alpha\beta}$ , ..., are called *scalars* (although in tensor calculus they are called scalars, vectors and tensors, respectively), whilst the objects  $\gamma_\alpha$ ,  $\gamma_\alpha \wedge \gamma_\beta$ , ..., are called *vectors*, *bivectors*, etc..

### 4.3 On the relation between the curvature of $C$ -space and the curvature of an ordinary space

Let us now consider the  $C$ -space curvature defined in eq.(75) The indices  $A, B$ , can be of vector, bivector, etc., type. It is instructive to consider a particular example.

$$A = [\mu\nu], B = [\alpha\beta], C = \gamma, D = \delta$$

$$\left( \left[ \frac{\partial}{\partial x^{\mu\nu}}, \frac{\partial}{\partial x^{\alpha\beta}} \right] \gamma_\gamma \right) \cdot \gamma^\delta = \mathcal{R}_{[\mu\nu][\alpha\beta]\gamma}{}^\delta \quad (107)$$

Using (84) we have

$$\frac{\partial}{\partial x^{\mu\nu}} \frac{\partial}{\partial x^{\alpha\beta}} \gamma_\gamma = \frac{\partial}{\partial x^{\mu\nu}} (R_{\alpha\beta\gamma}{}^\rho \gamma_\rho) = R_{\alpha\beta\gamma}{}^\rho R_{\mu\nu\rho}{}^\sigma \gamma_\sigma \quad (108)$$

where we have taken

$$\frac{\partial}{\partial x^{\mu\nu}} R_{\alpha\beta\gamma}{}^\rho = 0 \quad (109)$$

which is true in the case of vanishing torsion (see also an explanation that follows after the next paragraph). Inserting (108) into (107) we find

$$\mathcal{R}_{[\mu\nu][\alpha\beta]\gamma}{}^\delta = R_{\mu\nu\gamma}{}^\rho R_{\alpha\beta\rho}{}^\delta - R_{\alpha\beta\gamma}{}^\rho R_{\mu\nu\rho}{}^\delta \quad (110)$$

which is the product of two usual curvature tensors. We can proceed in analogous way to calculate the other components of  $\mathcal{R}_{ABC}{}^D$  such as  $\mathcal{R}_{[\alpha\beta\gamma\delta][\rho\sigma]}{}^\mu$ ,  $\mathcal{R}_{[\alpha\beta\gamma\delta][\rho\sigma\tau\kappa]}{}^{[\mu\nu]}$ , etc.. These contain higher powers of the curvature in an ordinary space. All this is true in our restricted  $C$ -space given by eqs.(81),(82) and the assumptions (i),(ii) bellow those equations. By releasing those restrictions we would have arrived at an even more involved situation which is beyond the scope of the present paper.

After performing the contractions of (110) and the corresponding higher order relations we obtain the expansion of the form

$$\mathcal{R} = R + \alpha_1 R^2 + \alpha_2 R_{\mu\nu} R^{\mu\nu} + \dots \quad (111)$$

So we have shown that the  $C$ -space curvature can be expressed as the sum of the products of the ordinary spacetime curvature. This bears a resemblance to the string effective action in curved spacetimes given by sums of powers of the curvature tensors based on the quantization of non-linear sigma models [110].

If one sets aside the algebraic convergence problems when working with Clifford algebras in infinite dimensions, one can consider the possibility of studying Quantum Gravity in a very large number of dimensions which has been revisited recently [79] in connection to a perturbative renormalizable quantum theory of gravity in infinite dimensions. Another interesting possibility is that an infinite series expansion of the powers of the scalar curvature could yield the recently proposed modified Lagrangians  $R + 1/R$  of gravity to accomodate the cosmological accelerated expansion of the Universe [123], after a judicious choice of the algebraic coefficients is taken. One may notice also that having a vanishing cosmological constant in  $C$ -space,  $\mathcal{R} = \Lambda = 0$  does not necessarily imply that one has a

vanishing cosmological constant in ordinary spacetime. For example, in the very special case of homogeneous symmetric spacetimes, like spheres and hyperboloids, where all the curvature tensors are proportional to suitable combinations of the metric tensor times the scalar curvature, it is possible to envision that the net combination of the sum of all the powers of the curvature tensors may cancel-out giving an overall zero value  $\mathcal{R} = 0$ . This possibility deserves investigation.

Let us now show that for vanishing torsion the curvature is independent of the bivector coordinates  $x^{\mu\nu}$ , as it was taken in eq.(109). Consider the basic relation

$$\gamma_\mu \cdot \gamma_\nu = g_{\mu\nu} \quad (112)$$

Differentiating with respect to  $x^{\alpha\beta}$  we have

$$\frac{\partial}{\partial x^{\alpha\beta}}(\gamma_\mu \cdot \gamma_\nu) = \frac{\partial \gamma_\mu}{\partial x^{\alpha\beta}} \cdot \gamma_\nu + \gamma_\mu \cdot \frac{\partial \gamma_\nu}{\partial x^{\alpha\beta}} = R_{\alpha\beta\mu\nu} + R_{\alpha\beta\nu\mu} = 0 \quad (113)$$

This implies that

$$\frac{\partial g_{\mu\nu}}{\partial \sigma_{\alpha\beta}} = [\partial_\alpha, \partial_\beta]g_{\mu\nu} = 0 \quad (114)$$

Hence the metric, in this particular case, is independent of the holographic (bivector) coordinates. Since the curvature tensor —when torsion is zero— can be written in terms of the metric tensor and its derivatives, we conclude that not only the metric, but also the curvature is independent of  $x^{\mu\nu}$ . In general, when the metric has a dependence on the holographic coordinates one expects further corrections to eq.(110) that would include torsion.

## 5 On the Quantization in $C$ -spaces

### 5.1 The momentum constraint in $C$ -space

A detailed discussion of the physical properties of all the components of the polymomentum  $P$  in four dimensions and the emergence of the physical mass in Minkowski spacetime has been provided in the book [15]. The polymomentum in  $D = 4$ , canonically conjugate to the position polyvector

$$X = \sigma + x^\mu \gamma_\mu + \gamma^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + \xi^\mu \gamma_5 \gamma_\mu + s \gamma_5 \quad (115)$$

can be written as:

$$P = \mu + p^\mu \gamma_\mu + S^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + \pi^\mu \gamma_5 \gamma_\mu + m \gamma_5. \quad (116)$$

where besides the vector components  $p^\mu$  we have the scalar component  $\mu$ , the 2-vector components  $S^{\mu\nu}$ , that are connected to the spin as shown by [14]; the pseudovector components  $\pi^\mu$  and the pseudoscalar component  $m$ .

The most salient feature of the polyparticle dynamics in  $C$ -spaces [15] is that one can start with a *constrained* action in  $C$ -space and arrive, nevertheless, at an *unconstrained*



Stuckelberg action in Minkowski space (a subspace of C-space) in which  $p_\mu p^\mu$  is a constant of motion. The true constraint in C-space is:

$$P_A P^A = \mu^2 + p_\mu p^\mu - 2S^{\mu\nu} S_{\mu\nu} + \pi_\mu \pi^\mu - m^2 = M^2. \quad (117)$$

where  $M$  is a *fixed* constant, the mass in  $C$ -space. The pseudoscalar component  $m$  is a variable, like  $\mu$ ,  $p_\mu$ ,  $S^{\mu\nu}$ , and  $\pi^\mu$ , which altogether are constrained according to eq.(117). It becomes the physical mass in Minkowski spacetime in the special case when other extra components vanish, i.e., when  $\mu = 0$ ,  $S^{\mu\nu} = 0$  and  $\pi^\mu = 0$ . This justifies using the notation  $m$  for mass. This is basically the distinction between the mass in Minkowski space which is a constant of motion  $p_\mu p^\mu$  and the fixed mass  $M$  in  $C$ -space. The variable  $m$  is canonically conjugate to  $s$  which acquires the role of the Stuckelberg evolution parameter  $s$  that allowed ref.[29, 15] to propose a natural solution of the problem of time in quantum gravity.

A derivation of a charge, mass, and spin relationship of a polyparticle can be obtained from the above polymomentum constraint in C-space if one relates the norm of the axial-momentum component  $\pi^\mu$  of the polymomentum  $P$  to the charge [76]. It agrees exactly with the recent charge-mass-spin relationship obtained by [44] based on the Kerr-Newman black hole metric solutions of the Einstein-Maxwell equations. The naked singularity Kerr-Newman solutions have been interpreted by [45] as Dirac particles. Further investigation is needed to understand better these relationships, in particular, the deep reasons behind the *charge* assignment to the norm of the axial-vector  $\pi^\mu$  component of the polymomentum which suggests that mass has a gravitational, electromagnetic and rotational aspects to it. In a Kaluza-Klein reduction from  $D = 5$  to  $D = 4$  it is well known that the electric charge is related to the  $p_5$  component of the momentum. Hence, charge bears a connection to an internal momentum.

## 5.2 C-space Klein-Gordon and Dirac Wave Equations

The ordinary Klein-Gordon equation can be easily obtained by implementing the on-shell constraint  $p^2 - m^2 = 0$  as an operator constraint on the physical states after replacing  $p_\mu$  for  $-i\partial/\partial x^\mu$  (we use units in which  $\hbar = 1, c = 1$ ):

$$\left( \frac{\partial^2}{\partial x^\mu \partial x_\mu} + m^2 \right) \phi = 0. \quad (118)$$

The C-space generalization follows from the  $P^2 - M^2 = 0$  condition by replacing

$$P_A \rightarrow -i \frac{\partial}{\partial X^A} = -i \left( \frac{\partial}{\partial \sigma}, \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^{\mu\nu}}, \dots \right) \quad (119)$$

$$\left( \frac{\partial^2}{\partial \sigma^2} + \frac{\partial^2}{\partial x^\mu \partial x_\mu} + \frac{\partial^2}{\partial x^{\mu\nu} \partial x_{\mu\nu}} + \dots + M^2 \right) \Phi = 0 \quad (120)$$

where we have set  $L = \hbar = c = 1$  for convenience purposes and the C-space scalar field  $\Phi(\sigma, x^\mu, x^{\mu\nu}, \dots)$  is a polyvector-valued *scalar* function of *all* the C-space variables. This is the Klein-Gordon equation associated with a free scalar polyparticle in C-space.

A wave equation for a generalized C-space harmonic oscillator requires to introduce the potential of the form  $V = \kappa X^2$  that admits straightforward solutions in terms of Gaussians and Hermite polynomials similar to the ordinary point-particle oscillator. There are no collective excitations of the Clifford-oscillator in terms of the number of Clifford-bits and which represent the quanta of areas, volumes, hypervolumes,..., associated with the p-loops oscillations in Planck scale units. The logarithm of the degeneracy of the first collective state of the C-space oscillator, as a function of the number of bits, bears the same functional form as the Bekenstein-Hawking black hole entropy, with the upshot that one recovers, in a natural way, the logarithmic corrections to the black-hole entropy as well, if one identifies the number of Clifford-bits with the number of area-quanta of the black hole horizon. For further details about this derivation and the emergence of the Schwarzschild horizon radius relation, the Hawking temperature, the maximal Planck temperature condition, etc., we refer to [21]. Perhaps the most important consequence of this latter view of black hole entropy is the possibility that there is a ground state of quantum spacetime, resulting from of a Bose-Einstein condensate of the C-space harmonic oscillator.

A C-space version of the Dirac Equation, representing the dynamics of spinning-polyparticles (theories of extended-spin, extended charges) is obtained via the square-root procedure of the Klein-Gordon equation:

$$-i \left( \frac{\partial}{\partial \sigma} + \gamma^\mu \frac{\partial}{\partial x_\mu} + \gamma^\mu \wedge \gamma^\nu \frac{\partial}{\partial x_{\mu\nu}} + \dots \right) \Psi = M \Psi \quad (121)$$

where  $\Psi(\sigma, x^\mu, x^{\mu\nu}, \dots)$  is a polyvector-valued function, a Clifford-number,  $\Psi = \Psi^A E_A$  of all the C-space variables. Ordinary spinors are nothing but left/right ideals of a Clifford algebra. So they are automatically contained in the polyvector valued wave function  $\Psi$ . The ordinary Dirac equation can be obtained when  $\Psi$  is independent of the extra variables associated with a polyvector-valued coordinates  $X$  (i.e., of  $x^{\mu\nu}$ ,  $x^{\mu\nu\rho}$ , ...). For details see [15].

Thus far we have written ordinary wave equations in C-space, that is, we considered the wave equations for a “point particle” in C-space. From the perspective of the 4-dimensional Minkowski spacetime the latter “point particle” has, of course, a much richer structure than a mere point: it is an extended object, modeled by coordinates  $x^\mu$ ,  $x^{\mu\nu}$ , .... But such modeling does not embrace all the details of an extended object. In order to provide a description with more details, one can consider not the “point particles” in C-space, but *branes* in C-space. They are described by the embeddings  $X = X(\Sigma)$ , that is  $X^M = X^M(\Sigma^A)$ , considered in sec.3.2. Quantization of such branes can employ wave functional equation, or other methods, including the second quantization formalism. For a more detailed study of the second quantization of extended objects using the tools of Clifford algebra see [15].

Without employing Clifford algebra a lot of illuminating work has been done in relation to description of branes in terms of p-loop coordinates [124]. A bosonic/fermionic p-brane wave-functional equation was presented in [12], generalizing the closed-string(loop) results in [13] and the quantum bosonic p-brane propagator, in the quenched-reduced minisuperspace approximation, was attained by [18]. In the latter work branes are described

in terms of the collective coordinates which are just the highest grade components in the expansion of a polyvector  $X$  given in eq(2). This work thus paved the way for the next logical step, that is, to consider other multivector components of  $X$  in a unified description of all branes.

Notice that the approach based on eqs.(120),(121) is different from that by Hestenes [1] who proposed an equation which is known as the Dirac-Hestenes equation. In the latter equation instead of the imaginary unit  $i$  there occurs the bivector  $\gamma_1\gamma_2$ . Its square is  $-1$  and it commutes with all the elements of the Dirac algebra which is just a desired property. But on the other hand the introduction of a bivector into an equation of motion implies a selection of a preferred orientation in spacetime. A question arises of why the preferred orientation defined by  $\gamma_1\gamma_2$  and not some other preferred orientation  $a \wedge b$  defined by orthonormal vectors  $a$  and  $b$  which are linear superpositions of  $\gamma_\mu$ ? How such a preferred orientation is determined? Is there some hidden dynamical principle which determines the preferred orientation? What is the more fundamental action which incorporates a kinetic term for the vectors  $a$  and  $b$ ?

Without such a more fundamental dynamical principle the Dirac-Hestenes equation is not covariant and the corresponding action is not invariant with respect to the *active* Lorentz transformations. Hestenes has argued that since in his equation there occur geometric objects, *vectors*, and not their *components*, the equation is covariant, because under the transformations both the components and the basis vectors change in such a way that the vectors remain the same. This is true, but only for *passive transformations*, and not for *active transformations* which affect either the components or the basis vectors. At this point one could argue that the Dirac-Hestenes equation is covariant under the active transformations as well, since instead of  $a \wedge b$  in the transformed equation there occurs  $a' \wedge b'$  and thus the form of equation is not changed. Such an argument would be fine if the vectors  $a$  and  $b$  were dynamical. But in the current formulation of the Dirac-Hestenes theory they are not dynamical, therefore the theory is not Lorentz invariant.

The approach in refs. [16, 15, 17, 8] is different. We start from the usual formulation of quantum theory and extend it to  $C$ -space. We retain the imaginary unit  $i$ . Next step is to give a geometric interpretation to  $i$ . Instead of trying to find a geometric origin of  $i$  in *spacetime* we adopt the interpretation proposed in [15] according to which the  $i$  is the bivector of the 2-dimensional *phase space* (whose direct product with the  $n$ -dimensional configuration space gives the  $2n$ -dimensional phase space).<sup>5</sup> This appears to be a natural assumption due to the fact that complex valued quantum mechanical wave functions involve momenta  $p_\mu$  and coordinates  $x^\mu$  (e.g., a plane wave is given by  $\exp[ip_\mu x^\mu]$ , and arbitrary wave packet is a superposition of plane waves).

## 6 Maximal-Acceleration Relativity in Phase-Spaces

In this section we shall discuss the maximal acceleration Relativity principle [68] based on Finsler geometry which does not destroy, nor deform, Lorentz invariance. Our discussion differs from the pseudo-complex Lorentz group description by Schuller [61] related to

---

<sup>5</sup>Yet another interpretation of the imaginary unit  $i$  present in the Heisenberg uncertainty relations has been undertaken by Finkelstein and collaborators [88].

the effects of maximal acceleration in Born-Infeld models that also maintains Lorentz invariance, in contrast to the approaches of Double Special Relativity (DSR). In addition one does not need to modify the energy-momentum addition (conservation) laws in the scattering of particles which break translational invariance. For a discussions on the open problems of Double Special Relativity theories based on kappa-deformed Poincare symmetries [63] and motivated by the anomalous Lorentz-violating dispersion relations in the ultra high energy cosmic rays [71, 72, 73], we refer to [70].

Related to the minimal Planck scale, an upper limit on the maximal acceleration principle in Nature was proposed by long ago Cainello [52]. This idea is a direct consequence of a suggestion made years earlier by Max Born on a Dual Relativity principle operating in phase spaces [49], [74] where there is an upper bound on the four-force (maximal string tension or tidal forces in the string case) acting on a particle as well as an upper bound in the particle velocity. One can combine the maximum speed of light with a minimum Planck scale into a maximal proper-accleration  $a = c^2/L =$  within the framework of Finsler geometry [56]. For a recent status of the geometries behind maximal-acceleration see [73]; its relation to the Double Special Relativity programs was studied by [55] and the possibility that Moyal deformations of Poincare algebras could be related to the kappa-deformed Poincare algebras was raised in [68]. A thorough study of Finsler geometry and Clifford algebras has been undertaken by Vacaru [77].

Other several new physical implications of the maximal acceleration principle in Nature, like neutrino oscillations and other phenomena, have been studied by [54], [67], [42]. Recently, the variations of the fine structure constant  $\alpha$  [64], with the cosmological accelerated expansion of the Universe, was recast as a renormalization group-like equation governing the cosmological reshift (Universe scale) variations of  $\alpha$  based on this maximal acceleration principle in Nature [68]. The fine structure constant was smaller in the past. Pushing the cutoff scale to the minimum Planck scale led to the intriguing result that the fine structure constant could have been extremely small (zero) in the early Universe and that all matter in the Universe could have emerged via the Unruh-Rindler-Hawking effect (creation of radiation/matter) due to the acceleration w.r.t the vacuum frame of reference. For reviews on the alledged variations of the fundamental constants in Nature see [65] and for more astonishing variations of  $\alpha$  driven by quintessence see [66].

## 6.1 Clifford algebras in Phase space

We shall employ the procedure described in [15] to construct the Phase Space Clifford algebra that allowed [119] to reproduce the sub-maximally accelerated particle action of [53].

For simplicity we will focus on a two-dim phase space. Let  $e_p, e_q$  be the Clifford-algebra basis elements in a two-dim phase space obeying the following relations [15]:

$$e_p \cdot e_q \equiv \frac{1}{2}(e_q e_p + e_p e_q) = 0. \quad (122)$$

and  $e_p \cdot e_p = e_q \cdot e_q = 1$ .

The Clifford product of  $e_p, e_q$  is by definition the sum of the scalar and the wedge

product:

$$e_p e_q = e_p \cdot e_q + e_p \wedge e_q = 0 + e_p \wedge e_q = i. \quad (123)$$

such that  $i^2 = e_p e_q e_p e_q = -1$ . Hence, the imaginary unit  $i$ ,  $i^2 = -1$  admits a very natural interpretation in terms of Clifford algebras, i.e., it is represented by the wedge product  $i = e_p \wedge e_q$ , a phase-space area element. Such imaginary unit allows us to express vectors in a C-phase space in the form:

$$\begin{aligned} Q &= Q = q e_q + p e_p \\ Q e_q &= q + p e_p e_q = q + i p = z \\ e_q Q &= q + p e_q e_p = q - i p = z^* \end{aligned} \quad (124)$$

which reminds us of the creation/annihilation operators used in the harmonic oscillator.

We shall now review the steps in [119] to reproduce the sub-maximally accelerated particle action [53]. The phase-space analog of the spacetime action is:

$$dQ \cdot dQ = (dq)^2 + (dp)^2 \Rightarrow S = m \int \sqrt{(dq)^2 + (dp)^2}. \quad (125)$$

Introducing the appropriate length/mass scale parameters in order to have consistent units yields:

$$S = m \int \sqrt{(dq)^2 + \left(\frac{L}{m}\right)^2 (dp)^2}. \quad (126)$$

where we have introduced the Planck scale  $L$  and have chosen the natural units  $\hbar = c = 1$ . A detailed physical discussion of the dilational invariant system of units  $\hbar = c = G = 4\pi\epsilon_0 = 1$  was presented in ref. [15].  $G$  is the Newton constant and  $\epsilon_0$  is the permittivity of the vacuum.

Extending this two-dim result to a  $2n$ -dim phase space result requires to have for Clifford basis the elements  $e_{p_\mu}, e_{q_\mu}$ , where  $\mu = 1, 2, 3, \dots, n$ . The action in the  $2n$ -dim phase space is:

$$S = m \int \sqrt{(dq^\mu dq_\mu) + \left(\frac{L}{m}\right)^2 (dp^\mu dp_\mu)} = m \int d\tau \sqrt{1 + \left(\frac{L}{m}\right)^2 (dp^\mu/d\tau)(dp_\mu/d\tau)}. \quad (127)$$

where we have factored-out of the square-root the infinitesimal proper-time displacement  $(d\tau)^2 = dq^\mu dq_\mu$ .

One can recognize the action (127), up to a numerical factor of  $m/a$ , where  $a$  is the proper acceleration, as the same action for a sub-maximally accelerated particle given by Nesterenko [53] by rewriting  $(dp^\mu/d\tau) = m(d^2x^\mu/d\tau^2)$ :

$$S = m \int d\tau \sqrt{1 + L^2 (d^2x^\mu/d\tau^2)(d^2x_\mu/d\tau^2)}. \quad (128)$$

Postulating that the maximal proper-acceleration is given in terms of the speed of light and the minimal Planck scale by  $a = c^2/L = 1/L$ , the action above gives the Nesterenko action, up to a numerical  $m/a$  factor:

$$S = m \int d\tau \sqrt{1 + a^{-2} (d^2x^\mu/d\tau^2)(d^2x_\mu/d\tau^2)}. \quad (129)$$

The proper-acceleration is orthogonal to the proper-velocity and this can be easily verified by differentiating the timelike proper-velocity squared:

$$V^2 = \frac{dx^\mu}{d\tau} \frac{dx_\mu}{d\tau} = V^\mu V_\mu = 1 > 0 \Rightarrow \frac{dV^\mu}{d\tau} V_\mu = \frac{d^2 x^\mu}{d\tau^2} V_\mu = 0. \quad (130)$$

which implies that the proper-acceleration is spacelike:

$$g^2(\tau) = -\frac{dx^\mu}{d\tau} \frac{dx_\mu}{d\tau} > 0 \Rightarrow S = m \int d\tau \sqrt{1 - \frac{g^2}{a^2}} = m \int d\omega. \quad (131)$$

where the analog of the Lorentz time-dilation factor for a sub-maximally accelerated particle is given by

$$d\omega = d\tau \sqrt{1 - \frac{g^2(\tau)}{a^2}}. \quad (132)$$

Therefore the dynamics of a sub-maximally accelerated particle can be reinterpreted as that of a particle moving in the spacetime tangent bundle whose Finsler-like metric is

$$(d\omega)^2 = g_{\mu\nu}(x^\mu, dx^\mu) dx^\mu dx^\nu = (d\tau)^2 \left(1 - \frac{g^2(\tau)}{a^2}\right). \quad (133)$$

The invariant time now is no longer the standard proper-time  $\tau$  but is given by the quantity  $\omega(\tau)$ . The deep connection between the physics of maximal acceleration and Finsler geometry has been analyzed by [56]. This sort of actions involving second derivatives have also been studied in the construction of actions associated with rigid particles (strings) [57], [58], [59], [60] among others.

The action is real-valued if, and only if,  $g^2 < a^2$  in the same fashion that the action in Minkowski spacetime is real-valued if, and only if,  $v^2 < c^2$ . This is the physical reason why there is an upper bound in the proper-acceleration. In the special case of uniformly-accelerated motion  $g(\tau) = g_o = \text{constant}$ , the trajectory of the particle in Minkowski spacetime is a hyperbola.

## 6.2 Invariance under the $U(1, 3)$ Group

In this section we will review in detail the principle of Maximal-acceleration Relativity [68] from the perspective of  $8D$  Phase Spaces and the  $U(1, 3)$  Group. The  $U(1, 3) = SU(1, 3) \otimes U(1)$  Group transformations, which leave invariant the phase-space intervals under rotations, velocity and acceleration boosts, were found by Low [74] and can be simplified drastically when the velocity/acceleration boosts are taken to lie in the  $z$ -direction, leaving the transverse directions  $x, y, p_x, p_y$  intact ; i.e., the  $U(1, 1) = SU(1, 1) \otimes U(1)$  subgroup transformations that leave invariant the phase-space interval are given by (in units of  $\hbar = c = 1$ )

$$\begin{aligned} (d\sigma)^2 &= (dT)^2 - (dX)^2 + \frac{(dE)^2 - (dP)^2}{b^2} = \\ (d\tau)^2 \left[ 1 + \frac{(dE/d\tau)^2 - (dP/d\tau)^2}{b^2} \right] &= (d\tau)^2 \left[ 1 - \frac{m^2 g^2(\tau)}{m_P^2 A_{max}^2} \right]. \end{aligned} \quad (134)$$

where we have factored out the proper time infinitesimal  $(d\tau)^2 = dT^2 - dX^2$  in eq.(134) and the maximal proper-force is set to be  $b \equiv m_P A_{max}$ .  $m_P$  is the Planck mass  $1/L_P$  so that  $b = (1/L_P)^2$ , may also be interpreted as the maximal string tension when  $L_P$  is the Planck scale.

The quantity  $g(\tau)$  is the proper four-acceleration of a particle of mass  $m$  in the  $z$ -direction which we take to be  $X$ . Notice that the invariant interval  $(d\sigma)^2$  in eq.(134) is not strictly the *same* as the interval  $(d\omega)^2$  of the Nesterenko action eq.(129), which was invariant under a pseudo-complexification of the Lorentz group [61]. Only when  $m = m_P$ , the two intervals agree. The interval  $(d\sigma)^2$  described by Low [74] is  $U(1, 3)$ -invariant for the most general transformations in the  $8D$  phase-space. These transformations are rather elaborate, so we refer to the references [74] for details. The analog of the Lorentz relativistic factor in eq.(134) involves the ratios of two proper *forces*. One variable force is given by  $ma$  and the maximal proper force sustained by an *elementary* particle of mass  $m_P$  (a *Planckton*) is assumed to be  $F_{max} = m_{Planck}c^2/L_P$ . When  $m = m_P$ , the ratio-squared of the forces appearing in the relativistic factor of eq.(134) becomes then  $g^2/A_{max}^2$ , and the phase space interval (134) coincides with the geometric interval of (129).

The transformations laws of the coordinates in that leave invariant the interval (134) are [74]:

$$T' = T \cosh \xi + (\xi_v X + \frac{\xi_a P}{b^2}) \frac{\sinh \xi}{\xi}. \quad (135)$$

$$E' = E \cosh \xi + (-\xi_a X + \xi_v P) \frac{\sinh \xi}{\xi}. \quad (136)$$

$$X' = X \cosh \xi + (\xi_v T - \frac{\xi_a E}{b^2}) \frac{\sinh \xi}{\xi}. \quad (137)$$

$$P' = P \cosh \xi + (\xi_v E + \xi_a T) \frac{\sinh \xi}{\xi}. \quad (138)$$

The  $\xi_v$  is velocity-boost rapidity parameter and the  $\xi_a$  is the force/acceleration-boost rapidity parameter of the primed-reference frame. They are defined respectively (in the special case when  $m = m_P$ ):

$$\begin{aligned} \tanh \frac{\xi_v}{c} &= \pm \frac{v}{c} \\ \tanh \frac{\xi_a}{b} &= \pm \frac{ma}{m_P A_{max}} = \frac{a}{A_{max}} \end{aligned} \quad (139)$$

The *effective* boost parameter  $\xi$  of the  $U(1, 1)$  subgroup transformations appearing in eqs.(135)–(138) is defined in terms of the velocity and acceleration boosts parameters  $\xi_v, \xi_a$  respectively as:

$$\xi \equiv \sqrt{\xi_v^2 + \frac{\xi_a^2}{b^2}}. \quad (140)$$

Our definition of the rapidity parameters are *different* than those in [74].

Straightforward algebra allows us to verify that these transformations leave the interval of eq.(134) in classical phase space invariant. They are fully consistent with Born's duality Relativity symmetry principle [49]  $(Q, P) \rightarrow (P, -Q)$ . By inspection we can see

that under Born duality, the transformations in eqs.(135)–(138) are *rotated* into each other, up to numerical  $b$  factors in order to match units. When one sets  $\xi_a = 0$  in (135)–(138) one recovers automatically the standard Lorentz transformations for the  $X, T$  and  $E, P$  variables *separately*, leaving invariant the intervals  $dT^2 - dX^2 = (d\tau)^2$  and  $(dE^2 - dP^2)/b^2$  separately.

When one sets  $\xi_v = 0$  we obtain the transformations rules of the events in Phase space, from one reference-frame into another *uniformly*-accelerated frame of reference,  $a = \text{constant}$ , whose acceleration-rapidity parameter is in this particular case:

$$\xi \equiv \frac{\xi_a}{b}. \quad \tanh \xi = \frac{a}{A_{max}}. \quad (141)$$

The transformations for pure acceleration-boosts in are:

$$T' = T \cosh \xi + \frac{P}{b} \sinh \xi. \quad (142)$$

$$E' = E \cosh \xi - bX \sinh \xi. \quad (143)$$

$$X' = X \cosh \xi - \frac{E}{b} \sinh \xi \quad (144)$$

$$P' = P \cosh \xi + bT \sinh \xi. \quad (145)$$

It is straightforward to verify that the transformations (142)–(144) leave invariant the fully phase space interval (134) but *does not* leave invariant the proper time interval  $(d\tau)^2 = dT^2 - dX^2$ . Only the *combination*:

$$(d\sigma)^2 = (d\tau)^2 \left(1 - \frac{m^2 g^2}{m_P^2 A_{max}^2}\right) \quad (146)$$

is truly left invariant under pure acceleration-boosts (142)–(144). One can verify as well that these transformations satisfy Born's duality symmetry principle:

$$(T, X) \rightarrow (E, P). \quad (E, P) \rightarrow (-T, -X). \quad (147)$$

and  $b \rightarrow \frac{1}{b}$ . The latter Born duality transformation is nothing but a manifestation of the large/small tension duality principle reminiscent of the  $T$ -duality symmetry in string theory; i.e. namely, a small/large radius duality, a winding modes/ Kaluza-Klein modes duality symmetry in string compactifications and the Ultraviolet/Infrared entanglement in Noncommutative Field Theories. Hence, Born's duality principle in exchanging coordinates for momenta could be the underlying physical reason behind  $T$ -duality in string theory.

The composition of two successive pure acceleration-boosts is another pure acceleration-boost with acceleration rapidity given by  $\xi'' = \xi + \xi'$ . The addition of *proper* accelerations follows the usual relativistic composition rule:

$$\tanh \xi'' = \tanh(\xi + \xi') = \frac{\tanh \xi + \tanh \xi'}{1 + \tanh \xi \tanh \xi'} \Rightarrow \frac{a''}{A} = \frac{\frac{a}{A} + \frac{a'}{A}}{1 + \frac{aa'}{A^2}}. \quad (148)$$



and in this fashion the upper limiting *proper* acceleration is never *surpassed* like it happens with the ordinary Special Relativistic addition of velocities.

The group properties of the full combination of velocity and acceleration boosts (135)–(138) requires much more algebra [68]. A careful study reveals that the composition *rule* of two successive full transformations is given by  $\xi'' = \xi + \xi'$  and the transformation laws are *preserved* if, and only if, the  $\xi; \xi'; \xi'' \dots$  parameters obeyed the suitable relations:

$$\frac{\xi_a}{\xi} = \frac{\xi'_a}{\xi'} = \frac{\xi''_a}{\xi''} = \frac{\xi''_a}{\xi + \xi'}. \quad (149)$$

$$\frac{\xi_v}{\xi} = \frac{\xi'_v}{\xi'} = \frac{\xi''_v}{\xi''} = \frac{\xi''_v}{\xi + \xi'}. \quad (150)$$

Finally we arrive at the composition law for the effective, velocity and acceleration boosts parameters  $\xi''; \xi''_v; \xi''_a$  respectively:

$$\xi''_v = \xi_v + \xi'_v. \quad (151)$$

$$\xi''_a = \xi_a + \xi'_a. \quad (152)$$

$$\xi'' = \xi + \xi'. \quad (153)$$

The relations (149, 150, 151, 152, 153) are required in order to prove the *group* composition law of the transformations of (135)–(138) and, consequently, in order to have a truly Maximal-Acceleration Phase Space Relativity theory resulting from a phase-space change of coordinates in the cotangent bundle of spacetime.

### 6.3 Planck-Scale Areas are Invariant under Acceleration Boosts

Having displayed explicitly the Group transformations rules of the coordinates in Phase space we will show why *infinite* acceleration-boosts (which is *not* the same as infinite proper acceleration) preserve Planck-Scale *Areas* [68] as a result of the fact that  $b = (1/L_P^2)$  equals the *maximal* invariant force, or string tension, if the units of  $\hbar = c = 1$  are used.

At Planck-scale  $L_P$  intervals/increments in one reference frame we have by definition (in units of  $\hbar = c = 1$ ):  $\Delta X = \Delta T = L_P$  and  $\Delta E = \Delta P = \frac{1}{L_P}$  where  $b \equiv \frac{1}{L_P^2}$  is the maximal tension. From eqs.(135)–(138) we get for the transformation rules of the finite intervals  $\Delta X, \Delta T, \Delta E, \Delta P$ , from one reference frame into another frame, in the *infinite* acceleration-boost limit  $\xi \rightarrow \infty$ ,

$$\Delta T' = L_P(\cosh\xi + \sinh\xi) \rightarrow \infty \quad (154)$$

$$\Delta E' = \frac{1}{L_P}(\cosh\xi - \sinh\xi) \rightarrow 0 \quad (155)$$

by a simple use of L'Hopital's rule or by noticing that both  $\cosh\xi$ ;  $\sinh\xi$  functions approach infinity at the same rate.

$$\Delta X' = L_P(\cosh\xi - \sinh\xi) \rightarrow 0. \quad (156)$$

$$\Delta P' = \frac{1}{L_P}(\cosh\xi + \sinh\xi) \rightarrow \infty \quad (157)$$

where the discrete displacements of two events in Phase Space are defined:  $\Delta X = X_2 - X_1 = L_P$ ,  $\Delta E = E_2 - E_1 = \frac{1}{L_P}$ ,  $\Delta T = T_2 - T_1 = L_P$  and  $\Delta P = P_2 - P_1 = \frac{1}{L_P}$ .

Due to the identity:

$$\infty \times 0 = (\cosh\xi + \sinh\xi)(\cosh\xi - \sinh\xi) = \cosh^2\xi - \sinh^2\xi = 1 \quad (158)$$

one can see from eqs.(154)–(157) that the Planck-scale *Areas* are truly *invariant* under *infinite* acceleration-boosts  $\xi = \infty$ :

$$\Delta X' \Delta P' = 0 \times \infty = \Delta X \Delta P (\cosh^2\xi - \sinh^2\xi) = \Delta X \Delta P = \frac{L_P}{L_P} = 1. \quad (159)$$

$$\Delta T' \Delta E' = \infty \times 0 = \Delta T \Delta E (\cosh^2\xi - \sinh^2\xi) = \Delta T \Delta E = \frac{L_P}{L_P} = 1. \quad (160)$$

$$\Delta X' \Delta T' = 0 \times \infty = \Delta X \Delta T (\cosh^2\xi - \sinh^2\xi) = \Delta X \Delta T = (L_P)^2. \quad (161)$$

$$\Delta P' \Delta E' = \infty \times 0 = \Delta P \Delta E (\cosh^2\xi - \sinh^2\xi) = \Delta P \Delta E = \frac{1}{L_P^2}. \quad (162)$$

It is important to emphasize that the invariance property of the minimal Planck-scale *Areas* (maximal Tension) is *not* an exclusive property of *infinite* acceleration boosts  $\xi = \infty$ , but, as a result of the identity  $\cosh^2\xi - \sinh^2\xi = 1$ , for all values of  $\xi$ , the minimal Planck-scale *Areas* are *always* invariant under *any* acceleration-boosts transformations. Meaning physically, in units of  $\hbar = c = 1$ , that the Maximal Tension (or maximal Force)  $b = \frac{1}{L_P^2}$  is a true physical *invariant* universal quantity. Also we notice that the Phase-space areas, or cells, in units of  $\hbar$ , are also invariant ! The pure-acceleration boosts transformations are " symplectic " .

The infinite acceleration-boosts are closely related to the infinite red-shift effects when light signals barely escape Black hole Horizons reaching an asymptotic observer with an infinite redshift factor. The important fact is that the Planck-scale *Areas* are truly maintained invariant under acceleration-boosts. This could reveal very important information about Black-holes Entropy and Holography. The logarithmic corrections to the Black-Hole Area-Entropy relation were obtained directly from Clifford-algebraic methods in C-spaces [21], in addition to the derivation of the maximal Planck temperature condition and the Schwarzschild radius in terms of the Thermodynamics of a gas of p-loop-oscillatorsquanta represented by area-bits, volume-bits, ... hyper-volume-bits in Planck scale units. Minimal loop-areas, in Planck units, is also one of the most important consequences found in Loop Quantum Gravity long ago [103].

# 7 Some Further Important Physical Applications Related to the $C$ -Space Physics

## 7.1 Relativity of signature

In previous sections we have seen how Clifford algebra can be used in the formulation of the point particle classical and quantum theory. The metric of spacetime was assumed, as usually, to have the Minkowski signature, and we have used the choice  $(+ - - -)$ . There were arguments in the literature of why the spacetime signature is of the Minkowski type [105, 43]. But there are also studies in which signature changes are admitted [104]. It has been found out [16, 15, 30] that within Clifford algebra the signature of the underlying space is a matter of choice of basis vectors amongst available Clifford numbers. We are now going to review those important topics.

Suppose we have a 4-dimensional space  $V_4$  with signature  $(+ + + +)$ . Let  $e_\mu$ ,  $\mu = 0, 1, 2, 3$ , be basis vectors satisfying

$$e_\mu \cdot e_\nu \equiv \frac{1}{2}(e_\mu e_\nu + e_\nu e_\mu) = \delta_{\mu\nu} , \quad (163)$$

where  $\delta_{\mu\nu}$  is the *Euclidean signature* of  $V_4$ . The vectors  $e_\mu$  can be used as generators of Clifford algebra  $\mathcal{C}_4$  over  $V_4$  with a generic Clifford number (also called polyvector or Clifford aggregate) expanded in term of  $e_J = (1, e_\mu, e_{\mu\nu}, e_{\mu\nu\alpha}, e_{\mu\nu\alpha\beta})$ ,  $\mu < \nu < \alpha < \beta$ ,

$$A = a^J e_J = a + a^\mu e_\mu + a^{\mu\nu} e_\mu e_\nu + a^{\mu\nu\alpha} e_\mu e_\nu e_\alpha + a^{\mu\nu\alpha\beta} e_\mu e_\nu e_\alpha e_\beta. \quad (164)$$

Let us consider the set of four Clifford numbers  $(e_0, e_i e_0)$ ,  $i = 1, 2, 3$ , and denote them as

$$\begin{aligned} e_0 &\equiv \gamma_0, \\ e_i e_0 &\equiv \gamma_i. \end{aligned} \quad (165)$$

The Clifford numbers  $\gamma_\mu$ ,  $\mu = 0, 1, 2, 3$ , satisfy

$$\frac{1}{2}(\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu) = \eta_{\mu\nu} , \quad (166)$$

where  $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$  is the *Minkowski tensor*. We see that the  $\gamma_\mu$  behave as basis vectors in a 4-dimensional space  $V_{1,3}$  with signature  $(+ - - -)$ . We can form a Clifford aggregate

$$\alpha = \alpha^\mu \gamma_\mu \quad (167)$$

which has the properties of a *vector* in  $V_{1,3}$ . From the point of view of the space  $V_4$  the same object  $\alpha$  is a linear combination of a vector and bivector:

$$\alpha = \alpha^0 e_0 + \alpha^i e_i e_0. \quad (168)$$

We may use  $\gamma_\mu$  as generators of the Clifford algebra  $\mathcal{C}_{1,3}$  defined over the pseudo-Euclidean space  $V_{1,3}$ . The basis elements of  $\mathcal{C}_{1,3}$  are  $\gamma_J = (1, \gamma_\mu, \gamma_{\mu\nu}, \gamma_{\mu\nu\alpha}, \gamma_{\mu\nu\alpha\beta})$ , with  $\mu < \nu < \alpha < \beta$ . A generic Clifford aggregate in  $\mathcal{C}_{1,3}$  is given by

$$B = b^J \gamma_J = b + b^\mu \gamma_\mu + b^{\mu\nu} \gamma_\mu \gamma_\nu + b^{\mu\nu\alpha} \gamma_\mu \gamma_\nu \gamma_\alpha + b^{\mu\nu\alpha\beta} \gamma_\mu \gamma_\nu \gamma_\alpha \gamma_\beta. \quad (169)$$

With suitable choice of the coefficients  $b^J = (b, b^\mu, b^{\mu\nu}, b^{\mu\nu\alpha}, b^{\mu\nu\alpha\beta})$  we have that  $B$  of eq.(169) is equal to  $A$  of eq.(164). Thus the same number  $A$  can be described either within  $\mathcal{C}_4$  or within  $\mathcal{C}_{1,3}$ . The expansions (169) and (164) exhaust all possible numbers of the Clifford algebras  $\mathcal{C}_{1,3}$  and  $\mathcal{C}_4$ . The algebra  $\mathcal{C}_{1,3}$  is isomorphic to the algebra  $\mathcal{C}_4$ , and actually they are just two different representations of the same set of Clifford numbers (also being called polyvectors or Clifford aggregates).

As an alternative to (165) we can choose

$$\begin{aligned} e_0 e_3 &\equiv \tilde{\gamma}_0, \\ e_i &\equiv \tilde{\gamma}_i, \end{aligned} \quad (170)$$

from which we have

$$\frac{1}{2}(\tilde{\gamma}_\mu \tilde{\gamma}_\nu + \tilde{\gamma}_\nu \tilde{\gamma}_\mu) = \tilde{\eta}_{\mu\nu} \quad (171)$$

with  $\tilde{\eta}_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ . Obviously  $\tilde{\gamma}_\mu$  are basis vectors of a pseudo-Euclidean space  $\tilde{V}_{1,3}$  and they generate the Clifford algebra over  $\tilde{V}_{1,3}$  which is yet another representation of the same set of objects (i.e., polyvectors). But the spaces  $V_4$ ,  $V_{1,3}$  and  $\tilde{V}_{1,3}$  are not the same and they span different subsets of polyvectors. In a similar way we can obtain spaces with signatures  $(+ - ++)$ ,  $(+ + - +)$ ,  $(+ + + -)$ ,  $(- + --)$ ,  $(- - +-)$ ,  $(- - - +)$  and corresponding higher dimensional analogs. But we cannot obtain signatures of the type  $(+ + --)$ ,  $(+ - +-)$ , etc. In order to obtain such signatures we proceed as follows.

*4-space.* First we observe that the bivector  $\bar{I} = e_3 e_4$  satisfies  $\bar{I}^2 = -1$ , commutes with  $e_1, e_2$  and anticommutes with  $e_3, e_4$ . So we obtain that the set of Clifford numbers  $\gamma_\mu = (e_1 \bar{I}, e_2 \bar{I}, e_3, e_4)$  satisfies

$$\gamma_\mu \cdot \gamma_\nu = \bar{\eta}_{\mu\nu}, \quad (172)$$

where  $\bar{\eta} = \text{diag}(-1, -1, 1, 1)$ .

*8-space.* Let  $e_A$  be basis vectors of 8-dimensional vector space with signature  $(+ + + + + +)$ . Let us decompose

$$\begin{aligned} e_A &= (e_\mu, e_{\bar{\mu}}), & \mu &= 0, 1, 2, 3, \\ & & \bar{\mu} &= \bar{0}, \bar{1}, \bar{2}, \bar{3}. \end{aligned} \quad (173)$$

The inner product of two basis vectors

$$e_A \cdot e_B = \delta_{AB}, \quad (174)$$

then splits into the following set of equations:

$$\begin{aligned} e_\mu \cdot e_\nu &= \delta_{\mu\nu}, \\ e_{\bar{\mu}} \cdot e_{\bar{\nu}} &= \delta_{\bar{\mu}\bar{\nu}}, \\ e_\mu \cdot e_{\bar{\nu}} &= 0. \end{aligned} \quad (175)$$

The number  $\bar{I} = e_{\bar{0}} e_{\bar{1}} e_{\bar{2}} e_{\bar{3}}$  has the properties

$$\begin{aligned} \bar{I}^2 &= 1, \\ \bar{I} e_\mu &= e_\mu \bar{I}, \\ \bar{I} e_{\bar{\mu}} &= -e_{\bar{\mu}} \bar{I}. \end{aligned} \quad (176)$$

The set of numbers

$$\begin{aligned}\gamma_\mu &= e_\mu, \\ \gamma_{\bar{\mu}} &= e_{\bar{\mu}}\bar{I}\end{aligned}\tag{177}$$

satisfies

$$\begin{aligned}\gamma_\mu \cdot \gamma_\nu &= \delta_{\mu\nu}, \\ \gamma_{\bar{\mu}} \cdot \gamma_{\bar{\nu}} &= -\delta_{\bar{\mu}\bar{\nu}}, \\ \gamma_\mu \cdot \gamma_{\bar{\mu}} &= 0.\end{aligned}\tag{178}$$

The numbers  $(\gamma_\mu, \gamma_{\bar{\mu}})$  thus form a set of basis vectors of a vector space  $V_{4,4}$  with signature  $(++++----)$ .

*10-space.* Let  $e_A = (e_\mu, e_{\bar{\mu}})$ ,  $\mu = 1, 2, 3, 4, 5$ ;  $\bar{\mu} = \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}$  be basis vectors of a 10-dimensional Euclidean space  $V_{10}$  with signature  $(++++\dots)$ . We introduce  $\bar{I} = e_{\bar{1}}e_{\bar{2}}e_{\bar{3}}e_{\bar{4}}e_{\bar{5}}$  which satisfies

$$\begin{aligned}\bar{I}^2 &= 1, \\ e_\mu\bar{I} &= -\bar{I}e_\mu, \\ e_{\bar{\mu}}\bar{I} &= \bar{I}e_{\bar{\mu}}.\end{aligned}\tag{179}$$

Then the Clifford numbers

$$\begin{aligned}\gamma_\mu &= e_\mu\bar{I}, \\ \gamma_{\bar{\mu}} &= e_{\bar{\mu}}\end{aligned}\tag{180}$$

satisfy

$$\begin{aligned}\gamma_\mu \cdot \gamma_\nu &= -\delta_{\mu\nu}, \\ \gamma_{\bar{\mu}} \cdot \gamma_{\bar{\nu}} &= \delta_{\bar{\mu}\bar{\nu}}, \\ \gamma_\mu \cdot \gamma_{\bar{\mu}} &= 0.\end{aligned}\tag{181}$$

The set  $\gamma_A = (\gamma_\mu, \gamma_{\bar{\mu}})$  therefore spans the vector space of signature  $(-----++++)$ .

The examples above demonstrate how vector spaces of various signatures are obtained within a given set of polyvectors. Namely, vector spaces of different signature are different subsets of polyvectors within the same Clifford algebra. In other words, vector spaces of different signature are different subspaces of  $C$ -space, i.e., different sections through  $C$ -space.

This has important physical implications. We have argued that physical quantities are polyvectors (Clifford numbers or Clifford aggregates). Physical space is then not simply a vector space (e.g., Minkowski space), but a space of polyvectors, called  $C$ -space, a pandimensional continuum of points, lines, planes, volumes, etc., altogether. Minkowski space is then just a subspace with pseudo-Euclidean signature. Other subspaces with other signatures also exist within the pandimensional continuum  $C$  and they all have physical significance. If we describe a particle as moving in Minkowski spacetime  $V_{1,3}$  we consider only certain physical aspects of the object considered. We have omitted its other physical properties like spin, charge, magnetic moment, etc.. We can as well describe the same object as moving in an Euclidean space  $V_4$ . Again such a description would reflect only a part of the underlying physical situation described by Clifford algebra.

## 7.2 Clifford space and the conformal group

### 7.2.1 Line element in $C$ -space of Minkowski spacetime

In 4-dimensional spacetime a polyvector and its square (1) can be written as

$$dX = d\sigma + dx^\mu \gamma_\mu + \frac{1}{2} dx^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + d\tilde{x}^\mu I \gamma_\mu + d\tilde{\sigma} I \quad (182)$$

$$|dX|^2 = d\sigma^2 + dx^\mu dx_\mu + \frac{1}{2} dx^{\mu\nu} dx_{\mu\nu} - d\tilde{x}^\mu d\tilde{x}_\mu - d\tilde{\sigma}^2 \quad (183)$$

The minus sign in the last two terms of the above quadratic form occurs because in 4-dimensional spacetime with signature  $(+---)$  we have  $I^2 = (\gamma_0 \gamma_1 \gamma_2 \gamma_3)(\gamma_0 \gamma_1 \gamma_2 \gamma_3) = -1$ , and  $I^\dagger I = (\gamma_3 \gamma_2 \gamma_1 \gamma_0)(\gamma_0 \gamma_1 \gamma_2 \gamma_3) = -1$ .

In eq.(183) the line element  $dx^\mu dx_\mu$  of the ordinary special or general relativity is replaced by the line element in Clifford space. A “square root” of such a generalized line element is  $dX$  of eq.(182). The latter object is a *polyvector*, a differential of the coordinate polyvector field

$$X = \sigma + x^\mu \gamma_\mu + \frac{1}{2} x^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + \tilde{x}^\mu I \gamma_\mu + \tilde{\sigma} I \quad (184)$$

whose square is

$$|X|^2 = \sigma^2 + x^\mu x_\mu + \frac{1}{2} x^{\mu\nu} x_{\mu\nu} - \tilde{x}^\mu \tilde{x}_\mu - \tilde{\sigma}^2 \quad (185)$$

The polyvector  $X$  contains not only the vector part  $x^\mu \gamma_\mu$ , but also a *scalar part*  $\sigma$ , *tensor part*  $x^{\mu\nu} \gamma_\mu \wedge \gamma_\nu$ , *pseudovector part*  $\tilde{x}^\mu I \gamma_\mu$  and pseudoscalar part  $\tilde{\sigma} I$ . Similarly for the differential  $dX$ .

When calculating the quadratic forms  $|X|^2$  and  $|dX|^2$  one obtains in 4-dimensional spacetime with pseudo euclidean signature  $(+---)$  the minus sign in front of the squares of the pseudovector and pseudoscalar terms. This is so, because in such a case the pseudoscalar unit square in flat spacetime is  $I^2 = I^\dagger I = -1$ . In 4-dimensions  $I^\dagger = I$  regardless of the signature.

Instead of Lorentz transformations—pseudo rotations in spacetime—which preserve  $x^\mu x_\mu$  and  $dx^\mu dx_\mu$  we have now more general rotations—rotations in  $C$ -space—which preserve  $|X|^2$  and  $|dX|^2$ .

### 7.2.2 $C$ -space and conformal transformations

From (183) and (185) we see [25] that a subgroup of the Clifford Group, or rotations in  $C$ -space is the group  $SO(4,2)$ . The transformations of the latter group rotate  $x^\mu$ ,  $\sigma$ ,  $\tilde{\sigma}$ , but leave  $x^{\mu\nu}$  and  $\tilde{x}^\mu$  unchanged. Although according to our assumption physics takes place in full  $C$ -space, it is very instructive to consider a subspace of  $C$ -space, that we shall call *conformal space* whose isometry group is  $SO(4,2)$ .

Coordinates can be given arbitrary symbols. Let us now use the symbol  $\eta^\mu$  instead of  $x^\mu$ , and  $\eta^5, \eta^6$  instead of  $\tilde{\sigma}, \sigma$ . In other words, instead of  $(x^\mu, \tilde{\sigma}, \sigma)$  we write  $(\eta^\mu, \eta^5, \eta^6) \equiv \eta^a$ ,  $\mu = 0, 1, 2, 3$ ,  $a = 0, 1, 2, 3, 5, 6$ . The quadratic form reads

$$\eta^a \eta_a = g_{ab} \eta^a \eta^b \quad (186)$$

with

$$g_{ab} = \text{diag}(1, -1, -1, -1, -1, 1) \quad (187)$$

being the diagonal metric of the flat 6-dimensional space, a subspace of  $C$ -space, parametrized by coordinates  $\eta^a$ . The transformations which preserve the quadratic form (186) belong to the group  $\text{SO}(4,2)$ . It is well known [38, 39] that the latter group, when taken on the cone

$$\eta^a \eta_a = 0 \quad (188)$$

is isomorphic to the 15-parameter group of conformal transformations in 4-dimensional spacetime [40].

Let us consider first the rotations of  $\eta^5$  and  $\eta^6$  which leave coordinates  $\eta^\mu$  unchanged. The transformations that leave  $-(\eta^5)^2 + (\eta^6)^2$  invariant are

$$\begin{aligned} \eta'^5 &= \eta^5 \cosh \alpha + \eta^6 \sinh \alpha \\ \eta'^6 &= \eta^5 \sinh \alpha + \eta^6 \cosh \alpha \end{aligned} \quad (189)$$

where  $\alpha$  is a parameter of such pseudo rotations.

Instead of the coordinates  $\eta^5, \eta^6$  we can introduce [38, 39] new coordinates  $\kappa, \lambda$  according to

$$\kappa = \eta^5 - \eta^6 \quad (190)$$

$$\lambda = \eta^5 + \eta^6 \quad (191)$$

In the new coordinates the quadratic form (186) reads

$$\eta^a \eta_a = \eta^\mu \eta_\mu - (\eta^5)^2 - (\eta^6)^2 = \eta^\mu \eta_\mu - \kappa \lambda \quad (192)$$

The transformation (189) becomes

$$\kappa' = \rho^{-1} \kappa \quad (193)$$

$$\lambda' = \rho \lambda \quad (194)$$

where  $\rho = e^\alpha$ . This is just a dilation of  $\kappa$  and the inverse dilation of  $\lambda$ .

Let us now introduce new coordinates  $x^\mu$  according  $x^\mu$  to<sup>6</sup>

$$\eta^\mu = \kappa x^\mu \quad (195)$$

Under the transformation (195) we have

$$\eta'^\mu = \eta^\mu \quad (196)$$

but

$$x'^\mu = \rho x^\mu \quad (197)$$

The latter transformation is *dilatation* of coordinates  $x^\mu$ .

Considering now a line element

$$d\eta^a d\eta_a = d\eta^\mu d\eta_\mu - d\kappa d\lambda \quad (198)$$

---

<sup>6</sup>These new coordinates  $x^\mu$  should not be confused with coordinate  $x^\mu$  used in Sec.2.

we find that *on the cone*  $\eta^a \eta_a = 0$  it is

$$d\eta^a d\eta_a = \kappa^2 dx^\mu dx_\mu \quad (199)$$

even if  $\kappa$  is not constant. Under the transformation (193) we have

$$d\eta'^a d\eta'_a = d\eta^a d\eta_a \quad (200)$$

$$dx'^\mu dx'_\mu = \rho^2 dx^\mu dx_\mu \quad (201)$$

The last relation is a *dilatation* of the 4-dimensional line element related to coordinates  $x^\mu$ . In a similar way also other transformations of the group  $\text{SO}(4,2)$  that preserve (188) and (200) we can rewrite in terms of the coordinates  $x^\mu$ . So we obtain—besides dilations—translations, Lorentz transformations, and special conformal transformations; altogether they are called *conformal transformations*. This is a well known old observation [38, 39] and we shall not discuss it further. What we wanted to point out here is that conformal group  $\text{SO}(4,2)$  is a subgroup of the Clifford group.

### 7.2.3 On the physical interpretation of the conformal group $\text{SO}(4,2)$

In order to understand the physical meaning of the transformations (195) from the coordinates  $\eta^\mu$  to the coordinates  $x^\mu$  let us consider the following transformation in 6-dimensional space  $V_6$ :

$$\begin{aligned} x^\mu &= \kappa^{-1} \eta^\mu \\ \alpha &= -\kappa^{-1} \\ \Lambda &= \lambda - \kappa^{-1} \eta^\mu \eta_\mu \end{aligned} \quad (202)$$

This is a transformation from the coordinates  $\eta^a = (\eta^\mu, \kappa, \lambda)$  to the new coordinates  $x^a = (x^\mu, \alpha, \Lambda)$ . No extra condition on coordinates, such as (188), is assumed now. If we calculate the line element in the coordinates  $\eta^a$  and  $x^a$ , respectively, we find the the following relation [27]

$$d\eta^\mu d\eta^\nu g_{\mu\nu} - d\kappa d\lambda = \alpha^{-2} (dx^\mu dx^\nu g_{\mu\nu} - d\alpha d\Lambda) \quad (203)$$

We can interpret a transformation of coordinates *passively* or *actively*. Geometric calculus clarifies significantly the meaning of passive and active transformations. Under a *passive transformation* a vector remains the same, but its components and basis vector change. For a vector  $d\eta = d\eta^a \gamma_a$  we have

$$d\eta' = d\eta'^a \gamma'_a = d\eta^a \gamma_a = d\eta \quad (204)$$

with

$$d\eta'^a = \frac{\partial \eta'^a}{\partial \eta^b} d\eta^b \quad (205)$$

and

$$\gamma'_a = \frac{\partial \eta^b}{\partial \eta'^a} \gamma_b \quad (206)$$



Since the vector is invariant, so it is its square:

$$d\eta'^2 = d\eta'^a \gamma'_a d\eta'^b \gamma'_b = d\eta'^a d\eta'^b g'_{ab} = d\eta^a d\eta^b g_{ab} \quad (207)$$

From (206) we read that the well known relation between new and old coordinates:

$$g'_{ab} = \frac{\partial \eta^c}{\partial \eta'^a} \frac{\partial \eta^d}{\partial \eta'^b} g_{cd} \quad (208)$$

Under an *active transformation* a vector changes. This means that in a fixed basis the components of a vector change:

$$d\eta' = d\eta'^a \gamma_a \quad (209)$$

with

$$d\eta'^a = \frac{\partial \eta'^a}{\partial \eta^b} d\eta^b \quad (210)$$

The transformed vector  $d\eta'$  is different from the original vector  $d\eta = d\eta^a \gamma_a$ . For the square we find

$$d\eta'^2 = d\eta'^a d\eta'^b g_{ab} = \frac{\partial \eta'^a}{\partial \eta^c} \frac{\partial \eta'^b}{\partial \eta^d} d\eta^c d\eta^d g_{ab} \quad (211)$$

i.e., the transformed line element  $d\eta'^2$  is different from the original line element.

Returning now to the coordinate transformation (202) with the identification  $\eta'^a = x^a$ , we can interpret eq. (203) passively or actively.

In the *passive interpretation* the metric tensor and the components  $d\eta^a$  change under a transformation, so that in our particular case the relation (207) becomes

$$dx^a dx^b g'_{ab} = \alpha^{-2} (dx^\mu dx^\nu g_{\mu\nu} - d\alpha d\Lambda) = d\eta^a d\eta^b g_{ab} = d\eta^\mu d\eta^\nu g_{\mu\nu} - d\kappa d\lambda \quad (212)$$

with

$$g'_{ab} = \alpha^{-2} \begin{pmatrix} g_{\mu\nu} & 0 & 0 \\ 0 & 0 & -\frac{1}{2} \\ 0 & -\frac{1}{2} & 0 \end{pmatrix}, \quad g_{ab} = \begin{pmatrix} g_{\mu\nu} & 0 & 0 \\ 0 & 0 & -\frac{1}{2} \\ 0 & -\frac{1}{2} & 0 \end{pmatrix} \quad (213)$$

In the above equation the same infinitesimal distance squared is expressed in two different coordinates  $\eta^a$  or  $x^a$ .

In *active interpretation*, only  $d\eta^a$  change, whilst the metric remains the same, so that the transformed element is

$$dx^a dx^b g_{ab} = dx^\mu dx^\nu g_{\mu\nu} - d\alpha d\Lambda = \kappa^{-2} d\eta^a d\eta^b g_{ab} = \kappa^{-2} (d\eta^\mu d\eta^\nu g_{\mu\nu} - d\kappa d\lambda) \quad (214)$$

The transformed line element  $dx^a dx_a$  is physically different from the original line element  $d\eta^a d\eta_a$  by a factor  $\alpha^2 = \kappa^{-2}$

A rotation (189) in the plane  $(\eta^5, \eta^6)$  (i.e., the transformation (193),(194) of  $(\kappa, \lambda)$ ) manifests in the new coordinates  $x^a$  as a *dilatation* of the line element  $dx^a dx_a = \kappa^{-2} d\eta^a d\eta_a$ :

$$dx'^a dx'_a = \rho^2 dx^a dx_a \quad (215)$$

All this is true in the full space  $V_6$ . On the cone  $\eta^a \eta_a = 0$  we have  $\Lambda = \lambda - \kappa \eta^\mu \eta_\mu = 0$ ,  $d\Lambda = 0$  so that  $dx^a dx_a = dx^\mu dx_\mu$  and we reproduce the relations (201) which is a dilatation

of the 4-dimensional line element. It can be interpreted either passively or actively. In general, the pseudo rotations in  $V_6$ , that is, the transformations of the 15-parameter group  $SO(4,2)$  when expressed in terms of coordinates  $x^a$ , assume on the cone  $\eta^a\eta_a = 0$  the form of the ordinary conformal transformations. They all can be given the active interpretation [27, 28].

We started from the new paradigm that physical phenomena actually occur not in spacetime, but in a larger space, the so called *Clifford space* or *C-space* which is a manifold associated with the Clifford algebra generated by the basis vectors  $\gamma_\mu$  of spacetime. An arbitrary element of Clifford algebra can be expanded in terms of the objects  $E_A$ ,  $A = 1, 2, \dots, 2^D$ , which include, when  $D = 4$ , the scalar unit  $\mathbf{1}$ , vectors  $\gamma_\mu$ , bivectors  $\gamma_\mu \wedge \gamma_\nu$ , pseudovectors  $I\gamma_\mu$  and the pseudoscalar unit  $I \equiv \gamma_5$ . *C-space* contains 6-dimensional subspace  $V_6$  spanned<sup>7</sup> by  $\mathbf{1}$ ,  $\gamma_\mu$ , and  $\gamma_5$ . The metric of  $V_6$  has the signature  $(+ - - - - +)$ . It is well known that the rotations in  $V_6$ , when taken on the conformal cone  $\eta^a\eta_a = 0$ , are isomorphic to the non linear transformations of the conformal group in spacetime. Thus we have found out that *C-space* contains —as a subspace— the 6-dimensional space  $V_6$  in which the conformal group acts linearly. From the physical point of view this is an important and, as far as we know, a novel finding, although it might look mathematically trivial. *So far it has not been clear what could be a physical interpretation of the 6 dimensional conformal space.* Now we see that it is just a subspace of Clifford space. The two extra dimensions, parametrized by  $\kappa$  and  $\lambda$ , are not the ordinary extra dimensions; they are coordinates of Clifford space  $C_4$  of the 4-dimensional Minkowski spacetime  $V_4$ .

We take *C-space* seriously as an arena in which physics takes place. The theory is a very natural, although not trivial, extension of the special relativity in spacetime. In special relativity the transformations that preserve the quadratic form are given an *active interpretation*: they relate the objects or the systems of reference in *relative translational motion*. Analogously also the transformations that preserve the quadratic form (183) or (185) in *C-space* should be given an active interpretation. We have found that among such transformations (rotations in *C-space*) there exist the transformations of the group  $SO(4,2)$ . Those transformations also should be given an active interpretation as the transformations that relate different physical objects or reference frames. Since in the ordinary relativity we do not impose any constraint on the coordinates of a freely moving object so we should not impose any constraint in *C-space*, or in the subspace  $V_6$ . However, by using the projective coordinate transformation (202), without any constraint such as  $\eta^a\eta_a = 0$ , we arrived at the relation (214) for the line elements. If in the coordinates  $\eta^a$  the line element is constant, then in the coordinates  $x^a$  the line element is changing by a scale factor  $\kappa$  which, in general, depends on the evolution parameter  $\tau$ . The line element need not be one associated between two events along a point particle's worldline: it can be between two arbitrary (space-like or time-like) events within an extended object. We may consider the line element ( $\equiv$  distance squared) between two infinitesimally separated events within an extended object such that both events have the same coordinate label

---

<sup>7</sup>It is a well known observation that the generators  $L_{ab}$  of  $SO(4,2)$  can be realized in terms of  $\mathbf{1}$ ,  $\gamma_\mu$ , and  $\gamma_5$ . Lorentz generators are  $M_{\mu\nu} = -\frac{i}{4}[\gamma_\mu, \gamma_\nu]$ , dilatations are generated by  $D = L_{65} = -\frac{1}{2}\gamma_5$ , translations by  $P_\mu = L_{5\mu} + L_{6\mu} = \frac{1}{2}\gamma_\mu(1 - i\gamma_5)$  and the special conformal transformations by  $L_{5\mu} - L_{6\mu} = \frac{1}{2}\gamma_\mu(1 + i\gamma_5)$ . This essentially means that the generators are  $L_{ab} = -\frac{i}{4}[e_a, e_b]$  with  $e_a = (\gamma_\mu, \gamma_5, \mathbf{1})$ , where care must be taken to replace commutators  $[\mathbf{1}, \gamma_5]$  and  $[\mathbf{1}, \gamma_\mu]$  with  $2\gamma_5$  and  $2\gamma_\mu$

$\Lambda$  so that  $d\Lambda = 0$ . Then the 6-dimensional line element  $dx^\mu dx^\nu g_{\mu\nu} - d\alpha d\Lambda$  becomes the 4-dimensional line element  $dx^\mu dx^\nu g_{\mu\nu}$  and, because of (214) it changes with  $\tau$  when  $\kappa$  does change. This means that the object changes its *size*, it is moving dilatationally [27, 28]. We have thus arrived at a very far reaching observation that the relativity in  $C$ -space implies *scale changes of physical objects as a result of free motion, without presence of any forces or such fields as assumed in Weyl theory*. This was advocated long time ago [27, 28], but without recourse to  $C$ -space. However, if we consider the full Clifford space  $C$  and not only the Minkowski spacetime section through  $C$ , then we arrive at a more general dilatational motion [17] related to the polyvector coordinates  $x^{\mu\nu}$ ,  $x^{\mu\nu\alpha}$  and  $x^{0123} \equiv \tilde{\sigma}$  (also denoted  $s$ ) as reviewed in section 3.

### 7.3 C-space Maxwell Electrodynamics

Finally, in this section we will review and complement the proposal of ref.[75] to generalize Maxwell Electrodynamics to  $C$ -spaces, namely, construct the Clifford algebra-valued extension of the Abelian field strength  $F = dA$  associated with ordinary vectors  $A_\mu$ . Using Clifford algebraic methods we shall describe how to generalize Maxwell's theory of Electrodynamics associated with ordinary point-charges to a generalized Maxwell theory in Clifford spaces involving *extended* charges and p-forms of arbitrary rank, not unlike the couplings of p-branes to antisymmetric tensor fields.

Based on the standard definition of the Abelian field strength  $F = dA$  we shall use the same definition in terms of polyvector-valued quantities and differential operators in  $C$ -space

$$A = A_N E^N = \phi \underline{1} + A_\mu \gamma^\mu + A_{\mu\nu} \gamma^\mu \wedge \gamma^\nu + \dots \quad (216)$$

The first component in the expansion  $\phi$  is a scalar field;  $A_\mu$  is the standard Maxwell field  $A_\mu$ , the third component  $A_{\mu\nu}$  is a rank two antisymmetric tensor field...and the last component of the expansion is a pseudo-scalar. The fact that a scalar and pseudo-scalar field appear very naturally in the expansion of the  $C$ -space polyvector valued field  $A_N$  suggests that one could attempt to identify the latter fields with a dilaton-like and axion-like field, respectively. Once again= , in order to match units in the expansion (216), it requires the introduction of suitable powers of a length scale parameter, the Planck scale which is conveniently set to unity.

The differential operator is the generalized Dirac operator

$$d = E^M \partial_M = \underline{1} \partial_\sigma + \gamma^\mu \partial_{x_\mu} + \gamma^\mu \wedge \gamma^\nu \partial_{x_{\mu\nu}} + \dots \quad (217)$$

the polyvector-valued indices  $M, N, \dots$  range from  $1, 2, \dots, 2^D$  since a Clifford algebra in  $D$ -dim has  $2^D$  basis elements. The generalized Maxwell field strength in  $C$ -space is

$$\begin{aligned} F = dA &= E^M \partial_M (E^N A_N) = E^M E^N \partial_M A_N = \frac{1}{2} \{E^M, E^N\} \partial_M A_N + \\ &\frac{1}{2} [E^M, E^N] \partial_M A_N = \frac{1}{2} F_{(MN)} \{E^M, E^N\} + \frac{1}{2} F_{[MN]} [E^M, E^N]. \end{aligned} \quad (218)$$

where one has *decomposed* the Field strength components into a symmetric plus antisymmetric piece by simply writing the Clifford geometric product of two polyvectors  $E^M E^N$  as the sum of an anticommutator plus a commutator piece respectively,

$$F_{(MN)} = \frac{1}{2}(\partial_M A_N + \partial_N A_M). \quad (219)$$

$$F_{[MN]} = \frac{1}{2}(\partial_M A_N - \partial_N A_M). \quad (220)$$

Let the C-space Maxwell action (up to a numerical factor) be given in terms of the antisymmetric part of the field strength:

$$I[A] = \int [DX] F_{[MN]} F^{[MN]}. \quad (221)$$

where  $[DX]$  is a C-space measure comprised of all the (holographic) coordinates degrees of freedom

$$[DX] \equiv (d\sigma)(dx^0 dx^1 \dots)(dx^{01} dx^{02} \dots) \dots (dx^{012 \dots D}). \quad (222)$$

Action (221) is invariant under the gauge transformations

$$A'_M = A_M + \partial_M \Lambda \quad (223)$$

The matter-field minimal coupling (interaction term) is:

$$\int A_M dX^M = \int [DX] J_M A^M, \quad (224)$$

where one has reabsorbed the coupling constant, the C-space analog of the electric charge, within the expression for the  $A$  field itself. Notice that this term (224) has the same form as the coupling of p-branes (whose world volume is  $p + 1$ -dimensional) to antisymmetric tensor fields of rank  $p + 1$ .

The open line integral in C-space of the matter-field interaction term in the action is taken from the polyparticle's proper time interval  $S$  ranging from  $-\infty$  to  $+\infty$  and can be recast via the Stokes law solely in terms of the antisymmetric part of the field strength. This requires closing off the integration contour by a semi-circle that starts at  $S = +\infty$ , goes all the way to C-space infinity, and comes back to the point  $S = -\infty$ . The field strength vanishes along the points of the semi-circle at infinity, and for this reason the net contribution to the contour integral is given by the open-line integral. Therefore, by rewriting the  $\int A_M dX^M$  via the Stokes law relation, it yields

$$\begin{aligned} \int A_M dX^M &= \int F_{[MN]} dS^{[MN]} = \int F_{[MN]} X^M dX^N = \\ &= \int dS F_{[MN]} X^M (dX^N / dS). \end{aligned} \quad (225)$$

where in order to go from the second term to the third term in the above equation we have integrated by parts and then used the Bianchi identity for the antisymmetric component  $F_{[MN]}$ .

The integration by parts permits us to go from a C-space domain integral, represented by the Clifford-value hypersurface  $S^{MN}$ , to = a C-space boundary-line integral

$$\int dS^{MN} = \frac{1}{2} \int (X^M dX^N - X^N dX^M). \quad (226)$$

The pure matter terms in the action are given by the analog of the proper time integral spanned by the motion of a particle in spacetime:

$$\kappa \int dS = \kappa \int dS \sqrt{\frac{dX^M}{dS} \frac{dX_M}{dS}}. \quad (227)$$

where  $\kappa$  is a parameter whose dimensions are  $(mass)^{p+1}$  and  $S$  is the polyparticle proper time in C-space.

The Lorentz force relation in C-space is directly obtained from a variation of

$$\int dS F_{[MN]} X^M (dX^N/dS). \quad (228)$$

and

$$\kappa \int dS = \kappa \int \sqrt{dX^M dX_M}. \quad (229)$$

with respect to the  $X^M$  variables:

$$\kappa \frac{d^2 X_M}{dS^2} = e F_{[MN]} \frac{dX^N}{dS}. \quad (230)$$

where we have re-introduced the C-space charge  $e$  back into the Lorentz force equation in C-space. A variation of the terms in the action w.r.t the  $A_M$  field furnishes the following equation of motion for the  $A$  fields:

$$\partial_M F^{[MN]} = J^N. \quad (231)$$

By taking derivatives on both sides of the last equation with respect to the  $X^N$  coordinate, one obtains due to the symmetry condition of  $\partial_M \partial_N$  versus the antisymmetry of  $F^{[MN]}$  that

$$\partial_N \partial_M F^{[MN]} = 0 = \partial_N J^N = 0. \quad (232)$$

which is precisely the continuity equation for the current.

The continuity equation is essential to ensure that the matter-field coupling term of the action  $\int A_M dX^M = \int [DX] J^M A_M$  is also gauge invariant, which can be readily verified after an integration by parts and setting the boundary terms to zero:

$$\delta \int [DX] J^M A_M = \int [DX] J^M \partial_M \Lambda = - \int [DX] (\partial_M J^M) \Lambda = 0. \quad (233)$$

Gauge invariance also ensures the conservation of the energy-momentum (via Noether's theorem) defined in terms of the Lagrangian density variation. We refer to [75] for further details.

The gauge invariant  $C$ -space Maxwell action as given in eq. (221) is in fact only a part of a more general action given by the expression

$$I[A] = \int [DX] F^\dagger * F \quad (234)$$

The latter action is not gauge invariant, since it contains not only the antisymmetric but also the symmetric part of  $F$ . Therefore the action of the form (234) contains a gauge fixing term.

A lesson that we have from these considerations is that the  $C$ -space Maxwell action written in the form (234) automatically contains a gauge fixing term. Analogous result for *ordinary* Maxwell field is known from Hestenes work [1], although formulated in a slightly different way, namely by directly considering the field equations without employing the action.

It remains to be seen if this construction of  $C$ -space generalized Maxwell Electrodynamics of  $p$ -forms can be generalized to the Nonabelian case when we replace ordinary derivatives by gauge-covariant ones:

$$F = dA \rightarrow F = DA = (dA + A \bullet A). \quad (235)$$

For example, one could define the graded-symmetric product  $E_M \bullet E_N$  based on the graded commutator of Superalgebras:

$$[A, B] = AB - (-1)^{s_A s_B} BA. \quad (236)$$

$s_A, s_B$  is the grade of  $A$  and  $B$  respectively. For bosons the grade is even and for fermions is odd. In this fashion the graded commutator captures both the anti-commutator of two fermions and the commutator of two bosons in one stroke. One may extend this graded bracket definition to the graded structure present in Clifford algebras, and define

$$E_M \bullet E_N = E_M E_N - (-1)^{s_M s_N} E_N E_M. \quad (237)$$

$s_M, s_N$  is the grade of  $E_M$  and  $E_N$  respectively. Even or odd depending on the grade of the basis elements.

One may generalize Maxwell's theory to Born-Infeld nonlinear Electrodynamics in  $C$ -spaces based on this extension of Maxwell Electrodynamics in  $C$ -spaces and to couple a  $C$ -space version of a Yang-Mills theory to  $C$ -space gravity, a higher derivative gravity with torsion, this will be left for a future publication. Clifford algebras have been used in the past [62] to study the Born-Infeld model in ordinary spacetime and to write a nonlinear version of the Dirac equation. The natural incorporation of monopoles in Maxwell's theory was investigated by [84] and a recent critical analysis of "unified" theories of gravity with electromagnetism has been presented by [85]. Most recently [22] has studied the covariance of Maxwell's theory from a Clifford algebraic point of view.

## 8 Concluding Remarks

We have presented a brief review of some of the most important features of the Extended Relativity theory in Clifford-spaces ( $C$ -spaces). The "coordinates"  $X$  are noncommuting

Clifford-valued quantities which incorporate the lines, areas, volumes,...degrees of freedom associated with the collective particle, string, membrane,... dynamics underlying the center-of-mass motion and holographic projections of the  $p$ -loops onto the embedding target spacetime backgrounds. C-space Relativity incorporates the idea of an invariant length, which upon quantization, should lead to the notion of minimal Planck scale [23]. Other relevant features are those of maximal acceleration [52], [49] ; the invariance of Planck-areas under acceleration boosts; the resolution of ordering ambiguities in QFT; supersymmetry ; holography [111]; the emergence of higher derivative gravity with torsion ;and the inclusion of variable dimensions/signatures that allows to study the dynamics of all (closed)  $p$ -branes, for all values of  $p$ , in one single unified footing, by starting with the C-space brane action constructed in this work.

The Conformal group construction presented in 7 , as a natural subgroup of the Clifford group in four-dimensions, needs to be generalized to other dimensions, in particular to two dimensions where the Conformal group is infinite-dimensional. Kinani [122] has shown that the Virasoro algebra can be obtained from generalized Clifford algebras. The construction of area-preserving diffs algebras, like  $w_\infty$  and  $su(\infty)$ , from Clifford algebras remains an open problem. Area-preserving diffs algebras are very important in the study of membranes and gravity since Higher-dim Gravity in  $m + n$ -dim has been shown a while ago to be equivalent to a lower  $m$ -dim Yang-Mills-like gauge theory of diffs of an internal  $n$ -dim space [112] and that amounts to another explanation of the holographic principle behind the *AdS/CFT* duality conjecture [113]. We have shown how C-space Relativity involves scale changes in the sizes of physical objects, in the absence of forces and Weyl'gauge field of dilations. The introduction of scale-motion degrees of freedom has recently been implemented in the wavelet-based regularization procedure of QFT by [82]. The connection to Penrose's Twistors program is another interesting project worthy of investigation.

The quantization and construction of QFTs in C-spaces remains a very daunting task since it may involve the construction of QM in Noncommutative spacetimes [94], braided Hopf quantum Clifford algebras [81], hypercomplex extensions of QM like quaternionic and octonionic QM [91], [89], [90], exceptional group extensions of the Standard Model [80],hyper-matrices and hyper-determinants [83], multi-symplectic mechanics, the de Donde-Weyl formulations of QFT [78], to cite a few, for example. The quantization program in C-spaces should share similar results as those in Loop Quantum Gravity [103], in particular the minimal Planck areas of the expectation values of the area-operator.

Spacetime at the Planck scale may be discrete, fractal, fuzzy, noncommutative... The original Scale Relativity theory in fractal spacetime [23] needs to be extended futher to incorporate the notion of fractal "manifolds". A scale-fractal calculus and a fractal-analysis construction that are esential in building the notion of a fractal "manifold" has been initiated in the past years by [121]. It remains yet to be proven that a scale-fractal calculus in fractal spacetimes is another realization of a Connes Noncommutative Geometry. Fractal strings/branes and their spectrum have been studied by [96] that may require generalized Statistics beyond the Boltzmann-Gibbs, Bose-Einstein and Fermi-Dirac, investigated by [97], [95], among others.

Non-Archimedean geometry has been recognized long ago as the natural one operating at the minimal Planck scale and requires the use  $p$ -adic numbers instead of ordinary

numbers [93]. By implementing the small/large scale, ultraviolet/infrared duality principle associated with QFTs in Noncommutative spaces, see [117] for a review, one would expect an upper maximum scale [23] and a maximum temperature [21] to be operating in Nature. Non-Archimedean Cosmologies based on an upper scale has been investigated by [86].

An upper/lower scale can be accomodated simultaneously and very naturally in the  $q$ -Gravity theory of [106], [69] based on bicovariant quantum group extensions of the Poincare, Conformal group, where the  $q$  deformation parameter could be equated to the quantity  $e^{\Lambda/L}$ , such that both  $\Lambda = 0$  and  $L = \infty$ , yield the same classical  $q = 1$  limit. For a review of  $q$ -deformations of Clifford algebras and their generalizations see [81], [120].

It was advocated long ago by Wheeler and others, that information theory [98], set theory and number theory, may be the ultimate physical theory. The important role of Clifford algebras in information theory have been known for some time [87]. Wheeler's spacetime foam at the Planck scale may be the background source generation of Noise in the Parisi-Wu stochastic qunatization [47] that is very relevant in Number theory [92]. The pre-geometry cellular-networks approach of [99] and the quantum-topos views based on gravitational quantum causal sets, noncommutative topology and category theory [101], [102], [116] deserves a futher study within the C-space Relativity framework, since the latter theory also invokes a Category point of view to the notion of dimensions. C-space is a pandimensional continuum [14], [8]. Dimensions are topological invariants and, since the dimensions of the extended objects change in C-space, topology-change is another ingredient that needs to be addressed in C-space Relativity and which may shed some light into the physical foundations of string/M theory [110]. It has been speculated that the universal symmetries of string theory [100] may be linked to Borcherds Vertex operator algebras (the Monstruous moonshine) that underline the deep interplay between Conformal Field Theories and Number theory. A lot remains to be done to bridge together these numerous branches of physics and mathematics. Many surprises may lie ahead of us.

#### Acknowledgements

We are indebted to Prof. D. Ahluwalia for his kind invitation to write this review article. C.C thanks T. Smith, D. Banerjee and C.Beck for discussions. The work of M. P has been supported by the Ministry of Education, Science and Sport of Slovenia.

## References

- [1] D. Hestenes, "Spacetime Algebra" Gordon and Breach, New York, 1996.  
D. Hestenes and G. Sobczyk, "Clifford Algebra to Geometric Calculus" D. Reidel Publishing Company, Dordrecht, 1984.
- [2] I. R. Porteous, Clifford Algebras and the Classical Groups, Cambridge University Press, 1995.
- [3] W. Baylis, Electrodynamics, A Modern Geometric Approach, Boston, Birkhauser, 1999.



- [4] G. Trayling and W. Baylis, J.Phys. A34 (2001) 3309-3324; Int.J.Mod.Phys. A16S1C (2001) 909-912.
- [5] B. Jancewicz, Multivectors and Clifford Algebra in Electrodynamics World Scientific, Singapore 1989.
- [6] "Clifford Algebras and their applications in Mathematical Physics Vol 1: Algebras and Physics. eds by R. Ablamowicz, B. Fauser. Vol 2: Clifford analysis. eds by J. Ryan, W. Sprosig Birkhauser, Boston 2000.
- [7] P. Lounesto, "Clifford Algebras and Spinors". Cambridge University Press. 1997.
- [8] C. Castro, Chaos, Solitons and Fractals **10** (1999) 295. Chaos, Solitons and Fractals **12** (2001) 1585. " The Search for the Origins of M Theory: Loop Quantum Mechanics, Loops/Strings and Bulk/Boundary Dualities " [arXiv: hep-th/9809102].
- [9] C. Castro *The Programs of the Extended Relativity in C-spaces, towards physical foundations of String Theory*, Advance NATO Workshop on the Nature of Time, Geometry and the Physics of Perception, Tatranksa Lomnica, Slovakia, May 2001. Kluwer Publishers 2003.
- [10] C. Castro, Chaos, Solitons and Fractals **11** (2000) 1663. Foundations of Physics **30** (2000) 1301.
- [11] D. Amati, M. Ciafaloni and G. Veneziano, Phys. Letts **B 197** (1987) 81. D. Gross, P. Mende, Phys. Letts **B 197** (1987) 129. M.Maggiore, Phys. Lett **B 304** (1993) 65.
- [12] S. Ansoldi, C.Castro and E. Spalluci, Class. Quant. Grav **18** (1999) 1833. C. Castro, Chaos, Solitons and Fractals **11** (2000) 1721
- [13] Y.Hosotani, Phys. Rev. Lett **55** (1985) 1719. L.Carson, C.H. Ho and Y. Hosotani, Phys. Rev. **D 37** (1988) 1519. C.H. Ho, Jour. Math. Phys **30** (1989) 2168.
- [14] W. Pezzaglia, "Physical Applications of a Generalized Geometric Calculus" [arXiv: gr-qc/9710027]. Dimensionally Democratic calculus and Principles of Polydimensional Physics [arXiv: gr-qc/9912025]. Classification of Multivector Theories and Modifications of the Postulates of Physics [arXiv: gr-qc/9306006] ; Physical Applications of Generalized Clifford Calculus: Papatetrou equations and Metamorphic Curvature [arXiv: gr-qc/9710027]. Classification of Multivector theories and modification of the postulates of Physics [arXiv: gr-qc/9306006].
- [15] M.Pavšič: "The Landscape of Theoretical Physics: A Global View; From Point Particle to the Brane World and Beyond, in Search of Unifying Principle", Kluwer Academic, Dordrecht 2001.
- [16] M.Pavšič, Foundations of Physics **31** (2001) 1185.
- [17] M.Pavšič, Foundations of Physics **33** (2003) 1277 gr-qc/0211085.

- [18] S. Ansoldi, A. Aurilia, C. Castro, E. Spallucci, Phys. Rev. **D 64** 026003 (2001)
- [19] A. Aurilia, S. Ansoldi and E. Spallucci, Class. Quant. Grav. **19** (2002) 3207 [arXiv:hep-th/0205028].
- [20] C. Castro, M. Pavšič, Phys. Lett. **B 539** (2002) 133.
- [21] C. Castro, Jour. of Entropy **3** (2001) 12-26. C. Castro and Alex Granik, Foundations of Physics **33** (3) (2003) 445.
- [22] T. Ivezić, Foundations of Physics Lett **15** (2002) 27. Invariant Relativistic Electrodynamics, Clifford Algebra Approach arXiv: hep-th/0207250].
- [23] L. Nottale: *Fractal Spacetime and Microphysics, towards the Theory of Scale Relativity*, World Scientific 1992. La Relativité dans tous ses états, Hachette Literature, Paris,
- [24] S. Ansoldi, A. Aurilia and E. Spallucci, Phys. Rev **D 56** no. 4 (1997) 2352. Chaos, Solitons and Fractals **10** (1999) 197.
- [25] C. Castro and M. Pavšič, Int. J. Theor. Phys. **42** (2003) 1693.
- [26] M. Pavšič, Class. Quant. Grav. **20** (2003) 2697 [arXiv:gr-qc/0111092].
- [27] M. Pavšič, Nuovo Cim. B **41** (1977) 397; *International Journal of Theoretical Physics* **14**, 299 (1975)
- [28] M. Pavšič, J. Phys. A **13** (1980) 1367.
- [29] M. Pavšič, Found. Phys. **26** (1996) 159 [arXiv:gr-qc/9506057].
- [30] M. Pavšič, arXiv:gr-qc/0210060.
- [31] J. P. Terletsky, *Doklady Akadm. Nauk SSSR*, **133**, 329 (1960); M. P. Bilaniuk, V. K. Deshpande and E.C. G. Sudarshan, *American Journal of Physics* **30**, 718 (1962); E. Recami and R. Mignani, *Rivista del Nuovo Cimento* **4**, 209 (1974); E. Recami, *Rivista del Nuovo Cimento* **9**, 1 (1986)
- [32] M. Pavšič, J. Phys. A **14** (1981) 3217.
- [33] E.C.G. Stueckelberg, *Helv. Phys. Acta*, **14**, 322 (1941); **14**, 588 (1941); **15**, 23 (1942)
- [34] L.P. Horwitz and C. Piron, *Helv. Phys. Acta*, **46**, 316 (1973); L.P. Horwitz and F. Rohrlich, *Physical Review D* **24**, 1528 (1981); **26**, 3452 (1982); L.P. Horwitz, R.I. Arshansky and A.C. Elitzur *Found. Phys* **18**, 1159 (1988); R.I. Arshansky, L.P. Horwitz and Y. Lavie, *Foundations of Physics* **13**, 1167 (1983)
- [35] M. Pavšič, *Found. Phys.* **21**, 1005 (1991); *Nuovo Cim.* **A104**, 1337 (1991); *Doga, Turkish Journ. Phys.* **17**, 768 (1993) *Found. Phys.* **25**, 819 (1995); *Nuovo Cimento A* **108**, 221 (1995); *Foundations of Physics* **26**, 159 (1996); M. Pavšič, *Nuovo Cimento A* **110**, 369 (1997); *Found. Phys.* **28** (1998); 1443. *Found. Phys.* **28** (1998) 1453.

- [36] C.W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, Freeman, San Francisco, 1973), p. 1215
- [37] A. Schild, *Phys.Rev.* **D16**, 1722 (1977)
- [38] H.A. Kastrup, *Annaled der Physik (Lpz.)* **7**, 388 (1962)
- [39] A.O. Barut and R. B. Haugen, *Annals of Physics* **71**, 519 (1972)
- [40] E. Cunningham, Proc. London Math. Soc. **8**, 77 (1909); H. Bateman, Proc. London Math. Soc. **8**, 223 (1910); T. Fulton, F. Rohrlich and L. Witten, Rev. Mod. Phys. **34**, 442 (1962); J. Wess, Nuovo Cim., **18**, 1086 (1960); G. Mack, Nucl. Phys. **B5**, 499 (1968); A.F. Grillo, Riv. Nuovo Cim. **3**, 146 (1973); J. Niederle and J. Tolar, Czech. J. Phys. **B23**, 871 (1973)
- [41] D. Hestenes and R. Ziegler, Projective Geometry with Clifford Algebra, Acta Applicandae Mathematicae **23** (1991) 25-63.
- [42] D. V Ahluwalia and C. Burgard, General Relativity and Gravitation **28** (10) (1996) 1163. D.V. Ahluwalia: General Relativity and Gravitation **29** (12) (1997) 1491. D.V. Ahluwalia, Phys. Letts **A 275** (2000) 31. G.Adunas, E. Rodriguez-Milla and D.V. Ahluwalia, Phys. Letts **B 485** (2000) 215.
- [43] F.D. Smith, Int. Jour. of Theoretical Physics **24** (1985) 155. Int. Jour. of Theoretical Physics **25** (1985) 355. From Sets to Quarks [arXiv: hep-ph/9708379]. <http://www.innerx.net/personal/tsmith/clfpq.html>.
- [44] F. I. Cooperstock, V. Faraoini, " Extended Planck Scale " [arXiv: hep-th/0302080].
- [45] H. I. Arcos and J. G. Pereira, " Kerr-Newman solutions as a Dirac particle" [arXiv: hep-th/0210103].
- [46] W. Smilga: Higher order terms in the contraction of  $S0(3,2)$  [arXiv: hep-th/0304137]. A. Wyler, C. R. Acad. Sc. Paris **269** (1969) 743-745.
- [47] C.Beck, *Spatio-Temporal Vacuum fluctuations of Quantized Fields* . World Scientific series in Advances in Nonlinear Dynamics, vol. 21 (2002).
- [48] G. Trayling and W. Baylis, J. Phys. **A 34** (2001) 3309. J. Chisholm and R. Farwell: J. Phys. **A 32** (1999) 2805. J. Chisholm and R. Farwell: Foundations of Physics **25** (1995) 1511.
- [49] M. Born: Proc. Royal Society **A 165** (1938) 291. Rev. Mod. Physics **21** (1949) 463.
- [50] *Clifford (Geometric) Algebras, with applications to Physics, Mathematics and Engineering* W. E. Baylis, editor; Birkhauser, Boston, 1997.

- [51] S. Somaro: Higher Spin and the Spacetime Algebra V. Dietrich et al. (eds), *Clifford Algebras and their Applications in Mathematical Physics* (1998) 347-368; Kluwer Academic Publishers, the Netherlands.
- [52] E. Caianiello, "Is there a maximal acceleration?", *Lett. Nuovo Cimento* **32** (1981) 65.
- [53] V. Nesterenko, *Class. Quant. Grav.* **9** (1992) 1101; *Phys. Lett.* **B 327** (1994) 50;
- [54] V. Bozza, A. Feoli, G. Lambiase, G. Papini and G. Scarpetta, *Phys. Lett.* **A 283** (2001) 53. V. Nesterenko, A. Feoli, G. Lambiase and G. Scarpetta, *Phys. Rev.* **D 60**, 065001 (1999).
- [55] K. Rama, "Classical velocity in kappa-deformed Poincare algebra and a maximal acceleration" [arXiv: hep-th/0209129].
- [56] H. Brandt: *Contemporary Mathematics* **196** (1996) 273. *Chaos, Solitons and Fractals* **10** (2-3) (1999) 267.
- [57] M. Pavšič: *Phys. Lett.* **B 205** (1988) 231 ; *Phys. Lett.* **B 221** (1989) 264. H. Arodz, A. Sitarz, P. Wegrzyn: *Acta Physica Polonica* **B 20** (1989) 921
- [58] M. Plyushchay, " Comment on the relativistic particle with curvature and torsion of world trajectory" [arXiv: hep-th/9810101]. *Mod. Phys. Lett.* **A10** (1995) 1463-1469
- [59] E. Ramos, J. Roca, *Nucl. Phys.* **B477** (1996) 606-622
- [60] H. Kleinert, A.M. Chervyakov, " Evidence for Negative Stiffness of QCD Strings " [arXiv: hep-th/9601030].
- [61] F. Schuller, *Annals of Phys.* **299** (2002) 174.
- [62] A. Chernitskii: Born-Infeld electrodynamics, Clifford numbers and spinor representations [arXiv: hep-th/0009121].
- [63] J. Lukierski, A. Nowicki, H. Ruegg, V. Tolstoy, *Phys. Lett.* **264** (1991) 331. J. Lukierski, H. Ruegg, W. Zakrzewski: *Ann. Phys.* **243** (1995) 90. J. Lukierski, A. Nowicki: Double Special Relativity versus kappa-deformed dynamics. [arXiv: hep-th/0203065].
- [64] J. Webb, M. Murphy, V. Flambaum, V. Dzuba, J. Barrow, C. Churchill, J. Prochaska, and A. Wolfe, " Further evidence for Cosmological Evolution of the Fine Structure Constant " *Monthly Notices of the Royal Astronomical Society* **327** (2001) 1208.
- [65] J.P. Uzan, " The fundamental constants and their variations: observational status and theoretical motivations" [arXiv: hep-ph/0205340].
- [66] L. Anchordogui, H. Goldberg, *Phys. Rev.* **D 68** (2003) 083513.

- [67] G. Lambiase, G.Papini. G. Scarpetta: Maximal Acceleration Corrections to the Lamb Shift of one Electron Atoms [ arXiv: hep-th/9702130]. G. Lambiase, G.Papini. G. Scarpetta, Phys. Lett **A 224** (1998) 349. G. Papini, " Shadows of a maximal acceleration" [arXiv: gr-qc/0211011].
- [68] C. Castro, Int. J. Mod. Phys. A **18** (2003) 5445 [arXiv: hep-th/0210061]
- [69] L. Castellani: Phys. Lett **B 327** (1994) 22. Comm. Math. Phys **171** (1995) 383.
- [70] G.Amelino-Camelia, Phys. Lett **B 510** (2001) 255. Int. J. Mod. Phys **D 11** (2002) 35. Int. J. Mod. Phys **D 11** (2002) 1643.
- [71] K. Greisen, Phys. Rev. Lett **16** (1966) 748. G.T. Zatsepin, V. Kurmin, Sov. Phys. JETP Lett **4** (1966) 78.
- [72] J. Ellis, N. Mavromatos and D. V. Nanopolous, Chaos, Solitons and Fractals, **10** (1999) 345.
- [73] M. Toller, "The Geometry of Maximal Acceleration" [ArXiv: hep-th/0312016].
- [74] S. Low: Jour. Phys **A Math. Gen 35** (2002) 5711.
- [75] C.Castro, Mod. Phys. Letts. **A 19** (2004) 19
- [76] C. Castro, " The charge-mass-spin relationship of a Clifford polyparticle, Kerr-Newman Black holes and the Fine structure Constant " submitted to Foundations of Physics.
- [77] S. Vacaru and N Vicol: Nonlinear Connections and Clifford Structures math.DG/0205190. S. Vacaru: " (Non) Commutative Finsler Geometry from String/M Theory " [arXiv: hep-th/0211068].
- [78] I.Kanatchikov, Rep. Math. Phys **43** (1999) 157. V. Kisil, [arXiv: quant-ph/0306101].
- [79] N. Bjerrus-Bohr, " Quantum Gravity at large number of dimensions" [arXiv: hep-th/0310263]. J.F. Donoghue, Phys. Rev. **D 50** (1994) 3874. A. Strominger, Phys. Rev. **D 24** (1981) 3082.
- [80] P. Ramond, " Exceptional Groups and Physics " [arXiv: hep-th/0301050]
- [81] . B. Fauser, " A treatise on Quantum Clifford Algebras" [ arXiv: math.QA/0202059.]
- [82] M. Altaisky, " Wavelet based regularization for Euclidean field theory and stochastic quantization " [arXiv: hep-th/0311048].
- [83] V. Tapia, " Polynomial identities for Hypermatrices" [ arXiv: math-ph/0208010].
- [84] M. Defaria-Rosa, E. Recami and W. Rodrigues, Phys. Letts **B 173** (1986) 233.

- [85] E. Capelas de Oliveira and W.A Rodrigues, " Clifford valued Differential Forms, Algebraic spinor fields, Gravitation, Electromagnetism and " unified " theories " [arXiv: math-ph/0311001].
- [86] K. Avinash and V. I. Rvachev, Foundations of Physics **30**(2000) 139.
- [87] D. Gottesman, "The Heisenberg Representation of Quantum Computers" [ArXiv: quant-ph/9807006].
- [88] J. Baugh, D. Finkelstein, A. Galiatdinov and M. Shiri-Garakai, Found.Phys. **33** (2003) 1267-1275
- [89] S.L. Adler, *Quaternionic Quantum Mechanics and Quantum Fields* Oxford Univ. Press, New York, 1995.
- [90] S. de Leo and K. Abdel-Khalek, " Towards an Octonionic World " [arXiv: hep-th/9905123]. S. de Leo, " Hypercomplex Group Theory " [arXiv: physics/9703033].
- [91] C.Castro, " Noncommutative QM and Geometry from quantization in C-spaces " [arXiv: hep-th/0206181].
- [92] M. Watkins, " Number theory and Physics " website, <http://www.maths.ex.ac.uk/mwatkins>
- [93] V. Vladimirov, I. Volovich and I. Zelenov, *p-adic Numbers in Mathematical Physics* World Scientific, 1994, Singapore. L. Brekke and P. Freund, Phys. Reports **231** (1991) 1. M. Pitkanen, *Topological Geometroynamics* <http://www.physics.helsinki.fi/matpitka/tgd.html>
- [94] C.N. Yang, Phys. Rev. **72** (1947) 874; H.S. Snyder, Phys. Rev. **71** (1947) 38. H.S. Snyder, Phys. Rev **72** (1947) 68. S. Tanaka, Yang's Quantized Spacetime Algebra and Holographic Hypothesis [arXiv: hep-th/0303105].
- [95] W. da Cruz: "Fractal von Neumann Entropy" [arXiv: cond-mat/0201489]. " Fractons and Fractal Statistics " arXiv: hep-th/9905229.
- [96] M. Lapidus and M. Frankenhuysen, *Fractal strings, complex dimensions and the zeros of the zeta function* Birkhauser, New York (2000). C.Castro, Chaos, Solitons and Fractals **14** (2002) 1341. **ibid** **15** (2003) 797.
- [97] J. Havrda and F. Charvat: Kybernetika **3** (1967) 30. C-Tsallis, Jour. of Statistical Physics **52** (1988) 479. C.Tsallis, " Non-extensive Statistical mechanics: A brief review of its present status " [arXiv: cond-mat/0205571]. C. Castro, " The Nonextensive Statistics of Fractal Strings and Branes " submitted to Physica A.
- [98] D. Frieden, *Physics from Fisher Information Theory*, Cambridge University Press 1998.
- [99] T. Nowotny, M. Requardt, Chaos, Solitons and Fractals **10** (1999) 469.

- [100] , F. Lizzi, R.J Szabo, Chaos, Solitons and Fractals **10** (1999) 445.
- [101] I. Raptis, " Quantum Space-Time as a Quantum Causal Set" [arXiv gr-qc/0201004].  
" Presheaves, Sheaves and their Topoi in Quantum Gravity and Quantum Logic"  
[arXiv: gr-qc/0110064]. " Non-Commutative Topology for Curved Quantum Causal-  
ity " [arXiv: gr-qc/0101082].
- [102] C. Isham and J. Butterfield, Found.Phys. **30** (2000) 1707-1735. A. K. Guts, E. B. Grinkevich, " Toposes in General Theory of Relativity " [arXiv: gr-qc/9610073].
- [103] A. Ashtekar, C. Rovelli and L. Smolin, Phys. Rev. Lett **69** (1992) 237. C. Rovelli,  
" A dialog on quantum gravity " [arXiv: hep-th/0310077] L. Freidel, E. Livine and  
Carlo Rovelli, Class.Quant.Grav. **20** (2003) 1463-1478. L. Smolin, " How far are we  
from the quantum theory of gravity? " [ arXiv:hep-th/0303185]
- [104] M.Saniga, Chaos, Solitons and Fractals **19** (2004) 739-741
- [105] Norma Mankoc Borstnik and H.B. Nielsen, J. Math. Phys. **44** (2003) 4817-4827
- [106] L. Castellani, Class. Quant. Grav. **17** (2000) 3377-3402 Phys.Lett. B327 (1994) 22-28
- [107] E. Guendelman, E. Nissimov and S. Pacheva, " Strings, p-Branes and Dp-Branes  
With Dynamical Tension ". hep-th/0304269
- [108] P. Bouwknegt, K. Schouetens, Phys. Reports **223** (1993) 183-276. E. Sezgin, "  
Aspects of  $W_\infty$  Symmetry " hep-th/9112025. X. Shen, Int.J.Mod.Phys. A7 (1992)  
6953-6994.
- [109] M. Vasiliev: " Higher Spin Gauge Theories, Star Product and AdS spaces " [arXiv:  
hep-th/9910096]. M. Vasiliev, S. Prokushkin: "Higher-Spin Gauge Theories with  
Matter". [arXiv: hep-th/9812242, hep-th/9806236].
- [110] Y. Ne'eman, E. Eizenberg, "Membranes and Other Extendons ( $p$ -branes) ", World  
Scientific Lecture Notes in Physics vol. **39** 1995. J. Polchinski, *Superstrings*. Cam-  
bridge University Press (2000). M. Green, J. Schwarz and E. Witten, *Superstring  
Theory*. Cambridge University Press (1986).
- [111] J. Maldacena, Adv. Theor. Math. Phys **2** (1998) 231.
- [112] Y.Cho, K, Soh, Q. Park, J. Yoon: Phys. Lets **B 286** (1992) 251. J. Yoon, Phys.  
Letts **B 308** (1993) 240; J. Yoon, Phys. Lett **A 292** (2001) 166. J. Yoon, Class.  
Quan. Grav **16** (1999) 1863.
- [113] C. Castro, Europhysics Letters **61** (4) (2003) 480. Class. Quant. Gravity **20** no. 16  
(2003) 3577.
- [114] A.B. Zamoldchikov, Teor. Fiz **65** (1985) 347 Pope et al C. Pope, Nucl.Phys. B413  
(1994) 413-432 C. Pope, L. Romans, X. Shen, Nuc. Phys, **B 339** (1990) 191. C.  
Pope, L. Romans, X. Shen, Phys. Letts **B 236** (1990) 173. C. Pope, L. Romans, X.  
Shen, Phys. Letts **B 242** (1990) 401.

- [115] B. Dolan, Tchrakian: *Phys. Letts B* **202** (2) (1988) 211.
- [116] E. Hawkins, F. Markopoulou and H. Sahlmann " Evolution in Quantum Causal Histories " [arXiv ; hep-th/0302111]. F. Markopolu, L. Smolin, " Quantum Theory from Quantum Gravity " [arXiv: gr-qc/0311059].
- [117] M.Douglas, N. Nekrasov, *Rev.Mod.Phys.* 73 (2001) 977-1029
- [118] T. Eguchi, *Phys. Rev. Lett.* **44**, 126 (1980)
- [119] C. Castro, " Maximal-acceleration phase space relativity from Clifford algebras " [arXiv: hep-th/0208138].
- [120] V. Abramov, R. Kerner and B. Le Roy, *J.Math.Phys.* 38 (1997) 1650-1669
- [121] J. Cresson, " Scale calculus and the Schrodinger equation " [arXiv: math.GM/0211071].
- [122] E.H. Kinani, " Between Quantum Virasoro Algebras and Generalize= d Clifford Algebras [arXiv: math-ph/0310044].
- [123] S. Capozziello, S. Carloni and A. Troisi, " Quintessence without scalar fields " [arXiv: astro-ph/0303041]. S. Carroll, V. Duvvuri, M. Trodden and M. Turner, " Is Cosmic Speed-Up Due to New Gravitational Physics? " [arXiv: astro-ph/0306438]. A. Lue, R. Scoccimarro and G. Strakman, " Differentiating between Modified Gravity and Dark Energy " [arXiv: astro-ph/0307034].
- [124] A. Aurilia, A. Smailagic and E. Spallucci, *Physical Review D* **47**,2536 (1993); A. Aurilia and E. Spallucci, *Classical and Quantum Gravity* **10**, 1217 (1993); A. Aurilia, E. Spallucci and I. Vanzetta, *Physical Review D* **50**, 6490 (1994); S. Ansoldi, A. Aurilia and E. Spallucci, *Physical Review D* **53**, 870 (1996); S. Ansoldi, A. Aurilia and E. Spallucci, *Physical Review D* **56**, 2352 (1997)