



OPEN

The fingerprints of climate warming on cereal crops phenology and adaptation options

Zartash Fatima^{1,11}, Mukhtar Ahmed^{2,3,11}✉, Mubshar Hussain^{1,4}, Ghulam Abbas^{1,11}, Sami Ul-Allah⁵, Shakeel Ahmad^{1,11}✉, Niaz Ahmed⁶, Muhammad Arif Ali⁶, Ghulam Sarwar⁷, Ehsan ul Haque⁸, Pakeeza Iqbal⁹ & Sajjad Hussain¹⁰

Growth and development of cereal crops are linked to weather, day length and growing degree-days (GDDs) which make them responsive to the specific environments in specific seasons. Global temperature is rising due to human activities such as burning of fossil fuels and clearance of woodlands for building construction. The rise in temperature disrupts crop growth and development. Disturbance mainly causes a shift in phenological development of crops and affects their economic yield. Scientists and farmers adapt to these phenological shifts, in part, by changing sowing time and cultivar shifts which may increase or decrease crop growth duration. Nonetheless, climate warming is a global phenomenon and cannot be avoided. In this scenario, food security can be ensured by improving cereal production through agronomic management, breeding of climate-adapted genotypes and increasing genetic biodiversity. In this review, climate warming, its impact and consequences are discussed with reference to their influences on phenological shifts. Furthermore, how different cereal crops adapt to climate warming by regulating their phenological development is elaborated. Based on the above mentioned discussion, different management strategies to cope with climate warming are suggested.

Cereal crops are typically grasses grown for their edible grains. Globally, cereal grains are produced in higher quantities than any other type of crop and deliver more food energy to human beings and livestock than other crops^{1,2} (Fig. 1). The cereal production of the top 20 countries in the world is presented in Table 1.

Wheat (*Triticum aestivum* L.), rice (*Oryza sativa* L.), maize (*Zea mays* L.), barley (*Hordeum vulgare* L.) and oat (*Avena sativa* L.) are important cereal crops in most countries. Other includes rye (*Secale cereal*), sorghum (*Sorghum bicolor*), and millet (*Pennisetum glaucum*). These crops belong to the family Poaceae or Gramineae subfamily Oryzaceae (Genus *Oryza*: Rice), Triticeae (Genus *Triticum*: Wheat, *Hordeum*: Barley, *Secale*: Rye), Aveneae (Genus *Avena*: Oat), Paniceae (Genus *Pennisetum*: Millet), Andropogoneae (Genus *Sorghum*, *Zea*: Maize, *Coix*: Job's tears) and Cynodonteae (Genus *Eleusine*, *Ragi*). They contribute over 50% of world's total production of cereals in million tonnes^{3–6}. Cereals as whole grain are a rich source of minerals, carbohydrates, vitamins, oils, fats, protein, fiber content and are preferred for consumption^{7–10} (Table 2).

Local and regional climatic conditions are a primary determinant of agronomic crop productivity. Plant metabolic processes are controlled by weather variables like maximum and minimum temperature, solar radiation, carbon dioxide concentration and availability of water^{11–17}. Cereal crop production is influenced by extreme climatic conditions, like heat waves, storms, drought, salinity and flooding^{17–22}. Similarly, Nicholson et al.²³ reported change in the seasonality of rainfall in large part of Africa. Similar kind of spatiotemporal changes in

¹Department of Agronomy, Bahauddin Zakariya University, Multan 60800, Pakistan. ²Department of Agricultural Research for Northern Sweden, Swedish University of Agricultural Sciences, 90183 Umeå, Sweden. ³Department of Agronomy, Pir Mehr Ali Shah, Arid Agriculture University, Rawalpindi 46300, Pakistan. ⁴Agriculture Discipline, College of Science Health, Engineering and Education, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia. ⁵College of Agriculture, Bahauddin Zakariya University, Bahadur Sub-campus, Layyah 31200, Pakistan. ⁶Department of Soil Science, Bahauddin Zakariya University, Multan 60800, Pakistan. ⁷Cotton Botanist, Cotton Research Station, Ayub Agricultural Research Institute, Faisalabad 38000, Pakistan. ⁸Citrus Research Institute Sargodha, Sargodha 40100, Pakistan. ⁹Department of Botany, University of Agriculture Faisalabad, Faisalabad, Pakistan. ¹⁰Department of Horticulture, Bahauddin Zakariya University, Multan, Pakistan. ¹¹These authors contributed equally: Zartash Fatima, Mukhtar Ahmed, Ghulam Abbas and Shakeel Ahmad. ✉email: mukhtar.ahmed@slu.se; shakeelahmad@bzu.edu.pk




Figure 1. Cereals grain production.

Sr. No	Countries & Years	2016	2015	2014	2013	2012
1	China	580,898,372	572,045,000	557,417,296	552,691,792	539,346,800
2	USA	475,983,881	431,865,458	442,849,090	434,308,450	356,210,124
3	India	294,711,871	284,333,000	296,010,000	294,909,510	293,290,000
4	Russia	117,749,733	102,450,556	103,141,153	90,369,572	68,757,301
5	Indonesia	97,667,060	95,010,276	89,854,891	89,791,565	88,443,150
6	Brazil	84,128,482	106,029,517	101,402,184	100,901,726	89,908,244
7	Argentina	67,024,441	55,650,667	55,876,126	50,703,913	47,449,657
8	Ukraine	65,211,480	59,623,480	63,378,190	62,679,750	45,742,860
9	Bangladesh	56,388,892	54,904,305	55,759,375	54,344,578	52,798,736
10	Canada	55,251,252	53,361,100	51,535,801	66,405,701	51,799,100
11	France	54,654,565	72,875,854	72,579,315	67,537,681	68,341,731
12	Vietnam	48,685,089	50,393,869	50,178,378	49,231,389	48,712,795
13	Germany	45,364,400	48,866,800	52,010,400	47,757,100	45,396,500
14	Pakistan	43,075,694	41,081,682	41,895,811	40,109,711	36,496,350
15	Mexico	38,466,082	34,704,514	36,526,602	33,210,301	33,614,212
16	Turkey	35,276,615	38,632,438	32,708,005	37,475,610	33,370,865
17	Australia	35,230,376	37,195,488	38,419,610	35,587,287	43,362,415
18	Thailand	30,420,758	32,786,514	37,751,337	41,962,227	43,367,009
19	Poland	29,849,223	28,002,726	31,945,433	28,455,154	28,543,870
20	Myanmar	28,109,023	28,634,939	28,792,954	28,626,124	28,344,271

Table 1. World cereal production of top 20 countries during 2012–2016 (metric tonnes). (Source: FAOSTAT).

aridity and shifts of dryland in Iran was concluded by Pour et al.²⁴ which could have severe effect on agricultural production and food security. The release of greenhouse gases due to different anthropogenic activities have affected agricultural activities across geographical regions⁷. Distant changes in air temperature resulted to significant change in plant phenology^{8,25–32}. During the previous century, but predominantly over the most recent decades, the earth has experienced noteworthy climate change, particularly warming trends across globe. The average air temperature has increased almost 0.95 °C from 1980 to 2018, and it is predicted to increase almost 3.0–5.0 °C (depending on region) by the end of this century. In the meantime, world population has grown considerably and will continue to grow at an increasing rate. It is expected that the world will need 70% more food by the middle of the current century^{15,33}.

Food security and climate change mitigation are major challenges in developing countries. Changes in management practices such as fertilizer use efficiency, use of organic fertilizers, use of legume with grasses, optimization of irrigation water, plant breeding and genetic modifications and selection of cultivars in field could be good mitigation strategies to cope climate change^{15,34,35}. These adaptive mitigation strategies could help to design resilience systems to future climate change. Similarly, promotion of climate smart agriculture (CSA) could help to mitigate the challenges of climate change. CSA aims to transform and redirect agricultural system to support sustainable development and food security under changing climate. It integrates three elements of sustainable development (economic, social, and environmental) by addressing food security and climate change. It has three main pillars (i) Sustainably increasing agricultural productivity and incomes (ii) Adapting and building

Nutritional values/Crops	Rice	Maize	Wheat	Barley	Oat	Rye	Sorghum	Millet
Water (g)	13.9	12.2	14	10.6	8.5	15	12	11
Energy (kJ)	1518	1517	1320	1535	1644	1268	1422	1468
Energy (kcal)	357	357	310	360	388	298	335	346
Carbohydrate (g)	81.3	77.2	63.9	83.6	69.4	65.9	69.9	68.1
Protein (g)	6.7	9.4	12.7	7.9	11.8	8.2	10.7	11.8
Fat (g)	2.8	3.3	2.2	1.7	9	2	3.3	4.8
Dietary fiber (g)	1.9	2.2	9	5.9	7	11.7	7.5	6.9
Essential amino acid (g/100 g protein)								
Phenylalanine	5.2	4.8	4.6	5.2	5.4	5	5.1	5.5
Histidine	2.5	2.9	2	2.1	2.4	2.4	2.1	2
Isoleucine	4.1	3.6	3	3.6	4.2	3.7	4.1	3.8
Leucine	8.6	12.4	6.3	6.6	7.5	6.4	14.2	10.9
Lysine	4.1	2.7	2.3	3.5	4.2	3.5	2.1	2.7
Methionine	2.4	1.9	1.2	2.2	2.3	1.6	1	2.5
Threonine	1	3.9	2.4	3.2	3.3	3.1	3.3	3.7
Tryptophan	1.4	0.5	2.4	1.5	–	0.8	1	1.3
Valine	5.8	4.9	3.6	5	5.8	4.9	5.4	5.5
Fatty acid profiles (g/100 g food)								
Total fat	2.8	–	1.2	1.7	9.2	2	–	–
Saturated fatty acids	0.74	–	0.16	0.29	1.61	0.27	–	–
Cis-monounsaturated fatty acids	0.66	–	0.13	0.14	3.34	0.21	–	–
Polyunsaturated fatty acids:								
Total cis	0.98	–	0.51	0.77	3.71	0.95	–	–
n–6 (as 18:2)	0.94	–	0.48	0.7	3.52	0.82	–	–
n–3	0.04	–	0.03	0.07	0.19	0.13	–	–
Nutrient	White	Brown	Wholemeal					
Typical Nutrient composition per 100 g of cereal bread								
Energy (kcal/kJ)	219/931	207/882	217/922					
Protein (g)	7.9	7.9	9.4					
Carbohydrate (g)	46.1	42.1	42					
Total sugars (g)	3.4	3.4	2.8					
Starch (g)	42.7	38.7	39.3					
Fat (g)	1.6	2	2.5					
Fibre (g)	1.9	3.5	5					
Thiamine (mg)	0.24	0.22	0.25					
Niacin equivalents (mg)	3.6	4.9	6.1					
Folate (µg)	25	45	40					
Iron (mg)	1.6	2.2	2.4					
Calcium (mg)	177	186	106					

Table 2. Nutritional values of cereals per 100 g. (Source: Price and Welch⁹).

resilience to climate change and (iii) Reducing and/or removing greenhouse gas emissions. Climate change can greatly influence overall food security by influencing cereal crop phenology and changing spatial allocation of crops. According to earlier work, it has been predicted that a 2.0 °C rise in average temperature can lead to a more than 20 to 40% reduction in cereal grain production, particularly in Asia and Africa. Furthermore, temperature rise can have either negative or positive impacts on crop productivity depending upon regions^{36–48}. Shift in the seasonality is the one of the big example of rise in temperature⁴⁹.

Temperature is the most important environmental attribute influencing growth and development, and hence, ultimate productivity of agronomic cereal crops. Rise in temperature also leads to higher evapotranspiration, more crop water and nutrients loss thus resulting to lower water use and nitrogen use efficiency. The effect of temperature on crop phenology could be documented by using GDD or heat units. Heat units, expressed in growing degree-days (GDD), are frequently used to describe the timing of biological processes. The basic equation used is

$$GDD = \left[\frac{(T_{maximum} - T_{minimum})}{2} \right] - T_{base}$$

Crops	T_b (°C)	T_{opt} (°C)	T_{max} (°C)	T_{C1} (°C)	T_{C2} (°C)
Barley	0.0–5.0	25.0–31.0	50.0	30.0	40.0
Maize	8.0	30.0	38.0	33.0	44.0
Millet	10.0	34.0	40.0	30.0	40.0
Oat	0.0–5.0	25.0–31.0	31.0–37.0	–	–
Rice	20.0	28.0	35.0	22.0	30.0
Rye	0.0–5.0	25.0–31.0	31.0–37.0	–	–
Sorghum	8.0	34.0	40.0	32.0	44.0
Wheat	0.0	13.2	35.0	34.0	40.0

Table 3. Cardinal (Base (T_b), optimum (T_{opt}) and maximum (T_{max}) and extreme temperature thresholds (Ceiling vegetative temperature (T_{C1}) and Ceiling reproductive temperature (T_{C2})) for cereal crops. (Source: Ramirez-Villegas et al⁶⁶).

Phenological stage/phase	T_b (°C)	T_{opt} (°C)	T_{max} (°C)
Leaf initiation	– 1.0	22.0	24.0
Shoot growth	3.0	20.3	> 20.9
Root growth	2.0	< 16.3	> 25.0
Sowing	5.0	20.0	30.0
Germination/emergence	0–4.5	24.0–28.0	35.0
Vernalization	– 5.0	0.0–12.0	> 12.0
Tillering	< 3.0	6.0–9.0	> 9.0
Double ridges	4.0	20.0	–
Spikelet Initiation	0.0	15.0	20.0–25.0
Terminal spikelet	3.0	8.0–12.0	–
Shoot elongation	< 12.0	15.0–22.0	> 40.0
Heading	3.9	24.3	–
Anthesis	< 10.0	18.0–24.0	> 32.0
Pollination	> 10.0	18.0–24.0	32.0
Grain-filling	12.0	20.0	35.0
Maturity	< 15.0	22.0–25.0	> 32.0
	T_{lmin}	T_{lmax}	
Lethal Limits	–17.2	47.5	

Table 4. Temperatures (Base (T_b), optimum (T_{opt}) and maximum (T_{max})) for different phenological phases and stages in wheat. (Source: Porter and Gawith⁵³).

where $T_{maximum}$ and $T_{minimum}$ are daily maximum and minimum air temperature, respectively, and T_{base} is the base temperature. It also controls the rate of development from emergence to physiological maturity^{50–54}. All cereal crops have a basic requirement of temperature for completion of a given phenological phase, or the entire life cycle^{55–58}. The daily maximum and minimum temperature play a significant role in determining the optimum sowing time or window and defining the seasonal duration, thus both can affect the achievable yield, produce quality, and along with their sustained productivity^{59–64}. Normally, the longer the optimum growing season, then higher the duration for maximum grain production. Luo⁶⁵, reviewed threshold temperature of different crops and stated that identification of threshold temperature is very important aspect of climate change risk assessment. Furthermore, cardinal T (Base $T = T_{base}$, optimum $T = T_{opt1}$ and T_{opt2} , and failure point $T = T_{fp}$) and lethal T (Lethal minimum $T = T_{lmin}$ and lethal maximum $T = T_{lmax}$) have strong association with crop production. Cardinal and extreme temperature thresholds for the major cereal crops were reviewed by Ramirez-Villegas et al.⁶⁶ as shown in Table 3. Threshold temperature for the different phenological stage/phase of the wheat crop have been presented in Table 4.

The predicted changes in temperature during the next 40 to 70 years are expected to be in the range of 2–3 °C in different regions. Intensity and duration of warming trends and heat wave events are predicted to become more extreme in future than at present and during last decade^{67–69}. Extreme temperature events have short-term spells of a few days with temperature increases over 5.0 °C above normal. Frost caused sterility and abortion of grain whilst extreme heat caused a decrease in the number of grains and reduced grain filling duration^{70–72}. Day-to-day minimum temperatures will rise more rapidly compared to daily maximum temperatures leading to increased mean temperatures. These variations have detrimental effects on yield^{52,73}. The work by Srivastava et al.⁷⁴ concluded that maximum temperature affects the maize grain yield more than the minimum temperature under variable CO₂ levels in both irrigated and rainfed conditions. Furthermore, their study confirmed that maize




Figure 2. A simple presentation of climate warming effects on growth and development of crop plants.

yield was low for RCP 2.6, 4.5, 6.0, and 8.5 under irrigated as compared to rainfed condition. If such changes in temperature will remain occurring in future, then adaptation strategies such as change in sowing date and cultivars shifts are needed to offset such impacts^{75–78}.

Crop phenology is the most significant feature of crop adaptation, and spatial and temporal changes in phenological stages and phases provide well-built indications of the biological influence of thermal trend. Change in phenological seasonality is having both direct and indirect effects on natural vegetation and cereal crop productivity^{25,26,79–81}. Measured climate-warming trends have been implicated as a reason for accelerated cereal crop growth rates and reduced growing durations. Since phenology is highly sensitive to climate change and has significant impact on carbon balance. Thus the asymmetric and opposing response of phenology to daytime and night-time was studied by Wang et al.⁸² and concluded that T_{max} and night-time T_{min} had opposite effects on the timing of start of the season, delayed end of the season and prolonged length of growing season. Furthermore, several earlier studies documented that the intensity of the response of phenological stages and phases of agricultural crops to warming trend is variable at spatio-temporal scales^{79,83,84}. Phenology is well known climate change indicator. However, there is a lack of critical analysis of climatic warming impacts on cereal phenology shift and management of climatic impacts to ensure future global food security. Therefore, a major objective of this review is to analyze the phenological fingerprinting of cereal crops in response to climate warming, and to discuss important management strategies to overcome the climate warming impacts on phenological shift of cereal crops.

Climate warming

The global environment is changing with increasing temperature and carbon dioxide concentration [CO_2]. Generally, burning of fossil fuels are the principle source of climate warming. Jackson et al.⁸⁵ reported that global CO_2 emissions by fossil fuel burning in 2017 were 36.2 billion tonnes, which reached to 37.1 billion tonnes in 2018⁸⁶. Additionally, CO_2 emissions due to biomass combustion were approximately 1/3rd of the fossil fuel emissions during 1997–2010⁸⁷. These CO_2 emissions contribute to global warming and the current fossil fuel reserves have the potential to increase the global temperature up to 2 °C by 2050. Phenology of crop is codetermined by environmental conditions (thermal time requirement) and agronomic practices (Sowing date and cultivar features). Detection of climate warming impact on phenology is not easy mainly because of change in sowing date and introduction of new cultivars. Warming accelerate crop development but it could result to delayed development for long duration cultivar or late sowing date. The impact of climate warming can be mitigated by use of new cultivars with longer thermal time requirement. It has been documented earlier that 1/3rd of the increased temperature impact on phenology was compensated by the introduction of new cultivars with altered temperature requirements. Globally, the climate warming and other stresses are contributing towards yield losses and advocating the necessity of changing the crop calendar (Fig. 2)^{57,88–91}. This necessity for changing the crop calendar is mainly because of changes in the timing of plant phenological stages or events that respond to temperature (Fig. 2). This change in cropping calendar might have positive or negative impacts on productivity, depending on type of crop and geographical locations⁹². He et al.³⁷ reported delayed sowing dates and use of longer duration cultivars are possible management adaptation options needs to be considered to adapt to climate change. Their results indicated that farmers have a wider sowing window in spring and can select frost tolerance and long growing season cultivars that can mitigate detrimental effects of climate warming. Thus, it is very significant to study the climate warming effect on cereal crops phenology.

Impact of climate warming on phenological shift

Phenological stages of the crop plant are set very strictly to the seasonality of the environment and are therefore influenced by changes in the environment. Furthermore, the duration of phenological stages is also associated with CO_2 assimilation, so a shift in phenology may affect crop yield. Tao et al.³⁸ explored the climate-crop-region relation for major crops, i.e., rice, maize and wheat over a period of 1951–2002 and reported a trend of 3.2×10^5 , -1.2×10^5 , and -21.2×10^5 t decade⁻¹ change in production of rice, maize and wheat respectively. They found different impacts of climate warming in different regions. For instance, in northeast China; rice production increased, and in northeast and north regions maize and wheat yield decreased in all studied regions. Wang

et al.⁷ reported shortening of vegetative and reproductive growth phases of double rice grown across southern China in response to temperature increase.

Blecharczyk et al.⁹³ investigated phenological and climate warming data of winter rye grown in Poland from 1957 to 2012 and reported an increase of 2 °C in average temperature. During this period, they observed significant delay in sowing dates and an advancement of 4 days decade⁻¹ in flowering initiation. These changes occurred due to the relative thermal requirement of the crops and cultivar effects. They also reported compensatory effects of cultivars and management practices on crop phenology in response to climate warming. Increased temperature speeds growth, which leads to reduced yield. Ahmad et al.⁹⁴, Huang and Ji⁹⁵ and Wang et al.⁹⁶ reported negative correlations of temperature rise with all the phenological stages besides cotton yield in central and lower Punjab regions of Pakistan. They further reported that about 30% of the negative consequences of climate warming are compensated by sowing new cotton cultivars with high thermal requirements.

Although climate warming is a long and continuous process, it has accelerated during the industrial revolution i.e., after the mid-twentieth century. With rapid industrialization, emission of CO₂ increased in the atmosphere and increasing population lead to deforestation to clear new residential area. Earlier crop flowering and maturity have been observed during the last few decades, but crop management also has impacted the effect of climate warming. Siebert and Ewert⁵² reported spatial differences in climatic effects in different regions of Germany due to temperature and day length effects. Compensation of climatic warming effects on cotton, wheat and rice (30, 21 and 35%, respectively) grown in Punjab, Pakistan, using new cultivars with higher thermal requirements are also reported^{88,94}. However, they reported spatio-temporal differences in the crop species for response to climate warming. Sorghum (*Sorghum bicolor* L.) is a very important crop in arid, marginal and warm regions throughout the world, and it is affected by climate warming. Srivastava et al.⁹⁷ analyzed the impact of climate change on sorghum using the InfoCrop-SORGHUM simulation model. They reported regional variation in response to climate warming and projected a 7% yield reduction up to 2020. Sultan et al.⁹⁸ assessed near-term climate warming impacts on sorghum yield using different crop models in West Africa. Based on 1961–1990 data, they projected an increase of 2.8 °C in mean temperature of 2031–2060 and robust change in rainfall pattern and projected 16–20% yield reduction in response to the climate changes. Shew et al.⁹⁹ studied climate-warming impact on 71 cultivars across 17 locations in South Africa from 1998 to 2014 through extensive regression models. The outcomes showed that heat drives wheat yield losses. The reduction was 12.5% with an additional one-day (24 h) exposure to temperature above 30 °C. Furthermore, 8.5%, 18.4% and 28.5% yield reduction was observed for uniform warming scenario of +1 °C, +2 and +3 °C respectively. They suggested that warming impact could be reduced by sharing gene pools with wheat breeding programs. Sonkar et al.¹⁰⁰ reported 7% decrease in wheat yield per 1 °C rise in the mean temperature. Thus, there is a dire need of further research especially on the effects of climate change on the optimum temperatures required for different crop development stages. In short, climate warming has mostly negative impacts on crop species by disturbing their phenology, but these impacts can be compensated or converted into positive effects by proper management, ideotype designing and breeding of fast-growing crops with higher thermal requirements^{37,88,101–106}.

Role of phenological stages and phases in cereal crop grain production

Cereal crop phenology is very sensitive to climate change⁸⁸ as compared to other agronomic crops; consequently, phenological changes are frequently used to measure how climate warming influences the agricultural ecosystems in a particular region^{107–109}. Numerous methods have been utilized to elucidate the intensity of the effect of climate change on agronomic cereal crops. The methods include analysis of satellite images on vegetation greenness, measurement of net primary production with Normalized Difference Vegetation Index (NDVI), and particularly, determination of temporal and spatial variations in phenological seasonality^{26,79,110–115}. The data on stages and phases of cereal crop phenology and climate meteorological data are collected from phenological networks or through designated observation stations.

Wheat phenology trend. Wheat phenology has been affected by climate warming worldwide. Identifying phenological responses to climate warming is tough due to regularly shifting sowing dates and the introduction of new cultivars. Wheat phenological trend from sowing to maturity is presented in Table 5. The results showed that it remained maximum in northwest China during 1983 to 2004 (sowing = 13.2 days decade⁻¹, emergence = 9.8 days decade⁻¹, anthesis = 11.0 days decade⁻¹ and maturity = 10.8 days decade⁻¹). However, highest trend in delaying of phenological stages was observed in north south regions of China during 1981–2010¹¹⁶. The highest reduction in phenological phases was also observed in northeast China¹¹. Xiao et al.¹¹⁷ observed that since 1981 to 2009, climate-warming in North China Plain caused dates of green-up after winter dormancy, anthesis, and winter wheat maturity to happen in an average of 1.1, 2.7, and 1.4 days earlier decade⁻¹, respectively. Wheat phenology was recorded at three variable sites of rainfed Pothwar and it depicted significant variability. The highest days to flowering and maturity was observed at Islamabad while lowest were at Attock which might be due to variability in climatic conditions during wheat growing cycle (Fig. 3). Ahmad et al.⁸⁸ reported that sowing (S), emergence (E), anthesis (A) and maturity (M) for wheat crop were delayed by 9.5, 1.3, 5.3 and 5.4 days decade⁻¹ while phenological-phases S-A, A-M along with S-M were reduced by 5.4, 5.5, 4.6 and 5.7 days decade⁻¹ in Pakistan during 1980 to 2014 (Fig. 3). The S-M phase was reduced by 7.2, 10.7, 5.4, 4.0 and 2.7 days decade⁻¹ in Spain, Australia, Argentina, Romania and Germany, respectively. In India, Russia and USA, crop phenology was also affected due to climate change^{118,119}.

Rice phenology trend. Phenological stages and phases of rice were affected by the worldwide thermal trend. Table 5 shows that sowing, transplanting (T), anthesis and maturity were earlier by 7.9, 6.6, 5.0 and 5.0 days decade⁻¹, respectively in Punjab, Pakistan (Fig. 4) during 1980 to 2014⁸⁸. Both the highest and lowest

Crop	Country	Period	Phenological stages (early/delay days/decade)				Phenological phases (reduction days/decade)			References
			Sowing	Emergence	Anthesis	Maturity	S-A	A-M	S-M	
Wheat	China	1983–2004	13.2 (E)	9.8 (E)	11.0 (E)	10.8 (E)	16.1	8.2	12.3	7
	Pakistan	1980–2014	9.5 (D)	1.3 (D)	5.3 (E)	5.4 (E)	5.5	4.6	5.7	88
	China	1981–2005	7.6 (E)	6.3 (E)	2.0 (E)	4.8 (E)	3.8	4.1	5.8	14
	Spain	1986–2012	3.8 (E)	2.6 (E)	5.2 (E)	2.9 (E)	4.6	5.1	7.2	81
	Australia	1995–2016	3.9 (E)	2.8 (E)	7.5 (E)	5.8 (E)	6.6	7.9	10.7	126
	China	1981–2009	1.2 (D)	1.3 (D)	3.7 (D)	3.1 (E)	5.0	3.1	4.3	127
	Argentina	1971–2000	3.0 (D)	2.9 (D)	4.2 (D)	4.9 (D)	7.5	6.9	5.4	25
	China	1980–2009	4.1 (D)	3.7 (D)	6.3 (D)	8.1 (D)	6.1	2.3	3.6	128
	Romania	1971–2006	3.5 (D)	2.5 (D)	2.2 (D)	3.0 (D)	2.3	3.2	4.0	61
	China	1981–2010	9.0 (D)	8.5 (D)	11.0 (D)	16.2 (D)	3.7	2.5	1.3	27
	China	1981–2009	1.5 (D)	1.7 (D)	2.1 (D)	2.5 (D)	2.0	1.8	3.1	11,28,42,50,128–133
China	1981–2000	3.4 (E)	2.9 (E)	3.0 (E)	3.3 (E)	0.4	0.3	1.0	38	
Germany	1952–2013	2.0 (E)	1.8 (E)	4.1 (E)	5.0 (E)	1.9	0.8	2.7	134,135	
Rice	Pakistan	1980–2014	7.9 (E)	6.6 (E)	5.0 (E)	5.0 (E)	1.4	4.1	6.4	88
	Madagascar	2008–2010	5.4 (E)	3.2 (E)	6.2 (E)	4.8 (E)	4.1	3.2	6.2	108
	China	1981–2006	1.0 (D)	1.4 (D)	2.7 (D)	3.1 (D)	3.3	1.2	4.1	70
	China	1981–2000	5.7 (E)	5.2 (E)	6.2 (E)	3.6 (E)	0.5	2.6	3.1	38
	China	1992–2013	2.2 (D)	1.9 (D)	2.8 (D)	3.4 (D)	0.8	1.7	2.4	136
	China	1981–2012	4.9 (D)	4.2 (D)	3.8 (D)	5.2 (D)	2.4	3.2	5.1	137
	China	1981–2009	3.7 (D)	3.0 (D)	2.0 (D)	4.0 (D)	2.8	1.9	2.2	138
	China	1981–2009	6.5 (D)	5.8 (D)	1.5 (D)	2.4 (D)	2.9	1.6	5.2	139
Maize	Pakistan	1980–2014	3.0 (D)	1.9 (D)	2.8 (D)	4.4 (D)	5.5	2.2	7.8	109
	Pakistan	1980–2014	4.6 (E)	3.7 (E)	7.1 (E)	9.2 (E)	2.4	1.9	4.6	109,140,141
	China	1981–2010	5.4 (D)	4.8 (D)	5.2 (D)	7.1 (D)	1.3	0.8	2.2	142
	China	1990–2012	10.0 (D)	9.4 (D)	10.5 (D)	5.6 (D)	4.1	3.9	5.7	143
	China	1981–2008	8.1 (D)	7.8 (D)	6.2 (D)	3.7 (D)	5.2	2.6	3.7	136
	USA	1981–2005	3.9 (E)	3.2 (E)	1.7 (E)	2.9 (E)	2.9	1.8	3.0	144
	Germany	1961–2000	4.5 (E)	4.1 (E)	5.6 (E)	8.8 (E)	3.1	7.2	4.9	8
	China	1981–2000	1.7 (D)	1.5 (D)	3.3 (D)	5.5 (D)	2.4	2.2	2.8	38
	China	1981–2009	5.0 (D)	3.1 (D)	4.0 (D)	6.7 (D)	1.0	2.7	3.5	27,102
	China	1992–2013	3.5 (D)	3.2 (D)	1.8 (D)	1.5 (D)	1.9	3.3	1.5	145,146
	China	1981–2010	8.7 (D)	6.9 (D)	4.6 (D)	2.2 (D)	4.6	2.4	6.2	146,147
China	1992–2013	1.3 (E)	1.0 (E)	4.1 (E)	2.7 (E)	1.1	2.0	3.0	145	
Oat	Germany	1959–2009	1.1 (E)	1.8 (E)	9.7 (E)	10.7 (E)	7.9	13.9	9.4	52
	Germany	1951–2004	1.5 (E)	1.2 (E)	4.9 (E)	6.4 (E)	3.4	1.5	4.9	110
Barley	Lithuania	1961–2015	1.7 (E)	2.8 (E)	1.1 (E)	0.4 (E)	1.0	1.6	2.2	43
	Spain	1986–2008	2.8 (D)	1.9 (D)	2.7 (D)	3.5 (D)	1.6	2.1	3.7	41
Rye	Poland	1957–2012	2.2 (D)	1.9 (D)	4.0 (D)	3.6 (D)	1.8	1.4	3.2	125
	Germany	1960–2013	1.0 (E)	1.2 (E)	1.8 (E)	1.6 (E)	2.9	3.1	4.5	60

Table 5. Cereal crops observed phenology trends across regions (*E* Early, *D* Delay, *S* Sowing, *T* Transplanting, *A* Anthesis, *M* Maturity).

transplanting-to-maturity phases were reduced in Pakistan and China, respectively. Shrestha et al.¹⁰⁸ reported that sowing, transplanting, anthesis and maturity were earlier by 5.4, 3.2, 6.2 and 4.8 days decade⁻¹ in Madagascar. The phenological phases of sowing to transplanting, transplanting to anthesis and A-M were reduced by 2.9, 1.6 and 5.2 days decade⁻¹, respectively in China during 1981 to 2009 as well as in other parts of world^{46,70,104,120–122}. Transplanting to maturity was more severely reduced than other phenological phases (Table 5).

Maize phenology trend. Phenological stages of maize were delayed in most countries (Table 5). Phenological phases of maize were also reduced in Pakistan (Fig. 4) during 1980 to 2014¹⁰⁹. Lowest reduction of phenological phases was in China¹⁰². In USA, phenological stages S, E, A and M were earlier by 3.9, 3.2, 1.7 and 2.9 days decade⁻¹, respectively, while, S-A, A-M and S-M were shortened by 2.9, 1.8 and 3.0 days decade⁻¹, respectively during 1981 to 2005^{123,124}. The S, E, A, and M were delayed by 3.0, 1.9, 2.8 and 4.4 days decade⁻¹ in Pakistan




Figure 3. Trend of phenological stages of wheat at three variable climatic sites of rainfed Pothwar.

Spring Maize																		
Months	Jan			Feb			Mar			Apr			May			Jun		
Previous				S	E		A								M			
Existing				S	E		A								M			
Fall Maize																		
Months	Jul			Aug			Sep			Oct			Nov			Dec		
Previous				S	E				A						M			
Existing				S	E				A						M			
Rice																		
Months	Jul			Aug			Sep			Oct			Nov			Dec		
Previous		T							PI			A			M			
Existing		T							PI			A			M			
Wheat																		
Months	Nov			Dec			Jan			Feb			Mar			Apr		
Previous	S	E										A						M
Existing	S	E										A						M

Figure 4. Spatial and temporal variability of spring maize, fall maize, rice and wheat crops phenological phases as effected by climate warming in Punjab, Pakistan. S Sowing, E Emergence, A Anthesis, M Maturity, T Transplanting, PI Panicle Initiation. Alphabets with green and red colors represent previous and prevailing trends. [Modified and adapted from Abbas et al.¹⁰⁹ and Ahmad et al.⁸⁸].

during 1980 to 2014¹⁰⁹. Phenological phases S-A, A-M and S-M were reduced by 3.1, 7.2 and 4.9 days decade⁻¹, respectively in Germany during 1961–2000⁸.

Oat phenology trend. Phenology of oat was affected due to climate warming. Table 5 shows that in Germany during 1959 to 2009, S, E, A and M of oat were earlier by 1.1, 1.8, 9.7 and 10.7 days decade⁻¹ and phases S-A, A-M and S-M were significantly reduced by 7.9, 13.9 and 9.4 days decade⁻¹, respectively. Estrella et al.¹¹⁰ reported that during 1951 to 2004, sowing, emergence, flowering and harvesting of oat were earlier by 1.5, 1.2, 4.9 and 6.4 days decade⁻¹ and S-A, A-M besides S-M phases were significantly shortened by 3.4, 1.5 and 4.9 days decade⁻¹, respectively.

Barley phenology trend. Phenological stages of barley were earlier in Lithuania and delayed in Spain with global warming (Table 5). Sujetovienė et al.⁴³ observed that S, E, A and M were delayed by 1.7, 2.8, 1.1 and 0.4 days decade⁻¹ and phases S-A, A-M and S-M were reduced by 1.0, 1.6 and 2.2 days decade⁻¹, respectively

during 1961 to 2015 in Lithuania (Table 5). Phenological phases were reduced by 1.6, 2.1 and 3.7 days decade⁻¹ for S-A, A-M and S-M, correspondingly in Spain during 1986 to 2008⁸¹.

Rye phenology trend. Phenological phases of rye were also reduced in Germany and minimally in Poland¹²⁵ (Table 5). The S, E, A and M were delayed by 2.2, 1.9, 4.0 and 3.6 days decade⁻¹, respectively and phases S-A (1.8 days decade⁻¹), A-M (1.4 days decade⁻¹) and S-M (3.2 days decade⁻¹) were reduced in Poland during 1957 to 2012. Phenological stages S, E, A and M were earlier by 1.0, 1.2, 1.8 and 1.6 days decade⁻¹ and phases S-A, A-M and S-M were shortened by 2.9, 3.1, 4.5 days decade⁻¹, respectively in Germany during 1960–2013 (Table 5).

Effect of phenological shifts on crop yield

Climate warming is shifting the phenological stages of the crop. Crop yield is very sensitive to the accumulation of heat units or GDD during specific phenological events, thus by shifting the phenology, crop yield is also affected. Several studies reported/projected reduction in crop yields with climate warming due to shortening of phenological events without considering management practices^{37,137,139,148}. The climate change adaptation strategies like cultivar shift and change of sowing dates could compensate warming effects. Although vegetative and reproductive phases are equally prone to climate warming^{37,145,149}, the impact of high temperature on pollen viability, fertilization and post-fertilization processes leads to a marked decrease in final yield^{16,145,150}. However, all phenological stages are not equally responsive to climate warming effects and do not have equal impacts to final yield^{12,103,127,151–153}. Tao et al.³⁸ investigated the effects of phenology shift on yield of wheat, maize and rice during 1981 to 2000. They reported that earlier planting of 3–6 days decade⁻¹ and earlier anthesis dates decreased yield up to 110 kg/ha/yr in wheat, 168 kg/ha/yr in maize, and 20 kg/ha/yr in rice. Zacharias et al.¹⁵⁴ investigated the effect of induced high temperature on phenology and yield of wheat and rice cultivars. They observed that high temperature advanced dates of anthesis and crop maturity, and reduced the duration of vegetative and maturity phases, and led to reduction in yield. They reported 26% reduction in grain yield of rice due to reduction in the anthesis and maturity dates of around seven to 8 days, respectively. The reduction in yield was primarily attributed to less time for photosynthesis and accumulation of assimilates. Warmer temperature during the reproductive phase led to pollen sterility and high evapotranspiration, which led to fewer grains and lower grain weight, reduced photosynthetic rate, and reduced duration of phenological events^{90,91}. Hatfield and Prueger⁹⁰ reported that temperature extremes during the reproductive phase could reduce the grain yield of maize up to 90% mainly due to pollen sterility and smaller grain size. Nahar et al.¹⁵⁰ investigated the impact of climate warming on five cultivars of wheat. In their studies heat stress imposed by late sowing of the crop resulted in significant reduction in days to booting, anthesis and maturity across all cultivars (although cultivar differences existed). About 8–13 days reduction in days to anthesis and maturity resulted in 53–73% yield losses in different cultivars. Meanwhile, Sadok and Jagadish¹⁵⁵ reported that nighttime warming poses a threat to global food security (6% decrease in winter wheat yield per 1 °C rise in night temperature while 4–7% reduction in spring wheat and rice yield per 1 °C rise in night temperature) as it is driving yield declines worldwide. Thus, they proposed ecophysiological framework as a guide to implement future research efforts to mitigate yield declines. They further elaborated that efforts should include integrated approaches i.e. physiology with crop modeling, breeding and management to intensify sustainable pathways for mitigation as intensity of climate change is becoming stronger and stronger day by day.

Climate warming significantly impacts the crop phenology, which in turn leads to a reduction in yield mainly due to shortened phenological events coupled with less production of assimilates, pollen sterility and biochemical irregularities¹⁵⁶. The climate warming impacts have been neutralized to some extent with shifting of sowing date and development of cultivars with longer duration of phenological events even under warming conditions. As climate warming is a continuous process, therefore, integrated research is required to understand the mitigation of climate warming impact for sustainable crop production to feed an ever-increasing world population.

Consequences of climate warming

Climate warming is threatening all living organisms including human beings. Rising temperatures at the poles and glaciers, increase the risk of floods, and many populated areas may be inundated, and fertile lands may become deserts^{157–161}. The main consequences linked with a rise in sea level and desertification are salinity, heat and drought stresses, and these factors are major threats to agricultural production including primary cereals (Fig. 5).

Adaptation strategies in response to climate change

Adaptation strategies in CSA are the only way to reduce negative effects of climate warming on cereal crops. Different adaptation approaches like better crop management practices, modern breeding, and biotechnological approaches can develop climate resilient cereals to help cope with climate warming^{56,105,106,111,127,162–165}. Revolutions in plant breeding (molecular and conventional for incorporation desired characteristics) and genetic engineering (integrated transgenic techniques like marker-trait associations and genome-wide selection) could assist to reduce overwhelming food security concerns against extreme-weather conditions, through producing climate adaptive cereal plants^{166–173}. Thus coping with climate warming is an urgent concern globally and in order to adapt cereals to changing climatic conditions, the following approaches are mandatory.

Crop management practices. Better and improved crop management practices can minimize negative impacts of temperature extremes. Under climate warming, all better crop management practices should be brought to the fields to get real benefits^{105,129,174–186}. Usage of quality seed are big concerns and quality seed




Figure 5. Effect of climate warming on productivity of primary cereal crops: wheat, rice and maize across various countries/continents. [Modified and adapted from Ishfaq et al.¹⁹²].

Crop	Continent	Adaptation strategies	References
Wheat	Asia, Spain, Australia, South America, Europe	Early Sowing, Sowing/planting dates adjustment	5,25,60,81,88,101,109,140,189,192–200
	Asia	Cultivars requiring higher GDDs; Improved cultivars; Use of heat tolerant cultivars;	14,109,127,140,198,201
	Asia, South America	Optimum plant population	178,198
Rice	Asia	Sowing date	88,144,202,203
	Asia	Varieties requiring higher GDDs	70,106,107,204–208
	Asia	Direct planting	209,210
	Asia	Early maturing, Climate ready rice	211,212
	Asia	System of rice intensification with alternate wetting and drying	213
	Asia	Improved variety and management	214
Maize	South Asia	Sowing date	78,92,140,141,145,149,215
	Asia, America	Early maturing cultivars; Varieties requiring higher GDDs	144,216
	Asia	Raised bed planting	216–220
	Asia	Precision nutrient management	221
Oat	Europe	Varieties with larger thermal time	52
Rye	Poland, Europe	Late maturing cultivars, Plant density	60,125,134,199,200
Millet	South Asia	Planting time; Plant spacing	222–224

Table 6. Observed changes in trend of phenology of agronomic crops in different countries.

availability to the farmers are under 20%. Good quality and genetically pure seed should be provided to farmers to minimize negative impacts of climate change and achieve sustainable nutrition security⁶. Farmers' access to good quality seed of cultivars adapted to diverse ranges is still obstructed by inadequate and incompetent seed production besides weak delivery systems, scanty seed policies, and excessive price of seed¹⁸⁷. There are many helpful adaptation strategies related to crop husbandry reported by scientists, comprising abiotic factors, for example, changing sowing and harvesting date, crop rotation, irrigation techniques, deficit irrigation, soil management practices, improvement of irrigation efficiency and variation in cropping patterns etc.^{5,14,25,188–191} (Table 6).

Raised bed planting method for maize and rye crop is useful to obtain maximum attainable grain yield. The most important approach is choosing the optimum sowing date for cereal crops under climate change. Changing sowing date to evade the harmful influences of high temperature at anthesis and during pollination and fertilization has been suggested as an adaptation tactic for climate warming. A change in sowing window, for example, earlier sowing of winter and spring cereal crops and delayed sowing of summer and autumn cereals to escape hot and dry periods in growing seasons, is one useful adaptation to climate uncertainty. Therefore, sowing of cereals has been adjusted according to optimum temperature conditions, resulting in substantial yield increases by escaping temperature stress during grain filling phase^{60,127}. Optimization of sowing time, planting density, and best irrigation techniques are dynamic strategies to confront weather-based stresses. Establishing innovative and precision irrigation water management might contribute to the mitigation of negative issues related to climate change. Better irrigation management can improve cereal crop productivity under climate warming. Water-saving interventions such as direct-planted rice, a modified system of rice intensification and alternate

wetting and drying (AWD) of rice were implemented in a cluster approach and resulted in enhanced water use efficiency^{225–229}. Optimum fertilization rates are also very effective to decrease the impact of global warming. Optimizing fertilizer use provides for better adaptability by enhancing energy capture by plants, by maintaining soil fertility and pesticide use, and increasing productivity^{230–233}. The significance of fertilizer in nourishing the world population is undeniable. Enhancing fertilizer use efficiency (minimizing losses like volatilization and leaching; ultimately nutrients availability to crops) is dynamic under climate change. Adaptation of integrated soil fertility management and precision nutrient management is helpful to minimize the negative influence of higher temperature^{221,234,235}. Laser land leveling (through enhancing water application efficiency, uniform distribution in fields and availability to crop plants; thus increasing water use efficiency) can enhance cereal grain yield under warming climate. Agroforestry techniques could help growers to diversify their crops, and their profits, along with enhancing climate resilience. Integrated pest management are useful approaches to cope with the thermal trend impact in cereal based cropping systems^{173,235,236}. Some important agronomic practices to cope with climate warming are described below in greater detail.

Sowing date. Adjustment of sowing date to accommodate the thermal trend is significant for proper phenology of cereal crops. Zhang et al.²³⁷ reported that delayed sowing might be an efficient way to cope with water stress. Climate warming accelerates crop growth and shortens duration of phenological phases of cereals. Physiological processes including photosynthesis, starch conversion and nutrient metabolism are negatively affected by changing phenology due to heat stress^{31,200,238–241}. The use of climate-smart cultivars, besides optimum planting dates, can reduce risk impacts due to warming^{149,242}. Hence, decisions about seed rate, planting density, spatial arrangement and other management practices including irrigation, nutrition and application of pesticides etc. are affected by planting date^{243–245}. However, to get real benefits sowing date should coincide with growth cycle to make the optimal use of possible environmental circumstances and managements^{149,202,203,246–248}.

Planting geometry. Optimum planting geometry can be employed for the reduction of harmful effects on phenology due to climate change. Plant phenology is also affected by planting geometry^{130,249}. Crop microclimate is changed by varying planting geometry like planting density, row spacing, seed rate etc. Management of planting geometry, e.g., decreased plant density, broader plant or row to row distance, and skip-row configurations are adaptation strategies that can be adapted in the dryland farming areas for better utilization of available soil-water²⁵⁰. Nevertheless, optimum planting geometry could improve resource use efficiencies in crops through proper crop phenology, producing more leaf area and ultimately more favorable yield components and yield in cereal crops in all climatic conditions^{251,252}. Planting geometry affects solar radiation interception, canopy coverage, crop growth rate and biomass accumulation^{215,253–258}. Furthermore, with optimum planting geometry, total dry matter production and ultimately grain yield increase was mainly due to more photo-assimilates²⁵². Planting practices have significant impact on maize productivity as concluded by Huang et al.²⁵⁹. They reported that soil temperature and growth period precipitation are determinant factors in dryland farming systems. Thus, they recommended that selection of suitable planting patterns according to the limiting factors can uplift dryland agriculture.

Supplementary irrigations. Supplemental irrigation is an important adaptation strategy for the reduction of harmful effects due to climate warming on cereal crops worldwide²⁶⁰. Supplemental irrigation is application of a limited amount of water at very critical stages for the improvement and stabilization of grain yield of cereal crops when rainfall is insufficient to provide adequate water for proper growth and development. Supplemental irrigation can minimize the impact of heat stress²⁶¹. During the previous decades, the main source of water was canal irrigation water, but nowadays, water shortage is severe, so supplemental irrigation can be helpful especially in arid environments. Supplemental irrigation, particularly during critical crop phenological stages and phases, can improve cereal crop yield as well as water efficiency in cereal based cropping systems²⁶². Supplementary irrigation practice is a simple, nevertheless, extremely effective practice that allows the farming community to grow and manage cereal crops by irrigating at the optimum time, without being at mercy of the unpredictable precipitation. Supplemental irrigation permits farmers to grow cereal crops during their optimum growing period, which can enhance grain yield and avoid crop exposure to lethal heat and drought stresses in warm areas, and frost in cooler areas worldwide²⁶³.

Plant breeding and genetic modifications. Plant breeding and genetics provide dynamic mechanisms for adaptation of cereal cropping systems to heat stress. The combination of molecular and conventional plant breeding and genetic methods can be helpful to recognize and develop eco-stable varieties with required genotype-environment combinations that will be beneficial in farming under changing climatic circumstances^{166,167,264–267}. The desirable characters like drought, and cold and heat stress resistance, resistance to pests, and ability to cope with water logging and salinity can be accomplished by integrating conventional, molecular and transgenic techniques^{169,170,268–270}. Zachariah et al.²⁷¹ recommended use of drought-resilient food crops to mitigate the immediate agrarian crisis. As according to their analysis, rainfall deficit is the primary cause of crop failure, as compared to rising temperatures. Comprehensive overview of the C5-MTase gene family members in wheat was presented by Gahlaut et al.²⁷². They identified 52 C5-MTases (cytosine-5 DNA methyltransferases) gene family members (4 sub-families i.e. CMTs (Chromomethylase), METs (Methyltransferase), DRMs (Domains Rearranged Methyltransferase) and DNMT2s (DNA methyltransferase homologue 2)) that could be used to develop drought/heat stress in wheat.

Molecular approaches, comprising the use of marker-trait associations and genome-wide selection, are providing opportunities for developing germplasm with tolerance to several abiotic and biotic stresses. The adverse

effects of heat stress can be mitigated by developing crops with enhanced thermo-tolerance. Boote et al.²⁷³ in their work concluded that peanut, soybean, pearl millet are more heat tolerant than sorghum, bean and chick-pea. Genetic diversity for climate change has previously been reported for cereals with the application of comparative biology, genetic diversity analysis, collaborative phenotyping and crop simulation modeling^{198,274,275}. Photoperiod insensitivity could be incorporated by breeders; accordingly seed-filling could start before onset of higher temperatures. The use of modern cereal crop varieties is dynamic adaptation strategy for obtaining higher grain yields under climate change²⁷⁴. Adoption of improved cultivars is an important approach to adapt to climate variability and change. Heat tolerant varieties for cereal crops are essential under higher temperature stress¹⁹⁸. Breeding and sowing of improved, resistant cultivars (against abiotic and biotic) are key strategies to adapt to climate change. Growing varieties with enhanced climate warming tolerance is important to maintain pollen viability, for example, grain number and grain protein^{163,275,276}. Late-maturing maize cultivars enhance grain yield due to longer grain filling duration. Improved cultivars with resistances and tolerances to abiotic and biotic stresses for cereal crops can increase resource use efficiency. Growing of genotypes requiring more GDDs can better resist thermal stress (Table 6). Drought tolerant varieties can increase radiation use efficiency under climate warming conditions. Avoiding drought stress by genetic improvements will have a major role for ensuring food security and reducing grower's exposure to drought-risk. Adapting cultivars with deep root-systems, mainly in areas that experience prolonged dry spells, could be a useful strategy under climate warming stress. Slow maturing cultivars also minimize the adverse effects of heat stress particularly at anthesis¹⁴⁴.

Biodiversity. Agricultural biodiversity is essential for adapting to climate change (i.e., multiple cropping vs. sole cropping; diversified/integrated farming vs. specialized farming). Growing different varieties of crop can be beneficial in tackling climate-warming risk^{277–279}. More diverse and longer-cycle crop rotations will need to combine sequences of annual row crops such as maize and soybean with close-drilled cereals, shallow-rooted with deep-rooted crops, summer crops with winter crops, and annuals with perennials in the same fields^{280–283}. Similarly, increasing diversity within crops may be a powerful way to reduce agricultural declines from climate change as concluded by Morales-Castilla et al.²⁸⁴. Multi cropping systems and crop rotation (including leguminous or green manuring crops in existing cereal-based cropping systems) are useful to combat the adverse influence of climate change. Resilience to unpredictable weather will also benefit from intercropping, with the creative arrangement of multiple interacting crop species to diversify the field and the landscape^{101,172,285,286}. Sloat et al.²⁸⁷ reported that warming impact on maize, wheat and rice could be moderated by migration of these crops over time and the expansion of irrigation. Multiple-cropping systems and strategies to integrate animals and crops will make more efficient use of natural resources and applied inputs; these include systems such as permaculture, agroforestry, alley cropping, intercropping and sowing C₄ crops compared to C₃ crops. Diverse cropping systems with spatial diversity, and adapted to specific fields, soil conditions, and unique agro-ecozones can minimize negative effects of heat stress²⁸⁸. Cereal crop production resilience can be achieved by planting diverse combinations of cropping schemes together in the same field, and economic resilience through production of a range of products that can be marketed by various channels^{215,253,289,290}. Multiple cropping systems can diversify the cropping systems. This comprised of sowing two or more cereal crops on the same field, either at the same time or one after another, are production intensification strategies^{215,253}. Such cropping systems have the benefits of decreasing the risk of whole crop failures, therefore ensuring a higher level of production stability for the farm community^{235,291}. Sufficient cropping systems, particularly in regions that are extremely influenced by the effects of climate warming, are professed as an important approach to adaptation. Grain yields in sequential cropping systems were higher than average grain yields in single cropping systems, suggesting that sequential cropping systems contributed towards reducing the negative impact of climate warming in comparison to single cropping systems^{83,292–294}. Resource use efficiency through crop diversification can further be improved by adapting advanced production technologies, physiological based resilience to climate change, crop rotations, improved water and nitrogen use efficiencies, crop modeling as well as planning before time through the use of forecasting skills to adapt to climate change under current and future climate scenarios^{163,196,286,290,295–314}.

Conclusion

This work has highlighted the negative impacts of climate warming on cereals crop phenology which will become frequent and severe in future. Climate warming is threatening global food security due to consequences like heat, drought, and salinity stresses. It has shown considerable impact on cereal production by shortening crop phenology. With climate warming, all phenological events occurred earlier and resulting phenological phases were shortened in various regions and countries across the globe, as a result of which plants got less time to assimilate CO₂, and this ultimately resulted in yield reduction. The earlier wheat stages were observed in northwest China, and delaying was recorded in north south regions in China and Pakistan. The reduction in phenological phases was also observed in northeast China, Pakistan, Spain, Australia, Argentina, Romania, Germany and Pakistan. Rice phenological stages and phases were affected worldwide. The rice stages were earlier in Punjab, Pakistan and India, and transplanting-to-maturity phases were reduced in Pakistan and China. Maize phenological stages were delayed in most countries. Phenological phases of maize were also shortened in Pakistan, China and USA. In Germany and in Lithuania, oat stages were earlier while delayed in Spain due to global warming. However, these phases were significantly reduced in Spain. Rye phenological stages were earlier, and phases were shortened in Poland and Germany.

Recommendations

This impact of climate warming can be mitigated by developing cereal crop cultivars well adapted to climate change, changes in management practices e.g. sowing dates zoning of crops and by increasing cereal crop biodiversity. Further research regarding crop plasticity is necessary to adapt cereal crops under a wide range of climatic conditions in order to avoid yield losses due to climate warming. Similarly, wider spectrum climate change adaptation options e.g. change in cropping patterns, use of early maturity and less water-consuming crop varieties, availability of stress tolerant crop varieties, modification in sowing date, crop insurance, implementation of location specific technologies, application of modern technology, intercropping, mixed cropping, socioeconomic and institutional interventions, crop modeling, whole farm modeling, integrated crop-livestock management and plantation of trees around the field are recommended to mitigate the impact of climate warming.

Received: 22 April 2020; Accepted: 6 October 2020

Published online: 22 October 2020

References

- Sarwar, M. H., Sarwar, M. F., Sarwar, M., Qadri, N. A. & Moghal, S. The importance of cereals (Poaceae: Gramineae) nutrition in human health: a review. *J. Cereals Oilseeds* **4**, 32–35 (2013).
- Ranjan, R. & Yadav, R. Targeting nitrogen use efficiency for sustained production of cereal crops. *J. Plant Nutr.* **42**, 1086–1113. <https://doi.org/10.1080/01904167.2019.1589497> (2019).
- Sofi, F. *et al.* Health and nutrition studies related to cereal biodiversity: a participatory multi-actor literature review approach. *Nutrients* **10**, 1207 (2018).
- Stewart, B. A. & Lal, R. In *Advances in Agronomy* Vol. 151 (ed Donald L. Sparks) 1–44 (Academic Press, 2018).
- Sadras, V. *et al.* In *Advances in Agronomy* Vol. 163 (ed Donald L. Sparks) 153–177 (Academic Press, 2020).
- Yu, S. & Tian, L. Breeding major cereal grains through the lens of nutrition sensitivity. *Mol. Plant* **11**, 23–30. <https://doi.org/10.1016/j.molp.2017.08.006> (2018).
- Wang, J., Vanga, S., Saxena, R., Orsat, V. & Raghavan, V. Effect of climate change on the yield of cereal crops: a review. *Climate* **6**, 41 (2018).
- Chmielewski, F.-M., Müller, A. & Bruns, E. Climate changes and trends in phenology of fruit trees and field crops in Germany, 1961–2000. *Agric. For. Meteorol.* **121**, 69–78. [https://doi.org/10.1016/S0168-1923\(03\)00161-8](https://doi.org/10.1016/S0168-1923(03)00161-8) (2004).
- Price, R. K. & Welch, R. W. In *Encyclopedia of Human Nutrition* (3rd Edn) (ed Benjamin Caballero) 307–316 (Academic Press, 2013).
- McKevith, B. Nutritional aspects of cereals. *Nutr. Bull.* **29**, 111–142. <https://doi.org/10.1111/j.1467-3010.2004.00418.x> (2004).
- Wang, H. L. *et al.* Phenological trends in winter wheat and spring cotton in response to climate changes in northwest China. *Agric. For. Meteorol.* **148**, 1242–1251. <https://doi.org/10.1016/j.agrformet.2008.03.003> (2008).
- Chen, X. *et al.* Does any phenological event defined by remote sensing deserve particular attention? An examination of spring phenology of winter wheat in Northern China. *Ecol. Ind.* **116**, 106456. <https://doi.org/10.1016/j.ecolind.2020.106456> (2020).
- Li, Y., Hou, R. & Tao, F. Interactive effects of different warming levels and tillage managements on winter wheat growth, physiological processes, grain yield and quality in the North China Plain. *Agr. Ecosyst. Environ.* **295**, 106923. <https://doi.org/10.1016/j.agee.2020.106923> (2020).
- Li, Z. *et al.* Response of maize phenology to climate warming in Northeast China between 1990 and 2012. *Reg. Environ. Change* **14**, 39–48. <https://doi.org/10.1007/s10113-013-0503-x> (2014).
- Rojas-Downing, M. M., Nejadhashemi, A. P., Harrigan, T. & Woznicki, S. A. Climate change and livestock: impacts, adaptation, and mitigation. *Clim. Risk Manag.* **16**, 145–163. <https://doi.org/10.1016/j.crm.2017.02.001> (2017).
- Akram, R. *et al.* In *Advances in Rice Research for Abiotic Stress Tolerance* (eds Hasanuzzaman, M. *et al.*) 69–85 (Woodhead Publishing, 2019).
- Farooq, M., Hussain, M., Wakeel, A. & Siddique, K. H. M. Salt stress in maize: effects, resistance mechanisms, and management. A review. *Agron. Sustain. Dev.* **35**, 461–481. <https://doi.org/10.1007/s13593-015-0287-0> (2015).
- Farooq, M., Hussain, M. & Siddique, K. H. M. Drought stress in wheat during flowering and grain-filling periods. *Crit. Rev. Plant Sci.* **33**, 331–349. <https://doi.org/10.1080/07352689.2014.875291> (2014).
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D. & Basra, S. M. A. Plant drought stress: effects, mechanisms and management. *Agron. Sustain. Dev.* **29**, 185–212. <https://doi.org/10.1051/agro:2008021> (2009).
- Hussain, J., Khaliq, T., Ahmad, A., Akhter, J. & Asseng, S. Wheat responses to climate change and its adaptations: a focus on arid and semi-arid environment. *Int. J. Environ. Res.* <https://doi.org/10.1007/s41742-018-0074-2> (2018).
- Hussain, M. *et al.* In *Advances in Agronomy* Vol. 148 (ed Donald L. Sparks) 231–287 (Academic Press, 2018).
- Dyer, G. A., López-Feldman, A., Yúnez-Naude, A. & Taylor, J. E. Genetic erosion in maize's center of origin. *Proc. Natl. Acad. Sci.* **111**, 14094–14099. <https://doi.org/10.1073/pnas.1407033111> (2014).
- Nicholson, S. E., Funk, C. & Fink, A. H. Rainfall over the African continent from the 19th through the 21st century. *Global Planet. Change* **165**, 114–127. <https://doi.org/10.1016/j.gloplacha.2017.12.014> (2018).
- Pour, S. H., Wahab, A. K. A. & Shahid, S. Spatiotemporal changes in aridity and the shift of drylands in Iran. *Atmos. Res.* **233**, 104704. <https://doi.org/10.1016/j.atmosres.2019.104704> (2020).
- Sadras, V. O. & Monzon, J. P. Modelled wheat phenology captures rising temperature trends: shortened time to flowering and maturity in Australia and Argentina. *Field Crops Res.* **99**, 136–146. <https://doi.org/10.1016/j.fcr.2006.04.003> (2006).
- Menzel, A. *et al.* European phenological response to climate change matches the warming pattern. *Glob. Change Biol.* **12**, 1969–1976. <https://doi.org/10.1111/j.1365-2486.2006.01193.x> (2006).
- Liu, Y., Qin, Y., Ge, Q., Dai, J. & Chen, Q. Responses and sensitivities of maize phenology to climate change from 1981 to 2009 in Henan Province, China. *J. Geogr. Sci.* **27**, 1072–1084. <https://doi.org/10.1007/s11442-017-1422-4> (2017).
- Liu, L. *et al.* Uncertainty in wheat phenology simulation induced by cultivar parameterization under climate warming. *Eur. J. Agron.* **94**, 46–53. <https://doi.org/10.1016/j.eja.2017.12.001> (2018).
- Ye, Z. *et al.* Impacts of 1.5°C and 2.0°C global warming above pre-industrial on potential winter wheat production of China. *Eur. J. Agron.* **120**, 126149. <https://doi.org/10.1016/j.eja.2020.126149> (2020).
- Kawakita, S., Takahashi, H. & Moriya, K. Prediction and parameter uncertainty for winter wheat phenology models depend on model and parameterization method differences. *Agric. For. Meteorol.* **290**, 107998. <https://doi.org/10.1016/j.agrformet.2020.107998> (2020).
- Ahmed, K., Shabbir, G., Ahmed, M. & Shah, K. N. Phenotyping for drought resistance in bread wheat using physiological and biochemical traits. *Sci. Total Environ.* **729**, 139082. <https://doi.org/10.1016/j.scitotenv.2020.139082> (2020).
- Ahmed, M., Aslam, M. A., Fayyaz-UI, H., Hayat, R. & Ahmad, S. Biochemical, physiological and agronomic response of wheat to changing climate of rainfed Pakistan. *Pak. J. Bot.* **51**, 535–551. [https://doi.org/10.30848/PJB2019-2\(10\)](https://doi.org/10.30848/PJB2019-2(10)) (2019).

33. Tamburino, L., Bravo, G., Clough, Y. & Nicholas, K. A. From population to production: 50 years of scientific literature on how to feed the world. *Global Food Secur.* **24**, 100346. <https://doi.org/10.1016/j.gfs.2019.100346> (2020).
34. Gomez-Zavaglia, A., Mejuto, J. C. & Simal-Gandara, J. Mitigation of emerging implications of climate change on food production systems. *Food Res. Int.* **134**, 109256. <https://doi.org/10.1016/j.foodres.2020.109256> (2020).
35. Wreford, A. & Topp, C. F. E. Impacts of climate change on livestock and possible adaptations: a case study of the United Kingdom. *Agric. Syst.* **178**, 102737. <https://doi.org/10.1016/j.agsy.2019.102737> (2020).
36. Hu, Q., Weiss, A., Feng, S. & Baenziger, P. S. Earlier winter wheat heading dates and warmer spring in the U.S. Great Plains. *Agric. For. Meteorol.* **135**, 284–290. <https://doi.org/10.1016/j.agrformet.2006.01.001> (2005).
37. He, L., Jin, N. & Yu, Q. Impacts of climate change and crop management practices on soybean phenology changes in China. *Sci. Total Environ.* **707**, 135638. <https://doi.org/10.1016/j.scitotenv.2019.135638> (2020).
38. Tao, F., Yokozawa, M., Liu, J. & Zhang, Z. Climate-crop yield relationships at provincial scales in China and the impacts of recent climate trends. *Climate Res.* **38**, 83–94 (2008).
39. Estrella, N., Sparks, T. H. & Menzel, A. Effects of temperature, phase type and timing, location, and human density on plant phenological responses in Europe. *Climate Res.* **39**, 235–248 (2009).
40. Ahmed, M., Fayyaz-ul-Hassan & Ahmad, S. In *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (eds Ahmed, M. & Claudio O. Stockle) 91–111 (Springer International Publishing, 2017).
41. García-Mozo, H., Mestre, A. & Galán, C. Phenological trends in southern Spain: a response to climate change. *Agric. For. Meteorol.* **150**, 575–580. <https://doi.org/10.1016/j.agrformet.2010.01.023> (2010).
42. Wu, D. *et al.* Measured phenology response of unchanged crop varieties to long-term historical climate change. *Int. J. Plant Prod.* **13**, 47–58. <https://doi.org/10.1007/s42106-018-0033-z> (2019).
43. Sujetovienė, G. *et al.* Climate-change-related long-term historical and projected changes to spring barley phenological development in Lithuania. *J. Agric. Sci.* **156**, 1061–1069. <https://doi.org/10.1017/S0021859618000904> (2019).
44. Ahmad, A. *et al.* In *Handbook of Climate Change and Agroecosystems* Vol. 3 (eds Rosenzweig, C. & Hillel, D.) 219–258 (World Scientific 2015).
45. Tsimba, R., Edmeades, G. O., Millner, J. P. & Kemp, P. D. The effect of planting date on maize: phenology, thermal time durations and growth rates in a cool temperate climate. *Field Crops Res.* **150**, 145–155. <https://doi.org/10.1016/j.fcr.2013.05.021> (2013).
46. Bai, H., Tao, F., Xiao, D., Liu, F. & Zhang, H. Attribution of yield change for rice-wheat rotation system in China to climate change, cultivars and agronomic management in the past three decades. *Clim. Change* **135**, 539–553. <https://doi.org/10.1007/s10584-015-1579-8> (2016).
47. Shim, D., Lee, K.-J. & Lee, B.-W. Response of phenology- and yield-related traits of maize to elevated temperature in a temperate region. *Crop J.* **5**, 305–316. <https://doi.org/10.1016/j.cj.2017.01.004> (2017).
48. Yang, J. *et al.* Yield-maturity relationships of summer maize from 2003 to 2017 in the Huanghuaihai plain of China. *Sci. Rep.* **9**, 11417. <https://doi.org/10.1038/s41598-019-47561-2> (2019).
49. Teller, A. S. Moving the conversation on climate change and inequality to the local: socio-ecological vulnerability in agricultural Tanzania. *Soc. Dev.* **2**, 25–50. <https://doi.org/10.1525/sod.2016.2.1.25> (2016).
50. Tao, F., Zhang, S. & Zhang, Z. Spatiotemporal changes of wheat phenology in China under the effects of temperature, day length and cultivar thermal characteristics. *Eur. J. Agron.* **43**, 201–212. <https://doi.org/10.1016/j.eja.2012.07.005> (2012).
51. Shi, W., Wang, M. & Liu, Y. Crop yield and production responses to climate disasters in China. *Sci. Total Environ.* **750**, 141147. <https://doi.org/10.1016/j.scitotenv.2020.141147> (2021).
52. Siebert, S. & Ewert, F. Spatio-temporal patterns of phenological development in Germany in relation to temperature and day length. *Agric. For. Meteorol.* **152**, 44–57. <https://doi.org/10.1016/j.agrformet.2011.08.007> (2012).
53. Porter, J. R. & Gawith, M. Temperatures and the growth and development of wheat: a review. *Eur. J. Agron.* **10**, 23–36. [https://doi.org/10.1016/S1161-0301\(98\)00047-1](https://doi.org/10.1016/S1161-0301(98)00047-1) (1999).
54. McMaster, G. S. & Wilhelm, W. W. Growing degree-days: one equation, two interpretations. *Agric. For. Meteorol.* **87**, 291–300. [https://doi.org/10.1016/S0168-1923\(97\)00027-0](https://doi.org/10.1016/S0168-1923(97)00027-0) (1997).
55. Cleland, E. E., Chuine, I., Menzel, A., Mooney, H. A. & Schwartz, M. D. Shifting plant phenology in response to global change. *Trends Ecol. Evol.* **22**, 357–365. <https://doi.org/10.1016/j.tree.2007.04.003> (2007).
56. Xiao, D. *et al.* Impact of warming climate and cultivar change on maize phenology in the last three decades in North China Plain. *Theor. Appl. Climatol.* **124**, 653–661. <https://doi.org/10.1007/s00704-015-1450-x> (2016).
57. Craufurd, P. Q. & Wheeler, T. R. Climate change and the flowering time of annual crops. *J. Exp. Bot.* **60**, 2529–2539. <https://doi.org/10.1093/jxb/erp196> (2009).
58. van Bussel, L. G. J., Ewert, F. & Leffelaar, P. A. Effects of data aggregation on simulations of crop phenology. *Agric. Ecosyst. Environ.* **142**, 75–84. <https://doi.org/10.1016/j.agee.2010.03.019> (2011).
59. van Oort, P. A. J., Zhang, T., de Vries, M. E., Heinemann, A. B. & Meinke, H. Correlation between temperature and phenology prediction error in rice (*Oryza sativa* L.). *Agric. For. Meteorol.* **151**, 1545–1555. <https://doi.org/10.1016/j.agrformet.2011.06.012> (2011).
60. Rezaei, E. E., Siebert, S., Hüging, H. & Ewert, F. Climate change effect on wheat phenology depends on cultivar change. *Sci. Rep.* **8**, 4891. <https://doi.org/10.1038/s41598-018-23101-2> (2018).
61. Croitoru, A.-E., Holobaca, I.-H., Lazar, C., Moldovan, F. & Imbroane, A. Air temperature trend and the impact on winter wheat phenology in Romania. *Clim. Change* **111**, 393–410. <https://doi.org/10.1007/s10584-011-0133-6> (2012).
62. Hussain, M., Shabir, G., Farooq, M., Jabran, K. & Farooq, S. Developmental and phenological responses of wheat to sowing dates. *Pak. J. Agri. Sci.* **49**, 459–468 (2012).
63. Zhang, S. & Tao, F. Modeling the response of rice phenology to climate change and variability in different climatic zones: comparisons of five models. *Eur. J. Agron.* **45**, 165–176. <https://doi.org/10.1016/j.eja.2012.10.005> (2013).
64. Yue, Y. *et al.* Prediction of maize growth stages based on deep learning. *Comput. Electron. Agric.* **172**, 105351. <https://doi.org/10.1016/j.compag.2020.105351> (2020).
65. Luo, Q. Temperature thresholds and crop production: a review. *Clim. Change* **109**, 583–598. <https://doi.org/10.1007/s10584-011-0028-6> (2011).
66. Ramirez-Villegas, J., Challinor, A. J., Thornton, P. K. & Jarvis, A. Implications of regional improvement in global climate models for agricultural impact research. *Environ. Res. Lett.* **8**, 024018. <https://doi.org/10.1088/1748-9326/8/2/024018> (2013).
67. Luo, Q., Bange, M. & Clancy, L. Cotton crop phenology in a new temperature regime. *Ecol. Model.* **285**, 22–29. <https://doi.org/10.1016/j.ecolmodel.2014.04.018> (2014).
68. Pulatov, B., Linderson, M.-L., Hall, K. & Jönsson, A. M. Modeling climate change impact on potato crop phenology, and risk of frost damage and heat stress in northern Europe. *Agric. For. Meteorol.* **214–215**, 281–292. <https://doi.org/10.1016/j.agrformet.2015.08.266> (2015).
69. Hatfield, J. L. & Dold, C. Climate change impacts on corn phenology and productivity. *Corn: Production and Human Health in Changing Climate*, 95 (2018).
70. Zhang, T., Huang, Y. & Yang, X. Climate warming over the past three decades has shortened rice growth duration in China and cultivar shifts have further accelerated the process for late rice. *Glob. Change Biol.* **19**, 563–570. <https://doi.org/10.1111/gcb.12057> (2013).
71. Lin, Y. *et al.* Potential impacts of climate change and adaptation on maize in northeast China. *Agron. J.* **109**, 1476–1490 (2017).

72. Wang, N. *et al.* Modelling maize phenology, biomass growth and yield under contrasting temperature conditions. *Agric. For. Meteorol.* **250–251**, 319–329. <https://doi.org/10.1016/j.agrformet.2018.01.005> (2018).
73. Ge, Q., Wang, H., Rutishauser, T. & Dai, J. Phenological response to climate change in China: a meta-analysis. *Glob. Change Biol.* **21**, 265–274 (2015).
74. Srivastava, R. K., Panda, R. K. & Chakraborty, A. Assessment of climate change impact on maize yield and yield attributes under different climate change scenarios in eastern India. *Ecol. Ind.* **120**, 106881. <https://doi.org/10.1016/j.ecolind.2020.106881> (2021).
75. Gordo, O. & Sanz, J. J. Phenology and climate change: a long-term study in a Mediterranean locality. *Oecologia* **146**, 484–495. <https://doi.org/10.1007/s00442-005-0240-z> (2005).
76. Brown, M. E., de Beurs, K. M. & Marshall, M. Global phenological response to climate change in crop areas using satellite remote sensing of vegetation, humidity and temperature over 26 years. *Remote Sens. Environ.* **126**, 174–183. <https://doi.org/10.1016/j.rse.2012.08.009> (2012).
77. Kim, Y.-U. & Lee, B.-W. Earlier planting offsets the adverse effect of global warming on spring potato in South Korea. *Sci. Total Environ.* **742**, 140667. <https://doi.org/10.1016/j.scitotenv.2020.140667> (2020).
78. Baum, M. E., Licht, M. A., Huber, I. & Archontoulis, S. V. Impacts of climate change on the optimum planting date of different maize cultivars in the central US Corn Belt. *Eur. J. Agron.* **119**, 126101. <https://doi.org/10.1016/j.eja.2020.126101> (2020).
79. Menzel, A. *et al.* Climate change fingerprints in recent European plant phenology. *Glob. Change Biol.* **26**, 2599–2612. <https://doi.org/10.1111/gcb.15000> (2020).
80. Menzel, A. Trends in phenological phases in Europe between 1951 and 1996. *Int. J. Biometeorol.* **44**, 76–81. <https://doi.org/10.1007/s004840000054> (2000).
81. Oteros, J., García-Mozo, H., Botey, R., Mestre, A. & Galán, C. Variations in cereal crop phenology in Spain over the last twenty-six years (1986–2012). *Clim. Change* **130**, 545–558. <https://doi.org/10.1007/s10584-015-1363-9> (2015).
82. Wang, Y., Luo, Y. & Shafeeque, M. Interpretation of vegetation phenology changes using daytime and night-time temperatures across the Yellow River Basin, China. *Sci. Total Environ.* **693**, 133553. <https://doi.org/10.1016/j.scitotenv.2019.07.359> (2019).
83. Moriondo, M. & Bindi, M. Impact of climate change on the phenology of typical Mediterranean crops. *Italian J. Agrometeorol.* **3**, 5–12 (2007).
84. Anwar, M. R. *et al.* Climate change impacts on phenology and yields of five broadacre crops at four climatologically distinct locations in Australia. *Agric. Syst.* **132**, 133–144. <https://doi.org/10.1016/j.agry.2014.09.010> (2015).
85. Jackson, R. B. *et al.* Global energy growth is outpacing decarbonization. *Environ. Res. Lett.* **13**, 120401. <https://doi.org/10.1088/1748-9326/aaf303> (2018).
86. Le Quéré, C. *et al.* Global carbon budget 2018. *Earth Syst. Sci. Data* **10**, 2141–2194. <https://doi.org/10.5194/essd-10-2141-2018> (2018).
87. Balch, J. K. *et al.* Global combustion: the connection between fossil fuel and biomass burning emissions (1997–2010). *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150177. <https://doi.org/10.1098/rstb.2015.0177> (2016).
88. Ahmad, S. *et al.* Climate warming and management impact on the change of phenology of the rice-wheat cropping system in Punjab, Pakistan. *Field Crops Res.* **230**, 46–61. <https://doi.org/10.1016/j.fcr.2018.10.008> (2019).
89. Ahmad, S. *et al.* Application of DSSAT Model for sowing date management of C 4 summer cereals for fodder and grain crops under irrigated arid environment. *Pakistan J. Life Soc. Sci.* **14** (2016).
90. Hatfield, J. L. & Prueger, J. H. Temperature extremes: effect on plant growth and development. *Weather Clim. Extremes* **10**, 4–10. <https://doi.org/10.1016/j.wace.2015.08.001> (2015).
91. Aslam, M. A. *et al.* Can growing degree days and photoperiod predict spring wheat phenology?. *Front. Environ. Sci.* <https://doi.org/10.3389/fenvs.2017.00057> (2017).
92. Zhang, L., Zhang, Z., Luo, Y., Cao, J. & Li, Z. Optimizing genotype-environment-management interactions for maize farmers to adapt to climate change in different agro-ecological zones across China. *Sci. Total Environ.* **728**, 138614. <https://doi.org/10.1016/j.scitotenv.2020.138614> (2020).
93. Tryjanowski, P. *et al.* Changing phenology of potato and of the treatment for its major pest (colorado potato beetle)—a long-term analysis. *Am. J. Potato Res.* <https://doi.org/10.1007/s12230-017-9611-3> (2017).
94. Ahmad, S. *et al.* Quantification of climate warming and crop management impacts on cotton phenology. *Plants (Basel)* **6**, 7. <https://doi.org/10.3390/plants6010007> (2017).
95. Huang, J. & Ji, F. Effects of climate change on phenological trends and seed cotton yields in oasis of arid regions. *Int. J. Biometeorol.* **59**, 877–888. <https://doi.org/10.1007/s00484-014-0904-7> (2015).
96. Wang, Z. *et al.* Response of cotton phenology to climate change on the North China Plain from 1981 to 2012. *Sci. Rep.* **7**, 6628. <https://doi.org/10.1038/s41598-017-07056-4> (2017).
97. Srivastava, A., NareshKumar, S. & Aggarwal, P. K. Assessment on vulnerability of sorghum to climate change in India. *Agric. Ecosyst. Environ.* **138**, 160–169. <https://doi.org/10.1016/j.agee.2010.04.012> (2010).
98. Sultan, B. *et al.* Robust features of future climate change impacts on sorghum yields in West Africa. *Environ. Res. Lett.* **9**, 104006. <https://doi.org/10.1088/1748-9326/9/10/104006> (2014).
99. Shew, A. M., Tack, J. B., Nalley, L. L. & Chaminuka, P. Yield reduction under climate warming varies among wheat cultivars in South Africa. *Nat. Commun.* **11**, 4408. <https://doi.org/10.1038/s41467-020-18317-8> (2020).
100. Sonkar, G. *et al.* Vulnerability of Indian wheat against rising temperature and aerosols. *Environ. Pollut.* **254**, 112946. <https://doi.org/10.1016/j.envpol.2019.07.114> (2019).
101. Fatima, Z. *et al.* Quantification of climate warming and crop management impacts on phenology of pulses-based cropping systems. *Int. J. Plant Prod.* <https://doi.org/10.1007/s42106-020-00112-6> (2020).
102. Liu, Y., Qin, Y., Wang, H., Lv, S. & Ge, Q. Trends in maize (*Zea mays* L.) phenology and sensitivity to climate factors in China from 1981 to 2010. *Int. J. Biometeorol.* **64**, 461–470. <https://doi.org/10.1007/s00484-019-01832-9> (2020).
103. Zhou, X. *et al.* Legacy effect of spring phenology on vegetation growth in temperate China. *Agric. For. Meteorol.* **281**, 107845. <https://doi.org/10.1016/j.agrformet.2019.107845> (2020).
104. Das, S., Kumar, A., Barman, M., Pal, S. & Bandopadhyay, P. In *Agronomic Crops: Volume 3: Stress Responses and Tolerance* (ed. Hasanuzzaman, M.) 13–28 (Springer Singapore, 2020).
105. Xiao, D., Liu, D. L., Wang, B., Feng, P. & Waters, C. Designing high-yielding maize ideotypes to adapt changing climate in the North China Plain. *Agric. Syst.* **181**, 102805. <https://doi.org/10.1016/j.agry.2020.102805> (2020).
106. Liu, Y. *et al.* Impacts of 1.5 and 2.0°C global warming on rice production across China. *Agric. For. Meteorol.* **284**, 107900. <https://doi.org/10.1016/j.agrformet.2020.107900> (2020).
107. Rani, B. A. & Maragatham, N. Effect of elevated temperature on rice phenology and yield. *Indian J. Sci. Technol.* **6**, 5095–5097 (2013).
108. Shrestha, S. *et al.* Phenological responses of upland rice grown along an altitudinal gradient. *Environ. Exp. Bot.* **89**, 1–10. <https://doi.org/10.1016/j.envexpbot.2012.12.007> (2013).
109. Abbas, G. *et al.* Quantification of the impacts of climate change and crop management on phenology of maize-based cropping system in Punjab, Pakistan. *Agric. For. Meteorol.* **247**, 42–55. <https://doi.org/10.1016/j.agrformet.2017.07.012> (2017).
110. Estrella, N., Sparks, T. H. & Menzel, A. Trends and temperature response in the phenology of crops in Germany. *Glob. Change Biol.* **13**, 1737–1747. <https://doi.org/10.1111/j.1365-2486.2007.01374.x> (2007).

111. Li, K. *et al.* Effects of changing climate and cultivar on the phenology and yield of winter wheat in the North China Plain. *Int. J. Biometeorol.* **60**, 21–32. <https://doi.org/10.1007/s00484-015-1002-1> (2016).
112. He, D. *et al.* Uncertainty in canola phenology modelling induced by cultivar parameterization and its impact on simulated yield. *Agric. For. Meteorol.* **232**, 163–175. <https://doi.org/10.1016/j.agrformet.2016.08.013> (2017).
113. Wei, W., Wu, W., Li, Z., Yang, P. & Zhou, Q. Selecting the optimal ndvi time-series reconstruction technique for crop phenology detection. *Intell. Autom. Soft Comput.* **22**, 237–247. <https://doi.org/10.1080/10798587.2015.1095482> (2016).
114. Chakraborty, A., Das, P. K., Sai, M. V. R. S. & Behera, G. Spatial pattern of temporal trend of crop phenology matrices over india using timeseries gimms NDVI data (1982‐2006). *ISPRS Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.* **3820**, 113–118 (2012).
115. Komal, C., Shi, W., Boori, M. S. & Corgne, S. Agriculture phenology monitoring using NDVI time series based on remote sensing satellites: a case study of Guangdong, China. *Opt. Mem. Neural Netw.* **28**, 204–214. <https://doi.org/10.3103/S1060992X19030093> (2019).
116. Liu, Y., Chen, Q., Ge, Q. & Dai, J. Spatiotemporal differentiation of changes in wheat phenology in China under climate change from 1981 to 2010. *Sci. China Earth Sci.* **61**, 1088–1097. <https://doi.org/10.1007/s11430-017-9149-0> (2018).
117. Xiao, D. *et al.* Observed changes in winter wheat phenology in the North China Plain for 1981–2009. *Int. J. Biometeorol.* **57**, 275–285. <https://doi.org/10.1007/s00484-012-0552-8> (2013).
118. Hossain, A., da Silva Teixeira, J. A., Lozovskaya, M. V. & Zvolinsky, V. P. High temperature combined with drought affect rainfed spring wheat and barley in South-Eastern Russia: I. Phenology and growth. *Saudi J. Biol. Sci.* **19**, 473–487. <https://doi.org/10.1016/j.sjbs.2012.07.005> (2012).
119. Martínez-Núñez, M. *et al.* The phenological growth stages of different amaranth species grown in restricted spaces based in BBCH code. *South African J. Bot.* **124**, 436–443. <https://doi.org/10.1016/j.sajb.2019.05.035> (2019).
120. Zhao, C. *et al.* Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci.* **114**, 9326–9331. <https://doi.org/10.1073/pnas.1701762114> (2017).
121. Li, T. *et al.* Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions. *Glob. Change Biol.* **21**, 1328–1341. <https://doi.org/10.1111/gcb.12758> (2015).
122. Subash, N. & Ram Mohan, H. S. Evaluation of the impact of climatic trends and variability in rice–wheat system productivity using Cropping System Model DSSAT over the Indo-Gangetic Plains of India. *Agric. For. Meteorol.* **164**, 71–81. <https://doi.org/10.1016/j.agrformet.2012.05.008> (2012).
123. Tian, D. *et al.* Does decadal climate variation influence wheat and maize production in the southeast USA?. *Agric. For. Meteorol.* **204**, 1–9. <https://doi.org/10.1016/j.agrformet.2015.01.013> (2015).
124. Quiring, S. M. & Legates, D. R. Application of CERES–Maize for within-season prediction of rainfed corn yields in Delaware, USA. *Agric. For. Meteorol.* **148**, 964–975. <https://doi.org/10.1016/j.agrformet.2008.01.009> (2008).
125. Bleharczyk, A., Sawinska, Z., Malecka, I., Sparks, T. H. & Tryjanowski, P. The phenology of winter rye in Poland: an analysis of long-term experimental data. *Int. J. Biometeorol.* **60**, 1341–1346. <https://doi.org/10.1007/s00484-015-1127-2> (2016).
126. Luo, Q., O’Leary, G., Cleverly, J. & Eamus, D. Effectiveness of time of sowing and cultivar choice for managing climate change: wheat crop phenology and water use efficiency. *Int. J. Biometeorol.* **62**, 1049–1061. <https://doi.org/10.1007/s00484-018-1508-4> (2018).
127. He, L. *et al.* Impacts of recent climate warming, cultivar changes, and crop management on winter wheat phenology across the Loess Plateau of China. *Agric. For. Meteorol.* **200**, 135–143. <https://doi.org/10.1016/j.agrformet.2014.09.011> (2015).
128. Wang, J., Wang, E., Feng, L., Yin, H. & Yu, W. Phenological trends of winter wheat in response to varietal and temperature changes in the North China Plain. *Field Crops Res.* **144**, 135–144. <https://doi.org/10.1016/j.fcr.2012.12.020> (2013).
129. Zheng, Z., Cai, H., Wang, Z. & Wang, X. Simulation of climate change impacts on phenology and production of winter wheat in Northwestern China using CERES-wheat model. *Atmosphere* **11**, 681 (2020).
130. Hyles, J., Bloomfield, M. T., Hunt, J. R., Trethowan, R. M. & Trevaskis, B. Phenology and related traits for wheat adaptation. *Heredity* <https://doi.org/10.1038/s41437-020-0320-1> (2020).
131. Li, Q.-Y. *et al.* Determination of optimum growing degree-days (GDD) range before winter for wheat cultivars with different growth characteristics in North China Plain. *J. Integr. Agric.* **11**, 405–415. [https://doi.org/10.1016/S2095-3119\(12\)60025-2](https://doi.org/10.1016/S2095-3119(12)60025-2) (2012).
132. Herndl, M., Shan, C.-G., Wang, P., Graeff, S. & Claupein, W. A model based ideotyping approach for wheat under different environmental conditions in North China Plain. *Agric. Sci. China* **6**, 1426–1436. [https://doi.org/10.1016/S1671-2927\(08\)60004-8](https://doi.org/10.1016/S1671-2927(08)60004-8) (2007).
133. Asseng, S., Turner, N. C., Ray, J. D. & Keating, B. A. A simulation analysis that predicts the influence of physiological traits on the potential yield of wheat. *Eur. J. Agron.* **17**, 123–141. [https://doi.org/10.1016/S1161-0301\(01\)00149-6](https://doi.org/10.1016/S1161-0301(01)00149-6) (2002).
134. Rezaei, E. E., Siebert, S. & Ewert, F. Intensity of heat stress in winter wheat—phenology compensates for the adverse effect of global warming. *Environ. Res. Lett.* **10**, 024012 (2015).
135. Eyshi Rezaei, E., Siebert, S. & Ewert, F. Climate and management interaction cause diverse crop phenology trends. *Agric. For. Meteorol.* **233**, 55–70. <https://doi.org/10.1016/j.agrformet.2016.11.003> (2017).
136. Wang, Y., Zhang, J., Song, G., Long, Z. & Chen, C. Impacts of recent temperatures rise on double-rice phenology across Southern China. *Int. J. Plant Prod.* **13**, 1–10. <https://doi.org/10.1007/s42106-018-0029-8> (2019).
137. Hu, X., Huang, Y., Sun, W. & Yu, L. Shifts in cultivar and planting date have regulated rice growth duration under climate warming in China since the early 1980s. *Agric. For. Meteorol.* **247**, 34–41. <https://doi.org/10.1016/j.agrformet.2017.07.014> (2017).
138. Zhang, S., Tao, F. & Zhang, Z. Rice reproductive growth duration increased despite of negative impacts of climate warming across China during 1981–2009. *Eur. J. Agron.* **54**, 70–83. <https://doi.org/10.1016/j.eja.2013.12.001> (2014).
139. Bai, H. & Xiao, D. Spatiotemporal changes of rice phenology in China during 1981–2010. *Theor Appl Climatol* **140**, 1483–1494. <https://doi.org/10.1007/s00704-020-03182-8> (2020).
140. Abbas, G. *et al.* Sowing date and hybrid choice matters production of maize–maize system. *Int. J. Plant Prod.* <https://doi.org/10.1007/s42106-020-00104-6> (2020).
141. Abbas, G. *et al.* Nitrogen rate and hybrid selection matters productivity of maize–maize cropping system under irrigated arid environment of Southern Punjab, Pakistan. *Int. J. Plant Prod.* **14**, 309–320. <https://doi.org/10.1007/s42106-020-00086-5> (2020).
142. Xiao, D., Zhao, Y., Bai, H., Hu, Y. & Cao, J. Impacts of climate warming and crop management on maize phenology in northern China. *J. Arid Land* **11**, 892–903. <https://doi.org/10.1007/s40333-019-0028-3> (2019).
143. Wang, Z. *et al.* Effects of climate change and cultivar on summer maize phenology. *International Journal of Plant Production* **10**, 509–525 (2016).
144. Sacks, W. J. & Kucharik, C. J. Crop management and phenology trends in the U.S. Corn Belt: Impacts on yields, evapotranspiration and energy balance. *Agricultural and Forest Meteorology* **151**, 882–894. <https://doi.org/10.1016/j.agrformet.2011.02.010> (2011).
145. Mo, F. *et al.* Phenological responses of spring wheat and maize to changes in crop management and rising temperatures from 1992 to 2013 across the Loess Plateau. *Field Crops Research* **196**, 337–347. <https://doi.org/10.1016/j.fcr.2016.06.024> (2016).
146. Wang, P. *et al.* Summer maize growth under different precipitation years in the Huang-Huai-Hai Plain of China. *Agric. For. Meteorol.* **285–286**, 107927. <https://doi.org/10.1016/j.agrformet.2020.107927> (2020).
147. Liu, Y., Qin, Y. & Ge, Q. Spatiotemporal differentiation of changes in maize phenology in China from 1981 to 2010. *J. Geog. Sci.* **29**, 351–362. <https://doi.org/10.1007/s11442-019-1602-5> (2019).

148. Chen, C. *et al.* Global warming and shifts in cropping systems together reduce China's rice production. *Global Food Security* **24**, 100359. <https://doi.org/10.1016/j.gfs.2020.100359> (2020).
149. Lv, Z., Li, F. & Lu, G. Adjusting sowing date and cultivar shift improve maize adaption to climate change in China. *Mitig Adapt Strat Glob Change* **25**, 87–106. <https://doi.org/10.1007/s11027-019-09861-w> (2020).
150. Nahar, K., Ahamed, K. U. & Fujita, M. Phenological variation and its relation with yield in several wheat (*Triticum aestivum* L.) cultivars under normal and late sowing mediated heat stress condition. *Notulae Scientia Biologicae* **2**, 51–56 (2010).
151. Raoufi, R. S. & Soufizadeh, S. Simulation of the impacts of climate change on phenology, growth, and yield of various rice genotypes in humid sub-tropical environments using AquaCrop-Rice. *Int J Biometeorol* <https://doi.org/10.1007/s00484-020-01946-5> (2020).
152. Karapinar, B. & Özertan, G. Yield implications of date and cultivar adaptation to wheat phenological shifts: a survey of farmers in Turkey. *Climatic Change* **158**, 453–472. <https://doi.org/10.1007/s10584-019-02532-4> (2020).
153. Ahmad, M. J., Iqbal, M. A. & Choi, K. S. Climate-driven constraints in sustaining future wheat yield and water productivity. *Agric. Water Manag.* **231**, 105991. <https://doi.org/10.1016/j.agwat.2019.105991> (2020).
154. Zacharias, M., Singh, S., Naresh Kumar, S., Harit, R. & Aggarwal, P. Impact of elevated temperature at different phenological stages on the growth and yield of wheat and rice. *Ind J Plant Physiol.* **15**, 350 (2010).
155. Sadok, W. & Jagadish, S. V. K. The Hidden Costs of Nighttime Warming on Yields. *Trends Plant Sci.* **25**, 644–651. <https://doi.org/10.1016/j.tplants.2020.02.003> (2020).
156. Kahiluoto, H. *et al.* Decline in climate resilience of European wheat. *Proc. Natl. Acad. Sci.* **116**, 123–128. <https://doi.org/10.1073/pnas.1804387115> (2019).
157. Lavee, H., Imeson, A. C. & Sarah, P. The impact of climate change on geomorphology and desertification along a mediterranean-arid transect. *Land Degrad. Dev.* **9**, 407–422. [https://doi.org/10.1002/\(sici\)1099-145x\(199809/10\)9:5%3c407::aid-ldr302%3e3.0.co;2-6](https://doi.org/10.1002/(sici)1099-145x(199809/10)9:5%3c407::aid-ldr302%3e3.0.co;2-6) (1998).
158. Traill, L. W. *et al.* Managing for change: wetland transitions under sea-level rise and outcomes for threatened species. *Divers. Distrib.* **17**, 1225–1233. <https://doi.org/10.1111/j.1472-4642.2011.00807.x> (2011).
159. Bellard, C., Leclerc, C. & Courchamp, F. Impact of sea level rise on the 10 insular biodiversity hotspots. *Glob. Ecol. Biogeogr.* **23**, 203–212. <https://doi.org/10.1111/geb.12093> (2014).
160. Feng, Q., Ma, H., Jiang, X., Wang, X. & Cao, S. What Has Caused Desertification in China?. *Scientific Reports* **5**, 15998. <https://doi.org/10.1038/srep15998> (2015).
161. Huang, J., Yu, H., Guan, X., Wang, G. & Guo, R. Accelerated dryland expansion under climate change. *Nature Climate Change* **6**, 166–171. <https://doi.org/10.1038/nclimate2837> (2016).
162. Liu, B. *et al.* Global wheat production with 1.5 and 2.0°C above pre-industrial warming. *Global Change Biology* **25**, 1428–1444. <https://doi.org/10.1111/gcb.14542> (2019).
163. Asseng, S. *et al.* Climate change impact and adaptation for wheat protein. *Glob. Change Biol.* **25**, 155–173. <https://doi.org/10.1111/gcb.14481> (2019).
164. Nayak, D. *et al.* Management opportunities to mitigate greenhouse gas emissions from Chinese agriculture. *Agr. Ecosyst. Environ.* **209**, 108–124. <https://doi.org/10.1016/j.agee.2015.04.035> (2015).
165. Kang, Y., Khan, S. & Ma, X. Climate change impacts on crop yield, crop water productivity and food security – A review. *Prog. Nat. Sci.* **19**, 1665–1674. <https://doi.org/10.1016/j.pnsc.2009.08.001> (2009).
166. Korres, N. E. *et al.* Cultivars to face climate change effects on crops and weeds: a review. *Agron. Sustain. Dev.* **36**, 12. <https://doi.org/10.1007/s13593-016-0350-5> (2016).
167. Ortiz, R. in *Food Security and Climate Change* 145–158 (2018).
168. Kumar, S. & Sidana, B. K. Farmers' perceptions and adaptation strategies to climate change in Punjab agriculture. *Indian J. Agric. Sci.* **88**, 1573–1581 (2018).
169. Burke, M. & Emerick, K. Adaptation to climate change: Evidence from US agriculture. *American Economic Journal: Economic Policy* **8**, 106–140 (2016).
170. Pradhan, A., Chan, C., Roul, P. K., Halbrendt, J. & Sipes, B. Potential of conservation agriculture (CA) for climate change adaptation and food security under rainfed uplands of India: A transdisciplinary approach. *Agric. Syst.* **163**, 27–35. <https://doi.org/10.1016/j.agsy.2017.01.002> (2018).
171. Bahri, H., Annabi, M., Cheikh M'Hamed, H. & Frija, A. Assessing the long-term impact of conservation agriculture on wheat-based systems in Tunisia using APSIM simulations under a climate change context. *Science of The Total Environment* **692**, 1223–1233. <https://doi.org/10.1016/j.scitotenv.2019.07.307> (2019).
172. Zampieri, M. *et al.* Estimating resilience of crop production systems: From theory to practice. *Sci. Total Environ.* **735**, 139378. <https://doi.org/10.1016/j.scitotenv.2020.139378> (2020).
173. Wiebe, K., Robinson, S. & Cattaneo, A. in *Sustainable Food and Agriculture* (eds Clayton Campanhola & Shivaji Pandey) 55–74 (Academic Press, 2019).
174. Abebe, T., Guenzi, A. C., Martin, B. & Cushman, J. C. Tolerance of mannitol-accumulating transgenic wheat to water stress and salinity. *Plant Physiol.* **131**, 1748–1755 (2003).
175. Lee, S.-H. *et al.* Identification and functional characterization of Siberian wild rye (*Elymus sibiricus* L.) small heat shock protein 16.9 gene (EsHsp16.9) conferring diverse stress tolerance in prokaryotic cells. *Biotechnology Letters* **37**, 881–890. <https://doi.org/10.1007/s10529-014-1749-1> (2015).
176. Yamakawa, H., Hirose, T., Kuroda, M. & Yamaguchi, T. Comprehensive Expression Profiling of Rice Grain Filling-Related Genes under High Temperature Using DNA Microarray. *Plant Physiol.* **144**, 258–277. <https://doi.org/10.1104/pp.107.098665> (2007).
177. Lehmann, N., Finger, R., Klein, T., Calanca, P. & Walter, A. Adapting crop management practices to climate change: Modeling optimal solutions at the field scale. *Agric. Syst.* **117**, 55–65. <https://doi.org/10.1016/j.agsy.2012.12.011> (2013).
178. Pimentel, A. J. B. *et al.* Characterization of heat tolerance in wheat cultivars and effects on production components. *Revista Ceres* **62**, 191–198 (2015).
179. Lin, B. B. Resilience in Agriculture through Crop Diversification: Adaptive Management for Environmental Change. *Bioscience* **61**, 183–193. <https://doi.org/10.1525/bio.2011.61.3.4> (2011).
180. Deryng, D., Sacks, W. J., Barford, C. C. & Ramankutty, N. Simulating the effects of climate and agricultural management practices on global crop yield. *Global Biogeochemical Cycles* **25**, n/a–n/a. <https://doi.org/10.1029/2009GB003765> (2011).
181. Mase, A. S., Gramig, B. M. & Prokopy, L. S. Climate change beliefs, risk perceptions, and adaptation behavior among Midwestern U.S. crop farmers. *Climate Risk Management* **15**, 8–17. <https://doi.org/10.1016/j.crm.2016.11.004> (2017).
182. Hernandez-Ochoa, I. M. *et al.* Adapting irrigated and rainfed wheat to climate change in semi-arid environments: Management, breeding options and land use change. *Eur. J. Agron.* **109**, 125915. <https://doi.org/10.1016/j.eja.2019.125915> (2019).
183. Meza, F. J. & Silva, D. Dynamic adaptation of maize and wheat production to climate change. *Climatic Change* **94**, 143–156. <https://doi.org/10.1007/s10584-009-9544-z> (2009).
184. Luo, Q., Bellotti, W., Williams, M. & Wang, E. Adaptation to climate change of wheat growing in South Australia: Analysis of management and breeding strategies. *Agr. Ecosyst. Environ.* **129**, 261–267. <https://doi.org/10.1016/j.agee.2008.09.010> (2009).
185. Wang, B. *et al.* Designing wheat ideotypes to cope with future changing climate in South-Eastern Australia. *Agric. Syst.* **170**, 9–18. <https://doi.org/10.1016/j.agsy.2018.12.005> (2019).

186. Li, Y. *et al.* Quantifying irrigation cooling benefits to maize yield in the US Midwest. *Glob. Change Biol.* **26**, 3065–3078. <https://doi.org/10.1111/gcb.15002> (2020).
187. Hampton, K. N. Persistent and pervasive community: New communication technologies and the future of community. *Am. Behav. Sci.* **60**, 101–124 (2016).
188. Asseng, S., Zhu, Y., Wang, E. & Zhang, W. in *Crop Physiology (Second Edition)* (ed Victor O. Sadras/Daniel F. Calderini) 505–546 (Academic Press, 2015).
189. Sadras, V. O., Vadez, V., Purushothaman, R., Lake, L. & Marrou, H. Unscrambling confounded effects of sowing date trials to screen for crop adaptation to high temperature. *Field Crops Research* **177**, 1–8. <https://doi.org/10.1016/j.fcr.2015.02.024> (2015).
190. Rodriguez, D. & Sadras, V. Opportunities from integrative approaches in farming systems design. *Field Crops Research* **124**, 131–141 (2011).
191. Cabezas, J. M. *et al.* Identifying adaptation strategies to climate change for Mediterranean olive orchards using impact response surfaces. *Agric. Syst.* **185**, 102937. <https://doi.org/10.1016/j.agry.2020.102937> (2020).
192. Ahmad, I., Ahmad, B., Boote, K. & Hoogenboom, G. Adaptation strategies for maize production under climate change for semi-arid environments. *Eur. J. Agron.* **115**, 126040. <https://doi.org/10.1016/j.eja.2020.126040> (2020).
193. Abbas, G. *et al.* in *Cotton Production and Uses: Agronomy, Crop Protection, and Postharvest Technologies* (eds Shakeel Ahmad & Mirza Hasanuzzaman) 429–445 (Springer Singapore, 2020).
194. Ahmed, M. *et al.* Novel multimodel ensemble approach to evaluate the sole effect of elevated CO₂ on winter wheat productivity. *Scientific Reports* **9**, 7813. <https://doi.org/10.1038/s41598-019-44251-x> (2019).
195. Ahmed, M. & Ahmad, S. in *Agronomic Crops: Volume 2: Management Practices* (ed Mirza Hasanuzzaman) 31–46 (Springer Singapore, 2019).
196. van Ogtrop, F., Ahmad, M. & Moeller, C. Principal components of sea surface temperatures as predictors of seasonal rainfall in rainfed wheat growing areas of Pakistan. *Meteorological Applications* **21**, 431–443. <https://doi.org/10.1002/met.1429> (2014).
197. Abedinpour, M. *et al.* Performance evaluation of AquaCrop model for maize crop in a semi-arid environment. *Agric. Water Manag.* **110**, 55–66. <https://doi.org/10.1016/j.agwat.2012.04.001> (2012).
198. Sareen, S. *et al.* Molecular genetic diversity analysis for heat tolerance of indigenous and exotic wheat genotypes. *J. Plant Biochem. Biotechnol.* **29**, 15–23. <https://doi.org/10.1007/s13562-019-00501-7> (2020).
199. Rezaei, E. E. *et al.* Quantifying the response of wheat yields to heat stress: The role of the experimental setup. *Field Crops Research* **217**, 93–103. <https://doi.org/10.1016/j.fcr.2017.12.015> (2018).
200. Mechanisms and modelling. Eyshi Rezaei, E., Webber, H., Gaiser, T., Naab, J. & Ewert, F. Heat stress in cereals. *Eur. J. Agron.* **64**, 98–113. <https://doi.org/10.1016/j.eja.2014.10.003> (2015).
201. Abbas, G. *et al.* Nitrogen rate and hybrid selection matters productivity of maize-maize cropping system under irrigated arid environment of Southern Punjab, Pakistan. *Int. J. Plant Prod.* <https://doi.org/10.1007/s42106-020-00086-5> (2020).
202. Jahan, M. A. H. S. *et al.* Optimizing sowing window for wheat cultivation in Bangladesh using CERES-wheat crop simulation model. *Agr. Ecosyst. Environ.* **258**, 23–29. <https://doi.org/10.1016/j.agee.2018.02.008> (2018).
203. Ahmed, S., Humphreys, E. & Chauhan, B. S. Optimum sowing date and cultivar duration of dry-seeded boro on the High Ganges River Floodplain of Bangladesh. *Field Crops Research* **190**, 91–102. <https://doi.org/10.1016/j.fcr.2015.12.004> (2016).
204. Basso, B., Liu, L. & Ritchie, J. T. in *Advances in Agronomy* Vol. 136 (ed Donald L. Sparks) 27–132 (Academic Press, 2016).
205. Xiong, W. *et al.* A calibration procedure to improve global rice yield simulations with EPIC. *Ecol. Model.* **273**, 128–139. <https://doi.org/10.1016/j.ecolmodel.2013.10.026> (2014).
206. Jalota, S. K., Vashisht, B. B., Kaur, H., Kaur, S. & Kaur, P. Location specific climate change scenario and its impact on rice and wheat in Central Indian Punjab. *Agric. Syst.* **131**, 77–86. <https://doi.org/10.1016/j.agry.2014.07.009> (2014).
207. Peng, S. *et al.* Rice yields decline with higher night temperature from global warming. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 9971–9975. <https://doi.org/10.1073/pnas.0403720101> (2004).
208. Ritchie, J. T., Singh, U., Godwin, D. C. & Bowen, W. T. in *Understanding Options for Agricultural Production* Vol. 7 *Systems Approaches for Sustainable Agricultural Development* (eds Gordon Y. Tsuji, Gerrit Hoogenboom, & Philip K. Thornton) Ch. 5, 79–98 (Springer Netherlands, 1998).
209. Nissanka, S. P. *et al.* Calibration of the phenology sub-model of APSIM-Oryza: Going beyond goodness of fit. *Environmental Modelling & Software* **70**, 128–137. <https://doi.org/10.1016/j.envsoft.2015.04.007> (2015).
210. Weerakoon, W. M. W. *et al.* Direct-seeded rice culture in Sri Lanka: Lessons from farmers. *Field Crops Research* **121**, 53–63. <https://doi.org/10.1016/j.fcr.2010.11.009> (2011).
211. Iftikharuddaula, K. M. *et al.* Development of early maturing submergence-tolerant rice varieties for Bangladesh. *Field Crops Research* **190**, 44–53. <https://doi.org/10.1016/j.fcr.2015.12.001> (2016).
212. Haefele, S. M., Kato, Y. & Singh, S. Climate ready rice: Augmenting drought tolerance with best management practices. *Field Crops Research* **190**, 60–69. <https://doi.org/10.1016/j.fcr.2016.02.001> (2016).
213. Latif, M. A., Islam, M. R., Ali, M. Y. & Saleque, M. A. Validation of the system of rice intensification (SRI) in Bangladesh. *Field Crops Research* **93**, 281–292. <https://doi.org/10.1016/j.fcr.2004.10.005> (2005).
214. Sarangi, S. K. *et al.* Using improved variety and management enhances rice productivity in stagnant flood-affected tropical coastal zones. *Field Crops Research* **190**, 70–81. <https://doi.org/10.1016/j.fcr.2015.10.024> (2016).
215. Raza, M. A. *et al.* Effect of planting patterns on yield, nutrient accumulation and distribution in maize and soybean under relay intercropping systems. *Scientific Reports* **9**, 4947. <https://doi.org/10.1038/s41598-019-41364-1> (2019).
216. Paudel, B., Khanal, R. C., KC, A., Bhatta, K. & Chaudhary, P. Climate-smart agriculture in Nepal. *Research program on Climate Change, Agriculture and food security* (2017).
217. Chauhan, B. S., Mahajan, G., Sardana, V., Timsina, J. & Jat, M. L. in *Advances in Agronomy* Vol. Volume 117 (ed L. Sparks Donald) 315–369 (Academic Press, 2012).
218. Yadvinder, S., Kukal, S. S., Jat, M. L. & Sidhu, H. S. in *Advances in Agronomy* Vol. 127 (ed Donald Sparks) 157–258 (Academic Press, 2014).
219. Jat, M. L. *et al.* in *Advances in Agronomy* Vol. 137 (ed Donald L. Sparks) 127–235 (Academic Press, 2016).
220. Kukal, S. S., Yadvinder, S., Jat, M. L. & Sidhu, H. S. in *Advances in Agronomy* Vol. Volume 127 (ed Sparks Donald) 157–258 (Academic Press, 2014).
221. Witt, C., Pasquin, J. & Dobermann, A. Towards a site-specific nutrient management approach for maize in Asia. *Better Crops* **90**, 28–31 (2006).
222. Ullah, A., Ahmad, A., Khaliq, T. & Akhtar, J. Recognizing production options for pearl millet in Pakistan under changing climate scenarios. *Journal of Integrative Agriculture* **16**, 762–773. [https://doi.org/10.1016/S2095-3119\(16\)61450-8](https://doi.org/10.1016/S2095-3119(16)61450-8) (2017).
223. Ullah, A., Salehnia, N., Kolsoumi, S., Ahmad, A. & Khaliq, T. Prediction of effective climate change indicators using statistical downscaling approach and impact assessment on pearl millet (*Pennisetum glaucum* L.) yield through Genetic Algorithm in Punjab, Pakistan. *Ecological Indicators* **90**, 569–576. <https://doi.org/10.1016/j.ecolind.2018.03.053> (2018).
224. Ausiku, A. P., Annandale, J. G., Steyn, J. M. & Sanewe, A. J. Improving Pearl Millet (*Pennisetum glaucum*) Productivity through Adaptive Management of Water and Nitrogen. *Water* **12**, 422 (2020).
225. Alauddin, M., Rashid Sarker, M. A., Islam, Z. & Tisdell, C. Adoption of alternate wetting and drying (AWD) irrigation as a water-saving technology in Bangladesh: Economic and environmental considerations. *Land Use Policy* **91**, 104430. <https://doi.org/10.1016/j.landusepol.2019.104430> (2020).

226. Rahman, M. H. u. *et al.* in *Cotton Production and Uses: Agronomy, Crop Protection, and Postharvest Technologies* (eds Shakeel Ahmad & Mirza Hasanuzzaman) 447–484 (Springer Singapore, 2020).
227. Ishfaq, M. *et al.* Alternate wetting and drying: A water-saving and ecofriendly rice production system. *Agric. Water Manag.* **241**, 106363. <https://doi.org/10.1016/j.agwat.2020.106363> (2020).
228. Rejesus, R. M., Palis, F. G., Rodriguez, D. G. P., Lampayan, R. M. & Bouman, B. A. M. Impact of the alternate wetting and drying (AWD) water-saving irrigation technique: Evidence from rice producers in the Philippines. *Food Policy* **36**, 280–288. <https://doi.org/10.1016/j.foodpol.2010.11.026> (2011).
229. Carrijo, D. R., Lundy, M. E. & Linquist, B. A. Rice yields and water use under alternate wetting and drying irrigation: A meta-analysis. *Field Crops Research* **203**, 173–180. <https://doi.org/10.1016/j.fcr.2016.12.002> (2017).
230. Stuart, D., Schewe, R. L. & McDermott, M. Reducing nitrogen fertilizer application as a climate change mitigation strategy: Understanding farmer decision-making and potential barriers to change in the US. *Land Use Policy* **36**, 210–218. <https://doi.org/10.1016/j.landusepol.2013.08.011> (2014).
231. Zheng, W., Luo, B. & Hu, X. The determinants of farmers' fertilizers and pesticides use behavior in China: An explanation based on label effect. *Journal of Cleaner Production* **272**, 123054. <https://doi.org/10.1016/j.jclepro.2020.123054> (2020).
232. Jiang, G. *et al.* Soil organic carbon sequestration in upland soils of northern China under variable fertilizer management and climate change scenarios. *Global Biogeochem. Cycles* **28**, 319–333. <https://doi.org/10.1002/2013gb004746> (2014).
233. Omotesho, A., Fakayode, S. & Tariya, Y. Curtailing fertilizer scarcity and climate change; an appraisal of factors affecting organic materials use option in Nigeria's agriculture. *Ethiopian Journal of Environmental Studies and Management* **5**, 281–290 (2012).
234. Raza, A. *et al.* Impact of Climate Change on Crops Adaptation and Strategies to Tackle Its Outcome: A Review. *Plants* **8**, 34 (2019).
235. Sloan, K. *et al.* in *The Climate-Smart Agriculture Papers: Investigating the Business of a Productive, Resilient and Low Emission Future* (eds Todd S. Rosenstock, Andreea Nowak, & Evan Girvetz) 227–233 (Springer International Publishing, 2019).
236. Keith, W. *et al.* Climate change impacts on agriculture in 2050 under a range of plausible socioeconomic and emissions scenarios. *Environ. Res. Lett.* **10**, 085010 (2015).
237. Zhang, Z., Yu, K., Siddique, K. H. M. & Nan, Z. Phenology and sowing time affect water use in four warm-season annual grasses under a semi-arid environment. *Agric. For. Meteorol.* **269–270**, 257–269. <https://doi.org/10.1016/j.agrformet.2019.02.027> (2019).
238. Dreccer, M. F., Fainges, J., Whish, J., Ogbonnaya, F. C. & Sadras, V. O. Comparison of sensitive stages of wheat, barley, canola, chickpea and field pea to temperature and water stress across Australia. *Agric. For. Meteorol.* **248**, 275–294. <https://doi.org/10.1016/j.agrformet.2017.10.006> (2018).
239. Abid, M. *et al.* Physiological and biochemical changes during drought and recovery periods at tillering and jointing stages in wheat (*Triticum aestivum* L.). *Scientific Reports* **8**, 4615. <https://doi.org/10.1038/s41598-018-21441-7> (2018).
240. Macabuhay, A. A. Physiological and biochemical responses of wheat to combined heat stress and elevated CO₂ during grain-filling (2016).
241. Stratonovitch, P. & Semenov, M. A. Heat tolerance around flowering in wheat identified as a key trait for increased yield potential in Europe under climate change. *J. Exp. Bot.* <https://doi.org/10.1093/jxb/erv070> (2015).
242. Hernández, F., Poverene, M., Mercer, K. L. & Presotto, A. Genetic variation for tolerance to extreme temperatures in wild and cultivated sunflower (*Helianthus annuus*) during early vegetative phases. *Crop and Pasture Science* **71**, 578–591. <https://doi.org/10.1071/CP20005> (2020).
243. Acharjee, T. K., van Halsema, G., Ludwig, F., Hellegers, P. & Supit, I. Shifting planting date of Boro rice as a climate change adaptation strategy to reduce water use. *Agric. Syst.* **168**, 131–143. <https://doi.org/10.1016/j.agry.2018.11.006> (2019).
244. Chibarabada, T. P., Modi, A. T. & Mabhaudhi, T. Options for improving water productivity: a case study of bambara groundnut and groundnut. *Phys. Chem. Earth Parts A/B/C* **115**, 102806. <https://doi.org/10.1016/j.pce.2019.10.003> (2020).
245. Islam, A. R. M. T., Shen, S., Yang, S., Hu, Z. & Atiqur Rahman, M. Spatiotemporal rice yield variations and potential agro-adaptation strategies in Bangladesh: A biophysical modeling approach. *Sustain. Prod. Consum.* **24**, 121–138. <https://doi.org/10.1016/j.spc.2020.07.005> (2020).
246. Tsegay, A. *et al.* Sowing and irrigation strategies for improving rainfed tef (*Eragrostis tef* (Zucc.) Trotter) production in the water scarce Tigray region, Ethiopia. *Agric. Water Manag.* **150**, 81–91. <https://doi.org/10.1016/j.agwat.2014.11.014> (2015).
247. van Oort, P. A. J., Timmermans, B. G. H. & van Swaaij, A. C. P. M. Why farmers' sowing dates hardly change when temperature rises. *Eur. J. Agron.* **40**, 102–111. <https://doi.org/10.1016/j.eja.2012.02.005> (2012).
248. Bassu, S., Asseng, S., Motzo, R. & Giunta, F. Optimising sowing date of durum wheat in a variable Mediterranean environment. *Field Crops Res.* **111**, 109–118. <https://doi.org/10.1016/j.fcr.2008.11.002> (2009).
249. Aasen, H., Kirchgessner, N., Walter, A. & Liebisch, F. PhenoCams for field phenotyping: using very high temporal resolution digital repeated photography to investigate interactions of growth, phenology, and harvest traits. *Front. Plant Sci.* <https://doi.org/10.3389/fpls.2020.00593> (2020).
250. Singh, S., Sandhu, S., Dhaliwal, L. & Singh, I. Effect of planting geometry on microclimate, growth and yield of mung-bean (*Vigna radiata* L.). *J. Agric. Phys.* **12**, 70–73 (2012).
251. van Etten, J. *et al.* Crop variety management for climate adaptation supported by citizen science. *Proc. Natl. Acad. Sci.* **116**, 4194–4199. <https://doi.org/10.1073/pnas.1813720116> (2019).
252. Mahato, M. & Adhikari, B. B. Effect of planting geometry on growth of rice varieties. *Int. J. Appl. Sci. Biotechnol.* **5**, 423–429 (2017).
253. Raza, M. A. *et al.* Optimum strip width increases dry matter, nutrient accumulation, and seed yield of intercrops under the relay intercropping system. *Food Energy Secur.* **9**, e199. <https://doi.org/10.1002/fes3.199> (2020).
254. Raza, M. A., van der Werf, W., Ahmed, M. & Yang, W. Removing top leaves increases yield and nutrient uptake in maize plants. *Nutr. Cycl. Agroecosyst.* **118**, 57–73. <https://doi.org/10.1007/s10705-020-10082-w> (2020).
255. Raza, M. A. *et al.* Effects of contrasting shade treatments on the carbon production and antioxidant activities of soybean plants. *Funct. Plant Biol. FPB* **47**, 342–354. <https://doi.org/10.1071/fp19213> (2020).
256. Raza, M. A. *et al.* Optimum leaf defoliation: a new agronomic approach for increasing nutrient uptake and land equivalent ratio of maize soybean relay intercropping system. *Field Crops Res.* **244**, 107647. <https://doi.org/10.1016/j.fcr.2019.107647> (2019).
257. Raza, M. A. *et al.* Growth and development of soybean under changing light environments in relay intercropping system. *PeerJ* **7**, e7262. <https://doi.org/10.7717/peerj.7262> (2019).
258. Raza, M. A. *et al.* Narrow-wide-row planting pattern increases the radiation use efficiency and seed yield of intercrop species in relay-intercropping system. *Food Energy Secur.* **8**, e170. <https://doi.org/10.1002/fes3.170> (2019).
259. Huang, F., Liu, Z., Zhang, P. & Jia, Z. Hydrothermal effects on maize productivity with different planting patterns in a rainfed farmland area. *Soil Tillage Res.* **205**, 104794. <https://doi.org/10.1016/j.still.2020.104794> (2021).
260. Rio, M., Rey, D., Prudhomme, C. & Holman, I. P. Evaluation of changing surface water abstraction reliability for supplemental irrigation under climate change. *Agric. Water Manag.* **206**, 200–208. <https://doi.org/10.1016/j.agwat.2018.05.005> (2018).
261. Muluneh, A., Stroosnijder, L., Keesstra, S. & Biazin, B. Adapting to climate change for food security in the Rift Valley dry lands of Ethiopia: supplemental irrigation, plant density and sowing date. *J. Agric. Sci.* **155**, 703–724. <https://doi.org/10.1017/S0021859616000897> (2016).
262. Ndhleve, S., Nakin, M. & Longo-Mbenza, B. Impacts of supplemental irrigation as a climate change adaptation strategy for maize production: a case of the Eastern Cape Province of South Africa. *Water SA* **43**, 222–228 (2017).

263. Bigelow, D. P. & Zhang, H. Supplemental irrigation water rights and climate change adaptation. *Ecol. Econ.* **154**, 156–167. <https://doi.org/10.1016/j.ecolecon.2018.07.015> (2018).
264. Trevaskis, B. Wheat gene for all seasons. *Proc. Natl. Acad. Sci.* **112**, 11991–11992. <https://doi.org/10.1073/pnas.1516398112> (2015).
265. Matthew, G., Pierre, M. & Ariel, O.-B. Negative impacts of climate change on cereal yields: statistical evidence from France. *Environ. Res. Lett.* **12**, 054007 (2017).
266. Ortiz, R. *et al.* Climate change: can wheat beat the heat?. *Agric. Ecosyst. Environ.* **126**, 46–58. <https://doi.org/10.1016/j.agee.2008.01.019> (2008).
267. Lobell, D. B. *et al.* Analysis of wheat yield and climatic trends in Mexico. *Field Crops Res.* **94**, 250–256. <https://doi.org/10.1016/j.fcr.2005.01.007> (2005).
268. Nazim Ud Dowla, M. A. N., Edwards, I., O'Hara, G., Islam, S. & Ma, W. Developing wheat for improved yield and adaptation under a changing climate: optimization of a few key genes. *Engineering* **4**, 514–522. <https://doi.org/10.1016/j.eng.2018.06.005> (2018).
269. Mohammadi, R. The use of a combination scoring index to improve durum productivity under drought stress. *Exp. Agric.* **56**, 161–170. <https://doi.org/10.1017/S0014479719000231> (2019).
270. Cui, L. *et al.* Development of perennial wheat through hybridization between wheat and wheatgrasses: a review. *Engineering* **4**, 507–513. <https://doi.org/10.1016/j.eng.2018.07.003> (2018).
271. Zachariah, M., Mondal, A., Das, M., AchuthaRao, K. M. & Ghosh, S. On the role of rainfall deficits and cropping choices in loss of agricultural yield in Marathwada, India. *Environ. Res. Lett.* <https://doi.org/10.1088/1748-9326/ab93fc> (2020).
272. Gahlaut, V., Samtani, H. & Khurana, P. Genome-wide identification and expression profiling of cytosine-5 DNA methyltransferases during drought and heat stress in wheat (*Triticum aestivum*). *Genomics* **112**, 4796–4807. <https://doi.org/10.1016/j.ygeno.2020.08.031> (2020).
273. Boote, K. J., Prasad, V., Allen, L. H., Singh, P. & Jones, J. W. Modeling sensitivity of grain yield to elevated temperature in the DSSAT crop models for peanut, soybean, dry bean, chickpea, sorghum, and millet. *Eur. J. Agron.* **100**, 99–109. <https://doi.org/10.1016/j.eja.2017.09.002> (2018).
274. Elbashir, A. A. E. *et al.* Genetic variation in heat tolerance-related traits in a population of wheat multiple synthetic derivatives. *Breed. Sci.* **67**, 483–492. <https://doi.org/10.1270/jsbbs.17048> (2017).
275. Reynolds, M. P. *et al.* An integrated approach to maintaining cereal productivity under climate change. *Global Food Secur.* **8**, 9–18. <https://doi.org/10.1016/j.gfs.2016.02.002> (2016).
276. Asseng, S. *et al.* Model-driven multidisciplinary global research to meet future needs: the case for “improving radiation use efficiency to increase yield”. *Crop Sci.* <https://doi.org/10.2135/cropsci2018.09.0562> (2019).
277. Smit, B. & Skinner, M. W. Adaptation options in agriculture to climate change: a typology. *Mitig. Adapt. Strat. Glob. Change* **7**, 85–114. <https://doi.org/10.1023/A:1015862228270> (2002).
278. Liu, Y., Chen, Q., Ge, Q., Dai, J. & Dou, Y. Effects of climate change and agronomic practice on changes in wheat phenology. *Clim. Change* **150**, 273–287. <https://doi.org/10.1007/s10584-018-2264-5> (2018).
279. Loboguerrero, A. M. *et al.* Food and earth systems: priorities for climate change adaptation and mitigation for agriculture and food systems. *Sustainability* **11**, 1372 (2019).
280. Howden, S. M. *et al.* Adapting agriculture to climate change. *Proc. Natl. Acad. Sci.* **104**, 19691–19696. <https://doi.org/10.1073/pnas.0701890104> (2007).
281. Bryan, E., Deressa, T. T., Gbetibouo, G. A. & Ringler, C. Adaptation to climate change in Ethiopia and South Africa: options and constraints. *Environ. Sci. Policy* **12**, 413–426. <https://doi.org/10.1016/j.envsci.2008.11.002> (2009).
282. Wongnaa, C. A. & Babu, S. Building resilience to shocks of climate change in Ghana's cocoa production and its effect on productivity and incomes. *Technol. Soc.* **62**, 101288. <https://doi.org/10.1016/j.techsoc.2020.101288> (2020).
283. Kumar, S., Mishra, A. K., Pramanik, S., Mamidanna, S. & Whitbread, A. Climate risk, vulnerability and resilience: supporting livelihood of smallholders in semiarid India. *Land Use Policy* **97**, 104729. <https://doi.org/10.1016/j.landusepol.2020.104729> (2020).
284. Morales-Castilla, I. *et al.* Diversity buffers winegrowing regions from climate change losses. *Proc. Natl. Acad. Sci.* **117**, 2864–2869. <https://doi.org/10.1073/pnas.1906731117> (2020).
285. Nelson, G. C. *et al.* *Climate change: Impact on agriculture and costs of adaptation*. Vol. 21 (Intl Food Policy Res Inst, 2009).
286. Olesen, J. E. *et al.* Impacts and adaptation of European crop production systems to climate change. *Eur. J. Agron.* **34**, 96–112. <https://doi.org/10.1016/j.eja.2010.11.003> (2011).
287. Sloat, L. L. *et al.* Climate adaptation by crop migration. *Nat. Commun.* **11**, 200. <https://doi.org/10.1038/s41467-020-15076-4> (2020).
288. Mertz, O., Mbow, C., Reenberg, A. & Diouf, A. Farmers' perceptions of climate change and agricultural adaptation strategies in Rural Sahel. *Environ. Manag.* **43**, 804–816. <https://doi.org/10.1007/s00267-008-9197-0> (2009).
289. Smit, B., Burton, I., Klein, R. J. T. & Wandel, J. An anatomy of adaptation to climate change and variability. *Clim. Change* **45**, 223–251. <https://doi.org/10.1023/A:1005661622966> (2000).
290. Thamo, T. *et al.* Climate change impacts and farm-level adaptation: economic analysis of a mixed cropping–livestock system. *Agric. Syst.* **150**, 99–108. <https://doi.org/10.1016/j.agsy.2016.10.013> (2017).
291. Challinor, A. J. *et al.* A meta-analysis of crop yield under climate change and adaptation. *Nat. Clim. Change* **4**, 287–291. <https://doi.org/10.1038/nclimate2153> (2014).
292. Reidsma, P., Janssen, S., Jansen, J. & van Ittersum, M. K. On the development and use of farm models for policy impact assessment in the European Union—a review. *Agric. Syst.* **159**, 111–125. <https://doi.org/10.1016/j.agsy.2017.10.012> (2018).
293. Reidsma, P. *et al.* Climate change impact and adaptation research requires integrated assessment and farming systems analysis: a case study in the Netherlands. *Environ. Res. Lett.* **10**, 045004 (2015).
294. Reidsma, P., Ewert, F., Lansink, A. O. & Leemans, R. Adaptation to climate change and climate variability in European agriculture: the importance of farm level responses. *Eur. J. Agron.* **32**, 91–102. <https://doi.org/10.1016/j.eja.2009.06.003> (2010).
295. Shahzad, A. N. & Ahmad, S. In *Agronomic Crops: Volume 2: Management Practices* (ed Hasanuzzaman, M.) 111–126 (Springer Singapore, 2019).
296. Ahmed, M. Introduction to Modern Climate Change. Andrew E. Dessler: Cambridge University Press, 2011, 252 pp, ISBN-10: 0521173159. *Science of The Total Environment* **734**, 139397. [10.1016/j.scitotenv.2020.139397](https://doi.org/10.1016/j.scitotenv.2020.139397) (2020).
297. Singh, S. Farmers' perception of climate change and adaptation decisions: a micro-level evidence from Bundelkhand Region, India. *Ecol. Ind.* **116**, 106475. <https://doi.org/10.1016/j.ecolind.2020.106475> (2020).
298. Wallach, D. *et al.* Multimodel ensembles improve predictions of crop–environment–management interactions. *Glob. Change Biol.* **24**, 5072–5083. <https://doi.org/10.1111/gcb.14411> (2018).
299. Aslam, M. U. *et al.* In *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (eds Ahmed, M. & Stockle, C.O.) 113–136 (Springer International Publishing, 2017).
300. Ijaz, W., Ahmed, M., Fayyaz-ul-Hassan, Asim, M. & Aslam, M. In *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (eds Ahmed, M. & Stockle, C.O.) 371–386 (Springer International Publishing, 2017).
301. Jabeen, M., Gabriel, H. F., Ahmed, M., Mahboob, M. A. & Iqbal, J. In *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (eds Ahmed, M. & Stockle, C.O.) 387–411 (Springer International Publishing, 2017).

302. Aslam, M. A., Ahmed, M., Fayyaz-ul-Hassan & Hayat, R. In *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (eds Ahmed, M. & Stockle, C.O.) 71–90 (Springer International Publishing, 2017).
303. Ahmed, M. *et al.* Calibration and validation of APSIM-Wheat and CERES-Wheat for spring wheat under rainfed conditions: models evaluation and application. *Comput. Electron. Agric.* **123**, 384–401. <https://doi.org/10.1016/j.compag.2016.03.015> (2016).
304. Ahmed, M. & Stockle, C. O. *Quantification of Climate Variability, Adaptation and Mitigation for Agricultural Sustainability* (Springer, Berlin, 2016).
305. Ahmed, M., Fayyaz Ul, H. & Van Ogtrop, F. F. Can models help to forecast rainwater dynamics for rainfed ecosystem?. *Weather Clim. Extremes* **5–6**, 48–55. <https://doi.org/10.1016/j.wace.2014.07.001> (2014).
306. Ahmed, M., Hassan, F., Aslam, M. A., Akram, M. N. & Akmal, M. Regression model for the study of sole and cumulative effect of temperature and solar radiation on wheat yield. *Afr. J. Biotech.* **10**, 9114–9121 (2011).
307. Ahmed, M. & Ahmad, S. In *Systems Modeling* (ed Ahmed, M.) 1–44 (Springer Singapore, 2020).
308. Tariq, M., Ahmed, M., Iqbal, P., Fatima, Z. & Ahmad, S. In *Systems Modeling* (ed Ahmed, M.) 45–60 (Springer Singapore, 2020).
309. Ahmed, M., Raza, M. A. & Hussain, T. In *Systems Modeling* (ed Ahmed, M.) 111–150 (Springer Singapore, 2020).
310. Ahmed, M. *et al.* In *Systems Modeling* (ed Ahmed, M.) 151–178 (Springer Singapore, 2020).
311. Kheir, A. M. S. *et al.* In *Systems Modeling* (ed Ahmed, M.) 179–202 (Springer Singapore, 2020).
312. Ahmad, S., & Hasanuzzaman, M. Cotton Production and Uses. Springer Nature Singapore Pte Ltd. (<https://link.springer.com/book/10.1007/978-981-15-1472-2>); doi: 10.1007/978-981-15-1472-2 (2020)
313. Khan, A., Ahmad, M., Shah, M. K. N. & Ahmed, M. Performance of wheat genotypes for Morpho-Physiological traits using multivariate analysis under terminal heat stress. *Pak. J. Bot.* **52**(6), 1981–1988. [https://doi.org/10.30848/PJB2020-6\(30\)\(2020\)](https://doi.org/10.30848/PJB2020-6(30)(2020)).
314. Khan, A., Ahmad, M., Shah, M. K. N., & Ahmed, M. Genetic manifestation of physio-morphic and yield related traits conferring thermotolerance in wheat. *Pak. J. Bot.* **52**(5), 1545–1552. [https://doi.org/10.30848/PJB2020-5\(27\) \(2020\)](https://doi.org/10.30848/PJB2020-5(27) (2020)).

Acknowledgements

Stewart Smock Higgins (SSH) from Washington State University Pullman USA is acknowledged for English editing.

Author contributions

Z.F., M.A., G.A., S.A. reviewed literature and collected data, M.A., M.H., S.U., N.A. designed this work and M.A., S.A. supervised the research work. M.A., M.H., G.A., S.U., S.A., N.A., M.A.A., G.S., E.H., P.I. and S.A. have written this work. S.S.H. served as an English editor. All authors read the manuscript before submission.

Funding

Open Access funding provided by Swedish University of Agricultural Sciences. This study was supported by Higher Education Commission (HEC) Pakistan, National Research Program for Universities (NRPU) (Project# NRPU-6132).

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.A. or S.A.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020