

# The Fitness Effects of Random Mutations in Single-Stranded DNA and RNA Bacteriophages

Pilar Domingo-Calap<sup>1</sup>, José M. Cuevas<sup>1</sup>, Rafael Sanjuán<sup>1,2\*</sup>

<sup>1</sup> Instituto Cavanilles de Biodiversidad y Biología Evolutiva, València, Spain, <sup>2</sup> Departamento de Genética, Universitat de València, València, Spain

## Abstract

Mutational fitness effects can be measured with relatively high accuracy in viruses due to their small genome size, which facilitates full-length sequencing and genetic manipulation. Previous work has shown that animal and plant RNA viruses are very sensitive to mutation. Here, we characterize mutational fitness effects in single-stranded (ss) DNA and ssRNA bacterial viruses. First, we performed a mutation-accumulation experiment in which we subjected three ssDNA ( $\Phi$ X174, G4, F1) and three ssRNA phages (Q $\beta$ , MS2, and SP) to plaque-to-plaque transfers and chemical mutagenesis. Genome sequencing and growth assays indicated that the average fitness effect of the accumulated mutations was similar in the two groups. Second, we used site-directed mutagenesis to obtain 45 clones of  $\Phi$ X174 and 42 clones of Q $\beta$  carrying random single-nucleotide substitutions and assayed them for fitness. In  $\Phi$ X174, 20% of such mutations were lethal, whereas viable ones reduced fitness by 13% on average. In Q $\beta$ , these figures were 29% and 10%, respectively. It seems therefore that high mutational sensitivity is a general property of viruses with small genomes, including those infecting animals, plants, and bacteria. Mutational fitness effects are important for understanding processes of fitness decline, but also of neutral evolution and adaptation. As such, these findings can contribute to explain the evolution of ssDNA and ssRNA viruses.

**Citation:** Domingo-Calap P, Cuevas JM, Sanjuán R (2009) The Fitness Effects of Random Mutations in Single-Stranded DNA and RNA Bacteriophages. *PLoS Genet* 5(11): e1000742. doi:10.1371/journal.pgen.1000742

**Editor:** David J. Begun, University of California Davis, United States of America

**Received:** July 22, 2009; **Accepted:** October 26, 2009; **Published:** November 26, 2009

**Copyright:** © 2009 Domingo-Calap et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was financially supported by grant BFU2008-03978/BMC from the Spanish MICINN ([www.micinn.es](http://www.micinn.es)) to RS, by the Juan de la Cierva and Ramon y Cajal research programs from the Spanish MICINN to JMC and RS, respectively, and by a fellowship from the Generalitat Valenciana ([www.gva.es](http://www.gva.es)) to PDC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: [rafael.sanjuan@uv.es](mailto:rafael.sanjuan@uv.es)

## Introduction

Mutational fitness effects are important for understanding the genetic variability of populations, the relative roles of natural selection and drift, the origin of sex and recombination, or the ability to produce evolutionary innovations, among other processes [1–4]. Further, they are of practical relevance to several fields, including complex human disease [5] and conservation genetics [6]. A simple parameter describing mutational fitness effects is the mean selection strength, defined as the average change in fitness caused by random mutations. However, it is also important to determine their variance and the shape of their statistical distribution. A classic approach to achieve these goals is the mutation-accumulation experiment, in which lines derived from a founder clone are propagated at the smallest possible population size to minimize selection, thereby allowing mutations to accumulate [7,8]. A more direct and powerful approach consists of genetically engineering random mutants, although this has been done far less often due to the greater difficulty of the task [9].

Some progress has been made in characterizing mutational fitness effects. For instance, in *Escherichia coli* more than 90% of gene knock-outs are viable [10] and random insertions caused by transposition reduce fitness by 3% on average [11]. In nematodes, the vast majority of nucleotide substitutions have very small fitness effects [12], whereas in humans few amino acid substitutions have effects greater than 10% and about 30% evolve neutrally [9]. However, the distribution of mutational fitness effects is complex and large differences between species may exist [9]. For example, in some

RNA viruses random nucleotide substitutions reduce fitness by nearly 50% on average and up to 40% are lethal [13,14]. This extreme mutational sensitivity contrasts with the greater robustness of cellular organisms. Mutational robustness is thought to emerge from the presence of alternative metabolic pathways, genetic redundancy, or modularity, but these buffering mechanisms are usually not found in RNA viruses due to their compact genomes [15].

Viruses are a unique experimental system for characterizing mutational fitness effects because it is relatively easy to engineer sets of random single-nucleotide substitutions. This has been done previously for single-stranded (ss) RNA viruses, but not for ssDNA viruses. These two types of viruses have similar genome sizes and therefore might be expected to be equally sensitive to mutation. On the other hand, given the obvious ecological [16], evolutionary [17], and genetic [18] differences between ssDNA and ssRNA viruses, differences in robustness would not be surprising. For instance, ssRNA viruses show higher mutation rates [19–21], possibly selecting for greater robustness [22–24], although this might have also promoted genome compression [25–27]. Second, replicase genes represent a large portion of the genome of RNA viruses but are not encoded by ssDNA viruses, which use host DNA polymerases. Since replicase genes are typically highly constrained, this might also lead to differences in robustness between ssRNA and ssDNA viruses. Finally, some ssDNA viruses encode two scaffolding proteins, apparently introducing some redundancy in the process of capsid assembly [28].

Here, we compare mutational fitness effects in ssDNA and ssRNA viruses using six phages which can infect the same *E. coli*

## Author Summary

The fitness effects of mutations are the raw material for natural selection. It has been shown that point mutations typically have strongly deleterious effects in plant and animal RNA viruses, whereas cellular organisms are comparatively more robust. Here, we characterize the fitness effects of random mutations in DNA viruses and compare them with those found in RNA viruses, using six phage species of similar genome sizes. To achieve this goal, we introduced mutations by chemical and site-directed mutagenesis, identified the genetic changes by sequencing, and quantified their fitness effects using growth-rate assays. In all cases, mutations had a strong average impact on fitness. We conclude that mutational sensitivity is a general property of viruses with small genomes and discuss the evolutionary implications of these findings.


strain under identical environmental conditions [29]: the microviruses  $\Phi$ X174 and G4, the inovirus F1 (ssDNA viruses), the alloviruses Q $\beta$  and SP, and the levivirus MS2 (ssRNA viruses)

(Figure 1). First, we carried out a mutation-accumulation experiment in which we subjected phages to plaque-to-plaque passages and chemical mutagenesis, quantified changes in fitness, and sequenced the ancestral and evolved genomes. Second, we used site-directed mutagenesis to determine the mean strength of selection more accurately and to characterize the distribution of fitness effects in  $\Phi$ X174 and Q $\beta$ . We demonstrate that ssDNA and ssRNA phage exhibit very similar mutational fitness effects.

## Results

### Mutation-accumulation

Preliminarily, we adapted each phage to our laboratory conditions by performing serial passages at high population sizes, until all lineages reached stable fitness values. For each phage, three independent mutation-accumulation lineages were initiated by picking single plaques at random from the adapted populations. Phage supernatants from plaques were subjected to chemical mutagenesis using nitrous acid and plated to isolate new plaques, a protocol that was continued until plaque sizes became drastically reduced. Under these conditions, selection is minimized and therefore, except for highly deleterious or lethal mutations,



**Figure 1. Genome structure of phages  $\Phi$ X174, G4, F1, Q $\beta$ , SP, and MS2, and location of the genetic changes fixed in mutation-accumulation lines.** Genome sizes are 5.4 kb, 5.6 kb, 6.4 kb, 4.2 kb, 4.3 kb, and 3.6 kb, respectively. Protein-coding regions represent between 91% (F1) and 96% ( $\Phi$ X174) of the total genome, respectively. Arrows indicate the location of each cistron and colors represent broad functional categories (blue: replication; green: particle assembly; yellow: coat; red: lysis/extrusion; grey: other/unknown). However, several genes are multifunctional; for instance A2 is involved in phage assembly and lysis. The location of each of the mutations fixed in the mutation-accumulation lines is shown within white bars, each bar corresponding to an independent line (see Table S1 for details). The three DNA phages have circular genomes but are shown in linear form for convenience. The genomes of  $\Phi$ X174 and G4 are shown collinearly to indicate gene homology. However, by convention, the first position of the  $\Phi$ X174 genome is assigned to the unique Pst I cleavage site (vertical dashed line).  
doi:10.1371/journal.pgen.1000742.g001

random genetic drift and mutational pressure are the only factors driving the fixation of mutations [9]. Growth rates were then assayed and the fitness values relative to the non-mutated genotype were obtained as growth rate ratios.

The change in fitness was homogeneous among species (Table 1, nested ANOVA:  $P=0.769$ ) and there were no significant differences between the DNA and RNA groups ( $P=0.605$ ). After sequencing the six ancestral and the 18 evolved genomes (GenBank accession numbers GQ153912-GQ153935) we found that DNA and RNA phages had accumulated  $157\pm 3$  and  $146\pm 1$  nucleotide substitutions in total, respectively (Table 1, Figure 1, nested ANOVA:  $P=0.802$ ). We calculated the per-mutation effect by dividing the fitness loss of each lineage by the number of mutations accumulated and again, this did not reveal significant differences between DNA and RNA phages ( $P=0.870$ ). We therefore conclude that selection strength is similar in the two groups. F1 appeared to show the highest level of robustness, but differences between species were non-significant ( $P=0.088$ ) and a Tukey's post-hoc test indicated that the six phages formed a single coherent group.

However, the number of passages required to reach similar fitness losses or mutation numbers was higher on average for DNA phages (Table 1, nested ANOVA,  $P=0.010$ ). This might be due to their lower spontaneous mutation rates or to lower susceptibility to the mutagen. In both groups, the proportion of transitions was high, and this excess was significantly more marked in DNA phages (94.9% versus 84.9%, Fisher's exact test:  $P=0.004$ ). Since nitrous acid induces mainly transitions, this suggests that, in DNA phages, most substitutions were caused by the mutagen, whereas in RNA phages there was a greater contribution of spontaneous mutation. This could explain why more passages were required in the former group, although differences in susceptibility to the mutagen or in the proportion of transitions among spontaneous mutations cannot be discarded. Importantly, the higher transition/transversion ratio of DNA phages did not result in a significantly higher fraction of synonymous substitutions (nested ANOVA,  $P=0.484$ ) and therefore is unlikely to have biased the above selection strength estimates.

### Site-directed mutagenesis

We constructed 45 clones of  $\Phi$ X174 and 42 clones of Q $\beta$  carrying single-nucleotide substitutions, choosing the target site and the substitution at random. First, we amplified the viral genomes by PCR using mutagenic primers, transfected *E. coli* with the PCR products, picked single plaques, and verified the presence

of the mutation by sequencing. Some transfected cultures failed to form plaques, suggesting that the engineered mutation was lethal for the virus. We first sequenced the region of the mutagenesis PCR product flanking the target site to verify that there were no additional changes. Then, control assays were carried out in which the entire mutagenesis protocol was repeated using PCR primers that did not carry the mutation. By comparing the numbers of plaques formed in mutagenesis and control assays, we showed that 9 and 12 mutations were lethal in  $\Phi$ X174 and Q $\beta$ , respectively (Figure S1), i.e. a lethal fraction of  $p_l=0.20$  and  $p_l=0.29$ , respectively. These two proportions did not differ significantly (Fisher's exact test,  $P=0.454$ ). All lethal mutations found in  $\Phi$ X174 produced amino-acid changes, whereas in the case of Q $\beta$ , one was synonymous (U2379A) and one intergenic (G1329A).

We measured the growth rate of viruses carrying viable substitutions to obtain the distribution of mutational fitness effects. Growth rates were also determined for the above control samples and corrected means and variances were obtained by subtracting the mean and variance of the control group from the grand mean and variance of the mutants. In  $\Phi$ X174, this yielded a mean selection strength of  $\bar{s}=-0.301$  including all (lethal or viable) mutations, with variance  $V(\bar{s})=0.162$ . For viable mutations only,  $\bar{s}_v=-0.126$  and  $V(\bar{s}_v)=0.047$ . In Q $\beta$ ,  $\bar{s}=-0.359$ ,  $V(\bar{s})=0.181$ ,  $\bar{s}_v=-0.103$  and  $V(\bar{s}_v)=0.018$  (Table 2). There were no significant differences in mean selection strength between the two phages (nested ANOVA for viable mutations:  $P=0.633$ ; Mann-Whitney test for all mutations:  $P=0.336$ ). The above correction using controls implies that our inferences of the mean and variance should be free of experimental error or bias resulting from the presence of additional mutations or changes in the assay conditions, although it is still possible that the fitness values of some individual mutants were altered by the presence of additional mutations.

The distribution of fitness effects of viable mutations was highly skewed, with deleterious substitutions of small effect being more abundant than those of large effect (Figure 2), a property shared by most model systems studied [9]. Focusing on viable mutations with negative  $s$ -values, we performed non-linear regression to characterize the shape of the distribution. In  $\Phi$ X174, an exponential model with an expected mean selection strength against deleterious mutations of  $\bar{s}_d=-0.186\pm 0.005$  described the data satisfactorily ( $R^2=0.989$ ) and was as accurate as a Gamma ( $R^2=0.989$ ) or a Beta ( $R^2=0.985$ ) model (partial  $F$ -test,  $P>0.25$  in both cases). Other statistical models were not considered here. In Q $\beta$ , the data were better described by the Gamma distribution ( $R^2=0.975$ ) with an expected  $\bar{s}_d=-0.136\pm 0.008$ . The Beta

**Table 1.** Summary of the results obtained in the mutation-accumulation experiment.

Genetic material	Phage <sup>a</sup>	Passage number	Relative fitness	Mutation number	Fitness effect per mutation <sup>b</sup>
ssDNA	$\Phi$ X174	$60.3\pm 3.0$	$0.235\pm 0.004$	$16.0\pm 3.0$	$-0.051\pm 0.008$
	G4	$34.0\pm 11.7$	$0.313\pm 0.103$	$10.7\pm 2.9$	$-0.072\pm 0.169$
	F1	$45.7\pm 9.9$	$0.342\pm 0.121$	$25.7\pm 2.8$	$-0.027\pm 0.007$
	Average	$46.7\pm 5.9$	$0.297\pm 0.049$	$17.4\pm 2.6$	$-0.050\pm 0.009$
ssRNA	Q $\beta$	$14.3\pm 3.7$	$0.295\pm 0.053$	$18.7\pm 2.4$	$-0.040\pm 0.008$
	SP	$7.0\pm 2.1$	$0.205\pm 0.089$	$15.0\pm 2.3$	$-0.057\pm 0.014$
	MS2	$7.3\pm 0.3$	$0.313\pm 0.081$	$15.0\pm 1.0$	$-0.046\pm 0.003$
	Average	$9.6\pm 1.7$	$0.284\pm 0.031$	$16.2\pm 1.2$	$-0.047\pm 0.053$

<sup>a</sup>For each phage, the average of three independent lines  $\pm$  SEM is shown.

<sup>b</sup>Calculated by subtracting one from the relative fitness and dividing by the number of mutations.

doi:10.1371/journal.pgen.1000742.t001

**Table 2.** Summary of the results obtained in the site-directed mutagenesis experiment<sup>a</sup>.

	$\Phi\text{X174}$	$\text{Q}\beta$
$p_l$	0.20	0.29
$\bar{s}$	-0.301	-0.359
$V(s)$	0.162	0.181
$\bar{s}_v$	-0.126	-0.103
$V(s_v)$	0.047	0.018
$\bar{s}_d$	-0.195/-0.176	-0.152/-0.121
$V(s_d)$	0.031/0.038	0.024/0.033
$p_d$	0.52/0.57	0.45/0.57
$p_n$	0.23/0.28	0.14/0.26


<sup>a</sup> $p_l$ ,  $p_d$ ,  $p_n$ : fraction of lethal, deleterious, and neutral mutations, respectively.  $\bar{s}$ : mean selection strength.  $V(s)$ : variance of mutational fitness effects. Sub-indices  $v$  and  $d$  refer to all viable mutations and deleterious (non-lethal) mutations, respectively.  $p_l$ ,  $\bar{s}$ ,  $V(s)$ ,  $\bar{s}_v$ , and  $V(s_v)$  were obtained directly from the data.  $\bar{s}_d$  and  $V(s_d)$  were estimated by non-linear regression from negative  $s$ -values only and used to infer  $p_d$  and  $p_n$  (95% confidence intervals are shown in these cases).  
doi:10.1371/journal.pgen.1000742.t002

model provided a similarly good fit ( $R^2 = 0.973$ ), whereas the exponential model was slightly worse ( $R^2 = 0.965$ ,  $P = 0.025$ ). Finally, we must emphasize that mutants with positive values were not used for these inferences, which implies that they were assumed to be neutral or beneficial. Neglecting beneficial mutations, the fraction of deleterious to total mutations should thus be  $p_d = (\bar{s} - p_l) / s_d$  and the fraction of neutral to total mutations simply  $p_n = 1 - p_l - p_d$  (Table 2). However, classifying mutations as neutral or deleterious is somewhat unnatural since the difference between a deleterious mutation of infinitesimally small effect and a neutral mutation is merely formal.

## Discussion

Previous site-directed mutagenesis studies have shown that most random nucleotide substitutions are strongly deleterious in animal and plant ssRNA viruses. In Vesicular stomatitis virus (VSV),  $p_l = 0.40$ ,  $\bar{s} = -0.48$  and  $\bar{s}_v = -0.13$  [14], whereas in the unrelated Tobacco etch virus (TEV),  $p_l = 0.41$ ,  $\bar{s} = -0.49$ , and  $\bar{s}_v = -0.13$  [13]. After performing similar experiments with phage  $\text{Q}\beta$ , we confirm that ssRNA viruses are extremely sensitive to mutation in general. Roughly speaking, the probability that a random single nucleotide substitution is lethal for an ssRNA virus is one third or higher, while viable mutations reduce fitness by 10–13% on average. The observed fraction of lethals was lower in  $\text{Q}\beta$  than in VSV or TEV, which might reflect real biological differences or may be a consequence of methodological issues. Concerning the shape of the distribution, viable mutations of small effect are more abundant than those of large effect. The specific statistical model that better describes the data varies, but the Gamma and the Beta distributions are generally accurate. However, larger mutant collections would be needed to increase the statistical power of inferences about the distribution of mutational fitness effects.

We analyzed  $\Phi\text{X174}$  using the same methodology and under the same environment as  $\text{Q}\beta$  to make the comparison between ssDNA and ssRNA viruses. The two phages did not significantly differ in the fraction of lethal mutations or the average strength of selection, suggesting that ssDNA and ssRNA viruses are similarly sensitive to mutation. The strong parallel shown by viruses as different as  $\Phi\text{X174}$  and  $\text{Q}\beta$  probably stems from the fact that both



**Figure 2.** Distribution of fitness effects caused by single-nucleotide substitutions in phages  $\Phi\text{X174}$  and  $\text{Q}\beta$ . Forty-five and 42 mutations, respectively, were obtained by site-directed mutagenesis and assayed for fitness. The effect of each individual mutation is provided in Table S2 and Table S3.  
doi:10.1371/journal.pgen.1000742.g002

have small genomes with few and short non-coding regions, several multifunctional proteins, and little genetic redundancy [15,30]. Differences between  $\Phi\text{X174}$  and  $\text{Q}\beta$  were only minor. For instance, deleterious (non-lethal) effects seemed to follow a simple exponential law in  $\Phi\text{X174}$  whereas  $\text{Q}\beta$  deviated from this model, probably because there were more mildly deleterious mutations and fewer strongly deleterious ones (Figure 2, Table 2), although the fraction of lethals might be higher in the latter.

The similar robustness shown by ssDNA and ssRNA viruses was also supported by the mutation-accumulation experiment, in which we examined three phage species of each group and found no significant differences in mean selection strength. Differences between species within the two groups were weak or absent, although F1 appeared to be the most robust phage. Related to this, it is worth noting that F1 has fewer overlapping genes. This might be related to the fact that inoviruses have filamentous capsids, which are structurally less constrained than icosahedral ones and can hence accommodate larger genomes [31,32]. However, further work is needed to elucidate whether F1 really differs from the other phages in terms of robustness.

It is possible to compare directly the mean selection strengths estimated by mutation-accumulation and site-directed mutagenesis, since the same viruses and the same environment were used. Because lethal mutations cannot be sampled during plaque-to-plaque passages we expected the former to be lower. However, even after excluding lethals, the  $\bar{s}$  values obtained by site-directed mutagenesis experiments were approximately twofold higher. The most likely explanation for this discrepancy is that, since the population size within plaques necessarily exceeds one, some degree of selection is inevitable in mutation-accumulation experiments. Also, given that several mutations accumulated in each lineage, the observed fitness values were dependent on genetic interactions. In RNA viruses, mutations tend to have weaker effects as they accumulate (antagonistic epistasis) and the same might hold for ssDNA viruses and small genomes in general [33], probably leading to an underestimate of the mean selection strength in mutation-accumulation experiments.

The fitness effects of random mutations are relevant to evolution in many ways. For instance, they determine the fraction of nucleotide sites that evolves neutrally and thus the rate at which populations diverge through random genetic drift. Further, neutral and deleterious mutations can also determine the rate of adaptive evolution indirectly. The observation that mutations tend to be highly deleterious indicates that there are no efficient buffering mechanisms, which might result in large phenotypic variation and strong selection for beneficial mutations too [34–36]. Consistent with this view, the fixation of big-benefit mutations has been reported in several phages [37–39] and low mutational robustness has been associated with faster adaptation in VSV [40]. Also, everything else being equal, greater effects of deleterious mutations might result in faster adaptation because strongly deleterious mutations are removed more efficiently from populations, favoring the spread of beneficial ones [41]. The connection between mutational robustness and evolvability is controversial, however. By reducing phenotypic variation, robustness should facilitate the accumulation of genetic variation and may foster evolutionary innovation upon changes in the environment or the genetic background over the long-term [4,36,42–44], a prediction that has also been supported experimentally [42,45].

The fitness effects of random mutations are not only relevant to neutral evolution and adaptation, but obviously, also to processes of fitness decline such as Muller's ratchet [46] or lethal mutagenesis [47]. Viruses experience frequent transmission bottlenecks during which random mutations may fix in the population. The consequences of this process for viral fitness are highly dependent on how deleterious are these mutations on average [46]. A substantial proportion of the variation observed in natural populations of RNA viruses comprises transient deleterious mutations, but most fail to reach fixation [48], possibly due to their strongly negative impact on fitness. Mutational fitness effects are also important for understanding viral extinction through mutagenesis [47] and the clinical use of mutagens as ribavirin to treat viral infections makes this subject of practical importance.

A conceptual dichotomy between DNA and RNA viruses has been established based on the observation that the latter evolve faster [49]. However, it has been shown more recently that some ssDNA viruses, including parvoviruses [50], anelloviruses [51] and geminiviruses [52] can match the evolutionary rates of RNA viruses and rapidly adapt to new hosts (although this has not been tested experimentally yet). Despite mutating faster than double-stranded DNA viruses or cellular organisms, ssDNA viruses are less error-prone than ssRNA viruses [19–21]. Therefore, the rate at which spontaneous mutations occur may not fully explain why ssRNA and ssDNA viruses evolve at similar rates. Additional

factors as, for instance, the fitness effects of mutations, should be considered.

## Materials and Methods

### Bacteriophages and cells

Bacteriophages  $\Phi$ X174, G4, F1, Q $\beta$ , SP, and MS2 and the *E. coli* C strain IJ1862 [29] were obtained from Prof. James J. Bull (University of Texas). The Q $\beta$  infectious clone pBRT7Q $\beta$  was provided by Dr. René C. Olsthoorn (Leiden University). General biology of the six phages can be found elsewhere [31].

### Preliminary adaptation

Each phage was serially passaged at high population sizes in IJ1862 cells. For each transfer,  $\sim 10^5$  particle forming units (pfu) were inoculated into 0.5 mL LB medium containing IJ1862 cells at their exponential growth phase. The appropriate cell density varied depending on the virus lytic activity and growth rate ( $OD_{600} = 0.7$  for  $\Phi$ X174 and G4, 0.15 for F1, Q $\beta$ , and MS2, and 0.05 for SP). Infected cultures were incubated in agitation (650 rpm) at 37°C in a Thermomixer 24-tube shaker (Eppendorf) and harvested during the late exponential growth phase of the virus ( $\sim 10^9$  pfu/mL). Cells were removed by centrifugation and supernatants were aliquoted, stored at  $-70^\circ\text{C}$ , and titrated using LB medium solidified with soft agar (0.7%). Initial and final titers were used to calculate growth rates and to adjust sampling times for the next passage accordingly. We performed 60–80 passages for each phage. In all cases, no significant changes in growth rate were observed during the last 30 passages, indicating that phages had reached a quasi-maximal fitness under these experimental conditions.

### Mutation-accumulation experiment

Three independent mutation-accumulation lines were seeded for each phage by picking random plaques from the high-fitness populations. Each plaque was resuspended in 50  $\mu\text{L}$  LB and stored at  $-70^\circ\text{C}$ . Lines were propagated plaque-to-plaque as previously described [53] and after each passage, phages were mutagenized with nitrous acid. To do so, four volumes of 0.3 M acetate buffer pH 4.3 were mixed with one volume of 5 M sodium nitrite and 50  $\mu\text{L}$  of this solution were immediately added to 4  $\mu\text{L}$  of phage-containing supernatant. The exposure time was adjusted to obtain a maximal titer loss. The mutagenesis reactions were quenched by adding 200  $\mu\text{L}$  of 1 M acetate buffer pH 4.3 containing 0.1 mg/mL BSA and 100  $\mu\text{L}$  of this final mix were plated without dilution. After incubation at 37°C, a single lysis plaque was picked, resuspended in 50  $\mu\text{L}$ , aliquoted, stored at  $-70^\circ\text{C}$ , titrated, and used for the next round of chemical mutagenesis.

### Site-directed mutagenesis

The High Pure Plasmid Isolation kit (Roche) was used to purify  $\Phi$ X174 DNA and the pBRT7 $\beta$  plasmid from partially lysed *E. coli* IJ1862 cells and from an overnight culture of *E. coli* K12 previously transformed with this cDNA, respectively. For PCR-based mutagenesis, full-length PCR amplicons were obtained from 500 pg of template using Phusion high-fidelity DNA polymerase (New England Biolabs, error rate provided by the manufacturer:  $4.4 \times 10^{-7}$  per base per replication round) and a pair of adjacent, divergent, 5' phosphorylated primers, one of which carried the desired nucleotide substitution. PCR products were circularized using the Quick T4 ligase (New England Biolabs) and *E. coli* IJ1862 competent cells were transfected by the heat-shock method (42 C, 30 s) in the presence of  $\text{CaCl}_2$  100 mM. The transfected cells were immediately plated onto LB plates using soft agar and



individual plaques were picked after 5–9 hours of incubation at 37°C, resuspended in LB, and stored at –70°C. To verify that the desired mutation had been introduced and that no additional changes were present in the region flanking the target site, (RT)-PCR was performed directly from the resuspended plaque. Moloney murine leukemia virus reverse transcriptase (Fermentas) was used for cDNA synthesis and Phusion DNA polymerase for PCR. The products were column-purified and sequenced using sequence-specific primers.

In cases where transfection yielded no plaques or phages recovered from plaques had not incorporated the mutation, the entire protocol was repeated and, after three consecutive failures, the mutation was classified as a candidate lethal. In all of these cases, the number of plaques obtained from transfection assays was abnormally low, suggesting that the mutation was lethal for the virus and that the few observed plaques came from the template wild-type DNA. To confirm lethality, we first sequenced the region of the PCR-based mutagenesis product flanking the target site to check that the mutation was present and that no additional changes had appeared in this region. Then, we designed new primers that did not carry the nucleotide substitution but were otherwise identical to those used for the mutagenesis reaction, and performed the PCR, circularization and transfection steps exactly as above. Lethality was assessed based on the comparison between the numbers of plaques recovered from mutagenesis versus control reactions (Figure S1).

Similar control assays were carried out to estimate the contribution of non-desired mutations and other sources of error to the inferred distribution of mutational fitness effects. For each phage, we transfected three of the above control PCR-based mutagenesis products and picked 11–12 random plaques from each, yielding 36 and 33 plaques in total for  $\Phi$ X174 and Q $\beta$ , respectively. The relative fitness of each clone was determined as above and we obtained the mean and variance for each phage. These values were subtracted from the means and variances obtained for the real mutants. This allowed us to account for experimental error and, in particular, to control for any potential mutations present in the DNA templates, arising during PCR amplification, or during plaque growth.

### Fitness assays

To measure growth rates,  $\sim 10^4$  pfu were inoculated into 0.5 mL of LB medium containing J11862 cells at their exponential growth phase. Infected cultures were incubated in agitation (650 rpm) at 37°C in a Thermomixer 24-tube shaker and harvested when the wild-type reached an estimated titer of  $10^8$  pfu/mL. Cells were removed by centrifugation and the supernatants were aliquoted, stored at –70°C and titrated. The growth rate ( $r$ ) was calculated as the increase in log-titer per hour. Relative fitness was obtained as  $W = r_i/r_0$ , where  $i$  and  $0$  refer to the mutant and wild-type, respectively (notice that this fitness measure is in log-scale), and the fitness effect (selection coefficient) was calculated as  $s = W - 1$ . In all cases, mutant and wild-type samples were assayed in the same experimental block, and experimental blocks were done in triplicate.

### Sequencing

Sequences were obtained using the Sanger method (Applied Biosystems). In general, we sequenced column-purified plaque-(RT)-PCRs directly. In cases where this yielded low-quality readings, we cloned the PCR fragments using the Zero Blunt TOPO PCR Cloning kit (Invitrogen) and sequenced the inserts with vector-based and internal primers, discarding mutations that were not present in at least 3/5 clones.

### Statistical analyses

For the mutation-accumulation experiment, we used a univariate linear model with two factors: Genetic material (G, main fixed factor) and Phage species (S, random factor nested within G), whereas the line was the experimental replicate. The variables analyzed were the number of passages, the number of mutations fixed, and the expected fitness effect per mutation. For each variable  $v$ , the model can be written as  $v_{ijk} = \mu + G_i + S_j + l_{ijk}$ , where  $\mu$  is the grand mean and  $l$  is the line (error term).

To test for differences in average fitness between the  $\Phi$ X174 and Q $\beta$  mutant collections, we used a univariate linear model with three factors: Phage species (S, main fixed factor), Assay type (A, site-directed mutagenesis or control, fixed factor nested within S), and Plaque (P, each of the mutant or control plaques, random factor nested within A). Hence, each experimental determination can be expressed as  $s_{ijkl} = \mu + S_i + A_j + P_{ijk} + \varepsilon_{ijkl}$ , where  $\mu$  is the grand mean,  $A_{ij}$  is positive for mutagenesis assays and negative for control assays, and  $\varepsilon$  is the error term.

Non-linear regressions were performed to estimate  $\bar{s}_d$  and  $V(s_d)$  using the Levenberg-Marquardt algorithm implemented in SPSS v12.

### Supporting Information

**Figure S1** Confirmation of lethal mutations. Site-directed mutagenesis reactions and control reactions (using non-mutagenic but otherwise identical primers) were loaded in 0.8% agarose gels and stained with ethidium bromide. For each candidate mutation, the left and right lanes contain the mutagenesis and control reactions, respectively. Bands of the expected size (5.4 kb for  $\Phi$ X174 and ca. 7.0 kb for Q $\beta$  insert plus the vector) were obtained in all cases (plus additional, non-specific, lower molecular weight bands which did not interfere with transfections). The number of plaques obtained after transfection of *E. coli* cells with these PCR products are shown below the brackets. (A) For  $\Phi$ X174, all mutations except A402T were confirmed as lethal. (B) For Q $\beta$ , all mutations except G780U, G1110A, and G2587U were confirmed. Found at: doi:10.1371/journal.pgen.1000742.s001 (0.43 MB PDF)

**Table S1** Genetic changes found in  $\Phi$ X174, G4, F1, Q $\beta$ , SP and MS2 mutation accumulation lines (three lines per phage). The final passage number, the relative fitness and its standard error (SEM), the nucleotide substitutions found (genomic position and substitution), the genes where they map, and the associated amino acid change (amino acid number within gene) are shown. Notice that some mutations fall at overlapping genes (the respective amino acid substitutions are indicated in these cases). Genomic positions in  $\Phi$ X174 are numbered following the convention that nucleotide 1 corresponds to the Pst I cleavage site. Sequences covered nearly the whole genome (>98%) in all cases except for line SP.1 in which we failed to sequence genomic regions 1–167, 653–823, 2277–2633, and 2985–3186, and line MS2.3 in which we failed to sequence region 3048–3569. Found at: doi:10.1371/journal.pgen.1000742.s002 (0.03 MB PDF)

**Table S2**  $\Phi$ X174 point mutants obtained by site-directed mutagenesis. For each mutant, the genomic position of the nucleotide substitution (relative to the Pst I cleavage site), the mutated gene, the amino acid change, and the relative fitness effect  $\pm$  SEM are shown. Assays were done in triplicate and fitness values were corrected to account for the presence of additional mutations and other sources of experimental error (see Materials and Methods). Notice that some mutations fall within regions at overlapping genes (the respective amino acid substitutions are indicated in these cases). Found at: doi:10.1371/journal.pgen.1000742.s003 (0.02 MB PDF)

**Table S3** Q $\beta$  point mutants obtained by site-directed mutagenesis. For each mutant, the genomic position of the nucleotide substitution, the mutated gene, the amino acid change, and the relative fitness effect  $\pm$  SEM are shown. Assays were done in triplicate and fitness values were corrected to account for the presence of additional mutations and other sources of experimental error (see Materials and Methods). Notice that some mutations fall within regions with overlapping genes (the respective amino acid substitutions are indicated in these cases).

Found at: doi:10.1371/journal.pgen.1000742.s004 (0.02 MB PDF)

## References

- de Visser JA, Hermisson J, Wagner GP, Ancel ML, Bagheri-Chaichian H, et al. (2003) Perspective: Evolution and detection of genetic robustness. *Evolution* 57: 1959–1972.
- Ohta T (1992) The nearly neutral theory of molecular evolution. *Annu Rev Ecol Syst* 23: 263–286.
- Peck JR, Barreau G, Heath SC (1997) Imperfect genes, Fisherian mutation and the evolution of sex. *Genetics* 145: 1171–1199.
- Wagner A (2005) Robustness and evolvability in living systems. New Jersey: Princeton University Press.
- Eyre-Walker A, Woolfit M, Phelps T (2006) The distribution of fitness effects of new deleterious amino acid mutations in humans. *Genetics* 173: 891–900.
- Schultz ST, Lynch M (1997) Mutation and extinction: the role of variable mutational effects, synergistic epistasis, beneficial mutations, and degree of outcrossing. *Evolution* 51: 1363–1371.
- Keightley PD (2004) Comparing analysis methods for mutation-accumulation data. *Genetics* 167: 551–553.
- Mukai T (1964) The genetic structure of natural populations of *Drosophila melanogaster*. I. spontaneous mutation rate of polygenes controlling viability. *Genetics* 50: 1–19.
- Eyre-Walker A, Keightley PD (2007) The distribution of fitness effects of new mutations. *Nat Rev Genet* 8: 610–618.
- Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, et al. (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* 2: 2006.
- Elena SF, Ekunwe L, Hajela N, Oden SA, Lenski RE (1998) Distribution of fitness effects caused by random insertion mutations in *Escherichia coli*. *Genetica* 102/103: 349–358.
- Davies EK, Peters AD, Keightley PD (1999) High frequency of cryptic deleterious mutations in *Caenorhabditis elegans*. *Science* 285: 1748–1751.
- Carrasco P, de la Iglesia F, Elena SF (2007) Distribution of fitness and virulence effects caused by single-nucleotide substitutions in *Tobacco etch virus*. *J Virol* 81: 12979–12984.
- Sanjuán R, Moya A, Elena SF (2004) The distribution of fitness effects caused by single-nucleotide substitutions in an RNA virus. *Proc Natl Acad Sci USA* 101: 8396–8401.
- Elena SF, Carrasco P, Darós JA, Sanjuán R (2006) Mechanisms of genetic robustness in RNA viruses. *EMBO Rep* 7: 168–173.
- Holmes EC (2008) Evolutionary history and phylogeography of human viruses. *Annu Rev Microbiol* 62: 307–328.
- Duffy S, Shackleton LA, Holmes EC (2008) Rates of evolutionary change in viruses: patterns and determinants. *Nat Rev Genet* 9: 267–276.
- Ahluquist P (2006) Parallels among positive-strand RNA viruses, reverse-transcribing viruses and double-stranded RNA viruses. *Nat Rev Microbiol* 4: 371–382.
- Cuevas JM, Duffy S, Sanjuán R (2009) Point mutation rate of bacteriophage  $\Phi$ X174. *Genetics*, in press.
- Drake JW, Charlesworth B, Charlesworth D, Crow JF (1998) Rates of spontaneous mutation. *Genetics* 148: 1667–1686.
- Sniegowski PD, Gerrish PJ, Johnson T, Shaver A (2000) The evolution of mutation rates: separating causes from consequences. *BioEssays* 22: 1057–1066.
- Sanjuán R, Cuevas JM, Furió V, Holmes EC, Moya A (2007) Selection for robustness in mutagenized RNA viruses. *PLoS Genet* 3: e93. doi:10.1371/journal.pgen.0030093.
- Schuster P, Swetina J (1988) Stationary mutant distributions and evolutionary optimization. *Bull Math Biol* 50: 635–660.
- Wilke CO, Wang JL, Ofria C, Lenski RE, Adami C (2001) Evolution of digital organisms at high mutation rates leads to survival of the flattest. *Nature* 412: 331–333.
- Belshaw R, Pybus OG, Rambaut A (2007) The evolution of genome compression and genomic novelty in RNA viruses. *Genome Res* 17: 1496–1504.
- Eigen M, McCaskill J, Schuster P (1988) Molecular quasi-species. *J Phys Chem* 92: 6881–6891.
- Holmes EC (2003) Error thresholds and the constraints to RNA virus evolution. *Trends Microbiol* 11: 543–546.
- Chen M, Uchiyama A, Fane BA (2007) Eliminating the requirement of an essential gene product in an already very small virus: scaffolding protein B-free  $\Phi$ X174, B-free. *J Mol Biol* 373: 308–314.
- Bull JJ, Badgett MR, Springman R, Molineux JJ (2004) Genome properties and the limits of adaptation in bacteriophages. *Evolution* 58: 692–701.
- Belshaw R, Gardner A, Rambaut A, Pybus OG (2008) Pacing a small cage: mutation and RNA viruses. *Trends Ecol Evol* 23: 188–193.
- Calendar R (2006) The bacteriophages. Oxford University Press.
- van der Ende A, Teertstra R, Weisbeek PJ (1982) Initiation and termination of the bacteriophage  $\Phi$ X174 rolling circle DNA replication *in vivo*: packaging of plasmid single-stranded DNA into bacteriophage  $\Phi$ X174 coats. *Nucl Acids Res* 10: 6849–6863.
- Sanjuán R, Elena SF (2006) Epistasis correlates to genomic complexity. *Proc Natl Acad Sci USA* 103: 14402–14405.
- Frank SA (2007) Maladaptation and the paradox of robustness in evolution. *PLoS ONE* 2: e1021. doi:10.1371/journal.pone.0001021.
- Lenski RE, Barrick JE, Ofria C (2006) Balancing robustness and evolvability. *PLoS Biol* 4: e428. doi:10.1371/journal.pbio.0040428.
- Wagner A (2008) Robustness and evolvability: a paradox resolved. *Proc Biol Sci* 275: 91–100.
- Bull JJ, Badgett MR, Wichman HA (2000) Big-benefit mutations in a bacteriophage inhibited with heat. *Mol Biol Evol* 17: 942–950.
- Rokyta DR, Joyce P, Caudle SB, Wichman HA (2005) An empirical test of the mutational landscape model of adaptation using a single-stranded DNA virus. *Nat Genet* 37: 441–444.
- Rokyta DR, Beisel CJ, Joyce P, Ferris MT, Burch CL, Wichman HA (2008) Beneficial fitness effects are not exponential for two viruses. *J Mol Evol* 67: 368–376.
- Cuevas JM, Moya A, Sanjuán R (2009) A genetic background with low mutational robustness is associated with increased adaptability to a novel host in an RNA virus. *J Evol Biol* 22: 2041–2048.
- Orr HA (2000) The rate of adaptation in asexuals. *Genetics* 155: 961–968.
- Bloom JD, Labthavikul ST, Otey CR, Arnold FH (2006) Protein stability promotes evolvability. *Proc Natl Acad Sci USA* 103: 5869–5874.
- Ciliberti S, Martin OC, Wagner A (2007) Innovation and robustness in complex regulatory gene networks. *Proc Natl Acad Sci USA* 104: 13591–13596.
- Kim Y (2007) Rate of adaptive peak shifts with partial genetic robustness. *Evolution*. 1847–1856.
- McBride RC, Ogbunugafor CB, Turner PE (2008) Robustness promotes evolvability of thermotolerance in an RNA virus. *BMC Evol Biol* 8: 231.
- Butcher D (1995) Muller's ratchet, epistasis and mutation effects. *Genetics* 141: 431–437.
- Bull JJ, Sanjuán R, Wilke CO (2007) Theory of lethal mutagenesis for viruses. *J Virol* 81: 2930–2939.
- Pybus OG, Rambaut A, Belshaw R, Freckleton RP, Drummond AJ, et al. (2007) Phylogenetic evidence for deleterious mutation load in RNA viruses and its contribution to viral evolution. *Mol Biol Evol* 24: 845–852.
- Holland JJ, Spindler K, Horodyski F, Grabau E, Nichol S, et al. (1982) Rapid evolution of RNA genomes. *Science* 215: 1577–1585.
- Shackleton LA, Parrish CR, Truyen U, Holmes EC (2005) High rate of viral evolution associated with the emergence of carnivore parvovirus. *Proc Natl Acad Sci USA* 102: 379–384.
- Umehura T, Tanaka Y, Kiyosawa K, Alter HJ, Shih JW (2002) Observation of positive selection within hypervariable regions of a newly identified DNA virus (SEN virus - 1). *FEBS Lett* 510: 171–174.
- Duffy S, Holmes EC (2008) Phylogenetic evidence for rapid rates of molecular evolution in the single-stranded DNA begomovirus Tomato yellow leaf curl virus. *J Virol* 82: 957–965.
- Chao L (1990) Fitness of RNA virus decreased by Muller's ratchet. *Nature* 348: 454–455.

## Acknowledgments

We thank Jim Bull and René C. Olsthoorn for supplying the viruses, cells, and molecular clones; Jeff Clune, Siobain Duffy, Holly Wichman, and Jim Bull for helpful comments; and Paqui de la Iglesia, Concha Hueso, and Loles Catalán for technical assistance.

## Author Contributions

Conceived and designed the experiments: RS. Performed the experiments: PDC JMC RS. Analyzed the data: RS. Wrote the paper: RS.