

# The Genetics of Autism Spectrum Disorders

John J.M. Connolly<sup>1</sup> and Hakon Hakonarson<sup>1,2</sup>

<sup>1</sup>*Center for Applied Genomics*

*The Children's Hospital of Philadelphia, Philadelphia*

<sup>2</sup>*Department of Pediatrics, University of Pennsylvania School of Medicine, Philadelphia  
USA*

## 1. Introduction

Autism is a neurodevelopmental disorder of complex etiology and is amongst the most heritable of neuropsychiatric disorders while sharing genetic liability with other neurodevelopmental disorders such as intellectual disability (ID). Autism spectrum disorders (ASDs) are defined more broadly and include autism, Asperger syndrome, childhood disintegrative disorder and pervasive developmental disorder not otherwise specified. Under the Diagnostic and Statistical Manual of Mental Disorders, 4<sup>th</sup> Edition Revised (DSM-IVTR), these disorders are grouped together with Rett syndrome ("Rett's disorder") as pervasive developmental disorders. However, Rett syndrome has a reportedly distinct pathophysiology, clinical course, and diagnostic strategy (Levy & Schultz, 2009) and will likely be removed in the impending publication of DSM-V (APA, 2010). The new diagnostic manual will formally adopt the single diagnostic category "ASDs", which is used here. Reported prevalence rates for ASDs range from 20 (Newschaffer *et al.* 2007) to 116 (Baird *et al.*, 2006) per 10,000 children, and vary in accordance with diagnostic, sampling, and screening criteria. The Centers for Disease Control and Prevention (CDC) suggest that in the United States, the prevalence of ASDs is 1 in 110 (1/70 in boys and 1/315 in girls) (ADDM, 2009). The three primary characteristics of ASDs are communication impairments, social impairments, and repetitive/stereotyped behaviors. The DSM-IVTR, ICD-10, and many other diagnostic instruments require impairment in each of these domains for a diagnosis of autistic disorder.

Within the last decade, a number of major technological developments have transformed our understanding of the genetic causes of autism, and the field continues to evolve rapidly. In this chapter, we will review three approaches to identifying genetic factors that contribute to the pathogenesis of ASDs: 1) common variants and genome-wide association studies (GWAS); 2) rare variants and copy number variation (CNV) studies, and 3) familial forms of autism and the role of next-generation sequencing (NGS) methods. Data from all three approaches underscores the conclusion that autism is a highly complex and heterogeneous disorder, involving a multifactorial etiology. Moreover, it is becoming increasingly apparent that autism is not a unitary disorder, and that the spectrum may consist of any number of different autisms that share similar symptoms or phenotypes. This conclusion has important implications for evaluation and treatment, which are discussed in the conclusion.

## 2. Heritability of ASDs

Although Skuse (2007) cautions that heritability estimates of ASDs may have been skewed by the co-inheritance of (low) intelligence or other variables, there is little doubt that genetic factors play a key role in autism. In the most widely-cited twin study, Bailey *et al.* (1995) report that monozygotic twins are 92% concordant on a broad spectrum of cognitive or social abnormalities, compared with only 10% for dizygotic twins. Parents and siblings of individuals with ASDs often exhibit subsyndromal levels of impairment (Piven *et al.*, 1997), and having an affected sibling is the single biggest risk factor for developing an ASD. In an analysis of 943,664 Danish children (Lauritsen *et al.*, 2005), the strongest predictors of autism were siblings with ASDs, who conferred a 22-fold increased risk, while Fombonne (2005) suggested that this risk may be even greater.

## 3. Early insight from Rett syndrome and fragile X

Early efforts to identify the genetic causes of ASDs utilized linkage and association approaches. Linkage studies, more prominent in the 1980s and 1990s, typically focus on families or larger pedigrees and is well powered to identify rare genetic variants. The most common linkage approach is the affected sib-pair design (see O'Roak & State, 2008), which examines the transmission of genomic segments through generations. Linkage studies helped define the locus containing *FMR1*, which is mutated in fragile X syndrome (e.g. Richards *et al.*, 1991), and is a common cause of autism in fragile X syndrome, affecting ~30% of children who are diagnosed with fragile X (Rogers *et al.*, 2001; Harris *et al.*, 2008). Similarly, this approach has been important to identifying *MECP2* as the major cause of Rett syndrome (e.g. Curtis *et al.*, 1993).

Association studies take the opposite approach, scanning the genome from the top down, with the goal of determining *post-hoc* whether identified variants are more or less common in affected individuals. Early association studies (i.e. pre HapMap) were complementary with the linkage approach, and in many designs, linkage primed target loci for this more fine-grained analysis. These early insights have played a significant role in shaping our current understanding of ASDs, and functional studies of *FMR1* and *MECP2* have highlighted the importance of synaptic dysfunction (Ramocki & Zoghbi, 2008) as a unifying factor that could extend into the more common forms of autism. This is significant because it provides a means of linking neural correlates with genomic data, as well as related clinical phenotypes such as seizures and cognitive deficits (Hagerman *et al.*, 2009). The Alzheimer's paradigm, which includes functional models of how molecular, biochemical, and neural systems interact is instructive in this regard (e.g. Cissé *et al.*, 2011).

## 4. Genome wide association and common variants

Aside from notable successes with fragile X and Rett syndrome, early linkage and association studies have been inconsistent in resolving more complex genetic correlates of ASDs, with candidate genes often not being replicated between studies. These challenges may in part accounted-for by their relatively low resolution, which makes it difficult to detect candidate loci other than those of major effect. In the past decade however, association studies have become increasingly more sophisticated, with the whole-genome approach, allowing us to examine thousands of individuals on a mass scale, using hundreds of thousands of markers.

Genome-wide association studies (GWAS) examine the frequency of single nucleotide polymorphisms (SNPs) in cases *versus* control populations and can adopt either a case-control or family-based approach. The former allows researchers to avoid the often complex process of acquiring diagnostic/phenotype data from a patient's family, and can incorporate very large numbers of control datasets that may be more readily available. The latter controls for the often confounding phenomenon of population stratification, where variants more common to specific racial groups may either be erroneously identified as causal, or obscure actual causal variants. A major caveat with family-based designs is the often unfounded assumption that unaffected family members do not share causal variants.

GWAS test for common variants (>1% population frequency), with the assumption that ASDs are at least in part caused by the coinheritance of multiple risk variants, each of small individual effect (odds ratios between 1:1 and 1:5). This assumption is known as the common disease-common variant (CDCV) model (Risch and Merikangas, 1996).

A 2009 paper by Wang *et al.* (2009) from our laboratory was the first to identify common variants for ASDs on a genome-wide scale. We examined 780 families (3,101 individuals) with affected children, a second group of 1,204 affected individuals, and 6,491 controls, all of whom were of European ancestry. We identified six genetic markers on chromosome 5 in the 5p14.1 region that confirmed susceptibility to ASDs. The region straddles two genes, *CDH9* and *CDH10*. Both genes encode type II classical cadherins, transmembrane proteins that promote cell adhesion. The association of cadherins is consistent with the cortical-disconnectivity model of autism (e.g. Gepner & Féron, 2009), which postulates that ASDs may result from an increase or decrease in functional connectivity and neuronal synchronization in relevant neural pathways. Functional studies suggest that under-activity between and within networks are correlated with social, communication, cognitive, and sensorimotor impairments (Müller *et al.*, 2011).

The study design utilized two independent replication cohorts and the key SNP at this locus has also been replicated in two additional independent cohort studies (Ma *et al.*, 2009; St Pourcain *et al.*, 2010), lending further support that genetic factors at the 5p14 locus, which is flanked by two relevant cadherin genes, represent strong candidates for aligning molecular function with known neural deficits in ASDs. The original report by Wang and colleagues, also demonstrated that there was an enrichment in Catherin associated genes in ASDs in general, based on gene-pathway analysis (Wang *et al.*, 2009). Cadherins represent a large family of transmembrane proteins that mediates calcium-dependent cell-cell adhesion and, *via* cell adhesion, has been shown to generate synaptic complexity in the developing brain (Redies, 2000). Other common GWAS variants reported have not been replicated in independent studies and will not be covered here.

#### 4.1 Replicated common variants from candidate gene studies

Other common variants from candidate gene studies include *CNTNAP2*, *EN2*, and *MET*, which are reviewed briefly below. A more in depth review of these genes can be derived from catalogs at <http://www.genome.gov/26525384> and <http://www.ncbi.nlm.nih.gov/omim/209850>.

Located on chromosome 7q35, Contactin Associated Protein 2 (*CNTNAP2*) was identified by Alarcón *et al.* (2002) as a candidate for the age at first word endophenotype. A subsequent follow-up by the same group (Alarcón *et al.* 2008) using linkage, association, and gene-expression analyses, found *CNTNAP2* to be the only autism-susceptibility gene to reach significance across all approaches. An independent linkage analysis by Arking *et al.* (2008) also highlighted *CNTNAP2* as a significant ASD candidate gene. *CNTNAP2* is part of the

neurexin family, which have repeatedly been associated with autism (see below). Interestingly, Vernes *et al.* (2008) showed that *CNTNAP2* binds to *FOXP2*, which is a well-established correlate of language and speech disorders (Lai *et al.*, 2001) – a common phenotype in ASDs.

Engrailed 2 (*EN2*) is a homeobox gene that is critical to the development of the midbrain and cerebellum. Like other homeobox genes, it regulates morphogenesis. *EN2* is a human homolog of the engrailed gene, which is found in *Drosophila*. *En2* mouse mutants have anatomic phenotypes in the cerebellum that resemble cerebellar abnormalities reported in autistic individuals (Cheng *et al.*, 2010). Benayed *et al.* (2005, 2009) have reported and replicated in three separate datasets a significant association with broad and narrow ASD phenotypes. Wang *et al.* (2008) also found an association between *EN2* and ASDs in a Chinese Han sample, although Zhong *et al.* (2003) failed to find evidence of an underlying association.

The oncogene *MET* is also strongly linked to ASD etiology, having been supported by a number of studies in the past decade (e.g. IMGSA, 2001; Campbell *et al.*, 2006, 2008; Sousa *et al.*, 2009). Recently, Eagleson *et al.* (2011) reported a role for *Met* signaling in cortical interneuron development in vitro in a mouse model.

#### 4.2 Unexplained variance

For the most significant discovery SNP identified in the Wang *et al.* study above (rs4307059), the risk allele frequency was 0.65 in cases with an odds ratio of 1.19, which is comparable with common variant discoveries in other psychiatric disorders including schizophrenia (Glessner & Hakonarson, 2009; Glessner *et al.*, 2010); bipolar disorder (Ferreira, 2008), and attention-deficit/hyperactivity disorder (Arcos-Burgos *et al.* 2010). While it is important not to undermine the significance of these findings, it should be noted that the predictive value of such ratios is relatively low (Dickson *et al.* 2010), often explaining less than 5% of the total risk (review at <http://www.genome.gov/26525384>). However, it is also possible that these common SNPs may be tagging a more rare causative variant (i.e., synthetic association), where the effect sizes may be markedly underestimated by the GWAS variant as we recently reported (Dickson *et al.* 2010). In one example, Wang *et al.* (2010), examined the *NOD2* locus as a cause of Crohn disease. Using resequencing data, they found that three causal variants explain > 5% of the genetic risk, where GWAS had estimated the risk at ~1%. Careful phenotyping of cohorts is important to ensure that the phenotypes produced by rare-variants are not being “filtered-out” and thereby missed as a consequence. A long range haplotype analysis of the GWAS data at the respective loci is therefore recommended in an attempt to enrich for individuals with rare-causative variants, who should be picked out of the cohort and subsequently sequenced for confirmation (Wang *et al.*, 2010). This is clearly a critical point, particularly in relation to psychiatric disorders, where diagnoses can be more contingent upon subjective observation than, for example, the genetics of height, which can utilize more intrinsically quantitative data.

The possibility that common variants are not the major cause of ASDs is also gaining increased support from the preponderance of copy number variation (CNV) studies, which are identifying rare variants with a stronger causal impact.

### 5. Copy number variation in ASDs

Copy number variations (CNVs) are insertions, deletions, or translocations in the human genome that are universal in the general population but more commonly found in genic

regions in individuals with neuropsychiatric disorders (e.g. Pinto *et al.*, 2010). CNVs can be detected by the same SNP arrays used in GWAS, and vary in length from many megabases to 1 kilobase or smaller. They are often not associated with any observable phenotype.

One of the most widely-known CNVs is Down syndrome, which is characterized by an extra chromosome 21. Rett syndrome is also caused by a CNV, which includes a deletion in *MECP2*. CNVs can be inherited or occur *de novo*, the cause of which is thus far unknown. Common disease-causing CNVs are infrequent but rare CNVs, with a frequency of less than 1%, have been identified for a range of disorders including ADHD (e.g. Williams *et al.*, 2010), schizophrenia (e.g. Glessner *et al.*, 2010; Levinson *et al.*, 2011), bipolar disorder (e.g. Chen *et al.*, 2010) and many others. A substantial portion of autism appears to be caused by rare CNVs. *De novo* CNVs that are greater than 100kb in size are more common in genic regions in individuals with ASDs than in the general population.

Sebat *et al.* (2007) provided some relevant early insights into the genomic features of CNVs. Firstly, they noted that *de novo* CNVs were individually rare - from 118 ASD cases, none of the identified variants were observed more than twice, with the majority seen just once. This confirmed the widely-held assumption that many different loci can contribute to the same ASD phenotype. Secondly, the authors affirmed the utility of population-study approaches that analyze sporadic and multiplex (i.e. more than one family member affected) families separately. The rate of *de novo* mutation in large (mostly genic) loci in multiplex families was significantly lower than for the sporadic cases ( $p = 0.04$ ). While this observation remains to be replicated in a larger study, the finding implies two mechanisms of genetic susceptibility - spontaneous mutation and inheritance. Finally, the sheer volume of loci identified by this approach (multiple loci on 20 chromosomes) affirms the extraordinarily complexity of ASDs.

A number of subsequent studies have greatly expanded the number of candidate loci. Our laboratory (Bucan *et al.* (2009)) reported 150+ CNVs in 912 ASD families that were not found in 1,488 controls. Critically, 27 of these loci were replicated in an independent cohort of 859 ASD cases and 1,051 controls. Some of the rare variants we identified had previously been associated with autism, including *NRXN1* and *UBE3A*, which are established ASD candidate genes (Guilmatre *et al.*, 2009). Samaco *et al.* (2005) previously identified significant deficits in *ube3a* expression in *mecp2*-deficient mice, suggesting a shared pathological pathway with Rett syndrome (as well as Angelman syndrome, and autism). Similarly, Kim *et al.* (2008) associated *NRXN1* with a balanced chromosomal abnormality at chromosome 2p16.3 in two unrelated ASD individuals. Rare variants in the coding region included two missense changes.

Glessner *et al.* (2009) identified and reported CNVs in two major gene networks, including neuronal cell adhesion molecules (such as *NRXN1*) and the ubiquitin gene family (such as *UBE3A*). Interestingly, four of the most prominent genes enriched by CNVs in ASD cases (*UBE3A*, *PARK2*, *RFWD2* and *FBXO40*) are all part of the ubiquitin gene family. Ubiquitination can alter protein function after translation, and degrade target proteins in conjunction with proteasomes. The ubiquitin-proteasome system operates at pre- and post-synapses, whose functions includes regulating neurotransmitter release, recycling synaptic vesicles in pre-synaptic terminals, and modulating changes in dendritic spines and post-synaptic density (Yi & Ehlers, 2005). As well as implicating an ubiquitination network in relation to ASDs, we also identified a second pathway involving *NRXN1*, *CNTN4*, *NLGN1*, and *ASTN2*. Genes in this group mediate neuronal cell-adhesion, and contribute to neurodevelopment by facilitating axon guidance, synapse formation and plasticity, and neuron-glia interactions. We also note that ubiquitins are involved in recycling cell-

adhesion molecules, which is a possible mechanism by which these two networks are cross linked.

In a similar approach, Pinto *et al.* (2010) further confirmed the importance of rare CNVs as causal factors for ASDs. Interestingly, the group did not observe a significant difference between cases and controls in terms of raw number of CNVs or estimated CNV size. However, the number of CNVs in genic regions was significantly greater in ASDs compared to controls. Again, loci enriched for CNVs include a number of genes known to be important for neurodevelopment and synaptic plasticity, such as *SHANK2*, *SYNGAP1*, and *DLGAP2*. Between 5.5% and 5.7% of ASD cases have at least one *de novo* CNV, further confirming the significance of *de novo* genetic events as risk factors for autism. Similar to the Glessner study, the Pinto group mapped CNVs to a series of networks involved in the development and regulation of the central nervous system functions. Implicated networks include neuronal cell adhesion, GTPase regulation (important for signal transduction and biosynthesis), and GTPase/Ras signaling, also involved in ubiquitination.

Finally, Gai *et al.* (2011) took a slightly different approach, focusing exclusively on inherited CNVs. While underlying loci were not necessarily common to those identified by the Glessner and Pinto groups, enrichment in pathways involving central nervous system development, synaptic functions and neuronal signaling processes was again confirmed. The Gai *et al.* study also emphasized the role of glutamate-mediated neuronal signals in ASDs.

Collectively, these CNV studies suggest that certain hotspots on the genome are particularly vulnerable to ASDs, which include loci on chromosomes 1q21, 3p26, 15q11-q13, 16p11, and 22q11. These hotspots are part of large gene networks that are important to neural signaling and neurodevelopment and have additionally been associated with other neuropsychiatric disorders.

In particular, a number of CNV studies in schizophrenia have highlighted structural mutations incorporating chromosomes 1q21, 15q13, and 22q11 (e.g. McClellan and King 2010; Glessner *et al.*, 2010), which are significantly enriched in cases versus controls, with *NRXN1* being a standout in this regard. From a phenotype perspective, autism and schizophrenia seem very different, both in behavioral manifestation and age of onset, and it may seem counter-intuitive that associated loci should overlap. Some authors have addressed this peculiarity by proposing that schizophrenia and autism may in fact be different poles of the same spectrum. Thus, Crespi and Braddock (2008) suggest that social cognition is underdeveloped in ASDs and over-developed in the psychotic spectrum, with a similar polarization of language and behavioral phenotypes. Although speculative, this hypothesis has gained some traction. In the next several years, genomic, imaging, and model-systems approaches will likely shed further light on the relationship between autism, schizophrenia and other neuropsychiatric disorders.

## 6. Sequencing familial forms of ASDs

To this point, we have focused primarily on the complex interactions of polygenic networks as the major cause of ASDs. However, this is not exclusively the case. Paralleling the recent spate of CNV is a renewed focus on rare disorders, including familial forms of complex diseases that potentially are monogenic or with less complex inheritance pattern. At the outset of this chapter, we emphasized the overlap with fragile X syndrome, where one third of cases are co-morbid for ASD. As mentioned, fragile X is caused by a failure to express the

protein coded by *FMR1*. However, mutations in *FMR1* do not always result in fragile-X and can result in a phenotype more representative of ASDs. Thus, Muhle *et al.* (2004) found that 7-8% of idiopathic ASD cases may have mutations at the *FMR1* locus. Likewise, although mutations in *MECP2* are the common cause of Rett syndrome, certain mutations at the same locus have been associated with idiopathic autism (Carney *et al.* (2003).

X-linked genes encoding neuroligins *NLGN3* and *NLGN4* and *SHANK3* (a neuroligin binding partner) are other prominent examples of distinct rare genetic causes, and a parallel can be drawn with these studies and mental retardation and epilepsy, which include many rare syndromes that collectively account for a substantial proportion of the two disorders (Morrow *et al.*, 2008). Indeed it is perhaps more than coincidence that autism is heavily comorbid with these two conditions, with >40% (Bölte *et al.*, 2009) and ~40% (Danielsson *et al.*, 2005) of ASD cases meeting diagnostic criteria for mental retardation and epilepsy respectively. It also is noteworthy that many of these monogenic-related genes are also major players in neurodevelopment and synapse activity. Other prominent examples include *TSC1*, *TSC2* (Osborne *et al.*, 1991; Franz, 1998), *NF1*, and *UBE3A* (see Morrow *et al.* (2008).

The identification of monogenic or possibly oligogenic autisms is likely to accelerate in the next several years as next-generation sequencing becomes more widely available. We recently encountered a family of two parents, six healthy siblings, and two siblings with severe autism suggestive of autosomal recessive inheritance. Unsuccessful attempts using linkage and CNV approaches failed to identify a causal locus, but whole-exome sequencing at 20x coverage identified four genes, including one with a non-synonymous SNP in the protocadherin alpha 4 isoform1 precursor (*PCDHA4*) gene, which presents a strong candidate gene, currently under validation. Protocadherins are part of the cadherin family that facilitates neuronal cell adhesion and this discovery is consistent genomically and neurobiologically with the findings addressed above in relation to *CDH9* and *CDH10*.

Known syndromes with ASD features include fragile-x, neurofibromatosis type 1, down syndrome, tuberous sclerosis, neurofibromatosis (which confers a 100-fold increased risk for ASDs Li *et al.* (2005), Angelman, Prader-Willi and related 15q syndromes, and at least several dozen others (see Zafeiriou *et al.*, 2007 for a comprehensive review). **Table 1** from Volkmar *et al.* (2005) lists the most commonly associated syndromes with median rate and range. It is likely that many more unidentified rare syndromes with Mendelian causes have ASD phenotypes. As of March 2011, the Online Mendelian Inheritance in Man (OMIM) database listed 6,727 known or suspected Mendelian diseases (MD), with 2,993 (44%) of these having an identified molecular basis. Since OMIM derives its data from published reports, these figures likely under-represent rare disorders, which may go unreported. It has been proposed that as many as 30,000 genetic disorders may exist, suggesting that many Mendelian disorders have no genetic etiology identified to date. Given the large-representation of autism phenotypes in known syndromes, we can assume a similar trend in unreported disease.

It remains to be determined whether rare variants will account for the majority of autisms. Irrespective, as with many other aspects of scientific inquiry, the study of rare variants will continue to play an important role in explicating the pathogenesis of ASDs. El-Fishawy and State (2010) point to hypercholesterolemia and hypertension (Brown, 1974; Lifton *et al.*, 2001) as examples where rare mutations have been successful in driving a molecular understanding of the disease as opposed to identifying risk factors in the general

population. Rare mutations, particularly when they are Mendelian, carry large effects and are typically in genic regions. These characteristics make the resolution of underlying networks distinctly less complex and, moreover, are amenable to modeling in other systems. Recent groundbreaking studies by Marchetto *et al.* (2010) and Muotri *et al.* (2010), who created a cell culture model of Rett syndrome, are potentially exciting developments in this regard. Here, the researchers used skin biopsies from four Rett's patients, each carrying a different *MeCP2* mutation, to culture induced pluripotent stem cells (iPS). Once the iPS cells developed into neurons, they showed a decreased number of neurons and dendritic spines, consistent with neurodevelopmental disruptions. Intervention with insulin-like growth factor 1 (*IGF1*), which is known to regulate neurodevelopment, was subsequently shown to reverse Rett-like symptoms in a mouse model of the disease. This innovative approach is an exciting model of how rare gene approaches can stimulate our understanding of the pathophysiology and potential reversibility of ASDs.

Syndrome	Number of Studies	Median Rate	Range %
Tuberous sclerosis	11	1.1	0-3.8
Fragile X	9	0.0	0-8.1
Down syndrome	12	0.7	0-16.7
Neurofibromatosis 1	6	0	0-1.4

Table 1. Associated disorders and their rate in autism (from Volkmar *et al.*, 2005 in Zafeiriou *et al.* 2007)

## 7. Conclusions

ASDs are clearly highly heritable disorders and advances in gene-finding technology in the past decade have rapidly accelerated gene discovery. As is typically the case, successive developments have made the problem more complex such that there are dozens of candidate genes, many of which remain to be replicated. In spite of this complexity, we can observe a number of patterns beginning to unfold 1) the relative scarcity of causal common variants, 2) the growing list of causal rare variants, and 3) the emergence of monogenic disorders with primary and secondary ASD phenotypes.

The monogenic autisms are particularly interesting from a treatment perspective, as they provide a mechanism for studying ASD phenotypes in model systems and an obvious target for drug intervention. They are also amenable to clinical testing and the decreasing cost of research technologies means that this capacity is more widely available to clinicians. In fact, as the resolution of clinical instruments becomes more sophisticated, it is likely that the clinic will become a primary workplace for syndromic discovery.

A key requirement in driving gene discovery is the necessity of high-quality phenotype data. ASDs are notoriously heterogeneous, and are fractionated in terms of symptoms and trajectory. Mandy & Skuse (2008) reviewed seven factor analysis studies of ASDs symptoms, and found that all but one dissociated social and non-social factors. In a non-clinical sample of 3,000 twin pairs, Happé *et al.* (2006) examined autistic-like traits and found consistently low correlations ( $r = 0.1-0.4$ ) between each of the core deficits on the autism spectrum. Endophenotypes, sub-components or sub-processes of the broader phenotype, may provide a productive avenue to disentangling some of this complexity. By filtering out all but a few discrete measures, we can theoretically increase the signal-to-noise ratio in genotype-phenotype associations. A number of endophenotypes for ASDs have been identified

associated with disease genes, including head circumference (associated with the *HOXA1* A218G polymorphism, Conciatori *et al.*, 2004), age at first word (associated with a quantitative trait locus on 7q35, Alarcón *et al.* 2005), delayed magnetoencephalography evoked responses to auditory stimuli (Roberts *et al.*, 2010), and enhanced perception (Mottron *et al.*, 2006). The endophenotype approach is arguably more consistent with rare-/mono-genic discovery, where a mutated network may not yield a diagnosis of autism *per se*, but nevertheless cause associated abnormalities. Note, this approach does not diminish the pleiotropic effects of genes involved in neurodevelopment, and only serves to make the point that the relevant genotype may associate with some but not all ASD features.

The converse, of course, is also true with a large number of candidate genes contributing to the majority of known ASDs. With ~80% of genes expressed in the brain it is likely that this number will continue to grow, and here again careful phenotyping is critical to identifying functional consequences. Ultimately, the primary goal is not to determine the frequency of variation/mutation in cases *versus* controls, but to determine the pathway(s) and gene networks that lead to pathology. This will be no mean feat, with other major players such as epigenetic factors, RNA regulatory elements, and environmental exposures also an important part of the equation. While daunting, the elucidation of these elements will doubtlessly take us closer to developing effective treatments for ASDs. Given the current rate of progress, we have cause for cautious optimism in this regard.

## 8. References

- Alarcon, M., Abrahams, B. S., Stone, J. L., Duvall, J. A., Perederiy, J. V., Bomar, J. M., Geschwind, D. H. (2008). Linkage, association, and gene-expression analyses identify CNTNAP2 as an autism-susceptibility gene. *Am J Hum Genet*, 82(1), 150-159. doi: 10.1016/j.ajhg.2007.09.005
- Alarcon, M., Yonan, A. L., Gilliam, T. C., Cantor, R. M., & Geschwind, D. H. (2005). Quantitative genome scan and Ordered-Subsets Analysis of autism endophenotypes support language QTLs. *Mol Psychiatry*, 10(8), 747-757. doi: 10.1038/sj.mp.4001666
- Alarcon, M., Cantor, R. M., Liu, J., Gilliam, T. C., & Geschwind, D. H. (2002). Evidence for a language quantitative trait locus on chromosome 7q in multiplex autism families. *Am J Hum Genet*, 70(1), 60-71. doi: 10.1086/338241
- Arcos-Burgos, M., Jain, M., Acosta, M. T., Shively, S., Stanescu, H., Wallis, D., Muenke, M. (2010). A common variant of the latrophilin 3 gene, LPHN3, confers susceptibility to ADHD and predicts effectiveness of stimulant medication. *Mol Psychiatry*, 15(11), 1053-1066. doi: 10.1038/mp.2010.6
- American Psychiatric Association, (2010, March 22, 2010). New Proposed Changes Posted for Leading Manual of Mental Disorders. Retrieved from <http://www.dsm5.org/Pages/Default.aspx>
- Arking, D. E., Cutler, D. J., Brune, C. W., Teslovich, T. M., West, K., Ikeda, M., Chakravarti, A. (2008). A common genetic variant in the neurexin superfamily member CNTNAP2 increases familial risk of autism. *Am J Hum Genet*, 82(1), 160-164. doi: 10.1016/j.ajhg.2007.09.015
- Autism and Developmental Disabilities Monitoring Network Surveillance Year 2006 (2009). Prevalence of autism spectrum disorders-Autism and Developmental Disabilities


- Monitoring Network *MMWR Surveillance Summary*, 58, 1-20. United States, 2006: Centers for Disease Control and Prevention.
- Bailey, A., Le Couteur, A., Gottesman, I., Bolton, P., Simonoff, E., Yuzda, E., & Rutter, M. (1995). Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol Med*, 25(1), 63-77.
- Baird, G., Simonoff, E., Pickles, A., Chandler, S., Loucas, T., Meldrum, D., & Charman, T. (2006). Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *Lancet*, 368(9531), 210-215. doi: 10.1016/s0140-6736(06)69041-7
- Benayed, R., Choi, J., Matteson, P. G., Gharani, N., Kamdar, S., Brzustowicz, L. M., & Millonig, J. H. (2009). Autism-associated haplotype affects the regulation of the homeobox gene, ENGRAILED 2. *Biol Psychiatry*, 66(10), 911-917. doi: 10.1016/j.biopsych.2009.05.027
- Benayed, R., Gharani, N., Rossman, I., Mancuso, V., Lazar, G., Kamdar, S., . Millonig, J. H. (2005). Support for the homeobox transcription factor gene ENGRAILED 2 as an autism spectrum disorder susceptibility locus. *Am J Hum Genet*, 77(5), 851-868. doi: 10.1086/497705
- Bolte, S., Dziobek, I., & Poustka, F. (2009). Brief report: The level and nature of autistic intelligence revisited. *J Autism Dev Disord*, 39(4), 678-682. doi: 10.1007/s10803-008-0667-2
- Brown, M. S., & Goldstein, J. L. (1974). Expression of the familial hypercholesterolemia gene in heterozygotes: mechanism for a dominant disorder in man. *Science*, 185(4145), 61-63.
- Bucan, M., Abrahams, B. S., Wang, K., Glessner, J. T., Herman, E. I., Sonnenblick, L. I., Hakonarson, H. (2009). Genome-wide analyses of exonic copy number variants in a family-based study point to novel autism susceptibility genes. *PLoS Genet*, 5(6), e1000536. doi: 10.1371/journal.pgen.1000536
- Campbell, D. B., Li, C., Sutcliffe, J. S., Persico, A. M., & Levitt, P. (2008). Genetic evidence implicating multiple genes in the MET receptor tyrosine kinase pathway in autism spectrum disorder. *Autism Res*, 1(3), 159-168. doi: 10.1002/aur.27
- Campbell, D. B., Sutcliffe, J. S., Ebert, P. J., Militerni, R., Bravaccio, C., Trillo, S., . . . Levitt, P. (2006). A genetic variant that disrupts MET transcription is associated with autism. *Proc Natl Acad Sci U S A*, 103(45), 16834-16839. doi: 10.1073/pnas.0605296103
- Carney, R. M., Wolpert, C. M., Ravan, S. A., Shahbazian, M., Ashley-Koch, A., Cuccaro, M. L., . . . Pericak-Vance, M. A. (2003). Identification of MeCP2 mutations in a series of females with autistic disorder. *Pediatr Neurol*, 28(3), 205-211.
- Chen, X., Li, X., Wang, P., Liu, Y., Zhang, Z., Zhao, G., Kong, X. (2010). Novel association strategy with copy number variation for identifying new risk Loci of human diseases. *PLoS One*, 5(8), e12185. doi: 10.1371/journal.pone.0012185
- Cheng, Y., Sudarov, A., Szulc, K. U., Sgaier, S. K., Stephen, D., Turnbull, D. H., & Joyner, A. L. (2010). The Engrailed homeobox genes determine the different foliation patterns in the vermis and hemispheres of the mammalian cerebellum. *Development*, 137(3), 519-529. doi: 10.1242/dev.027045
- Cisse, M., Halabisky, B., Harris, J., Devidze, N., Dubal, D. B., Sun, B., Mucke, L. (2011). Reversing EphB2 depletion rescues cognitive functions in Alzheimer model. *Nature*, 469(7328), 47-52. doi: 10.1038/nature09635

- Conciatori, M., Stodgell, C. J., Hyman, S. L., O'Bara, M., Militerni, R., Bravaccio, C., .Persico, A. M. (2004). Association between the HOXA1 A218G polymorphism and increased head circumference in patients with autism. *Biol Psychiatry*, 55(4), 413-419. doi: 10.1016/j.biopsych.2003.10.005
- Crespi, B., & Badcock, C. (2008). Psychosis and autism as diametrical disorders of the social brain. *Behav Brain Sci*, 31(3), 241-261; discussion 261-320. doi: 10.1017/s0140525x08004214
- Curtis, A. R., Headland, S., Lindsay, S., Thomas, N. S., Boye, E., Kamakari, S., et al. (1993). X chromosome linkage studies in familial Rett syndrome. *Hum Genet*, 90(5), 551-555.
- Danielsson, S., Gillberg, I. C., Billstedt, E., Gillberg, C., & Olsson, I. (2005). Epilepsy in young adults with autism: a prospective population-based follow-up study of 120 individuals diagnosed in childhood. *Epilepsia*, 46(6), 918-923. doi: 10.1111/j.1528-1167.2005.57504.x
- Dickson, S. P., Wang, K., Krantz, I., Hakonarson, H., & Goldstein, D. B. (2010). Rare variants create synthetic genome-wide associations. *PLoS Biol*, 8(1), e1000294. doi: 10.1371/journal.pbio.1000294
- Eagleson, K. L., Campbell, D. B., Thompson, B. L., Bergman, M. Y., & Levitt, P. (2011). The autism risk genes MET and PLAU8 differentially impact cortical development. *Autism Res*, 4(1), 68-83. doi: 10.1002/aur.172
- El-Fishawy, P., & State, M. W. (2010). The genetics of autism: key issues, recent findings, and clinical implications. *Psychiatr Clin North Am*, 33(1), 83-105. doi: 10.1016/j.psc.2009.12.002
- Ferreira, M. A., O'Donovan, M. C., Meng, Y. A., Jones, I. R., Ruderfer, D. M., Jones, L., Craddock, N. (2008). Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat Genet*, 40(9), 1056-1058. doi: 10.1038/ng.209
- Fombonne, E. (2009). Epidemiology of pervasive developmental disorders. *Pediatr Res*, 65(6), 591-598. doi: 10.1203/PDR.0b013e31819e7203
- Franz, D. N. (1998). Diagnosis and management of tuberous sclerosis complex. *Semin Pediatr Neurol*, 5(4), 253-268.
- Gai, X., Xie, H. M., Perin, J. C., Takahashi, N., Murphy, K., Wenocur, A. S., .White, P. S. (2011). Rare structural variation of synapse and neurotransmission genes in autism. *Mol Psychiatry*. doi: 10.1038/mp.2011.10
- Gepner, B., & Feron, F. (2009). Autism: a world changing too fast for a mis-wired brain? *Neurosci Biobehav Rev*, 33(8), 1227-1242. doi: 10.1016/j.neubiorev.2009.06.006
- Glessner, J. T., & Hakonarson, H. (2009). Common variants in polygenic schizophrenia. *Genome Biol*, 10(9), 236. doi: 10.1186/gb-2009-10-9-236
- Glessner, J. T., Reilly, M. P., Kim, C. E., Takahashi, N., Albano, A., Hou, C., Hakonarson, H. (2010). Strong synaptic transmission impact by copy number variations in schizophrenia. *Proc Natl Acad Sci U S A*, 107(23), 10584-10589. doi: 10.1073/pnas.1000274107
- Glessner, J. T., Wang, K., Cai, G., Korvatska, O., Kim, C. E., Wood, S., Hakonarson, H. (2009). Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature*, 459(7246), 569-573. doi: 10.1038/nature07953
- Guilmatre, A., Dubourg, C., Mosca, A. L., Legallic, S., Goldenberg, A., Drouin-Garraud, V., Campion, D. (2009). Recurrent rearrangements in synaptic and

- neurodevelopmental genes and shared biologic pathways in schizophrenia, autism, and mental retardation. *Arch Gen Psychiatry*, 66(9), 947-956. doi: 10.1001/archgenpsychiatry.2009.80
- Hagerman, R. J., Berry-Kravis, E., Kaufmann, W. E., Ono, M. Y., Tartaglia, N., Lachiewicz, A., Tranfaglia, M. (2009). Advances in the treatment of fragile X syndrome. *Pediatrics*, 123(1), 378-390. doi: 10.1542/peds.2008-0317
- Happé, F., Ronald, A., & Plomin, R. (2006). Time to give up on a single explanation for autism. *Nat Neurosci*, 9(10), 1218-1220. doi: 10.1038/nn1770
- Harris, S. W., Hessel, D., Goodlin-Jones, B., Ferranti, J., Bacalman, S., Barbato, I., Hagerman, R. J. (2008). Autism profiles of males with fragile X syndrome. *Am J Ment Retard*, 113(6), 427-438.
- International Molecular Genetic Study of Autism Consortium (IMGSAC). A genomewide screen for autism: strong evidence for linkage to chromosomes 2q, 7q, and 16p. (2001). *Am J Hum Genet*, 69(3), 570-581. doi: 10.1086/323264
- Kim, H. G., Kishikawa, S., Higgins, A. W., Seong, I. S., Donovan, D. J., Shen, Y., Gusella, J. F. (2008). Disruption of neurexin 1 associated with autism spectrum disorder. *Am J Hum Genet*, 82(1), 199-207. doi: 10.1016/j.ajhg.2007.09.011
- Lai, C. S., Fisher, S. E., Hurst, J. A., Vargha-Khadem, F., & Monaco, A. P. (2001). A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature*, 413(6855), 519-523. doi: 10.1038/35097076
- Lauritsen, M. B., Pedersen, C. B., & Mortensen, P. B. (2005). Effects of familial risk factors and place of birth on the risk of autism: a nationwide register-based study. *J Child Psychol Psychiatry*, 46(9), 963-971. doi: 10.1111/j.1469-7610.2004.00391.x
- Levinson, D. F., Duan, J., Oh, S., Wang, K., Sanders, A. R., Shi, J., Gejman, P. V. (2011). Copy Number Variants in Schizophrenia: Confirmation of Five Previous Findings and New Evidence for 3q29 Microdeletions and VIPR2 Duplications. *Am J Psychiatry*, 168(3), 302-316. doi: 10.1176/appi.ajp.2010.10060876
- Levy, S. E., Mandell, D. S., & Schultz, R. T. (2009). Autism. *Lancet*, 374(9701), 1627-1638. doi: 10.1016/s0140-6736(09)61376-3
- Li, W., Cui, Y., Kushner, S. A., Brown, R. A., Jentsch, J. D., Frankland, P. W., Silva, A. J. (2005). The HMG-CoA reductase inhibitor lovastatin reverses the learning and attention deficits in a mouse model of neurofibromatosis type 1. *Curr Biol*, 15(21), 1961-1967. doi: 10.1016/j.cub.2005.09.043
- Lifton, R. P., Gharavi, A. G., & Geller, D. S. (2001). Molecular mechanisms of human hypertension. *Cell*, 104(4), 545-556.
- Ma, D., Salyakina, D., Jaworski, J.M., Konidari, I., Whitehead, P.L., Andersen, A.N., ...Pericak-Vance MA (2009). *Ann. Hum. Genet.* 73, 263-273.
- Mandy, W. P., & Skuse, D. H. (2008). Research review: What is the association between the social-communication element of autism and repetitive interests, behaviours and activities? *J Child Psychol Psychiatry*, 49(8), 795-808. doi: 10.1111/j.1469-7610.2008.01911.x
- Marchetto, M. C., Carroumeu, C., Acab, A., Yu, D., Yeo, G. W., Mu, Y., Muotri, A. R. (2010). A model for neural development and treatment of Rett syndrome using human induced pluripotent stem cells. *Cell*, 143(4), 527-539. doi: 10.1016/j.cell.2010.10.016
- McClellan, J., & King, M. C. (2010). Genetic heterogeneity in human disease. *Cell*, 141(2), 210-217. doi: 10.1016/j.cell.2010.03.032

- Morrow, E. M., Yoo, S. Y., Flavell, S. W., Kim, T. K., Lin, Y., Hill, R. S., Walsh, C. A. (2008). Identifying autism loci and genes by tracing recent shared ancestry. *Science*, 321(5886), 218-223. doi: 10.1126/science.1157657
- Mottron, L., Dawson, M., Soulières, I., Hubert, B., & Burack, J. (2006). Enhanced perceptual functioning in autism: an update, and eight principles of autistic perception. *J Autism Dev Disord*, 36(1), 27-43. doi: 10.1007/s10803-005-0040-7
- Muhle, R., Trentacoste, S. V., & Rapin, I. (2004). The genetics of autism. *Pediatrics*, 113(5), e472-486.
- Muller, R. A., Shih, P., Keehn, B., Deyoe, J. R., Leyden, K. M., & Shukla, D. K. (2011). Underconnected, but How? A Survey of Functional Connectivity MRI Studies in Autism Spectrum Disorders. *Cereb Cortex*. doi: 10.1093/cercor/bhq296
- Muotri, A. R., Marchetto, M. C., Coufal, N. G., Oefner, R., Yeo, G., Nakashima, K., & Gage, F. H. (2010). L1 retrotransposition in neurons is modulated by MeCP2. *Nature*, 468(7322), 443-446. doi: 10.1038/nature09544
- Newschaffer, C. J., Croen, L. A., Daniels, J., Giarelli, E., Grether, J. K., Levy, S. E., . Windham, G. C. (2007). The epidemiology of autism spectrum disorders. *Annu Rev Public Health*, 28, 235-258. doi: 10.1146/annurev.publhealth.28.021406.144007
- O'Roak, B. J., & State, M. W. (2008). Autism genetics: strategies, challenges, and opportunities. *Autism Res*, 1(1), 4-17. doi: 10.1002/aur.3
- Osborne, J. P., Fryer, A., & Webb, D. (1991). Epidemiology of tuberous sclerosis. *Ann N Y Acad Sci*, 615, 125-127.
- Pinto, D., Pagnamenta, A. T., Klei, L., Anney, R., Merico, D., Regan, R., Betancur, C. (2010). Functional impact of global rare copy number variation in autism spectrum disorders. *Nature*, 466(7304), 368-372. doi: 10.1038/nature09146
- Piven, J., Palmer, P., Jacobi, D., Childress, D., & Arndt, S. (1997). Broader autism phenotype: evidence from a family history study of multiple-incidence autism families. *Am J Psychiatry*, 154(2), 185-190.
- Ramocki, M. B., & Zoghbi, H. Y. (2008). Failure of neuronal homeostasis results in common neuropsychiatric phenotypes. *Nature*, 455(7215), 912-918. doi: 10.1038/nature07457
- Redies, C. (2000). Cadherins in the central nervous system. *Prog Neurobiol*, 61(6), 611-648.
- Richards, R. I., Holman, K., Kozman, H., Kremer, E., Lynch, M., Pritchard, M., Sutherland, G. R. (1991). Fragile X syndrome: genetic localisation by linkage mapping of two microsatellite repeats FRAXAC1 and FRAXAC2 which immediately flank the fragile site. *J Med Genet*, 28(12), 818-823.
- Risch, N., & Merikangas, K. (1996). The future of genetic studies of complex human diseases. *Science*, 273(5281), 1516-1517.
- Roberts, T. P., Khan, S. Y., Rey, M., Monroe, J. F., Cannon, K., Blaskey, L., .Edgar, J. C. (2010). MEG detection of delayed auditory evoked responses in autism spectrum disorders: towards an imaging biomarker for autism. *Autism Res*, 3(1), 8-18. doi: 10.1002/aur.111
- Rogers, S. J., Wehner, D. E., & Hagerman, R. (2001). The behavioral phenotype in fragile X: symptoms of autism in very young children with fragile X syndrome, idiopathic autism, and other developmental disorders. *J Dev Behav Pediatr*, 22(6), 409-417.
- Samaco, R. C., Hogart, A., & LaSalle, J. M. (2005). Epigenetic overlap in autism-spectrum neurodevelopmental disorders: MECP2 deficiency causes reduced expression of UBE3A and GABRB3. *Hum Mol Genet*, 14(4), 483-492. doi: 10.1093/hmg/ddi045

- Sebat, J., Lakshmi, B., Malhotra, D., Troge, J., Lese-Martin, C., Walsh, T., Wigler, M. (2007). Strong association of de novo copy number mutations with autism. *Science*, 316(5823), 445-449. doi: 10.1126/science.1138659
- Skuse, D. H. (2007). Rethinking the nature of genetic vulnerability to autistic spectrum disorders. *Trends Genet*, 23(8), 387-395. doi: 10.1016/j.tig.2007.06.003
- Sousa, I., Clark, T. G., Toma, C., Kobayashi, K., Choma, M., Holt, R., Monaco, A. P. (2009). MET and autism susceptibility: family and case-control studies. *Eur J Hum Genet*, 17(6), 749-758. doi: 10.1038/ejhg.2008.215
- St Pourcain, B., Wang, K., Glessner, J.T., Golding, J., Steer, C., Ring, S.M., Skuse, D.H., Grant, S.F., Hakonarson, H., and Smith, G.D. (2010). Am. J. Psychiatry. Published online July 15, 2010. 10.1176/appi.ajp.2010.09121789.
- Vernes, S. C., Newbury, D. F., Abrahams, B. S., Winchester, L., Nicod, J., Groszer, M., Fisher, S. E. (2008). A functional genetic link between distinct developmental language disorders. *N Engl J Med*, 359(22), 2337-2345. doi: 10.1056/NEJMoa0802828
- Volkmar, F. R. (2005). *Handbook of Autism and Pervasive Developmental Disorders* (3 ed.). New Jersey: John Wiley & Sons.
- Wang, K., Zhang, H., Ma, D., Bucan, M., Glessner, J. T., Abrahams, B. S., Hakonarson, H. (2009). Common genetic variants on 5p14.1 associate with autism spectrum disorders. *Nature*, 459(7246), 528-533. doi: 10.1038/nature07999
- Wang K, Dickson SP, Stolle CA, Krantz ID, Goldstein DB, Hakonarson H. Interpretation of association signals and identification of causal variants from genome-wide association studies. *Am J Hum Genet*. 2010 May 14;86(5):730-42. Epub 2010 Apr 29.
- Wang, L., Jia, M., Yue, W., Tang, F., Qu, M., Ruan, Y., Zhang, D. (2008). Association of the ENGRAILED 2 (EN2) gene with autism in Chinese Han population. *Am J Med Genet B Neuropsychiatr Genet*, 147B(4), 434-438. doi: 10.1002/ajmg.b.30623
- Yi, J. J., & Ehlers, M. D. (2005). Ubiquitin and protein turnover in synapse function. *Neuron*, 47(5), 629-632. doi: 10.1016/j.neuron.2005.07.008
- Zafeiriou, D. I., Ververi, A., & Vargiami, E. (2007). Childhood autism and associated comorbidities. *Brain Dev*, 29(5), 257-272. doi: 10.1016/j.braindev.2006.09.003
- Zhong, H., Serajee, F. J., Nabi, R., & Huq, A. H. (2003). No association between the EN2 gene and autistic disorder. *J Med Genet*, 40(1), e4.



## **Autism Spectrum Disorders: The Role of Genetics in Diagnosis and Treatment**

Edited by Prof. Stephen Deutsch

ISBN 978-953-307-495-5

Hard cover, 198 pages

**Publisher** InTech

**Published online** 01, August, 2011

**Published in print edition** August, 2011

Estimated prevalence rates of autism spectrum disorders (ASDs) have increased at an alarming rate over the past decade; current estimates stand as high as 1 in 110 persons in the population with a higher ratio of affected males to females. In addition to their emotional impact on the affected persons and their family members (in fact, the latter are often unrecognized unaffected "patients" themselves), the economic and social impacts of ASDs on society are staggering. Persons with ASDs will need interdisciplinary approaches to complex treatment and life planning, including, but not limited to, special education, speech and language therapy, vocational skills training and rehabilitation, social skills training and cognitive remediation, in addition to pharmacotherapy. The current book highlights some of the recent research on nosology, etiology, and pathophysiology. Additionally, the book touches on the implications of new research for treatment and genetic counseling. Importantly, because the field is advancing rapidly, no book can be considered the final word or finished product; thus, the availability of open access rapid publication is a mechanism that will help to assure that readers remain current and up-to-date.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

John J.M. Connolly and Hakon Hakonarson (2011). The Genetics of Autism Spectrum Disorders, Autism Spectrum Disorders: The Role of Genetics in Diagnosis and Treatment, Prof. Stephen Deutsch (Ed.), ISBN: 978-953-307-495-5, InTech, Available from: <http://www.intechopen.com/books/autism-spectrum-disorders-the-role-of-genetics-in-diagnosis-and-treatment/the-genetics-of-autism-spectrum-disorders>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.