

# The geometry of right angled Artin subgroups of mapping class groups

Matt T. Clay\*, Christopher J. Leininger† and Johanna Mangahas‡

July 30, 2010

## Abstract

We describe sufficient conditions which guarantee that a finite set of mapping classes generate a right-angled Artin group quasi-isometrically embedded in the mapping class group. Moreover, under these conditions, the orbit map to Teichmüller space is a quasi-isometric embedding for both of the standard metrics. As a consequence, we produce infinitely many genus  $h$  surfaces (for any  $h$  at least 2) in the moduli space of genus  $g$  surfaces (for any  $g$  at least 3) for which the universal covers are quasi-isometrically embedded in the Teichmüller space.

## 1 Introduction

Let  $S$  denote a surface and  $\text{Mod}(S)$  its mapping class group. Given independent pseudo-Anosov mapping classes  $f_1, \dots, f_n \in \text{Mod}(S)$ , McCarthy [35] and Ivanov [21] proved that by passing to sufficiently high powers, these mapping classes generate a free subgroup. This is the primary ingredient in the proof that  $\text{Mod}(S)$  satisfies the “Tits alternative”; see also [14, 32] for quantitative versions of this. Farb and Mosher [13] defined a notion of convex cocompactness for subgroups of  $\text{Mod}(S)$  by way of analogy with Kleinian groups, and proved that  $f_1, \dots, f_n$  could be raised to sufficiently high powers to further guarantee that the subgroup they generate is convex cocompact; see also [37, 22, 18].

Given an arbitrary set of elements  $f_1, \dots, f_n \in \text{Mod}(S)$ , we cannot expect that they generate a free group upon raising to sufficiently high powers. However, Koberda [25] has recently proven that the powers do generate a right-angled Artin group; see also [9, 12, 8] for partial results in this direction.

In this paper, we are interested in geometric properties of right-angled Artin subgroups of the mapping class group. As convex cocompact subgroups are necessarily Gromov hyperbolic, we must consider other geometric properties for non-free right-angled Artin subgroups of  $\text{Mod}(S)$ . For example, Crisp and Wiest [12] produced quasi-isometric embeddings of certain right-angled Artin groups into braid groups (and hence also mapping class groups). In this paper we show that this is possible in much greater generality, and furthermore, one can often conclude even stronger geometric statements for the corresponding subgroups. Here we state

---

\*Partially supported by NSF grant DMS-1006898.

†Partially supported by NSF grant DMS-0905748.

‡Partially supported by NSF RTG grant 0602191.

our main theorem, and refer the reader to Section 2 for necessary terminology and a more precise statement (Theorem 2.2).

**Theorem 1.1.** *Suppose  $f_1, \dots, f_n \in \text{Mod}(S)$  are fully supported on overlapping nonannular subsurfaces. Then after raising to sufficiently high powers, these elements generate a quasi-isometrically embedded right-angled Artin subgroup of  $\text{Mod}(S)$ . Furthermore, the orbit map to the Teichmüller space is a quasi-isometric embedding for both of the standard metrics, namely the Teichmüller and Weil–Petersson metrics.*

**Remarks.**

1. We note that for the second statement to hold, the assumption that the support of each  $f_i$  is not an annulus is necessary. On the other hand, it seems likely that the homomorphism to  $\text{Mod}(S)$  is a quasi-isometry without this additional assumption.
2. There are a number of other “natural” metrics on Teichmüller space besides the two we have mentioned; the Bergman metric, Carathéodory metric, McMullen metric, Kähler–Einstein metric, Ricci metric and perturbed Ricci metric. However, each of these is quasi-isometric to the Teichmüller metric (see [36, 42, 29, 30]), and so the conclusion of Theorem 1.1 also holds for any of these metrics.

In section 6 we use the ideas from the proof of this theorem to describe the Thurston type of any element in the right-angled Artin subgroup of  $\text{Mod}(S)$  we construct, and we see that it is pseudo-Anosov on the largest possible subsurface. In particular, we describe exactly which elements are pseudo-Anosov on  $S$ ; see Theorem 6.1.

The hypotheses in Theorem 1.1 are general enough to easily provide quasi-isometric embeddings of any right-angled Artin group into some mapping class group (see the end of Section 2.4). In particular we have the following.

**Corollary 1.2.** *Any right-angled Artin group admits a homomorphism to some mapping class group which is a quasi-isometric embedding, and for which the orbit map to Teichmüller space is a quasi-isometric embedding with respect to either of the standard metrics.  $\square$*

The fundamental group of a closed orientable surface (of genus  $h \geq 2$ ) is called a (*genus  $h$  surface subgroup*). Many right-angled Artin groups contain quasi-isometrically embedded surface subgroups; see [40, 11] (though the question of exactly which right-angled Artin groups contain surface subgroups is still open; see for example [16, 23, 24, 10, 39]). There are also constructions of surface subgroups of the mapping class group [1, 26, 15]. In [27], infinitely many nonconjugate surface subgroups were constructed with geometric properties akin to geometric finiteness in the setting of Kleinian groups. From an explicit version of Corollary 1.2, and the aforementioned examples of surface subgroups of right-angled Artin groups, we obtain the following. See Section 5 for the proof.

**Corollary 1.3.** *For any closed surface  $S$  of genus at least 3 and any  $h \geq 2$ , there exist infinitely many nonconjugate genus  $h$  surface subgroups of  $\text{Mod}(S)$ , each of which act cocompactly on some quasi-isometrically embedded hyperbolic plane in the Teichmüller space  $\mathcal{T}(S)$ , with either of the standard metrics.*

This corollary is in contrast to the work of Bowditch [5] who proves finiteness, for any fixed  $h \geq 2$ , for the number of conjugacy classes of genus  $h$  surface subgroups of  $\text{Mod}(S)$  which are purely pseudo-Anosov (we note that surface subgroups of the mapping class group which arise

as subgroups of right-angled Artin groups can never be purely pseudo-Anosov; see Proposition 7.1 below). While these surface subgroups are not purely pseudo-Anosov, by the corollary, they do have the closely related property that every nontrivial element has positive translation length on  $\mathcal{T}(S)$ .

Finally, we remark that while Bowditch’s result mentioned above is an example of a kind of rank-1 phenomenon for  $\text{Mod}(S)$ , our examples illustrate higher rank behavior. Specifically, we could compare our results with those of Wang [41], who finds infinitely many conjugacy classes of discrete, faithful representations of right-angled Artin groups (hence surface subgroups) into higher rank Lie groups. Furthermore, Long, Reid and Thistlethwaite [31], find infinitely many conjugacy classes of Zariski dense, purely semi-simple representations of a surface group into  $SL(3, \mathbb{Z})$ . In fact, these surface groups are very closely related to the ones we study, in the sense that every nontrivial element has positive translation length on the associated symmetric space.

## 1.1 Plan of the paper

We begin in Section 2 by setting up the relevant definitions and notation we will use throughout. The section ends with a more precise version of our main theorem (Theorem 2.2). In Section 3 we describe an alternative space on which  $\text{Mod}(S)$  acts, namely Masur and Minsky’s graph of *markings* [33]. We also state the required *distance formulas* (Theorems 3.1, 3.2 and 3.3) which provide the coarse estimates for the distances in the desired spaces,  $\text{Mod}(S)$  and  $\mathcal{T}(S)$ , in terms of sums of “local distances” between pairs of markings. These local distances are precisely the *subsurface distances*, also described in this section.

The idea of the proof of Theorem 2.2 is as follows. The hypothesis implies that each of the generators of the right-angled Artin group corresponds to a mapping class which makes progress in some subsurface—that is, it contributes nontrivially to some local distance. A geodesic in the Cayley graph of the right-angled Artin group determines a sequence of mapping classes, each of which makes progress in some subsurface. We need only ensure that this progress accumulates (that is, we need to avoid cancellation of local distances). This is verified by Theorem 5.2, which relates a partial order on the set of syllables in a minimal length representative for an element of the right-angled Artin group (see Section 4) with the partial order from [3] on the set of subsurfaces “between” a marking and its image under the associated mapping class (see Section 3.4). The details of the proof of Theorem 5.2 are carried out in Section 5, followed by the proof of Theorem 2.2.

In Section 6 we find the Thurston type of each element in the right-angled Artin subgroups of  $\text{Mod}(S)$  we are considering. We show that by conjugating to use the minimal number of generators to represent the element, it will be pseudo-Anosov on the smallest subsurface filled by the supports of the generators. For this, we use Masur and Minsky’s *Bounded Geodesic Image Theorem* [33] to prove that the element acts with positive translation distance on the curve complex of the appropriate subsurface.

We end with a discussion of surface subgroups and the proofs of Corollary 1.3 and Proposition 7.1.

**Acknowledgements.** We would like to thank Richard Kent, Alan Reid, Thomas Koberda and Jason Behrstock for helpful conversations, and Sergio Fenley for carefully reading and commenting on an earlier version of this paper. We would also like to thank the *Hausdorff Research Institute for Mathematics* in Bonn, Germany, for its hospitality while this work was

being completed.

## 2 Notation and terminology

### 2.1 Quasi-isometries

Given  $A \geq 1$  and  $B \geq 0$ , we write  $x \stackrel{A,B}{\asymp} y$  to mean

$$\frac{y - B}{A} \leq x \leq Ay + B$$

If  $(\mathfrak{X}_1, d_1)$  and  $(\mathfrak{X}_2, d_2)$  are metric spaces and  $A \geq 1, B \geq 0$ , then an  $(A, B)$ -quasi-isometric embedding from  $\mathfrak{X}_1$  to  $\mathfrak{X}_2$  is a map

$$F : \mathfrak{X}_1 \rightarrow \mathfrak{X}_2$$

with the property that for all  $x, y \in \mathfrak{X}_1$ , we have

$$d_1(x, y) \stackrel{A,B}{\asymp} d_2(F(x), F(y)).$$

If  $F$  is an  $(A, B)$ -quasi-isometric embedding for some  $A$  and  $B$ , then we will say that  $F$  is a *quasi-isometric embedding*.

If  $F : \mathfrak{X}_1 \rightarrow \mathfrak{X}_2$  is a quasi-isometric embedding and there is a constant  $D > 0$  so that any point of  $\mathfrak{X}_2$  is within  $D$  of some point of  $F(\mathfrak{X}_1)$ , then  $F$  is called a *quasi-isometry*.

### 2.2 Right angled Artin groups

Let  $\Gamma$  be a graph with vertex set  $\{s_1, \dots, s_n\}$ . The associated *right-angled Artin group*  $G = G(\Gamma)$ , is defined to be the group with presentation

$$G = \langle s_1, \dots, s_n \mid [s_i, s_j] = 1 \text{ if } \{s_i, s_j\} \text{ is an edge of } \Gamma \rangle.$$

We will always work with the word metric on  $G$  with respect to this generating set, and will denote it  $d_G$ .

Examples of right-angled Artin groups are free groups, and direct products of free groups (in particular, free abelian groups). A simple example of a right-angled Artin group which is neither free nor a product of free groups is  $G(\Gamma)$  where  $\Gamma$  is the cyclic graph with 5 vertices shown in Figure 1.

### 2.3 Surfaces

Given a connected surface  $S$  of genus  $g$  with  $n$  punctures, the *complexity* is defined to be  $\xi(S) = 3g - 3 + n$ . Unless otherwise stated, we will assume throughout that  $\xi(S) > 0$ . The *mapping class group* of  $S$  is the group of isotopy classes of orientation preserving homeomorphisms of  $S$  and is denoted  $\text{Mod}(S)$ . By a *curve* in  $S$ , we mean the isotopy class of an essential (non-null-homotopic and non-peripheral) simple closed curve. A *pants decomposition* of  $S$  is a maximal collection of pairwise disjoint curves in  $S$ . Since  $\xi(S) > 0$ , a nonempty pants decomposition exists and has precisely  $\xi(S)$  curves in it.

A subsurface  $X \subset S$  is *essential* if it is either a regular neighborhood of an essential simple closed curve, or else a component of the complement of an open regular neighborhood of a




Figure 1: The cyclic graph with 5 vertices.

(possibly empty) union of pairwise disjoint essential simple closed curves. In particular, we assume that essential subsurfaces are connected. We will generally not distinguish between punctures and boundary components, and if  $X \subset S$  has genus  $h$  with  $k$  punctures and  $b$  boundary components, then we will write  $\xi(X) = 3g - 3 + k + b$ . Finally, we will assume that an essential subsurface  $X$  has  $\xi(X) \neq 0$ , thus excluding a pair of pants as an essential subsurface. The set of all isotopy classes of essential subsurfaces  $X$  of  $S$  with  $\xi(X) \neq 0$  will be denoted  $\Omega(S)$ .




Figure 2: A genus 2 surface with 1 puncture  $S$ , a subsurface  $X$  (shaded) and a curve  $\gamma$ .

We will often refer to the isotopy class of an essential subsurface simply as a *subsurface*. Furthermore, we will choose nice representative for each curve and each subsurface, and will not distinguish between a representative and its isotopy class when it is convenient. To be precise, we choose representatives as follows (annuli will play essentially no role in our discussion, so we do not bother describing their preferred representatives).

Fix a complete hyperbolic metric on  $S$ , and realize each curve by its unique geodesic representative. These representatives minimize the number of intersections (that is, they realize geometric intersection number). For each curve  $\alpha$ , we may choose some  $\epsilon_\alpha$ -neighborhood  $N(\alpha)$  so that for any curves  $\alpha$  and  $\beta$ , the intersections of  $N(\alpha)$  and  $N(\beta)$  correspond precisely to the intersections of  $\alpha$  and  $\beta$ , and each such intersection is a “product square” (see Figure 4). For any nonannular subsurface  $X$ , which is a component of the complement of an open regular neighborhood of  $\alpha_1 \cup \dots \cup \alpha_k$ , we take its representative to be defined as the corresponding component of the complement of the interior of the neighborhood  $N(\alpha_1) \cup \dots \cup N(\alpha_k)$ .

Suppose  $X, Y \subsetneq S$  are representative subsurfaces. Observe that  $X \cap Y = \emptyset$  if and only if  $X$  and  $Y$  cannot be isotoped to be disjoint. In this case, we say that  $X$  and  $Y$  are *overlapping*, and write  $X \pitchfork Y$  if  $X \not\subseteq Y$  and  $Y \not\subseteq X$ . One can check that this notion of overlapping agrees with that defined in [3], which is to say that  $X \pitchfork Y$  if and only if some component of  $\partial X$  cannot be isotoped disjoint from  $Y$  and some component of  $\partial Y$  cannot be isotoped disjoint from  $X$ .

## 2.4 Realizing a graph

Given a graph  $\Gamma$ , a surface  $S$ , and a collection of nonannular subsurfaces  $X_1, \dots, X_n \subset S$ , we say that  $\mathbb{X} = \{X_1, \dots, X_n\}$  *realizes*  $\Gamma$  *nicely* in  $S$  if

- (1)  $X_i \cap X_j = \emptyset$  if and only if  $\{s_i, s_j\}$  is an edge of  $\Gamma$ , and
- (2) whenever  $X_i \cap X_j \neq \emptyset$ , then  $X_i \pitchfork X_j$ .

As Figure 3 indicates, there is a nice realization of the cyclic graph of length 5 in a genus 3 surface obtained from a branched cover of the sphere, branched over 8 points. By adding more points to this picture and taking a branched cover, we can produce nice realizations of this graph in any surface of genus  $g \geq 3$ . Moreover, given any graph it is easy to find some surface and a collection of subsurfaces which provide a nice realization (see [9, 11] for this kind of construction). We sketch one such construction here.




Figure 3: This figure represents a sphere with 8 punctures containing five curves, each of which bounds a disk with 3 punctures. These five 3-punctured disks provide a nonannular realization of the cyclic graph with 5 vertices. Taking a two-fold branched cover over the 8 points, we obtain a nonannular realization on a genus 3 surface (by 1-holed tori).

Starting with a graph  $\Gamma$ , we take a disjoint union of annuli, one for each vertex of  $\Gamma$ . Next, glue together the annuli along product squares whenever the associated vertices of  $\Gamma$  are not connected by an edge. In each annulus, remove a disk and glue in a 1-holed torus. Finally, cap off the boundary components of the resulting surface with disks. See Figure 4 for a particular example.

If  $X$  is a nonannular subsurface of  $S$  and  $f \in \text{Mod}(S)$  is the identity outside  $X$ , we say that  $f$  is *supported on*  $X$ . We say that  $f$  is *fully supported on*  $X$  if we also have that  $f$  is




Figure 4: A graph  $\Gamma$  and the associated annuli glued together along product squares as prescribed by  $\Gamma$ . At the bottom, we glue in a 1-holed torus to an annulus with a disk removed.

pseudo-Anosov on  $X$ . If  $f$  is supported on  $X$ , then  $f$  acts on  $\mathcal{C}(X)$ , the *curve complex of  $X$* , and we let  $\tau_X(f)$  denote the *translation length of  $f$  on  $\mathcal{C}(X)$* . This is defined by

$$\tau_X(f) = \lim_{k \rightarrow \infty} \frac{d_X(\alpha, f^k(\alpha))}{k},$$

where  $\alpha$  is any curve in  $X$ , and  $d_X$  is the distance in  $\mathcal{C}(X)$ —see Section 3.3. The following result of Masur and Minsky [34][Proposition 4.6] says that if  $f$  is supported on  $X$ , then it is fully supported on  $X$  if and only if  $\tau_X(f) > 0$ .

**Theorem 2.1** (Masur-Minsky). *Given  $X$  there exists  $c > 0$  so that if  $f \in \text{Mod}(X)$  is pseudo-Anosov, then for all  $\alpha \in \mathcal{C}(X)$*

$$d_X(\alpha, f^k(\alpha)) \geq ck.$$

## 2.5 Homomorphisms

Suppose now that  $\mathbb{X} = \{X_1, \dots, X_n\}$  nicely realizes  $\Gamma$  in  $S$  and that  $\mathbb{F} = \{f_1, \dots, f_n\} \subset \text{Mod}(S)$  are mapping classes. We say that  $\mathbb{F}$  is (fully) supported on  $\mathbb{X}$  if  $f_i$  is (fully) supported on  $X_i$  for each  $i = 1, \dots, n$ . Since homeomorphisms on disjoint subsurfaces commute, there is a unique homomorphism

$$\phi_{\mathbb{F}} : G \rightarrow \text{Mod}(S)$$

defined by  $\phi_{\mathbb{F}}(s_i) = f_i$ .

We now state a more precise version of our main theorem. We write  $\mathcal{T}(S)$  for the Teichmüller space, and we denote its two standard metrics by  $d_{\mathcal{T}}$  for the Teichmüller metric and  $d_{\text{WP}}$  for the Weil–Petersson metric.

**Theorem 2.2.** *Given a graph  $\Gamma$  and a nice realization  $\mathbb{X} = \{X_1, \dots, X_n\}$  of  $\Gamma$  in  $S$ , there exists a constant  $C > 0$  with the following property. If  $\mathbb{F} = \{f_1, \dots, f_n\}$  is fully supported on  $\mathbb{X}$  and  $\tau_{X_i}(f_i) \geq C$  for all  $i = 1, \dots, n$ , then the associated homomorphism*

$$\phi_{\mathbb{F}} : G(\Gamma) \rightarrow \text{Mod}(S)$$

*is a quasi-isometric embedding. Furthermore, the orbit map  $G \rightarrow \mathcal{J}(S)$  is a quasi-isometric embedding for both  $d_{\mathcal{J}}$  and  $d_{\text{VP}}$ .*

**Remark.** We reiterate for the casual reader that the subsurfaces  $X_i$  are assumed to be essential, connected, and nonannular.

The proof of Theorem 2.2 will be carried out in Section 5. Theorem 2.2 easily implies Theorem 1.1.

### 3 Projections and distance estimates

Our proof of Theorem 2.2 uses results from [33], [6], [38] and [3]. The main construction we will use is that of *subsurface projection*, which we now briefly recall.

#### 3.1 Projections




Figure 5: The projection  $\pi_X(\gamma)$ , where  $S$ ,  $X$  and  $\gamma$  are as in Figure 2.

Given a nonannular subsurface  $X$  of  $S$  and a curve  $\gamma$ , we define the *projection of  $\gamma$  to  $X$* , denoted  $\pi_X(\gamma)$ , to be the subset of  $\mathcal{C}(X)$  constructed as follows. If  $\gamma \cap X \neq \emptyset$ , then either  $\gamma$  is an essential simple closed curve in  $X$ , and we define  $\pi_X(\gamma) = \{\gamma\}$ , or else  $\gamma \cap X$  is a disjoint union of essential arcs in  $X$ . For each arc, consider  $N$ , the regular neighborhood of the arc union the boundary components of  $X$  which the arc meets. Then the boundary of  $N$  is a union of curves in  $X$  (and components of  $\partial X$ ), and we define  $\pi_X(\gamma)$  to be the set of all such curves in  $X$ , over all arcs of  $\gamma \cap X$ . See Figure 5. In general, the curves in  $\pi_X(\gamma)$  need not be disjoint, but the set has diameter at most 2; see [33].

When  $X$  is an annulus and  $\gamma$  a curve, there is also a notion of a projection to  $X$ , which assigns to  $\gamma$  a diameter one subset of the arc complex of  $X$ , denoted  $\mathcal{C}(X)$ , and again we denote this by  $\pi_X(\gamma)$ . For our purposes, simply the existence of this projection will suffice, so for the details of its definition, we refer the reader to [33].



If  $\gamma$  is a disjoint union of curves  $\gamma_1 \cup \dots \cup \gamma_k$ , then we define  $\pi_X(\gamma)$  to be the union  $\bigcup_i \pi_X(\gamma_i)$ . This set also has diameter at most 2. If  $\gamma \cap X = \emptyset$ , then  $\pi_X(\gamma) = \emptyset$ .

### 3.2 Markings

Another object we will need is a *marking*. For us, this will mean a *complete clean marking* in the sense of Masur and Minsky [33]. More precisely, a marking  $\mu$  is a pants decomposition called the *base of  $\mu$*

$$\text{base}(\mu) = \{\alpha_1, \dots, \alpha_{\xi(S)}\},$$

together with a *transversal* for each curve  $\alpha_i \in \text{base}(\mu)$ : this is a diameter at most one subset of  $\mathcal{C}(X_i)$ , where  $X_i$  is the annular neighborhood of  $\alpha_i$ , together with some additional properties which we will not need descriptions for; see [33] for a discussion.

Masur and Minsky [33] identify the set of all markings with the vertex set of a graph  $\tilde{\mathcal{M}}(S)$  called the *marking graph of  $S$* . The edges of this graph correspond to certain *elementary moves* one can perform on a marking. We denote the resulting path metric on  $\tilde{\mathcal{M}}(S)$  by  $d_{\tilde{\mathcal{M}}}$ . The graph  $\tilde{\mathcal{M}}(S)$  is locally finite, and  $\text{Mod}(S)$  acts by isometries on it. In particular, the orbit map of this action is a quasi-isometry. We will use  $\tilde{\mathcal{M}}(S)$  as a model for  $\text{Mod}(S)$ .

Any marking  $\mu$  can be projected to a subsurface. If  $X$  is a nonannular subsurface, then  $\pi_X(\mu)$  is defined to be  $\pi_X(\text{base}(\mu))$ . For annuli, the projection is defined differently; see [33].

### 3.3 Distances

Given a subsurface  $X$  and curves or markings  $\mu$  and  $\mu'$ , we define their *distance in  $X$*  to be

$$d_X(\mu, \mu') = \text{diam}(\pi_X(\mu) \cup \pi_X(\mu'))$$

where the diameter is computed in  $\mathcal{C}(X)$ .

A trivial observation is that if  $\mu, \mu'$  are curves or markings on  $S$ ,  $f \in \text{Mod}(S)$  is supported on  $X$ , and  $Y$  is a nonannular subsurface disjoint from  $X$  such that  $\mu$  and  $\mu'$  have nonempty projection to  $Y$  then

$$d_Y(\mu, f(\mu')) = d_Y(\mu, \mu').$$

**Remark.** We note that the validity of this observation relies on the assumption that  $Y$  is nonannular.

Given  $K > 0$  and  $\mu, \mu' \in \tilde{\mathcal{M}}(S)$ , define

$$\Omega(K, \mu, \mu') = \{X \subseteq S \mid \xi(X) \geq 1 \text{ or } X \text{ is an annulus, and } d_X(\mu, \mu') \geq K\}.$$

It is convenient to decompose  $\Omega(K, \mu, \mu')$  into the annular subsurfaces  $\Omega_a(K, \mu, \mu')$  and the nonannular subsurfaces  $\Omega_n(K, \mu, \mu')$ .

The following theorem is proven in [33].

**Theorem 3.1** (Masur-Minsky). *There exists  $K_0 > 0$  (depending on  $S$ ) so that if  $K \geq K_0$ , then there exists  $A \geq 1$ ,  $B \geq 0$  with the following property. Given  $\mu, \mu' \in \tilde{\mathcal{M}}(S)$  then*

$$d_{\tilde{\mathcal{M}}}(\mu, \mu') \stackrel{A, B}{\asymp} \sum_{X \in \Omega(K, \mu, \mu')} d_X(\mu, \mu')$$

A theorem of Brock [6] states that the Weil–Petersson metric on Teichmüller space is quasi-isometric to the *pants graph*. In [33], Masur and Minsky give a formula similar to that of the previous formula for distance in the pants graph. In particular combining these two results one obtains the following.

**Theorem 3.2** (Brock, Masur-Minsky). *There exists  $K_0 > 0$  (depending on  $S$ ) so that if  $K \geq K_0$ , then there exists  $A \geq 1$ ,  $B \geq 0$  with the following property. If  $\mu, \mu' \in \tilde{\mathcal{M}}(S)$  are shortest markings for  $m, m' \in \mathcal{T}(S)$ , respectively, then*

$$d_{\text{WP}}(m, m') \stackrel{A, B}{\asymp} \sum_{X \in \Omega_n(K, \mu, \mu')} d_X(\mu, \mu')$$

A *shortest marking* for  $m$  is just a marking for which the pants decomposition has the shortest total length among all pants decompositions, and the transversals are projections of the shortest curves among those which can be used for transversals. For this theorem, the transversals are unimportant.

The analogous result for the Teichmüller metric was proven by Rafi in [38].

**Theorem 3.3** (Rafi). *There exists  $K_0 > 0$  (depending on  $S$ ) so that if  $K \geq K_0$  and  $\epsilon > 0$  then there exists  $A \geq 1$ ,  $B \geq 0$  with the following property. If  $\mu, \mu' \in \tilde{\mathcal{M}}(S)$  are shortest markings for  $m, m'$  in the  $\epsilon$ -thick part of  $\mathcal{T}(S)$ , respectively, then*

$$d_{\mathcal{T}}(m, m') \stackrel{A, B}{\asymp} \sum_{X \in \Omega_n(K, \mu, \mu')} d_X(\mu, \mu') + \sum_{X \in \Omega_a(K, \mu, \mu')} \log(d_X(\mu, \mu'))$$

**Remarks.** (1) The special case of Theorem 2.2 in which  $G(\Gamma)$  is abelian now follows immediately from the preceding three theorems and Theorem 2.1. Our proof is an extension of this idea.

(2) Strictly speaking, Theorems 3.1 and 3.2 would be sufficient for our purposes since, up to a constant,  $d_{\text{WP}}$  provides a lower bound for  $d_{\mathcal{T}}$  by a result of Linch [28], and the lower bound on distortion is the only nontrivial inequality we need to prove. However, it seems worthwhile to include Theorem 3.3 as this illustrates a common interpretation for all of the metric spaces  $\tilde{\mathcal{M}}(S)$  (or  $\text{Mod}(S)$ ),  $(\mathcal{T}(S), d_{\mathcal{T}})$ , and  $(\mathcal{T}(S), d_{\text{WP}})$ .

One final result about distances and subsurface projections which we will need is the following Bounded Geodesic Image Theorem [33].

**Theorem 3.4** (Masur-Minsky). *There exists  $K_0 > 0$  (depending on  $S$ ) so that if  $\{v_1, \dots, v_n\}$  is a geodesic in  $\mathcal{C}(S)$  and  $X \in \Omega(S)$ , then either  $\pi_X(v_j) = \emptyset$  for some  $j$  or else*

$$\text{diam}_X(\{\pi_X(v_1), \dots, \pi_X(v_n)\}) < K_0.$$

In particular, note that if  $v, v' \in \mathcal{C}(S)$  are two curves with  $d_X(v, v') \geq K_0$ , then any geodesic between  $v$  and  $v'$  in  $\mathcal{C}(S)$  must pass through a curve  $v''$  disjoint from  $X$  (for example, it may pass through a curve in  $\partial X$ ).

For simplicity, we will assume, as we may, that  $K_0$  is the same constant in all of the theorems in this section.

### 3.4 Partial order on subsurfaces

In [3], Behrstock, Kleiner, Minsky and Mosher defined a partial order on  $\Omega(K, \mu, \mu')$  (for  $K$  sufficiently large) which is closely related to the time-order constructed in [33] (see also [4]). However, as is noted in [3], while the time-order in [33] (which is defined on geodesics in hierarchies) requires a fair amount of the hierarchy machinery to describe it, the partial order on  $\Omega(K, \mu, \mu')$  is completely elementary. As this is the basic tool we will use, we include the construction and verification of the necessary properties of this partial order, for the sake of completeness.

The starting point is the ‘‘Behrstock inequality’’ [2] (see also [32], Lemma 2.5, for the version stated here).

**Proposition 3.5** (Behrstock). *Suppose  $X$  and  $Y$  are overlapping subsurfaces of  $S$  and  $\mu$  is a marking on  $S$ . Then*

$$d_X(\partial Y, \mu) \geq 10 \Rightarrow d_Y(\partial X, \mu) \leq 4.$$

Suppose  $K \geq 20$  and we define the partial order as follows. Given  $X, Y \in \Omega(K, \mu, \mu')$  with  $X \pitchfork Y$ , then we write  $X \prec Y$  if

$$d_X(\mu, \partial Y) \geq 10. \tag{1}$$

That this is a strict partial order is a consequence of the following useful description of  $\prec$ .

**Proposition 3.6** (Behrstock-Kleiner-Minsky-Mosher). *Suppose  $K \geq 20$  and  $X, Y \in \Omega(K, \mu, \mu')$  with  $X \pitchfork Y$ . Then  $X$  and  $Y$  are ordered and the following are equivalent*

- |                                       |  |
|---------------------------------------|--|
| (1) $X \prec Y$                       | (5) $d_Y(\mu', \partial X) \geq 10$    |
| (2) $d_X(\mu, \partial Y) \geq 10$    | (6) $d_Y(\mu', \partial X) \geq K - 4$ |
| (3) $d_X(\mu, \partial Y) \geq K - 4$ | (7) $d_Y(\mu, \partial X) \leq 4$      |
| (4) $d_X(\mu', \partial Y) \leq 4$    |  |

*Proof.* Assume the hypothesis of the proposition. Since  $X \pitchfork Y$ , we know that  $\pi_X(\partial Y) \neq \emptyset$  and  $\pi_Y(\partial X) \neq \emptyset$ . To verify the equivalences, first observe that (1) and (2) are equivalent by definition, and since  $K - 4 > 10$ , (3) implies (2) and (6) implies (5). Next, since  $d_X(\mu, \mu'), d_Y(\mu, \mu') \geq K$ , the triangle inequality guarantees that (4) implies (3) and (7) implies (6). Furthermore, since  $K - 4 > 10$ , Proposition 3.5 tells us that (2) implies (7) and (5) implies (4). This proves all the required implications.

Finally, we prove that  $X$  and  $Y$  are ordered. By the triangle inequality we have

$$20 \leq K \leq d_X(\mu, \mu') \leq d_X(\mu, \partial Y) + d_X(\mu', \partial Y).$$

and so one of  $d_X(\mu, \partial Y)$  or  $d_X(\mu', \partial Y)$  is at least 10. If  $d_X(\mu, \partial Y) \geq 10$  then  $X \prec Y$ . If  $d_X(\mu', \partial Y) \geq 10$ , then reversing the roles of  $X$  and  $Y$  in each of the 7 equivalent statements we see that  $Y \prec X$ , as required.  $\square$

**Corollary 3.7** (Behrstock-Kleiner-Minsky-Mosher). *Suppose  $K \geq 20$ . Then the relation  $\prec$  is a strict partial order.*

*Proof.* Since we never have  $X \pitchfork X$ , it follows that  $\prec$  is non-reflexive. Furthermore, the equivalence of (2) and (7) in Proposition 3.6 means that  $X \prec Y$  implies  $Y \not\prec X$ , so  $\prec$  is antisymmetric. Finally, if  $X \prec Y$  and  $Y \prec Z$  then we know  $\pi_Y(\partial X)$  and  $\pi_Y(\partial Z)$  are nonempty, and appealing to Proposition 3.6 and the triangle inequality we have

$$20 \leq K \leq d_Y(\mu, \mu') \leq d_Y(\mu, \partial X) + d_Y(\partial X, \partial Z) + d_Y(\mu', \partial Z) \leq d_Y(\partial X, \partial Z) + 8$$

and so

$$d_Y(\partial X, \partial Z) \geq 12 > 10.$$

In this case,  $\partial X$  and  $\partial Z$  intersect nontrivially in  $Y$ , so in particular,  $X \pitchfork Z$ .

Now we apply Proposition 3.5 to the preceding inequality to obtain

$$d_X(\partial Y, \partial Z) \leq 4$$

and hence by the triangle inequality

$$16 \leq K - 4 \leq d_X(\partial Y, \mu) \leq d_X(\partial Y, \partial Z) + d_X(\partial Z, \mu) \leq 4 + d_X(\partial Z, \mu).$$

Therefore,  $d_X(\partial Z, \mu) \geq 12 > 10$ , and  $X \prec Z$ . □

## 4 Normal forms in right-angled Artin groups

Here we describe the normal forms in  $G = G(\Gamma)$  as defined by Green [17], and Hermiller and Meier's procedure for obtaining these normal forms [19]. We refer the reader to Charney's survey article [7] for a discussion.

Suppose  $w = x_1^{e_1} \cdots x_k^{e_k}$  is a word in the generators:  $x_i \in \{s_1, \dots, s_n\}$  and  $e_i \in \mathbb{Z}$ . Each  $x_i^{e_i}$  is called a *syllable* of  $w$ . We consider the following moves which can be applied to  $w$  (see also [20]):

1. Remove a syllable  $x_i^{e_i}$  if  $e_i = 0$ .
2. If  $x_i = x_{i+1}$ , then replace consecutive syllables  $x_i^{e_i} x_{i+1}^{e_{i+1}}$  by  $x_i^{e_i + e_{i+1}}$ .
3. If  $[x_i, x_{i+1}] = 1$ , then replace  $x_i^{e_i} x_{i+1}^{e_{i+1}}$  with  $x_{i+1}^{e_{i+1}} x_i^{e_i}$ .

Let  $\text{Min}(\sigma)$  be the set of words representing  $\sigma \in G$  with the fewest number of syllables. Green's normal form for  $\sigma$  is a certain type of element of  $\text{Min}(\sigma)$  obtained by stringing together, from left to right, maximal collections of commuting syllables. For us, we will consider any element of  $\text{Min}(\sigma)$  as a normal form, and we will shortly impose some additional structure on the set of syllables. First, we state the following from [19].

**Theorem 4.1** (Hermiller-Meier). *Any word representing  $\sigma \in G$  can be transformed to any element of  $\text{Min}(\sigma)$  by applying a sequence of the moves above. In particular, in any such sequence, the number of syllables and the length does not increase.*

It follows that the words in  $\text{Min}(\sigma)$  determine geodesics in (the Cayley graph of)  $G$  with respect to  $s_1, \dots, s_n$ . Moreover, note that any two elements of  $\text{Min}(\sigma)$  differ by moves of type (3).

Let  $w = x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$  and consider the set of syllables  $\text{syl}(w) = \{x_i^{e_i}\}_{i=1}^k$ . We consider this as a set of  $k$  distinct elements: for example, we can artificially write this as

$\{(x_i^{e_i}, i)\}_{i=1}^k$ . If we have two elements  $w, w' \in \text{Min}(\sigma)$  that differ by a single application of move (3) above, then there is an obvious bijection between  $\text{syl}(w)$  and  $\text{syl}(w')$ . Moreover, any sequence of these types of moves results in a sequence of bijections between the syllables of consecutive words in  $\text{Min}(\sigma)$ . Observe that any such bijection between  $\text{syl}(w)$  and  $\text{syl}(w')$  sends a syllable of  $w$  to one of  $w'$  representing the same element of  $G$ .

From this it follows that if any such sequence of moves ever brings a word  $w$  back to itself, then the bijection from  $\text{syl}(w)$  to itself is the identity. The reason is that if  $x_i^{e_i}$  and  $x_j^{e_j}$  are syllables of  $w$  which represent the same element in  $G$  (so  $x_i = x_j$ ,  $e_i = e_j$ ), then if  $x_i^{e_i}$  precedes  $x_j^{e_j}$  in  $w$ , any of the bijections will preserve this property: a sequence of type (3) moves which would theoretically accomplish a swap of their positions making  $x_j^{e_j}$  precede  $x_i^{e_i}$  would require a move where  $x_i^{e_i}$  and  $x_j^{e_j}$  are adjacent, at which time a type (2) move could be applied to reduce the number of syllables, and this is impossible. We use these bijections to identify the syllables of any two words  $w, w' \in \text{Min}(\sigma)$ , and simply write  $\text{syl}(\sigma)$  for this set of syllables.

We can define a strict partial order on this set of syllables, denoted  $\text{syl}(\sigma)$ , by declaring  $x_i^{e_i} \prec x_j^{e_j}$  if and only if  $x_i^{e_i}$  precedes  $x_j^{e_j}$  in *every* word  $w \in \text{Min}(\sigma)$ . So for any  $w \in \text{Min}(\sigma)$ , the order of the syllables is a refinement of the partial order (and the partial order is the largest partial order having this property for every  $w \in \text{Min}(\sigma)$ ).

## 5 The proof of Theorem 2.2.

Throughout this section, we will assume  $\mathbb{X} = \{X_1, \dots, X_n\}$  realizes  $\Gamma$  nicely in  $S$ ,  $\mathbb{F} = \{f_1, \dots, f_n\}$  is fully supported on  $\mathbb{X}$ , and  $\phi_{\mathbb{F}} : G = G(\Gamma) \rightarrow \text{Mod}(S)$  is the associated homomorphism.

Given a word  $x_1^{e_1} \cdots x_k^{e_k}$  with  $x_i \in \{s_1, \dots, s_n\}$  for all  $i$ , let  $J(i) \in \{1, \dots, n\}$  be the unique number for which  $x_i = s_{J(i)}$ . For any  $\sigma \in G$  and  $w = x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$ , set

$$X^w(x_i^{e_i}) = \phi_{\mathbb{F}}(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}})(X_{J(i)})$$

for  $i = 2, \dots, k$  and define  $X^w(x_1^{e_1}) = X_{J(1)}$ . We think of this as defining a map

$$X^w : \text{syl}(\sigma) \rightarrow \Omega(S).$$

**Lemma 5.1.** *Suppose  $\Gamma$ ,  $\mathbb{X}$  and  $\mathbb{F}$  are as above. If  $\sigma \in G(\Gamma)$  and  $w, w' \in \text{Min}(\sigma)$ , then  $X^w = X^{w'} : \text{syl}(\sigma) \rightarrow \Omega(S)$ .*

*Proof.* Since any two words  $w, w' \in \text{Min}(\sigma)$  differ by a sequence of moves of type (3), that is, in which adjacent commuting syllables are exchanged, it suffices to verify the lemma in the case that  $w$  and  $w'$  differ by such a move:

$$w = x_1^{e_1} \cdots x_i^{e_i} x_{i+1}^{e_{i+1}} \cdots x_n^{e_n} \text{ and } w' = x_1^{e_1} \cdots x_{i+1}^{e_{i+1}} x_i^{e_i} \cdots x_n^{e_n}.$$

For  $j \neq i$  or  $i+1$ , we clearly have  $X^w(x_j^{e_j}) = X^{w'}(x_j^{e_j})$ , and so we must show

$$X^w(x_i^{e_i}) = X^{w'}(x_i^{e_i}) \quad \text{and} \quad X^w(x_{i+1}^{e_{i+1}}) = X^{w'}(x_{i+1}^{e_{i+1}}).$$

Interchanging the roles of  $w$  and  $w'$ , it suffices to prove just one of these equations, say  $X^w(x_i^{e_i}) = X^{w'}(x_i^{e_i})$ .

We have

$$X^w(x_i^{e_i}) = \phi_{\mathbb{F}}(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}})(X_{J(i)})$$

whereas

$$X^{w'}(x_i^{e_i}) = \phi_{\mathbb{F}}(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}} x_{i+1}^{e_{i+1}})(X_{J(i)}) = \phi_{\mathbb{F}}(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}}) \phi_{\mathbb{F}}(x_{i+1}^{e_{i+1}})(X_{J(i)}).$$

Since  $x_i^{e_i}$  and  $x_{i+1}^{e_{i+1}}$  commute,  $X_{J(i+1)}$ , the support of  $\phi_{\mathbb{F}}(x_{i+1}^{e_{i+1}}) = f_{J(i+1)}^{e_{i+1}}$  is disjoint from  $X_{J(i)}$ . Therefore,

$$\phi_{\mathbb{F}}(x_{i+1}^{e_{i+1}})(X_{J(i)}) = X_{J(i)}$$

and the lemma follows.  $\square$

By this lemma we can unambiguously define  $X^\sigma = X^w$ , independent of the choice of  $w \in \text{Min}(\sigma)$ .

The main technical theorem we prove is the following. From this, together with Theorems 3.1, 3.2 and 3.3, our Theorem 2.2 (and hence also Theorem 1.1) follows easily.

**Theorem 5.2.** *Suppose  $\Gamma$  and  $\mathbb{X}$  are as above and  $\mu \in \tilde{\mathcal{M}}(S)$ . Then there exists a constant  $K \geq K_0$  with the following property.*

*Suppose that  $\mathbb{F} = \{f_1, \dots, f_n\}$  is fully supported on  $\mathbb{X}$  and that  $\tau_{X_j}(f_j) \geq 2K$  for all  $1 \leq j \leq n$ , and let  $\phi_{\mathbb{F}} : G \rightarrow \text{Mod}(S)$  be the associated homomorphism. Then, for any  $\sigma \in G$  with  $x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$  we have*

1.  $d_{X^\sigma(x_i^{e_i})}(\mu, \phi(\sigma)\mu) \geq K|e_i|$  for each  $i = 1, \dots, k$ . Consequently,

$$X^\sigma(\text{syl}(\sigma)) \subset \Omega(K, \mu, \phi(\sigma)\mu).$$

2.  $X^\sigma(\cdot) : \text{syl}(\sigma) \rightarrow \Omega(K, \mu, \phi(\sigma)\mu)$  is an order-preserving injection.

*Proof.* Let

$$K = K_0 + 20 + 2 \cdot \max\{d_{X_j}(\partial X_i, \mu) : i \neq j\}.$$

Throughout the proof, we let  $\phi = \phi_{\mathbb{F}}$ .

In what follows, we prove statements 1 and 2 separately. For both, the proof is by induction on the number of syllables in  $w \in \text{Min}(\sigma)$ .

*Proof of Statement 1.* To make the ideas in the proof more transparent, we introduce simplified notation. Given  $w = x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$ , define

$$g_i = \phi(x_i^{e_i}) = f_{J(i)}^{e_i} \quad \text{and} \quad Y_i = X_{J(i)}.$$

Then  $X^\sigma(x_1^{e_1}) = Y_1$ ,  $X^\sigma(x_2^{e_2}) = g_1 Y_2$ , and in general  $X^\sigma(x_i^{e_i}) = g_1 g_2 \cdots g_{i-1} Y_i$ . In this notation, statement 1 claims that

$$d_{g_1 \cdots g_{i-1} Y_i}(\mu, g_1 \cdots g_k \mu) \geq K|e_i|,$$

for  $i = 2, \dots, k$ , and also  $d_{Y_1}(\mu, g_1 \cdots g_k \mu) \geq K|e_1|$ .

Suppose  $w$  has only one syllable. Then the claim only states that  $d_{Y_1}(\mu, g_1 \mu) \geq K|e_1|$ , which holds because, letting  $j = J(1)$ , we know

$$\begin{aligned} d_{Y_1}(\mu, g_1 \mu) &= d_{X_j}(\mu, f_j^{e_1}(\mu)) \\ &= \text{diam}_{X_j}(\pi_{X_j}(\mu) \cup \pi_{X_j}((f_j^{e_1})\mu)) \\ &= \text{diam}_{X_j}(\pi_{X_j}(\mu) \cup f_j^{e_1}(\pi_{X_j}(\mu))) \\ &\geq \tau_{X_j}(f_j^{e_1}) \\ &\geq 2K|e_1|. \end{aligned}$$

Now suppose we have proved the claim for elements of  $G = G(\Gamma)$  whose minimal representatives have at most  $k - 1$  syllables. Let  $\sigma \in G$  with  $w = x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$  having  $k$  syllables. Define  $g_i$  and  $Y_i$  as above. Our next step is to separate the product  $g_1 \cdots g_k$  into subproducts, as illustrated below, with the additional possibility that  $a$ ,  $b$ , or  $c$  might be the empty word:

$$\overbrace{g_1 \cdots g_\ell}^a \overbrace{g_{\ell+1} \cdots g_{i-1}}^b \overbrace{g_i g_{i+1} \cdots g_k}^c,$$

The subproducts  $a$ ,  $b$ , and  $c$  are defined as follows. By Lemma 5.1, we may assume that either  $g_i$  and  $g_{i+1}$  fail to commute, or by replacing  $w$  with another word in  $\text{Min}(\sigma)$ , that  $i = k$ . In the first case, let  $c = g_{i+1} \cdots g_k$ ; in the latter case, let  $c$  be the identity. If there exists some syllable to the left of  $g_i$  which does not commute with  $g_i$ , let  $\ell$  be the largest index such that  $g_i$  and  $g_\ell$  do not commute, and let  $a = g_1 \cdots g_\ell$ . Otherwise let  $a$  be the identity. Let  $b = g_{\ell+1} \cdots g_{i-1}$  if  $\ell + 1 < i$ , and otherwise let  $b$  be the identity; observe that by construction,  $b$  commutes with  $g_i$ .

Because  $g_1 \cdots g_k = abg_i c$ , we have

$$d_{g_1 \cdots g_{i-1} Y_i}(\mu, g_1 \cdots g_k \mu) = d_{abY_i}(\mu, abg_i c \mu) = d_{Y_i}(b^{-1} a^{-1} \mu, g_i c \mu).$$

By the triangle inequality and the fact that  $d_{Y_i}(g_i c \mu, c \mu) \geq 2K|e_i|$ ,

$$d_{Y_i}(b^{-1} a^{-1} \mu, g_i c \mu) \geq 2K|e_i| - d_{Y_i}(b^{-1} a^{-1} \mu, c \mu).$$

To control the last term we again employ the triangle inequality:

$$d_{Y_i}(b^{-1} a^{-1} \mu, c \mu) \leq d_{Y_i}(b^{-1} a^{-1} \mu, \mu) + d_{Y_i}(\mu, c \mu).$$

Because  $b$  is the (possibly empty) product of syllables  $g_j$  that commute with  $g_i$ ,  $b$  acts as the identity on  $Y_i$ . Therefore we have

$$\begin{aligned} d_{Y_i}(b^{-1} a^{-1} \mu, \mu) &= d_{Y_i}(a^{-1} \mu, b \mu) \\ &= \text{diam}_{Y_i}(\pi_{Y_i}(a^{-1} \mu) \cup \pi_{Y_i}(b \mu)) \\ &= \text{diam}_{Y_i}(\pi_{Y_i}(a^{-1} \mu) \cup \pi_{Y_i}(\mu)) \\ &= d_{Y_i}(a^{-1} \mu, \mu). \end{aligned}$$

So far, we have shown

$$d_{g_1 \cdots g_{i-1} Y_i}(\mu, g_1 \cdots g_k \mu) \geq 2K|e_i| - d_{Y_i}(a^{-1} \mu, \mu) - d_{Y_i}(\mu, c \mu).$$

To finish, we prove that the last two terms on the right are each less than  $K/2$ . Since the sign of the  $e_i$  never comes into play, the proof is very similar for either term, so we focus on  $d_{Y_i}(\mu, c \mu)$ . If  $c$  is the identity, then  $d_{Y_i}(\mu, c \mu) = \text{diam}_{Y_i}(\mu) \leq 2 \leq K/2$ . Otherwise  $c = g_{i+1} \cdots g_k$ . Because subwords of minimal words are also minimal,  $c = \phi(\sigma_c)$  for some  $\sigma_c \in G$  with minimal word  $x_{i+1}^{e_{i+1}} \cdots x_k^{e_k}$ , which has strictly less than  $k$  syllables. Let  $x'_1 = x_{i+1}^{e_{i+1}}$  be the first syllable. Applying the induction hypothesis, we have

$$\begin{aligned} d_{Y_{i+1}}(\mu, g_{i+1} \cdots g_k \mu) &= d_{Y_{i+1}}(\mu, c \mu) \\ &= d_{X^{\sigma_c}(x'_1)}(\mu, \phi(\sigma_c) \mu) \\ &\geq K|e_{i+1}|. \end{aligned}$$

By our choice of  $K$ ,  $d_{Y_{i+1}}(\mu, \partial Y_i) = d_{X_{J(i+1)}}(\mu, \partial X_{J(i)}) \leq K/2$ . Since  $g_i$  and  $g_{i+1}$  do not commute,  $Y_i \pitchfork Y_{i+1}$ , so we may apply the triangle inequality to obtain

$$\begin{aligned} d_{Y_{i+1}}(\partial Y_i, g_{i+1} \cdots g_k \mu) &\geq K|e_{i+1}| - d_{Y_{i+1}}(\mu, \partial Y_i) \\ &\geq K|e_{i+1}| - K/2 \geq K/2 \geq 20/2 = 10. \end{aligned}$$

On the other hand, appealing to Proposition 3.5, we know that  $d_{Y_i}(\partial Y_{i+1}, g_{i+1} \cdots g_k \mu) \leq 4$ . By our choice of  $K$ ,  $d_{Y_i}(\mu, \partial Y_{i+1}) = d_{X_{J(i)}}(\mu, \partial X_{J(i+1)}) \leq K/2 - 4$ , so combining, we have

$$\begin{aligned} d_{Y_i}(\mu, c\mu) &\leq d_{Y_i}(\mu, \partial Y_{i+1}) + d_{Y_i}(\partial Y_{i+1}, c\mu) \\ &= d_{Y_i}(\mu, \partial Y_{i+1}) + d_{Y_i}(\partial Y_{i+1}, g_{i+1} \cdots g_k \mu) \\ &\leq K/2 - 4 + 4 = K/2. \end{aligned}$$

The entire argument can be mirrored for  $d_{Y_i}(a^{-1}\mu, \mu)$ , starting with the observation that either  $a^{-1}$  is the identity or  $a^{-1} = g_\ell^{-1} \cdots g_1^{-1}$ , where  $g_\ell$  and  $g_i$  do not commute.

To summarize, we have shown

$$d_{g_1 \cdots g_{i-1} Y_i}(\mu, g_1 \cdots g_k \mu) \geq 2K|e_i| - K/2 - K/2 \geq K|e_i|,$$

completing the induction for statement 1.  $\square$

*Proof of Statement 2.* We will now show that  $X^\sigma$  is an order-preserving injection. The base case for the induction is when  $\sigma$  has one syllable, and then the conclusion is trivially verified. We assume that the conclusion holds for elements  $\sigma$  with at most  $k-1$  syllables, and prove that it also holds for elements with  $k$  syllables.

The subwords  $w_{init} = x_1^{e_1} \cdots x_{k-1}^{e_{k-1}}$  and  $w_{term} = x_2^{e_2} \cdots x_k^{e_k}$  of  $w$  are clearly minimal representatives of the elements  $\sigma_{init}, \sigma_{term} \in G$  they represent. Furthermore, the partial order on the syllables of  $w_{init}$  and  $w_{term}$  is the restriction of the partial order on the syllables of  $w$ .

By the inductive hypothesis, the conclusion of the theorem holds for  $\sigma_{init}$  and  $\sigma_{term}$ . By construction we have

$$X^\sigma(x_i^{e_i}) = \begin{cases} X^{\sigma_{init}}(x_i^{e_i}) & \text{if } i \neq k \\ \phi(x_1^{e_1})(X^{\sigma_{term}}(x_i^{e_i})) & \text{if } i \neq 1 \end{cases}.$$

If  $i$  is neither 1 nor  $k$ , then the two defining expressions are indeed equal. In particular, notice that to establish injectivity, we need only show that  $X^\sigma(x_1^{e_1}) \neq X^\sigma(x_k^{e_k})$ .

Suppose  $x_i^{e_i} \prec x_j^{e_j}$  for two syllables of  $\sigma$ . If  $j \neq k$ , then both are syllables of  $\sigma_{init}$  and hence by induction  $X^{\sigma_{init}}(x_i^{e_i}) \prec X^{\sigma_{init}}(x_j^{e_j})$ . Thus  $X^{\sigma_{init}}(x_i^{e_i}) \pitchfork X^{\sigma_{init}}(x_j^{e_j})$  and

$$d_{X^{\sigma_{init}}(x_i^{e_i})}(\mu, \partial X^{\sigma_{init}}(x_j^{e_j})) \geq 10.$$

Since  $X^{\sigma_{init}}(x_i^{e_i}) = X^\sigma(x_i^{e_i})$  and  $X^{\sigma_{init}}(x_j^{e_j}) = X^\sigma(x_j^{e_j})$ , we have  $X^\sigma(x_i^{e_i}) \prec X^\sigma(x_j^{e_j})$ . If  $i \neq 1$ , then we can make a similar argument using the induction hypothesis applied to  $\sigma_{term}$ . In this case we appeal to condition (4) of Proposition 3.6 to conclude that

$$d_{X^{\sigma_{term}}(x_i^{e_i})}(\partial X^{\sigma_{term}}(x_j^{e_j}), \phi(\sigma_{term})\mu) \leq 4,$$

so that applying  $\phi(x_1^{e_1})$ , this becomes

$$d_{X^\sigma(x_i^{e_i})}(\partial X^\sigma(x_j^{e_j}), \phi(\sigma)\mu) \leq 4,$$



and so  $X^\sigma(x_i^{e_i}) \prec X^\sigma(x_j^{e_j})$ , as required.

Now suppose  $x_1^{e_1} \prec x_k^{e_k}$ . There are two cases.

**Case 1.** There is a syllable  $x_i^{e_i}$  such that  $x_1^{e_1} \prec x_i^{e_i} \prec x_k^{e_k}$ .

Arguing as above, by induction we have  $X^\sigma(x_1^{e_1}) \prec X^\sigma(x_i^{e_i})$  and  $X^\sigma(x_i^{e_i}) \prec X^\sigma(x_k^{e_k})$  in  $\Omega(K, \mu, \phi(\mu))$  and hence by Corollary 3.7 we have that  $X^\sigma(x_1^{e_1}) \prec X^\sigma(x_k^{e_k})$ .

**Case 2.** There is no syllable  $x_i^{e_i}$  such that  $x_1^{e_1} \prec x_i^{e_i} \prec x_k^{e_k}$ .

Then there is a word  $w \in \text{Min}(\sigma)$  of the form:

$$w = w_1 x_1^{e_1} x_k^{e_k} w_2$$

where  $[w_1, x_1] = 1$  and  $[x_k, w_2] = 1$  (and either or both of  $w_i$  may be the empty word). Now:

$$\begin{aligned} d_{X_{J(1)}}(\mu, \partial X^\sigma(x_k^{e_k})) &= d_{X_{J(1)}}(\mu, \phi(w_1)\phi(x_1^{e_1})\partial X_{J(k)}) \\ &= d_{X_{J(1)}}(\mu, \phi(x_1^{e_1})\partial X_{J(k)}) \\ &\geq d_{X_{J(1)}}(\mu, \phi(x_1^{e_1})\mu) - d_{X_{J(1)}}(\phi(x_1^{e_1})\mu, \phi(x_1^{e_1})\partial X_{J(k)}) \\ &\geq 2K|e_1| - (K - 20) \geq 10 \end{aligned}$$

where the second equality comes from the fact that  $w_1$  commutes with  $x_1$ , and so  $\phi(w_1)$  is the identity on  $X_{J(1)}$ . Thus, by Proposition 3.6 (2), we have  $X^\sigma(x_1^{e_1}) \prec X^\sigma(x_k^{e_k})$ .

In particular, if  $x_1^{e_1} \prec x_k^{e_k}$  then  $X^\sigma$  is injective.

All that remains now is to show that if  $x_1^{e_1}$  and  $x_k^{e_k}$  are not comparable by  $\prec$ , then  $X^\sigma(x_1^{e_1}) \neq X^\sigma(x_k^{e_k})$ . If they are not comparable, then  $\sigma$  is represented by a word of the form  $w = w_1 x_1^{e_1} x_k^{e_k} w_2$ , where as above  $[w_1, x_1] = 1$  and  $[x_k, w_2] = 1$ . Furthermore,  $x_1^{e_1}$  and  $x_k^{e_k}$  must commute, and hence it is clear that  $X^\sigma(x_k^{e_k})$  is disjoint from  $X^\sigma(x_1^{e_1})$ . In particular,  $X^\sigma(x_1^{e_1}) \neq X^\sigma(x_k^{e_k})$ .  $\square$

This completes the proof of the Theorem.  $\square$

We can now prove the main theorem.

**Theorem 2.2.** *Given a graph  $\Gamma$  and a nice realization  $\mathbb{X} = \{X_1, \dots, X_n\}$  of  $\Gamma$  in  $S$ , there exists a constant  $C > 0$  with the following property. If  $\mathbb{F} = \{f_1, \dots, f_n\}$  is fully supported on  $\mathbb{X}$  and  $\tau_{X_i}(f_i) \geq C$  for all  $i = 1, \dots, n$ , then the associated homomorphism*

$$\phi_{\mathbb{F}} : G(\Gamma) \rightarrow \text{Mod}(S)$$

*is a quasi-isometric embedding. Furthermore, the orbit map  $G \rightarrow \mathcal{J}(S)$  is a quasi-isometric embedding for both  $d_{\mathcal{J}}$  and  $d_{\text{VP}}$ .*

*Proof.* We first prove that given  $\mu \in \tilde{\mathcal{M}}(S)$ , we can choose  $C > 0$  so that if  $\tau_{X_i}(f_i) \geq C$  for each  $i$ , then the orbit map  $G(\Gamma) \rightarrow \tilde{\mathcal{M}}(S)$  given by  $\sigma \mapsto \phi_{\mathbb{F}}(\sigma)\mu$  is a quasi-isometric embedding. Since the orbit map  $\text{Mod}(S) \rightarrow \tilde{\mathcal{M}}(S)$  is a quasi-isometry, this will suffice to prove the first statement. Let  $K > 0$  be as in the proof of Theorem 5.2 and let  $C = 2K$ .

First observe that for *any* metric space  $(\mathfrak{X}, d)$ , any  $x \in \mathfrak{X}$  and any  $\sigma, \tau \in G(\Gamma)$ , the triangle inequality implies

$$d(\sigma \cdot x, \tau \cdot x) \leq A d_G(\sigma, \tau)$$

as long as  $A \geq \max\{d(s_i \cdot x, x)\}_{i=1}^n$  (here  $s_i$  are the generators for  $G$ ).

In particular, given  $\mu \in \tilde{\mathcal{M}}(S)$ , to prove that the orbit map to  $\tilde{\mathcal{M}}(S)$  is a quasi-isometry, it suffices to find  $A \geq 1$  and  $B \geq 0$  so that

$$d_G(1, \sigma) \leq A d_{\tilde{\mathcal{M}}}(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)) + B$$

for all  $\sigma \in G$  (then we further increase  $A$  if necessary so that  $A \geq \max\{d_{\tilde{\mathcal{M}}}(\mu, \phi_{\mathbb{F}}(s_i)\mu)\}_{i=1}^n$ ).

Since  $K \geq K_0$ , from Theorem 3.1 there exists  $A$  and  $B$  so that for all  $\sigma \in G$ ,

$$\sum_{X \in \Omega(K, \mu, \phi_{\mathbb{F}}(\sigma)(\mu))} d_X(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)) \leq A d_{\tilde{\mathcal{M}}}(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)) + B.$$

On the other hand, if we let  $w = x_1^{e_1} \dots x_k^{e_k} \in \text{Min}(\sigma)$  then by Theorem 5.2, since  $\tau_{X_i}(f_i) \geq C = 2K$  (and since  $K \geq 1$ ) we have

$$\begin{aligned} d_G(1, \sigma) &= \sum_{i=1}^k |e_i| \leq \sum_{i=1}^k K |e_i| \\ &\leq \sum_{i=1}^k d_{X^\sigma(x_i)}(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)) \\ &\leq \sum_{X \in \Omega(K, \mu, \phi_{\mathbb{F}}(\mu))} d_X(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)). \\ &\leq A d_{\tilde{\mathcal{M}}}(\mu, \phi_{\mathbb{F}}(\sigma)(\mu)) + B \end{aligned}$$

which completes the proof of the first statement.

The proof of the statements regarding Teichmüller space are essentially identical. For this, we observe that the topological types of the surfaces in  $X^\sigma(\text{syl}(\sigma))$  are the same as those of  $\mathbb{X}$ , and hence all are nonannular. That is,  $X^\sigma(\text{syl}(\sigma)) \subset \Omega_n(K, \mu, \phi_{\mathbb{F}}(\mu))$ . So, the above proof can be carried out replacing the use of Theorem 3.1 with the use of Theorems 3.2 and 3.3.  $\square$

## 6 Elements of the Constructed Subgroups

We now assume the hypothesis of Theorem 2.2 (and hence also Theorem 5.2) on  $\Gamma$ ,  $\mathbb{X} = \{X_1, \dots, X_n\}$ ,  $C = 2K > 0$ , and  $\mathbb{F} = \{f_1, \dots, f_n\}$ , and let

$$\phi_{\mathbb{F}} : G(\Gamma) \rightarrow \text{Mod}(S)$$

denote the associated homomorphism. In this section we describe, in terms of the Thurston classification, the mapping class image of any  $\sigma \in G(\Gamma)$ . In particular, we identify all pseudo-Anosov elements in the image.

Conjugate elements in  $G(\Gamma)$  map to conjugate elements in  $\text{Mod}(S)$ , and conjugation preserves mapping class type, displacing the support of a mapping class by the homeomorphism corresponding to the conjugating element. Therefore to understand the image of  $\sigma \in G(\Gamma)$ , we may assume it is an element with the minimal number of syllables among members of its conjugacy class (there may be more than one such element, but we just pick one). We represent  $\sigma$  by a word  $w \in \text{Min}(\sigma)$ . By changing the indices if necessary, we can assume that  $w$  is a word

in the first  $r = r(\sigma)$  generators  $s_1, \dots, s_r$ , and  $r$  is the least number of generators needed to write  $w$  (note that a conjugate also having the minimal number of syllables will be written in terms of this same set of generators).

**Remark.** In what follows, we will always assume that the indices on the generators are of this type for the particular element  $\sigma$  we are interested in.

We write  $\text{Fill}(X_1, \dots, X_r)$  to denote the minimal union of subsurfaces, ordered by inclusion, which contains all of the subsurfaces  $X_1, \dots, X_r$ . Alternatively,  $\text{Fill}(X_1, \dots, X_r)$  is the unique union of subsurfaces containing  $X_1 \cup \dots \cup X_r$  with the property that for every essential curve  $\gamma$  contained in it, the projection to at least one of  $X_1, \dots, X_r$  is nontrivial. Write  $\text{Fill}_{\mathbb{X}}(\sigma) = \text{Fill}(X_1, \dots, X_r)$ .

Now, if  $\sigma' = \sigma_0 \sigma \sigma_0^{-1}$ , then we define  $\text{Fill}_{\mathbb{F}}(\sigma') = \phi_{\mathbb{F}}(\sigma_0)(\text{Fill}_{\mathbb{X}}(\sigma))$ . Note that if  $\sigma_0$  is the identity so  $\sigma' = \sigma$ , then  $\text{Fill}_{\mathbb{F}}(\sigma) = \text{Fill}_{\mathbb{X}}(\sigma)$  depends only on  $\mathbb{X}$ , whereas otherwise, it may also depend on  $\mathbb{F}$ .

It follows easily from the discussion above that for any  $\sigma$ ,  $\phi_{\mathbb{F}}(\sigma)$  is supported on  $\text{Fill}_{\mathbb{F}}(\sigma)$ . That is,  $\phi_{\mathbb{F}}(\sigma)$  is represented by a homeomorphism which is the identity outside  $\text{Fill}_{\mathbb{F}}(\sigma)$ . In this section, we prove the following.

**Theorem 6.1.** *Suppose  $\Gamma$ ,  $\mathbb{X} = \{X_1, \dots, X_n\}$ ,  $C = 2K > 0$ ,  $\mathbb{F} = \{f_1, \dots, f_n\}$  satisfy the hypotheses of Theorem 2.2, and let*

$$\phi_{\mathbb{F}} = \phi : G(\Gamma) \rightarrow \text{Mod}(S)$$

*denote the associated homomorphism. Then  $\phi(\sigma)$  is pseudo-Anosov on each component of  $\text{Fill}_{\mathbb{F}}(\sigma)$ . In particular,  $\phi(\sigma)$  is pseudo-Anosov if and only if  $\text{Fill}_{\mathbb{F}}(\sigma) = S$ .*

Before we begin the proof, we explain a few reductions which will greatly simplify the exposition. First, as remarked above, we need only consider the case that  $\sigma$  has the minimal number of syllables among all its conjugates, so we assume this is the case from now on. Therefore  $\text{Fill}_{\mathbb{F}}(\sigma) = \text{Fill}_{\mathbb{X}}(\sigma)$ .

Next, we wish to further reduce to the case that  $\text{Fill}_{\mathbb{X}}(\sigma)$  is connected. To describe this reduction, first let  $\Gamma'$  denote the subgraph spanned by  $s_1, \dots, s_r$ . Since the other generators play no role in this discussion, we assume as we may, that  $\Gamma' = \Gamma$ . Let  $\Gamma^c$  be the *complement* of  $\Gamma$ . That is,  $\Gamma^c$  is the graph with the same vertex set as  $\Gamma$  and where two vertices span an edge in  $\Gamma^c$  if and only if they do not span an edge in  $\Gamma$ . Note that generators/vertices  $s_i$  and  $s_j$  in different components of  $\Gamma^c$  commute. Therefore, we may write  $\sigma = \sigma_1 \cdots \sigma_\ell$ , where  $\ell$  is the number of components of  $\Gamma^c$  and each  $\sigma_i$  is in the group generated by vertices in a single component of  $\Gamma^c$ . In particular,  $[\sigma_i, \sigma_j] = 1$  for all  $i$  and  $j$ .

Now observe that the vertices of a path in  $\Gamma^c$  corresponds to a chain of overlapping subsurfaces in  $S$ , and hence the components of  $\text{Fill}_{\mathbb{X}}(\sigma)$  correspond precisely to the components of  $\Gamma^c$ . In fact, one easily checks that each  $\sigma_i$  also has the least number of syllables in its conjugacy class, and  $\{\text{Fill}_{\mathbb{X}}(\sigma_1), \dots, \text{Fill}_{\mathbb{X}}(\sigma_\ell)\}$  is precisely the set of components of  $\text{Fill}_{\mathbb{X}}(\sigma)$ . So, restricting attention to one of the subwords  $\sigma_i$ , we may assume that  $\text{Fill}_{\mathbb{X}}(\sigma)$  is connected.

Finally, we note that we can in fact restrict to the case that  $\text{Fill}_{\mathbb{X}}(\sigma) = S$ . To see that this is possible, note that  $\phi(G(\Gamma))$  is the identity outside  $S' = \text{Fill}_{\mathbb{X}}(\sigma)$ . So,  $\phi$  “restricts” to a homomorphism

$$\hat{\phi} : G(\Gamma) \rightarrow \text{Mod}(S')$$

and  $\hat{\phi}(\sigma)$  is pseudo-Anosov if and only if  $\phi(\sigma)$  is pseudo-Anosov on  $S' = \text{Fill}_{\mathbb{X}}(\sigma)$ .

We now set out to prove Theorem 6.1 assuming (1) that  $\sigma$  has the least number of syllables in its conjugacy class and (2)  $\text{Fill}_{\mathbb{X}}(\sigma) = S$ . The proof makes use of the partial order on  $\text{syl}(\sigma)$  and  $\text{syl}(\sigma^n)$ , and the order-preserving injection  $X^\sigma(\cdot)$  of the previous section. Regarding these, let us set down a series of lemmas.

**Lemma 6.2.** *For  $\sigma$  as above,  $\text{Fill}(X^\sigma(\text{syl}(\sigma))) = S$ .*

*Proof.* Fix a minimal word  $x_1^{e_1} \cdots x_k^{e_k} \in \text{Min}(\sigma)$ . Given a curve  $\gamma \subset S$ , we must show that  $\gamma$  intersects some  $X^\sigma(x_i^{e_i}) = \phi(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}})X_{J(i)}$ . As  $\text{Fill}_{\mathbb{X}}(\sigma) = \text{Fill}(X_1, \dots, X_r) = S$ , the curve  $\gamma$  intersects some  $X_j$ . Let  $i$  be the minimal index such that  $\gamma$  intersects  $X_{J(i)}$ .

If  $i = 1$ , then  $\gamma$  intersects  $X_{J(1)} = X^\sigma(x_1^{e_1})$  and the lemma holds. Else, notice that  $\phi(x_1^{e_1} \cdots x_{i-1}^{e_{i-1}})(\gamma) = \gamma$ . Hence  $\gamma$  intersects  $X^\sigma(x_i^{e_i})$ .  $\square$

**Lemma 6.3.** *For  $\sigma$  as above,  $w \in \text{Min}(\sigma)$ , and  $n \in \mathbb{Z}$  we have  $w^n \in \text{Min}(\sigma^n)$ .*

*Proof.* Clearly  $w^n$  represents the element  $\sigma^n$ ; what needs to be shown is that this word is minimal.

Write  $w = x_1^{e_1} \cdots x_k^{e_k}$  and assume that  $w^n$  is not a minimal word representing  $\sigma^n$ . Thus we have a sequence of the three moves described in Section 4 which reduces the number of syllables in  $w^k$ . We can label the syllables of  $w^n$  by  $x_{1,1}^{e_1} \cdots x_{1,k}^{e_k} x_{2,1}^{e_1} \cdots x_{n,k}^{e_k}$  where each block  $x_{j,1}^{e_1} \cdots x_{j,k}^{e_k} = w$ . As  $w$  is minimal, each of the  $e_i \neq 0$ , hence in order to reduce the number of syllables of  $w^k$  we have some sequence of type (3) moves followed by a type (2) move.

Also as  $w$  is minimal, the type (2) cannot occur between syllables of the form  $x_{j,i}^{e_i}$  and  $x_{j,i'}^{e_{i'}}$ . Therefore, after applying some type (3) moves, we have a type (2) move between syllables of the form  $x_{j,i}^{e_i}$  and  $x_{j',i'}^{e_{i'}}$  where  $j < j'$ . We claim that we can assume that  $j' = j + 1$ . For if not, then  $[x_i, x_\ell] = 1$  for all  $\ell$  and hence after a sequence of type (3) moves we could apply the move  $x_i^{e_i} x_{i'}^{e_{i'}} \mapsto x_i^{e_i + e_{i'}}$ , contradicting the fact that  $w$  is minimal.

As the set of indices  $\ell$  such that  $[x_{j,i}, x_\ell] = 1$  and the set of indices  $\ell$  such that  $[x_{j',i'}, x_\ell] = 1$  are the same, the above assumptions give a sequence of type (3) moves on  $w$  such that brings  $w$  to a word of the form  $x_{i'}^{e_{i'}} x_1^{e_1} \cdots x_k^{e_k} x_i^{e_i}$ . But now conjugating  $\sigma$  by  $x_{i'}^{e_{i'}}$  results in an element with fewer syllables than  $\sigma$  which contradicts our assumption that  $|\text{syl}(\sigma)|$  is minimal among conjugates of  $\sigma$ . Thus  $w^n \in \text{Min}(\sigma^n)$ .  $\square$

The above lemma allows us to define syllable shift maps  $\sigma_n: \text{syl}(\sigma^n) \rightarrow \text{syl}(\sigma^{n+1})$  by  $\sigma_n(x_{j,i}^{e_i}) = x_{j+1,i}^{e_i}$  using the notation from the proof of the lemma. Notice, the maps  $\sigma_n$  preserve the partial order. For  $n > m$  we use the notation  $\sigma_{m,n} = \sigma_{n-1} \cdots \sigma_m$ . The map  $\sigma_{m,n}: \text{syl}(\sigma^m) \rightarrow \text{syl}(\sigma^n)$  shifts syllables by  $n - m$  blocks and also preserves the partial order. The lemma also allows us to view  $\text{syl}(\sigma^m)$  as a subset of  $\text{syl}(\sigma^n)$ , if  $m \leq n$ , via the obvious inclusion of  $w^m$  as a prefix of  $w^n$ .

Under the assumption that  $\text{Fill}(X_1, \dots, X_r) = S$ , we have that  $\Gamma^c$  is connected. In particular for any syllable  $x_i^{e_i}$ , there is another syllable  $x_j^{e_j}$  such that  $[x_i, x_j] \neq 1$ .

**Lemma 6.4.** *For  $\sigma$  as above, and all  $x_i^{e_i} \in \text{syl}(\sigma)$ , we have  $x_i^{e_i} \prec \sigma_{1,2}(x_i^{e_i}) \in \text{syl}(\sigma^2)$ .*

*Proof.* This is similar to the proof of Lemma 6.3. If the conclusion is wrong, then  $[x_i, x_j] = 1$  for all syllables  $x_j^{e_j} \in \text{syl}(\sigma)$ . This contradicts the fact that  $\Gamma^c$  is connected.  $\square$

**Lemma 6.5.** *For  $\sigma$  as above, and all  $x_i^{e_i}, x_j^{e_j} \in \text{syl}(\sigma)$ , there exists  $n$ ,  $1 \leq n \leq r+1$ , such that  $x_i^{e_i} \prec \sigma_{1,n}(x_j^{e_j})$ . In particular, for all syllables  $x_i^{e_i}, x_j^{e_j} \in \text{syl}(\sigma)$  we have  $x_i^{e_i} \prec \sigma_{1,r+1}(x_j^{e_j})$ .*

*Proof.* Fix a minimal sequence of generators  $x_i = x_{i_1}, \dots, x_{i_m} = x_j$  such that  $[x_{i_\ell}, x_{i_{\ell+1}}] \neq 1$ . Such a sequence exists as  $\Gamma^c$  is connected. Further, notice that  $m \leq r$ . We will prove the lemma by induction on  $m$ . Specifically, we will prove that if there is a path of length  $m$  between  $v_{J(i)}$  and  $v_{J(j)}$  in  $\Gamma^c$ , then  $x_i^{e_i} \prec \sigma_{1,m+1}(x_j^{e_j})$ .

Suppose  $m = 1$ , hence as generators  $x_i = x_j$ . The case when  $x_i^{e_i} = x_j^{e_j}$  as syllables in  $\text{syl}(\sigma)$  is covered by Lemma 6.4. Else, we must have that  $x_i^{e_i} \prec x_j^{e_j}$  or  $x_j^{e_j} \prec x_i^{e_i}$ . In the first case using Lemma 6.4 we have  $x_i^{e_i} \prec x_j^{e_j} \prec \sigma_{1,2}(x_j^{e_j})$ . In the second case, if  $x_i^{e_i} \not\prec \sigma_{1,2}(x_j^{e_j})$  we can argue as in the proof of Lemma 6.3 that  $|\text{syl}(\sigma)|$  is not minimal among conjugates of  $\sigma$ .

Now by induction, we have that  $x_i^{e_i} \prec \sigma_{1,m}(x_{i_{m-1}}^{e_{i_{m-1}}})$ . Since  $[x_{i_{m-1}}, x_j] \neq 1$ , we must have  $\sigma_{1,m}(x_{i_{m-1}}^{e_{i_{m-1}}}) \prec \sigma_{1,m+1}(x_j^{e_j})$ . Hence  $x_i^{e_i} \prec \sigma_{1,m}(x_{i_{m-1}}^{e_{i_{m-1}}}) \prec \sigma_{1,m+1}(x_j^{e_j})$ . This completes the proof of the lemma.  $\square$

*Proof of Theorem 6.1.* We assume  $\sigma$  is as above, so  $\text{Fill}_{\mathbb{X}}(\sigma) = S$ , and prove  $\phi(\sigma)$  is pseudo-Anosov. For this, it suffices to prove the following.

**Claim.** *For every integer  $N > 0$  we have*

$$d_S(\partial X_{J(1)}, \phi(\sigma^{N(2r+1)})(\partial X_{J(1)})) \geq N.$$

Indeed, this claim says that  $\phi(\sigma)$  acts with positive translation length on  $\mathcal{C}(S)$  as required.

*Proof of claim.* According to Lemma 6.5 we have

$$x_1^{e_1} \prec \sigma_{1,r+1}(x_j^{e_j}) \prec \sigma_{r+1,2r+1}(\sigma_{1,r+1}(x_1^{e_1})) = \sigma_{1,2r+1}(x_1^{e_1})$$

Now, from the definitions, we see that  $X^{\sigma^n} \circ \sigma_{1,n} = \phi(\sigma^{n-1}) \circ X^\sigma$  for all  $n > 1$ , and since  $X^\sigma$  and  $X^{\sigma^n}$  are order preserving by Theorem 5.2 we also have

$$X_{J(1)} \prec \phi(\sigma^{r+1})(X^\sigma(x_j^{e_j})) \prec \phi(\sigma^{2r+1})(X_{J(1)})$$

This implies that no curve  $\gamma \subset S$  is disjoint from both  $\partial X_{J(1)}$  and  $\phi(\sigma^{2r+1})\partial X_{J(1)}$ . Indeed, suppose otherwise. According to Lemma 6.2, the collection of subsurfaces  $\phi(\sigma^{r+1})X^\sigma(\text{syl}(\sigma))$  fill  $S$ , so there is some subsurface, say  $\phi(\sigma^{r+1})X^\sigma(x_j^{e_j})$  where  $\gamma$  has nonempty projection. Hence,

$$d_{\phi(\sigma^{r+1})X^\sigma(x_j^{e_j})}(\partial X_{J(1)}, \phi(\sigma^{2r+1})\partial X_{J(1)}) \leq 4.$$

However, since  $X_{J(1)} \prec \phi(\sigma^{r+1})(X^\sigma(x_j^{e_j})) \prec \phi(\sigma^{2r+1})(X_{J(1)})$  it follows from Proposition 3.6 and the triangle inequality that

$$d_{\phi(\sigma^{r+1})X^\sigma(x_j^{e_j})}(\partial X_{J(1)}, \phi(\sigma^{2r+1})\partial X_{J(1)}) \geq K - 8 > 4$$

which is a contradiction.

By the same reasoning, no curve  $\gamma$  can be disjoint from more than one of the following sets of curves

$$\{\partial X_{J(1)}, \phi(\sigma^{2r+1})(\partial X_{J(1)}), \dots, \phi(\sigma^{N(2r+1)})(\partial X_{J(1)})\}.$$

On the other hand, since  $X_{J(1)} \prec \phi(\sigma^{\ell(r+1)}(X_{J(1)})) \prec \phi(\sigma^{N(2r+1)}(X_{J(1)}))$  for all  $0 < \ell < N$ , Proposition 3.6 and the triangle inequality again imply

$$d_{\phi(\sigma^{\ell(r+1)}(X_{J(1)}))}(\partial X_{J(1)}, \phi(\sigma^{N(2r+1)}(X_{J(1)}))\partial X_{J(1)}) \geq K - 8 \geq K_0$$

where the last inequality comes from the choice of  $K$  in the proof of Theorem 5.2.

Now, according to Theorem 3.4 any geodesic in  $\mathcal{C}(S)$  from  $\partial X_{J(1)}$  to  $\phi(\sigma^{N(2r+1)}(X_{J(1)}))\partial X_{J(1)}$  must contain a curve disjoint from each  $\phi(\sigma^{\ell(2r+1)}(X_{J(1)}))\partial X_{J(1)}$ , for each  $\ell = 0, \dots, N$ . Since these curves must all be distinct by the previous paragraph, we see that this geodesic contains at least  $N + 1$  vertices, so

$$d_S(\partial X_{J(1)}, \phi(\sigma^{N(2r+1)}(X_{J(1)}))\partial X_{J(1)}) \geq N$$

as required.

This completes the proof of the claim, and also the proof of the Theorem. □

□

## 7 Surface subgroups

In this final section we prove the following corollary of Theorem 2.2 and briefly discuss surface subgroups of right-angled Artin subgroups of the mapping class group.

**Corollary 1.3.** *For any closed surface  $S$  of genus at least 3 and any  $h \geq 2$ , there exist infinitely many nonconjugate genus  $h$  surface subgroups of  $\text{Mod}(S)$ , each of which act cocompactly on some quasi-isometrically embedded hyperbolic plane in the Teichmüller space  $\mathcal{T}(S)$ , with either of the standard metrics.*

*Proof.* Let  $\Gamma$  be the cyclic graph of length 5 and  $G(\Gamma)$  the associated right-angled Artin group. It was shown in [11] that  $G(\Gamma)$  contains a quasi-isometrically embedded genus 2 surface subgroup, and hence surface subgroups of all genus  $h \geq 2$  (it had been previously shown to contain a genus 5 surface subgroup in [40]). As described in [11], this example has a nice description as follows.

Suppose the generators of  $G(\Gamma)$  are  $a, b, c, d, e$  with  $[e, a] = [a, b] = [b, c] = [c, d] = [d, e] = 1$ . Then the homomorphism from the fundamental group of a genus two surface to  $G(\Gamma)$  is described by Figure 6 as follows. The figure shows a system of curves on the surface with labels from the set  $\{a, b, c, d, e\}$  and transverse orientations. Choosing a basepoint in the complement of the curve system shown, a loop will cross the curves in the system, and one reads off an element of  $G(\Gamma)$  according to the curves one crosses, and in which direction (crossing in the direction opposite the given transverse orientation, one should read an inverse of the generator); see [11] for more details.

In Section 2.4 we observed that  $\Gamma$  has a nice realization  $\mathbb{X} = \{X_1, \dots, X_5\}$  in any closed surface  $S$  of genus  $g \geq 3$ . Let  $C > 0$  be the constant from Theorem 2.2 and  $\mathbb{F} = \{f_1, \dots, f_5\}$  be any mapping classes fully supported on  $\mathbb{X}$  with  $\tau_{X_i}(f_i) \geq C$ . For every  $n > 0$  let  $\mathbb{F}^n = \{f_1^n, \dots, f_5^n\}$  so that we also have  $\tau_{X_i}(f_i^n) \geq nC$ . The family of right-angled Artin subgroups  $\phi_{\mathbb{F}^n}(G(\Gamma))$  necessarily contains infinitely many distinct conjugacy classes—observe that the proof of Theorem 2.2 implies that the minimal translation length on  $\mathcal{T}(S)$  of any element of  $\phi_{\mathbb{F}^n}(G(\Gamma))$  is tending to infinity as  $n \rightarrow \infty$ . Similarly, the set of surface subgroups described above, thought of as subgroups of  $\text{Mod}(S)$  via the homomorphisms  $\phi_{\mathbb{F}^n}$ , have minimal translation length on  $\mathcal{T}(S)$




Figure 6: A curve system on a genus 2 surface which defines an embedding into  $G(\Gamma)$ , where  $\Gamma$  is the cyclic graph of length 5.

tending to infinity as  $n \rightarrow \infty$ . Consequently, there are infinitely many pairwise nonconjugate genus  $h$  surface subgroups.

That each of these stabilizes a quasi-isometrically embedded hyperbolic plane  $\mathbb{H} \subset \mathcal{T}(S)$  follows from the fact that the surface group itself is quasi-isometric to  $\mathbb{H}$ , and the orbit map defines a quasi-isometric embedding by Theorem 2.2. The surface group clearly acts cocompactly on this plane.  $\square$

It follows that these surface groups all have positive translation length on Teichmüller space. However, as we have already mentioned, they cannot be purely pseudo-Anosov. In fact, for surface subgroups of right-angled Artin groups, this is always the case.

**Proposition 7.1.** *Suppose  $G(\Gamma) < \text{Mod}(S)$  is a right-angled Artin subgroup and  $\pi_1(\Sigma) < G(\Gamma)$  is a surface subgroup. Then as a subgroup of  $\text{Mod}(S)$ ,  $\pi_1(\Sigma)$  contains a nontrivial reducible element.*

*Proof.* As was shown in [11], every homomorphism from a surface group  $\pi_1(\Sigma)$  into a right-angled Artin group  $G(\Gamma)$  arises as in the proof of the previous corollary. That is, there is a curve system on  $\Sigma$ , each curve is endowed with a transverse orientation, the components are labeled by generators of  $G(\Gamma)$ , and the homomorphism is obtained by taking a loop and reading off the generators as one crosses the curves in the system.

Furthermore, one may assume that each of the curves in the system is essential, and if the homomorphism  $H \rightarrow G(\Gamma)$  is injective then these curves cut  $\Sigma$  into disks. Now, consider a loop  $\gamma$  which runs parallel to, without crossing, one of the curves in the system. Call this curve  $\eta_1$  and suppose the associated generator of  $G(\Gamma)$  is  $s_1$ . The loop  $\gamma$  crosses other curves  $\eta_{i_1}, \dots, \eta_{i_k}$  and so determines some word  $s_{i_1}^{\pm 1} \cdots s_{i_k}^{\pm 1}$ , which is the image of  $\gamma$  in  $G(\Gamma)$ . Choosing  $\gamma$  to run very close to  $\eta_1$ , we can assume that the curves  $\eta_{i_1}, \dots, \eta_{i_k}$  which  $\gamma$  crosses are also nontrivially intersected by  $\eta_1$ . As noted in [11], each of the associated generators  $s_{i_1}, \dots, s_{i_k}$  must commute with  $s_1$ , and be different from it.

Now we are essentially done. The image of  $\gamma$  is an element which commutes with  $s_1$ , and in fact, the image of  $\gamma$  and  $s_1$  in  $G(\Gamma)$  generate a subgroup isomorphic to  $\mathbb{Z}^2$ . If  $G(\Gamma) < \text{Mod}(S)$ , then the image of  $\gamma$  in  $\text{Mod}(S)$  has centralizer which contains  $\mathbb{Z}^2$ . As is well known, the image of  $\gamma$  cannot be pseudo-Anosov; see [21].  $\square$

**Remark.** In fact, the assumption that  $\pi_1(\Sigma)$  is a surface group can be relaxed considerably. Indeed, a similar proof shows that any finitely presented 1-ended subgroup of a right-angled

Artin group  $G(\Gamma) < \text{Mod}(S)$  contains a reducible element.

In [25], Koberda observes that  $\text{Mod}(S_g)$  is not commensurable with a right-angled Artin group if  $g \geq 3$  (in fact, he proves the stronger statement that  $\text{Mod}(S_g)$  cannot virtually embed in a right-angled Artin group). This is also true for genus 2 as the following shows.

**Proposition 7.2.** *The group  $\text{Mod}(S_2)$  is not commensurable with a right-angled Artin group.*

*Proof.* Suppose  $\text{Mod}(S_2)$  is commensurable with  $G(\Gamma)$ , with  $\Lambda$  isomorphic to a finite index subgroup of both. Let  $\pi_1(\Sigma) < \text{Mod}(S_2)$  be a surface subgroup as constructed in [27]. In this surface group, there is exactly one element of  $\pi_1(\Sigma)$ , up to conjugacy and powers, which is not pseudo-Anosov. Moreover, this one element represents a simple closed curve  $\alpha$  on  $\Sigma$ .

Now,  $\pi_1(\Sigma) \cap \Lambda$  is a finite index subgroup of  $\pi_1(\Sigma)$  and so corresponds to a covering space  $p : \tilde{\Sigma} \rightarrow \Sigma$ , and we write  $\pi_1(\tilde{\Sigma}) < \pi_1(\Sigma)$  for the image under  $p_*$ . Note that the reducible elements of  $\pi_1(\tilde{\Sigma})$  in  $\text{Mod}(S_2)$  represent a finite set of pairwise disjoint simple closed curves on  $\tilde{\Sigma}$ , namely  $p^{-1}(\alpha)$ .

On the other hand, a closer inspection of the proof of the previous proposition shows that, viewing  $\Lambda < G(\Gamma)$ , there are actually *two* elements  $\gamma_1, \gamma_2 \in \pi_1(\tilde{\Sigma})$  which represent curves on  $\tilde{\Sigma}$  which nontrivially intersect, whose centralizers in  $G(\Gamma)$  contain  $\mathbb{Z}^2$ . These must represent reducible elements in  $\text{Mod}(S)$ , and this is a contradiction.  $\square$

**Remark.** This same proof also works to show that the mapping class group of an  $n$ -punctured sphere, with  $n \geq 6$  is not commensurable with any right-angled Artin group. The point is that the examples from [27] can be chosen to descend to the quotient by the hyper-elliptic involution, and then one of the punctures can be erased (with the exception of the genus 2 case).

## References

- [1] M. F. Atiyah. The signature of fibre-bundles. In *Global Analysis (Papers in Honor of K. Kodaira)*, pages 73–84. Univ. Tokyo Press, Tokyo, 1969.
- [2] Jason Behrstock. Asymptotic geometry of the mapping class group and teichmueller space. *Geom. Topol.*, 10:1523–1578, 2006.
- [3] Jason Behrstock, Bruce Kleiner, Yair Minsky, and Lee Mosher. Geometry and rigidity of mapping class groups. Preprint, arXiv:0801.2006, 2008.
- [4] Jason Behrstock and Yair Minsky. Centroids and the rapid decay property in mapping class groups. Preprint, arXiv:0810.1969, 2008.
- [5] Brian H. Bowditch. Atoroidal surface bundles over surfaces. *Geom. Funct. Anal.*, 19(4):943–988, 2009.
- [6] Jeffrey F. Brock. The Weil-Petersson metric and volumes of 3-dimensional hyperbolic convex cores. *J. Amer. Math. Soc.*, 16(3):495–535 (electronic), 2003.
- [7] Ruth Charney. An introduction to right-angled Artin groups. *Geom. Dedicata*, 125:141–158, 2007.



- [8] John Crisp and Benson Farb. The prevalence of surface subgroups in mapping class groups. In preparation.
- [9] John Crisp and Luis Paris. The solution to a conjecture of Tits on the subgroup generated by the squares of the generators of an Artin group. *Invent. Math.*, 145(1):19–36, 2001.
- [10] John Crisp, Michah Sageev, and Mark Sapir. Surface subgroups of right-angled Artin groups. *Internat. J. Algebra Comput.*, 18(3):443–491, 2008.
- [11] John Crisp and Bert Wiest. Embeddings of graph braid and surface groups in right-angled Artin groups and braid groups. *Algebr. Geom. Topol.*, 4:439–472, 2004.
- [12] John Crisp and Bert Wiest. Quasi-isometrically embedded subgroups of braid and diffeomorphism groups. *Trans. Amer. Math. Soc.*, 359(11):5485–5503, 2007.
- [13] Benson Farb and Lee Mosher. Convex cocompact subgroups of mapping class groups. *Geom. Topol.*, 6:91–152 (electronic), 2002.
- [14] Koji Fujiwara. Subgroups generated by two pseudo-Anosov elements in a mapping class group. I. Uniform exponential growth. In *Groups of diffeomorphisms*, volume 52 of *Adv. Stud. Pure Math.*, pages 283–296. Math. Soc. Japan, Tokyo, 2008.
- [15] G. González-Díez and W. J. Harvey. Surface groups inside mapping class groups. *Topology*, 38(1):57–69, 1999.
- [16] C. McA. Gordon, D. D. Long, and A. W. Reid. Surface subgroups of Coxeter and Artin groups. *J. Pure Appl. Algebra*, 189(1-3):135–148, 2004.
- [17] E. Green. Graph Products of Groups, Thesis. The University of Leeds (1990).
- [18] Ursula Hamenstädt. Word hyperbolic extensions of surface groups. Preprint, arXiv:math.GT/0505244, 2005.
- [19] Susan Hermiller and John Meier. Algorithms and geometry for graph products of groups. *J. Algebra*, 171(1):230–257, 1995.
- [20] Tim Hsu and Daniel T. Wise. On linear and residual properties of graph products. *Michigan Math. J.*, 46(2):251–259, 1999.
- [21] Nikolai V. Ivanov. *Subgroups of Teichmüller modular groups*, volume 115 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1992. Translated from the Russian by E. J. F. Primrose and revised by the author.
- [22] Richard P. Kent, IV and Christopher J. Leininger. Shadows of mapping class groups: capturing convex cocompactness. *Geom. Funct. Anal.*, 18(4):1270–1325, 2008.
- [23] Sang-hyun Kim. Co-contractions of graphs and right-angled Artin groups. *Algebr. Geom. Topol.*, 8(2):849–868, 2008.
- [24] Sang-hyun Kim. On right-angled Artin groups without surface subgroups. *Groups Geom. Dyn.*, 4(2):275–307, 2010.
- [25] Thomas Koberda. Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups. Preprint.

- [26] K. Kodaira. A certain type of irregular algebraic surfaces. *J. Analyse Math.*, 19:207–215, 1967.
- [27] C. J. Leininger and A. W. Reid. A combination theorem for Veech subgroups of the mapping class group. *Geom. Funct. Anal.*, 16(2):403–436, 2006.
- [28] Michele Linch. A comparison of metrics on Teichmüller space. *Proc. Amer. Math. Soc.*, 43:349–352, 1974.
- [29] Kefeng Liu, Xiaofeng Sun, and Shing-Tung Yau. Canonical metrics on the moduli space of Riemann surfaces. I. *J. Differential Geom.*, 68(3):571–637, 2004.
- [30] Kefeng Liu, Xiaofeng Sun, and Shing-Tung Yau. Canonical metrics on the moduli space of Riemann surfaces. II. *J. Differential Geom.*, 69(1):163–216, 2005.
- [31] Darren Long, Alan Reid, and Morwen Thistlethwaite. Zariski dense surface subgroups in  $SL(3, \mathbf{Z})$ . Preprint.
- [32] Johanna Mangahas. Uniform uniform exponential growth of subgroups of the mapping class group. *Geom. Funct. Anal.*, 19(5):1468–1480, 2010.
- [33] H. A. Masur and Y. N. Minsky. Geometry of the complex of curves. II. Hierarchical structure. *Geom. Funct. Anal.*, 10(4):902–974, 2000.
- [34] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.
- [35] John McCarthy. A “Tits-alternative” for subgroups of surface mapping class groups. *Trans. Amer. Math. Soc.*, 291(2):583–612, 1985.
- [36] Curtis T. McMullen. The moduli space of Riemann surfaces is Kähler hyperbolic. *Ann. of Math. (2)*, 151(1):327–357, 2000.
- [37] Lee Mosher. A hyperbolic-by-hyperbolic hyperbolic group. *Proc. Amer. Math. Soc.*, 125(12):3447–3455, 1997.
- [38] Kasra Rafi. A combinatorial model for the Teichmüller metric. *Geom. Funct. Anal.*, 17(3):936–959, 2007.
- [39] Claas E. Röver. On subgroups of the pentagon group. *Math. Proc. R. Ir. Acad.*, 107(1):11–13 (electronic), 2007.
- [40] Herman Servatius, Carl Droms, and Brigitte Servatius. Surface subgroups of graph groups. *Proc. Amer. Math. Soc.*, 106(3):573–578, 1989.
- [41] Stephen Wang. Representations of surface groups and right-angled Artin groups in higher rank. *Algebr. Geom. Topol.*, 7:1099–1117, 2007.
- [42] Sai-Kee Yeung. Quasi-isometry of metrics on Teichmüller spaces. *Int. Math. Res. Not.*, (4):239–255, 2005.

Department of Mathematics  
Allegheny College  
Meadvile, PA 16335  
E-mail: [mclay@allegheny.edu](mailto:mclay@allegheny.edu)

Department of Mathematics  
University of Illinois at Urbana-Champaign  
Urbana, IL 61801  
E-mail: [clein@math.uiuc.edu](mailto:clein@math.uiuc.edu)

Department of Mathematics  
University of Michigan  
Ann Arbor, MI 48109  
E-mail: [mangahas@umich.edu](mailto:mangahas@umich.edu)